Format: Article

Prevalent human gut _Enterocloster_ spp. produce urolithin A from its dietary precursor

using a molybdenum-dependent dehydroxylase

Reilly Pidgeon[1], Sacha Mitchell[1], Michael Shamash[2], Layan Suleiman[1], Lharbi Dridi[1], Corinne

Maurice[2], Bastien Castagner[1*]

[1] Department of Pharmacology & Therapeutics, McGill University, 3655 Prom. Sir-William-

Osler, Montreal, Quebec, H3G 1Y6, Canada.

[2] Department of Microbiology & Immunology, McGill University, 3775 University Street,

Montreal, Quebec, H3A 2B4, Canada

Abstract

Urolithin A is a polyphenol derived from the multi-step metabolism of dietary ellagitannins by the human gut microbiota that can affect host health. Most, but not all, individuals harbor a microbiota capable of urolithin A production; however, the enzymes that dehydroxylate its dietary precursor (urolithin C) are unknown. Here, we use a combination of untargeted transcriptomics, proteomics, and comparative genomics to uncover an inducible urolithin C dehydroxylase (*ucd*) operon in *Lachnospiraceae*. Biochemical characterization and structural modeling of proteins encoded by *ucd* demonstrated that urolithin C dehydroxylation was NADH-dependent, required strict anaerobic conditions, and used urolithin C as an electron acceptor. By combining functional assays in complex fecal communities with publicly available metagenomic data, our study reveals that both urolithin C-metabolizing bacteria and *ucd* operon genes are widely distributed in gut metagenomes and likely comprise keystone species in the metabolism of urolithins by the human gut microbiota.

Main

The human gut microbiota is a collection of trillions of microorganisms that colonize the intestinal tract and play a pivotal role in host health and disease [1]. Gut bacteria help maintain homeostasis by regulating host immune cell activity, gut barrier integrity, and nutrient availability [2]. One of the main mediators of microbiota-host interactions are microbial metabolites. Gut bacteria possess an immense metabolic repertoire (nearly 1000-fold more protein coding sequences than the human genome [3]) to perform four main classes of reactions: hydrolysis, conjugation, cleavage and reduction [4-7]. These reactions are ubiquitous in the gut environment have been linked to microbiota-dependent activities whereby bacteria can alter the availability and efficacy of therapeutic drugs [8-10], modify host bile acids [6,11,12], and metabolize diet-derived compounds [13-15].

Diet is a strong modulator of the composition and function of the gut microbiota [16-19]. Diet-derived polyphenols are a diverse class of plant secondary metabolites found in fruits, vegetables, and nuts (reviewed in [20]) that are poorly absorbed by the host and reach the large intestine relatively intact [7,21]. Ellagitannins are a sub-group of polyphenols that belong to the family of hydrolysable tannins and are characterized by a central glucose (open-chain or cyclic forms) linked to a variety of pyrogallol-like moieties [20]. Depending on microbiota composition, ellagitannins can be hydrolyzed and reduced by bacteria into bioactive metabolites (e.g., ellagic acid, urolithins, nasutins) according to different metabolic phenotypes characterized by the terminal metabolites observed in biological fluids [22] (Extended Data Fig. 1).

Urolithin A (uroA) is the most common terminal metabolite of ellagitannin metabolism [23]. Clinical trials in healthy individuals have demonstrated that uroA is safe, bioavailable, and can be detected in its aglycone, glucuronidated, and sulfated forms in plasma [23,24]. Once absorbed,

uroA can impact host health by improving mitochondrial function (mitophagy) [24-26], protecting

gut barrier integrity [27,28], and inhibiting inflammation [29]. Most notably, uroA modulates immune

function through the activation of Pink1 and aryl hydrocarbon receptor (AhR) signaling

pathways, thus showing promise as a postbiotic to treat age-related conditions [25,27,30,31].

While urolithin metabolism is prevalent in human populations, only few gut bacteria have

reported urolithin metabolism activity [32-34]. Most urolithin metabolizers belong to the

Eggerthellaceae family (*Gordonibacter urolithinfaciens*, *Gordonibacter pamelaeae*, *Ellagibacter*

isourolithinifaciens) and can perform multiple metabolic steps in the urolithin metabolism

pathway, yielding either uroC or isourolithin A (isouroA) from ellagic acid [35]. Recently, certain

members of the *Enterocloster* spp. (*Lachnospiraceae* family) were reported to dehydroxylate

urolithin C (uroC) to uroA and isouroA to urolithin B (uroB) both *in vitro* and *in vivo* [36,37]. These

findings shed light on the minimal bacterial community required for complete metabolism of

ellagic acid to uroA; however, the genes and enzymes that catalyze these dehydroxylation

reactions remain unknown (Fig. 1A).

Here, we used a multi-omics enzyme identification framework to uncover uroC dehydroxylase

(*ucd*) genes and enzymes in *Enterocloster* spp. and their relative, *Lachnoclostridium pacaense*.

We found that the Ucd enzyme complex specifically dehydroxylates 9-hydroxy urolithins and

that both metabolizing species and *ucd* genes are prevalent and actively transcribed in human

feces during metabolism. Our study sheds light on the genetic basis underlying the complex

reciprocal interactions between urolithins, the gut microbiota, and the host.

Results

A subset of *Enterocloster* species converts urolithin C to urolithin A *in vitro*.

Members of the *Enterocloster* spp. have previously been shown to metabolize

(dehydroxylate) uroC *in vitro* [36] and *in vivo* [37] under anaerobic conditions (Fig. 1A, full metabolic

pathway in Extended Data Fig. 1). To determine the prevalence of uroC metabolism within this

genus, we incubated all available *Enterocloster* spp. type strains (Methods) with uroC and

quantified urolithin concentrations by liquid chromatography-mass spectrometry (LC-MS). Of

the tested bacteria, only *E. asparagiformis*, *E. bolteae*, and *E. citroniae* dehydroxylated uroC to

produce uroA (Fig. 1B). Interestingly, uroC metabolism was not predicted by phylogeny, as

uroC-metabolizing species did not cluster based on 16S rRNA genes, genomes, or proteomes

(Fig. 1C, Extended Data Fig. 2A,B), suggesting gain or loss of metabolic gene clusters (MGCs)

throughout the evolution of *Enterocloster* spp. Based on these results, we chose to perform

more in-depth analysis on *E. asparagiformis* and *E. bolteae* to identify the MGCs involved in

uroC dehydroxylation.

Figure 1. Urolithin C metabolism by *Enterocloster* spp. is not predicted by phylogeny.

A) Reaction scheme of uroC dehydroxylation by gut resident *Enterocloster* spp. via unknown
enzymes. **B)** LC-MS screen of *Enterocloster* spp. type strains for dehydroxylation activity. UroC

(100 μM) was added to cultures (in mABB+H media) at the start of growth and urolithins were

extracted after 24 h anaerobic incubation, then analyzed by LC-MS. Left: Representative

chromatograms (λ = 305 nm) for each experimental group (from one representative biological

replicate). The same scale was used for each chromatogram. Right: Quantification of urolithin

peak areas relative to a salicylic acid internal standard (IS) (n = 3 biological replicates). Data

are represented as mean ± SEM. **C)** Phylogenetic tree of tested *Enterocloster* spp. type strain

16S rRNA sequences constructed using the Genome-to-Genome Distance Calculator (GGDC)

Phylogeny Server [55]. Maximum likelihood (ML) tree inferred under the GTR+GAMMA model

and rooted by midpoint-rooting. The branches are scaled in terms of the expected number of

substitutions per site. The numbers above the branches are support values when larger than

60% from ML (left) and maximum parsimony (right) bootstrapping. The GenBank accession

numbers are provided to the right of each taxon. Source data and statistical details are provided

as a Source data file.

A putative urolithin C dehydroxylase metabolic gene cluster is upregulated upon

urolithin C treatment.

To understand when uroC metabolism machinery was being expressed, we first sought

to characterize the kinetics of uroC dehydroxylation. Therefore, a simultaneous growth and

metabolism experiment was designed, whereby uroC was spiked into cultures during the

exponential phase of growth and metabolites were measured by LC-MS (Fig. 2A). Treatment

with uroC did not affect growth of either bacterium compared to the DMSO control (Fig. 2B). In

both bacteria, quantitative conversion of uroC to uroA occurred within 4 h post-spike-in (Fig.

2C,D), demonstrating that metabolism in rich media is fast and robust.

We next sought to determine whether uroC metabolism is inducible or constitutive. To test for

inducibility, both bacteria were treated with DMSO or uroC during exponential growth, then

washed and resuspended in PBS, yielding resting cells unable to synthesize new proteins.

Metabolism of uroC to uroA was inducible as only resting cells originating from bacteria grown

with uroC were capable of uroA production (Fig. 2E). We therefore performed RNA-sequencing

to compare gene expression in DMSO and uroC-treated cultures of *E. asparagiformis* and *E.*

bolteae. Since uroA was detected in both bacterial cultures 2 h post spike-in (Fig. 2C), this

timepoint was selected to isolate mRNA. RNA sequencing of uroC-induced cultures revealed a

distinct MGC, which we term the uroC dehydroxylase (*ucd*) cluster, that was highly and

differentially expressed ($\log_2$FC > 8) in both *E. asparagiformis* (Fig. 2F) and *E. bolteae* (Fig.

2G). In both bacteria, these MGCs contained adjacent genes that were expressed to similar

$\log_2$FC values: a xanthine dehydrogenase family protein subunit M, a (2Fe-2S)-binding protein,

and a molybdopterin-dependent oxidoreductase (Fig. 2F,G insets). These genes, which belong

to the xanthine dehydrogenase family of proteins, will hereafter be referred to as *ucdC* (for

coenzyme), *ucdF* (for ferredoxin), and *ucdO* (for oxidoreductase), respectively.

Figure 2. Urolithin C treatment upregulates a putative dehydroxylase metabolic gene cluster.

A) Experimental design of uroC (100 µM) spike-in experiments during the exponential phase of growth. For each biological replicate in this design, growth (B), metabolism (C,D), and RNA-

seq (F-G) results are matched. **B)** Growth curve (optical density (OD) at 620 nm) of DMSO or uroC-spiked *E. asparagiformis* (*Ea*) and *E. bolteae* (*Eb*) type strain cultures according to the design in (A). 200 μL of culture were sampled at each timepoint and OD_{620} was measured in a 96-well plate (n = 4 biological replicates). The same sampled culture was then frozen and extracted for analysis by LC-MS (C,D). **C)** Representative chromatograms (λ = 305 nm) of cultures sampled 2 h post-spike-in (from one representative biological replicate). The same scale was used for each chromatogram. **D)** Quantification of urolithin concentrations from peak areas relative to a salicylic acid internal standard (IS) over 4 h in uroC-spiked *Ea* and *Eb* type strain cultures (n = 4 biological replicates). **E)** Quantification of urolithin A concentrations in DMSO- or uroC-treated *Ea* and *Eb* resting cell suspensions. Resting cell suspensions were prepared from *Ea* and *Eb* cells grown with either DMSO or 50 μM uroC. The cells were washed and resuspended in PBS to halt the production of new enzymes, then treated with DMSO or uroC (n = 3 biological replicates). **F,G)** Manhattan plots of genes altered by uroC treatment in **F)** *Ea* and **G)** *Eb* based on DESeq2 analysis (n = 4 biological replicates). Data points are colored according to their adjusted p-value (based on the Benjamini-Hochberg-corrected Wald statistic). Grey, p-adj ≥ 0.05. Red or blue, p-adj < 0.05 for *Ea* and *Eb*, respectively. The most highly and differentially expressed genes by RNA-seq are depicted in the insets, which show the genomic organization around the differentially expressed genes (generated from the NCBI Sequence Viewer). Genes in the insets are colored according to their log_2FC values. NCBI accessions for select proteins encoded by highlighted genes are provided. Data are represented as mean ± SEM in (B,D,E). FC, fold change (uroC/DMSO); Source data and statistical details are provided as a Source data file.

Presence of *ucd* metabolic gene cluster homologs in genomes predicts urolithin C metabolism by gut bacteria.

We wondered whether novel metabolizers of uroC could be discovered based on nucleotide sequence homology to the *ucd* MGC. Homology searches using the *E. bolteae ucd* MGC sequence confirmed that uroC-metabolizing *Enterocloster* spp. (*E. asparagiformis*, *E. bolteae*, and *E. citroniae*) possessed homologs of the *ucdCFO* genes with a similar organization (Extended Data Fig. 3). In addition, the gut bacterium *Lachnoclostridium pacaense* (or *Clostridium pacaense*) [38] was identified as a hit (Extended Data Fig. 3). The type strain of this bacterium (CCUG 71489T = Marseille-P3100) was closely related to *Enterocloster* spp. based on 16S rRNA, whole genome, and whole proteome phylogenies (Fig. 3A, Extended Data Fig. 2A,B, respectively). *L. pacaense* possessed genomic sequences with high homology (86.5% nucleotide identity) and identical functional annotations to the *E. bolteae ucd* MGC sequence

(Fig. 3B). When incubated with uroC, *L. pacaense* CCUG 71489T quantitatively produced uroA

(Fig. 3C,D). Other urolithin and catechol-metabolizing bacteria belonging to the

Eggerthellaceae family lack uroC dehydroxylase activity [33,35], which correlates with the absence

of *ucd*-like operons in their genomes (Fig. 3B). These comparative genomics data indicate that

the presence of a *ucd* MGC in genomes predicts uroC metabolism by gut bacteria.

Figure 3. Urolithin C metabolism correlates with *ucd* metabolic gene cluster prevalence

in gut bacteria.

A) Phylogenetic tree of *Enterocloster* spp., *Lachnoclostridium pacaense*, and catechol-
metabolizing *Eggerthellaceae* type strain 16S rRNA sequences constructed using the Genome-
to-Genome Distance Calculator (GGDC) Phylogeny Server [55]. Maximum likelihood (ML) tree
inferred under the GTR+GAMMA model and rooted by midpoint-rooting. The branches are
scaled in terms of the expected number of substitutions per site. The numbers above the
branches are support values when larger than 60% from ML (left) and maximum parsimony
(right) bootstrapping. Bacteria that convert uroC to uroA are labeled with grey squares. **B)** NBCI
Multiple Sequence Aligner viewer hits for BLASTn searches using the *E. bolteae* DSM 15670
ucd MGC nucleotide sequence as a query against the NCBI refseq_genomes database. Only
hits with ≥ 90 % query coverage and species-level taxonomic resolution are displayed with %
identity to the query sequence. Domain annotations for each gene are denoted below according
to InterPro annotations for corresponding proteins. ND, Not detected; Moco, Molybdenum
cofactor. **C,D)** *In vitro* metabolism of uroC by *Lachnoclostridium pacaense* CCUG 71489T.
UroC (100 µM) was added to cultures (in mABB+H media) at the start of growth and urolithins
were extracted after 24 h anaerobic incubation, then analyzed by LC-MS. **C)** Representative
chromatograms (λ = 305 nm) for each experimental group (from one representative biological
replicate). The same scale was used for each chromatogram. **D)** Quantification of urolithin peak
areas relative to a salicylic acid internal standard (IS) (n = 3 biological replicates). Data are

represented as mean ± SEM. Source data and statistical details are provided as a Source data
file.

The *E. bolteae ucd* metabolic gene cluster is organized in an operon.

We next sought to characterize the genomic organization of the *ucd* MGC in *E. bolteae*.

Based on the proximity, sense, and expression levels of each of the three genes by RNA-seq

(Fig. 2G), we hypothesized that all three genes in the MGC were organized in an operon. We

designed a gene-specific RT-PCR assay that would enable the detection of full-length

polycistronic *ucdCFO* genes using cDNA from DMSO- or uroC-treated *E. bolteae* as a template

(Fig. 4A). An amplicon of the expected size (~3.6 kb) was detected only in cDNA derived from

uroC-treated *E. bolteae*, validating the inducibility of these genes (Fig. 4B). Long-read

sequencing of the obtained amplicon yielded a sequence corresponding to the *E. bolteae*

ucdCFO MGC with 100% identity (Supplementary Information). Using an independent set of *E.*

bolteae cultures, we then performed RT-qPCR on DMSO- or uroC-treated *E. bolteae* cultures

with all three genes in the putative operon as targets. Similar to our RNA-seq results, all three

genes were significantly induced (Fig. 4C) relative to DMSO controls and were expressed at

the same levels (Fig. 4D). These results indicate that the *ucdCFO* genes are transcribed as a

single polycistronic mRNA and therefore form a uroC-inducible operon.

The *E. bolteae ucd* operon is induced by other 9-hydroxy urolithins

Next, we aimed to determine the substrate scope of the *ucd* operon. Multiple urolithins

possess pyrogallol, catechol, and hydroxyl structural motifs that are dehydroxylated at various

positions by gut bacteria (Fig. 4E, Extended Data Fig. 1). Interestingly, *E. bolteae* only

metabolizes the 9-position hydroxyl group of urolithins and does not require adjacent hydroxyl

groups for dehydroxylation (Fig. 4F,G, Extended Data Fig. 4A,B). Since dehydroxylation of

urolithins in *E. bolteae* is position specific, we hypothesized that the *ucd* operon would also be

induced by these substrates. We performed RT-qPCR on DMSO-, uroM6-, uroC-, or isouroA-

treated *E. bolteae* cultures using the *ucdO* gene as a target. While each urolithin significantly

induced the expression of the *ucd* operon compared to DMSO (Fig. 4H), no significant

difference in expression was observed between urolithin treatments (Fig. 4I). In addition, *E.*

bolteae resting cell suspensions induced with uroC were capable of dehydroxylating uroM6 and

isouroA, indicating that the same proteins induced by uroC can metabolize structurally similar

9-hydroxy urolithins (Extended Data Fig. 4C,D). Thus, it is likely that the same metabolic

enzymes, encoded by the *E. bolteae ucd* operon, are acting on 9-hydroxy urolithins.

Figure 4. The *E. bolteae ucd* is organized in a three-gene operon that is induced by 9-hydroxy urolithins.

A) Primer design for RT-PCR (B) experiment targeting the *E. bolteae* (*Eb*) *ucd* MGC. The same reverse primer was used for both the reverse transcription step and the subsequent PCR reaction. **B)** 1% agarose gel image of RT-PCR amplicons using primers (A) that span the full-length *Eb ucd* MGC (from one biological replicate). Growing *Eb* cultures were treated with DMSO or uroC (100 µM) for 2 h before RNA isolation and reverse transcription. NTC, no template control. **C,D)** RT-qPCR expression of each gene in the *Eb ucd* MGC: molybdopterin-dependent oxidoreductase (*ucdO*), (2Fe-2S)-binding protein (*ucdF*), xanthine dehydrogenase family protein subunit M (*ucdC*). Growing *Eb* cultures were treated with DMSO or uroC (100

μM) for 2 h before RNA isolation and reverse transcription (n = 3 biological replicates). **C)** Differential gene expression comparing DMSO vs. uroC is displayed as target-specific ΔCt (C_t Target Gene - C_t dnaK Reference Gene) values. Data are presented as individual ΔCt values with lines connecting paired biological replicates; multiple paired t tests with Benjamini, Krieger, and Yekutieli FDR correction (Q set to 0.05); *, q < 0.05. **D)** Gene expression profiles of each target gene in the *Eb ucd* MGC displayed as $\log_2$FC (equivalent to -ΔΔCt, where ΔΔCt = $ΔC_t$ uroC - $ΔC_t$ DMSO). Data are presented as individual $\log_2$FC values with lines connecting paired biological replicates; repeated-measures one-way ANOVA with Tukey's multiple comparisons test; ns, not significant. **E)** Chemical structures of urolithin M-6, urolithin C, and isourolithin A highlighting the common 9-position hydroxyl group that gets dehydroxylated. **F)** Quantification of extracted ion chromatogram (EIC) peak areas from *Eb* cultures sampled after 24 h of growth with 100 μM uroM6 (n = 3 biological replicates). **G)** Quantification of urolithin peak areas from *Eb* cultures sampled after 24 h of growth with 100 μM isouroA (n = 3 biological replicates). **H,I)** RT-qPCR expression of the *Eb ucdO* gene. Growing *Eb* cultures were treated with DMSO, uroM6, uroC, or isouroA (100 μM) for 2 h before RNA isolation and reverse transcription (n = 3 biological replicates). **H)** Differential *Eb ucdO* gene expression comparing DMSO, uroM6, uroC, and isouroA is displayed as target-specific ΔCt (C_t MoO Gene - C_t dnaK Reference Gene) values. Data are presented as individual ΔCt values with lines connecting paired biological replicates (from the same pre-spike culture); repeated-measures one-way ANOVA with Dunnett's multiple comparisons test; **, p < 0.01; ***, p < 0.001. **I)** Gene expression profile of the *Eb ucdO* gene in different urolithin treatment groups displayed as $\log_2$FC (equivalent to -ΔΔCt, where ΔΔCt = $ΔC_t$ urolithin - $ΔC_t$ DMSO). Data are presented as individual $\log_2$FC values with lines connecting paired biological replicates; repeated-measures one-way ANOVA with Tukey's multiple comparisons test. ns, not significant; FC, fold change; FDR, false discovery rate. Source data and statistical details are provided as a Source data file. The uncropped and unannotated gel image is provided in the Supplementary Information.

A molybdopterin cofactor biosynthetic gene cluster is upregulated upon urolithin C treatment.

Molybdoenzymes are organized into 3 main superfamilies (DMSO reductases, sulfite oxidases, and xanthine oxidases) that have specific cofactor requirements (bis-MGD, Moco, MCD, respectively) [39]. In addition to the three genes in the *ucd* operon, we observed a significant increase ($\log_2$FC ≥ 2.6) in 9 molybdopterin cytosine dinucleotide cofactor (MCD) biosynthesis genes upon uroC treatment (Extended Data Fig. 5A). These 9 genes, which recapitulate the function of 10 genes in *E. coli* [39], are involved in molybdenum cofactor biosynthesis (*moaAC, mogA, moeA*), molybdate ion transport (*modABCE*), cytosine addition to the molybdenum cofactor (*mocA*), and MCD cofactor insertion into the active site (*xdhC*)

(Extended Data Fig. 5B) [40]. All 9 genes cluster in the genomes of *E. asparagiformis* and *E.*

bolteae and are organized into 2 adjacent operons (Extended Data Fig. 5C) that are induced

upon uroC treatment. Based on sequence homology to *E. coli* oxidoreductases, proteins

encoded by the *ucd* operon belong to the xanthine dehydrogenase (Xdh) family, which are

reported to bind an MCD cofactor in the oxidoreductase subunit (XdhB) of the active XdhAB

enzyme complex [40]. These findings imply that uroC dehydroxylation is MCD-dependent, which

differs from the bis-MGD requirement of catechol dehydroxylases in *Eggerthellaceae* [10,14].

The UcdCFO complex enables anaerobic electron transport from NADH to uroC

Since oxidoreductases utilise a variety of cofactors and coenzymes for catalytic activity,

we sought to determine the redox coenzymes and conditions necessary for uroC

dehydroxylation. Therefore, we performed metabolism assays using crude lysates from uroC-

induced *E. bolteae*. As crude lysates alone did not metabolize uroC (Fig.5A), various redox

coenzymes (NADPH, NADH, and FAD) were added to lysates to promote uroC

dehydroxylation. Only NADH-treated lysates yielded quantitative dehydroxylation of uroC to

uroA compared to the no cofactor control (Fig. 5A). Interestingly, the addition of free FAD

partially inhibited uroC dehydroxylation in NADH-treated lysates (Fig. 5A), likely by decreasing

the free NADH pool. NADPH, which differs from NADH by a phosphate group on the 2'-OH

group of the adenosine moiety, was unable to promote uroC dehydroxylation, indicating some

specificity in the redox cofactors necessary to sustain catalysis. Aerobic incubation of crude

lysates supplemented with NADH completely inhibited uroC dehydroxylation (Fig. 5B),

indicating that the active enzyme complex requires a strictly anaerobic environment for

dehydroxylation, as has been demonstrated for various metalloenzymes [41].

We attempted heterologous expression of the *E. bolteae ucd* operon in *E. coli* to validate

its function; however, all expression experiments were unsuccessful due to inclusion body

formation (data not shown). This lack of solubility and activity likely results from the choice of

heterologous host and from the complex assembly of active molybdoenzymes [40]. To confirm

that *ucd* operon-encoded proteins were expressed in *E. bolteae* crude lysates, we performed

untargeted proteomics and compared protein expression upon DMSO or uroC treatment.

Indeed, all 3 proteins encoded by the *ucd* operon (UcdC, UcdF, and UcdO) were the most

differentially expressed proteins in the uroC treatment group (Fig. 5C). In addition, proteins

involved in MCD biosynthesis were also strongly increased upon uroC treatment (Fig. 5C,D,

Extended Data Fig. 5A), pointing to the coordination between MCD biosynthesis and active

oxidoreductase assembly. These multi-omics datasets implicate all three genes and their

products in the metabolism of uroC to uroA, as demonstrated by the strong positive correlation

between transcript and protein differential expression (Fig. 5D).

To gain an understanding of the structural organization of proteins encoded by the *ucd*

operon, we performed modeling using AlphaFold2 [42,43]. Structures of each protein encoded by

the *E. bolteae ucd* operon (Fig. 5E) were superposed onto published X-ray crystal structures of

oxidoreductases with similar folds (using the UcdO amino acid sequence as a query) [44], yet

from different taxonomic domains: carbon monoxide dehydrogenase from *Afipia*

carboxidovorans [45] and xanthine dehydrogenase from *Bos taurus* [46]. The 3 proteins encoded

by the *ucd* operon formed subunits in an oxidoreductase complex with a similar quaternary

structure to the published crystal structures (Fig. 5F, Extended Data Fig. 6A,B). The predicted

quaternary structure of the UcdCFO enzyme complex supported a complete electron transport

chain whereby electrons would flow from reduced FAD to two 2Fe-2S clusters, then to the MCD

cofactor, and finally to uroC as the terminal electron acceptor (Fig. 5G, Extended Data Fig.

6C,D). Using homology modeling, we further identified the putative uroC binding site in UcdO,

which overlaps with the salicylic acid ligand in the *Bos taurus* xanthine dehydrogenase

(Extended Data Fig. 6E). This putative uroC binding site contains multiple tyrosine (Y375, Y538,

Y624, Y632), tryptophan (W345), and phenylalanine (F458, F464) residues that could form π-

π stacking interactions with uroC (Extended Data Fig. 6F), orienting it towards the molybdenum

cofactor.

Figure 5. The UcdCFO complex enables anaerobic electron transport from NADH to

uroC.

A) Quantification of urolithin peak concentrations in crude uroC-induced *Eb* lysates re-treated
with DMSO or uroC (350 µM) and various coenzymes (n = 3 biological replicates). Data are
represented as mean ± SEM. NADPH, nicotinamide adenine dinucleotide phosphate; NADH,
nicotinamide adenine dinucleotide; FAD, flavin adenine dinucleotide. **B)** Quantification of
urolithin peak concentrations in crude uroC-induced *Eb* lysates re-treated with DMSO or uroC
(350 µM) and NADH in anaerobic or aerobic environments (n = 3 biological replicates). **C)**
Volcano plot of untargeted proteomics analysis on DMSO or uroC-treated *Eb* (n = 3 biological
replicates). Data points are colored according to their significance (Fisher's exact test with
Benjamini-Hochberg correction for multiple comparisons). Grey, p-adj ≥ cutoff p-value
(0.00048). Blue, p-adj < cutoff p-value (0.00048). **D)** Scatter plot showing the correlation

between gene and protein expression ($\log_2$FC values) induced in uroC-treated *Eb* using the datasets in Fig. 2G and Fig. 5C, respectively. The non-parametric Spearman rank correlation test was used for statistical analysis. **E)** Domains of genes in the *ucd* operon based on InterPro annotations. **F)** Quaternary structure prediction of the proteins encoded by the *Eb ucd* operon. AlphaFold2 structures for each protein were superposed onto the X-ray crystal structure of PDB 1ZXI (carbon monoxide dehydrogenase from *Afipia carboxidovorans* OM5). **G)** Small molecule ligands from PDB 1ZXI in the superposed UcdCFO model form a complete electron transport chain from FAD to two 2Fe-2S clusters to a molybdopterin cytosine dinucleotide cofactor, which can then reduce uroC (terminal electron acceptor). Source data and statistical details are provided as a Source data file.

UroC-metabolizing species and *ucd* genes are prevalent and correlate with uroC

metabolism in human fecal samples.

We next wondered whether uroC-metabolizing *Enterocloster* spp. and their *ucd* operons were prevalent and active in human fecal samples. We first utilized uniformly processed metagenomic data from the curatedMetagenomicData R package [47]. After filtering for fecal samples (86 studies, n = 21,030 subjects), we counted the prevalence of at least one uroC-metabolizing species and at least one *ucd* gene homolog (Methods). The prevalence of both features was variable across studies (Fig. 6A for studies with >200 participants, Extended Data Fig. 7A for all studies). Combining all studies, the prevalence of at least one uroC-metabolizing species and at least one *ucdCFO* gene homolog was 9,343/21,030 (44.9%) and 4,356/21,030 (20.7%), respectively. *E. bolteae* was the most prevalent and abundant uroC-metabolizing species detected in gut metagenomes (Extended Data Fig. 7B,C). These findings suggest that uroC-metabolizing *Enterocloster* spp. and *ucd* operon genes are prevalent in human fecal metagenomic samples and reflect the variable urolithin metabolism profiles (metabotypes) in the general population [23,48].

Next, we performed *ex vivo* metabolism assays to determine whether *Enterocloster* spp. could metabolize uroC in the context of a complex community. Fecal slurries from 10 healthy individuals were first profiled according to their uroC metabotypes (Fig. 6B) [22]. Individuals

clustered into metabotypes A (only uroA produced), B (uroA and isouroA/uroB), and 0 (no terminal urolithin metabolites). While we observed metabotypes A and B in uroC-metabolizing fecal slurries, all slurries produced some amount of uroA from uroC. Stools JL73, TR06, and YE96 displayed variable metabolism patterns and did not metabolize uroC in some experiments, likely reflecting differences in activity between aliquots of feces (Fig. 6B). We then repeated metabolism assays using fecal slurries from all 10 healthy individuals and extracted urolithins, DNA, and RNA from each culture. In this experiment, only 5/10 fecal slurries metabolized uroC to uroA (Fig. 6C). We hypothesized that differences in metabotypes could be explained by microbial composition. Therefore, V1-V9 16S rRNA sequencing was performed on fecal slurries. Both DMSO- and uroC-treated fecal slurries within individuals had similar microbial compositions but showed differences in composition between individuals (Extended Data Fig. 8A,B). Surprisingly, all samples contained 16S rRNA sequences mapping to *E. bolteae*, and many non-metabolizing fecal slurries contained *E. asparagiformis* (Fig. 6C, Extended Data Fig. 8C). We then assayed the treated fecal slurries for the presence of the *ucd* operon by PCR and found that 10/10 individuals (19/20 conditions) yielded a detectable amplicon of the expected size (~3.6 kb) (Extended Data Fig. 8D,E). These data indicate that the prevalence of uroC-metabolizing *Enterocloster* spp. 16S rRNA and *ucd* operon genes does not predict metabolism in fecal samples .

We then surmised that the *ucd* operon would be transcribed only in fecal slurries actively metabolizing uroC. Using a gene-specific reverse primer that binds to *ucdO*, the full-length *ucd* operon was reverse transcribed and amplified in RNA extracted from DMSO- and uroC-treated fecal slurries. An amplicon (~3.6 kb) corresponding to the *ucd* operon was only detected in uroC-metabolizing fecal slurries when treated with uroC and entirely absent from non-metabolizing slurries (Fig. 6D). This amplicon was absent in no reverse transcriptase controls,

387 indicating no gDNA contamination (Extended Data Fig. 8F,G). These data demonstrate that

388 *ucd* transcription correlates with uroC metabolism in complex fecal communities and that *E.*

389 *bolteae* is keystone species involved in urolithin A production.

Figure 6. The *ucd* operon is prevalent in metagenomes and actively transcribed in

urolithin C-metabolizing human fecal samples.

A) Prevalence of *ucd* operon (at least one gene) and of a uroC-metabolizing species (at least
one species) in fecal metagenomes from the CuratedMetagenomicData R package. Only
studies with ≥200 participants are depicted. All 86 studies are available in Extended Data Fig.
7. **B)** Summary of urolithin concentrations in fecal slurries (n = 10 healthy donors) incubated
with 100 µM uroC for 48 h. Data are representative of 3-6 experimental replicates. **C)** Summary
of urolithin concentrations in fecal slurries (n = 10 healthy donors) incubated with 100 µM uroC
for 48 h. Presence of uroC-metabolizing species and of the *ucd* operon is denoted above the
graph if the bacterium or operon was detected in the uroC-treated fecal slurry (see Extended
Data Fig. 8C,D). Data are representative of 1 replicate where DNA and RNA was also extracted
from fecal slurries. **D,E)** *ucd* gene-specific RT-PCR on fecal microbiota communities from 10
healthy donors using the primer set described in Fig. 4A. Samples are matched to the urolithin
metabolism data in C) and 16S rRNA sequencing in Extended Data Fig. 8A,C. 1% agarose gel
of amplicons derived from **D)** uroC-metabolizing and **E)** non-metabolizing fecal slurries. Bands
corresponding to the *E. bolteae ucd* operon (~3.6 kb) are labeled with red arrows. The no
template control (NTC) is the same for D,E). See Extended Data Fig. 8D,E for the no reverse
transcriptase control PCR reactions on the same samples. Source data are provided as a
Source data file. The uncropped and unannotated gel images are provided in the
Supplementary Information.

Discussion

Here, we identified genes and proteins that are essential for the metabolism of urolithins by gut resident *Lachnospiraceae* through a combination of transcriptomics, comparative genomics, and untargeted proteomics. Our study reveals a novel multi-subunit molybdoenzyme (urolithin C dehydroxylase, Ucd) that catalyzes the dehydroxylation of 9-hydroxy urolithins like uroM6, uroC, and isouroA. Importantly, prevalence analysis in published data and *ex vivo* transcriptomics established *E. bolteae* as a keystone urolithin-metabolizing member of the gut microbiota.

Catechol dehydroxylases are widespread in *Eggerthella lenta* and *Gordonibacter pamelaeae* [10,14,49]. These molybdoenzymes, which belong to the DMSO reductase superfamily (bis-MGD cofactor) [50], dehydroxylate substrates like catechol lignan (Cldh), DOPAC (Dodh), dopamine (Dadh), hydrocaffeic acid (Hcdh), and caffeic acid (Cadh). The UcdCFO enzyme complex differs from catechol dehydroxylases in *Eggerthellaceae* in important ways as it 1) does not require a catechol structural motif for activity, 2) belongs to the xanthine oxidase superfamily (MCD cofactor), and 3) is composed of 3 subunits (UcdCFO) that each lack signal sequences. A recent survey of reductases in gut bacteria established that most respiratory reductases contain N-terminal signal sequences and are translocated across the cytoplasmic membrane, while non-respiratory reductases, which lack signal sequences, remain in the cytoplasm [51]. Based on the absence of signal sequences and the cytoplasmic localization of xanthine dehydrogenases, the *ucd* operon encodes for a non-respiratory reductase (see [51] Supplementary Table 2).

In the host, UroA has reported pharmacological activities both within the gut environment and systemically following absorption. In the gut, uroA can attenuate colitis by increasing the

expression of epithelial tight junction proteins [27] *via* the activation of AhR-Nrf2 pathways [28].

Additionally, uroA can enhance immunotherapy in colorectal cancer models by triggering Pink1-

dependent mitophagy pathways in T cells, improving anti-tumor CD8+ T cell immunity [25]. Once

absorbed by the host, uroA can trigger mitophagy in muscle cells, improving muscle function in

animal models of ageing and Duchenne muscular dystrophy [39,52]. Overall, uroA can enhance

gut barrier integrity, reduce inflammation, and promote mitochondrial health in the host.

While uroA is the most common terminal metabolite following ellagitannin consumption

in humans, its production varies widely [34]. Interestingly, the ability of a fecal sample to produce

uroA from uroC did not correlate with the presence of widespread uroC-metabolizing

Enterocloster spp. or a *ucd* gene homologue, likely owing to poor viability or dead bacteria in

fecal samples. However, the active transcription of the *ucd* operon correlated perfectly with

metabotypes. By identifying the genetic basis for metabolism of uroC, we found a novel

metabolizing species that could not have been predicted based on phylogeny alone. Our data

suggests that *ucd*-containing *Enterocloster* spp., and the closely related *L. pacaense*, are the

main drivers of urolithin A production in the gut microbiota based on their prevalence in

metagenomes and activity in fecal samples. These findings further emphasize the importance

of functional assays such as transcriptomics and *ex vivo* metabolism to understand the

metabolism of xenobiotics by the gut microbiota.

Our study has several limitations. Similar to other molybdoenzyme studies [14], we were

unable to express a soluble, active UcdCFO enzyme complex in a heterologous host to validate

enzyme function. In addition, while we demonstrate that *Enterocloster* spp. actively

dehydroxylate uroC in human fecal samples, we cannot conclude that they are solely

responsible for this activity. Rare, strain-specific urolithin A production has been reported for

Bifidobacterium pseudocatenulatum INIA P815 [34], *Streptococcus thermophilus* FUA329 [53], and

Enterococcus faecium FUA027 [54], which may be a result of horizontal gene transfer since it is

not shared by other members of the taxa.

In conclusion, our studies reveal the genetic basis for urolithin A production by gut

bacteria and broaden our understanding of the molecular mechanisms underlying urolithin

metabotypes in human populations. Since diet can modulate gut microbiota function and host

health, elucidating the xenobiotic metabolism genes encoded by gut bacteria will be key to

developing dietary interventions targeting the gut microbiota.

 Extended Data Figures

Extended Data Figure 1. Ellagitannins are metabolized by gut bacteria.

Reaction scheme of dietary ellagitannin metabolism by the human gut microbiota. Larger ellagitannin structures are hydrolyzed during gut transit, releasing hexahydroxydiphenic acid, which spontaneously lactonizes into ellagic acid. Once in the gut lumen, members of the *Gordonibacter* spp. and *Ellagibacter isourolithinifaciens* can decarboxylate ellagic acid, forming urolithin M-5. The resulting urolithin M-5 can be further dehydroxylated (at the 4,10- or 4,8,10-positions) to uroC or isourolithin A by *Gordonibacter* spp. or *Ellagibacter isourolithinifaciens*, respectively. Compounds colored in light gray are urolithin metabolites that are rarely observed during *ex vivo* metabolism assays on ellagitannins. Once uroC or isouroA are produced, *Enterocloster* spp. can further dehydroxylate the 9-position, yielding uroA or uroB, respectively. The cartoon was generated in BioRender.

Extended Data Figure 2. *Lachnoclostridium pacaense* **clusters with** *Enterocloster* **spp.**

in whole genome and proteome phylogenetic trees.

A) Whole genome phylogenetic tree of *Enterocloster* spp. The tree was inferred with FastME 2.1.6.1 from Genome BLAST Distance Phylogeny (GBDP) distances calculated from genome sequences. The branch lengths are scaled in terms of GBDP distance formula d5. The numbers above branches are GBDP pseudo-bootstrap support values > 60 % from 100 replications, with an average branch support of 36.8 %. The tree was rooted at the midpoint. **B)** Whole proteome phylogenetic tree of *Enterocloster* spp. The tree was inferred with FastME 2.1.6.1 from whole-proteome-based GBDP distances. The branch lengths are scaled via GBDP distance formula d5. Branch values are GBDP pseudo-bootstrap support values > 60 % from 100 replications, with an average branch support of 100.0 %. The tree was rooted at the midpoint.

Extended Data Figure 3. BLASTn searches using the *E. bolteae ucd* MGC genomic

sequence identifies homologues in gut bacteria.

NBCI Multiple Sequence Aligner viewer hits for BLASTn searches using the *E. bolteae* DSM
15670 *ucd* MGC nucleotide sequence as a query against the NCBI refseq_genomes database
(limited to Bacteria). Vertical lines in the sequence alignment represent nucleotide differences
(show differences option) and insertions relative to the query sequence.

Extended Data Figure 4. *E. bolteae* **dehydroxylates urolithins at the 9-position.**

A) Representative extracted ion chromatograms (EIC) of *Eb* cultures sampled after 24 h of growth with 100 µM uroM6 (from one representative biological replicate). The same scale was used for each chromatogram. Analytes of interest and their respective [M-H]⁻ m/z values are provided above the figure. **B)** Representative chromatograms (λ = 305 nm) of *Eb* cultures sampled after 24 h of growth with 100 µM uroC or isouroA (from one biological replicate). The same scale was used for each chromatogram. **C)** Quantification of extracted ion chromatogram (EIC) peak areas in DMSO- or uroC-treated *Eb* resting cell suspensions. Resting cell suspensions were prepared from *Eb* cells grown with either DMSO or 50 µM uroC. The cells were washed and resuspended in PBS to halt the production of new enzymes, then treated with 100 µM uroM6 (n = 3 biological replicates). **D)** Quantification of urolithin concentrations in DMSO- or uroC-treated *Eb* resting cell suspensions. Resting cell suspensions were prepared as described in C) (n = 3 biological replicates). For C-D, data are represented as mean ± SEM. ND, not detected.

A

B

Protein annotation	Role(s)	*Ea* NCBI Accession	*Eb* NCBI Accession	#AA in *Eb*	Domain(s) [positions]
GTP 3',8-cyclase	**MoaA**	WP_007712811.1	WP_002565229.1	355	Radical SAM core [7-238]
Cyclic pyranopterin monophosphate synthase	**MoaC**	WP_007712808.1	WP_002565230.1	164	Molbdopterin cofactor biosynthesis C (MoaC) [14-149]
MOSC domain-containing protein	**MogA**	WP_051409812.1	WP_002565228.1	322	Molbdenum cofactor sulfurase C-terminal (MOSC) [28-153]
Molybdopterin-binding protein / Molybdopterin molybdenumtransferase	**MoeA**	WP_007712805.1	WP_002565231.1	352	MoaB/Mog [177-317]
NTP-transferase domain-containing protein / LysR family transcriptional regulator	**MocA/ModE**	WP_007713105.1	WP_002565224.1	313	MobA-like NTP transferase [5-160] Helix-turn-helix lysR-type [222-277]
ATP-binding cassette domain-containing protein	**ModC**	WP_117778241.1	WP_002565225.1	382	ABC transporter [2-259]
Molybdate ABC transporter permease subunit	**ModB**	WP_007712823.1	WP_002565226.1	223	ABC transmembrane type-1 [6-215]
Molybdate ABC transporter substrate-binding protein	**ModA**	WP_007712821.1	WP_002565227.1	294	Signal [1-25] Molybdate ABC transporter substrate-binding protein [26-294]
XdhC family protein	**XdhC**	WP_007713106.1	WP_002565223.1	343	XdhC- CoxI [12-70] XdhC Rossmann [189-331]

C

Extended Data Figure 5. Urolithin C treatment upregulates molybdopterin cytosine

dinucleotide cofactor biosynthetic gene clusters.

A) Molybdopterin cytosine dinucleotide (MCD) cofactor biosynthetic pathway. The $\log_2$FC
values of MCD cofactor biosynthetic genes upregulated by uroC in both transcriptomics (T) and
proteomics (P) datasets (when available) are provided to the left of the figure for both *Ea* and
Eb. **B)** Table of molybdenum cofactor biosynthetic genes found in the genomes of *Ea* and *Eb*.
Annotations are based on NCBI (GTF files) and UniProt gene/protein names. Roles are
assigned based on required proteins for molybdenum cofactors in the xanthine
oxidase/dehydrogenase family of enzymes. Accessions, primary sequence length, and
annotated domains (with positions within the primary sequence) are also provided. **C)** Genomic
organization of the MCD cofactor biosynthetic genes for *Ea* and *Eb* (generated from the NCBI
Sequence Viewer).

Extended Data Figure 6. The AlphaFold2 model of the *E. bolteae* UcdCFO enzyme

complex has a similar quaternary structure to xanthine dehydrogenase superfamily

crystal structures.

A,B) Structural superposition of AlphaFold2-predicted *Eb* Ucd proteins (right) onto the X-ray crystal structures of **A)** PDB 1ZXI (carbon monoxide dehydrogenase from *Oligotropha carboxidovorans* OM5) X-ray crystal structure (left) and **B)** PDB 3UNI (bovine milk xanthine

dehydrogenase with NADH bound). PDB 1ZXI is colored in shades of green (according to chain
ID), PDB 3UNI is colored in gold (single chain A shown), and the *Eb* Ucd enzyme complex is
colored according to its per-residue confidence score as indicated in the legend. The various
ligands (cofactors, coenzymes, ions, small molecules) of PDB 1ZXI and PDB 3UNI are included
in the respective *Eb* Ucd enzyme models. **C,D)** Cofactors from X-ray crystal structures **C)** PDB
1ZXI and **D)** PDB 3UNI modeled into the AlphaFold2 *Eb* Ucd enzyme complex, showing a
complete electron transport chain from a bound FAD molecule to a molybdopterin cofactor *via*
two 2Fe-2S clusters. **E)** Xanthine dehydrogenase active site from PDB 3UNI. Side chains within
8 Å of the salicylic acid ligand are colored in gold. Residues important for substrate (purine)
binding and catalysis are labeled with their one letter amino acid code and sequence position.
Position E1261* is catalytically important (acts as a general base) and is conserved in
XDH/xanthine oxidase enzymes [56]. **F)** *Eb* Ucd enzyme complex active site modeling. Side
chains within 8 Å of the salicylic acid ligand from PDB 3UNI are colored on the superposed *Eb*
Ucd enzyme complex according to their per-residue confidence score as indicated in the
legend. Residues surrounding the predicted active site are labeled with their one letter amino
acid code and sequence position (in the *Eb* MoO protein). The predicted urolithin binding site
is depicted by the surface (mesh) created by the active site residues. The surface was rendered
using the cavities and pockets only (culled) setting with a cavity detection cutoff of 5 solvent
radii in PyMOL.

Extended Data Figure 7. Urolithin C-metabolizing species and genes are prevalent and correlate with *ucdO* gene abundance in human gut metagenomes.

A) Related to Fig. 6A. Prevalence of at least one *ucd* operon gene (blue bars) and uroC-metabolizing species (white bars) in fecal metagenomes across all 86 studies (in reverse alphabetical order). The number of participants in each study are represented to the right of the

prevalence plot as a heatmap. Details on the study populations (from the
curatedMetagenomicData R package) can be found in the Source Data file. **B)** Prevalence of
uroC-metabolizing species in human fecal metagenomes from the curatedMetagenomicData R
package. Data are reported for 86 studies (N=21,030 individuals) and are colored according to
the species. **C)** Violin plot of the $\log_{10}$(relative abundance) of uroC-metabolizing species in
human fecal metagenomes. The solid horizontal line corresponds to the median and the dashed
white lines correspond to the first and third quartiles. The percentage of zeroes are denoted
below the plotted distributions. Differences between groups were determined using the Kruskal-
Wallis test on untransformed relative abundance values. Significant differences between
groups are denoted by a different lowercase letter above each plot. **D)** Correlation between the
ucdO gene abundance in reads per kilobase per million mapped reads (RPKM) and the relative
abundance of each uroC-metabolizing species in fecal metagenomes. Both values are
illustrated on a $\log_{10}$ scale. Spearman rho (ρ) values are denoted above the scatter plots.

Extended Data Figure 8. Urolithin C-metabolizing species and the *ucd* operon are prevalent in fecal slurries, but only *ucd* transcription correlates with urolithin C metabolism.

A) Stacked bar plot of bacterial percent relative abundance (based on V1-V9 16S rRNA gene sequencing) in DMSO- or uroC-treated fecal slurries from 10 healthy donors (from one experimental replicate). Bars are colored according to the phylum (bold) and order. **B)** Principal coordinate analyses of dissimilarities between 16S rRNA compositions based on the (left) weighted UniFrac and (right) Bray-Curtis distance methods. **C)** Heatmap of the percent relative abundance of uroC-metabolizing *Enterocloster* spp. in A). **D,E)** PCR with *ucd*-specific primers on gDNA extracted from an *E. bolteae* isolate and fecal microbiota communities from 10 healthy donors (from one experimental replicate). 1% agarose gel of amplicons derived from **D)** uroC-metabolizing and **E)** non-metabolizing fecal slurries. Samples are matched to the urolithin metabolism data in Fig. 6C and 16S rRNA sequencing in A) and C). The no template control (NTC) and *E. bolteae* positive control are the same for both gels. **F,G)** 1% agarose gel of amplicons from Fig. 6D,E including the no reverse transcriptase (-RT) in **F)** uroC-metabolizing and **G)** non-metabolizing fecal slurries. The NTC is derived from the final lane of the gel in F). The uncropped and unannotated gel image is provided in the Supplementary Information.

 Supplementary Information

593 cDNA sequence reverse transcribed and amplified from uroC-induced *E. bolteae* RNA with *ucd*-
594 specific primers, 3605 bp (band can be seen on 1% agarose gel in Fig. 4B)

595 >Eb_ucd_RT-PCR_band coverage: 4.94e+03x

```
596 AAGCACAGCCAGGGTTATGGCAGGCGCAACGGATCTGATTCCGCCAATGAAAGACAAGGTTATATCACCAGAGTATATCATTG
597 ACCTTAAGAAAATTCCAGGTTTGGATTATCTGGAATACGATGACAGGGAAGGTTTAAAAATAGGGGCGCTGACAACACTGCGT
598 ACCATAGAGACATCTCCTCTGGTTAAAGAAAAGAATCCGGCAGTGGCTCATGCCGCCAAGGTAGTGGCATCCACACAAATCAG
599 GACAAAGGGCACCATGGCAGGCAATATCTGCAATGCGTCCCCATCCTGCGATACAGCCCCCAACCTACTGGCTCAGGGCGCTA
600 AGATATTGGTACAGGGTCCCAACAAGGACCGGGTCATTCAAATCGAAGACTTTTTCCTGGGTGTCAAAAAGACTTCCCTGGAG
601 CCGGGAGAAATCGTGACAGGCATTGTGATTCCGCCGCTGGCTGAGAATGAGAGGGCTGCATACATAAAACACGCGGTCAGAAA
602 GGCGATGGACCTGGCCATCATCGGCGTCGCAGTCAAGATTAAGGTGGAAGACGGCGTCTGCACAGACGCCAGGATTGCCCTGG
603 GCGCGGTTGCGGCCACGCCGGTACGGGCACCCGGGGCGGAGGAAGCCCTGATTGGCAAAGAACTTACCGATGAAGTGATTGTG
604 AAGGCATCGGAGGAAGCAATGAATTCCTGTCATCCAATCTCTGATATCCGCGCATCGGCCGAATACAGGAAGGATATGATCCG
605 CGTGTTTACAAAACGTGCCATTAAGCAGGCCATGGAATGCTATAACTAGGAGGGAGACAGGAATGAAACATAGTATAAGTTTA
606 ATCATCAACGGAGATCCGGTTGATGCTATTGTGAAGGATAACCTGACGCTTCTGGATTTTCTTCGTGACCAGTTGTTTCTGAC
607 AGGCACGAAAAAGGGGTGTGAGGAAGGGGAGTGTGGCGCCTGTACCGTGATGCTTGACGGTAAGCCGGTTAACTCCTGCTGTA
608 CCCTGGCCGTGGAGTGCGACGGCCATGAGATCATAACCGTGGAGGGAATTGCCAGGGAGGGGATGCTTCATCCCATTCAGAAG
609 CAGTTTATAGAGAAGTGGGCCATGCAGTGCGGCTACTGTACTCCGGGTATGATTATGTCCGCCAAAGCTCTTCTGGATGTGAA
610 TAAACATCCTACGGAACTGGAAATCCGGGAAGCCATTGAGGGCAATCTGTGCCGCTGCACCGGTTATGCAAAGATTGTGGAGG
611 CAATCCAGGCAGCCGCTGCGCAGATGAACTGGGAGGAGGAAGCAAAGAATGCATAAAGACTGTGACAAACATTATTTTAAAAA
612 ACCGGAATTTTTACCGTCTGACCGGCGAGAACAATTATGTCAGGATTGACGCTGAAGACAAGGTAACAGGACACGGCCAGTATG
613 TGGGTGACATCATGTTTCCGGATATGCTTACCGGAAAAATGGTCAGAAGCCCTTATGCATCTGCAAAAATCCTGTCCATTGAC
614 ACCAGCGAGGCAGAGAAGCTTCCAGGTGTAAAGTGCATCCTCACTGCCAGGGACTTTGAGTGGAAGTCCCTGGTGGGAAATGG
615 AGAATTTGCAGCTGAGTTCGCGGACAAGGAAGTATTGTGTTCTGAGAAGGTGCGCCAGGTTGGTGACGATGTGGCAGCCGTGG
616 CAGCCGTGGATGAGGAAACCGCCCAGCGGGCCGCGGATCTGATTAAGGTAGAGTACCAGGTGCTTCCGGGGGGTGTTTGACCCC
617 TTTGAGGCAATGGAGGAGAATGCCCCCGAGGTAAACTGGGAGGGCAAAGGCATACACAATATCGGCATGCAGTCCGTGATGAA
618 GGCGGGCACGGATATTGATGAGGAATTTGACCGCGCATCCTATGTGCAGCACAGGGATTACAAGACCCACCGTATGGTACATG
619 CGGCCATGGAGCCCCACGGTGCGGTGGCTACCTACAGGAATGGGACCTACACCATCTGGATGTCCACCCAGATGTCCTTTGTG
620 GACCAGTTCTGGTATGCCCGCTGCCTTGGCGTTGGAGAGAACCAGGTGCGGGTAATCAAGCCTCTTGTGGGCGGCGGTTTCGG
621 CGGCAAGCTGGATTCCTATTCCTTTGGCCTTTGCGCTGCCAAGATGGCGGAGATGACCGGACGCCCGGTACGCATGATTCTGT
622 CCCGCGAAGAAGTATTCCAGACCACGCGCAACCGCCACCCCATTTACATGCATATTGACACTGCCTTTGGGACAGACGGTAAG
623 CTGCTGGCAAAGAAATGCTACCATGTGCTGGACGGCGGGCCTTACGGCGGGTCCGGTGTTGCGGCCTGCGCCCAGTCCACATT
624 GTGGGCCAACTTCCCTTACAAGATGAATTCCGTGGATTTCCTGGCAAGGCGTGTGTATACCAATAACCCTTCTGCAGGAGCCA
625 TGAGAGGATATACGGCATGCCAGGTGCATTTTGCCCATGACCTGAACATGCAGTTCGCGGCTGACCAGATGGGCATTGACCCG
626 GTTGAGTTCCGCAAGATAAGCGCGGCAGACCCCGGATACGTGGCTCCGGCCGGACTGGCAATCACCAGCTGCGCATACAAGGA
627 GACACTGGATACGGCTGCAAAGGAAATTGGCTGGTATGAGAAGAAGGATAAGCTGAAAAAGGGAGAGGGCATTGGTTTTGCGG
628 GCACCGGCTTTGTATCGGGCACCGGATTCGCGGTGCTGGAAGCGCCCAACCAGAGCTCTGCGTGCGTGACCCTGCGCATGAAC
629 AAGCGGGGCATGGCCACCCTGTACATCGGCTCCCACGACATAGGACAGGGTTCCGATACGGTTATGACGGCCATTGTGGCCGA
630 GGAACTGGGGCTTCCCATGGATATGGTCAAGACCTTTATGTCCGACACCTTCCTGACGCCCTGGGACTCCGGCTCTTACGGCA
631 GCCGTGTTACCTTCCTGGCCGGCAATGCGGCCCGCCGCGCTGCCGTGGATGCAAAACGCCAGCTGTTTGAGGTCATTGCGCCT
632 ATGTGGGGGGTGATGCCGGAGACATTAGAGTGTCTGGACGGAAAGGTAATCAGCAAGGAGAAGGCAGAGTATCAGATGTCCAT
633 TGGAGACGCCATGTTCAAGTATATGACGGTCAAGGGCGGCGATGAACTGATTGGCGTGGGATCCTATTACCACCGTACCGACA
634 ATTCACAGTATAACGGCAACAATACCACCAACTACGCGCCTGCATACAGCTTCTCCACCGGAGCAGCCCATCTGACTGTGGAC
635 GAGGAGACCGGCGTCCTGGATATTGATGAATTTGTATTTGCCCATGACTGCGGCCGCGCACTGAATAAGAGGGCCGTGGAAGG
636 CCAGCTGGAGGGATCCATCGGCATGGGCCTGGGCTATGCTGTCTATGAGCACAATGTAACCAGGGAAGGAAAGATTCTCAATC
637 CCAACTTCCGCGATTACCGGCTGCCCACTGCCCTTGATATGCCGAAGATGCGCACCTTTTACGACTTTACACCGGACGAGGAA
638 GGACCTCTGGGCGCAAAGGAAGCTGGCGAGGGCTCCGCAGCACCTGTGGCGCCTGCCATTGCCAATGCGGTCAACATGGCAAC
639 CGGCGTGTACTTCACAGAGCTGCCTCTGGACCCGGA
```

Full uncropped gel image from Fig. 4B:

The ladder used here is the Quick-Load® Purple 1 kb Plus DNA Ladder (NEB, N0550).

Full uncropped gel image from Fig. 6D and Extended Data Fig. 8F:

The ladder used here is the Quick-Load® Purple 1 kb Plus DNA Ladder (NEB, N0550).

Full uncropped gel image from Fig. 6E and Extended Data Fig. 8G:

The ladder used here is the Quick-Load® Purple 1 kb Plus DNA Ladder (NEB, N0550).

Full uncropped gel image from Extended Data Fig. 8D:

The ladder used here is the Quick-Load® Purple 1 kb Plus DNA Ladder (NEB, N0550).

Full uncropped gel image from Extended Data Fig. 8E:

The ladder used here is the Quick-Load® Purple 1 kb Plus DNA Ladder (NEB, N0550).

Online Methods

Resources table

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background	*Enterocloster aldenensis* DSM 19262	DSMZ	DSM 19262	Type strain
Strain, strain background	*Enterocloster asparagiformis* DSM 15981	DSMZ	DSM 15981	Type strain
Strain, strain background	*Enterocloster bolteae* DSM 15670	DSMZ	DSM 15670	Type strain
Strain, strain background	*Enterocloster citroniae* DSM 19261	DSMZ	DSM 19261	Type strain
Strain, strain background	*Enterocloster clostridioformis* DSM 933	DSMZ	DSM 933	Type strain
Strain, strain background	*Enterocloster lavalensis* DSM 19851	DSMZ	DSM 19851	Type strain
Strain, strain background	*Lachnoclostridium pacaense* CCUG 71489T	CCUG	CCUG 71489T	Type strain
Commercial assay or kit	One-4-All Genomic DNA Minipreps Kit	BioBasic	BS88503	
Commercial assay or kit	Direct-zol RNA Miniprep Kit	Zymo Research	R2051	
Commercial assay or kit	ZR BashingBead Lysis Tubes (0.1 & 0.5 mm)	Zymo Research	S6012-50	
Commercial assay or kit	Ambion *DNA-free* DNA Removal Kit	Invitrogen	AM1906	
Commercial assay or kit	*OneStep* PCR Inhibitor Removal Kit	Zymo Research	D6030	
Commercial assay or kit	iScript Reverse Transcription Supermix	Bio-Rad	1708840	RT-PCR with random hexamers
Commercial assay or kit	LunaScript RT Master Mix Kit (Primer-free)	NEB	E3025	For gene-specific RT-PCR
Commercial assay or kit	Luna Universal qPCR Master Mix	NEB	M3003	
Commercial assay or kit	Monarch PCR & DNA Cleanup Kit	NEB	T1030	
Commercial assay or kit	Q5 High-Fidelity Polymerase	NEB	M0491	For 16S PCR and cloning
Commercial assay or kit	One*Taq* Quick-Load 2X Master Mix with Standard Buffer	NEB	M0486	For colony PCR
Commercial assay or kit	One*Taq* 2X Master Mix with Standard Buffer	NEB	M0482	For *ucd*-specific PCR
Commercial assay or kit	Platinum SuperFi II Green PCR Master Mix	Invitrogen	12369010	For ONT 16S PCR on fecal slurries
Commercial assay or kit	dNTP Mixture (10 mM)	BioBasic	DD0056	
Commercial assay or kit	Qubit dsDNA High Sensitivity (HS) Assay Kit	Invitrogen	Q32851	

Commercial assay or kit	Qubit RNA Broad Range (BR) Assay Kit	Invitrogen	Q10210	
Commercial assay or kit	Amplicon barcoding kit AB01-24	ONT	SQK-AMB111.24	
Commercial assay or kit	Thermolabile proteinase K	NEB	P8111	
Commercial assay or kit	Agencourt AMPure XP	Beckman Coulter	A63880	
Commercial assay or kit	Flongle Flow Cell (R9.4.1)	ONT	FLO-FLG001	
Plasmids and primers	Custom oligonucleotides (Desalted, Dried Form)	Invitrogen	10336022	See Primer sequences table
Commercial assay or kit	ExcelBand 100 bp+3K DNA Ladder	SMOBIO	DM2300	0.1-3 kb
Commercial assay or kit	Quick-Load Purple 1 kb Plus DNA Ladder	NEB	N0550	0.1-10 kb
Chemical compound, drug	Acetonitrile	Sigma	34998	CAS 75-05-8
Chemical compound, drug	Ethyl acetate	Sigma	34858	CAS 141-78-6
Chemical compound, drug	Methanol	Sigma	34860	CAS 67-56-1
Chemical compound, drug	Ethanol	Commercial Alcohols	P016EAAN	CAS 64-17-5
Chemical compound, drug	Formic acid	Sigma	F0507	CAS 64-18-6
Chemical compound, drug	Peptone	BioShop	PEP403.1	
Chemical compound, drug	Yeast extract	BioShop	YEX401.500	CAS 8013-01-2
Chemical compound, drug	Sodium chloride	BioShop	SOD004.5	CAS 7647-14-5
Chemical compound, drug	Starch (from potato)	Sigma	S2004-500G	CAS 9005-25-8
Chemical compound, drug	D-glucose monohydrate	BioShop	GLU601.1	CAS 14431-43-7
Chemical compound, drug	Sodium pyruvate	BioShop	PYR302.100	CAS 113-24-6
Chemical compound, drug	Sodium succinate dibasic	Sigma	14160-100G	CAS 150-90-3
Chemical compound, drug	Sodium thioglycolate	TCI	M0053	CAS 367-51-1
Chemical compound, drug	Agar	Fisher	BP1423-500	CAS 9002-18-0
Chemical compound, drug	L-arginine HCl	BioShop	ARG006.100	CAS 1119-34-2
Chemical compound, drug	L-cysteine	BioShop	CYS555.100	CAS 52-90-4
Chemical compound, drug	Sodium bicarbonate	BioShop	SOB308.500	CAS 144-55-8
Chemical compound, drug	Haemin	Sigma	H9039-1G	CAS 16009-13-5
Chemical compound, drug	Vitamin K1	Sigma	95271-250MG	CAS 84-80-0
Chemical compound, drug	Dithiothreitol (DTT)	Fisher	BP172-25	CAS 3483-12-3

Chemical compound, drug	Tris base	Sigma	10708976001	CAS 77-86-1
Chemical compound, drug	Magnesium sulfate heptahydrate	BioShop	MAG511.1	CAS 10034-99-8
Chemical compound, drug	Calcium chloride dihydrate	Fisher	BP510-500	CAS 10035-04-8
Chemical compound, drug	SIGMAFAST Protease Inhibitor Cocktail, EDTA-free	Sigma	S8830-2TAB	
Chemical compound, drug	Urolithin C	TRC	U847015	CAS 165393-06-6
Chemical compound, drug	Urolithin A	TRC	U847000	CAS 1143-70-0
Chemical compound, drug	Urolithin A	Sigma	SML1791-5MG	CAS 1143-70-0
Chemical compound, drug	Urolithin M-6	TRC	U847040	CAS 1006683-97-1
Chemical compound, drug	Isourolithin A	TRC	I917520	CAS 174023-48-4
Chemical compound, drug	Urolithin B	TRC	U847005	CAS 1139-83-9
Chemical compound, drug	Salicylic acid	Sigma	84210-100G	CAS 69-72-7
Chemical compound, drug	TRI reagent	Zymo Research	R2050-1-50	
Chemical compound, drug	Agarose	Fisher	BP160-500	CAS 9012-36-6
Chemical compound, drug	SafeView Classic	Abm	G108	
Chemical compound, drug	PBS pH 7.4 (10X)	Gibco	70011-044	
Chemical compound, drug	Nuclease-free water	Invitrogen	10977-015	
Lab equipment, instruments	AnaeroPack System	Mitsubishi Gas Chemical Company	10-01	
Lab equipment, instruments	Vinyl anaerobic chambers	Coy Laboratory Products	Type A	
Lab equipment, instruments	Multiscan FC Microplate Photometer	Thermo Fisher	51119100	
Lab equipment, instruments	Costar 96-Well Clear Round Bottom	Corning	3788	
Lab equipment, instruments	LifeECO Thermal Cycler	BIOER	TC-96	
Lab equipment, instruments	Nanodrop 2000c	Thermo Fisher	ND-2000C	
Lab equipment, instruments	Mini Beadbeater 16	Biospec	607	
Lab equipment, instruments	Viia7 qPCR machine	Applied Biosystems	4453535	
Lab equipment, instruments	MicroAmp Fast 96-Well Reaction Plates (0.1 mL)	Applied Biosystems	4346907	
Lab equipment, instruments	PlateSeal Film, Clear Polyester, X-Large, Non-Sterile	PlateSeal	PS-PETSTXL-100	
Lab equipment, instruments	LC-MS (Agilent 1260 Infinity II LC connected to a 6120 Quadrupole MS)	Agilent		

Lab equipment, instruments	Poroshell 120 EC-C18 column (4.6x50 mm, 2.7 μm)	Agilent	699975-902	
Lab equipment, instruments	Poroshell 120 EC-C18 column (4.6x5 mm, 2.7 μm) guard column	Agilent	820750-911	
Lab equipment, instruments	Vacuum concentrator	Heto Lab	VR-1	
Lab equipment, instruments	Sonicator	Misonix	S3000	
Lab equipment, instruments	Rotary evaporator	Buchi	R-300	
Software	Agilent OpenLab CDS	Agilent	N/A	
Software	GraphPad Prism 10	GraphPad by Dotmatics	N/A	
Software	Mascot 2.6.2	Matrix Science	N/A	
Software	Scaffold 5	Proteome Software Inc	N/A	
Software	R Studio 2023.06.0+421	Posit Software	N/A	
Software	Affinity Designer	Serif Ltd	N/A	

658 NEB = New England Biolabs, ONT = Oxford Nanopore Technologies, TRC = Toronto Research
659 Chemicals

660 **Primer sequences table**

Designation	Purpose	Source	5'-Sequence-3'
16S_V1_27_f	16S PCR		AGAGTTTGATCMTGGCTAG
16S_V9_1492_r	16S PCR		TACGGYTACCTTGTTAYGACTT
ONT_16S_27_f_GGK	16S PCR		**GGK**AGRGTTYGATYMTGGCTCAG
ONT_16S_1492_r_GGK	16S PCR		**GGK**CCGGYTACCTTGTTACGACTT
Eb_ucdO_qPCR_f	qPCR	This work	GTGGATGAGGAAACCGCCCAGC
Eb_ucdO_qPCR_r	qPCR	This work	TGCCCTCCCAGTTTACCTCGGG
Eb_ucdF_qPCR_f	qPCR	This work	GCGCCTGTACCGTGATGCTTGA
Eb_ucdF_qPCR_r	qPCR	This work	CCCTCCCTGGCAATTCCCTCCA
Eb_ucdC_qPCR_f	qPCR	This work	AGCCCCCAACCTACTGGCTCAG
Eb_ucdC_qPCR_r	qPCR	This work	GCCTGTCACGATTTCTCCCGGC
Eb_dnaK_Ref_qPCR_f	qPCR	This work	GGTGCTGTTGGTAGGCGGTTCC
Eb_dnaK_Ref_qPCR_r	qPCR	This work	GGGTCTTGCTGGGTTCCTTGCC
ucdCFO_RT-PCR_f	RT-PCR	This work	GCAATCTGTTCCTYGAGCTTGG
ucdCFO_RT-PCR_r	RT-PCR	This work	CGTGSAGCGCTCTCCAGATATG

661 **Adapter sites** are bolded.

Bacterial strains and culturing conditions

Bacterial strains used in this study are listed in the *Resources table*. All bacterial stains were validated by sequencing the 16S rRNA gene (see Genomic DNA extraction and 16S rRNA sequencing of bacterial isolates). The same culture used for validation was used to make 25% glycerol stocks. Strains were grown from glycerol stocks on mABB+H agar plates (recipe below) for 48-72 h at 37 °C in a vinyl anaerobic chamber, which was maintained with a gas mixture of 3% H_2, 10% CO_2, 87% N_2. To make overnight cultures, a single colony was inoculated into 5 mL of liquid mABB+H medium and incubated at 37 °C between 16-24 h.

Human fecal sample collection

Human fecal samples were collected under the approval of protocol A04-M27-15B by the McGill Faculty of Medicine Institutional Review Board. Informed written consent was received from the participants for the use of human samples. Eligibility criteria for the healthy participants were as follows: body mass index between 18.5–30, no diagnosed gastrointestinal disease, no ongoing therapeutic treatment, and no antibiotic usage 3 months prior to the start of the study. Subject information was recorded at the time of sampling. The age of donors ranged from 21–40 years. Fresh fecal samples were collected and placed immediately in an anaerobic chamber, aliquoted, and stored at −70 °C until use.

Modified anaerobe basal broth (mABB and mABB+H)

For 1 L of modified anaerobe basal broth (mABB), the following components were dissolved in MilliQ water, then autoclaved: 16 g peptone, 7 g yeast extract, 5 g sodium chloride, 1 g starch, 1 g D-glucose monohydrate, 1 g sodium pyruvate, 0.5 g sodium succinate, 1 g sodium thioglycolate, 15 g agar (for plates). The autoclaved solution was allowed to cool, then the following filter-sterilized solutions were added aseptically: 10 mL of 100 mg/mL L-arginine-HCl, 10 mL of 50 mg/mL L-cysteine, 8 mL of 50 mg/mL sodium bicarbonate, 50 µL of 10 mg/mL

vitamin K1, 20 mL of 50 mg/mL dithiothreitol, and, for mABB+H, 10 mL of 0.5 mg/mL haemin.

The media was then placed in the anaerobic chamber and allowed to reduce for at least 24 h

prior to its use in experiments.

Genomic DNA extraction of isolates and identity validation

The identities of all bacteria in this study were validated by full-length (V1-V9) 16S rRNA

sequencing. DNA was first extracted from 0.5-1 mL of overnight culture using the One-4-All

Genomic DNA Miniprep Kit (BioBasic) according to the manufacturer's instructions. The purified

genomic DNA (2 µL, ~ 20 ng) was used as a template for PCR reactions (25 µL reaction volume)

using the Q5 High-Fidelity polymerase (NEB). PCR tubes were placed in a thermal cycler and

targets (~1.5 kb) were amplified according to the following cycling conditions: 30 s at 98 °C, 30

cycles (10 s at 98 °C, 20 s at 60 °C, 45 s at 72 °C), 2 min at 72 °C, and hold at 10 °C. 5 µL of

the reaction was mixed with 6X loading buffer and loaded onto a 1% agarose gel (made with

1X TAE buffer) containing SafeView Classic (Abm). PCR product sizes were compared to the

ExcelBand 100 bp+3K DNA Ladder (SMOBIO).

PCR products (~1.5 kb) were purified using the Monarch PCR & DNA Cleanup Kit (NEB)

according to the manufacturer's instructions for products < 2 kb. Purified 16S PCR products

were eluted in nuclease-free water, quantified using the Qubit dsDNA HS assay kit (Invitrogen),

and adjusted to 30 ng/µL. Samples were submitted to Plasmidsaurus for long-read sequencing

using Oxford Nanopore Technologies (v14 library preparation chemistry, R10.4.1 flow cells).

Urolithin extraction from fecal slurries or bacterial cultures

Frozen (-70 °C) fecal slurries or bacterial cultures were thawed at room temperature. For

quantification of urolithin concentrations, urolithin standards (stock 10 mM in DMSO) were

spiked into media immediately before extraction.

Extraction Method A: Salicylic acid (3 mg/mL in DMSO) was spiked-in as an internal standard

at a final concentration of 50 µg/mL. The cultures and standards were then extracted with 3

volumes of ethyl acetate + 1% formic acid (e.g., 600 µL solvent to 200 µL thawed culture). The

organic phase (top) was transferred to a new tube and dried in a vacuum concentrator (Heto

Lab) connected to a rotary evaporator (Buchi). After solvent removal, samples were redissolved

in 0.5 volumes (relative to the starting culture) of 50% MeOH:H_2O. Samples were centrifuged

at 20,000 x g for 5 min to pellet insoluble material, then transferred to LC-MS vials. Urolithins

were then analyzed by LC-MS.

Extraction Method B: The cultures, suspensions, or supernatants were diluted with an equal

volume of MeOH, vortexed briefly, and incubated at room temperature for 10 min. Samples

were centrifuged at 20,000 x g for 5 min to pellet insoluble material, then transferred to LC-MS

vials. Urolithins were then analyzed by LC-MS.

LC-MS method to quantify urolithins

Samples (10 µL) were injected into a 1260 Infinity II Single Quadrupole LC/MS system

(Agilent) fitted with a Poroshell 120 EC-C18 4.6x50 mm, 2.7 µm column (Agilent). The mobile

phase was composed of MilliQ water + 0.1% formic acid (solvent A) and acetonitrile + 0.1%

formic acid (solvent B). The flow rate was set to 0.7 mL/min. The gradient was as follows: 0-8

min: 10-30 %B, 8-10 min: 30-100 %B, 10-13.5 min: 100 %B isocratic, 13.5-13.6 min: 100-10

%B, then 13.6-15.5 min: 10 %B. The multiple wavelength detector was set to monitor

absorbance at 305 nm. The mass spectrometer was run in negative mode in both selected ion

monitoring (SIM) and scan (100-1000 m/z) modes to validate peak identities. Peaks were

validated based on retention times compared to spike-in standards and mass-to-charge ratios.

To quantify urolithins, peak areas for the compounds of interest were compared to spike-in

standards of known concentration(s). When standards were not available (urolithin M7), the

extracted ion chromatogram was used ($[M-H]^- = 243$).

Isolate treatments with urolithins

Treatment during growth (spike-in): Overnight cultures of *Enterocloster* spp. were diluted

1/50 into fresh mABB+H and incubated at 37 °C. After 5 hours of incubation (~OD_{620} from a

200 µL sample ~ 0.4), 10 mM urolithins (or an equivalent volume of DMSO) were added to the

growing cultures at a final concentration of 50 or 100 µM for protein expression or RNA

expression, respectively. For protein expression analyses and inducibility tests, the cultures

were incubated for an additional 4 h. For RNA expression analyses, the cultures were incubated

for an additional 2 h.

Treatment prior to growth: Overnight cultures of *Enterocloster* spp. were diluted 1/50 into fresh

mABB+H. Separately, treatments (10 mM stocks solutions, dissolved in DMSO) were prepared

in fresh mABB+H medium to a final concentration of 111 µM. In a 96-well plate, 180 µL of

treated media were combined with 20 µL of 1/5 culture. These were plated in technical

triplicates. The final concentration of treatment was 100 µM (unless otherwise specified in

concentration-response experiments) and the final dilution of bacteria was 1/100.

In both cases, urolithins were extracted from cultures using *Extraction Method A*.

Phylogenetic tree construction

Phylogenetic trees based on the 16S rRNA gene were constructed using the DSMZ

single-gene phylogeny server (https://ggdc.dsmz.de/phylogeny-service.php#)[57] with GenBank

16S rRNA sequence accessions: *E. aldenensis* (DQ279736), *E. asparagiformis* (AJ582080), *E.*

bolteae (AJ508452), *E. citroniae* (HM245936), *E. clostridioformis* (M59089), *E. lavalensis*

(EF564277), *L. pacaense* (LT799004), *G. pamelaeae* (AM886059), *G. urolithinfaciens*

(HG000667), *E. isourolithinifaciens* (MF322780).

Phylogenetic trees based on whole genomes and proteomes were constructed using the

Type (Strain) Genome Server (TYGS, https://tygs.dsmz.de)[55,58] with the following GenBank

genome accessions: *E. aldenensis* (GCA_003467385.1), *E. asparagiformis*

(GCA_025149125), *E. bolteae* (GCA_000154365), *E. clostridioformis* (GCA_900113155), *E.*

lavalensis (GCA_003024655), *L. pacaense* (GCA_900566185). For *E. citroniae*, the Integrated

Microbial Genomes ObjectID was used: *E. citroniae* (2928404274). Further information on

nomenclature and taxonomy was obtained from the List of Prokaryotic names with Standing in

Nomenclature (LPSN, available at https://lpsn.dsmz.de).

Growth curves

Bacterial cultures were prepared as described above (see *Treatment during growth*

(spike-in) in the *Isolate treatments with urolithins section*). At each timepoint, 200 μL of culture

were transferred to a Costar 96-Well Clear Round Bottom (Corning) and the OD_{620} was

measured using a Multiscan FC Microplate Photometer (Thermo Fisher).

Resting cell suspension assay to test inducibility

Bacteria (10 mL growing cultures in mABB+H media) were grown with 50 μM uroC (or

an equivalent volume of DMSO) as detailed in the *Treatment during growth (spike-in)* section

above and incubated for 4 h at 37 °C. Cultures were then pelleted at 6,500 x g for 3 min and

the supernatants were discarded. The cells were washed with 10 mL of pre-reduced PBS

(placed in the anaerobic chamber 24 h before), re-pelleted, and resuspended in 2 mL of pre-

reduced PBS, yielding resting cell suspensions. For each condition tested, a 200 μL aliquot of

the resting cell suspension was transferred into a sterile 1.5 mL tube, and 10 mM urolithins

(uroM6, uroC, isouroA, or DMSO) were added at a final concentration of 100 µM. Resting cell

suspensions were briefly vortexed and incubated at room temperature in the anaerobic

chamber for 16h prior to freezing and urolithin extraction using *Extraction Method B*.

RNA extraction from isolates

A volume of 1.5 mL of treated (100 µM urolithin C for 2 h) *Enterocloster* spp. culture (see

Enterocloster spp. urolithin C treatments) was pelleted (6,500 g for 3 min) and the supernatant

was removed for later LC-MS analysis. The pellet (suspended in 200 µL of media) was then

mixed with 800 µL TRI reagent (Zymo Research) and transferred to a ZR BashingBead lysis

tube (Zymo Research). Samples were lysed in a Mini Beadbeater 16 (Biospec) according to

the following sequence: 1 min ON, 5 min OFF. For RNA-sequencing, bead beating was done

for a total of 5 min ON. For RT-(q)PCR, bead beating was done for a total of 2 min ON to

preserve longer transcripts. RNA isolation was then performed using the Direct-zol RNA

Miniprep Kit (Zymo Research) according to the manufacturer's instructions (including an on-

column DNase digestion). To ensure complete DNA removal, an additional DNA digestion step

was performed on the isolated RNA using the Ambion *DNA-free* DNA Removal Kit (Invitrogen)

according to the manufacturer's instructions. The DNA-free RNA was then cleaned up using

the *OneStep* PCR Inhibitor Removal Kit (Zymo Research). RNA concentration and quality were

initially verified by NanoDrop and 1 % agarose gel electrophoresis. For RNA-sequencing, RNA

integrity was assessed by Génome Québec using a Bioanalyzer 2100 (Agilent). RNA integrity

(RIN) values ranged between 7.5-7.8 for *E. asparagiformis* DSM 15981 and 7.0-7.3 for *E.*

bolteae DSM 15670.

RNA-sequencing of urolithin C-treated *E. bolteae* and *E. asparagiformis* isolates

Total RNA was sent to Génome Québec for library preparation and RNA-sequencing.

Briefly, total RNA was prepared for Illumina sequencing using the NEBNext rRNA Depletion Kit

(Bacteria) (NEB) kit to remove rRNA and using the NEBNext Multiplex Oligos for Illumina (NEB)

kit (stranded/directional). Prepared libraries were quality checked with a Bioanalyzer 2100

(Agilent) prior to sequencing. Sequencing was performed on a NovaSeq 6000 (Illumina) with

the following flow cell/settings: S4 flow cell, 100 bp, 25 M reads, paired end.

Analysis of RNA-seq reads was done using Galaxy bioinformatics cloud computing

(https://usegalaxy.org/) hosted by Compute Canada Genetics and Genomics Analysis Platform

(GenAP) (https://www.genap.ca/). Genomes and annotations were fetched from the NCBI

genome browser: *E. bolteae* (ASM223457v2) (accessed 2022/05/11) and *E. asparagiformis*

(ASM2514912v1) (accessed 2023/09/19). Raw reads were first verified for quality using

FastQC (v0.73, https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) with default

parameters. FastQC reports were aggregated into MultiQC (v1.11) [59]. The mean sequence

quality scored were above 35 for all samples. Raw reads were then trimmed using Cutadapt

(v3.7) [60] to trim adapter sequences (R1 sequence:

AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC, and R2 sequence:

AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT) that were not removed after sequencing

using default parameters for paired end reads. Trimmed reads were then aligned to reference

genomes for each bacterium using HISAT2 (v2.2.1) [61] with paired-end parameters and reverse

strandedness (RF). Aligned read counts were assigned to features in annotation files (.gtf)

using featureCounts (v2.0.1) [62] with the following parameters: reverse strandedness, count

fragments instead of reads, GFF feature type filter = "gene", multi-mapping and multi-

overlapping features included (-M -O), minimum mapping quality per read of 0, and the rest of

the parameters were kept as default. Differential gene expression analysis was then performed

using DESeq2 (v2.11.40.7) [63] using default parameters. Differential expression tables were

annotated with the Annotate DESeq2/DEXSeq output tables tool (v1.1.0) in Galaxy to include

the following: GFF feature type = "CDS", GFF feature identifier = "gene_id", GFF transcript

identifier = "transcript_id", GFF attribute to include = "protein_id, product". The "protein_id" was

used to query the NCBI database and the NCBI Sequence Viewer was used to investigate the

genomic context surrounding genes of interest.

Comparative genomics

The nucleotide sequence for the *Enterocloster bolteae* DSM 15670 *ucd* operon (NCBI

NZ_CP022464 REGION: complement(4417875..4421605)) was used as a query for BLASTn

(megablast) searches using the refseq_genomes database limited to Bacteria (taxid:2). The

NCBI multiple sequence alignment (MSA) viewer was used to download alignment figures.

RT-PCR analysis of *E. bolteae* to determine *ucd* operon structure

Isolated RNA samples (500 ng) were reverse transcribed using the LunaScript® RT

Master Mix Kit (Primer-free) (NEB) in a reaction volume of 10 µL containing the ucdCFO_RT-

PCR_r primer at a final concentration of 1 µM. The No-RT Control included in the kit was used

as a no-enzyme control for reverse transcription. The reaction mixtures were incubated in a

thermal cycler: 10 min at 55°C, 1 min at 95°C. PCR reactions were conducted using the OneTaq

2X Master Mix with Standard Buffer (NEB). The ucdCFO_RT-PCR primer pair was added to

the master mix (to a final concentration of 0.2 µM) and 1 µL of template (cDNA, -RT, no

template, or gDNA) was added for a total reaction volume of 20 µL. PCR tubes were placed in

a thermal cycler and targets were amplified according to the following conditions: 20s at 94°C,

31 cycles (20s at 94°C, 30s at 62°C, 3 min at 68°C), 5 min at 68°C. A volume of 5 µL of reaction

was directly loaded onto a 1% agarose gel (made with 1X TAE buffer) containing SafeView

Classic (Abm). PCR product sizes were compared to the Quick-Load® Purple 1 kb Plus DNA

Ladder (NEB). The rest of the PCR product was then run on a 1% agarose gel and bands

corresponding to the desired products were cut out and purified using the Monarch DNA Gel

Extraction kit (NEB). DNA was quantified using the Qubit dsDNA HS assay kit (Invitrogen) and

submitted to Plasmidsaurus for long-read sequencing using Oxford Nanopore Technologies

(Supplementary Information).

RT-qPCR analysis of *E. bolteae ucd* operon genes

Isolated RNA samples (500 ng) were reverse transcribed using the iScript Reverse

Transcription Supermix (Bio-Rad) in a reaction volume of 10 μL. The iScript No-RT Control

Supermix was used as a no enzyme control for reverse transcription (-RT). The reaction

mixtures were incubated in a thermal cycler: 5 min at 25 °C, 20 min at 48 °C, and 1 min at 95

°C. Both cDNA and -RT controls were diluted 1/20 in nuclease-free water before use. qPCR

reactions were conducted using the Luna Universal qPCR Master Mix kit (NEB). The

Eb_ucdO_qPCR, Eb_ucdF_qPCR, Eb_ucdC_qPCR, and Eb_dnaK_Ref_qPCR primer pairs

were added to their respective master mixes (final primer concentration of 250 nM) and 6.6 or

4.4 μL of diluted template (cDNA, -RT, no template) were added to 26.4 or 17.6 μL of master

mix for triplicates or duplicates, respectively. All cDNA samples were run in technical triplicates,

while other sample types were run in technical duplicates. Replicate mixes were pipetted (10

μL/well) into a MicroAmp Fast 96-Well Reaction Plate (Applied Biosystems) and the plates were

sealed, then spun down for 2 min to eliminate air bubbles. The qPCR detection parameters

were as follows: SYBR Green detection, ROX reference dye, 10 μL reaction volume. The

thermal cycling conditions were: 1 min at 95 °C, 40 cycles (15 s at 95 °C, 30 s at 60 °C), then

melt analysis (60-95 °C). Data were analyzed according to the $2^{-\Delta\Delta Ct}$ method [64] with the dnaK

gene serving as the reference gene (*E. bolteae* dnaK RNA-seq $\log_2$FC = 0.122).

Protein extraction

All steps other than sonication were carried out under anaerobic conditions. To extract

proteins, 10 mL of treated (50 μM urolithin C for 4 h) *Enterocloster* spp. culture (see

Enterocloster spp. urolithin C treatments) were pelleted (6,500 g for 3 min) and the supernatant

was discarded. The pellet was washed with 10 mL of pre-reduced PBS, pelleted again, and

resuspended in 0.5 mL of pre-reduced lysis buffer (20 mM Tris, pH 7.5, 500 mM NaCl, 10 mM

$MgSO_4$, 10 mM $CaCl_2$, and 1 tablet/100 mL SIGMAFAST protease inhibitor (EDTA-free)). The

resuspended pellet was then sonicated on ice using a Misonix Sonicator 3000 set to power

level 2/10 according to the following sequence (aerobically, in a cold room): 20 s ON, 40 s OFF,

for a total of 2 min ON. Tubes were centrifuged at 20,000 x g for 2 min to pellet insoluble

particles and 0.4 mL of lysate was transferred to a new 1.5 mL tube (kept on ice). Lysates used

in metabolism assays were transported to the anaerobic chamber in a sealed plastic bag

containing an anaerobic gas generating system to minimize loss in activity.

Urolithin metabolism assays using crude lysates from uroC-induced *E. bolteae*

Protein lysates (described above) were aliquoted (50 µL aliquots) into 1.5 mL tubes, then

treated with DMSO or urolithin C (10 mM stock) at a final concentration of 350 µM. Cofactors

(NADPH, NADP, NADH, and FAD, each dissolved to a final concentration of 30 mM (in lysis

buffer immediately before the assay was run) were added individually to the lysates at a final

concentration of 1 mM. The lysates were incubated at room temperature in an anaerobic

chamber for 20 h prior to freezing at -70 °C. Samples were then extracted using *Extraction*

Method B.

To assess the oxygen sensitivity of crude lysates from uroC-induced *E. bolteae*, samples were

prepared as described above. After adding DMSO or uroC and NADH (under anaerobic

conditions), tubes were either incubated at room temperature inside the anaerobic chamber or

just outside of the chamber for 20 h. Afterwards, samples were frozen at -70 °C and then

extracted using *Extraction Method B*.

Proteomics analysis of uroC-treated *E. bolteae*

Extracted proteins were submitted for proteomic analysis at the RI-MUHC. For each sample, protein lysates were loaded onto a single stacking gel band to remove lipids, detergents, and salts. The gel band was reduced with DTT, alkylated with iodoacetic acid, and digested with trypsin. Extracted peptides were re-solubilized in 0.1% aqueous formic acid and loaded onto a Thermo Acclaim Pepmap (Thermo, 75 um ID X 2 cm C18 3 um beads) precolumn and then onto an Acclaim Pepmap Easyspray (Thermo, 75 um ID X 15 cm with 2 um C18 beads) analytical column separation using a Dionex Ultimate 3000 uHPLC at 250 nL/min with a gradient of 2-35% organic (0.1% formic acid in acetonitrile) over 3 hours. Peptides were analyzed using a Thermo Orbitrap Fusion mass spectrometer operating at 120,000 resolution (FWHM in MS1) with HCD sequencing (15,000 resolution) at top speed for all peptides with a charge of 2+ or greater. The raw data were converted into *.mgf format (Mascot generic format) for searching using the Mascot 2.6.2 search engine (Matrix Science) against *Enterocloster bolteae* DSM 15670 proteins (NCBI assembly GCF_002234575.2) and a database of common contaminant proteins. Mascot was searched with a fragment ion mass tolerance of 0.100 Da and a parent ion tolerance of 5.0 ppm. O-63 of pyrrolysine, carboxymethyl of cysteine and j+66 of leucine/isoleucine indecision were specified in Mascot as fixed modifications. Deamidation of asparagine and glutamine and oxidation of methionine were specified in Mascot as variable modifications.

The database search results were loaded into Scaffold Q+ Scaffold_5.0.1 (Proteome Sciences) for statistical treatment and data visualization. Scaffold (v5.3.0) was used to validate MS/MS based peptide and protein identifications. Peptide identifications were accepted if they could be established at greater than 95.0% probability by the Peptide Prophet algorithm [65] with Scaffold delta-mass correction. Protein identifications were accepted if they could be established at

greater than 99.0% probability and contained at least 2 identified peptides. Protein probabilities

were assigned by the Protein Prophet algorithm [66]. Proteins that contained similar peptides and

could not be differentiated based on MS/MS analysis alone were grouped to satisfy the

principles of parsimony. Proteins sharing significant peptide evidence were grouped into

clusters. Protein quantification and differential expression were determined in Scaffold using

the following parameters: Quantitative method was set to total spectra, the minimum value was

set to 0.5 in case proteins were not detected in one condition, and statistical tests were

performed using Fisher's exact test with the Benjamini-Hochberg multiple test correction at a

significance level set to 0.05.

Protein structures

The protein FASTA sequences (NCBI RefSeq accessions for UcdC: WP_002569575.1,

UcdF: WP_002569574.1, UcdO: WP_002569573.1) for the *Enterocloster bolteae* DSM 15670

ucd operon were used as a query for BLASTp searches against the UniProtKB reference

proteomes + Swiss-Prot databases. The AlphaFold2 protein structures for matches (UniProt

accessions for UcdC: A8RZR5, UcdF: A8RZR3, UcdO: A8RZR2) of each protein were

downloaded and imported into PyMOL (v2.4.1). Foldseek (in 3Di/AA mode) was used to

generate a list of proteins with similar structures from solved crystal structures in the Protein

Data Bank (PDB) [44].

Hits of published X-ray crystal structures (PDB: 1ZXI (from *Afipia carboxidovorans* OM5) and

PDB: 3UNI (from *Bos taurus*)) were fetched from the PDB and imported into PyMOL. The

AlphaFold2 structures for each Ucd protein were aligned to the following chains in the published

PDBs using the "super" command: PDB 1ZXI: UcdC to chain C, UcdF to chain A, UcdO to chain

B; PDB 3UNI: UcdC, UcdF, and UcdO to chain A.

curatedMetagenomicData meta-analysis of *Enterocloster ucd* operon in human fecal metagenome datasets

All 93 metagenomic studies (22,588 samples and their metadata) available in the curatedMetagenomicData R package [47] (v3.8.0) were downloaded locally (ExperimentHub snapshotDate(): 2023-04-24, accessed on 2023-06-06) and transferred to the Narval cluster hosted by the Digital Research Alliance of Canada. Metagenomic data for urolithin C-metabolizing *Enterocloster* spp. were obtained by querying the "relative_abundance" (pre-processed using MetaPhlAn3 and "gene_families" (pre-processed using HUMAnN3) entries in individual study datasets [67]. For individual taxa (containing partial strings "bolteae", "citroniae", "asparagiformis", "asparagiforme", "Enterocloster", or "47FAA" (corresponding to *L. pacaense*)), relative abundance (%) was extracted from the rows of the "relative_abundance" datasets using the stringr R package (v1.5.0, https://github.com/tidyverse/stringr). Prevalence (relative abundance in sample > 0) was then calculated for each sample.

For specific genes, the NCBI protein accessions for each gene of the *ucd* operon (*ucdO*, *ucdF*, *ucdC*) was used to search the UniProt database. UniRef90 accession numbers corresponding to hits (C5EGQ4, G5HFF3, A8RZR5, respectively) were then extracted from the rows of the "gene_families" datasets using the stringr R package. Prevalence (abundance in sample > 0) was then calculated for each sample. R scripts and RData files are available in Zenodo (see Data Availability).

Fecal slurry preparation and treatment

Frozen (-70 °C) fecal samples were brought into the anaerobic chamber and allowed to thaw. The samples were suspended in 1 mL mABB medium per 0.1 g feces and homogenized by breaking apart large pieces with a sterile loop and by vortexing. Large particles were pelleted by centrifuging the tubes at 700 x g for 3 min. The supernatants (containing bacteria) were

transferred to new tubes and centrifuged at 6,500 x g for 5 min to pellet the cells. The

supernatants were discarded, and the cell pellets were washed with 5 mL of fresh media. The

cell suspensions were once again centrifuged at 6,500 x g for 5 min and the resulting cell pellets

were resuspended in 600 µL per 0.1 g feces. Resuspended cells were treated with either 100

µM urolithin C or an equivalent volume DMSO and incubated at 37°C anaerobically for 48 h.

200-300 µL volumes were removed from the batch cultures and immediately frozen at -70°C

for later extraction of urolithins (using *Extraction Method A*), DNA, and RNA.

Genomic DNA extraction from microbial communities

A 300 µL fecal slurry aliquot (between 50-100 mg wet weight) was pelleted (10,000 g for

5 min) and the supernatant was removed for later LC-MS analysis. The pellet was then mixed

with 750 µL of ZymoBIOMICS lysis solution (Zymo Research) and transferred to a ZR

BashingBead lysis tube (Zymo Research). Samples were lysed in a Mini Beadbeater 16

(Biospec) according to the following sequence: 1 min ON, 5 min OFF for a total of 5 min ON.

DNA was then purified using the ZymoBIOMICS DNA Miniprep Kit (Zymo Research) according

to the manufacturer's instructions (including the *OneStep* PCR Inhibitor Removal step). Purified

DNA samples were quantified using the Qubit dsDNA HS assay kit (Invitrogen).

Long read 16S rRNA sequencing of microbial communities

Long read 16S PCR reactions were conducted using the Platinum SuperFi II Green PCR

Master Mix (Invitrogen). The ONT_16S_27F_GGK and ONT_16S_1492R_GGK primer pairs

(see *Primer sequences table*) were added to their respective master mixes (final primer

concentration of 0.2 µM) and 1 µL of template (~10 ng) was added (for a total reaction volume

of 25 µL). PCR tubes were placed in a thermal cycler and targets were amplified according to

the following cycling conditions: 30 s at 98 °C, 30 cycles (10 s at 98 °C, 10 s at 60 °C, 30 s at

72 °C), 5 min at 72 °C, and hold at 4 °C. Amplicons were quantified using the Qubit dsDNA HS

assay kit (Invitrogen) to verify that amplicon concentrations were reasonably balanced (range

= 18.36-24.00 ng/μL). Barcoding of amplicons was performed with 2 μL of PCR reaction

according to the manufacturer's instructions (for ONT kit SQK-AMB111-24). Barcoding

reactions were incubated in a thermal cycler for 10 min at 65 °C, then for 2 min at 80 °C. 10 μL

of each barcoding reaction were pooled and proteins were digested using heat-labile proteinase

K (NEB) by incubating the pooled library for 15 min at 37 °C, followed by heat inactivation for

10 min at 55 °C. Amplicons were purified using Agencourt AMPure XP beads (Beckman

Coulter) using 0.7 volumes of beads-to-library. Following 70% EtOH washes and drying steps,

the library was eluted using 15 μL of the provided elution buffer (EB), yielding a library with a

concentration of 30 ng/μL using the Qubit dsDNA HS assay kit (Invitrogen). 11 μL of the eluted

DNA library were transferred to a new tube and combined with 1 μL of Rapid Adapter T (RAP

T). This mixture was incubated at room temperature for 10 min. Since the library was

concentrated, it was diluted 1:2 in EB before combining with SB II and LB II, then loaded into a

primed Flongle Flow Cell (R9.4.1) in a MinION device following the manufacturer's instructions.

Sequencing was allowed to proceed for ~20 h until pore exhaustion or enough reads were

obtained. Base calling & demultiplexing was performed using Guppy (v6.4.6) using the "SUP"

super high accuracy model for R9.4.1 flow cells. The raw reads were filtered for a length

between 1500 ± 200 bp. Filtered reads were assigned to taxa using Emu [68] (v3.4.4, GitLab

Project ID: 19618062) by mapping 16S rRNA sequences to the emu_database database

(based on the NCBI 16S RefSeq with the entry for *E. asparagiformis* changed to the sequence

obtained by ONT sequencing (GenBank accession PP280819) since the RefSeq sequence for

this bacterium contained multiple N nucleotides that biased the assignment of *E. asparagiformis*

to *E. lavalensis*). Count tables were then used to create a phyloseq (v1.44.0,

https://github.com/joey711/phyloseq) object in R [69]. Stacked bar plots were generated using

ggnested (v0.1.0, https://github.com/gmteunisse/ggnested) and fantaxtic (v0.2.0,

https://github.com/gmteunisse/Fantaxtic), and β-diversity analysis was performed using the

phyloseq ordinate() and plot_ordination() functions.

Total RNA extraction from microbial communities

A 300 μL fecal slurry aliquot (treated with either 100 μM urolithin C or an equivalent

volume DMSO for 48 h) was thawed and pelleted (6,500 g for 3 min). The pellet (in 200 μL

media) was then mixed with 800 μL TRI reagent (Zymo Research). Samples were lysed in a

Mini Beadbeater 16 (Biospec) according to the following sequence: 1 min ON, 5 min OFF for a

total of 5 min ON. RNA isolation was then performed using the Direct-zol RNA Miniprep Kit

(Zymo Research) according to the manufacturer's instructions (including an on-column DNase

digestion). To ensure complete DNA removal, an additional DNA digestion step was performed

on the isolated RNA using the Ambion DNA-free DNA Removal Kit (Invitrogen) according to the

manufacturer's instructions. The DNA-free RNA was then cleaned up using the OneStep PCR

Inhibitor Removal Kit (Zymo Research). RNA concentration and quality were verified by Qubit

RNA BR assay kit (Invitrogen) and 1% agarose gel electrophoresis.

RT-PCR analysis of the *ucd* operon in microbial communities

Total RNA was extracted from frozen fecal slurries as previously described (see Total

RNA extraction from microbial communities), and subsequently reverse transcribed as

described above (see *RT-PCR analysis of E. bolteae ucd operon structure*) in a reaction volume

of 5 μL. PCR reactions were conducted using the OneTaq 2X Master Mix with Standard Buffer

(NEB). The ucdCFO_RT-PCR primer pair was added to the master mix (to a final concentration

of 0.2 μM) and 1 μL of template (cDNA, -RT, or no template) was added for a total reaction

volume of 20 μL. PCR tubes were placed in a thermal cycler and targets were amplified

according to the following conditions: 30s at 94°C, 45 cycles (30s at 94°C, 1 min at 61°C, 4 min

at 68°C), 5 min at 68°C. A volume of 10 µL of reaction was directly loaded onto a 1% agarose

gel (made with 1X TAE buffer) containing SafeView Classic (Abm). PCR product sizes were

compared to the Quick-Load® Purple 1 kb Plus DNA Ladder (NEB).

PCR analysis of the *ucd* operon in microbial communities

1043 Genomic DNA (gDNA) was extracted from frozen fecal slurries as previously described

1044 (see Genomic DNA extraction from microbial communities). PCR reactions and product

1045 visualization was conducted on the gDNA as described above (see RT-PCR analysis of the *ucd*

1046 operon in microbial communities). In this case, 5 µL of PCR product was loaded onto the gels

1047 instead of 10 µL.

Statistical analyses and graphing

Statistical methods were not used to determine sample sizes, experiments were not

randomized, and the investigators were not blinded. Data points related to uroC metabolism

and RT-qPCR were assumed to be normally distributed, though this was not formally tested.

Correlation analyses were performed using the non-parametric Spearman rank correlation (ρ).

Statistical tests on bacterial relative abundances were performed using the Kruskal-Wallis test

on untransformed relative abundance values, which are skewed towards 0. Statistical analyses

for large datasets are detailed in the relevant methods sections. Details related to each test

performed are supplied in the figure legends. In all cases, $\alpha = 0.05$ and tests were two-tailed.

Data were plotted in GraphPad Prism (v10.0.0) or using the ggplot2 (v3.4.2) R package. Figures

were assembled in Affinity Designer (v1.10.6.1665).

Data Availability

RNA-seq reads were deposited in the NCBI SRA BioProject ID PRJNA996126 under BioSample accession codes SAMN36514640 (*Enterocloster bolteae* DSM 15670) and SAMN36514641 (*Enterocloster asparagiformis* DSM 15981). Reviewer link: https://dataview.ncbi.nlm.nih.gov/object/PRJNA996126?reviewer=fpbuj6eeuebv6mij3pme3mp8ep.

Untargeted proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD048514 and 10.6019/PXD048514. [70]

Oxford Nanopore 16S rRNA sequencing reads of healthy human fecal slurries were deposited in the NCBI SRA BioProject ID PRJNA1073957. Reviewer link: https://dataview.ncbi.nlm.nih.gov/object/PRJNA1073957?reviewer=nh1a04lg59enlti318qs23s6ot.

The 16S rRNA sequence for *E. asparagiformis* DSM 15670 used in the Emu database search was deposited in GenBank under accession PP280819.

All original code, tables, and RData files obtained from the analysis of curatedMetagenomicData were deposited in Zenodo (https://doi.org/10.5281/zenodo.8302320).

References

Knight, R. *et al.* The Microbiome and Human Biology. *Annu Rev Genomics Hum Genet* **18**, 65-86 (2017). https://doi.org/10.1146/annurev-genom-083115-022438

Taguer, M. & Maurice, C. F. The complex interplay of diet, xenobiotics, and microbial metabolism in the gut: Implications for clinical outcomes. *Clin Pharmacol Ther* **99**, 588-599 (2016). https://doi.org/10.1002/cpt.366

Almeida, A. *et al.* A unified catalog of 204,938 reference genomes from the human gut microbiome. *Nat Biotechnol* **39**, 105-114 (2021). https://doi.org/10.1038/s41587-020-0603-3

Spanogiannopoulos, P., Bess, E. N., Carmody, R. N. & Turnbaugh, P. J. The microbial pharmacists within us: a metagenomic view of xenobiotic metabolism. *Nat Rev Microbiol* **14**, 273-287 (2016). https://doi.org/10.1038/nrmicro.2016.17

Ervin, S. M. *et al.* Structural Insights into Endobiotic Reactivation by Human Gut Microbiome-Encoded Sulfatases. *Biochemistry* **59**, 3939-3950 (2020). https://doi.org/10.1021/acs.biochem.0c00711

Yao, L. *et al.* A biosynthetic pathway for the selective sulfonation of steroidal metabolites by human gut bacteria. *Nat Microbiol* **7**, 1404-1418 (2022). https://doi.org/10.1038/s41564-022-01176-y

Tomas-Barberan, F. A. & Espin, J. C. Effect of Food Structure and Processing on (Poly)phenol-Gut Microbiota Interactions and the Effects on Human Health. *Annu Rev Food Sci Technol* **10**, 221-238 (2019). https://doi.org/10.1146/annurev-food-032818-121615

Maier, L. *et al.* Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature* **555**, 623-628 (2018). https://doi.org/10.1038/nature25979

Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R. & Goodman, A. L. Mapping human microbiome drug metabolism by gut bacteria and their genes. *Nature* **570**, 462-467 (2019). https://doi.org/10.1038/s41586-019-1291-3

Maini Rekdal, V., Bess, E. N., Bisanz, J. E., Turnbaugh, P. J. & Balskus, E. P. Discovery and inhibition of an interspecies gut bacterial pathway for Levodopa metabolism. *Science* **364**, eaau6323 (2019). https://doi.org/10.1126/science.aau6323

Campbell, C. *et al.* Bacterial metabolism of bile acids promotes generation of peripheral regulatory T cells. *Nature* **581**, 475-479 (2020). https://doi.org/10.1038/s41586-020-2193-0

Paik, D. *et al.* Human gut bacteria produce Tau(Eta)17-modulating bile acid metabolites. *Nature* **603**, 907-912 (2022). https://doi.org/10.1038/s41586-022-04480-z

Cerqueira, F. M., Photenhauer, A. L., Pollet, R. M., Brown, H. A. & Koropatkin, N. M. Starch Digestion by Gut Bacteria: Crowdsourcing for Carbs. *Trends Microbiol* **28**, 95-108 (2020). https://doi.org/10.1016/j.tim.2019.09.004

Maini Rekdal, V. *et al.* A widely distributed metalloenzyme class enables gut microbial metabolism of host- and diet-derived catechols. *Elife* **9** (2020). https://doi.org/10.7554/eLife.50845

Garcia-Villalba, R., Selma, M. V., Espin, J. C. & Tomas-Barberan, F. A. Identification of Novel Urolithin Metabolites in Human Feces and Urine after the Intake of a Pomegranate Extract. *J Agric Food Chem* **67**, 11099-11107 (2019). https://doi.org/10.1021/acs.jafc.9b04435

Carmody, R. N. *et al.* Diet dominates host genotype in shaping the murine gut microbiota. *Cell Host Microbe* **17**, 72-84 (2015). https://doi.org/10.1016/j.chom.2014.11.010

17 Carmody, R. N. *et al.* Cooking shapes the structure and function of the gut microbiome.
Nat Microbiol **4**, 2052-2063 (2019). https://doi.org/10.1038/s41564-019-0569-4
18 Sonnenburg, E. D. *et al.* Diet-induced extinctions in the gut microbiota compound over
generations. *Nature* **529**, 212-215 (2016). https://doi.org/10.1038/nature16504
19 Rothschild, D. *et al.* Environment dominates over host genetics in shaping human gut
microbiota. *Nature* **555**, 210-215 (2018). https://doi.org/10.1038/nature25973
20 Quideau, S., Deffieux, D., Douat-Casassus, C. & Pouysegu, L. Plant polyphenols:
chemical properties, biological activities, and synthesis. *Angew Chem Int Ed Engl* **50**,
586-621 (2011). https://doi.org/10.1002/anie.201000044
21 van Duynhoven, J. *et al.* Metabolic fate of polyphenols in the human superorganism.
Proc Natl Acad Sci U S A **108 Suppl 1**, 4531-4538 (2011).
https://doi.org/10.1073/pnas.1000098107
22 Garcia-Villalba, R. *et al.* Urolithins: a Comprehensive Update on their Metabolism,
Bioactivity, and Associated Gut Microbiota. *Mol Nutr Food Res* **66**, e2101019 (2022).
https://doi.org/10.1002/mnfr.202101019
23 Singh, A. *et al.* Direct supplementation with Urolithin A overcomes limitations of dietary
exposure and gut microbiome variability in healthy adults to achieve consistent levels
across the population. *Eur J Clin Nutr* **76**, 297-308 (2022).
https://doi.org/10.1038/s41430-021-00950-1
24 Andreux, P. A. *et al.* The mitophagy activator urolithin A is safe and induces a molecular
signature of improved mitochondrial and cellular health in humans. *Nat Metab* **1**, 595-
603 (2019). https://doi.org/10.1038/s42255-019-0073-4
25 Denk, D. *et al.* Expansion of T memory stem cells with superior anti-tumor immunity by
Urolithin A-induced mitophagy. *Immunity* **55**, 2059-2073 e2058 (2022).
https://doi.org/10.1016/j.immuni.2022.09.014
26 Ryu, D. *et al.* Urolithin A induces mitophagy and prolongs lifespan in C. elegans and
increases muscle function in rodents. *Nat Med* **22**, 879-888 (2016).
https://doi.org/10.1038/nm.4132
27 Ghosh, S., Banerjee, M., Haribabu, B. & Jala, V. R. Urolithin A attenuates arsenic-
induced gut barrier dysfunction. *Arch Toxicol* **96**, 987-1007 (2022).
https://doi.org/10.1007/s00204-022-03232-2
28 Singh, R. *et al.* Enhancement of the gut barrier integrity by a microbial metabolite through
the Nrf2 pathway. *Nat Commun* **10**, 89 (2019). https://doi.org/10.1038/s41467-018-
07859-7
29 Shen, P. X. *et al.* Urolithin A ameliorates experimental autoimmune encephalomyelitis
by targeting aryl hydrocarbon receptor. *EBioMedicine* **64**, 103227 (2021).
https://doi.org/10.1016/j.ebiom.2021.103227
30 Ghosh, S., Haribabu, B. & Jala, V. R. Microbial metabolite Urolithin A induces expansion
of Regulatory T Cells in aryl hydrocarbon receptor (AhR)-dependent manner. *Journal of
Immunology* **206**, 113.103-113.103 (2021).
31 Xia, B. *et al.* Urolithin A exerts antiobesity effects through enhancing adipose tissue
thermogenesis in mice. *PLoS Biol* **18**, e3000688 (2020).
https://doi.org/10.1371/journal.pbio.3000688
32 Selma, M. V. *et al.* Isolation of Human Intestinal Bacteria Capable of Producing the
Bioactive Metabolite Isourolithin A from Ellagic Acid. *Front Microbiol* **8**, 1521 (2017).
https://doi.org/10.3389/fmicb.2017.01521
33 Selma, M. V., Beltran, D., Garcia-Villalba, R., Espin, J. C. & Tomas-Barberan, F. A.
Description of urolithin production capacity from ellagic acid of two human intestinal

Gordonibacter species. *Food Funct* **5**, 1779-1784 (2014). https://doi.org/10.1039/c4fo00092g

Gaya, P., Peirotén, A., Medina, M., Alvarez, I. & Landete, J. M. Bifidobacterium pseudocatenulatum INIA P815: The first bacterium able to produce urolithins A and B from ellagic acid. *Journal of Functional Foods* **45**, 95-99 (2018). https://doi.org/10.1016/j.jff.2018.03.040

Garcia-Villalba, R. *et al.* Metabolism of different dietary phenolic compounds by the urolithin-producing human-gut bacteria Gordonibacter urolithinfaciens and Ellagibacter isourolithinifaciens. *Food Funct* **11**, 7012-7022 (2020). https://doi.org/10.1039/d0fo01649g

Iglesias-Aguirre, C. E. *et al.* Gut Bacteria Involved in Ellagic Acid Metabolism To Yield Human Urolithin Metabotypes Revealed. *J Agric Food Chem* **71**, 4029-4035 (2023). https://doi.org/10.1021/acs.jafc.2c08889

Iglesias-Aguirre, C. E. *et al.* In vivo administration of gut bacterial consortia replicates urolithin metabotypes A and B in a non-urolithin-producing rat model. *Food Funct* **14**, 2657-2667 (2023). https://doi.org/10.1039/d2fo03957e

Hosny, M. *et al.* Clostridium pacaense: a new species within the genus Clostridium. *New Microbes New Infect* **28**, 6-10 (2019). https://doi.org/10.1016/j.nmni.2018.12.003

Zupok, A., Iobbi-Nivol, C., Mejean, V. & Leimkuhler, S. The regulation of Moco biosynthesis and molybdoenzyme gene expression by molybdenum and iron in bacteria. *Metallomics* **11**, 1602-1624 (2019). https://doi.org/10.1039/c9mt00186g

Leimkuhler, S. & Iobbi-Nivol, C. Bacterial molybdoenzymes: old enzymes for new purposes. *FEMS Microbiol Rev* **40**, 1-18 (2016). https://doi.org/10.1093/femsre/fuv043

Koppel, N., Bisanz, J. E., Pandelia, M. E., Turnbaugh, P. J. & Balskus, E. P. Discovery and characterization of a prevalent human gut bacterial enzyme sufficient for the inactivation of a family of plant toxins. *Elife* **7** (2018). https://doi.org/10.7554/eLife.33953

Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021). https://doi.org/10.1038/s41586-021-03819-2

Varadi, M. *et al.* AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**, D439-D444 (2022). https://doi.org/10.1093/nar/gkab1061

van Kempen, M. *et al.* Fast and accurate protein structure search with Foldseek. *Nat Biotechnol* (2023). https://doi.org/10.1038/s41587-023-01773-0

Resch, M., Dobbek, H. & Meyer, O. Structural and functional reconstruction in situ of the [CuSMoO2] active site of carbon monoxide dehydrogenase from the carbon monoxide oxidizing eubacterium Oligotropha carboxidovorans. *J Biol Inorg Chem* **10**, 518-528 (2005). https://doi.org/10.1007/s00775-005-0006-4

Ishikita, H., Eger, B. T., Okamoto, K., Nishino, T. & Pai, E. F. Protein conformational gating of enzymatic activity in xanthine oxidoreductase. *J Am Chem Soc* **134**, 999-1009 (2012). https://doi.org/10.1021/ja207173p

Pasolli, E. *et al.* Accessible, curated metagenomic data through ExperimentHub. *Nat Methods* **14**, 1023-1024 (2017). https://doi.org/10.1038/nmeth.4468

Cortes-Martin, A. *et al.* The gut microbiota urolithin metabotypes revisited: the human metabolism of ellagic acid is mainly determined by aging. *Food Funct* **9**, 4100-4106 (2018). https://doi.org/10.1039/c8fo00956b

Bess, E. N. *et al.* Genetic basis for the cooperative bioactivation of plant lignans by Eggerthella lenta and other human gut bacteria. *Nat Microbiol* **5**, 56-66 (2020). https://doi.org/10.1038/s41564-019-0596-1

50   Le, C. C., Bae, M., Kiamehr, S. & Balskus, E. P. Emerging Chemical Diversity and Potential Applications of Enzymes in the DMSO Reductase Superfamily. *Annu Rev Biochem* **91**, 475-504 (2022). https://doi.org/10.1146/annurev-biochem-032620-110804

51   Little, A. S. *et al.* Dietary- and host-derived metabolites are used by diverse gut bacteria for anaerobic respiration. *Nat Microbiol* **9**, 55-69 (2024). https://doi.org/10.1038/s41564-023-01560-2

52   Pinske, C., Bonn, M., Kruger, S., Lindenstrauss, U. & Sawers, R. G. Metabolic deficiences revealed in the biotechnologically important model bacterium Escherichia coli BL21(DE3). *PLoS One* **6**, e22830 (2011). https://doi.org/10.1371/journal.pone.0022830

53   Xia, M. *et al.* Improvement of Urolithin A Yield by In Vitro Cofermentation of Streptococcus thermophilus FUA329 with Human Gut Microbiota from Different Urolithin Metabotypes. *J Agric Food Chem* (2024). https://doi.org/10.1021/acs.jafc.3c09734

54   Zhang, X. *et al.* Isolation and characterization of a novel human intestinal Enterococcus faecium FUA027 capable of producing urolithin A from ellagic acid. *Front Nutr* **9**, 1039697 (2022). https://doi.org/10.3389/fnut.2022.1039697

55   Meier-Kolthoff, J. P., Carbasse, J. S., Peinado-Olarte, R. L. & Goker, M. TYGS and LPSN: a database tandem for fast and reliable genome-based classification and nomenclature of prokaryotes. *Nucleic Acids Res* **50**, D801-D807 (2022). https://doi.org/10.1093/nar/gkab902

56   Pauff, J. M., Cao, H. & Hille, R. Substrate Orientation and Catalysis at the Molybdenum Site in Xanthine Oxidase: CRYSTAL STRUCTURES IN COMPLEX WITH XANTHINE AND LUMAZINE. *J Biol Chem* **284**, 8760-8767 (2009). https://doi.org/10.1074/jbc.M804517200

57   Meier-Kolthoff, J. P., Goker, M., Sproer, C. & Klenk, H. P. When should a DDH experiment be mandatory in microbial taxonomy? *Arch Microbiol* **195**, 413-418 (2013). https://doi.org/10.1007/s00203-013-0888-4

58   Meier-Kolthoff, J. P. & Goker, M. TYGS is an automated high-throughput platform for state-of-the-art genome-based taxonomy. *Nat Commun* **10**, 2182 (2019). https://doi.org/10.1038/s41467-019-10210-3

59   Ewels, P., Magnusson, M., Lundin, S. & Kaller, M. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047-3048 (2016). https://doi.org/10.1093/bioinformatics/btw354

60   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10 (2011). https://doi.org/10.14806/ej.17.1.200

61   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907-915 (2019). https://doi.org/10.1038/s41587-019-0201-4

62   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014). https://doi.org/10.1093/bioinformatics/btt656

63   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014). https://doi.org/10.1186/s13059-014-0550-8

64   Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* **3**, 1101-1108 (2008). https://doi.org/10.1038/nprot.2008.73

65   Keller, A., Nesvizhskii, A. I., Kolker, E. & Aebersold, R. Empirical statistical model to
estimate the accuracy of peptide identifications made by MS/MS and database search.
Anal Chem **74**, 5383-5392 (2002). https://doi.org/10.1021/ac025747h
66   Nesvizhskii, A. I., Keller, A., Kolker, E. & Aebersold, R. A statistical model for identifying
proteins by tandem mass spectrometry. *Anal Chem* **75**, 4646-4658 (2003).
https://doi.org/10.1021/ac0341261
67   Beghini, F. *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse
microbial communities with bioBakery 3. *Elife* **10**, e65088 (2021).
https://doi.org/10.7554/eLife.65088
68   Curry, K. D. *et al.* Emu: species-level microbial community profiling of full-length 16S
rRNA Oxford Nanopore sequencing data. *Nat Methods* **19**, 845-853 (2022).
https://doi.org/10.1038/s41592-022-01520-4
69   McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).
https://doi.org/10.1371/journal.pone.0061217
70   Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass
spectrometry-based proteomics evidences. *Nucleic Acids Res* **50**, D543-D552 (2022).
https://doi.org/10.1093/nar/gkab1038

1283

Acknowledgements

This research was funded by the Canadian Institutes of Health Research (CIHR) grant PJT-437944 to B.C. and C.F.M. and a Weston Family Microbiome Initiative's 2021 Transformational Research program to B.C. B.C. holds a tier II Canada Research Chair (CRC) in Therapeutic Chemistry. C.F.M. holds a tier II CRC in Gut Microbial Physiology and is a Azrieli Global Scholar in the Humans & Microbiome program. R.P. is supported by the CIHR Canada Graduate Scholarship-Doctoral (#493808 for 2023-2026) and by the Fonds de Recherche du Québec-Santé : Bourse de formation au doctorat (#316063 for 2022-2023). M.S. is supported by the CIHR Canada Graduate Scholarship to Honor Nelson Mandela (#DF2-187718 for 2023-2026), the Fonds de Recherche du Québec-Santé: Bourse de formation au doctorat (#311071 for 2022-2023).

RNA sample preparation and sequencing was performed by Génome Québec. Untargeted proteomics sample preparation and analysis was performed by the Research Institute of the McGill University Health Centre (RI-MUHC) Proteomics and Molecular Analysis Platform. Small-molecule MALDI-TOF mass spectrometry was performed by Mark Hancock at the McGill SPR-MS Facility. Molecular biology experiments were enabled by the McGill University Imaging and Molecular Biology Platform (IMBP). High-performance cloud computing was enabled by Calcul Québec (https://www.calculquebec.ca/) and the Digital Research Alliance of Canada (https://alliancecan.ca/). RNA-seq data analysis was performed using two instances of Galaxy: (https://usegalaxy.org/) and Compute Canada Genetics and Genomics Analysis Platform (GenAP) (https://www.genap.ca/).

Author information

Authors and Affiliations

Department of Pharmacology & Therapeutics, McGill University, 3655 Prom. Sir-William-Osler,

Montreal, Quebec, H3G 1Y6, Canada

Reilly Pidgeon, Sacha Mitchell, Layan Suleiman, Lharbi Dridi & Bastien Castagner

Department of Microbiology & Immunology, McGill University, 3775 University Street, Montreal,

Quebec, H3A 2B4, Canada

Michael Shamash & Corinne F. Maurice

Contributions

R.P. designed the study, performed microbiology experiments, performed bioinformatics analyses, analyzed data, created figures, and wrote the initial manuscript with B.C. **S.M.** performed experiments on microbial communities, analyzed data, and created figures. **M.S.** performed Nanopore sequencing and processed raw sequencing reads. **L.S.** performed microbiology experiments. **L.D.** consulted on experimental design and methodology. **C.M.** supervised research, obtained ethical approval for the use of human fecal samples, and obtained research funding. **B.C.** designed the study, supervised the research, obtained research funding, and wrote the initial manuscript with R.P. All authors reviewed and edited the manuscript.

Corresponding authors

Correspondence to Bastien Castagner (bastien.castagner@mcgill.ca).

Ethics declarations

The authors declare no competing interests.