The dietary ellagitannin metabolite urolithin A is produced by a molybdenum-dependent dehydroxylase encoded by prevalent human gut *Enterocloster* spp.

Reilly Pidgeon[1], Sacha Mitchell[1], Michael Shamash[2], Layan Suleiman[1], Lharbi Dridi[1], Corinne F. Maurice[2, 3], Bastien Castagner[1*]

[1] Department of Pharmacology & Therapeutics, McGill University, 3655 Prom. Sir-William-Osler, Montreal, Quebec, H3G 1Y6, Canada.

[2] Department of Microbiology & Immunology, McGill University, 3775 University Street, Montreal, Quebec, H3A 2B4, Canada

[3] McGill Centre for Microbiome Research, Montreal, Quebec, Canada

Abstract

Abstract

Urolithin A (uroA) is a polyphenol derived from the multi-step metabolism of dietary ellagitannins

by the human gut microbiota that can affect host health by stimulating mitophagy. Most

individuals harbor a microbiota capable of uroA production; however, the mechanisms

underlying the dehydroxylation of its catechol-containing dietary precursor (uroC) are unknown.

Here, we use a combination of untargeted bacterial transcriptomics, proteomics, and

comparative genomics to uncover an inducible uroC dehydroxylase (*ucd*) operon in

Enterocloster spp. We show that *Enterocloster* spp. are sensitive to iron chelation by uroC, and

dehydroxylation to uroA rescues growth by disrupting the iron-binding catechol. Importantly,

only microbiota samples actively transcribing *ucd* could produce uroA, establishing *ucd*-

containing *Enterocloster* spp. as keystone urolithin metabolizers. Overall, this work identifies

Enterocloster spp. and the *ucd* operon as main contributors to uroA production and establishes

a multi-omics framework to further our mechanistic understanding of polyphenol metabolism by

the human gut microbiota.

Main

The human gut microbiota is a collection of trillions of microorganisms that colonize the gastrointestinal tract and play pivotal roles in host health and disease [1]. Gut bacteria help maintain homeostasis by regulating host immune cell activity, gut barrier integrity, and nutrient availability [2]. One of the main mediators of microbiota-host interactions are microbial metabolites. Gut bacteria possess an immense metabolic repertoire (nearly 1000-fold more protein coding sequences than the human genome [3]) to perform four main classes of reactions: hydrolysis, conjugation, cleavage, and reduction [4-7]. These ubiquitous reactions have been linked to microbiota-dependent metabolism of therapeutic drugs [8-10], host bile acids [6,11,12], and diet-derived compounds [13-15].

Diet is a strong modulator of the composition and function of the gut microbiota [16-19]. Diet-derived polyphenols are a diverse class of plant secondary metabolites found in fruits, vegetables, and nuts (reviewed in [20]) that are poorly absorbed by the host and reach the large intestine relatively intact [7,21]. Ellagitannins are a large sub-group of polyphenols that belong to the family of hydrolysable tannins and are characterized by a central glucose (open-chain or pyranose forms) linked to diverse pyrogallol-like moieties [20]. Camu camu, a berry rich in the ellagitannin castalagin, has been shown to impact anti-cancer immunity via the gut microbiome, and is currently in clinical trials (NCT05303493, NCT06049576) in combination with immune checkpoint inhibitors [22,23]. Depending on microbiota composition, ellagitannins can be hydrolyzed and reduced by gut bacteria into bioactive metabolites (ellagic acid, urolithins, nasutins) according to different metabolic phenotypes characterized by the terminal metabolites observed in biological fluids [24] (Supplementary Fig. 1).

Urolithin A (uroA) is the most common terminal metabolite of ellagitannin metabolism and has

reported pharmacological activities both within the gut environment and systemically following

absorption [25]. In the gut, uroA can attenuate colitis by increasing the expression of epithelial

tight junction proteins [26-28] *via* the activation of aryl hydrocarbon receptor (AhR)-Nrf2 pathways

[29,30]. Additionally, uroA can enhance immunotherapy in colorectal cancer models by activating

Pink1-dependent mitophagy pathways in T cells, improving anti-tumor CD8+ T cell immunity [31].

Clinical trials in healthy individuals have demonstrated that uroA is safe, bioavailable, and can

be detected in its aglycone, glucuronidated, and sulfated forms in plasma [25,27]. Once absorbed

by the host, uroA can trigger mitophagy in muscle cells, improving muscle function in animal

models of ageing and Duchenne muscular dystrophy [26,29,32,33]. Overall, uroA can enhance gut

barrier integrity, modulate the immune system, and promote mitochondrial health in the host,

thus showing promise as a postbiotic to treat age-related conditions [34-36].

While urolithin metabolism is prevalent in human populations, few gut bacteria have been

reported to metabolize urolithins [34-36]. Most known urolithin metabolizers belong to the

Eggerthellaceae family (*Gordonibacter urolithinfaciens*, *Gordonibacter pamelaeae*, *Ellagibacter*

isourolithinifaciens) and can perform multiple metabolic steps in the urolithin metabolism

pathway, yielding either urolithin C (uroC) or isourolithin A (isouroA) from ellagic acid [37].

Recently, certain members of the *Enterocloster* spp. (*Lachnospiraceae* family) were reported

to dehydroxylate uroC to uroA and isouroA to urolithin B (uroB) both *in vitro* and *in vivo* [38,39].

These findings shed light on the minimal bacterial community required for the complete

metabolism of ellagic acid to uroA; however, the genes and enzymes responsible for these

dehydroxylation reactions remain unknown (Fig. 1A).

Here, we use a multi-omics enzyme identification framework to uncover uroC dehydroxylase

(*ucd*) genes and enzymes in *Enterocloster* spp. and their relative, *Lachnoclostridium pacaense*.

We find that the UcdCFO enzyme complex specifically dehydroxylates 9-hydroxy urolithins and

that both metabolizing species and *ucd* genes are prevalent and actively transcribed in human

feces during *ex vivo* metabolism. We further demonstrate that *Enterocloster* spp. growth is

delayed by uroC and that dehydroxylation may be a mechanism to inactivate its iron-binding

catechol moiety. Our study sheds light on the genetic and chemical basis underlying the

complex reciprocal interactions between urolithins and the gut microbiota.

Results

A subset of *Enterocloster* species converts urolithin C to urolithin A *in vitro*.

Members of the *Enterocloster* spp. have previously been shown to dehydroxylate uroC

in vitro [38] and *in vivo* [39] under anaerobic conditions (Fig. 1A, full metabolic pathway in

Supplementary Fig. 1). To determine the prevalence of uroC metabolism within this genus, we

incubated all available *Enterocloster* spp. type strains (Methods) with uroC and quantified

urolithin concentrations by liquid chromatography-mass spectrometry (LC-MS). Of the tested

bacteria, only *E. asparagiformis* , *E. bolteae* , and *E. citroniae* dehydroxylated uroC to produce

uroA (Fig. 1B). Interestingly, uroC metabolism was not predicted by phylogeny, as uroC-

metabolizing species did not cluster based on 16S rRNA genes, genomes, or proteomes (Fig.

1C, Supplementary Fig. 2A,B, respectively), suggesting gain or loss of metabolic gene clusters

throughout the evolution of *Enterocloster* spp. Based on these results, we chose to perform

more in-depth analysis on *E. asparagiformis* and *E. bolteae* to identify the metabolic gene

clusters involved in uroC dehydroxylation.

Figure 1. Urolithin C metabolism by *Enterocloster* spp. is not predicted by phylogeny.

A) Reaction scheme of uroC dehydroxylation by gut resident *Enterocloster* spp. via unknown enzymes. **B)** LC-MS screen of *Enterocloster* spp. type strains for dehydroxylation activity. UroC (100 μM) was added to cultures (in mABB+H media) at the start of growth and urolithins were extracted after 24 h anaerobic incubation, then analyzed by LC-MS. Left: Representative chromatograms (λ = 305 nm) for each experimental group (from one representative biological replicate). The same scale was used for each chromatogram. Right: Quantification of urolithin peak areas relative to a salicylic acid internal standard (IS) (n = 3 biological replicates). Data are represented as mean ± SEM. **C)** Phylogenetic tree of tested *Enterocloster* spp. type strain 16S rRNA sequences constructed using the Genome-to-Genome Distance Calculator (GGDC) Phylogeny Server [40]. Maximum likelihood (ML) tree inferred under the GTR+GAMMA model and rooted by midpoint-rooting. The branches are scaled in terms of the expected number of substitutions per site. The numbers above the branches are support values when larger than 60% from ML (left) and maximum parsimony (right) bootstrapping. The GenBank accession numbers are provided to the right of each taxon. Source data and statistical details are provided as a Source data file.

A putative urolithin C dehydroxylase metabolic gene cluster is upregulated upon urolithin C treatment.

To understand when uroC metabolism machinery was being expressed, we first sought to characterize the kinetics of uroC dehydroxylation in rich media (mABB+H). Therefore, a simultaneous growth and metabolism experiment was designed, whereby uroC was spiked into *E. asparagiformis* and *E. boltea*e cultures during the exponential phase of growth and metabolites were measured by LC-MS (Fig. 2A). Treatment with uroC during the exponential phase did not affect growth of either bacterium compared to the DMSO control (Fig. 2B). In both bacteria, quantitative conversion of uroC to uroA occurred within 4 h post-spike-in (Fig. 2C,D), demonstrating that metabolism in rich media is fast and robust.

We next sought to determine whether uroC metabolism is inducible or constitutive. To test for inducibility, both bacteria were treated with DMSO or uroC during exponential growth, then washed and resuspended in PBS, yielding cell suspensions unable to synthesize new proteins. Metabolism of uroC to uroA was inducible as only cells originating from bacteria grown with

uroC were capable of uroA production (Fig. 2E). Consequently, we performed RNA-sequencing

to compare gene expression in DMSO and uroC-treated cultures of *E. asparagiformis* and *E.*

bolteae. Since uroA was detected in both bacterial cultures as soon as 2 h post spike-in (Fig.

2C), this timepoint was selected to isolate mRNA.

RNA sequencing of uroC-induced cultures revealed a distinct gene cluster, which we term uroC

dehydroxylase (*ucd*), that was highly and differentially expressed ($\log_2$FC > 8) in both *E.*

asparagiformis (Fig. 2F) and *E. bolteae* (Fig. 2G). In both bacteria, these clusters contained

adjacent genes that were expressed to similar $\log_2$FC values: a xanthine dehydrogenase family

protein subunit M, a (2Fe-2S)-binding protein, and a molybdopterin-dependent oxidoreductase

(Fig. 2F,G). These genes will hereafter be referred to as *ucdC* (for coenzyme), *ucdF* (for

ferredoxin), and *ucdO* (for oxidoreductase), respectively. Interestingly, we also observed an

upregulation of genes involved in efflux (MepA-like multidrug and toxin extrusion (MATE)

transporters) and iron transport (FecCD-like) (Fig. 2F,G), suggesting a link between uroC

metabolism, iron uptake, and efflux.

Figure 2. Urolithin C treatment upregulates a putative dehydroxylase operon.

A) Experimental design of uroC (100 μM) spike-in experiments during the exponential phase of growth. For each biological replicate in this design, growth (B), metabolism (C,D), and RNA-seq (F-G) results are matched. **B)** Growth curve (optical density (OD) at 620 nm) of DMSO or uroC-spiked *E. asparagiformis* (*Ea*) and *E. bolteae* (*Eb*) type strain cultures according to the design in (A). 200 μL of culture were sampled at each timepoint and OD_{620} was measured in a 96-well plate (n = 4 biological replicates). The same sampled culture was then frozen and extracted for analysis by LC-MS (C,D). **C)** Representative chromatograms (λ = 305 nm) of cultures sampled 2 h post-spike-in (from one representative biological replicate). The same scale was used for each chromatogram. **D)** Quantification of urolithin concentrations from peak areas relative to a salicylic acid internal standard (IS) over 4 h in uroC-spiked *Ea* and *Eb* type strain cultures (n = 4 biological replicates). **E)** Quantification of urolithin A concentrations in DMSO- or uroC-treated *Ea* and *Eb* cell suspensions. Cell suspensions were prepared from *Ea* and *Eb* cells grown with either DMSO or 50 μM uroC. The cells were washed and resuspended in PBS to halt the production of new enzymes, then treated with DMSO or 100 μM uroC (n = 3 biological replicates). **F,G)** Manhattan plots of genes altered by uroC treatment in *Ea* (F) and *Eb* (G) based on DESeq2 analysis (n = 4 biological replicates). Data points are colored according to their adjusted p-value (based on the Benjamini-Hochberg-corrected Wald statistic). Grey, p-adj ≥ 0.05. Red or blue, p-adj < 0.05 for *Ea* and *Eb*, respectively. The genomic organization around the differentially expressed genes (generated from the NCBI Sequence

Viewer) is depicted above Manhattan plots, which show the most highly and differentially
expressed genes by RNA-seq. Genes are colored according to their $\log_2$FC values. NCBI
accessions for select proteins encoded by highlighted genes are provided. **H)** Primer design for
RT-PCR (I) experiment targeting the *Eb ucd* gene cluster. The same reverse primer was used
for both the reverse transcription step and the subsequent PCR reaction. **I)** 1% agarose gel
image of RT-PCR amplicons using primers (H) that span the full-length *Eb ucd* gene cluster
(from one biological replicate). NTC, no template control. **J)** RT-qPCR expression of each gene
in the *Eb ucd* operon. Growing *Eb* cultures were treated with DMSO or uroC (100 µM) for 2 h
before RNA isolation and reverse transcription (n = 3 biological replicates). Gene expression
profiles of each target gene in the *Eb ucd* gene cluster displayed as $\log_2$FC (equivalent to -
$\Delta\Delta C_t$, where $\Delta\Delta C_t = \Delta C_{t\ uroC} - \Delta C_{t\ DMSO}$) with lines connecting paired biological replicates;
repeated-measures one-way ANOVA with Tukey's multiple comparisons test; ns, not
significant. Data are represented as mean ± SEM (behind symbols) in (B,D,E). FC, fold change
(uroC/DMSO); Source data and statistical details are provided as a Source data file.

The *ucd* metabolic gene cluster is organized in an operon.

We next sought to characterize the *ucd* metabolic gene cluster in *E. bolteae* since this

bacterium is considered a core species of the gut microbiome [8]. Based on the proximity, sense,

and expression levels of each of the three genes by RNA-seq (Fig. 2G), we hypothesized that

all three genes in the cluster were organized in an operon. We designed a gene-specific RT-

PCR assay that would enable the detection of full-length polycistronic *ucdCFO* genes using

cDNA from DMSO- or uroC-treated *E. bolteae* as a template (Fig. 2H). An amplicon of the

expected size (~3.6 kb) was detected only in cDNA derived from uroC-treated *E. bolteae*,

validating the inducibility of these genes (Fig. 2I). Long-read sequencing of the obtained

amplicon yielded a sequence corresponding to the *E. bolteae ucdCFO* metabolic gene cluster

with 100% identity (Supplementary Sequence 1). Using an independent set of *E. bolteae*

cultures, we then performed RT-qPCR on DMSO- or uroC-treated *E. bolteae* with all three

genes in the putative operon as targets. Similar to our RNA-seq results, all three genes were

highly induced (mean $\log_2$FC ≥ 9.7 for all three *ucd* genes) relative to DMSO controls and were

expressed at the same level (Fig. 2J). These results indicate that the *ucdCFO* genes are

transcribed as a single polycistronic mRNA and therefore form a uroC-inducible operon.

The *ucd* operon is induced by 9-hydroxy urolithins.

Next, we aimed to determine the substrate scope of the *ucd* operon. Multiple urolithins possess pyrogallol, catechol, and phenol structural motifs that are dehydroxylated at various positions by gut bacteria (Supplementary Fig. 1). Interestingly, *E. bolteae* only metabolizes the 9-position hydroxyl group of urolithins (Supplementary Fig. 3A-D) and does not require adjacent hydroxyl groups since isouroA is dehydroxylated to uroB (Supplementary Fig. 3D) [38]. Since dehydroxylation of urolithins in *E. bolteae* is position specific, we hypothesized that the *ucd* operon would be induced by other 9-hydroxy urolithins (uroM6 and isouroA). Therefore, we performed RT-qPCR on DMSO-, uroM6-, uroC-, or isouroA-treated *E. bolteae* cultures using the *ucdO* gene as a target. Each urolithin significantly induced the expression of the *ucd* operon to a similar extent (Supplementary Fig. 3E-F). In addition, *E. bolteae* cell suspensions induced with uroC were capable of dehydroxylating uroM6 (Supplementary Fig. 3G) and isouroA (Supplementary Fig. 3H), indicating that the same proteins induced by uroC can metabolize structurally similar 9-hydroxy urolithins. Thus, it is likely that the same metabolic enzymes, encoded by the *E. bolteae ucd* operon, are acting on 9-hydroxy urolithins.

Presence of *ucd* operon homologs in genomes predicts urolithin C metabolism by gut bacteria.

We wondered whether novel metabolizers of uroC could be discovered based on nucleotide sequence homology to the *ucd* operon. Homology searches using the *E. bolteae ucd* operon sequence confirmed that only uroC-metabolizing *Enterocloster* spp. (*E. asparagiformis*, *E. bolteae*, and *E. citroniae*) possessed homologs of the *ucdCFO* genes with a similar organization (Supplementary Fig. 4). In addition, the gut bacterium *Lachnoclostridium pacaense* [41] was identified as another hit (Supplementary Fig. 4). The type strain of this bacterium (CCUG 71489T = Marseille-P3100) was closely related to *Enterocloster* spp. based

on 16S rRNA, whole genome, and whole proteome phylogenies (Fig. 3A, Supplementary Fig.

2A,B, respectively). *L. pacaense* possessed genomic sequences with high homology (86.5%

nucleotide identity) and identical functional annotations to the *E. bolteae ucd* operon sequence

(Fig. 3B). When incubated with uroC, *L. pacaense* CCUG 71489T quantitatively produced uroA

(Fig. 3C,D). We searched for homologs of the *E. bolteae ucd* in the genomes of urolithin- and

catechol-metabolizing bacteria belonging to the *Eggerthellaceae* but could not identify any hits.

Notably, *Eggerthellaceae* lack 9-hydroxy urolithin dehydroxylase activity [35,37], which correlates

with an absence of *ucd*-like operons in their genomes (Fig. 3B). These comparative genomics

data indicate that the presence of a *ucd* operon in genomes predicts uroC metabolism by gut

bacteria.

Figure 3. Urolithin C metabolism correlates with *ucd* operon prevalence in gut bacteria.

A) Phylogenetic tree of *Enterocloster* spp., *Lachnoclostridium pacaense* (*Lp*), and catechol-
metabolizing *Eggerthellaceae* type strain 16S rRNA sequences constructed using the Genome-
to-Genome Distance Calculator (GGDC) Phylogeny Server [40]. Maximum likelihood (ML) tree
inferred under the GTR+GAMMA model and rooted by midpoint-rooting. The branches are
scaled in terms of the expected number of substitutions per site. The numbers above the
branches are support values when larger than 60% from ML (left) and maximum parsimony
(right) bootstrapping. Bacteria that convert uroC to uroA are labeled with grey squares. **B)** NBCI
Multiple Sequence Aligner viewer hits for BLASTn searches using the *E. bolteae* DSM 15670
ucd operon nucleotide sequence as a query against the NCBI refseq_genomes database. Only

hits with ≥ 90 % query coverage and species-level taxonomic resolution are displayed with % identity to the query sequence. Domain annotations for each gene are denoted below according to InterPro annotations for corresponding proteins. ND, Not detected; Moco, Molybdenum cofactor. **C,D)** *In vitro* metabolism of uroC by *Lp*. UroC (100 µM) was added to cultures (in mABB+H media) at the start of growth and urolithins were extracted after 24 h anaerobic incubation, then analyzed by LC-MS. **C)** Representative chromatograms (λ = 305 nm) for each experimental group (from one representative biological replicate). The same scale was used for each chromatogram. **D)** Quantification of urolithin peak areas relative to a salicylic acid internal standard (IS) (n = 3 biological replicates). Data are represented as mean ± SEM. Source data and statistical details are provided as a Source data file.

A molybdopterin cofactor biosynthetic gene cluster is upregulated upon urolithin C treatment.

In addition to the three genes in the *ucd* operon, we observed a significant increase ($\log_2$FC ≥ 2.6) in 9 molybdopterin cytosine dinucleotide (MCD) biosynthesis genes upon uroC treatment (Fig. 2F,G, Supplementary Fig. 5A). These 9 genes, which recapitulate the function of 10 genes in *E. coli* [42], are involved in molybdenum cofactor biosynthesis (*moaAC*, *mogA*, *moeA*), molybdate ion transport (*modABCE*), cytosine addition to the molybdenum cofactor (*mocA*), and MCD cofactor insertion into the active site (*xdhC*) (Supplementary Fig. 5B) [42]. All 9 genes cluster in the genomes of *E. asparagiformis* and *E. bolteae* and are organized into 2 adjacent operons (Supplementary Fig. 5C) that are induced upon uroC treatment. Based on sequence homology to *E. coli* oxidoreductases and MCD biosynthetic machinery, proteins encoded by the *ucd* operon belong to the xanthine dehydrogenase family [43]. These findings imply that uroC dehydroxylation is MCD-dependent, which differs from the bis-molybdopterin guanine dinucleotide requirement of catechol dehydroxylases in *Eggerthellaceae* [10,14].

The UcdCFO complex enables anaerobic electron transport from NADH to uroC

Since oxidoreductases utilise a variety of cofactors and coenzymes for catalytic activity, we sought to determine the redox coenzymes and conditions necessary for uroC

dehydroxylation. Therefore, we performed metabolism assays using crude lysates from uroC-induced *E. bolteae*. As crude lysates alone did not metabolize uroC, various redox coenzymes (NADPH, NADH, and FAD) were added to lysates to promote uroC dehydroxylation (Fig. 4A). Only NADH-treated lysates yielded quantitative dehydroxylation of uroC to uroA compared to the no cofactor control (Fig. 4A). Interestingly, the addition of free FAD partially inhibited uroC dehydroxylation in NADH-treated lysates (Fig. 4A), likely by decreasing the free NADH pool. NADPH, which differs from NADH by a phosphate group on the 2'-OH group of the adenosine moiety, was unable to promote uroC dehydroxylation, indicating some specificity in the redox cofactors necessary for dehydroxylation. Aerobic incubation of crude lysates supplemented with NADH completely inhibited uroC dehydroxylation (Fig. 4B), indicating that the active enzyme complex requires a strictly anaerobic environment for dehydroxylation, as has been demonstrated for various metalloenzymes [44].

To confirm that *ucd* operon-encoded proteins were expressed in *E. bolteae* crude lysates, we performed untargeted proteomics and compared protein expression upon DMSO or uroC treatment. Indeed, all 3 proteins encoded by the *ucd* operon (UcdC, UcdF, and UcdO) were the most differentially expressed proteins in the uroC treatment group (Fig. 4C). In addition, proteins involved in MCD biosynthesis were also strongly increased upon uroC treatment (Fig. 4C,D, Supplementary Fig. 5A), pointing to the coordination between MCD biosynthesis and active UcdCFO oxidoreductase assembly. These multi-omics datasets implicate all three *ucdCFO* genes and MCD biosynthesis genes in the metabolism of uroC to uroA, as demonstrated by the strong positive correlation between transcript and protein differential expression (Fig. 4D).

To validate the function of the *E. bolteae ucd* operon, we attempted heterologous expression of *E. bolteae* UcdCFO in *E. coli*; however, all expression and activity assays were unsuccessful

despite the inclusion of *mocA* and *xdhC* genes involved in MCD maturation in our expression plasmids. This lack of activity likely resulted from the choice of heterologous host and from the complex assembly of active molybdoenzymes [43]. We therefore attempted to express UcdCFO in the phenol-degrading soil bacterium *Rhodococcus erythropolis* using a thiostrepton-inducible expression system [45] (pTipQC2-*ucdCFO*, Supplementary Fig. 6A,B), previously used to express the anaerobic *E. lenta* Cgr2 protein [44]. Despite the poor yield of soluble protein (Supplementary Fig. 6C,D), we were able to observe uroM6 and uroC dehydroxylation at the 9-position in crude lysates of *R. erythropolis* transformed with pTipQC2-*ucdCFO*, but not in the no insert control (pTipQC2) (Fig. 4E, Supplementary Fig. 6E-G), thus confirming that the *ucd* operon confers 9-hydroxy urolithin dehydroxylase activity.

To gain an understanding of the structural organization of proteins encoded by the *ucd* operon, we performed modeling using AlphaFold2 [46,47]. Structures of each protein encoded by the *E. bolteae ucd* operon (Fig. 4F) were superposed onto published X-ray crystal structures of xanthine dehydrogenase family enzymes with similar folds [48], yet from different taxonomic domains: *Afipia carboxidovorans* carbon monoxide dehydrogenase [49] and *Bos taurus* xanthine dehydrogenase [50]. The 3 proteins encoded by the *ucd* operon formed subunits in an oxidoreductase complex with a similar quaternary structure to the published crystal structures (Fig. 4G, Supplementary Fig. 7A,B). The predicted quaternary structure of the UcdCFO enzyme complex supported a complete electron transport chain whereby electrons would flow from reduced FAD to two 2Fe-2S clusters, then to the MCD cofactor, and finally to uroC as the terminal electron acceptor (Fig. 4H, Supplementary Fig. 7C,D). This model supports our findings in crude lysates whereby NADH serves as an electron donor to reduce UcdC-bound FAD (Fig. 4A). Using homology modeling, we further identified the putative uroC binding site in UcdO, which overlaps with the salicylic acid ligand in the *Bos taurus* xanthine dehydrogenase

structure (Supplementary Fig. 7E). This putative uroC binding site contains multiple tyrosine (Y375, Y538, Y624, Y632), tryptophan (W345), and phenylalanine (F458, F464) residues that could form π-π stacking interactions with uroC (Supplementary Fig. 7F), orienting it towards the molybdenum cofactor.

Figure 4. The UcdCFO complex enables anaerobic electron transport from NADH to uroC.

A) Quantification of urolithin peak concentrations in crude uroC-induced *Eb* lysates re-treated with DMSO or uroC (350 µM) and various coenzymes (n = 3 biological replicates). NADPH, nicotinamide adenine dinucleotide phosphate; NADH, nicotinamide adenine dinucleotide; FAD, flavin adenine dinucleotide. **B)** Quantification of urolithin peak concentrations in crude uroC-induced *Eb* lysates re-treated with DMSO or uroC (350 µM) and NADH in anaerobic or aerobic environments (n = 3 biological replicates). Data are represented as mean ± SEM for (A,B). **C)** Volcano plot of untargeted proteomics analysis on DMSO or uroC-treated *Eb* (n = 3 biological

replicates). Data points are colored according to their significance (Fisher's exact test with Benjamini-Hochberg correction for multiple comparisons). Grey, p-adj ≥ cutoff p-value (0.00048). Blue, p-adj < cutoff p-value (0.00048). **D)** Scatter plot showing the correlation between gene and protein expression ($\log_2$FC values) induced in uroC-treated *Eb* using the datasets in Fig. 2G and Fig. 5C, respectively. The non-parametric Spearman rank correlation test was used for statistical analysis. **E)** LC-MS extracted ion chromatograms (EIC) of uroC ([M-H]$^-$ = 243) and uroA ([M-H]$^-$ = 227) from a representative anaerobic uroC dehydroxylation assay using crude lysates of *R. erythropolis* harboring either pTipQC2 (no insert) or pTipQC2-*ucdCFO* plasmids. **F)** Domains of genes in the *ucd* operon based on InterPro annotations. **G)** Quaternary structure prediction of the proteins encoded by the *Eb ucd* operon. AlphaFold2 structures for each protein were superposed onto the X-ray crystal structure of PDB 1ZXI (carbon monoxide dehydrogenase from *Afipia carboxidovorans* OM5). **H)** Small molecule ligands from PDB 1ZXI in the superposed UcdCFO model form a complete electron transport chain from FAD to two 2Fe-2S clusters to a molybdopterin cytosine dinucleotide cofactor, which can then reduce uroC (terminal electron acceptor). Source data and statistical details are provided as a Source data file.

Disruption of the urolithin C catechol moiety rescues growth delay in iron-limited media

To gain an understanding of why *Enterocloster* spp. and *L. pacaense* metabolize 9-hydroxy urolithins, we performed growth experiments in different media conditions. When uroC was added prior to growth in rich medium containing hemin (mABB+H), a concentration-dependent increase in lag time was observed for all uroC-metabolizing bacteria (Fig. 5A,B). As catechols are common structural motifs in iron-binding siderophores [51], we hypothesized that uroC was delaying growth by altering iron availability in the growth medium *via* its catechol moiety. Incubation of *Enterocloster* spp. and *L. pacaense* in medium lacking added iron (mABB) exacerbated the growth delay by uroC (Fig. 5C, Supplementary Fig. 8A,B); however, this growth delay was partially rescued upon supplementation of different iron sources (hemin, Fe(II)SO$_4$, or Fe(III) pyrophosphate) (Fig. 5C, Supplementary Fig. 8C). To validate that iron chelation could extend the lag time of *Enterocloster* spp. and *L. pacaense*, we incubated all four uroC-metabolizers with 2,2'-bipyridyl (biP) in mABB media. As observed with uroC, biP delayed the growth of all tested bacteria, but supplementation of Fe(II)SO$_4$, or Fe(III) pyrophosphate could partially rescue growth delay (Supplementary Fig. 8C). Interestingly,

uroA, which lacks a catechol moiety, did not impact the growth of the tested bacteria in either mABB or mABB+H media compared to uroC (Fig. 5C, Supplementary Fig. 8A,B). To confirm that the catechol moiety of uroC was responsible for delaying growth, we synthesized a methylated analogue of uroC (8,9-di-O-methyl-uroC, Fig. 5D, Supplementary Fig. 9A-F), which is unable to bind iron [52], and tested its effect on growth in mABB. Like uroA, 8,9-di-O-methyl-uroC did not delay growth of uroC-metabolizing bacteria (Fig. 5E,F). These data demonstrate that both the catechol moiety of uroC and iron availability are essential to uroC-mediated lag phase extension.

Dehydroxylation of catechols by gut bacteria has been observed for diverse classes of compounds like neurotransmitters, therapeutic drugs, and diet-derived polyphenols [14]. Although catechol dehydroxylation can promote growth in some species [14], we hypothesized that dehydroxylation could be a mechanism used by gut bacteria to inactivate catechol-containing compounds that affect their fitness. To determine whether diverse catechols can delay growth, uroC-metabolizing bacteria and the dopamine-metabolizing *Eggerthella lenta* A2 were incubated with catechol-containing compounds and their dehydroxylated counterparts: uroC (uroA), entacapone, dopamine (m-tyramine), caffeic acid (m-coumaric acid) (Fig. 5D). Surprisingly, neither dopamine nor caffeic acid (and their dehydroxylated counterparts) delayed the growth of the tested bacteria (Fig. 5G). On the other hand, both uroC and the nitrocatechol-containing Parkinson's drug entacapone delayed the growth of *Enterocloster spp*. and *L. pacaense* but did not affect *E. lenta* A2 (Fig. 5G) [8]. Thus, catechol-containing compounds show differential effects on the growth of gut bacteria, depending on their structure. These results prompted us to investigate the effect of uroC on a more diverse panel of gut bacteria including *E. aldenensis, E. clostridioformis*, and *E. lavalensis*, which do not metabolize uroC, along with *Gordonibacter* spp., which produce uroC from dietary ellagic acid [35]. Treatment with uroC

381 delayed the growth of *E. aldenensis* , *E. clostridioformis,* and *E. lavalensis* to varying extents

382 (Supplementary Fig. 10A); however, there was no difference in growth between the DMSO-,

383 uroC-, and uroA-treated cultures of *Enterococcus faecium* and *Gordonibacter* spp.

384 (Supplementary Fig. 10B,C). Thus, all *Enterocloster* spp. tested showed sensitivity to uroC-

385 mediated lag time extension, while other bacteria were insensitive to its effects on growth.

Figure 5. The catechol moiety of uroC delays *Enterocloster* spp. growth in an iron-dependent manner.

A) Growth curves (optical density (OD) at 620 nm) of uroC-metabolizing *Enterocloster* spp. and *L. pacaense* treated with increasing concentrations of uroC in rich mABB+H media (7.7 μM hemin) (n = 3 biological replicates). Data are represented as mean ± SEM. **B)** Quantification of the difference in lag time compared to the DMSO control for growth curves in (A). Data are represented as mean ± SEM; lines were fitted using simple linear regression. **C)** Quantification of the difference in lag time of *Enterocloster* spp. and *L. pacaense* grown in mABB (no added iron) or mABB+H (7.7 μM hemin) compared to respective DMSO controls for growth curves in Supplementary Fig. 8A,B. Data are represented as mean ± SEM; repeated measures two-way ANOVA (matching by biological replicate) with Tukey's multiple comparisons test. Significant differences between treatments for individual bacteria are denoted by a different lowercase letter above each plot. **D)** Structures of tested catechols and their derivatives. **E)** Growth curves (OD at 620 nm) of uroC-metabolizing *Enterocloster* spp. and *L. pacaense* grown in mABB media and treated with DMSO (vehicle), 100 μM of uroC, uroA, or 8,9-di-O-methyl-uroC (n = 3 biological replicates). Data are represented as mean ± SEM. **F)** Quantification of the difference in lag time compared to the DMSO control for growth curves in (E). Data are represented as mean ± SEM; repeated measures two-way ANOVA (matching by biological replicate) with Tukey's multiple comparisons test. Significant differences between treatments for individual bacteria are denoted by a different lowercase letter above each plot. **G)** Growth curves (OD at 620 nm) of uroC-metabolizing *Enterocloster* spp., *L. pacaense*, and *E. lenta* A2 grown in mABB media and treated with DMSO (vehicle), 100 μM of uroC, uroA, entacapone, dopamine, m-tyramine, caffeic acid or m-coumaric acid. Data are represented as mean ± SEM.

UroC-metabolizing species and *ucd* genes are prevalent and correlate with uroC metabolism in human fecal samples.

We next wondered whether uroC-metabolizing *Enterocloster* spp. and their *ucd* operons were prevalent and active in human fecal samples. We first utilized uniformly processed metagenomic data from the curatedMetagenomicData R package [53]. After filtering for fecal samples (86 studies, n = 21,030 subjects), we counted the prevalence of at least one uroC-metabolizing species and at least one *ucd* gene homolog (Methods). The prevalence of both features was variable across studies (Fig. 6A for studies with >200 participants, Supplementary Fig. 11A for all studies). Combining all studies, the prevalence of at least one uroC-metabolizing species and at least one *ucdCFO* gene homolog was 9,343/21,030 (44.9%) and 4,356/21,030

(20.7%), respectively. *E. bolteae* was the most prevalent and abundant uroC-metabolizing

species detected in gut metagenomes (Supplementary Fig. 11B,C) and correlated strongly with

ucd abundance (Supplementary Fig. 11D). These findings suggest that uroC-metabolizing

Enterocloster spp. and *ucd* operon genes are prevalent in human fecal metagenomic samples

and reflect the variable urolithin metabolism profiles (metabotypes) in the general population

[25,54].

Next, we performed *ex vivo* metabolism assays to determine whether *Enterocloster* spp. could

metabolize uroC in the context of a complex community. Fecal slurries from 10 healthy

individuals were first profiled according to their uroC metabotypes (Fig. 6B) [24]. Individuals

clustered into metabotypes A (only uroA produced), B (uroA and isouroA/uroB), and 0 (no

terminal urolithin metabolites). While we observed metabotypes A and B in uroC-metabolizing

fecal slurries, all slurries produced some amount of uroA from uroC. Stools JL73, TR06, and

YE96 displayed variable metabolism patterns and did not metabolize uroC in some

experiments, likely reflecting differences in activity between aliquots of feces (Fig. 6B). We then

repeated metabolism assays using fecal slurries from all 10 healthy individuals and extracted

urolithins, DNA, and RNA from each culture. In this experiment, only 5/10 fecal slurries

metabolized uroC to uroA (Fig. 6C). We hypothesized that differences in metabotypes could be

explained by microbial composition. Therefore, long-read V1-V9 16S rRNA sequencing was

performed on fecal slurries. Both DMSO- and uroC-treated fecal slurries within individuals had

similar microbial compositions and diversity metrics (Supplementary Fig. 12A-D) but showed

differences in composition between individuals and metabolism status (Supplementary Fig.

12A,E). Surprisingly, all samples contained 16S rRNA sequences mapping to *E. bolteae*, and

many non-metabolizing fecal slurries contained *E. asparagiformis* (Fig. 6C, Supplementary Fig.

12B). We then assayed genomic DNA from treated fecal slurries for the presence of the *ucd*

operon by PCR and found that 10/10 individuals (19/20 conditions) yielded a detectable

amplicon of the expected size (~3.6 kb) (Supplementary Fig. 12F,G). These data indicate that

the prevalence of uroC-metabolizing *Enterocloster* spp. 16S rRNA and *ucd* operon genes does

not predict metabolism in fecal samples.

We then surmised that the *ucd* operon would be transcribed only in fecal slurries actively

metabolizing uroC. Using a gene-specific reverse primer that binds to *ucdO* (Fig. 2H), the full-

length *ucd* operon was reverse transcribed and amplified in RNA extracted from DMSO- and

uroC-treated fecal slurries. An amplicon (~3.6 kb) corresponding to the *ucd* operon was only

detected in uroC-metabolizing fecal slurries (Fig. 6D) when treated with uroC and entirely

absent from non-metabolizing slurries (Fig. 6E). This amplicon was absent in no reverse

transcriptase controls, indicating no gDNA contamination (Supplementary Fig. 12H,I). These

data demonstrate that *ucd* transcription correlates with uroC metabolism in complex fecal

communities and that *E. bolteae* is keystone species involved in urolithin A production.

Figure 6. The *ucd* operon is prevalent in metagenomes and actively transcribed in urolithin C-metabolizing human fecal samples.

A) Prevalence of *ucd* operon (at least one gene) and of a uroC-metabolizing species (at least one species) in fecal metagenomes from the CuratedMetagenomicData R package. Only studies with ≥200 participants are depicted. All 86 studies are available in Supplementary Fig. 11. **B)** Summary of urolithin concentrations in fecal slurries (n = 10 healthy donors) incubated with 100 µM uroC for 48 h. Data are represented as mean ± SEM (n = 3-6 experimental replicates). **C)** Summary of urolithin concentrations in fecal slurries (n = 10 healthy donors) incubated with 100 µM uroC for 48 h. Presence of uroC-metabolizing species and of the *ucd* operon is denoted above the graph if the bacterium or operon was detected in the uroC-treated fecal slurry (Supplementary Fig. 12B,F,G). Data are representative of 1 replicate where DNA and RNA was also extracted from fecal slurries. **D,E)** *ucd* gene-specific RT-PCR on fecal microbiota communities from 10 healthy donors using the primer set described in Fig. 2H. Samples are matched to the urolithin metabolism data in C) and 16S rRNA sequencing in Supplementary Fig. 12A-E. 1% agarose gel of amplicons derived from **D)** uroC-metabolizing and **E)** non-metabolizing fecal slurries. Bands corresponding to the *E. bolteae ucd* operon (~3.6 kb) are labeled with red arrows. The no template control (NTC) is the same for (D,E). See Supplementary Fig. 12H,I for the no reverse transcriptase control PCR reactions on the same samples. Source data are provided as a Source data file.

Discussion

We identified genes and proteins that are essential for the metabolism of urolithins by gut resident *Lachnospiraceae* through a combination of transcriptomics, comparative genomics, and untargeted proteomics. Our study reveals a novel multi-subunit molybdoenzyme (urolithin C dehydroxylase, Ucd) that catalyzes the dehydroxylation of 9-hydroxy urolithins including uroM6, uroC, and isouroA. Importantly, prevalence analysis in published data and *ex vivo* transcriptomics established *E. bolteae* as a keystone urolithin-metabolizing member of the gut microbiota.

Catechol dehydroxylases are widespread in gut resident *Eggerthella lenta* and *Gordonibacter* spp. [10,14,55]. These molybdoenzymes, which belong to the DMSO reductase superfamily [56], dehydroxylate substrates like catechol lignan (Cldh), dopamine (Dadh), DOPAC

(Dodh), hydrocaffeic acid (Hcdh), and caffeic acid (Cadh), which can promote growth by using

these substrates as alternative electron acceptors [14]. A recent survey of reductases in gut

bacteria established that most respiratory reductases contain N-terminal signal sequences and

are translocated across the cytoplasmic membrane, while non-respiratory reductases, which

lack signal sequences, remain in the cytoplasm [57]. The UcdCFO enzyme complex we found in

Enterocloster spp. differs from catechol dehydroxylases in *Eggerthellaceae* in important ways

as it does not require a catechol structural motif for activity, belongs to the xanthine oxidase

superfamily, and is composed of 3 subunits that each lack signal sequences. Based on the

absence of signal sequences and the cytoplasmic localization of xanthine dehydrogenases, the

ucd operon likely encodes for a non-respiratory reductase serving a different role than

previously characterized catechol dehydroxylases [57].

In rich media conditions, uroC, but not uroA, extended the lag phase of growth in both

uroC-metabolizing and non-metabolizing *Enterocloster* spp. This growth delay was not

observed for other taxa, suggesting that *Enterocloster* spp. are especially sensitive to uroC-

mediated iron chelation. In addition to the *ucd* operon, uroC-treated *E. bolteae* and *E.*

asparagiformis upregulated gene clusters related to efflux (MepA-like MATE family efflux

transporters) and iron/siderophore transport (FecCD-like ABC transporter). These responses

are analogous to antimicrobial resistance mechanisms raised against entacapone and other

non-antibiotic drugs [8]. This suggests that non-respiratory uroC dehydroxylation could serve as

an additional strategy that evolved in *E. asparagiformis*, *E. bolteae*, *E. citroniae*, and *L.*

pacaense to overcome catechol-mediated iron chelation.

While uroA is the most common terminal metabolite following ellagitannin consumption

in humans, its production varies widely [36]. Interestingly, the ability of a fecal sample to produce

uroA from uroC did not correlate with the presence of widespread uroC-metabolizing

Enterocloster spp. or a *ucd* gene homologue, likely owing to poor viability or dead bacteria in

fecal samples. However, the active transcription of the *ucd* operon correlated perfectly with

metabotypes. These findings further emphasize the importance of functional assays such as

transcriptomics and *ex vivo* metabolism to understand the metabolism of xenobiotics by the gut

microbiota.

By identifying the genetic basis for metabolism of uroC, we found a novel metabolizing

species that could not have been predicted based on phylogeny alone. Our data suggests that

ucd-containing *Enterocloster* spp., and the closely related *L. pacaense*, are the main drivers of

urolithin A production in the gut microbiota based on their prevalence in metagenomes and

activity in fecal samples. However, we cannot conclude that they are solely responsible for this

activity. Rare, strain-specific urolithin A production has been reported for *Bifidobacterium*

pseudocatenulatum INIA P815 [36], *Streptococcus thermophilus* FUA329 [58], and *Enterococcus*

faecium FUA027 [58], which may be a result of horizontal gene transfer since it is not shared by

other members of the taxa. Thus, further enzyme discovery efforts are necessary to understand

urolithin production in these bacteria.

In conclusion, our studies reveal the genetic and chemical basis for urolithin A production

by gut bacteria and broaden our understanding of the molecular mechanisms underlying

urolithin metabotypes in human populations. Since diet can modulate gut microbiota function

and host health, elucidating the xenobiotic metabolism genes encoded by gut bacteria will be

key to developing dietary interventions targeting the gut microbiota.

Supplementary Figures

Supplementary Figure 1. Ellagitannins are metabolized by gut bacteria.

Reaction scheme of dietary ellagitannin metabolism by the human gut microbiota. Larger ellagitannin structures are hydrolyzed during gut transit, releasing hexahydroxydiphenic acid, which spontaneously lactonizes into ellagic acid. Once in the gut lumen, members of the *Gordonibacter* spp. and *Ellagibacter isourolithinifaciens* can decarboxylate ellagic acid, forming urolithin M-5. The resulting urolithin M-5 can be further dehydroxylated (at the 4,10- or 4,8,10- positions) to uroC or isourolithin A by *Gordonibacter* spp. or *Ellagibacter isourolithinifaciens*, respectively. Compounds colored in light gray are urolithin metabolites that are rarely observed during *ex vivo* metabolism assays on ellagitannins. Once uroC or isouroA are produced, *Enterocloster* spp. can further dehydroxylate the 9-position, yielding uroA or uroB, respectively. The cartoon was generated in BioRender.

Supplementary Figure 2. *Enterocloster* **spp. whole genome and proteome phylogenetic**

trees.

A) Whole genome phylogenetic tree of *Enterocloster* spp. The tree was inferred with FastME 2.1.6.1 from Genome BLAST Distance Phylogeny (GBDP) distances calculated from genome sequences. The branch lengths are scaled in terms of GBDP distance formula d5. The numbers above branches are GBDP pseudo-bootstrap support values > 60 % from 100 replications, with an average branch support of 36.8 %. The tree was rooted at the midpoint. **B)** Whole proteome phylogenetic tree of *Enterocloster* spp. The tree was inferred with FastME 2.1.6.1 from whole-proteome-based GBDP distances. The branch lengths are scaled via GBDP distance formula d5. Branch values are GBDP pseudo-bootstrap support values > 60 % from 100 replications, with an average branch support of 100.0 %. The tree was rooted at the midpoint. Source data are provided as a Source data file.

Supplementary Figure 3. *E. bolteae* dehydroxylates urolithins at the 9-position.

A) Chemical structures of urolithin M-6, urolithin C, and isourolithin A along with their dehydroxylated counterparts urolithin M7, urolithin A, and urolithin B. **B)** Quantification of extracted ion chromatogram (EIC) peak areas from *Eb* cultures sampled after 24 h of growth with 100 µM uroM6 (n = 3 biological replicates) with representative extracted ion chromatograms (EIC) to the right (from one representative biological replicate). The same scale was used for each chromatogram. **C)** Quantification of urolithin peak areas from *Eb* cultures sampled after 24 h of growth with 100 µM uroC (n = 3 biological replicates) with representative chromatograms (λ = 305 nm) to the right (from one biological replicate). The same scale was used for each chromatogram. **D)** Quantification of urolithin peak areas from *Eb* cultures sampled after 24 h of growth with 100 µM isouroA (n = 3 biological replicates) with representative chromatograms (λ = 305 nm) to the right (from one biological replicate). The

same scale was used for each chromatogram. **E,F)** RT-qPCR expression of the *Eb ucdO* gene.
Growing *Eb* cultures were treated with DMSO, uroM6, uroC, or isouroA (100 µM) for 2 h before
RNA isolation and reverse transcription (n = 3 biological replicates). **E)** Differential *Eb ucdO*
gene expression comparing DMSO, uroM6, uroC, and isouroA is displayed as target-specific
ΔCt ($C_{t\ \text{MoO Gene}}$ - $C_{t\ \text{dnaK Reference Gene}}$) values. Data are presented as individual ΔCt values with
lines connecting paired biological replicates (from the same pre-spike culture); repeated-
measures one-way ANOVA with Dunnett's multiple comparisons test; **, $p < 0.01$; ***, $p <$
0.001. **F)** Gene expression profile of the *Eb ucdO* gene in different urolithin treatment groups
displayed as $\log_2$FC (equivalent to $-\Delta\Delta C_t$, where $\Delta\Delta C_t = \Delta C_{t\ \text{urolithin}} - \Delta C_{t\ \text{DMSO}}$). Data are
presented as individual $\log_2$FC values with lines connecting paired biological replicates;
repeated-measures one-way ANOVA with Tukey's multiple comparisons test. **G)** Quantification
of extracted ion chromatogram (EIC) peak areas in DMSO- or uroC-treated *Eb* cell
suspensions. Cell suspensions were prepared from *Eb* cells grown with either DMSO or 50 µM
uroC. The cells were washed and resuspended in PBS to halt the production of new enzymes,
then treated with 100 µM uroM6 (n = 3 biological replicates). **H)** Quantification of urolithin
concentrations in DMSO- or uroC-treated *Eb* cell suspensions (n = 3 biological replicates). For
B-D and G-H, data are represented as mean ± SEM. ND, not detected; ns, not significant; FC,
fold change; FDR, false discovery rate. Source data and statistical details are provided as a
Source data file.

Note: Enterocloster clostridioformis AM07-19 (NZ_QVEN01000003.1) is mis-assigned
(closest match type strain is *E. bolteae*).
See https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_003433765.1/

Supplementary Figure 4. BLASTn searches using the *E. bolteae ucd* operon genomic

sequence identifies homologues in gut bacteria.

NBCI Multiple Sequence Aligner viewer hits for BLASTn searches using the *E. bolteae* DSM
15670 *ucd* operon nucleotide sequence as a query against the NCBI refseq_genomes
database (limited to Bacteria). Vertical lines in the sequence alignment represent nucleotide
differences (show differences option) and insertions relative to the query sequence. Source
data are provided as a Source data file.

A

B

Protein annotation	Role(s)	*Ea* NCBI Accession	*Eb* NCBI Accession	#AA in *Eb*	Domain(s) [positions]
GTP 3',8-cyclase	MoaA	WP_007712811.1	WP_002565229.1	355	Radical SAM core [7-238]
Cyclic pyranopterin monophosphate synthase	MoaC	WP_007712808.1	WP_002565230.1	164	Molbdopterin cofactor biosynthesis C (MoaC) [14-149]
MOSC domain-containing protein	MogA	WP_051409812.1	WP_002565228.1	322	Molbdenum cofactor sulfurase C-terminal (MOSC) [28-153]
Molybdopterin-binding protein / Molybdopterin molybdenumtransferase	MoeA	WP_007712805.1	WP_002565231.1	352	MoaB/Mog [177-317]
NTP-transferase domain-containing protein / LysR family transcriptional regulator	MocA/ModE	WP_007713105.1	WP_002565224.1	313	MobA-like NTP transferase [5-160] Helix-turn-helix lysr-type [222-277]
ATP-binding cassette domain-containing protein	ModC	WP_117778241.1	WP_002565225.1	382	ABC transporter [2-259]
Molybdate ABC transporter permease subunit	ModB	WP_007712823.1	WP_002565226.1	223	ABC transmembrane type-1 [6-215]
Molybdate ABC transporter substrate-binding protein	ModA	WP_007712821.1	WP_002565227.1	294	Signal [1-25] Molybdate ABC transporter substrate-binding protein [26-294]
XdhC family protein	XdhC	WP_007713106.1	WP_002565223.1	343	XdhC- CoxI [12-70] XdhC Rossmann [189-331]

C

Supplementary Figure 5. Urolithin C treatment upregulates molybdopterin cytosine

dinucleotide cofactor biosynthetic gene clusters.

A) Molybdopterin cytosine dinucleotide (MCD) cofactor biosynthetic pathway. The $\log_2$FC
values of MCD cofactor biosynthetic genes upregulated by uroC in both transcriptomics (T) and
proteomics (P) datasets (when available) are provided to the left of the figure for both *Ea* and
Eb. **B)** Table of molybdenum cofactor biosynthetic genes found in the genomes of *Ea* and *Eb*.
Annotations are based on NCBI (GTF files) and UniProt gene/protein names. Roles are
assigned based on required proteins for molybdenum cofactors in the xanthine
oxidase/dehydrogenase family of enzymes. Accessions, primary sequence length, and
annotated domains (with positions within the primary sequence) are also provided. **C)** Genomic
organization of the MCD cofactor biosynthetic genes for *Ea* and *Eb* (generated from the NCBI
Sequence Viewer). Source data are provided as a Source data file.

Supplementary Figure 6. Heterologous expression of *E. bolteae ucdCFO* genes in *R. erythropolis*.

A) Map of the *E. coli – R. erythropolis* pTipQC2-*ucdCFO* shuttle plasmid generated in Benchling. **B)** Genomic organization of the wild-type *E. bolteae ucd* operon with intergenic

regions highlighted. Shine-Dalgarno consensus sequences (bold) are denoted between
translational stop and start (underlined) sites for *ucdC* and *ucdF* (Intergenic Region I) and for
ucdF and *ucdO* (Intergenic Region II). **C,D)** SDS-PAGE gels (10% bis-tris) stained with colloidal
Coomassie dye of the insoluble (C) and soluble (D) fractions from thiostrepton-induced (1
µg/mL) *R. erythropolis* harboring pTipQC2 no insert or pTipQC2-*ucdCFO* plasmids (n = 3
biological replicates). UcdCFO complex proteins are labeled on the right side of each gel image.
E,F) Quantification of extracted ion chromatogram (EIC) peak areas from crude lysates of
thiostrepton-induced *R. erythropolis* harboring pTipQC2 no insert or pTipQC2-*ucdCFO*
plasmids (n = 3 biological replicates). Crude lysates were incubated anaerobically with 2 mM
NADH and 357 µM uroC (E) or 357 µM uroM6 (F) for 72 h before extraction and analysis by
LC-MS. Data are represented as mean ± SEM. **G)** LC-MS extracted ion chromatograms (EIC)
of uroM6 ($[M-H]^- = 259$) and uroM7 ($[M-H]^- = 243$) from a representative anaerobic uroC
dehydroxylation assay using crude lysates of *R. erythropolis* harboring either pTipQC2 (no
insert) or pTipQC2-*ucdCFO* plasmids. Source data are provided as a Source data file.

Supplementary Figure 7. The AlphaFold2 model of the *E. bolteae* UcdCFO enzyme complex has a similar quaternary structure to xanthine dehydrogenase superfamily crystal structures.

A,B) Structural superposition of AlphaFold2-predicted *Eb* Ucd proteins (right) onto the X-ray crystal structures of **A)** PDB 1ZXI (carbon monoxide dehydrogenase from *Oligotropha carboxidovorans* OM5) X-ray crystal structure (left) and **B)** PDB 3UNI (bovine milk xanthine

dehydrogenase with NADH bound). PDB 1ZXI is colored in shades of green (according to chain
ID), PDB 3UNI is colored in gold (single chain A shown), and the *Eb* Ucd enzyme complex is
colored according to its per-residue confidence score as indicated in the legend. The various
ligands (cofactors, coenzymes, ions, small molecules) of PDB 1ZXI and PDB 3UNI are included
in the respective *Eb* Ucd enzyme models. **C,D)** Cofactors from X-ray crystal structures **C)** PDB
1ZXI and **D)** PDB 3UNI modeled into the AlphaFold2 *Eb* Ucd enzyme complex, showing a
complete electron transport chain from a bound FAD molecule to a molybdopterin cofactor *via*
two 2Fe-2S clusters. **E)** Xanthine dehydrogenase active site from PDB 3UNI. Side chains within
8 Å of the salicylic acid ligand are colored in gold. Residues important for substrate (purine)
binding and catalysis are labeled with their one letter amino acid code and sequence position.
Position E1261* is catalytically important (acts as a general base) and is conserved in
XDH/xanthine oxidase enzymes [59]. **F)** *Eb* Ucd enzyme complex active site modeling. Side
chains within 8 Å of the salicylic acid ligand from PDB 3UNI are colored on the superposed *Eb*
Ucd enzyme complex according to their per-residue confidence score as indicated in the
legend. Residues surrounding the predicted active site are labeled with their one letter amino
acid code and sequence position (in the *Eb* MoO protein). The predicted urolithin binding site
is depicted by the surface (mesh) created by the active site residues. The surface was rendered
using the cavities and pockets only (culled) setting with a cavity detection cutoff of 5 solvent
radii in PyMOL. Source data are provided as a Source data file.

Supplementary Figure 8. Iron supplementation rescues lag time extension by uroC and

2,2'-bipyridyl.

A,B) Growth curves (optical density (OD) at 620 nm) of uroC-metabolizing *Enterocloster* spp. and *L. pacaense* treated with H_2O, DMSO (vehicle), or 100 µM of uroC or uroA in mABB medium (lacking added iron) (A) or mABB+H (B) medium (containing 7.7 µM hemin) (n = 3 biological replicates). Data are represented as mean ± SEM. **C)** Growth curves (optical density (OD) at 620 nm) of uroC-metabolizing *Enterocloster* spp. and *L. pacaense* treated with DMSO (vehicle), 100 µM of uroC, or 2,2'-bipyridyl (biP) in mABB media (lacking added iron) supplemented with solutions containing no iron (H_2O), 7.7 µM Fe^{2+} (Fe(II)SO_4), or 7.7 µM Fe^{3+} (Fe(III) pyrophosphate) (n = 3 biological replicates). Data are represented as means without error bars for clarity. Source data are provided as a Source data file.

 Supplementary Figure 9. Synthesis and characterization of 8,9-di-O-methyl-urolithin C

A) Reaction scheme for the synthesis of 8,9-di-O-methy-urolithin C. **B)** Aligned reversed-phase
(C18) HPLC chromatograms (λ = 305 nm) of 10 µL injections of the following solutions: MeOH
blank, urolithin C 100 µM, 8,9-di-O-methy-urolithin C 100 µM. **C,D)** Negative ESI-MS spectra
of urolithin C (C) and 8,9-di-O-methy-urolithin C (D) following chromatographic separation (B).
E) ^{1}H NMR spectrum (600 MHz, (CD$_3$)$_2$SO)) of 8,9-di-O-methy-urolithin C. **F)** COSY NMR
spectrum (600 MHz, (CD$_3$)$_2$SO)) of 8,9-di-O-methy-urolithin C. The grey diagonal line denotes
self correlation between protons. The right panel corresponds to the area in the red box.
Coupling between protons is shown on the 8,9-di-O-methy-urolithin C using bold lines and bold
arrows (for long range coupling).

Supplementary Figure 10. Urolithin C differentially affects the growth of gut bacteria *in*

***vitro*.**

A,B,C) Growth curves (optical density (OD) at 620 nm) of non-uroC metabolizing *Enterocloster*
spp. (A), *Enterococcus faecium* (B) and *Gordonibacter* spp. (C) treated with DMSO (vehicle),
or 100 µM of uroC or uroA in mABB medium (n = 3 biological replicates for *Enterocloster* spp.
and *E. faecium*, and n = 4 biological replicates for *Gordonibacter* spp.). Data are represented
as mean ± SEM.

Supplementary Figure 11. Urolithin C-metabolizing species and genes are prevalent and correlate with *ucdO* gene abundance in human gut metagenomes.

A) Related to Fig. 6A. Prevalence of at least one *ucd* operon gene (blue bars) and uroC-metabolizing species (white bars) in fecal metagenomes across all 86 studies (in reverse alphabetical order). The number of participants in each study are represented to the right of the

prevalence plot as a heatmap. Details on the study populations (from the
curatedMetagenomicData R package) can be found in the Source Data file. **B)** Prevalence of
uroC-metabolizing species in human fecal metagenomes from the curatedMetagenomicData R
package. Data are reported for 86 studies (N=21,030 individuals) and are colored according to
the species. **C)** Violin plot of the $\log_{10}$(relative abundance) of uroC-metabolizing species in
human fecal metagenomes. The solid horizontal line corresponds to the median and the dashed
white lines correspond to the first and third quartiles. The percentage of zeroes are denoted
below the plotted distributions. Differences between groups were determined using the Kruskal-
Wallis test on untransformed relative abundance values. Significant differences between
groups are denoted by a different lowercase letter above each plot. **D)** Correlation between the
ucdO gene abundance in reads per kilobase per million mapped reads (RPKM) and the relative
abundance of each uroC-metabolizing species in fecal metagenomes. Both values are
illustrated on a $\log_{10}$ scale. Spearman rho (ρ) values are denoted above the scatter plots. All
correlations were significant, P < 0.0001. Source data are provided as a Source data file.

Supplementary Figure 12. Urolithin C-metabolizing species and the *ucd* operon are prevalent in fecal slurries, but only *ucd* transcription correlates with urolithin C metabolism.

A) Stacked bar plot of bacterial percent relative abundance (based on V1-V9 16S rRNA gene sequencing) in DMSO- or uroC-treated fecal slurries from 10 healthy donors (from one experimental replicate). Bars are colored according to the phylum (bold) and order. **B)** Heatmap of the percent relative abundance of uroC-metabolizing *Enterocloster* spp. in A). **C)** Alpha diversity plots between DMSO- and uroC-treated fecal slurries according to Chao1, Shannon, and Simpson diversity metrics. Lines between data points connect paired biological replicates; Wilcoxon test; ns, not significant. **D,E)** Principal coordinate analyses of dissimilarities between 16S rRNA compositions based on the Bray-Curtis and weighted UniFrac distance methods. Data points are colored according to the treatment used and the metabolism status of the fecal slurry; PERMANOVA test according to the treatment (D) for all fecal slurries or uroC metabolism status (E) for uroC-treated slurries. **F,G)** PCR with *ucd*-specific primers on gDNA extracted from an *E. bolteae* isolate and fecal microbiota communities from 10 healthy donors (from one experimental replicate). 1% agarose gel of amplicons derived from uroC-metabolizing (F) and non-metabolizing (G) fecal slurries. Samples are matched to the urolithin metabolism data in Fig. 6C and 16S rRNA sequencing in A) and B). The no template control (NTC) and *E. bolteae* positive control are the same for both gels. **H,I)** 1% agarose gel of amplicons from Fig. 6D,E including the no reverse transcriptase (-RT) in uroC-metabolizing (H) and non-metabolizing (I) fecal slurries. The NTC is derived from the final lane of the gel in H). Source data are provided as a Source data file.

Supplementary Sequence 1. cDNA sequence of *E. bolteae ucdCFO* transcript, 3605 bp

(band in Fig. 2I)

>Eb_ucd_RT-PCR_band coverage: 4.94e+03x

```
AAGCACAGCCAGGGTTATGGCAGGCGCAACGGATCTGATTCCGCCAATGAAAGACAAGGTTATATCACCAGAGTATATCATTG
ACCTTAAGAAAATTCCAGGTTTGGATTATCTGGAATACGATGACAGGGAAGGTTTAAAAATAGGGGCGCTGACAACACTGCGT
ACCATAGAGACATCTCCTCTGGTTAAAGAAAAGAATCCGGCAGTGGCTCATGCCGCCAAGGTAGTGGCATCCACACAAATCAG
GACAAAGGGCACCATGGCAGGCAATATCTGCAATGCGTCCCCATCCTGCGATACAGCCCCCAACCTACTGGCTCAGGGCGCTA
AGATATTGGTACAGGGTCCCAACAAGGACCGGGTCATTCAAATCGAAGACTTTTTCCTGGGTGTCAAAAAGACTTCCCTGGAG
CCGGGAGAAATCGTGACAGGCATTGTGATTCCGCCGCTGGCTGAGAATGAGAGGGCTGCATACATAAAACACGCGGTCAGAAA
GGCGATGGACCTGGCCATCATCGGCGTCGCAGTCAAGATTAAGGTGGAAGACGGCGTCTGCACAGACGCCAGGATTGCCCTGG
GCGCGGTTGCGGCCACGCCGGTACGGGCACCCGGGGCGGAGGAAGCCCTGATTGGCAAAGAACTTACCGATGAAGTGATTGTG
AAGGCATCGGAGGAAGCAATGAATTCCTGTCATCCAATCTCTGATATCCGCGCATCGGCCGAATACAGGAAGGATATGATCCG
CGTGTTTACAAAACGTGCCATTAAGCAGGCCATGGAATGCTATAACTAGGAGGGAGACAGGAATGAAACATAGTATAAGTTTA
ATCATCAACGGAGATCCGGTTGATGCTATTGTGAAGGATAACCTGACGCTTCTGGATTTTCTTCGTGACCAGTTGTTTCTGAC
AGGCACGAAAAAGGGGTGTGAGGAAGGGGAGTGTGGCGCCTGTACCGTGATGCTTGACGGTAAGCCGGTTAACTCCTGCTGTA
CCCTGGCCGTGGAGTGCGACGGCCATGAGATCATAACCGTGGAGGGAATTGCCAGGGAGGGGATGCTTCATCCCATTCAGAAG
CAGTTTATAGAGAAGTGGGCCATGCAGTGCGGCTACTGTACTCCGGGTATGATTATGTCCGCCAAAGCTCTTCTGGATGTGAA
TAAACATCCTACGGAACTGGAAATCCGGGAAGCCATTGAGGGCAATCTGTGCCGCTGCACCGGTTATGCAAAGATTGTGGAGG
CAATCCAGGCAGCCGCTGCGCAGATGAACTGGGAGGAGGAAGCAAAGAATGCATAAAGACTGTGACAAACATTATTTTAAAAA
ACCGGAATTTTACCGTCTGACCGGCGAGAACAATTATGTCAGGATTGACGCTGAAGACAAGGTAACAGGACACGGCCAGTATG
TGGGTGACATCATGTTTCCGGATATGCTTACCGGAAAAATGGTCAGAAGCCCTTATGCATCTGCAAAAATCCTGTCCATTGAC
ACCAGCGAGGCAGAGAAGCTTCCAGGTGTAAAGTGCATCCTCACTGCCAGGGACTTTGAGTGGAAGTCCCTGGTGGGAAATGG
AGAATTTGCAGCTGAGTTCGCGGACAAGGAAGTATTGTGTTCTGAGAAGGTGCGCCAGGTTGGTGACGATGTGGCAGCCGTGG
CAGCCGTGGATGAGGAAACCGCCCAGCGGGCCGCGGATCTGATTAAGGTAGAGTACCAGGTGCTTCCGGGGGTGTTTGACCCC
TTTGAGGCAATGGAGGAGAATGCCCCCGAGGTAAACTGGGAGGGCAAAGGCATACACAATATCGGCATGCAGTCCGTGATGAA
GGCGGGCACGGATATTGATGAGGAATTTGACCGCGCATCCTATGTGCAGCACAGGGATTACAAGACCCACCGTATGGTACATG
CGGCCATGGAGCCCCACGGTGCGGTGGCTACCTACAGGAATGGGACCTACACCATCTGGATGTCCACCCAGATGTCCTTTGTG
GACCAGTTCTGGTATGCCCGCTGCCTTGGCGTTGGAGAGAACCAGGTGCGGGTAATCAAGCCTCTTGTGGGCGGCGGTTTCGG
CGGCAAGCTGGATTCCTATTCCTTTGGCCTTTGCGCTGCCAAGATGGCGGAGATGACCGGACGCCCGGTACGCATGATTCTGT
CCCGCGAAGAAGTATTCCAGACCACGCGCAACCGCCACCCCATTTACATGCATATTGACACTGCCTTTGGGACAGACGGTAAG
CTGCTGGCAAAGAAATGCTACCATGTGCTGGACGGCGGGCCTTACGGCGGGTCCGGTGTTGCGGCCTGCGCCCAGTCCACATT
GTGGGCCAACTTCCCTTACAAGATGAATTCCGTGGATTTCCTGGCAAGGCGTGTGTATACCAATAACCCTTCTGCAGGAGCCA
TGAGAGGATATACGGCATGCCAGGTGCATTTTGCCCATGACCTGAACATGCAGTTCGCGGCTGACCAGATGGGCATTGACCCG
GTTGAGTTCCGCAAGATAAGCGCGGCAGACCCCGGATACGTGGCTCCGGCCGGACTGGCAATCACCAGCTGCGCATACAAGGA
GACACTGGATACGGCTGCAAAGGAAATTGGCTGGTATGAGAAGAAGGATAAGCTGAAAAAGGGAGAGGGCATTGGTTTTGCGG
GCACCGGCTTTGTATCGGGCACCGGATTCGCGGTGCTGGAAGCGCCCAACCAGAGCTCTGCGTGCGTGACCCTGCGCATGAAC
AAGCGGGGCATGGCCACCCTGTACATCGGCTCCCACGACATAGGACAGGGTTCCGATACGGTTATGACGGCCATTGTGGCCGA
GGAACTGGGGCTTCCCATGGATATGGTCAAGACCTTTATGTCCGACACCTTCCTGACGCCCTGGGACTCCGGCTCTTACGGCA
GCCGTGTTACCTTCCTGGCCGGCAATGCGGCCCGCCGCGCTGCCGTGGATGCAAAACGCCAGCTGTTTGAGGTCATTGCGCCT
ATGTGGGGGGTGATGCCGGAGACATTAGAGTGTCTGGACGGAAAGGTAATCAGCAAGGAGAAGGCAGAGTATCAGATGTCCAT
TGGAGACGCCATGTTCAAGTATATGACGGTCAAGGGCGGCGATGAACTGATTGGCGTGGGATCCTATTACCACCGTACCGACA
ATTCACAGTATAACGGCAACAATACCACCAACTACGCGCCTGCATACAGCTTCTCCACCGGAGCAGCCCATCTGACTGTGGAC
GAGGAGACCGGCGTCCTGGATATTGATGAATTTGTATTTGCCCATGACTGCGGCCGCGCACTGAATAAGAGGGCCGTGGAAGG
CCAGCTGGAGGGATCCATCGGCATGGGCCTGGGCTATGCTGTCTATGAGCACAATGTAACCAGGGAAGGAAAGATTCTCAATC
CCAACTTCCGCGATTACCGGCTGCCCACTGCCCTTGATATGCCGAAGATGCGCACCTTTTACGACTTTACACCGGACGAGGAA
GGACCTCTGGGCGCAAAGGAAGCTGGCGAGGGCTCCGCAGCACCTGTGGCGCCTGCCATTGCCAATGCGGTCAACATGGCAAC
CGGCGTGTACTTCACAGAGCTGCCTCTGGACCCGGA
```

Supplementary Sequence 2. pTipQC2-*ucdCFO*

>pTipQC2-ucdCFO coverage: 717x

```
AAAAAAACCACCGCTACCAGCGGTGGTTTGTTTGCCGGATCAAGAGCTACCAACTCTTTTTCCGAAGGTAACTGGCTTCAGCA
GAGCGCAGATACCAAATACTGTTCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTACATAC
CTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTT
ACCGGATAAGGCGCAGCGGTCGGGCTGAACGGGGGGTTCGTGCACACAGCCCAGCTTGGAGCGAACGACCTACACCGAACTGA
GATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTC
GGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCTGTCGGGTTTCGCCACCTCTGACT
TGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGCGGCCTTTTTACGGTTCCTGG
CCTTTTGCTGGCCTTTTGCTCACATGTTCTTTCCTGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGA
GCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGCGGAAGAGCGCCCAATACGCAAACC
GCCTCTCCCCGCGCGTTGGCCGATTCATTAATGCAGCTGGCACGACTAGAGTCCCGCTGAGGCGGCGTAGCAGGTCAGCCGCC
CCAGCGGTGGTCACCAACCGGGGTGGAACGGCGCCGGTATCGGGTGTGTCCGTGGCGCTCATTCCAACCTCCGTGTGTTTGTG
CAGGTTTCGCGTGTTGCAGTCCCTCGCACCGGCACCCGCAGCGAGGGGCTCACGGGTGCCGGTGGGTCGACTAGTTCAGTGAT
GGTGATGGTGATGctcgagCTGCCT**TTATTTTGAATTCCTGTCATCCTTCATGCCGTGCAGCGCTCTCCAGATATGCTCCGGG**
**TCCAGAGGCAGCTCTGTGAAGTACACGCCGGTTGCCATGTTGACCGCATTGGCAATGGCAGGCGCCACAGGTGCTGCGGAGCC**
**CTCGCCAGCTTCCTTTGCGCCCAGAGGTCCTTCCTCGTCCGGTGTAAAGTCGTAAAAGGTGCGCATCTTCGGCATATCAAGGG**
**CAGTGGGCAGCCGGTAATCGCGGAAGTTGGGATTGAGAATCTTTCCTTCCCTGGTTACATTGTGCTCATAGACAGCATAGCCC**
**AGGCCCATGCCGATGGATCCCTCCAGCTGGCCTTCCACGGCCCTCTTATTCAGTGCGCGGCCGCAGTCATGGGCAAATACAAA**
**TTCATCAATATCCAGGACGCCGGTCTCCTCGTCCACAGTCAGATGGGCTGCTCCGGTGGAGAAGCTGTATGCAGGCGCGTAGT**
**TGGTGGTATTGTTGCCGTTATACTGTGAATTGTCGGTACGGTGGTAATAGGATCCCACGCCAATCAGTTCATCGCCGCCCTTG**
**ACCGTCATATACTTGAACATGGCGTCTCCAATGGACATCTGATACTCTGCCTTCTCCTTGCTGATTACCTTTCCGTCCAGACA**
**CTCTAATGTCTCCGGCATCACCCCCCACATAGGCGCAATGACCTCAAACAGCTGGCGTTTTGCATCCACGGCAGCGCGGCGGG**
**CCGCATTGCCGGCCAGGAAGGTAACACGGCTGCCGTAAGAGCCGGAGTCCCAGGGCGTCAGGAAGGTGTCGGACATAAAGGTC**
**TTGACCATATCCATGGGAAGCCCCAGTTCCTCGGCCACAATGGCCGTCATAACCGTATCGGAACCCTGTCCTATGTCGTGGGA**
**GCCGATGTACAGGGTGGCCATGCCCCGCTTGTTCATGCGCAGGGTCACGCACGCAGAGCTCTGGTTGGGCGCTTCCAGCACCG**
**CGAATCCGGTGCCCGATACAAAGCCGGTGCCCGCAAAACCAATGCCCTCTCCCTTTTTCAGCTTATCCTTCTTCTCATACCAG**
**CCAATTTCCTTTGCAGCCGTATCCAGTGTCTCCTTGTATGCGCAGCTGGTGATTGCCAGTCCGGCCGGAGCCACGTATCCGGG**
**GTCTGCCGCGCTTATCTTGCGGAACTCAACCGGGTCAATGCCCATCTGGTCAGCCGCGAACTGCATGTTCAGGTCATGGGCAA**
**AATGCACCTGGCATGCCGTATATCCTCTCATGGCTCCTGCAGAAGGGTTATTGGTATACACACGCCTTGCCAGGAAATCCACG**
**GAATTCATCTTGTAAGGGAAGTTGGCCCACAATGTGGACTGGGCGCAGGCCGCAACACCGGACCCGCCGTAAGGCCCGCCGTC**
**CAGCACATGGTAGCATTTCTTTGCCAGCAGCTTACCGTCTGTCCCAAAGGCAGTGTCAATATGCATGTAAATGGGGTGGCGGT**
**TGCGCGTGGTCTGGAATACTTCTTCGCGGGACAGAATCATGCGTACCGGGCGTCCGGTCATCTCCGCCATCTTGGCAGCGCAA**
**AGGCCAAAGGAATAGGAATCCAGCTTGCCGCCGAAACCGCCGCCCACAAGAGGCTTGATTACCCGCACCTGGTTCTCTCCAAC**
**GCCAAGGCAGCGGGCATACCAGAACTGGTCCACAAAGGACATCTGGGTGGACATCCAGATGGTGTAGGTCCCATTCCTGTAGG**
**TAGCCACCGCACCGTGGGGCTCCATGGCCGCATGTACCATACGGTGGGTCTTGTAATCCCTGTGCTGCACATAGGATGCGCGG**
**TCAAATTCCTCATCAATATCCGTGCCCGCCTTCATCACGGACTGCATGCCGATATTGTGTATGCCTTTGCCCTCCCAGTTTAC**
**CTCGGGGGCATTCTCCTCCATTGCCTCAAAGGGGTCAAACACCCCCGGAAGCACCTGGTACTCTACCTTAATCAGATCCGCGG**
**CCCGCTGGGCGGTTTCCTCATCCACGGCTGCCACGGCTGCCACATCGTCACCAACCTGGCGCACCTTCTCAGAACACAATACT**
**TCCTTGTCCGCGAACTCAGCTGCAAATTCTCCATTTCCCACCAGGGACTTCCACTCAAAGTCCCTGGCAGTGAGGATGCACTT**
**TACACCTGGAAGCTTCTCTGCCTCGCTGGTGTCAATGGACAGGATTTTTGCAGATGCATAAGGGCTTCTGACCATTTTTCCGG**
**TAAGCATATCCGGAAACATGATGTCACCCACATACTGGCCGTGTCCTGTTACCTTGTCTTCAGCGTCAATCCTGACATAATTG**
**TTCTCGCCGGTCAGACGGTAAAATTCCGGTTTTTTAAAATAATGTTTGTCACAGTCTTTATGCATTCTTTGCTTCCTCCTCCC**
**AGTTCATCTGCGCAGCGGCTGCCTGGATTGCCTCCACAATCTTTGCATAACCGGTGCAGCGGCACAGATTGCCCTCAATGGCT**
**TCCCGGATTTCCAGTTCCGTAGGATGTTTATTCACATCCAGAAGAGCTTTGGCGGACATAATCATACCCGGAGTACAGTAGCC**
**GCACTGCATGGCCCACTTCTCTATAAACTGCTTCTGAATGGGATGAAGCATCCCCTCCCTGGCAATTCCCTCCACGGTTATGA**
**TCTCATGGCCGTCGCACTCCACGGCCAGGGTACAGCAGGAGTTAACCGGCTTACCGTCAAGCATCACGGTACAGGCGCCACAC**
**TCCCCTTCCTCACACCCCTTTTTCGTGCCTGTCAGAAACAACTGGTCACGAAGAAAATCCAGAAGCGTCAGGTTATCCTTCAC**
**AATAGCATCAACCGGATCTCCGTTGATGATTAAACTTATACTATGTTTCATTCCTGTCTCCCTCCTAGTTATAGCATTCCATG**
**GCCTGCTTAATGGCACGTTTTGTAAACACGCGGATCATATCCTTCCTGTATTCGGCCGATGCGCGGATATCAGAGATTGGATG**
**ACAGGAATTCATTGCTTCCTCCGATGCCTTCACAATCACTTCATCGGTAAGTTCTTTGCCAATCAGGGCTTCCTCCGCCCCGG**
**GTGCCCGTACCGGCGTGGCCGCAACCGCGCCCAGGGCAATCCTGGCGTCTGTGCAGACGCCGTCTTCCACCTTAATCTTGACT**
**GCGACGCCGATGATGGCCAGGTCCATCGCCTTTCTGACCGCGTGTTTTATGTATGCAGCCCTCTCATTCTCAGCCAGCGGCGG**
**AATCACAATGCCTGTCACGATTTCTCCCGGCTCCAGGGAAGTCTTTTTGACACCCAGGAAAAAGTCTTCGATTTGAATGACCC**
**GGTCCTTGTTGGGACCCTGTACCAATATCTTAGCGCCCTGAGCCAGTAGGTTGGGGGCTGTATCGCAGGATGGGGACGCATTG**
**CAGATATTGCCTGCCATGGTGCCCTTTGTCCTGATTTGTGTGGATGCCACTACCTTGGCGGCATGAGCCACTGCCGGATTCTT**
```

TTCTTTAACCAGAGGAGATGTCTCTATGGTACGCAGTGTTGTCAGCGCCCCTATTTTTAAACCTTCCCTGTCATCGTATTCCA
GATAATCCAAACCTGGAATTTTCTTAAGGTCAATGATATACTCTGGTGATATAACCTTGTCTTTCATTGGCGGAATCAGATCC
GTTGCGCCTGCCCATAACCCTGGCTGTGCTTCCAAGCTCAAGGAACAGATTGCACGCCTCCCCGATGGTCTTAGGAGCAAGATA
CTCGAATTGAGGTAATACCATcatatgTATATCTCCTTCTTAAAGTTAAACAAAATTATTTCTAGACGCCGTCCACGCTGCCT
CCTCACGTGACGTGAGGTGCAAGCCCGGACGTTCCGCGTGCCACGCCGTGAGCCGCCGCGTGCCGTCGGCTCCCTCAGCCCGG
GCGGCCGTGGGAGCCCGCCTCGATATGTACACCCGAGAAGCTCCCAGCGTCCTCCTGGGCCGCGATACTCGACCACCACGCAC
GCACACCGCACTAACGATTCGGCCGGCGCTCGATTCGGCCGGCGCTCGATTCGGCCGGCGCTCGATTCGGCCGGCGCTCGATT
CGGCCGGCGCTCGATTCGGCCGAGCAGAAGAGTGAACAACCACCGACCACGCTTCCGCTCTGCGCGCCGTACCCGACCTACCT
CCCGCAGCTCGAAGCAGCTCCCGGGAGTACCGCCGTACTCACCCGCCTGTGCTCACCATCCACCGACGCAAAGCCCAACCCGA
GCACACCTCTTGCACCAAGGTGCCGACCGTGGCTTTCCGCTCGCAGGGTTCCAGAAGAAATCGAACGATCCAGCGCGGCAAGG
TTCAAAAAGCAGGGGTTGGTGGGGAGGAGGTTTTGGGGGGTGTCGCCGGGATACCTGATATGGCTTTGTTTTGCGTAGTCGAA
TAATTTTCCATATAGCCTCGGCGCGTCGGACTCGAATAGTTGATGTGGGCGGGCACAGTTGCCCCATGAAATCCGCAACGGGG
GGCGTGCTGAGCGATCGGCAATGGGCGGATGCGGTGTTGCTTCCGCACCGGCCGTTCGCGACGAACAACCTCCAACGAGGTCA
GTACCGGATGAGCCGCGACGACGCATTGGCAATGCGGTACGTCGAGCATTCACCGCACGCGTTGCTCGGATCTATCGTCATCG
ACTGCGATCACGTTGACGCCGCGATGCGCGCATTCGAGCAACCATCCGACCATCCGGCGCCGAACTGGGTCGCACAATCGCCG
TCCGGCCGCGCACACATCGGATGGTGGCTCGGCCCCAACCACGTGTGCCGCACCGACAGCGCCCGACTGACGCCACTGCGCTA
CGCCCACCGCATCGAAACCGGCCTCAAGATCAGCGTCGGCGGCGATTTCGCGTATGGCGGGCAACTGACCAAAAACCCGATTC
ACCCCGATTGGGAGACGATCTACGCCCGGCCACCCCGTACACATTGCGGCAGCTGGCCACCATCCACACACCCCGGCAGATG
CCGCGTCGGCCCGATCGGGCCGTGGGCCTGGGCCGCAACGTCACCATGTTCGACGCCACCCGGCGATGGGCATACCCGCAGTG
GTGGCAACACCGAAACGGAACCGGCCGCGACTGGGACCATCTCGTCCTGCAGCACTGCCACGCCGTCAACACCGAGTTCACGA
CACCACTGCCGTTCACCGAAGTACGCGCCACCGCGCAATCCATCTCCAAATGGATCTGGCGCAATTTCACCGAAGAACAGTAC
CGAGCCCGACAAGCGCATCTCGGTCAAAAAGGCGGCAAGGCAACGACACTCGCCAAACAAGAAGCCGTCCGAAACAATGCAAG
AAAGTACGACGAACATACGATGCGAGAGGCGATTATCTGATGGGCGGAGCCAAAAATCCGGTGCGCCGAAAGATGACGGCAGC
AGCAGCAGCCGAAAAATTCGGTGCCTCCACTCGCACAATCCAACGCTTGTTTGCTGAGCCGCGTGACGATTACCTCGGCCGTG
CGAAAGCTCGCCGTGACAAAGCTGTCGAGCTGCGGAAGCAGGGGTTGAAGTACCGGGAAATCGCCGAAGCGATGGAACTCTCG
ACCGGGATCGTCGGCCGATTACTGCACGACGCCCGCAGGCACGGCGAGATTTCAGCGGAGGATCTGTCGGCGTAACCAAGTCA
GCGGGTTGTCGGGTTCCGGCCGGCGCTCGGCACTCGGACCGGCCGGCGGATGGTGTTCTGCCTCTGGCGCAGCGTCAGCTACC
GCCGAAGGCCTGTCATCGACCGGCTTCGACTGAAGTATGAGCAACGTCACAGCCTGTGATTGGATGATCCGCTCACGCTCGAC
CGCTACCTGTTCAGCTGCCGCCCGCTGGGCATGAGCAACGGCCAACTCTCGTTCAAGAGCTCGACCGCGCGGGTCCCGGACGG
GGAAGAGCGGGGAGCTTTGCCAGAGAGCGACGACTTCCCCTTGCGTTGGTGATTGCCGGTCAGGGCAGCCATCCGCCATCGTC
GCGTAGGGTGTCACACCCCAGGAATCGCGTCACTGAACACAGCAGCCGGTAGGACGACCATGACTGAGTTGGACACCATCGCA
AATCCGTCCGATCCCGCGGTGCAGCGGATCATCGATGTCACCAAGCCGTCACGATCCAACATAAAGACAACGTTGATCGAGGA
CGTCGAGCCCCTCATGCACAGCATCGCGGCCGGGGTGGAGTTCATCGAGGTCTACGGCAGCGACAGCAGTCCTTTTCCATCTG
AGTTGCTGGATCTGTGCGGGCGGCAGAACATACCGGTCCGCCTCATCGACTCCTCGATCGTCAACCAGTTGTTCAAGGGGGAG
CGGAAGGCCAAGACATTCGGCATCGCCCGCGTCCCTCGCCCGGCCAGGTTCGGCGATATCGCGAGCCGGCGTGGGGACGTCGT
CGTTCTCGACGGGGTGAAGATCGTCGGGAACATCGGCGCGATAGTACGCACGTCGCTCGCGCTCGGAGCGTCGGGGATCATCC
TGGTGGACAGTGACATCACCAGCATCGCGGACCGGCGTCTCCAAAGGGCCAGCCGAGGTTACGTCTTCTCCCTTCCCGTCGTT
CTCTCCGGTCGCGAGGAGGCCATCGCCTTCATTCGGGACAGCGGTATGCAGCTGATGACGCTCAAGGCGGATGGCGACATTTC
CGTGAAGGAACTCGGGGACAATCCGGATCGGCTGGCCTTGCTGTTCGGCAGCGAAAAGGGTGGGCCTTCCGACCTGTTCGAGG
AGGCGTCTTCCGCCTCGGTTTCCATCCCCATGATGAGCCAGACCGAGTCTCTCAACGTTTCCGTTTCCCTCGGAATCGCGCTG
CACGAGAGGATCGACAGGAATCTCGCGGCCAACCGATAAGCGCCTCTGTTCCTCGGACGCTCGGTTCCTCGACCTCGATTCGT
CAGTGATGATCACCTCACACGGCAGCGATCACCACTGACATATCGAGGTCAACGGTCGTGGTCCGGCGGGCACTCCTCGAAG
GCGCGGCCGACGCCCTTGAACGACTCGATGACTCTAGACGCATCCGAAACCTCCACCCCACTCACCTAGTCCGACATCCGTAC
CTTGGAAACCGACCTGTATTGGCATTTCAGTTGGACATCGACCAGTGGCGTTGCTAGGTTCAAGACCATGTCCAGCCCGAAGG
CGTCCAGACTCTAGCCACCGGAGGTAGTCCGGTGGCCACATCCCGTCGCGCCCGAACGTCACGCTCTTGTGTGGCCTTCCCTT
GTTGTTTGCGATCAGTGGCACACCTCTACCGTCTGAATTTCGAGTCTGGCCTCGGCTGCGCACATCTCGCACTGTGACGCTGT
CAGGTCACCCGCTTCGCGGCTACCAGTTCCTTTCATCGAATCGAGCTTCCGGTGCCGCCGCGCAGCCTCCCTGACCATCCTCA
GATTTTATGGAGTCTCGCAGTGCCTTTCGCTATCTACGTCCTCGGGCTTGCTGTCTTCGCCCAGGGCACATCCGAGTTCATGT
TGTCCGGACTCATACCGGACATGGCCCGTGACCTCGGGGTTTCGGTCCCCGCCGCCGGACTCCTCACCTCCGCCTTCGCGGTC
GGGATGATCATCGGCGCTCCGTTGATGGCTATCGCCAGCATGCGGTGGCCCCGGCGACGCGCCCTTCTGACATTCCTCATCAC
GTTCATGCTGGTCCACGTCATCGGCGCGCTCACCAGCAGCTTCGAGGTCTTGCTGGTCACACGCATCGTGGGAGCCCTCGCCA
ATGCCGGATTCTTGGCAGTGGCCCTGGGGGCGGCGATGGCGATGGTGCCCGCCGACATGAAAGGGCGCGCCACGTCCGTCCTC
CTCGGCGGTGTCACGATCGCATGTGTAGCCGGTGTTCCCGGGGGCGCCTTCCTGGGTGAAATGTGGGGCTGGCGTGCAGCGTT
CTGGGCTGTCGTCGTCATCTCCGCCCCTGCAGTGGTGGCGATTATGTTCGCCACCCCGGCCGAGCCGCTTGCAGAGTCCACAC
CGAATGCCAAGCGTGAACTGTCCTCACTGCGCTCACGCAAGCTCCAGCTCATGCTTGTCCTCGGGGCGCTGATCAACGGCGCA
ACGTTCTGTTCGTTCACGTACATGGCGCCCACGCTCACCGACATCTCCGGTTTCGACTCCCGTTGGATTCCGTTGCTGCTGGG
GCTGTTCGGGCTCGGATCGTTCATCGGTGTCAGCGTCGGAGGCAGGCTCGCCGACACCCGGCCGTTCCAACTGCTCGCTGTCG
GGTCCGCAGCACTGTTGACGGGATGGATCGTCTTCGCTCTCACGGCATCCCACCCCGCGGTGACATTGGTGATGCTGTTCGTG
CAGGGCGCTTTGTCCTTCGCGGTCGGCTCGACTTTGATCTCCCAGGTGCTCTACGCCGCCGACGCGGCACCGACCTTGGGTGG

```
908   ATCGTTCGCGACGGCCGCGTTCAACGTCGGTGCTGCACTGGGACCGGCGCTCGGCGGGTTGGCGATCGGCATGGGTCTGAGCT
909   ACCGCGCCCCGCTCTGGACGAGCGCCGCGCTGGTGACACTCGCGATCGTCATCGGCGCAGCCACCTTGTCTCTGTGGCGGCGA
910   CCAGCGTCTGTCCACGAATCTGTCCCCGCCTGACCAGAAACCAGGATCTGTGAGTGTGGTGACTGATCTGTGCACGCTCAGCA
911   GTCACCGCGCGCTCGCGTCGTACCGAGGGCCAGCGCCAACAGGTGTGTGGAGCTCTGCCCCTGCCTCTTTCACGCGAACTCAC
912   TGTTCAGTGCGGCGATACGTGCTCGGTGAGTTCCACTACAGCGACCATGACTAGAATTGATCTCCTCGACCGCCAATTGGGCA
913   TCTGAGAATCATCTGCGTTTCTCGCACGCAACGTACTTGCAACGTTGCAACTCCTAGTGTTGTGAATCACACCCCACCGGGGG
914   GTGGGATTGCAGTCACCGATTTGGTGGGTGCGCCCAGGAAGATCACGTTTACATAGGAGCTTGCAATGAGCTACTCCGTGGGA
915   CAGGTGGCCGGCTTCGCCGGAGTGACGGTGCGCACGCTGCACCACTACGACGACATCGGCCTGCTCGTACCGAGCGAGCGCAG
916   CCACGCGGGCCACCGGCGCTACAGCGACGCCGACCTCGACCGGCTGCAGCAGATCCTGTTCTACCGGGAGCTGGGCTTCCCGC
917   TCGACGAGGTCGCCGCCCTGCTCGACGACCCGGCCGCGGACCCGCGCGCGCACCTGCGCCGCCAGCACGAGCTGCTGTCCGCC
918   CGGATCGGGAAACTGCAGAAGATGGCGGCGGCCGTGGAGCAGGCGATGGAGGCACGCAGCATGGGAATCAACCTCACCCCGGA
919   GGAGAAGTTCGAGGTCTTCGGCGACTTCGACCCCGACCAGTACGAGGAGGAGGTCCGGGAACGCTGGGGGAACACCGACGCCT
920   ACCGCCAGTCCAAGGAGAAGACCGCCTCGTACACCAAGGAGGACTGGCAGCGCATCCAGGACGAGGCCGACGAGCTCACCCGG
921   CGCTTCGTCGCCCTGATGGACGCGGGTGAGCCCGCCGACTCCGAGGGGGCGATGGACGCCGCCGAGGACCACCGGCAGGGCAT
922   CGCCCGCAACCACTACGACTGCGGGTACGAGATGCACACCTGCCTGGGCGAGATGTACGTGTCCGACGAACGTTTCACGCGAA
923   ACATCGACGCCGCCAAGCCGGGCCTCGCCGCCTACATGCGCGACGCGATCCTCGCCAACGCCGTCCGGCACACCCCCTGAGCG
924   GTGGTCGTGGCCCGGGTCTCCCGCCCGGTCTCACCCCACGGCTCACTCCCGGGCCACGACCACCGCCGTCCCGTACGCGCACA
925   CCTCGGTGCCCACGTCCGCCGCCTCCGTCACGTCGAAACGGAAGATCCCCGGGTACCGAGCTCGTCAGGTGGCACTTTTCGGG
926   GAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAA
927   ATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTG
928   CCTTCCTGTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTTACATCG
929   AACTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTTCTG
930   CTATGTGGCGCGGTATTATCCCGTATTGACGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGT
931   TGAGTACTCACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCATGAGTG
932   ATAACACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCAT
933   GTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTAGCAAT
934   GGCAACAACGTTGCGCAAACTATTAACTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGG
935   ATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGGG
936   TCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAAC
937   TATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTCAGACCAAGTTTACTCAT
938   ATATACTTTAGATTGATTTAAAACTTCATTTTTAATTTAAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAA
939   ATCCCTTAACGTGAGTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCT
940   GCGCGTAATCTGCTGCTTGCAAAC

941

942   NdeI and XhoI restriction sites are lowercase and the *ucdCFO* insert is in bold.
```

943 **Online Methods**

944 **Resources table**

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background	*Enterocloster aldenensis* DSM 19262	DSMZ	DSM 19262	Type strain
Strain, strain background	*Enterocloster asparagiformis* DSM 15981	DSMZ	DSM 15981	Type strain
Strain, strain background	*Enterocloster bolteae* DSM 15670	DSMZ	DSM 15670	Type strain
Strain, strain background	*Enterocloster citroniae* DSM 19261	DSMZ	DSM 19261	Type strain
Strain, strain background	*Enterocloster clostridioformis* DSM 933	DSMZ	DSM 933	Type strain
Strain, strain background	*Enterocloster lavalensis* DSM 19851	DSMZ	DSM 19851	Type strain
Strain, strain background	*Lachnoclostridium pacaense* CCUG 71489T	CCUG	CCUG 71489T	Type strain
Strain, strain background	*Eggerthella lenta* A2	DSMZ	DSM 110911	
Strain, strain background	*Enterococcus faecium* DSM 20477	DSMZ	DSM 20477	Type strain
Strain, strain background	*Gordonibacter pamelaeae* DSM 19378	DSMZ	DSM 19378	Type strain
Strain, strain background	*Gordonibacter pamelaeae* DSM 110924	DSMZ	DSM 110924	
Strain, strain background	*Gordonibacter urolithinfaciens* DSM 27213	DSMZ	DSM 27213	Type strain
Strain, strain background	*Rhodococcus erythropolis* DSM 43066	DSMZ	DSM 43066	Type strain
Strain, strain background	*Escherichia coli* NEB10β	NEB	C3019	Cloning strain
Strain, strain background	*Escherichia coli* NEB10β pTipQC2	This study		
Strain, strain background	*Escherichia coli* NEB10β pTipQC2-*ucdCFO*	This study		
Strain, strain background	*Rhodococcus erythropolis* DSM 43066 pTipQC2	This study		
Strain, strain background	*Rhodococcus erythropolis* DSM 43066 pTipQC2-*ucdCFO*	This study		

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Commercial assay or kit	One-4-All Genomic DNA Minipreps Kit	BioBasic	BS88503	
Commercial assay or kit	EZ-10 Spin Column Plasmid DNA Miniprep Kit	BioBasic	BS414	
Commercial assay or kit	Direct-zol RNA Miniprep Kit	Zymo Research	R2051	
Commercial assay or kit	ZR BashingBead Lysis Tubes (0.1 & 0.5 mm)	Zymo Research	S6012-50	
Commercial assay or kit	Ambion *DNA-free* DNA Removal Kit	Invitrogen	AM1906	
Commercial assay or kit	*OneStep* PCR Inhibitor Removal Kit	Zymo Research	D6030	
Commercial assay or kit	iScript Reverse Transcription Supermix	Bio-Rad	1708840	RT-PCR with random hexamers
Commercial assay or kit	LunaScript RT Master Mix Kit (Primer-free)	NEB	E3025	For gene-specific RT-PCR
Commercial assay or kit	Luna Universal qPCR Master Mix	NEB	M3003	
Commercial assay or kit	Monarch PCR & DNA Cleanup Kit	NEB	T1030	
Commercial assay or kit	Monarch DNA Gel Extraction Kit	NEB	T1020	
Commercial assay or kit	Q5 High-Fidelity Polymerase	NEB	M0491	For 16S PCR and cloning
Commercial assay or kit	One*Taq* Quick-Load 2X Master Mix with Standard Buffer	NEB	M0486	For colony PCR
Commercial assay or kit	One*Taq* 2X Master Mix with Standard Buffer	NEB	M0482	For *ucd*-specific PCR
Commercial assay or kit	Platinum SuperFi II Green PCR Master Mix	Invitrogen	12369010	For ONT 16S PCR on fecal slurries
Commercial assay or kit	dNTP Mixture (10 mM)	BioBasic	DD0056	
Commercial assay or kit	Qubit dsDNA High Sensitivity (HS) Assay Kit	Invitrogen	Q32851	
Commercial assay or kit	Qubit RNA Broad Range (BR) Assay Kit	Invitrogen	Q10210	
Commercial assay or kit	GelCode Blue Stain Reagent	Fisher	24590	
Commercial assay or kit	XhoI Restriction Enzyme	NEB	R0146	
Commercial assay or kit	NdeI Restriction Enzyme	NEB	R0111	
Commercial assay or kit	Hi-T4 DNA Ligase	NEB	M2622	

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Plasmids and primers	pTipQC2	Hokkaido System Science Co.	RE-0006	GenBank AB127591.1
Plasmids and primers	pTipQC2-*ucdCFO*	This work	This work	See Supplementary Sequence 2
Commercial assay or kit	Amplicon barcoding kit AB01-24	ONT	SQK-AMB111.24	
Commercial assay or kit	Thermolabile proteinase K	NEB	P8111	
Commercial assay or kit	Agencourt AMPure XP	Beckman Coulter	A63880	
Commercial assay or kit	Flongle Flow Cell (R9.4.1)	ONT	FLO-FLG001	
Plasmids and primers	Custom oligonucleotides (Desalted, Dried Form)	Invitrogen	10336022	See Primer sequences table
Commercial assay or kit	ExcelBand 100 bp+3K DNA Ladder	SMOBIO	DM2300	0.1-3 kb
Commercial assay or kit	Quick-Load Purple 1 kb Plus DNA Ladder	NEB	N0550	0.1-10 kb
Commercial assay or kit	BLUelf Prestained Protein Ladder	FroggaBio	PM008-0500	5-235 kDa
Chemical compound, drug	SIGMAFAST Protease Inhibitor Cocktail, EDTA-free	Sigma	S8830-2TAB	
Chemical compound, drug	Acetonitrile	Sigma	34998	CAS 75-05-8
Chemical compound, drug	Ethyl acetate	Sigma	34858	CAS 141-78-6
Chemical compound, drug	Methanol	Sigma	34860	CAS 67-56-1
Chemical compound, drug	Ethanol	Commercial Alcohols	P016EAAN	CAS 64-17-5
Chemical compound, drug	Formic acid	Sigma	F0507	CAS 64-18-6
Chemical compound, drug	Peptone	BioShop	PEP403.1	
Chemical compound, drug	Yeast extract	BioShop	YEX401.500	CAS 8013-01-2
Chemical compound, drug	Sodium chloride	BioShop	SOD004.5	CAS 7647-14-5
Chemical compound, drug	Starch (from potato)	Sigma	S2004-500G	CAS 9005-25-8

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Chemical compound, drug	D-glucose monohydrate	BioShop	GLU601.1	CAS 14431-43-7
Chemical compound, drug	Sodium pyruvate	BioShop	PYR302.100	CAS 113-24-6
Chemical compound, drug	Sodium succinate dibasic	Sigma	14160-100G	CAS 150-90-3
Chemical compound, drug	Sodium thioglycolate	TCI	M0053	CAS 367-51-1
Chemical compound, drug	Agar	Fisher	BP1423-500	CAS 9002-18-0
Chemical compound, drug	L-arginine HCl	BioShop	ARG006.100	CAS 1119-34-2
Chemical compound, drug	L-cysteine	BioShop	CYS555.100	CAS 52-90-4
Chemical compound, drug	Sodium bicarbonate	BioShop	SOB308.500	CAS 144-55-8
Chemical compound, drug	Haemin	Sigma	H9039-1G	CAS 16009-13-5
Chemical compound, drug	Vitamin K1	Sigma	95271-250MG	CAS 84-80-0
Chemical compound, drug	Dithiothreitol (DTT)	Fisher	BP172-25	CAS 3483-12-3
Chemical compound, drug	Ferrous sulfate heptahydrate	BioShop	FER005	CAS 7782-63-0
Chemical compound, drug	Iron(III) pyrophosphate	Sigma	P6526-100G	CAS 10058-44-3
Chemical compound, drug	LB Broth (Miller)	BioShop	LBL407.1	
Chemical compound, drug	Tris base	Sigma	10708976001	CAS 77-86-1
Chemical compound, drug	Magnesium sulfate heptahydrate	BioShop	MAG511.1	CAS 10034-99-8
Chemical compound, drug	Calcium chloride dihydrate	Fisher	BP510-500	CAS 10035-04-8

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Chemical compound, drug	Lysozyme	BioShop	LYS702.10	CAS 12650-88-3
Chemical compound, drug	Triton X-100	Sigma	X100-100ML	CAS 9036-19-5
Chemical compound, drug	Glycerol	Sigma	G2025	CAS 56-81-5
Chemical compound, drug	Sodium dodecyl sulfate (SDS)	BioShop	SDS001	CAS 151-21-3
Chemical compound, drug	Bromophenol blue	BioShop	BRO777	CAS 6262-5-28-9
Chemical compound, drug	Urolithin C	TRC	U847015	CAS 165393-06-6
Chemical compound, drug	Urolithin A	TRC	U847000	CAS 1143-70-0
Chemical compound, drug	Urolithin A	Sigma	SML1791-5MG	CAS 1143-70-0
Chemical compound, drug	Urolithin M-6	TRC	U847040	CAS 1006683-97-1
Chemical compound, drug	Isourolithin A	TRC	I917520	CAS 174023-48-4
Chemical compound, drug	Urolithin B	TRC	U847005	CAS 1139-83-9
Chemical compound, drug	Salicylic acid	Sigma	84210-100G	CAS 69-72-7
Chemical compound, drug	Caffeic acid	Sigma	C0625-2G	CAS 331-39-5
Chemical compound, drug	m-coumaric acid	MedChemExpress	HY-113357	CAS 588-30-7
Chemical compound, drug	Dopamine hydrochloride	Sigma	H8502-5G	CAS 62-31-7
Chemical compound, drug	m-tyramine	MedChemExpress	HY-128975	CAS 38449-59-1
Chemical compound, drug	Entacapone	Sigma	SML0654-10MG	CAS 130929-57-6

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Chemical compound, drug	2,2'-bipyridyl	Sigma	D216305-10G	CAS 366-18-7
Chemical compound, drug	Resorcinol	Sigma	398047-100G	CAS 108-46-3
Chemical compound, drug	2-bromo-4,5-dimethoxybenzoic acid	Sigma	441074-5G	CAS 6286-46-0
Chemical compound, drug	Sodium hydroxide	Sigma	221465-500G	CAS 1310-73-2
Chemical compound, drug	Copper (II) sulfate pentahydrate	Sigma	209198-100G	CAS 7758-99-8
Chemical compound, drug	TRI reagent	Zymo Research	R2050-1-50	
Chemical compound, drug	NADH disodium salt, trihydrate, reduced	BioBasic	NB0642.SIZE.1g	CAS 606-68-8
Chemical compound, drug	NADPH	BioBasic	Y4433000.SIZE.100mg	CAS 2646-71-1
Chemical compound, drug	FAD disodium salt hydrate	Sigma	F6625-100MG	CAS 84366-81-4
Chemical compound, drug	Agarose	Fisher	BP160-500	CAS 9012-36-6
Chemical compound, drug	SafeView Classic	Abm	G108	
Chemical compound, drug	PBS pH 7.4 (10X)	Gibco	70011-044	
Chemical compound, drug	Nuclease-free water	Invitrogen	10977-015	
Lab equipment, instruments	MilliQ water system	Millipore	IQ7000	
Lab equipment, instruments	Gene Pulser Cuvette (0.1 cm gap)	Bio-Rad	1652089	
Lab equipment, instruments	Gene Pulser Xcell Total Electroporation System	Bio-Rad	1652660	
Lab equipment, instruments	AnaeroPack System	Mitsubishi Gas Chemical Company	10-01	

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Lab equipment, instruments	Vinyl anaerobic chambers	Coy Laboratory Products	Type A	
Lab equipment, instruments	Multiscan FC Microplate Photometer	Thermo Fisher	51119100	
Lab equipment, instruments	BioTek Epoch 2 Microplate Spectrophotometer	Agilent	EPOCH2NS-SN	
Lab equipment, instruments	Costar 96-Well Clear Round Bottom	Corning	3788	
Lab equipment, instruments	Breathe-Easy Sealing Membranes	Sigma	Z380059-1PAK	
Lab equipment, instruments	LifeECO Thermal Cycler	BIOER	TC-96	
Lab equipment, instruments	Nanodrop 2000c	Thermo Fisher	ND-2000C	
Lab equipment, instruments	Mini Beadbeater 16	Biospec	607	
Lab equipment, instruments	Viia7 qPCR machine	Applied Biosystems	4453535	
Lab equipment, instruments	MicroAmp Fast 96-Well Reaction Plates (0.1 mL)	Applied Biosystems	4346907	
Lab equipment, instruments	PlateSeal Film, Clear Polyester, X-Large, Non-Sterile	PlateSeal	PS-PETSTXL-100	
Lab equipment, instruments	LC-MS (Agilent 1260 Infinity II LC connected to a 6120 Quadrupole MS)	Agilent		
Lab equipment, instruments	Poroshell 120 EC-C18 column (4.6x50 mm, 2.7 µm)	Agilent	699975-902	
Lab equipment, instruments	Poroshell 120 EC-C18 column (4.6x5 mm, 2.7 µm) guard column	Agilent	820750-911	
Lab equipment, instruments	Vacuum concentrator	Heto Lab	VR-1	
Lab equipment, instruments	Sonicator	Misonix	S3000	
Lab equipment, instruments	Rotary evaporator	Buchi	R-300	
Software	Agilent OpenLab CDS	Agilent	N/A	

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Software	GraphPad Prism 10	GraphPad by Dotmatics	N/A	
Software	Mascot 2.6.2	Matrix Science	N/A	
Software	Scaffold 5	Proteome Software Inc	N/A	
Software	BioTek Gen6 v1.03.01	Agilent	N/A	
Software	R Studio 2023.06.0+421	Posit Software	N/A	
Software	Affinity Designer	Serif Ltd	N/A	

945 NEB = New England Biolabs, ONT = Oxford Nanopore Technologies, TRC = Toronto Research
946 Chemicals

947 **Primer sequences table**

Designation	Purpose	Source	5'-Sequence-3'
16S_V1_27_f	16S PCR		AGAGTTTGATCMTGGCTAG
16S_V9_1492_r	16S PCR		TACGGYTACCTTGTTAYGACTT
ONT_16S_27_f_GGK	16S PCR		**GGK**AGRGTTYGATYMTGGCTCAG
ONT_16S_1492_r_GGK	16S PCR		**GGK**CCGGYTACCTTGTTACGACTT
Eb_ucdO_qPCR_f	qPCR	This work	GTGGATGAGGAAACCGCCCAGC
Eb_ucdO_qPCR_r	qPCR	This work	TGCCCTCCCAGTTTACCTCGGG
Eb_ucdF_qPCR_f	qPCR	This work	GCGCCTGTACCGTGATGCTTGA
Eb_ucdF_qPCR_r	qPCR	This work	CCCTCCCTGGCAATTCCCTCCA
Eb_ucdC_qPCR_f	qPCR	This work	AGCCCCCAACCTACTGGCTCAG
Eb_ucdC_qPCR_r	qPCR	This work	GCCTGTCACGATTTCTCCCGGC
Eb_dnaK_Ref_qPCR_f	qPCR	This work	GGTGCTGTTGGTAGGCGGTTCC
Eb_dnaK_Ref_qPCR_r	qPCR	This work	GGGTCTTGCTGGGTTCCTTGCC
ucdCFO_RT-PCR_f	RT-PCR	This work	GCAATCTGTTCCTYGAGCTTGG
ucdCFO_RT-PCR_r	RT-PCR	This work	CGTGSAGCGCTCTCCAGATATG
Eb_ucdCFO_NdeI_f	Cloning	This work	CCCTGGcatatgATGGTATTACCTCAATTCGAG
Eb_ucdCFO_XhoI_r	Cloning	This work	CTGTCTctcgagCTGCCTTTATTTTGAATTCCTG
pTipQC2-ucdCFO_cPCR_f	cPCR	This work	CGAGGGAGCTTCCAGGGGGAAA
pTipQC2-ucdCFO_cPCR_r	cPCR	This work	GAAGGCCAGCTGGAGGGATCCA

948 **Adapter sites** are bolded. Restriction sites are lowercase.

Anaerobic bacterial strains and culturing conditions

Bacterial strains used in this study are listed in the *Resources table*. All bacterial stains were validated by sequencing the 16S rRNA gene (see Genomic DNA extraction and 16S rRNA sequencing of bacterial isolates). The same culture used for validation was used to make 25% glycerol stocks. Anaerobic strains were grown from glycerol stocks on mABB+H agar plates (recipe below) for 48-72 h at 37 °C in a vinyl anaerobic chamber, which was maintained with a gas mixture of 3% H_2, 10% CO_2, 87% N_2. To make overnight cultures, a single colony was inoculated into 5 mL of liquid mABB or mABB+H medium and incubated at 37 °C between 16-48h, depending on the bacterium (16-24 h for *Enterocloster* spp. and *E. faecium*, and 48 h for *E. lenta* and *Gordonibacter* spp.).

Human fecal sample collection

Human fecal samples were collected under the approval of protocol A04-M27-15B by the McGill Faculty of Medicine Institutional Review Board. Informed written consent was received from the participants for the use of human samples. Eligibility criteria for the healthy participants were as follows: body mass index between 18.5–30, no diagnosed gastrointestinal disease, no ongoing therapeutic treatment, and no antibiotic usage 3 months prior to the start of the study. Subject information was recorded at the time of sampling. The age of donors ranged from 21–40 years. Fresh fecal samples were collected and placed immediately in an anaerobic chamber, aliquoted, and stored at −70 °C until use.

Modified anaerobe basal broth (mABB and mABB+H)

For 1 L of modified anaerobe basal broth (mABB), the following components were dissolved in MilliQ water, then autoclaved: 16 g peptone, 7 g yeast extract, 5 g sodium chloride, 1 g starch, 1 g D-glucose monohydrate, 1 g sodium pyruvate, 0.5 g sodium succinate, 1 g sodium thioglycolate, 15 g agar (for plates). The autoclaved solution was allowed to cool, then

the following filter-sterilized solutions were added aseptically: 10 mL of 100 mg/mL L-arginine-

HCl, 10 mL of 50 mg/mL L-cysteine, 8 mL of 50 mg/mL sodium bicarbonate, 50 µL of 10 mg/mL

vitamin K1, 20 mL of 50 mg/mL dithiothreitol, and, for mABB+H, 10 mL of 0.5 mg/mL haemin.

The media was then placed in the anaerobic chamber and allowed to reduce for at least 24 h

prior to its use in experiments.

Genomic DNA extraction of isolates and identity validation

The identities of all bacteria in this study were validated by full-length (V1-V9) 16S rRNA

sequencing. DNA was first extracted from 0.5-1 mL of overnight culture using the One-4-All

Genomic DNA Miniprep Kit (BioBasic) according to the manufacturer's instructions. The purified

genomic DNA (2 µL, ~ 20 ng) was used as a template for PCR reactions (25 µL reaction volume)

using the Q5 High-Fidelity polymerase (NEB). PCR tubes were placed in a thermal cycler and

targets (~1.5 kb) were amplified according to the following cycling conditions: 30 s at 98 °C, 30

cycles (10 s at 98 °C, 20 s at 60 °C, 45 s at 72 °C), 2 min at 72 °C, and hold at 10 °C. 5 µL of

the reaction was mixed with 6X loading buffer and loaded onto a 1% agarose gel (made with

1X TAE buffer) containing SafeView Classic (Abm). PCR product sizes were compared to the

ExcelBand 100 bp+3K DNA Ladder (SMOBIO).

PCR products (~1.5 kb) were purified using the Monarch PCR & DNA Cleanup Kit (NEB)

according to the manufacturer's instructions for products < 2 kb. Purified 16S PCR products

were eluted in nuclease-free water, quantified using the Qubit dsDNA HS assay kit (Invitrogen),

and adjusted to 30 ng/µL. Samples were submitted to Plasmidsaurus for long-read sequencing

using Oxford Nanopore Technologies (v14 library preparation chemistry, R10.4.1 flow cells).

Treatments with urolithins and other catechols

994   All treatments used in this study (urolithin C, urolithin M6, urolithin A, isourolithin A, 8,9-di-O-methyl-urolithin C, dopamine, m-tyramine, caffeic acid, m-coumaric acid, entacapone, and 2,2'-bipyridyl) were dissolved in DMSO to a concentration of 10 mM.

Treatment prior to growth (metabolism only): Overnight cultures of bacteria were diluted 1/50 into fresh mABB+H. Treatments (10 mM stocks solutions, dissolved in DMSO) were added to the diluted bacterial suspension to a final concentration of 100 µM and samples were incubated for 24 h at 37 °C in an anaerobic chamber.

Treatment during growth (spike-in): Overnight cultures of *Enterocloster* spp. were diluted 1/50 into fresh mABB+H and incubated at 37 °C in an anaerobic chamber. After 5 hours of incubation (~OD_{620} from a 200 µL sample ~ 0.4), 10 mM urolithins (or an equivalent volume of DMSO) were added to the growing cultures at a final concentration of 50 or 100 µM for protein expression or RNA expression, respectively. For protein expression analyses and inducibility tests, the cultures were incubated for an additional 4 h. For RNA expression analyses, the cultures were incubated for an additional 2 h.

Treatment prior to growth (growth curves): Overnight cultures of *Enterocloster* spp. were diluted 1/25 into mABB (with or without added 15.4 µM iron source), depending on the experimental design. Separately, treatments (10 mM stocks solutions, dissolved in DMSO) were prepared in mABB (with or without added 15.4 µM iron source) to a final concentration of 200 µM. In each well of a 96-well plate, 100 µL of 1/25 bacteria and 200 µM treatment were combined. These were plated in technical duplicates. The final concentration of treatment was 100 µM (unless otherwise specified in concentration-response experiments) and the final dilution of bacteria was 1/100.

Urolithin extraction from fecal slurries or bacterial cultures

Frozen (-70 °C) fecal slurries or bacterial cultures were thawed at room temperature. For quantification of urolithin concentrations, urolithin standards (stock 10 mM in DMSO) were spiked into separate media aliquots immediately before extraction.

Extraction Method A: This method was used for cultures. Salicylic acid (3 mg/mL in DMSO) was spiked-in as an internal standard at a final concentration of 50 µg/mL. The cultures and standards were then extracted with 3 volumes of ethyl acetate + 1% formic acid (e.g., 600 µL solvent to 200 µL thawed culture). The organic phase (top) was transferred to a new tube and dried in a vacuum concentrator (Heto Lab) connected to a rotary evaporator (Buchi). After solvent removal, samples were redissolved in 0.5 volumes (relative to the starting culture) of 50% MeOH:H_2O. Samples were centrifuged at 20,000 x g for 5 min to pellet insoluble material, then transferred to LC-MS vials. Urolithins were then analyzed by LC-MS.

Extraction Method B: This method was used for cultures and pre-induced cell suspensions. Samples were diluted with an equal volume of MeOH, vortexed briefly, and incubated at room temperature for 10 min. Samples were centrifuged at 20,000 x g for 5 min to pellet insoluble material, then transferred to LC-MS vials. Urolithins were then analyzed by LC-MS.

Extraction Method C: This method was used to extract urolithins from crude bacterial lysates. Lysates were diluted with 3 volumes of MeOH, vortexed briefly, and incubated at room temperature for 10 min. Samples were centrifuged at 20,000 x g for 5 min to pellet insoluble material, then transferred to LC-MS vials. Urolithins were then analyzed by LC-MS.

LC-MS method to quantify urolithins

Samples (10 µL) were injected into a 1260 Infinity II Single Quadrupole LC/MS system (Agilent) fitted with a Poroshell 120 EC-C18 4.6x50 mm, 2.7 µm column (Agilent). The mobile

phase was composed of MilliQ water + 0.1% formic acid (solvent A) and acetonitrile + 0.1%

formic acid (solvent B). The flow rate was set to 0.7 mL/min. The gradient was as follows: 0-8

min: 10-30 %B, 8-10 min: 30-100 %B, 10-13.5 min: 100 %B isocratic, 13.5-13.6 min: 100-10

%B, then 13.6-15.5 min: 10 %B. The multiple wavelength detector was set to monitor

absorbance at 305 nm. The mass spectrometer was run in negative mode in both selected ion

monitoring (SIM) and scan (100-1000 m/z) modes to validate peak identities. Peaks were

validated based on retention times compared to spike-in standards and mass-to-charge ratios.

To quantify urolithins, peak areas for the compounds of interest were compared to spike-in

standards of known concentration(s). When standards were not available (urolithin M7), the

extracted ion chromatogram was used ($[M-H]^- = 243$).

Synthesis of di-O-methyl-urolithin C

Di-O-methyl-urolithin C (3-Hydroxy-8,9-dimethoxy-6H-dibenzo[b,d]pyran-6-one, CAS

126438-35-5) was synthesized based on previously described Ullmann-type coupling

conditions for urolithin derivatives [60]. Resorcinol (213 mg, 2 mmol) and 2-bromo-4,5-

dimethoxybenzoic acid (261 mg, 1 mmol) were dissolved in 1 mL of 8% w/v NaOH (in MilliQ

H_2O) and heated in a thermo-shaker set to 100 °C for 20 min (in 1.7 mL tube). Then, 200 µL of

a 10% w/v Cu(II)SO$_4$ pentahydrate solution was added and the reaction was heated at 100 °C

for 1 h. The reaction solution (pink-red coloration) contained an insoluble precipitate which was

collected by centrifugation (20,000 x g for 30 s). The insoluble pellet was washed 7 times with

1 mL of MilliQ H_2O until the pH of the wash solution was equal to the pH of MilliQ H_2O (~pH 6).

The pellet was dried by lyophilization for 16 h (0.0010 mbar, -90 °C) and the product was

recovered as a pale pink solid (94 mg, 35% yield).

^{1}H NMR (600 MHz, (CD$_3$)$_2$SO)): δ = 10.22 (s, 1H), 8.21 (d, J = 8.76 Hz, 1H), 7.69 (s, 1H), 7.54

(s, 1H), 6.83 (dd, J = 8.67, 2.37 Hz, 1H), 6.74 (d, J = 2.34 Hz, 1H), 4.02 (s, 3H), 3.89 (s, 3H);

HRMS: m/z [M+Na]$^+$ calculated for C$_{15}$H$_{12}$NaO$_5$: 295.0577, found: 295.0585.

Phylogenetic tree construction

1065

Phylogenetic trees based on the 16S rRNA gene were constructed using the DSMZ

single-gene phylogeny server (https://ggdc.dsmz.de/phylogeny-service.php#)[61] with GenBank

16S rRNA sequence accessions: *E. aldenensis* (DQ279736), *E. asparagiformis* (AJ582080), *E.*

bolteae (AJ508452), *E. citroniae* (HM245936), *E. clostridioformis* (M59089), *E. lavalensis*

(EF564277), *L. pacaense* (LT799004), *G. pamelaeae* (AM886059), *G. urolithinfaciens*

(HG000667), *E. isourolithinifaciens* (MF322780).

Phylogenetic trees based on whole genomes and proteomes were constructed using the

Type (Strain) Genome Server (TYGS, https://tygs.dsmz.de)[40,62] with the following GenBank

genome accessions: *E. aldenensis* (GCA_003467385.1), *E. asparagiformis*

(GCA_025149125), *E. bolteae* (GCA_000154365), *E. clostridioformis* (GCA_900113155), *E.*

lavalensis (GCA_003024655), *L. pacaense* (GCA_900566185). For *E. citroniae*, the Integrated

Microbial Genomes ObjectID was used: *E. citroniae* (2928404274). Further information on

nomenclature and taxonomy was obtained from the List of Prokaryotic names with Standing in

Nomenclature (LPSN, available at https://lpsn.dsmz.de).

Cell suspension assay to test inducibility

1080

Bacteria (10 mL growing cultures in mABB+H media) were grown with 50 µM uroC (or

an equivalent volume of DMSO) as detailed in the *Treatment during growth (spike-in)* section

above and incubated for 4 h at 37 °C. Cultures were then pelleted at 6,500 x g for 3 min and

the supernatants were discarded. The cells were washed with 10 mL of pre-reduced PBS

(placed in the anaerobic chamber 24 h before), re-pelleted, and resuspended in 2 mL of pre-

reduced PBS. For each condition tested, a 200 µL aliquot of the cell suspension was transferred

into a sterile 1.5 mL tube, and 10 mM urolithins (uroM6, uroC, isouroA, or DMSO) were added

at a final concentration of 100 µM. Cell suspensions were briefly vortexed and incubated at

room temperature in the anaerobic chamber for 16h prior to freezing and urolithin extraction

using *Extraction Method B*.

RNA extraction from isolates

A volume of 1.5 mL of treated (100 µM urolithin C for 2 h) *Enterocloster* spp. culture (see

Enterocloster spp. urolithin C treatments) was pelleted (6,500 g for 3 min) and the supernatant

was removed for later LC-MS analysis. The pellet (suspended in 200 µL of media) was then

mixed with 800 µL TRI reagent (Zymo Research) and transferred to a ZR BashingBead lysis

tube (Zymo Research). Samples were lysed in a Mini Beadbeater 16 (Biospec) according to

the following sequence: 1 min ON, 5 min OFF. For RNA-sequencing, bead beating was done

for a total of 5 min ON. For RT-(q)PCR, bead beating was done for a total of 2 min ON to

preserve longer transcripts. RNA isolation was then performed using the Direct-zol RNA

Miniprep Kit (Zymo Research) according to the manufacturer's instructions (including an on-

column DNase digestion). To ensure complete DNA removal, an additional DNA digestion step

was performed on the isolated RNA using the Ambion *DNA-free* DNA Removal Kit (Invitrogen)

according to the manufacturer's instructions. The DNA-free RNA was then cleaned up using

the *OneStep* PCR Inhibitor Removal Kit (Zymo Research). RNA concentration and quality were

initially verified by NanoDrop and 1 % agarose gel electrophoresis. For RNA-sequencing, RNA

integrity was assessed by Génome Québec using a Bioanalyzer 2100 (Agilent). RNA integrity

(RIN) values ranged between 7.5-7.8 for *E. asparagiformis* DSM 15981 and 7.0-7.3 for *E.*

bolteae DSM 15670.

RNA-sequencing of urolithin C-treated *E. bolteae* and *E. asparagiformis* isolates

Total RNA was sent to Génome Québec for library preparation and RNA-sequencing. Briefly, total RNA was prepared for Illumina sequencing using the NEBNext rRNA Depletion Kit (Bacteria) (NEB) kit to remove rRNA and using the NEBNext Multiplex Oligos for Illumina (NEB) kit (stranded/directional). Prepared libraries were quality checked with a Bioanalyzer 2100 (Agilent) prior to sequencing. Sequencing was performed on a NovaSeq 6000 (Illumina) with the following flow cell/settings: S4 flow cell, 100 bp, 25 M reads, paired end.

Analysis of RNA-seq reads was done using Galaxy bioinformatics cloud computing (https://usegalaxy.org/) hosted by Compute Canada Genetics and Genomics Analysis Platform (GenAP) (https://www.genap.ca/). Genomes and annotations were fetched from the NCBI genome browser: *E. bolteae* (ASM223457v2) (accessed 2022/05/11) and *E. asparagiformis* (ASM2514912v1) (accessed 2023/09/19). Raw reads were first verified for quality using FastQC (v0.73, https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) with default parameters. FastQC reports were aggregated into MultiQC (v1.11) [63]. The mean sequence quality scored were above 35 for all samples. Raw reads were then trimmed using Cutadapt (v3.7) [64] to trim adapter sequences (R1 sequence: AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC, and R2 sequence: AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT) that were not removed after sequencing using default parameters for paired end reads. Trimmed reads were then aligned to reference genomes for each bacterium using HISAT2 (v2.2.1) [65] with paired-end parameters and reverse strandedness (RF). Aligned read counts were assigned to features in annotation files (.gtf) using featureCounts (v2.0.1) [66] with the following parameters: reverse strandedness, count fragments instead of reads, GFF feature type filter = "gene", multi-mapping and multi-overlapping features included (-M -O), minimum mapping quality per read of 0, and the rest of

the parameters were kept as default. Differential gene expression analysis was then performed

using DESeq2 (v2.11.40.7) [67] using default parameters. Differential expression tables were

annotated with the Annotate DESeq2/DEXSeq output tables tool (v1.1.0) in Galaxy to include

the following: GFF feature type = "CDS", GFF feature identifier = "gene_id", GFF transcript

identifier = "transcript_id", GFF attribute to include = "protein_id, product". The "protein_id" was

used to query the NCBI database and the NCBI Sequence Viewer was used to investigate the

genomic context surrounding genes of interest.

Comparative genomics

The nucleotide sequence for the *Enterocloster bolteae* DSM 15670 *ucd* operon (NCBI

NZ_CP022464 REGION: complement(4417875..4421605)) was used as a query for BLASTn

(megablast) searches using the refseq_genomes database limited to Bacteria (taxid:2). The

NCBI multiple sequence alignment (MSA) viewer was used to download alignment figures.

RT-PCR analysis of *E. bolteae* to determine *ucd* operon structure

Isolated RNA samples (500 ng) were reverse transcribed using the LunaScript® RT

Master Mix Kit (Primer-free) (NEB) in a reaction volume of 10 µL containing the ucdCFO_RT-

PCR_r primer at a final concentration of 1 µM. The No-RT Control included in the kit was used

as a no-enzyme control for reverse transcription. The reaction mixtures were incubated in a

thermal cycler: 10 min at 55°C, 1 min at 95°C. PCR reactions were conducted using the OneTaq

2X Master Mix with Standard Buffer (NEB). The ucdCFO_RT-PCR primer pair was added to

the master mix (to a final concentration of 0.2 µM) and 1 µL of template (cDNA, -RT, no

template, or gDNA) was added for a total reaction volume of 20 µL. PCR tubes were placed in

a thermal cycler and targets were amplified according to the following conditions: 20s at 94°C,

31 cycles (20s at 94°C, 30s at 62°C, 3 min at 68°C), 5 min at 68°C. A volume of 5 µL of reaction

was directly loaded onto a 1% agarose gel (made with 1X TAE buffer) containing SafeView

Classic (Abm). PCR product sizes were compared to the Quick-Load® Purple 1 kb Plus DNA

Ladder (NEB). The rest of the PCR product was then run on a 1% agarose gel and bands

corresponding to the desired products were cut out and purified using the Monarch DNA Gel

Extraction kit (NEB). DNA was quantified using the Qubit dsDNA HS assay kit (Invitrogen) and

submitted to Plasmidsaurus for long-read sequencing using Oxford Nanopore Technologies

(Supplementary Sequence 1).

RT-qPCR analysis of *E. bolteae ucd* operon genes

Isolated RNA samples (500 ng) were reverse transcribed using the iScript Reverse

Transcription Supermix (Bio-Rad) in a reaction volume of 10 μL. The iScript No-RT Control

Supermix was used as a no enzyme control for reverse transcription (-RT). The reaction

mixtures were incubated in a thermal cycler: 5 min at 25 °C, 20 min at 48 °C, and 1 min at 95

°C. Both cDNA and -RT controls were diluted 1/20 in nuclease-free water before use. qPCR

reactions were conducted using the Luna Universal qPCR Master Mix kit (NEB). The

Eb_ucdO_qPCR, Eb_ucdF_qPCR, Eb_ucdC_qPCR, and Eb_dnaK_Ref_qPCR primer pairs

were added to their respective master mixes (final primer concentration of 250 nM) and 6.6 or

4.4 μL of diluted template (cDNA, -RT, no template) were added to 26.4 or 17.6 μL of master

mix for triplicates or duplicates, respectively. All cDNA samples were run in technical triplicates,

while other sample types were run in technical duplicates. Replicate mixes were pipetted (10

μL/well) into a MicroAmp Fast 96-Well Reaction Plate (Applied Biosystems) and the plates were

sealed, then spun down for 2 min to eliminate air bubbles. The qPCR detection parameters

were as follows: SYBR Green detection, ROX reference dye, 10 μL reaction volume. The

thermal cycling conditions were: 1 min at 95 °C, 40 cycles (15 s at 95 °C, 30 s at 60 °C), then

melt analysis (60-95 °C). Data were analyzed according to the $2^{-\Delta\Delta Ct}$ method [68] with the dnaK

gene serving as the reference gene (*E. bolteae* dnaK RNA-seq $log_2FC = 0.122$).

Protein extraction from *Enterocloster* spp.

All steps other than sonication were carried out under anaerobic conditions. To extract proteins, 10 mL of treated (50 μM urolithin C for 4 h) *Enterocloster* spp. culture (see *Enterocloster* spp. urolithin C treatments) were pelleted (6,500 g for 3 min) and the supernatant was discarded. The pellet was washed with 10 mL of pre-reduced PBS, pelleted again, and resuspended in 0.5 mL of pre-reduced lysis buffer (20 mM Tris, pH 7.5, 500 mM NaCl, 10 mM MgSO$_4$, 10 mM CaCl$_2$, and 1 tablet/100 mL SIGMAFAST protease inhibitor (EDTA-free)). The resuspended pellet was then sonicated on ice using a Misonix Sonicator 3000 set to power level 2/10 according to the following sequence (aerobically, in a cold room): 20 s ON, 40 s OFF, for a total of 2 min ON. Tubes were centrifuged at 20,000 x g for 2 min to pellet insoluble particles and 0.4 mL of lysate was transferred to a new 1.5 mL tube (kept on ice). Lysates used in metabolism assays were transported to the anaerobic chamber in a sealed plastic bag containing an anaerobic gas generating system to minimize loss in activity.

Urolithin metabolism assays using crude lysates from uroC-induced *E. bolteae*

Protein lysates (described above) were aliquoted (50 μL aliquots) into 1.5 mL tubes, then treated with DMSO or urolithin C (10 mM stock) at a final concentration of 350 μM. Cofactors (NADPH, NADH, and FAD, each dissolved to a final concentration of 30 mM (in lysis buffer immediately before the assay was run) were added individually to the lysates at a final concentration of 1 mM. The lysates were incubated at room temperature in an anaerobic chamber for 20 h prior to freezing at -70 °C. Samples were then extracted using *Extraction Method C*.

To assess the oxygen sensitivity of crude lysates from uroC-induced *E. bolteae*, samples were prepared as described above. After adding DMSO or uroC and NADH (under anaerobic conditions), tubes were either incubated at room temperature inside the anaerobic chamber or

just outside of the chamber for 20 h. Afterwards, samples were frozen at -70 °C and then

extracted using *Extraction Method C*.

Proteomics analysis of uroC-treated *E. bolteae*

Extracted proteins were submitted for proteomic analysis at the RI-MUHC. For each

sample, protein lysates were loaded onto a single stacking gel band to remove lipids,

detergents, and salts. The gel band was reduced with DTT, alkylated with iodoacetic acid, and

digested with trypsin. Extracted peptides were re-solubilized in 0.1% aqueous formic acid and

loaded onto a Thermo Acclaim Pepmap (Thermo, 75 um ID X 2 cm C18 3 um beads) precolumn

and then onto an Acclaim Pepmap Easyspray (Thermo, 75 um ID X 15 cm with 2 um C18

beads) analytical column separation using a Dionex Ultimate 3000 uHPLC at 250 nL/min with

a gradient of 2-35% organic (0.1% formic acid in acetonitrile) over 3 hours. Peptides were

analyzed using a Thermo Orbitrap Fusion mass spectrometer operating at 120,000 resolution

(FWHM in MS1) with HCD sequencing (15,000 resolution) at top speed for all peptides with a

charge of 2+ or greater. The raw data were converted into *.mgf format (Mascot generic format)

for searching using the Mascot 2.6.2 search engine (Matrix Science) against *Enterocloster*

bolteae DSM 15670 proteins (NCBI assembly GCF_002234575.2) and a database of common

contaminant proteins. Mascot was searched with a fragment ion mass tolerance of 0.100 Da

and a parent ion tolerance of 5.0 ppm. O-63 of pyrrolysine, carboxymethyl of cysteine and j+66

of leucine/isoleucine indecision were specified in Mascot as fixed modifications. Deamidation

of asparagine and glutamine and oxidation of methionine were specified in Mascot as variable

modifications.

The database search results were loaded into Scaffold Q+ Scaffold_5.0.1 (Proteome Sciences)

for statistical treatment and data visualization. Scaffold (v5.3.0) was used to validate MS/MS

based peptide and protein identifications. Peptide identifications were accepted if they could be

established at greater than 95.0% probability by the Peptide Prophet algorithm [69] with Scaffold

delta-mass correction. Protein identifications were accepted if they could be established at

greater than 99.0% probability and contained at least 2 identified peptides. Protein probabilities

were assigned by the Protein Prophet algorithm [70]. Proteins that contained similar peptides and

could not be differentiated based on MS/MS analysis alone were grouped to satisfy the

principles of parsimony. Proteins sharing significant peptide evidence were grouped into

clusters. Protein quantification and differential expression were determined in Scaffold using

the following parameters: Quantitative method was set to total spectra, the minimum value was

set to 0.5 in case proteins were not detected in one condition, and statistical tests were

performed using Fisher's exact test with the Benjamini-Hochberg multiple test correction at a

significance level set to 0.05.

Protein structures

The protein FASTA sequences (NCBI RefSeq accessions for UcdC: WP_002569575.1,

UcdF: WP_002569574.1, UcdO: WP_002569573.1) for the *Enterocloster bolteae* DSM 15670

ucd operon were used as a query for BLASTp searches against the UniProtKB reference

proteomes + Swiss-Prot databases. The AlphaFold2 protein structures for matches (UniProt

accessions for UcdC: A8RZR5, UcdF: A8RZR3, UcdO: A8RZR2) of each protein were

downloaded and imported into PyMOL (v2.4.1). Foldseek (in 3Di/AA mode) was used to

generate a list of proteins with similar structures from solved crystal structures in the Protein

Data Bank (PDB) [48].

Hits of published X-ray crystal structures (PDB: 1ZXI (from *Afipia carboxidovorans* OM5) and

PDB: 3UNI (from *Bos taurus*)) were fetched from the PDB and imported into PyMOL. The

AlphaFold2 structures for each Ucd protein were aligned to the following chains in the published

PDBs using the "super" command: PDB 1ZXI: UcdC to chain C, UcdF to chain A, UcdO to chain

B; PDB 3UNI: UcdC, UcdF, and UcdO to chain A.

1254 pTipQC2-*ucdCFO* Plasmid Construction, Purification, and Transformation

Plasmid construction in E. coli NEB10β: Primers flanking the *E. bolteae ucd* operon

(NCBI NZ_CP022464 REGION: complement(4417875..4421605)) were designed in Benchling

using the Primer3 tool. Tails including 6 random bases, followed by restriction sites for NdeI

and XhoI were included on the forward and reverse primers, respectively (Eb_ucdCFO_NdeI_f

and Eb_ucdCFO_XhoI_r). PCR was performed using the Q5 High-Fidelity polymerase (NEB)

with *E. bolteae* DSM 15670 genomic DNA as a template. The target was amplified according

to the following cycling conditions: 30 s at 98 °C, 30 cycles (10 s at 98 °C, 20 s at 60 °C, 80 s

at 72 °C), 2 min at 72 °C. The *ucdCFO* PCR product was purified using the Monarch PCR &

DNA Cleanup Kit (NEB) according to the manufacturer's instructions for products ≥ 2 kb. The

resulting purified PCR product and the pTipQC2 plasmid (Hokkaido Systems Science Co.) were

digested overnight (16 h) with NdeI and XhoI (both from NEB) in rCutSmart buffer according to

the manufacturer's instructions (~600-1000 ng DNA per 50 µL reaction). Double digested DNA

was migrated on a 0.6% agarose gel and bands corresponding to the desired products were

cut out and purified using the Monarch DNA Gel Extraction kit (NEB). The purified products

were ligated using the Hi-T4 DNA Ligase (NEB): a ~3:1 insert:plasmid molar ratio ligation

reaction was set up on ice, then incubated at room temperature for 2 h. The ligation mixture (2

µL) was electroporated (1.8 kV, 25 µF, 200 Ω) into 40 µL electrocompetent *E. coli* NEB10β cells

(according to the Quick-n'-Dirty Electrocompetent *E. coli* protocol

(dx.doi.org/10.17504/protocols.io.bjpykmpw) using 0.1 cm gap cuvettes (Bio-Rad). The cuvette

was immediately filled with 1 mL pre-warmed LB post-shock and cells were allowed to recover

at 37 °C for 30 min before plating on LB + 100 µg/mL ampicillin. After an overnight incubation

at 37 °C, colonies were picked and grown in selective LB + 100 µg/mL ampicillin. Plasmids

were purified using the Plasmid DNA Miniprep Kit (BioBasic) and size was confirmed with a

diagnostic restriction digest (10 µL reactions). The final plasmid construct (pTipQC2-*ucdCFO*)

was submitted to Plasmidsaurus for long-read sequencing using Oxford Nanopore

Technologies (v14 library preparation chemistry, R10.4.1 flow cells) (Supplementary Sequence

2).

pTipQC2-ucdh transformation into Rhodococcus erythropolis DSM 43066:

Electrocompetent *R. erythropolis* DSM 43066 were prepared according to a modified protocol

from P. Lessard 2002. Briefly, 50 mL LB were inoculated with 1 mL of a stationary phase (48-

72 h growth from a single colony) *R. erythropolis* DSM 43066 culture and grown aerobically for

16 h at 30 °C with shaking at 200 RPM. The next day, cells were pelleted at 5,000 x g for 10

min at 4 °C and washed according to the following sequence: 2 washes of (10 mL of ice cold

sterile MilliQ water), 10 mL of ice cold sterile 10% glycerol. The final pellet was then

resuspended in 5 mL of ice cold sterile 10% glycerol. The resuspended electrocompetent *R.*

erythropolis DSM 43066 were aliquoted (50 µL/aliquot), then 3 µL (~0.5-1 µg) of pTipQC2-

ucdCFO plasmid was added to appropriate tubes and incubated for 30 min on ice. Cells with

plasmid were transferred to 0.1 cm gap cuvettes (Bio-Rad) and electroporated (1.8 kV, 25 µF,

200 Ω). Time constants were between 4.3-4.6 ms. The cuvette was immediately filled with 1

mL LB post shock and cells were allowed to recover at 30 °C for 2.5 h before plating 100 µL

dilutions (1/10 dilution, undiluted, and concentrated recovery culture) on LB + 30 µg/mL

chloramphenicol at 30 °C. After 2-3 days of incubation, colonies were picked and grown in

selective liquid LB + 30 µg/mL chloramphenicol at 30 °C with shaking at 200 RPM. Plasmid-

positive colonies were identified by colony PCR using the pTipQC2-*ucdCFO*_cPCR primer set

and validated by diagnostic restriction digests and whole-plasmid sequencing.

Heterologous expression of UcdCFO enzymes in *Rhodococcus erythropolis*:

All growth steps below were performed in selective media (LB + 30 µg/mL chloramphenicol) in aerobic conditions at 30 °C with shaking at 200 RPM, unless otherwise specified. Single colonies of *R. erythropolis* DSM 43066 harboring the pTipQC2 (empty plasmid) or pTipQC2-*ucdCFO* were inoculated into 5 mL selective media and grown for 72 h to produce overnight cultures. Overnight cultures were then thoroughly resuspended and diluted 1:10 into 25 mL fresh selective media and grown for ~8 h until OD_{600} values reached ~0.6. Thiostrepton (5 mg/mL in DMSO) was added to a final concentration of 1 µg/mL and cultures were incubated aerobically for 16 h at 25 °C to induce protein expression. The next morning, cultures were pelleted and resuspended in 0.2 volumes of lysis buffer (20 mM Tris, pH 7.5, 500 mM NaCl, 10 mM $MgSO_4$, 10 mM $CaCl_2$, 2 mM DTT, 1% Triton X-100, 2 mg/mL lysozyme, and 1 tablet/100 mL SIGMAFAST protease inhibitor (EDTA-free)). The resuspended cells in lysis buffer were incubated on ice for 1 h with shaking, then sonicated on ice using a Misonix Sonicator 3000 set to power level 2/10 according to the following sequence (aerobically, in a cold room): 20 s ON, 40 s OFF, for a total of 4 min ON. Crude lysates were transported to the anaerobic chamber in a sealed plastic bag containing an anaerobic gas generating system to minimize loss in activity and treated in the same manner detailed in *Urolithin metabolism assays using crude lysates from uroC-induced E. bolteae*.

SDS-PAGE analysis of UcdCFO proteins in crude lysates

Crude lysates described above (*Heterologous expression of UcdCFO enzymes in Rhodococcus erythropolis*) were centrifuged for 2 min at 20,000 x g. The insoluble pellet was separated from the soluble supernatant. The insoluble pellet (from 100 µL of crude lysate) was resuspended in 100 µL of 1X reducing loading dye (62.5 mM Tris-HCl (pH 6.8), 2% (w/v) SDS, 10% glycerol, 0.01% (w/v) bromophenol blue, 38 mM DTT). The soluble fraction was diluted

with 3X reducing loading dye to a final concentration of 1X. All samples were heated at 95 °C

for 5 min, then 10 µL were loaded onto a 10% bis-tris polyacrylamide protein gel. Gels were

fixed and stained with GelCode Blue Stain Reagent (Fisher) according to the manufacturer's

instructions.

Growth curves with catechols in different iron-containing media conditions

Overnight cultures of *Enterocloster* spp. were treated as described in the *Treatment prior*

to growth (growth curves) sub-section of *Treatments with urolithins and other catechols*.

Once plated, 96-well plates were sealed with a Breathe-Easy membrane and placed in a pre-

warmed plate reader inside the anaerobic chamber (BioTek Epoch 2). The optical density at

620 nm was recorded every 30 min for 48 h. Kinetic analysis was performed in BioTek Gen6

Software using the built-in kinetic analysis.

curatedMetagenomicData meta-analysis of *Enterocloster ucd* operon in human fecal

metagenome datasets

All 93 metagenomic studies (22,588 samples and their metadata) available in the

curatedMetagenomicData R package [53] (v3.8.0) were downloaded locally (ExperimentHub

snapshotDate(): 2023-04-24, accessed on 2023-06-06) and transferred to the Narval cluster

hosted by the Digital Research Alliance of Canada. Metagenomic data for urolithin C-

metabolizing *Enterocloster* spp. were obtained by querying the "relative_abundance" (pre-

processed using MetaPhlAn3 and "gene_families" (pre-processed using HUMAnN3) entries in

individual study datasets [71]. For individual taxa (containing partial strings "bolteae", "citroniae",

"asparagiformis", "asparagiforme", "Enterocloster", or "47FAA" (corresponding to *L.*

pacaense)), relative abundance (%) was extracted from the rows of the "relative_abundance"

datasets using the stringr R package (v1.5.0, https://github.com/tidyverse/stringr). Prevalence

(relative abundance in sample > 0) was then calculated for each sample.

For specific genes, the NCBI protein accessions for each gene of the *ucd* operon (*ucdO, ucdF,*

ucdC) was used to search the UniProt database. UniRef90 accession numbers corresponding

to hits (C5EGQ4, G5HFF3, A8RZR5, respectively) were then extracted from the rows of the

"gene_families" datasets using the stringr R package. Prevalence (abundance in sample > 0)

was then calculated for each sample. R scripts and RData files are available in Zenodo (see

Data Availability).

Fecal slurry preparation and treatment

Frozen (-70 °C) fecal samples were brought into the anaerobic chamber and allowed to

thaw. The samples were suspended in 1 mL mABB medium per 0.1 g feces and homogenized

by breaking apart large pieces with a sterile loop and by vortexing. Large particles were pelleted

by centrifuging the tubes at 700 x g for 3 min. The supernatants (containing bacteria) were

transferred to new tubes and centrifuged at 6,500 x g for 5 min to pellet the cells. The

supernatants were discarded, and the cell pellets were washed with 5 mL of fresh media. The

cell suspensions were once again centrifuged at 6,500 x g for 5 min and the resulting cell pellets

were resuspended in 600 µL media per 0.1 g feces. Resuspended cells were treated with either

100 µM urolithin C or an equivalent volume of DMSO and incubated at 37˚C anaerobically for

48 h. 200-300 µL volumes were removed from the batch cultures and immediately frozen at -

70˚C for later extraction of urolithins (using *Extraction Method A*), DNA, and RNA.

Genomic DNA extraction from fecal slurries

A 300 µL fecal slurry aliquot (between 50-100 mg wet weight) was pelleted (10,000 g for

5 min) and the supernatant was removed for later LC-MS analysis. The pellet was then mixed

with 750 µL of ZymoBIOMICS lysis solution (Zymo Research) and transferred to a ZR

BashingBead lysis tube (Zymo Research). Samples were lysed in a Mini Beadbeater 16

(Biospec) according to the following sequence: 1 min ON, 5 min OFF for a total of 5 min ON.

DNA was then purified using the ZymoBIOMICS DNA Miniprep Kit (Zymo Research) according

to the manufacturer's instructions (including the *OneStep* PCR Inhibitor Removal step). Purified

DNA samples were quantified using the Qubit dsDNA HS assay kit (Invitrogen).

Long read 16S rRNA sequencing of microbial communities in fecal slurries

Long read 16S PCR reactions were conducted using the Platinum SuperFi II Green PCR

Master Mix (Invitrogen). The ONT_16S_27F_GGK and ONT_16S_1492R_GGK primer pairs

(see *Primer sequences table*) were added to their respective master mixes (final primer

concentration of 0.2 µM) and 1 µL of template (~10 ng) was added (for a total reaction volume

of 25 µL). PCR tubes were placed in a thermal cycler and targets were amplified according to

the following cycling conditions: 30 s at 98 °C, 30 cycles (10 s at 98 °C, 10 s at 60 °C, 30 s at

72 °C), 5 min at 72 °C, and hold at 4 °C. Amplicons were quantified using the Qubit dsDNA HS

assay kit (Invitrogen) to verify that amplicon concentrations were reasonably balanced (range

= 18.36-24.00 ng/µL). Barcoding of amplicons was performed with 2 µL of PCR reaction

according to the manufacturer's instructions (for ONT kit SQK-AMB111-24). Barcoding

reactions were incubated in a thermal cycler for 10 min at 65 °C, then for 2 min at 80 °C. 10 µL

of each barcoding reaction were pooled and proteins were digested using heat-labile proteinase

K (NEB) by incubating the pooled library for 15 min at 37 °C, followed by heat inactivation for

10 min at 55 °C. Amplicons were purified using Agencourt AMPure XP beads (Beckman

Coulter) using 0.7 volumes of beads-to-library. Following 70% EtOH washes and drying steps,

the library was eluted using 15 µL of the provided elution buffer (EB), yielding a library with a

concentration of 30 ng/µL using the Qubit dsDNA HS assay kit (Invitrogen). 11 µL of the eluted

DNA library were transferred to a new tube and combined with 1 μL of Rapid Adapter T (RAP

T). This mixture was incubated at room temperature for 10 min. Since the library was

concentrated, it was diluted 1:2 in EB before combining with SB II and LB II, then loaded into a

primed Flongle Flow Cell (R9.4.1) in a MinION device following the manufacturer's instructions.

Sequencing was allowed to proceed for ~20 h until pore exhaustion or enough reads were

obtained. Base calling & demultiplexing was performed using Guppy (v6.4.6) using the "SUP"

super high accuracy model for R9.4.1 flow cells. The raw reads were filtered for a length

between 1500 ± 200 bp. Filtered reads were assigned to taxa using Emu [72] (v3.4.4, GitLab

Project ID: 19618062) by mapping 16S rRNA sequences to the emu_database database

(based on the NCBI 16S RefSeq with the entry for *E. asparagiformis* changed to the sequence

obtained by ONT sequencing (GenBank accession PP280819) since the RefSeq sequence for

this bacterium contained multiple N nucleotides that biased the assignment of *E. asparagiformis*

to *E. lavalensis*). Data were not rarefied or scaled. Count tables were then used to create a

phyloseq (v1.44.0, https://github.com/joey711/phyloseq) object in R [73]. Stacked bar plots were

generated using ggnested (v0.1.0, https://github.com/gmteunisse/ggnested) and fantaxtic

(v0.2.0, https://github.com/gmteunisse/Fantaxtic). Diversity analyses were performed using

Microbiome Analyst (https://www.microbiomeanalyst.ca/) [74].

Total RNA extraction from fecal slurries

A 300 μL fecal slurry aliquot (treated with either 100 μM urolithin C or an equivalent

volume DMSO for 48 h) was thawed and pelleted (6,500 g for 3 min). The pellet (in 200 μL

media) was then mixed with 800 μL TRI reagent (Zymo Research). Samples were lysed in a

Mini Beadbeater 16 (Biospec) according to the following sequence: 1 min ON, 5 min OFF for a

total of 5 min ON. RNA isolation was then performed using the Direct-zol RNA Miniprep Kit

(Zymo Research) according to the manufacturer's instructions (including an on-column DNase

digestion). To ensure complete DNA removal, an additional DNA digestion step was performed

on the isolated RNA using the Ambion DNA-free DNA Removal Kit (Invitrogen) according to the

manufacturer's instructions. The DNA-free RNA was then cleaned up using the OneStep PCR

Inhibitor Removal Kit (Zymo Research). RNA concentration and quality were verified by Qubit

RNA BR assay kit (Invitrogen) and 1% agarose gel electrophoresis.

RT-PCR analysis of the *ucd* operon in fecal slurries

Total RNA was extracted from frozen fecal slurries as previously described (see Total

RNA extraction from microbial communities), and subsequently reverse transcribed as

described above (see *RT-PCR analysis of E. bolteae ucd operon structure*) in a reaction volume

of 5 µL. PCR reactions were conducted using the OneTaq 2X Master Mix with Standard Buffer

(NEB). The ucdCFO_RT-PCR primer pair was added to the master mix (to a final concentration

of 0.2 µM) and 1 µL of template (cDNA, -RT, or no template) was added for a total reaction

volume of 20 µL. PCR tubes were placed in a thermal cycler and targets were amplified

according to the following conditions: 30s at 94˚C, 45 cycles (30s at 94˚C, 1 min at 61˚C, 4 min

at 68˚C), 5 min at 68˚C. A volume of 10 µL of reaction was directly loaded onto a 1% agarose

gel (made with 1X TAE buffer) containing SafeView Classic (Abm). PCR product sizes were

compared to the Quick-Load® Purple 1 kb Plus DNA Ladder (NEB).

PCR analysis of the *ucd* operon prevalence in fecal slurries

Genomic DNA (gDNA) was extracted from frozen fecal slurries as previously described

(see Genomic DNA extraction from microbial communities). PCR reactions and product

visualization was conducted on the gDNA as described above (see RT-PCR analysis of the *ucd*

operon in microbial communities). In this case, 5 µL of PCR product was loaded onto the gels

instead of 10 µL.

Statistical analyses and graphing

1440

Statistical methods were not used to determine sample sizes, experiments were not

randomized, and the investigators were not blinded. Data points related to uroC metabolism

and RT-qPCR were assumed to be normally distributed, though this was not formally tested.

Correlation analyses were performed using the non-parametric Spearman rank correlation (ρ).

Statistical tests on bacterial relative abundances were performed using the Kruskal-Wallis test

on untransformed relative abundance values, which are skewed towards 0. Statistical analyses

for large datasets are detailed in the relevant methods sections. Details related to each test

performed are supplied in the figure legends. In all cases, $\alpha = 0.05$ and tests were two-tailed.

Data were plotted in GraphPad Prism (v10.0.0) or using the ggplot2 (v3.4.2) R package. Figures

were assembled in Affinity Designer (v1.10.6.1665).

Data Availability

RNA-seq reads were deposited in the NCBI SRA BioProject ID PRJNA996126 under BioSample accession codes SAMN36514640 (*Enterocloster bolteae* DSM 15670) and SAMN36514641 (*Enterocloster asparagiformis* DSM 15981). Reviewer link: https://dataview.ncbi.nlm.nih.gov/object/PRJNA996126?reviewer=fpbuj6eeuebv6mij3pme3mp8ep. Untargeted proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD048514 and 10.6019/PXD048514 [75]. Oxford Nanopore 16S rRNA sequencing reads of healthy human fecal slurries were deposited in the NCBI SRA BioProject ID PRJNA1073957. Reviewer link: https://dataview.ncbi.nlm.nih.gov/object/PRJNA1073957?reviewer=nh1a04lg59enlti318qs23s6ot. The 16S rRNA sequence for *E. asparagiformis* DSM 15670 used in the Emu database search was deposited in GenBank under accession PP280819. All original code, tables, and RData files obtained from the analysis of curatedMetagenomicData were deposited in Zenodo (https://doi.org/10.5281/zenodo.8302320).

References

1.   Knight, R. *et al.* The Microbiome and Human Biology. *Annu Rev Genomics Hum Genet* **18**, 65-86 (2017). https://doi.org/10.1146/annurev-genom-083115-022438

2.   Taguer, M. & Maurice, C. F. The complex interplay of diet, xenobiotics, and microbial metabolism in the gut: Implications for clinical outcomes. *Clin Pharmacol Ther* **99**, 588-599 (2016). https://doi.org/10.1002/cpt.366

3.   Almeida, A. *et al.* A unified catalog of 204,938 reference genomes from the human gut microbiome. *Nat Biotechnol* **39**, 105-114 (2021). https://doi.org/10.1038/s41587-020-0603-3

4.   Spanogiannopoulos, P., Bess, E. N., Carmody, R. N. & Turnbaugh, P. J. The microbial pharmacists within us: a metagenomic view of xenobiotic metabolism. *Nat Rev Microbiol* **14**, 273-287 (2016). https://doi.org/10.1038/nrmicro.2016.17

5.   Ervin, S. M. *et al.* Structural Insights into Endobiotic Reactivation by Human Gut Microbiome-Encoded Sulfatases. *Biochemistry* **59**, 3939-3950 (2020). https://doi.org/10.1021/acs.biochem.0c00711

6.   Yao, L. *et al.* A biosynthetic pathway for the selective sulfonation of steroidal metabolites by human gut bacteria. *Nat Microbiol* **7**, 1404-1418 (2022). https://doi.org/10.1038/s41564-022-01176-y

7.   Tomas-Barberan, F. A. & Espin, J. C. Effect of Food Structure and Processing on (Poly)phenol-Gut Microbiota Interactions and the Effects on Human Health. *Annu Rev Food Sci Technol* **10**, 221-238 (2019). https://doi.org/10.1146/annurev-food-032818-121615

8.   Maier, L. *et al.* Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature* **555**, 623-628 (2018). https://doi.org/10.1038/nature25979

9.   Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R. & Goodman, A. L. Mapping human microbiome drug metabolism by gut bacteria and their genes. *Nature* **570**, 462-467 (2019). https://doi.org/10.1038/s41586-019-1291-3

10.   Maini Rekdal, V., Bess, E. N., Bisanz, J. E., Turnbaugh, P. J. & Balskus, E. P. Discovery and inhibition of an interspecies gut bacterial pathway for Levodopa metabolism. *Science* **364**, eaau6323 (2019). https://doi.org/10.1126/science.aau6323

11.   Campbell, C. *et al.* Bacterial metabolism of bile acids promotes generation of peripheral regulatory T cells. *Nature* **581**, 475-479 (2020). https://doi.org/10.1038/s41586-020-2193-0

12.   Paik, D. *et al.* Human gut bacteria produce Tau(Eta)17-modulating bile acid metabolites. *Nature* **603**, 907-912 (2022). https://doi.org/10.1038/s41586-022-04480-z

13.   Cerqueira, F. M., Photenhauer, A. L., Pollet, R. M., Brown, H. A. & Koropatkin, N. M. Starch Digestion by Gut Bacteria: Crowdsourcing for Carbs. *Trends Microbiol* **28**, 95-108 (2020). https://doi.org/10.1016/j.tim.2019.09.004

14.   Maini Rekdal, V. *et al.* A widely distributed metalloenzyme class enables gut microbial metabolism of host- and diet-derived catechols. *Elife* **9** (2020). https://doi.org/10.7554/eLife.50845

15.   Garcia-Villalba, R., Selma, M. V., Espin, J. C. & Tomas-Barberan, F. A. Identification of Novel Urolithin Metabolites in Human Feces and Urine after the Intake of a Pomegranate Extract. *J Agric Food Chem* **67**, 11099-11107 (2019). https://doi.org/10.1021/acs.jafc.9b04435

16.   Carmody, R. N. *et al.* Diet dominates host genotype in shaping the murine gut microbiota. *Cell Host Microbe* **17**, 72-84 (2015). https://doi.org/10.1016/j.chom.2014.11.010

17   Carmody, R. N. *et al.* Cooking shapes the structure and function of the gut microbiome.
Nat Microbiol **4**, 2052-2063 (2019). https://doi.org/10.1038/s41564-019-0569-4
18   Sonnenburg, E. D. *et al.* Diet-induced extinctions in the gut microbiota compound over
generations. *Nature* **529**, 212-215 (2016). https://doi.org/10.1038/nature16504
19   Rothschild, D. *et al.* Environment dominates over host genetics in shaping human gut
microbiota. *Nature* **555**, 210-215 (2018). https://doi.org/10.1038/nature25973
20   Quideau, S., Deffieux, D., Douat-Casassus, C. & Pouysegu, L. Plant polyphenols:
chemical properties, biological activities, and synthesis. *Angew Chem Int Ed Engl* **50**,
586-621 (2011). https://doi.org/10.1002/anie.201000044
21   van Duynhoven, J. *et al.* Metabolic fate of polyphenols in the human superorganism.
Proc Natl Acad Sci U S A **108 Suppl 1**, 4531-4538 (2011).
https://doi.org/10.1073/pnas.1000098107
22   Messaoudene, M. *et al.* A Natural Polyphenol Exerts Antitumor Activity and Circumvents
Anti-PD-1 Resistance through Effects on the Gut Microbiota. *Cancer Discov* **12**, 1070-
1087 (2022). https://doi.org/10.1158/2159-8290.CD-21-0808
23   Pang, S. A., Elkrief, A., Capella, M. P. & Miller, W. H., Jr. Two Cases of Durable and
Deep Responses to Immune Checkpoint Inhibition-Refractory Metastatic Melanoma
after Addition of Camu Camu Prebiotic. *Curr Oncol* **30**, 7852-7859 (2023).
https://doi.org/10.3390/curroncol30090570
24   Garcia-Villalba, R. *et al.* Urolithins: a Comprehensive Update on their Metabolism,
Bioactivity, and Associated Gut Microbiota. *Mol Nutr Food Res* **66**, e2101019 (2022).
https://doi.org/10.1002/mnfr.202101019
25   Singh, A. *et al.* Direct supplementation with Urolithin A overcomes limitations of dietary
exposure and gut microbiome variability in healthy adults to achieve consistent levels
across the population. *Eur J Clin Nutr* **76**, 297-308 (2022).
https://doi.org/10.1038/s41430-021-00950-1
26   Denk, D. *et al.* Expansion of T memory stem cells with superior anti-tumor immunity by
Urolithin A-induced mitophagy. *Immunity* **55**, 2059-2073 e2058 (2022).
https://doi.org/10.1016/j.immuni.2022.09.014
27   Andreux, P. A. *et al.* The mitophagy activator urolithin A is safe and induces a molecular
signature of improved mitochondrial and cellular health in humans. *Nat Metab* **1**, 595-
603 (2019). https://doi.org/10.1038/s42255-019-0073-4
28   Ryu, D. *et al.* Urolithin A induces mitophagy and prolongs lifespan in C. elegans and
increases muscle function in rodents. *Nat Med* **22**, 879-888 (2016).
https://doi.org/10.1038/nm.4132
29   Ghosh, S., Banerjee, M., Haribabu, B. & Jala, V. R. Urolithin A attenuates arsenic-
induced gut barrier dysfunction. *Arch Toxicol* **96**, 987-1007 (2022).
https://doi.org/10.1007/s00204-022-03232-2
30   Singh, R. *et al.* Enhancement of the gut barrier integrity by a microbial metabolite through
the Nrf2 pathway. *Nat Commun* **10**, 89 (2019). https://doi.org/10.1038/s41467-018-
07859-7
31   Shen, P. X. *et al.* Urolithin A ameliorates experimental autoimmune encephalomyelitis
by targeting aryl hydrocarbon receptor. *EBioMedicine* **64**, 103227 (2021).
https://doi.org/10.1016/j.ebiom.2021.103227
32   Ghosh, S., Haribabu, B. & Jala, V. R. Microbial metabolite Urolithin A induces expansion
of Regulatory T Cells in aryl hydrocarbon receptor (AhR)-dependent manner. *Journal of*
Immunology **206**, 113.103-113.103 (2021).

33   Xia, B. *et al.* Urolithin A exerts antiobesity effects through enhancing adipose tissue
thermogenesis in mice. *PLoS Biol* **18**, e3000688 (2020).
https://doi.org/10.1371/journal.pbio.3000688

34   Selma, M. V. *et al.* Isolation of Human Intestinal Bacteria Capable of Producing the
Bioactive Metabolite Isourolithin A from Ellagic Acid. *Front Microbiol* **8**, 1521 (2017).
https://doi.org/10.3389/fmicb.2017.01521

35   Selma, M. V., Beltran, D., Garcia-Villalba, R., Espin, J. C. & Tomas-Barberan, F. A.
Description of urolithin production capacity from ellagic acid of two human intestinal
Gordonibacter species. *Food Funct* **5**, 1779-1784 (2014).
https://doi.org/10.1039/c4fo00092g

36   Gaya, P., Peirotén, A., Medina, M., Alvarez, I. & Landete, J. M. Bifidobacterium
pseudocatenulatum INIA P815: The first bacterium able to produce urolithins A and B
from ellagic acid. *Journal of Functional Foods* **45**, 95-99 (2018).
https://doi.org/10.1016/j.jff.2018.03.040

37   Garcia-Villalba, R. *et al.* Metabolism of different dietary phenolic compounds by the
urolithin-producing human-gut bacteria Gordonibacter urolithinfaciens and Ellagibacter
isourolithinifaciens. *Food Funct* **11**, 7012-7022 (2020).
https://doi.org/10.1039/d0fo01649g

38   Iglesias-Aguirre, C. E. *et al.* Gut Bacteria Involved in Ellagic Acid Metabolism To Yield
Human Urolithin Metabotypes Revealed. *J Agric Food Chem* **71**, 4029-4035 (2023).
https://doi.org/10.1021/acs.jafc.2c08889

39   Iglesias-Aguirre, C. E. *et al.* In vivo administration of gut bacterial consortia replicates
urolithin metabotypes A and B in a non-urolithin-producing rat model. *Food Funct* **14**,
2657-2667 (2023). https://doi.org/10.1039/d2fo03957e

40   Meier-Kolthoff, J. P., Carbasse, J. S., Peinado-Olarte, R. L. & Goker, M. TYGS and
LPSN: a database tandem for fast and reliable genome-based classification and
nomenclature of prokaryotes. *Nucleic Acids Res* **50**, D801-D807 (2022).
https://doi.org/10.1093/nar/gkab902

41   Hosny, M. *et al.* Clostridium pacaense: a new species within the genus Clostridium. *New
Microbes New Infect* **28**, 6-10 (2019). https://doi.org/10.1016/j.nmni.2018.12.003

42   Zupok, A., Iobbi-Nivol, C., Mejean, V. & Leimkuhler, S. The regulation of Moco
biosynthesis and molybdoenzyme gene expression by molybdenum and iron in bacteria.
Metallomics **11**, 1602-1624 (2019). https://doi.org/10.1039/c9mt00186g

43   Leimkuhler, S. & Iobbi-Nivol, C. Bacterial molybdoenzymes: old enzymes for new
purposes. *FEMS Microbiol Rev* **40**, 1-18 (2016). https://doi.org/10.1093/femsre/fuv043

44   Koppel, N., Bisanz, J. E., Pandelia, M. E., Turnbaugh, P. J. & Balskus, E. P. Discovery
and characterization of a prevalent human gut bacterial enzyme sufficient for the
inactivation of a family of plant toxins. *Elife* **7** (2018). https://doi.org/10.7554/eLife.33953

45   Nakashima, N. & Tamura, T. Isolation and characterization of a rolling-circle-type
plasmid from *Rhodococcus erythropolis* and application of the plasmid to multiple-
recombinant-protein expression. *Appl Environ Microbiol* **70**, 5557-5568 (2004).
https://doi.org/10.1128/AEM.70.9.5557-5568.2004

46   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021). https://doi.org/10.1038/s41586-021-03819-2

47   Varadi, M. *et al.* AlphaFold Protein Structure Database: massively expanding the
structural coverage of protein-sequence space with high-accuracy models. *Nucleic
Acids Res* **50**, D439-D444 (2022). https://doi.org/10.1093/nar/gkab1061

48 van Kempen, M. *et al.* Fast and accurate protein structure search with Foldseek. *Nat*
Biotechnol (2023). https://doi.org/10.1038/s41587-023-01773-0

49 Resch, M., Dobbek, H. & Meyer, O. Structural and functional reconstruction in situ of the
[CuSMoO2] active site of carbon monoxide dehydrogenase from the carbon monoxide
oxidizing eubacterium Oligotropha carboxidovorans. *J Biol Inorg Chem* **10**, 518-528
(2005). https://doi.org/10.1007/s00775-005-0006-4

50 Ishikita, H., Eger, B. T., Okamoto, K., Nishino, T. & Pai, E. F. Protein conformational
gating of enzymatic activity in xanthine oxidoreductase. *J Am Chem Soc* **134**, 999-1009
(2012). https://doi.org/10.1021/ja207173p

51 Kramer, J., Özkaya, Ö. & Kümmerli, R. Bacterial siderophores in community and host
interactions. *Nature Reviews Microbiology* **18**, 152-163 (2020).
https://doi.org/10.1038/s41579-019-0284-4

52 Pals, M. J., Wijnberg, L., Yildiz, C. & Velema, W. A. Catechol-Siderophore Mimics
Convey Nucleic Acid Therapeutics into Bacteria. *Angew Chem Int Ed Engl* **n/a**,
e202402405 (2024). https://doi.org/10.1002/anie.202402405

53 Pasolli, E. *et al.* Accessible, curated metagenomic data through ExperimentHub. *Nat*
Methods **14**, 1023-1024 (2017). https://doi.org/10.1038/nmeth.4468

54 Cortes-Martin, A. *et al.* The gut microbiota urolithin metabotypes revisited: the human
metabolism of ellagic acid is mainly determined by aging. *Food Funct* **9**, 4100-4106
(2018). https://doi.org/10.1039/c8fo00956b

55 Bess, E. N. *et al.* Genetic basis for the cooperative bioactivation of plant lignans by
Eggerthella lenta and other human gut bacteria. *Nat Microbiol* **5**, 56-66 (2020).
https://doi.org/10.1038/s41564-019-0596-1

56 Le, C. C., Bae, M., Kiamehr, S. & Balskus, E. P. Emerging Chemical Diversity and
Potential Applications of Enzymes in the DMSO Reductase Superfamily. *Annu Rev*
Biochem **91**, 475-504 (2022). https://doi.org/10.1146/annurev-biochem-032620-110804

57 Little, A. S. *et al.* Dietary- and host-derived metabolites are used by diverse gut bacteria
for anaerobic respiration. *Nat Microbiol* **9**, 55-69 (2024). https://doi.org/10.1038/s41564-
023-01560-2

58 Zhang, X. *et al.* Isolation and characterization of a novel human intestinal Enterococcus
faecium FUA027 capable of producing urolithin A from ellagic acid. *Front Nutr* **9**,
1039697 (2022). https://doi.org/10.3389/fnut.2022.1039697

59 Pauff, J. M., Cao, H. & Hille, R. Substrate Orientation and Catalysis at the Molybdenum
Site in Xanthine Oxidase: CRYSTAL STRUCTURES IN COMPLEX WITH XANTHINE
AND LUMAZINE. *J Biol Chem* **284**, 8760-8767 (2009).
https://doi.org/10.1074/jbc.M804517200

60 Cozza, G. *et al.* Urolithin as a Converging Scaffold Linking Ellagic acid and Coumarin
Analogues: Design of Potent Protein Kinase CK2 Inhibitors. *ChemMedChem* **6**, 2273-
2286 (2011). https://doi.org/10.1002/cmdc.201100338

61 Meier-Kolthoff, J. P., Goker, M., Sproer, C. & Klenk, H. P. When should a DDH
experiment be mandatory in microbial taxonomy? *Arch Microbiol* **195**, 413-418 (2013).
https://doi.org/10.1007/s00203-013-0888-4

62 Meier-Kolthoff, J. P. & Goker, M. TYGS is an automated high-throughput platform for
state-of-the-art genome-based taxonomy. *Nat Commun* **10**, 2182 (2019).
https://doi.org/10.1038/s41467-019-10210-3

63 Ewels, P., Magnusson, M., Lundin, S. & Kaller, M. MultiQC: summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047-3048 (2016).
https://doi.org/10.1093/bioinformatics/btw354

64   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
1655   reads. *EMBnet.journal* **17**, 10 (2011). https://doi.org/10.14806/ej.17.1.200
1656   65   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
1657   alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907-
1658   915 (2019). https://doi.org/10.1038/s41587-019-0201-4
1659   66   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
1660   assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
1661   https://doi.org/10.1093/bioinformatics/btt656
1662   67   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
1663   for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
1664   https://doi.org/10.1186/s13059-014-0550-8
1665   68   Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative C(T)
1666   method. *Nat Protoc* **3**, 1101-1108 (2008). https://doi.org/10.1038/nprot.2008.73
1667   69   Keller, A., Nesvizhskii, A. I., Kolker, E. & Aebersold, R. Empirical statistical model to
1668   estimate the accuracy of peptide identifications made by MS/MS and database search.
1669   *Anal Chem* **74**, 5383-5392 (2002). https://doi.org/10.1021/ac025747h
1670   70   Nesvizhskii, A. I., Keller, A., Kolker, E. & Aebersold, R. A statistical model for identifying
1671   proteins by tandem mass spectrometry. *Anal Chem* **75**, 4646-4658 (2003).
1672   https://doi.org/10.1021/ac0341261
1673   71   Beghini, F. *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse
1674   microbial communities with bioBakery 3. *Elife* **10**, e65088 (2021).
1675   https://doi.org/10.7554/eLife.65088
1676   72   Curry, K. D. *et al.* Emu: species-level microbial community profiling of full-length 16S
1677   rRNA Oxford Nanopore sequencing data. *Nat Methods* **19**, 845-853 (2022).
1678   https://doi.org/10.1038/s41592-022-01520-4
1679   73   McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive
1680   analysis and graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).
1681   https://doi.org/10.1371/journal.pone.0061217
1682   74   Lu, Y. *et al.* MicrobiomeAnalyst 2.0: comprehensive statistical, functional and integrative
1683   analysis of microbiome data. *Nucleic Acids Research* **51**, W310-W318 (2023).
1684   https://doi.org/10.1093/nar/gkad407
1685   75   Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass
1686   spectrometry-based proteomics evidences. *Nucleic Acids Res* **50**, D543-D552 (2022).
1687   https://doi.org/10.1093/nar/gkab1038

1688

Acknowledgements

This research was funded by the Canadian Institutes of Health Research (CIHR) grant PJT-437944 to B.C. and C.F.M. and a Weston Family Microbiome Initiative's 2021 Transformational Research program to B.C. B.C. holds a tier II Canada Research Chair (CRC) in Therapeutic Chemistry. C.F.M. holds a tier II CRC in Gut Microbial Physiology and is a Azrieli Global Scholar in the Humans & Microbiome program. R.P. is supported by the CIHR Canada Graduate Scholarship-Doctoral (#493808 for 2023-2026) and by the Fonds de Recherche du Québec-Santé : Bourse de formation au doctorat (#316063 for 2022-2023). M.S. is supported by the CIHR Canada Graduate Scholarship to Honor Nelson Mandela (#DF2-187718 for 2023-2026), the Fonds de Recherche du Québec-Santé: Bourse de formation au doctorat (#311071 for 2022-2023).

RNA sample preparation and sequencing was performed by Génome Québec. Untargeted proteomics sample preparation and analysis was performed by the Research Institute of the McGill University Health Centre (RI-MUHC) Proteomics and Molecular Analysis Platform. Preliminary small-molecule MALDI-TOF mass spectrometry was performed by Mark Hancock at the McGill SPR-MS Facility. High resolution mass spectrometry was performed by the McGill Chemistry Characterization Facility. Molecular biology experiments were enabled by the McGill University Imaging and Molecular Biology Platform (IMBP). High-performance cloud computing was enabled by Calcul Québec (https://www.calculquebec.ca/) and the Digital Research Alliance of Canada (https://alliancecan.ca/). RNA-seq data analysis was performed using two instances of Galaxy: (https://usegalaxy.org/) and Compute Canada Genetics and Genomics Analysis Platform (GenAP) (https://www.genap.ca/).

Author information

Authors and Affiliations

Department of Pharmacology & Therapeutics, McGill University, 3655 Prom. Sir-William-Osler,

Montreal, Quebec, H3G 1Y6, Canada

Reilly Pidgeon, Sacha Mitchell, Layan Suleiman, Lharbi Dridi & Bastien Castagner

Department of Microbiology & Immunology, McGill University, 3775 University Street, Montreal,

Quebec, H3A 2B4, Canada

Michael Shamash & Corinne F. Maurice

Contributions

R.P. designed the study, performed microbiology experiments, performed bioinformatics analyses, analyzed data, created figures, and wrote the initial manuscript with B.C. **S.M.** performed experiments on microbial communities, analyzed data, and created figures. **M.S.** performed Nanopore sequencing and processed raw sequencing reads. **L.S.** performed microbiology experiments. **L.D.** consulted on experimental design and methodology. **C.M.** supervised research, obtained ethical approval for the use of human fecal samples, and obtained research funding. **B.C.** designed the study, supervised the research, obtained research funding, and wrote the initial manuscript with R.P. All authors reviewed and edited the manuscript.

Corresponding authors

Correspondence to Bastien Castagner (bastien.castagner@mcgill.ca).

Ethics declarations

The authors declare no competing interests.