Ungulate substrate selection in fauna passages

Jan Olof Helldin[1]* & Milla Niemi[2]

1) SLU Swedish Biodiversity Centre, Dept of Urban and Rural Development, Swedish University of Agricultural Sciences, Uppsala, Sweden

2) Latvasilmu osk, Hankasalmi, Finland

*) Corresponding author: j-o.helldin@slu.se

Abstract

Fauna passages are increasingly constructed at major roads and railways to mitigate the negative effects of infrastructure and traffic on wildlife. The function of such passages depends on their design, including the construction materials, soil and vegetation. Providing "naturalness" in fauna passages may entail significant costs, yet its benefits are incompletely understood. In particular, there is a demand for better knowledge support regarding combining passages for large fauna with minor roads, trails and watercourses. By using camera trap data collected across the passages, we examined ungulate substrate selection in seven passages serving both fauna and local roads (overpasses) and fauna and watercourses (underpasses) in boreal Sweden and Finland. While all substrates were used, during snow-free periods, ungulates used smoother surfaces (fine-grained topsoil, grass, artificial fiber mat, and dirt road) more than expected based on their availability. Coarser surface (stony/rocky ground), shrub, and water were used less than expected. The results for road and water were however inconsistent between passages; in one overpass road was instead used less than expected, and in one underpass the water section was used particularly during winter (ice cover) but also by moose wading or swimming through in summer and autumn. The general patterns of use largely remained when we analysed data on species level, although these analyses were restricted because of limited sample sizes. While our study has several limitations, it offers valuable insights for the planning and construction of fauna passages. To our knowledge, this study was the first of its kind to study the substrate preferences of ungulates in fauna passages, and we suggest to conduct further research in the field.

Introduction

Transport agencies worldwide are increasingly constructing large fauna passages at major roads and railways to reduce both wildlife-vehicle collisions and the barrier effects of these transportation routes on wildlife (Iuell et al. 2003; Grilo et al. 2010; Smith et al. 2015a; Hlavac et al. 2019; Denneboom et al. 2021). Such fauna passages may be vegetated bridges, viaducts or culverts of different sizes, construction materials and shapes, but have in common that they are constructed or adapted to meet the requirements of the target species or taxa. Many practical guidelines exist to aid the construction of fauna passages (e.g., Clevenger & Ford 2010; Queensland Government 2010; Smith et al. 2015b; Hlavac et al. 2019; Chrétien et al. 2022; see also national guidelines accessed thought Transport Ecology Guidelines Portal 2024). Most of these advocate for, *inter alia*, large openness (maximizing width and height for underpasses, while minimizing length), using natural materials and substrate, providing shelter and connected habitat structures through the passage, screening from traffic disturbance, as well as minimizing other human disturbances.

While the dimensions of the constructions is generally the most cost-driving factor for fauna
passages (Sijtsma et al. 2020, Helldin 2022), also adaptations to create "naturalness" in the passage
may entail significant costs, for example to plan and establish vegetation and natural soil conditions.
The degree to which non-wildlife structures such as roads, hiking trails or watercourses can be
accepted in fauna passages may also be a cost-driving factor as it impacts decisions to construct
passages with a combined function for fauna and humans or fauna and drainage, and therefore
decisions of total number of bridges and fauna passages needed.

Regarding the ground surface or substrate in large fauna passages, the current Swedish construction
guideline (Trafikverket 2022) recommend natural vegetation on a bed of natural soil. Minor unpaved
roads with limited traffic can be accepted, and a dry bank ≥20 cm above high water level should be
kept along watercourses. Planners however still express uncertainties about combining passages for
larger fauna with local roads or streams, due to the risk of human disturbance and of losing effective
passage width to road or water. For ungulates (*Artiodactyla*), which are the main target taxon for
larger fauna passages in northern Europe, results of avoidance or preference of certain substrates in
fauna passages are sparse and inconclusive (Denneboom et al. 2021). To assess the function for
ungulates of non-wildlife over- and underpasses in Sweden, Seiler et al. (2015) proposed to exclude
road and water surfaces when calculating the effective passage width, thereby implying that these
surfaces would not be used by ungulates.

The aim of this study was to describe the preferences of ungulates for substrates in fauna passages,
as revealed by the animal trajectories through the passage. Particular focus was put on the potential
avoidance of roads and streams going through the passage. Studies were conducted year-round at
three underpasses (viaducts) and four overpasses (green bridges) in boreal Sweden and Finland.
While these passages may be used by many different species, we perceive that the passages were
primarily adapted to meet the requirements by ungulates. While acknowledging some limitations of
our study, we discuss how the results may guide future construction of passages built or adapted for
boreal ungulates.

Method

Fauna passages included in the study
We selected seven fauna passages to include in the study (Table 1), out of a total of ca 35 under- and
overpasses that were monitored for ungulate use within larger research projects in Sweden
(Håkansson 2020, Raud Westberg & Ellvin 2021) and Finland (Niemi 2021) during 2018–2022. The
selection was arbitrarily based on the following criteria:
• Minimum width 20 m and containing a variety of substrates (≥2), to allow animals to actively
select substrate.
• Monitored ≥1 year.
• Camera trap set up allowed equal (un-biassed) coverage of the entire passage width.
In one case, however, a bridge was included (Loviisa) despite a 5m strip behind a railing not being
reliably covered by the camera. We judged the strip being of less interest, and limited the analyses to
the rest of the passage width.

By this selection, four passages (three viaducts and one overpass) along the Haparandabanan railway
in northern Sweden and three over passages along motorway E18 in southern Finland were included
in the analyses (Fig. 1). All seven passages are defined by the respective infrastructure administration
as multifunctional fauna passages, i.e., the intended function is for both fauna and any other purpose
(in these cases watercourses or local, low-traffic roads, and including recreational use).

Study areas

The study area in northern Sweden is situated at 65°N 23°E, in northern boreal zone dominated by a mix of forest and bogs, and with stable winter conditions (snow layer and freezing temperatures) during at least December-March. Ungulate species that occur in the area are moose (*Alces alces*), roe deer (*Capreolus capreolus*) and reindeer (*Rangifer tarandus tarandus*); the latter being semi-domestic but free-ranging most of the year. The study area in southern Finland is situated at 60°N ~25°E, in southern boreal zone dominated by a mix of forest and farmland, with winter conditions being more erratic. Ungulate species occurring in this area are moose, roe deer, white-tailed deer (*Odocoileus virginianus*; non-native) and wild boar (*Sus scrofa*).

Camera trapping

We monitored ungulates in the passages using motion-triggered cameras (Browning 2017 Spec Ops Advantage Trail in the Swedish passages, Uovision UM785-3G in the Finnish) with infrared (IR) motion sensor, IR flash and IR night vision, and with the detection range of motion sensor and flash specified by manufacturer as minimum 16 m (Uovision) or 24 m (Browning). We set the cameras to rapid fire 3-5 images when being triggered, thereafter with a delay of 5 sec (Browning) or 60 sec (Uovision) to the next triggering.

We mounted the cameras in the middle of the passages (some of the passages also had cameras distributed outside the passage, but these cameras where not used in the present study), at approximately 1 m above ground, attached to pre-existing fence poles or in few cases new poles. We placed the cameras at one or both sides of the passage and facing the passage center (Fig. 2), and in one case we also mounted cameras near a center pillar and facing the sides, accordingly aiming at covering the entire width of the passage. We did some careful vegetation cutting in front of cameras to minimize false "wind triggers". We revisited cameras once per 1–3 months in order to check functionality and download images. For the aim of this study, ungulate use was recorded for one full year (Table 1).

Image handling

We combined all consecutive images of ungulates at a passage within a period of 10 min into one crossing event. We used only events that we judged showing animals successfully crossing the passage, i.e., coming in from one side and leaving towards the other. For each event, we noted the ungulate species, the summed number of individuals, date and time, snow cover (Y/N) and direction of movement (south/north). We excluded a small number of crossing events where species could not be well recognized (roe/white-tailed deer; Finnish data). For each first crossing individual in one event ("the group leader"), we recorded the substrate where it crossed a predefined imaginary line over the passage in front of the cameras (record line; Fig. 2A–B). We classified substrates into six types:

- Grassy/sandy (spontaneous herbaceous or graminoid vegetation on sandy soil)
- Stony/rocky (sometimes with sparse herbaceous vegetation)
- Fiber mat (artificial coco liner ca 5 cm thick, rolled out on stony/rocky substrate)
- Shrub (woody vegetation ca 0.5-2 m height, mainly with ground covered by low vegetation)
- Water (river or stream)
- Road (local gravel/dirt road)

Substrates grassy/sandy and road were only represented in the overpasses while stony/rocky, fiber mat and water were only represented in the underpasses (Table 2).

As a complement, we mapped each trajectory on a spatial map (Fig. 2A–B).

Data analyses

We compared the substrates used by ungulates (when crossing the imaginary line) with available
substrates using a chi-square (χ^2) goodness-of-fit analysis conducted in Excel, separate for each fauna
passage and season (snow-free/snow covered ground), and separate for each ungulate species plus
all species taken together. For the sake of liability of results (minimum sample size), we excluded
analyses with any cell with an expected value lower than 5 (Siegel & Castellan 1988). For all
significant analyses ($p < 0.05$), we recorded the relative contribution (proportion of total χ^2) of each
substrate type, and pointed out preferences (observed > expected) and avoidances (observed <
expected).

For the particular focus on watercourses and roads through the passage, we made additional chi-
square goodness-of-fit analyses of these two substrates compared to other substrates pooled,
separate for each fauna passage and season (snow-free/snow covered ground), for each ungulate
species plus all species taken together, and using the same analysis exclusion criterion (i.e., any cell
with an expected value lower than 5).

We separated seasons by records of snow covered ground on images, which in the Swedish passages
corresponded to mid-October to early May (ca 200 days) and in the Finnish a number of shorter
periods occurring from end of November through early April (a total of <15 days).

Results

A total of 983 ungulate crossing events were recorded in the study (Table 3), with moose and roe
deer recorded in all passages, reindeer only in the Swedish, white-tailed deer and wild boar only in
the Finnish; wild boar only at very few occasions. A majority of crossing events (719) were from the
snow-free season. From the period of snow covered ground, most (90%) crossing events were from
moose or reindeer. Only the Swedish passages allowed analyses from snowy period.

Although all types of substrates were used, the extent varied, and ungulates displayed a non-random
use of substrate in all analyses (Table 4), except for the passage Loviisa.
- In the **snow-free season** (Table 4A), the general pattern was that fiber mat and grassy/sandy
ground (only available in underpasses) and road (only in overpasses) were used more than
randomly; this was most pronounced for fiber mat (large contribution to χ^2). Conversely, water
and stony/rocky ground (in underpasses) and shrub (available in both under- and overpasses)
was used less than randomly, however not well pronounced for stony/rocky ground (≤5%
contribution to χ^2). For road, the results were inconsistent, with much more use than random in
two passages but much less in one.
When analyzed at the species level, these general patterns mainly persisted. However, roe deer
and reindeer utilized stony/rocky ground more than randomly in the Kvarnbäcken passage (still
with low contribution to χ^2).
- In the period of **snow covered ground** (and, notably, also ice covered water), water and road
were used more than randomly, while stony/rocky ground, fiber mat and shrub were used less
than randomly (Table 4B). However, only two passages allowed analysis for snowy season – the
underpass at Kvarnbäcken and the overpass at Sangijärvi – and the result was to a large extent
attributed to moose in the former case and to reindeer in the latter.

The particular analyses of water and road, respectively, allowed analysing a few more combinations
of passages and species (Table 5A–B), and largely underlined results above.
- **Water** was used less than randomly during snow-free (and ice-free) season, in all cases except for
moose in one case (Kvarnbäcken) where water was used according to availability. In the period

with snow and ice, the use of water differed between passages; more than randomly in one case
(Kvarnbäcken) and less in the other two cases (Aavajoki and Keräsjoki).

• For **road**, the results remained inconsistent, with road used more than randomly in two passages
(Sangijärvi and Sammatti, for the former also with snow-covered ground) but less in one
(Kärmekorpi) and according to availability in one (Loviisa).

Discussion

Our study indicated that fine-grained topsoil, grass, and fiber mat were preferred substrates for
ungulates in fauna passages during snow-free conditions (Fig. 3A–B). Although we can only speculate
about the reason for this selection, we acknowledge that these surfaces are smoother, and may be
perceived by animals as more comfortable for walking or running. It has been argued that fine-
grained soil and smooth surfaces should be used in fauna passages, rather than coarse materials such
as macadam or riprap (Smith et al. 2015b, Trafikverket 2022). In line with that, we found that
ungulates used coarser surfaces (Fig. 3C) slightly less than expected based on its availability. This was
true also in snowy conditions.

Smooth and easy-to-walk-on surface may apply also to road (local gravel or dirt road), which was
strongly selected for in two of the overpasses in our study (of which one included data also from
snowy conditions; Fig. 3D), and used according to availability in one. The analyses on species level
showed that this selection of roads was attributed to roe deer, reindeer, and white-tailed deer. In the
overpass where road was instead avoided, the alternative substrate was grassy/sandy, i.e., another
smooth surface, but also the site where moose was the most frequent species. It has been proposed
that ungulates (reindeer) may use minor, low-traffic roads as transportation corridors (e.g. Strand et
al. 2018). Although some studies indicate that ungulates in forest landscapes avoid minor roads
(Laurian et al. 2008, Mathisen et al. 2018), this is likely due more to higher risk of hunting mortality
and predation by large carnivores than to the character of the road surface.

Furthermore, the results indicated that shrub as a substrate (or habitat) in fauna passages was
avoided by ungulates, although shrub should appear natural and provide shelter and even food
(depending, of course, on species), and although shrub in the current cases forms more or less
continuous habitat corridors through these passages and thereby connects habitats across the
infrastructure. In contrast to shrub, all other solid-ground substrates (grassy/sandy, stony/rocky,
fiber mat, road) are more or less open habitats, and may be preferred for that reason.

As a general conclusion, we propose that ungulates may require overview rather than shelter when
crossing through fauna passages, and accordingly avoid denser habitats such as shrub. Moving
through a fauna passage is likely to be an aggravated situation for wildlife, with an elevated risk of
predation or strife (Mata et al. 2015), and open areas near passages may facilitate predator
avoidance and escape (Clevenger & Waltho 2005, Denneboom et al. 2021). However, we cannot
exclude the possibility that the shrub in our cases may have provided shelter from the side (for
example from traffic disturbance) or functioned as a guiding habitat structure, and therefore may
have served a purpose for ungulates even when not used directly.

Water was generally avoided by ungulates in fauna passages, but with the prominent exception of
Kvarnbäcken underpass where moose used the water section year-round and reindeer in winter (Fig.
4A–C). In this passage, much of its span is dedicated to a gently flowing, deep but narrow river (ca 10
m wide). In winter, the river froze and the ice created a solid floor used by the vast majority of moose
and reindeer that crossed in the six winter months (November-April). However, also in the rest of the
year, moose used the river for crossing, either by wading along the river bank or swimming through

the deeper part. We recorded the same behaviors in the other two underpasses, however less
frequently. Interestingly, as far as we could observe (by additional records from cameras outside of
the passages), in all these occasions moose approached on land, went into the river some 10 m
before the passage, swam or waded through, and got up again on land shortly after crossing.

Our study had a number of limitations, related to the study setup and confounding factors, so
conclusions must be drawn with care. We could only include a limited number of fauna passages in
our study. These passages varied in type, size and substrate composition, and were located in two
areas with different species abundance. Therefore, the observed patterns might be influenced by
these local circumstances and may not necessarily be universal. Also other structural factors than
substrate could potentially have impacted the trajectory along which ungulates crossed through the
passages, for example if they approached and entered the passage along preferred guiding structures
in the surroundings (for example perimeter fencing or established wildlife trails), or if they avoided
moving in the direct proximity to technical elements in the passage such as the concrete bridge
abutments.

We also acknowledge the risk of pseudo-replication (non-independent data) within passages,
deriving from a smaller number of individuals returning to a passage, and therefore that results may
reflect individual choices or habits. We did not systematically record individuals but find it most likely
that local residents use fauna passages repeatedly, and we could incidentally recognize a few
returning individual roe deer bucks by their antler shape and reindeer by their fur patterns. Another
aspect to pseudo-replication is that we expect ungulates to follow each other´s trails based on scent
and tracks, probably even after much longer period than the 10 min we set to define a separate
crossing event.

Despite these limitations, our work provides insight in a topic on which surprisingly little research has
been conducted. Given the significant implications for planning and costs associated with fauna
passages, there's a clear need for more applied research in this area. The importance of substrate
and microhabitats in fauna passages have been studied for smaller vertebrate species such as small
mammals, reptiles and amphibians (Georgii et al. 2011, Connolly-Newman 2013, Andrews et al.
2015), but much less so for large mammals (Denneboom et al. 2021). While authors have stressed
the importance for ungulates of vegetation, natural soil, level ground surface, and dry pathways
along watercourses through fauna passages, this has usually been addressed by analyzing crossing
rates vs. crossing failures (Smith 2003, Eco-Kare International 2020, Denneboom et al. 2021), hence
comparisons on a between-passage level. Our study may serve as a complement to these previous
studies, by illustrating also how ungulates use, and possibly select, substrates within passages.

Conclusions and recommendations

Our results largely support the current, broad recommendation to provide natural soil and
vegetation in fauna passages (Väre et al. 2003, Chrétien et al. 2022, Rosell et al. 2022, Trafikverket
2022). However, we point out some opportunities for clarifications on this matter. The results
indicate that open habitats are more important for ungulates than previously acknowledged, and
accordingly, plantations should be used restrictively in the passages and encroaching woody
vegetation may need to be managed. We are aware that dense vegetation in fauna passages may
benefit smaller species and thus make the passages functional for a broader array of species. This
should, however, not be at the cost of the effectiveness for the primary target species. More
emphasis may be put in the guidelines on the value of a smooth ground surface, such as that created
by fine-grained topsoil and fiber mats, and on the potential value of minor roads with low traffic, as
they can function as conduits of ungulate movement. We also recommend to consider the function

of watercourses for some ungulate species, not only as a large scale guiding structure but also as a
unique substrate providing the opportunity to wade or swim through the passage, and to walk
through in winter in regions with stable winter cold.

All these conclusions are tentative, and we hope that our study will open up for further research of
vegetation, soil and structures in and around fauna passages. We suggest to plan study setups that
allow recording of entire trajectories of animals through the fauna passage, taking into consideration
the substrates, guiding structures and microhabitats both in the passages and in their direct vicinity;
i.e., in the area planned and managed by the road or railway agency. We find this field particularly
suitable for experimental analyses, for example by modifying substrates within a passage and
monitoring any changes in the use by animals. Further study of watercourses and minor roads should
aim at sorting out whether these facilitate or obstruct animal movements, since both our results and
previous research (as reviewed in Denneboom et al. 2021) are inconclusive on these points.

Acknowledgements

275 We are grateful to the many colleagues and assistants involved in data collection and analysis (Ida
276 Anomaa, Christine Godeau, Charlotte Hansson, Emma Håkansson, Jan-Erik Innala, Victor Johansson,
277 Fabian Knufinke, Torbjörn Nilsson, Lars-Gunnar Nyström, Mattias Olsson, Andreas Seiler, Andreas
278 Öhlund). The study was financed by the Swedish Transport Administration (research project TRIEKOL;
279 https://triekol.se/) and the Finnish Transport Infrastructure Agency.

Data supplements

281 Data supplement no. 1: Crossing events used in analyses.
282 Data supplement no. 2: Spatial maps of individual trajectories separated by site, species and season.

References

Andrews K, Langen TA, Struijk RPJH (2015) Reptiles: overlooked but often at risk from roads. Chapter
32 (pp. 271-280) in van der Ree R, Smith DJ, Grilo C (eds.) Handbook of Road Ecology. John
Wiley & Sons, Ltd.

Clevenger AP, Waltho N (2005) Performance indices to identify attributes of highway crossing
structures facilitating movement of large mammals. Biological Conservation 121: 453–464.
https://doi.org/10.1016/j.biocon.2004.04.025

Clevenger AP, Ford AT (2010) Wildlife crossing structures, fencing, and other highway design
considerations. In: Beckman JP, Clevenger AP, Huijser MP, Hilty JA (eds.) Safe passages:
highways, wildlife, and habitat connectivity. Island Press, Washington DC, 17–49.

Connolly-Newman HR (2013) Effects of cover on small mammal abundance and movement through
wildlife underpasses. Graduate Student Theses, Dissertations, & Professional Papers 350,
University of Montana, USA.

Chrétien L, Guinard E, Nowicki F, Righetti A, Rosell C, Trocmé M, Fernández LM, Böttcher M, Eicher C,
Elstrom M, Morand A, Petrovan S., Schwab T, Suter S, Zumbach S (2022) Solutions to mitigate
transport infrastructure impacts on wildlife. Chapter 7 update of: Iuell et al. (eds.) IENE Wildlife
and Traffic. A European Handbook for Identifying Conflicts and Designing Solutions. KNNV
Publishers, Brussels.

Denneboom D, Bar-Massada A, Shwartz A (2021) Factors affecting usage of crossing structures by wildlife – A systematic review and meta-analysis. Science of the Total Environment 777: 146061.

Eco-Kare International (2020) Effectiveness monitoring of wildlife mitigation measures for large- and mid-sized animals on Highway 69 in Northeastern Ontario: September 2011 to September 2019. Submitted to the Ontario Ministry of Transportation, North Bay, Ontario, Canada, 73 pp.

Georgii B, Keller V, Pfister HP, Reck H, Peters-Ostenberg E, Henneberg M, Herrmann M, Mueller-Stiess H, Bach L (2011) Use of wildlife passages by invertebrate and vertebrate species. Wildlife passages in Germany 2011.

Grilo C, Bissonette JA, Cramer PC (2010) Mitigation measures to reduce impacts on biodiversity. Pp 73-114 in: Jones SR Highways: Construction, Management, and Maintenance. Nova Science Publishers, Inc.

Håkansson E (2020) Effectivity of road and railway crossing structures for wild mammals. MSc thesis, Dept. of Biological and Environmental Sciences, Gothenburg University. https://triekol.se/project/effectivity-of-road-and-railway-crossing-structures/

Helldin J.O. (2022) Are several small wildlife crossing structures better than a single large? Arguments from the perspective of large wildlife conservation. Nature Conservation 47: 197-213. https://natureconservation.pensoft.net/article/67979/

Hlaváč V, Anděl P, Matoušová J, Dostál I, Strnad M, Immerová B, Kadlečík J, Meyer H, Moţ R, Pavelko A, Hahn E, Georgiadis L (2019) Wildlife and Traffic in the Carpathians. Guidelines how to minimize impact of transport infrastructure development on nature in the Carpathian countries. Danube Transnational Programme TRANSGREEN Project, The State Nature Conservancy of the Slovak Republic.

Iuell B, Bekker HGJ, Cuperus R, Dufek J, Fry G, Hicks C, Hlaváč V, Keller V, Rosell C, Sangwine T, Tørsløv N, Le Maire Wandall B (2020) COST 341–Wildlife and Traffic: a European handbook for identifying conflicts and designing solutions. KNNV Publishers, Brussels.

Laurian C, Dussault C, Ouellet J-P, Courtois R, Poulin M, Breton L (2008) Behavior of Moose Relative to a Road Network. Journal of Wildlife Management, 72(7) : 1550-1557 https://doi.org/10.2193/2008-063

Mata C, Bencini R, Chambers BK, Malo JE (2015) Predator-prey interactions at wildlife crossing structures: Between myth and reality. In: van der Ree R, Smith DJ, Grilo C (Eds) Handbook of Road Ecology. John Wiley and Sons Ltd, UK, 190–197.

Mathisen KM, Wójcicki A, Borowski Z (2018) Effects of forest roads on oak trees via cervid habitat use and browsing. Forest Ecology and Management 424: 378–386.

Niemi M (2021) Vihersillat eläinten kulkureittinä tien yli - Eläinyhteyksien riistakameramonitorointi [Green bridges as crossing structures for animals]. Publications of the FTIA 26/2021, Finnish Transport Infrastructure Agency, Helsinki. 56 pp. https://urn.fi/URN:ISBN:978-952-317-864-9

Queensland Government (2010) Fauna sensitive road design manual Vol. 2. https://www.tmr.qld.gov.au/business-industry/technical-standards-publications/fauna-sensitive-road-design-volume-2

Raud Westberg A & Ellvin A (2021) Uppföljning av viltets användning av broar vid E4 Sundsvall – med särskilt fokus på smala broar och mänskliga störningar. Report from Swedish Transport Administration dated 2021-10-24. https://triekol.se/project/uppfoljning-av-viltets-anvandning-av-broar-vid-e4-sundsvall/

Rosell C, Torrellas C, Colomer J, Reck H, Navàs F, Bil M (2020) Maintenance of ecological assets on transport linear infrastructure. Chapter 10 update of: Iuell et al. (eds.) IENE Wildlife and Traffic. A European Handbook for Identifying Conflicts and Designing Solutions. KNNV Publishers, Brussels.

Seiler A, Olsson M, Lindqvist M (2015) Deficiencies in permeability of roads and railroads for ungulates [Analys av infrastrukturens permeabilitet för klövdjur]. Swedish Transport Administration publication 2015:254.

Siegel S & Castellan NJ Jr (1988) Nonparametric statistics for the behavioral sciences (2nd ed.). Mcgraw-Hill Book Company.

Sijtsma FJ, van der Veen E, van Hinsberg A, Pouwels R, Bekker R, van Dijk R, Grutters M, Klaassen R, Krijn M, Mouissie M, Wymenga E (2020) Ecological impact and cost-effectiveness of wildlife crossings in a highly fragmented landscape: A multi-method approach. Landscape Ecology 35(7): 1701–1720.

Smith DJ (2003) Monitoring wildlife use and determining standards for culvert design. Final Report, Florida Dept. of Transportation. 82 pp.

Smith DJ, van der Ree R, Rosell C (2015a) Wildlife crossing structures: an effective strategy to restore or maintain wildlife connectivity across roads. Chapter 21 (pp. 172-183) in van der Ree R, Smith DJ, Grilo C (eds.) Handbook of Road Ecology. John Wiley & Sons, Ltd.

Smith DJ, Kintsch J, Cramer P, Jacobson SL, Tonjes S (2015b) Modifying structures on existing roads to enhance wildlife passage. Chapter 10 (pp. 208-228) in Andrews KM, Nanjappa P, Riley SPD (eds.) Roads and ecological infrastructure; Concepts and applications for small animals. John Hopkins University Press.

Trafikverket (2022) Krav - VGU, Vägars och gators utformning. Report 2022:001, Trafikverket, Borlänge, Sweden, 365 pp. http://trafikverket.diva-portal.org/smash/get/diva2:1621114/FULLTEXT02.pdf

Transport Ecology Guidelines Portal (2024). IENE – Infrastructure & Ecology Network Europe. https://handbookwildlifetraffic.info/transport-ecology-guidelines-portal/

Väre S, Huhta M, Martin A (2003) Eläinten kulkujärjestelyt tiealueen poikki. Reports of the Finnish Road Administration 36/2003. https://www.doria.fi/handle/10024/139129

Tables

Table 1. Fauna passages monitored for selection of substrate by ungulates, with basic design and monitoring period.

	Passage type	Infra-structure	Area	Monitoring start	Monitoring end
Kvarnbäcken	Underpass/viaduct	Railway	Sweden	July 2019	July 2020
Aavajoki	Underpass/viaduct	Railway	Sweden	Nov 2019	Nov 2020
Keräsjoki	Underpass/viaduct	Railway	Sweden	July 2019	July 2020
Sangijärvi	Overpass/green bridge	Railway	Sweden	Nov 2019	Oct 2020
Sammatti	Overpass/green bridge	Motorway	Finland	Dec 2019	Nov 2020
Loviisa[a]	Overpass/green bridge	Motorway	Finland	Dec 2019	Nov 2020
Kärmekorpi	Overpass/green bridge	Motorway	Finland	Dec 2019	Nov 2020

[a] Labelled "Loviisa 2" in Niemi (2021).

Table 2. Substrates available in the fauna passages, in width (m) along the record line.

	Grassy/sandy	Stony/rocky	Fiber mat	Shrub	Water	Road	*Total*
Kvarnbäcken	0	10.0	6.4	0	9.6	0	*26.0*
Aavajoki	0	27.1	5.4	2.5	5.0	0	*40.0*
Keräsjoki	0	16.0	2.8	5.0	30.2	0	*54.0*
Sangijärvi	0	0	0	17.5	0	2.5	*20.0*
Sammatti	4.2	0	0	13.7	0	6.1	*24.0*
Loviisa[a]	0	0	0	16.1	0	5.8	*21.9*
Kärmekorpi	22.2	0	0	0	0	6.8	*29.0*

[a] Total width of bridge was 27m but 5m behind a railing was excluded from the analysis as it was not reliably covered by the camera.

Table 3. Number of ungulate crossing events included in the study, separated by season (snow-free/snow covered ground) and species.

	Snow-free					Snow covered ground				
	Moose	Roe deer	Reindeer	White-tailed deer	Wild boar	Moose	Roe deer	Reindeer	White-tailed deer	Wild boar
Kvarnbäcken	20	38	21	-	-	53	1	19	-	-
Aavajoki	96	6	65	-	-	31	0	11	-	-
Keräsjoki	22	59	1	-	-	13	9	10	-	-
Sangijärvi	5	18	59	-	-	20	2	80	-	-
Sammatti	9	37	-	172	0	0	0	-	10	0
Loviisa	21	4	-	2	3	1	0	-	2	0
Kärmekorpi	41	5	-	13	2	2	0	-	0	0

Table 4A-B. Results from analyses of ungulate substrate use in relation to availability, during snow-
free season and snow covered ground, respectively. U = underpass, O = overpass. Values in columns
4-9 are the relative contribution to the χ^2 in column 2; green cells = selected substrate (observed >
expected), red cells = avoided substrate (observed < expected), a = substrate used according to
availablility, empty cells = substrate not available. Only analyses reaching required sample size (all
expected values >5) are presented.

1	2	3	4	5	6	7	8	9
A) Snow-free	χ^2	p	Grassy/ sandy	Stony/ rocky	Fiber mat	Shrub	Water	Road
All species								
Kvarnbäcken U	31.07	<0.001		0.002	0.634		0.364	
Aavajoki U	178.1	<0.001		0.052	0.833	0.039	0.077	
Sangijärvi O	473.1	<0.001				0.127		0.873
Sammatti O	332.8	<0.001	0.015			0.356		0.629
Loviisa O	0.189	0.664				a		a
Kärmekorpi O	6.335	0.012	0.235					0.765
Moose								
Aavajoki U	65.27	<0.001		0.04	0.813	0.065	0.082	
Loviisa O	0.046	0.83				a		a
Kärmekorpi O	10.12	0.002	0.235					0.765
Roe deer								
Kvarnbäcken U	19.09	<0.001		0.041	0.419		0.54	
Sammatti O	56.35	<0.001	0.006			0.34		0.653
Reindeer								
Kvarnbäcken U	16.89	<0.001		0.006	0.535		0.459	
Sangijärvi O	390.0	<0.001				0.127		0.873
White-tailed deer								
Sammatti O	278.4	<0.001	0.006			0.339		0.656

B) Snow covered ground	χ^2	p	Grassy/ sandy	Stony/ rocky	Fiber mat	Shrub	Water	Road
All species								
Kvarnbäcken U	89.82	<0.001		0.211	0.158		0.631	
Sangijärvi O	552.5	<0.001				0.127		0.873
Moose								
Kvarnbäcken U	75.10	<0.001		0.221	0.148		0.631	
Reindeer								
Sangijärvi O	503.7	<0.001				0.127		0.873

392   *Table 5A-B. Results from analyses of ungulate use of water and road in relation to other substrates,*
393   *during snow-free season and snow covered ground, respectively. Green cells = selected substrate*
394   *(observed > expected), red cells = avoided substrate (observed < expected). Only analyses reaching*
395   *required sample size (all expected values >5) are presented.*

	Snow-free		Snow covered	
A) Water (underpasses)	χ^2	*p*	χ^2	*p*
All species				
Kvarnbäcken	17.91	<0.001	89.77	<0.001
Aavajoki	15.59	<0.001	6.0	0.014
Keräsjoki	95.07	<0.001	36.15	<0.001
Moose				
Kvarnbäcken	0.564	0.45	75.10	<0.001
Aavajoki	6.095	0.014	16.48	<0.001
Keräsjoki	19.55	<0.001		
Roe deer				
Kvarnbäcken	16.33	<0.001		
Keräsjoki	74.79	<0.001		
Reindeer				
Kvarnbäcken	12.28	<0.001	18.26	<0.001
Aavajoki	9.286	0.002		
	Snow-free		Snow covered	
B) Road (overpasses)	χ^2	*p*	χ^2	*p*
All species				
Sangijärvi	930.2	<0.001	715.2	<0.001
Sammatti	380.4	<0.001		
Loviisa	0.189	0.66		
Kärmekorpi	6.335	0.012		
Moose				
Loviisa	0.046	0.83		
Kärmekorpi	10.12	0.001		
Roe deer				
Sammatti	49.36	<0.001		
Reindeer				
Sangijärvi	826.4	<0.001	620.4	<0.001
White-tailed deer				
Sammatti	244.7	<0.001		

397 Figures

398 *Figure 1. Location of fauna passages included in the study.*

Figure 2A-B. Examples of ungulate trajectories through fauna passages. A: Kvarnbäcken underpass, B: Sammatti overpass, both at snow free conditions. Red = moose, blue = reindeer, green = roe deer, orange = white-tailed deer. Dotted line = record line. Background images show camera placement and direction, perimeter fencing, and available substrates.

Figure 3A-D. Examples
of records of ungulates
crossing through fauna
passages on different
substrates. A: moose
on grassy/sandy
substrate (Kärmekorpi,
12 Jun, 2020); B: roe
deer on fiber mat
(Kvarnbäcken, 2 Jun
2020), C: reindeer on
stony/rocky substrate
(Aavajoki, 25 Jun 2020),
D: white-tailed deer on
road (Sammatti, May 6
2020).

Figure 4A-C. Ungulates crossing fauna passages via a watercourse. A: reindeer walking on river ice
(Kvarnbäcken, 31 Dec 2019); B: moose wading along river bank (Kvarnbäcken, 24 Oct 2019). C: moose
swimming along river (Aavajoki, Oct 13 2020).