Research Article

Augmented-reality-based multi-person exercise has more beneficial effects on mood state and

oxytocin secretion than standard solitary exercise

Takeru Shima[1*], Junpei Iijima[1], Hirotaka Sutoh[1], Chiho Terashima[1], Yuki Matsuura[2]

[1] Department of Health and Physical Education, Cooperative Faculty of Education, Gunma

University, 4-2 Aramaki-machi, Maebashi, Gunma 371-8510, Japan

[2] Department of Health and Physical Education, Cooperative Faculty of Education, Utsunomiya

University, 350 Mine-machi, Utsunomiya, Tochigi 321-8505, Japan

ORCID: Takeru Shima, 0000-0003-4209-1564; Yuki Matsuura, 0000-0003-1660-5680

*Corresponding author: Takeru Shima, Department of Health and Physical Education,

Cooperative Faculty of Education, Gunma University, 4-2 Aramaki-machi, Maebashi, Gunma, 371-

8510, Japan.

E-mail: ta-shima@gunma-u.ac.jp

Abstract

Exercise is known to have positive effects on psychological well-being, with team sports often

associated with superior mental health compared to individual sports. Augmented reality (AR)

technology has the potential to convert solitary exercise into multi-person exercise. Given the role

of oxytocin in mediating the psychological benefits of exercise and sports, this study aimed to

investigate the impact of AR-based multi-person exercise on mood and salivary oxytocin levels.

Fourteen participants underwent three distinct regimens: non-exercise (Rest), standard solitary

cycling exercise (Ex), and AR-based multi-person cycling exercise (Ex+AR). In both exercise

conditions (Ex and Ex+AR), participants engaged in cycling at a self-regulated pace to maintain a

Rating of Perceived Exertion of 10. In the Ex+AR condition, participants' avatars were projected

onto a tablet screen, allowing them to cycle alongside ten other virtual avatars in an AR

environment. Mood states (assessed using POMS2) and saliva samples were collected before and

immediately after each 10-minute regimen. Subsequently, the levels of salivary oxytocin were

measured. Participants exhibited higher cycling speeds during Ex+AR compared to Ex, despite

comparable levels of self-reported fatigue between the two groups. Notably, only the Ex+AR

condition significantly improved mood states associated with depression-dejection and exhibited a

trend toward suppressing anger-hostility in participants. Moreover, the Ex+AR condition led to a

significant elevation in salivary oxytocin levels, while the Ex condition showed a trend toward an

increase. However, changes in salivary oxytocin did not show a significant correlation with changes

in mood states. These findings suggest that Ex+AR enhances mood states and promotes oxytocin

release. AR-based multi-person exercise may offer greater psychological benefits compared to

standard solitary exercise, although the relationship between oxytocin and mood changes remains

inconclusive.

Highlights

- Exercise, especially team sports, has positive effects on mood state, and oxytocin partially supports these effects.

- Augmented reality (AR) technology can potentially convert standard solitary exercise into multi-person exercise.

- AR-based multi-person exercise (Ex+AR) significantly ameliorated depression-dejection and increased salivary oxytocin levels, but standard solitary exercise did not.

- Ex+AR may provide more significant psychological benefits than standard solitary exercise.

Keywords

Exercise, Augmented reality, Mood state, Oxytocin

1. Introduction

Numerous studies have underscored the advantageous impacts of exercise on both physiological and psychological well-being. Consequently, exercise is viewed as a therapeutic intervention not only for metabolic disturbances but also for mental disorders (Deslandes et al., 2009; Hwang et al., 2023; Laird et al., 2023; Viana and de Lira, 2020). The prevalence of mental disorders has been on the rise in recent decades (ten Have et al., 2023), with children and young adults under 24 years of age being particularly affected by the adverse mental health effects of the COVID-19 pandemic (Kauhanen et al., 2023). While team sport athletes typically exhibit lower levels of anxiety and depression compared to individual sport athletes (Pluhar et al., 2019), the mental well-being of team sport athletes is susceptible to the impact of the COVID-19 pandemic due to the lack of social team activities (Salles et al., 2022). Consequently, exercise modalities resembling team sports that can be practiced under any environmental condition, such as during the COVID-19 pandemic, are deemed crucial for sustaining and improving human mental health.

Extended Reality (XR) technology holds promise for facilitating exercise under various environmental conditions. XR technology encompasses a spectrum of natural and virtual environments, including Virtual Reality (VR), Augmented Reality (AR), and Mixed Reality (MR) technologies. XR-based communication is likely effective in enhancing interaction (Dong et al., 2013; Rogers et al., 2022), and its applications to enhance human capabilities are burgeoning across

various fields, such as sports and medical science (Barteit et al., 2021; Janssen et al., 2023; Le

Noury et al., 2023). Among XR technologies, AR technology is particularly suited for sports

applications as it can be utilized without the need for a head-mounted display. Leveraging AR has

the potential to convert standard solitary exercise into multi-person exercise. Nevertheless, research

investigating the effectiveness of AR technology for multi-person exercise is still in its nascent

stages.

Oxytocin is implicated in the psychological benefits derived from exercise and sports.

Oxytocin is a nine-amino acid peptide synthesized in the paraventricular nuclei and supraoptic

nuclei of the hypothalamus, subsequently secreted from the posterior pituitary gland into circulation.

Social behaviors such as affectionate touch and physical proximity are known to stimulate oxytocin

secretion (Burenkova et al., 2023; Caicedo Mera et al., 2021; Holt-Lunstad et al., 2008; Landgraf

and Neumann, 2004). Light-intensity acute exercise has been shown to enhance mood and prosocial

skills in healthy individuals (Basso and Suzuki, 2017; Edwards et al., 2017; Ludyga et al., 2022),

with oxytocin believed to partially mediate these effects (Ludyga et al., 2022). Moreover, even brief

10-minute exercise sessions have been shown to elevate salivary oxytocin levels (de Jong et al.,

2015). Given that variations in oxytocin secretion are influenced by the type of exercise (Rassovsky

et al., 2019), disparities in oxytocin secretion may contribute to the psychological benefits observed

in team sports, such as reduced anxiety and depression levels compared to individual sports (Pluhar

et al., 2019).

Here, to investigate the beneficial effects of multi-person exercises using AR technology,

we examined the alterations in mood state and salivary oxytocin levels across three experimental

conditions: 1. AR-based multi-person cycling exercise (Ex+AR), 2. standard solitary cycling

exercise (Ex), and 3. Rest.

2. Methods and materials

2.1. Participants

Fourteen Japanese college students (8 men and 6 women, age: 20.5 ± 1.7), who had received

instructions and provided consent, participated in the study. Informed consent was obtained from all

individual participants prior to their involvement. The Utsunomiya University Ethical Review

Board approved the current investigation for Medical Research Involving Human Subjects (ethical

number: H23-0012). Participants had no history of neurological, psychiatric, or respiratory

disorders, diabetes, anemia, or other medical conditions. Additionally, all participants underwent

Ex+AR training prior to the main experiment. Post hoc sensitivity analysis conducted on the current

sample, with 80% power and $\alpha = 0.05$, revealed adequate sensitivity to detect repeated-measures

effects exceeding $f = 0.63$ and paired t-test differences exceeding $d = 0.81$ (with a two-tailed α), as

calculated using G*Power ver. 3.1.9.6 (The G*Power Team).

2.2. Experimental procedure

All participants were randomly assigned to undergo all three experimental conditions at

one-week intervals: rest, Ex, and Ex+AR. Participants were instructed to refrain from exhaustive

exercise, as well as from consuming alcohol and caffeine, for at least 24 hours prior to each

experiment. The experimental procedure is shown in Figure 1. Paired participants were

simultaneously exposed to the experiment under the same conditions in separate rooms. Participants

completed the POMS2 questionnaire, and 1 ml saliva samples were collected after a 10-minute rest

period. Following another 10-minute rest, participants engaged in a 10-minute sedentary period, Ex,

or Ex+AR. Subsequently, participants completed the POMS2 questionnaire again, and saliva

samples were collected. Heart rate (HR) was measured using an HR sensor (H10; Polar Electro,

Japan).

2.3. Exercise and augmented reality system

121   In both exercise conditions (Ex and Ex+AR), participants engaged in self-paced cycling

122 exercises to maintain a Rating of Perceived Exertion (RPE) of 10. Information regarding time

123 remaining and confirming RPE remained at 10 were received, with no other information or external

124 encouragement provided. Mean speed (km/h) was recorded throughout the 10-minute cycling

session using the SmarT Turbo KAGURA (LST9200, MINOURA Co., Ltd., Japan) and the

ROUVY application (VirtualTraining Co., Ltd., Czech Republic). During the Ex condition,

participants cycled in front of a flipped tablet. In contrast, during the Ex+AR condition, participants

cycled in front of a tablet (11-in) running the ROUVY application. In the Ex+AR condition,

participants' avatars were projected onto the tablet screen as they cycled in augmented reality

alongside ten other avatars, including the avatar of a paired participant (refer to Fig. 1). Notably,

participants did not utilize a headset with the current AR system.

2.4. Questionnaire

To assess the seven subscales of mood states (anger-hostility, confusion-bewilderment,

depression-ejection, fatigue-inertia, tension-anxiety, vigor-activity, and friendliness), as well as

total mood disturbance, participants completed the short form of POMS2 (Kaneko Shobo, Japan)

before and immediately after each experimental condition. The short form of POMS2 comprises 35

items, with participants responding to a 5-point Likert scale ranging from 1 (strongly disagree) to 5

(strongly agree).

2.5. Saliva collection and measurement of oxytocin concentration

Saliva was chosen as the medium for evaluating oxytocin levels as it provides a reliable

indicator of changes associated with interventions (Burenkova et al., 2023). Saliva samples were

collected from participants before and immediately after engaging in each experimental condition

using the Saliva Collection Aid (Salimetrics, LLC., USA) and Cryovial (Salimetrics, LLC., USA).

These samples were stored at -20°C for subsequent analysis. Oxytocin concentrations in the saliva

samples were measured in duplicate and calculated following the manufacturer's instructions

utilizing the Oxytocin ELISA Kit Wako (FUJIFILM Wako Pure Chemicals Corp., Japan) and a

plate reader (EnSpire® Multimode Plate Reader, PerkinElmer, Inc., Massachusetts, USA).

2.6. Statistical analysis

The data are presented as mean ± SD and were analyzed using Prism version 10.1.1 (MDF,

Japan). Before the analysis for group comparisons, we checked sphericity of the data. If the

sphericity assumption was confirmed, group comparisons were conducted using repeated measures

two-way ANOVA (factor 1: time points [pre vs. post]; factor 2: groups [rest vs. Ex vs. Ex+AR])

with Šídák's multiple comparisons test. When the sphericity assumption was not confirmed,

Geisser-Greenhouse correction was applied. A comparison between Ex and Ex+AR was performed

using a paired t-test with checking the normality of raw data distribution using histograms (for

analyzing mean speed). The correlation of mean speeds and changes in mood states with the

changes in salivary oxytocin levels (baseline-normalized changes) were assessed using Pearson

correlation. Changes in mood states were calculated by subtracting the scores at pre-exercise from

the scores at post-exercise. Baseline-normalized changes in salivary oxytocin levels were calculated

by subtracting the scores at pre from the scores at poste and then dividing by the scores at pre-

exercise. Statistical significance was set at $p < 0.05$.

3. Results

3.1. Heart rate and mean speed during cycling exercise

Both the Ex and Ex+AR conditions demonstrated a significant increase in participants' heart

rate post-exercise compared to pre-exercise (Fig. 2A, Ex: $p < 0.0001$, Ex+AR: $p < 0.0001$; main

effects of time points: $F_{(1.00, 13.00)} = 129.0$, $p < 0.0001$, main effects of group: $F_{(1.90, 24.70)} = 40.6$, $p <$

0.0001, interaction: $F_{(1.52, 19.71)} = 79.8$, $p < 0.0001$). Conversely, participants' heart rate remained

unchanged during the rest condition (Fig. 2A, $p = 0.9984$). Despite efforts to maintain an RPE of 10

in both the Ex and Ex+AR conditions, the mean cycling speed was higher in the Ex+AR condition

compared to the Ex condition (Fig. 2B; $t = 2.39$, $p = 0.033$).

3.2. The changes in mood state

Only Ex+AR significantly suppressed the depression-ejection score (Fig. 3C, $p = 0.0314$;

main effects of time points: $F_{(1.00, 13.00)} = 6.92$, $p = 0.0208$, main effects of group: $F_{(1.89, 24.57)} = 1.53$,

$p = 0.2370$, interaction: $F_{(1.61, 20.92)} = 0.72$, $p = 0.4686$) and exhibited a trend towards a decrease in

the scores of anger-hostility and total mood disturbance (Fig. 3A, H; anger-hostility: main effects of

time points: $F_{(1.00, 13.00)} = 8.87$, $p = 0.0107$, main effects of group: $F_{(1.46, 18.95)} = 0.04$, $p = 0.9240$,

interaction: $F_{(1.63, 21.20)} = 1.26$, $p = 0.2960$; total mood disturbance: main effects of time points: $F_{(1.00,}$

$_{13.00)} = 7.08$, $p = 0.0196$, main effects of group: $F_{(1.77, 23.05)} = 0.95$, $p = 0.3907$, interaction: $F_{(1.70, 22.04)}$

$= 1.32$, $p = 0.2830$). Both Ex and Ex+AR showed a trend towards a decrease in the tension-anxiety

score (Fig. 3E; main effects of time points: $F_{(1.00, 13.00)} = 15.0$, $p = 0.0019$, main effects of group:

$F_{(1.88, 24.41)} = 1.01$, $p = 0.3747$, interaction: $F_{(1.65, 21.46)} = 1.27$, $p = 0.2941$). None of the subscales of

mood state or total mood disturbance showed alterations in the rest condition (Fig. 3A-H).

3.3. The changes in salivary oxytocin

Ex+AR significantly increased salivary oxytocin levels (Fig. 4A, $p = 0.0005$; main effects of

time points: $F_{(1.00, 13.00)} = 24.8$, $p = 0.0002$, main effects of group: $F_{(1.84, 23.93)} = 2.30$, $p = 0.1259$,

interaction: $F_{(1.94, 25.16)} = 10.0$, $p = 0.0007$), whereas Ex exhibited a trend towards an increase (Fig.

4A, $p = 0.0868$). Salivary oxytocin levels remained unchanged in the rest condition (Fig. 4A, $p =$

0.5238). On the other hand, no significant correlation was observed between cycling speeds and

baseline-normalized changes in salivary oxytocin (Fig. 4B; $r = 0.0411$, $p = 0.8354$). Additionally,

baseline-normalized changes in salivary oxytocin showed no significant correlation with changes in

any mood states (Fig. 5A-H; all $p > 0.05$)

4. Discussion

In the current study, we investigated the effects of Ex+AR on psychological state and

salivary oxytocin concentration compared to Ex. Our findings demonstrated that, despite

comparable levels of effort to Ex, Ex+AR significantly improved depression-dejection scores and

elevated salivary oxytocin levels.

Although participants engaged in cycling exercises at a self-regulated pace to maintain a

RPE of 10 in the current study, the mean speed during cycling exercise in the Ex+AR condition was

significantly faster with no difference in heart rate than in the Ex condition (Fig. 2A and B). Prior

research has indicated that utilizing Extended Reality during exercise can mitigate human fatigue

(Berezina et al., 2022; Plante et al., 2003). The current results showed no difference in fatigue

evaluated by POMS2 among the groups (Fig. 3D). Therefore, utilizing Augmented Reality during

exercise could potentially increase exercise intensity while maintaining similar levels of fatigue as

standard exercise. It's also worth considering that using an AR system without a head-mounted

display may affect perceptual functions differently (Livingston et al., 2013). Since our study did not

incorporate a head-mounted display in the AR system, future research should examine the

comparative effects of the presence or absence of such a head-mounted display on the effectiveness

of AR-based exercises.

Only the Ex+AR condition significantly suppressed the depression-ejection score (Fig. 3C)

and showed a trend towards reducing anger-hostility and total mood disturbance scores (Fig. 3A, H).

Considering that team sport athletes often exhibit a lower risk of anxiety and depression compared

to individual sport athletes (Pluhar et al., 2019), the online exercise with multiple participants (ghost

riders) in our Ex+AR condition may have induced effects similar to those observed in team sports.

On the other hand, both the Ex and Ex+AR conditions showed a trend towards decreasing tension-

anxiety scores (Fig. 3E), indicating the effectiveness of not only Ex+AR but also Ex, consistent

with previous studies (Basso and Suzuki, 2017; Edwards et al., 2017). based on our findings,

Ex+AR appears to have more beneficial effects on psychological aspects than Ex. Therefore, AR-

based multi-person exercise may contribute to maintaining better mental health compared to

standard solitary exercise.

Salivary oxytocin levels increased significantly only in the Ex+AR condition, while the Ex

condition exhibited a trend towards an increase (Fig. 4A). Since baseline-normalized changes in

salivary oxytocin levels did not correlate with cycling speeds (Fig. 4B), it suggests that the use of

the AR system could enhance oxytocin secretion during exercise. Previous research has

demonstrated that XR-based communication facilitates social interaction (Dong et al., 2013; Rogers

et al., 2022) and oxytocin secretion (Dekker et al., 2021). Although we could not investigate

enhancements in the social interaction of subjects by the Ex+AR condition, AR-based multi-person

exercise would enhance not only mood states but also salivary oxytocin levels.

Baseline-normalized changes in salivary oxytocin levels did not correlate with changes in

any mood states (Fig. 5A-H). While oxytocin has been implicated in reducing depressed mood,

anger expression, and anxiety state (Domes et al., 2007; Galbally et al., 2021; Goodin et al., 2015;

Heinrichs et al., 2003; Yoon and Kim, 2022), our findings suggest that changes in oxytocin

secretion with exercise may not directly relate to exercise-induced changes in mood states.

Oxytocin is also associated with empathy (Meyer-Lindenberg et al., 2011), and several studies

have reported that oxytocin administration enhances empathy and related behaviors in humans

(Auyeung et al., 2015; Geng et al., 2018). Considering that physical activities also contribute to

enhancing empathy (Shima et al., 2022a, 2022b, 2021c, 2021a, 2021b), it is conceivable that

exercise-induced oxytocin secretion plays a role in enhancing empathy. Therefore, AR-based

multi-person exercise may have more beneficial effects on enhancing empathy than standard

exercise, warranting further investigation.

The current study has several limitations. Firstly, the current study exclusively involved

healthy participants. Given that standard exercise has been shown to prevent and treat psychiatric

disorders (Deslandes et al., 2009; Hwang et al., 2023; Laird et al., 2023; Viana and de Lira, 2020),

there is a possibility that the current Ex+AR approach could serve as a more efficient therapy for

such disorders compared to standard exercise alone. Future studies should explore this possibility

Secondly, our investigation focused solely on the acute effects of a 10-minute regimen. Prior

research suggests that prolonged exposure to AR (>20 min) is associated with cybersickness

(Hughes et al., 2020). Therefore, further study is warranted to determine the optimal duration for

AR-based multi-person exercise. Thirdly, the effects of oxytocin and exercise may vary among

individuals (Shima et al., 2022a; Sikich et al., 2021). Therefore, further research is needed to

determine whether the current Ex+AR condition effectively enhances human well-being. Lastly, the

assessment of social interaction provided by the AR system was not evaluated. Future studies would

benefit from analyzing the quality of social interaction using established methods (Lopes et al.,

2004; Reis and Wheeler, 1991).

In conclusion, our findings exhibit that Ex+AR significantly enhances mood states and

increases salivary oxytocin levels. AR-based multi-person exercise may indeed provide greater

psychological benefits compared to standard solitary exercise, although the relationship between

oxytocin and mood changes remains inconclusive.

Funding:

This work was supported by the Public Health Research Foundation and the Foundation

for the Fusion of Science and Technology.

Data availability:

The datasets in the current study are available from the corresponding author on

reasonable request.

Declaration of competing interest

The authors inform no conflicts of interest.

Author contribution

TS: Conceptualization, Methodology, Resources, Investigation, Writing–original draft,

Writing–review & editing, Funding acquisition; JI: Investigation, Writing–review & editing, HS:

Investigation, Writing–review & editing, CT: Investigation, Writing–review & editing, YM:

Conceptualization, Methodology, Writing–review & editing.

References

Auyeung, B., Lombardo, M. V., Heinrichs, M., Chakrabarti, B., Sule, A., Deakin, J.B., Bethlehem, R.A.I., Dickens, L., Mooney, N., Sipple, J.A.N., Thiemann, P., Baron-Cohen, S., 2015. Oxytocin increases eye contact during a real-time, naturalistic social interaction in males with and without autism. Transl Psychiatry 5. https://doi.org/10.1038/tp.2014.146

Barteit, S., Lanfermann, L., Bärnighausen, T., Neuhann, F., Beiersmann, C., 2021. Augmented, mixed, and virtual reality-based head-mounted devices for medical education: Systematic review. JMIR Serious Games 9. https://doi.org/10.2196/29080

Basso, J.C., Suzuki, W.A., 2017. The Effects of Acute Exercise on Mood, Cognition, Neurophysiology, and Neurochemical Pathways: A Review. Brain Plasticity 2, 127–152. https://doi.org/10.3233/BPL-160040

Berezina, T., Temirkanova, A., Litvinova, A., Kokurin, A., 2022. Using Virtual Reality Techniques to Alleviate Cognitive Fatigue in Graduate Students Working while in College. European Journal of Contemporary Education 11, 36–46. https://doi.org/10.13187/ejced.2022.1.36

Burenkova, O. V., Dolgorukova, T.A., An, I., Kustova, T.A., Podturkin, A.A., Shurdova, E.M., Talantseva, O.I., Zhukova, M.A., Grigorenko, E.L., 2023. Endogenous oxytocin and human social interactions: A systematic review and meta-analysis., Psychological Bulletin. https://doi.org/10.1037/bul0000402

Caicedo Mera, J.C., Cárdenas Molano, M.A., García López, C.C., Acevedo Triana, C., Martínez

Cotrina, J., 2021. Discussions and perspectives regarding oxytocin as a biomarker in human

investigations. Heliyon 7. https://doi.org/10.1016/j.heliyon.2021.e08289

de Jong, T.R., Menon, R., Bludau, A., Grund, T., Biermeier, V., Klampfl, S.M., Jurek, B., Bosch,

O.J., Hellhammer, J., Neumann, I.D., 2015. Salivary oxytocin concentrations in response to

running, sexual self-stimulation, breastfeeding and the TSST: The Regensburg Oxytocin

Challenge (ROC) study. Psychoneuroendocrinology 62, 381–388.

https://doi.org/10.1016/j.psyneuen.2015.08.027

Dekker, A., Wenzlaff, F., Biedermann, S. V., Briken, P., Fuss, J., 2021. VR Porn as "Empathy

Machine"? Perception of Self and Others in Virtual Reality Pornography. The Journal of Sex

Research 58, 273–278. https://doi.org/10.1080/00224499.2020.1856316

Deslandes, A., Moraes, H., Ferreira, C., Veiga, H., Silveira, H., Mouta, R., Pompeu, F.A.M.S.,

Coutinho, E.S.F., Laks, J., 2009. Exercise and Mental Health: Many Reasons to Move.

Neuropsychobiology 59, 191–198. https://doi.org/10.1159/000223730

Domes, G., Heinrichs, M., Gläscher, J., Büchel, C., Braus, D.F., Herpertz, S.C., 2007. Oxytocin

Attenuates Amygdala Responses to Emotional Faces Regardless of Valence. Biol Psychiatry

62, 1187–1190. https://doi.org/10.1016/j.biopsych.2007.03.025

Dong, S., Behzadan, A.H., Chen, F., Kamat, V.R., 2013. Collaborative visualization of engineering

processes using tabletop augmented reality. Advances in Engineering Software 55, 45–55.

https://doi.org/10.1016/j.advengsoft.2012.09.001

Edwards, M.K., Rhodes, R.E., Loprinzi, P.D., 2017. A Randomized Control Intervention

Investigating the Effects of Acute Exercise on Emotional Regulation. Am J Health Behav 41,

534–543. https://doi.org/10.5993/AJHB.41.5.2

Galbally, M., Watson, S.J., Keelan, J.A., Spigset, O., Lewis, A., 2021. The relationship between

oxytocin blood concentrations and antidepressants over pregnancy and the postpartum. Prog

Neuropsychopharmacol Biol Psychiatry 109, 110218.

https://doi.org/10.1016/j.pnpbp.2020.110218

Geng, Y., Zhao, W., Zhou, F., Ma, X., Yao, S., Hurlemann, R., Becker, B., Kendrick, K.M., 2018.

Oxytocin Enhancement of Emotional Empathy: Generalization Across Cultures and Effects on

Amygdala Activity. Front Neurosci 12, 1–14. https://doi.org/10.3389/fnins.2018.00512

Goodin, B.R., Anderson, A.J.B., Freeman, E.L., Bulls, H.W., Robbins, M.T., Ness, T.J., 2015.

Intranasal Oxytocin Administration is Associated With Enhanced Endogenous Pain Inhibition

and Reduced Negative Mood States. Clin J Pain 31, 757–767.

https://doi.org/10.1097/AJP.0000000000000166

Heinrichs, M., Baumgartner, T., Kirschbaum, C., Ehlert, U., 2003. Social support and oxytocin

interact to suppress cortisol and subjective responses to psychosocial stress. Biol Psychiatry 54,

1389–1398. https://doi.org/10.1016/S0006-3223(03)00465-7

Holt-Lunstad, J., Birmingham, W.A., Light, K.C., 2008. Influence of a "Warm Touch" Support

Enhancement Intervention Among Married Couples on Ambulatory Blood Pressure, Oxytocin,

Alpha Amylase, and Cortisol. Psychosom Med 70, 976–985.

https://doi.org/10.1097/PSY.0b013e318187aef7

Hughes, C.L., Fidopiastis, C., Stanney, K.M., Bailey, P.S., Ruiz, E., 2020. The Psychometrics of

Cybersickness in Augmented Reality. Front Virtual Real 1, 1–12.

https://doi.org/10.3389/frvir.2020.602954

Hwang, D.J., Koo, J.H., Kim, T.K., Jang, Y.C., Hyun, A.H., Yook, J.S., Yoon, C.S., Cho, J.Y.,

2023. Exercise as an antidepressant: exploring its therapeutic potential. Front Psychiatry 14, 1–

11. https://doi.org/10.3389/fpsyt.2023.1259711

Janssen, T., Müller, D., Mann, D.L., 2023. From Natural Towards Representative Decision Making

in Sports: A Framework for Decision Making in Virtual and Augmented Environments. Sports

Medicine 53, 1851–1864. https://doi.org/10.1007/s40279-023-01884-3

Kauhanen, L., Wan Mohd Yunus, W.M.A., Lempinen, L., Peltonen, K., Gyllenberg, D., Mishina,

K., Gilbert, S., Bastola, K., Brown, J.S.L., Sourander, A., 2023. A systematic review of the

mental health changes of children and young people before and during the COVID-19

pandemic. Eur Child Adolesc Psychiatry 32, 995–1013. https://doi.org/10.1007/s00787-022-

02060-0

Laird, E., Rasmussen, C.L., Kenny, R.A., Herring, M.P., 2023. Physical Activity Dose and

Depression in a Cohort of Older Adults in The Irish Longitudinal Study on Ageing. JAMA

Netw Open 6, e2322489. https://doi.org/10.1001/jamanetworkopen.2023.22489

Landgraf, R., Neumann, I.D., 2004. Vasopressin and oxytocin release within the brain: a dynamic

concept of multiple and variable modes of neuropeptide communication. Front

Neuroendocrinol 25, 150–176. https://doi.org/10.1016/j.yfrne.2004.05.001

Le Noury, P.J., Polman, R.C., Maloney, M.A., Gorman, A.D., 2023. XR Programmers Give Their

Perspective on How XR Technology can be Effectively Utilised in High-Performance Sport.

Sports Med Open 9. https://doi.org/10.1186/s40798-023-00593-5

Livingston, M.A., Gabbard, J.L., Swan, J.E., Sibley, C.M., Barrow, J.H., 2013. Basic Perception in

Head-Worn Augmented Reality Displays, in: Human Factors in Augmented Reality

Environments. Springer New York, New York, NY, pp. 35–65. https://doi.org/10.1007/978-1-

4614-4205-9_3

365 Lopes, P.N., Brackett, M.A., Nezlek, J.B., Schütz, A., Sellin, I., Salovey, P., 2004. Emotional

366   Intelligence and Social Interaction. Pers Soc Psychol Bull 30, 1018–1034.

367   https://doi.org/10.1177/0146167204264762

368 Ludyga, S., Ishihara, T., Kamijo, K., 2022. The Nervous System as a Pathway for Exercise to

369   Improve Social Cognition. Exerc Sport Sci Rev 50, 203–212.

370   https://doi.org/10.1249/JES.0000000000000300

371 Meyer-Lindenberg, A., Domes, G., Kirsch, P., Heinrichs, M., 2011. Oxytocin and vasopressin in

372   the human brain: social neuropeptides for translational medicine. Nat Rev Neurosci 12, 524–

373   538. https://doi.org/10.1038/nrn3044

374 Plante, T.G., Aldridge, A., Bogden, R., Hanelin, C., 2003. Might virtual reality promote the mood

375   benefits of exercise? Comput Human Behav 19, 495–509. https://doi.org/10.1016/S0747-

376   5632(02)00074-2

377 Pluhar, E., McCracken, C., Griffith, K.L., Christino, M.A., Sugimoto, D., Meehan, W.P., 2019.

378   Team Sport Athletes May Be Less Likely To Suffer Anxiety or Depression than Individual

379   Sport Athletes. J Sports Sci Med 18, 490–496.

380 Rassovsky, Y., Harwood, A., Zagoory-sharon, O., Feldman, R., 2019. Martial arts increase oxytocin

381   production. Sci Rep 1–8. https://doi.org/10.1038/s41598-019-49620-0

Reis, H.T., Wheeler, L., 1991. Studying Social Interaction with the Rochester Interaction Record.

Adv Exp Soc Psychol 24, 269–318. https://doi.org/10.1016/S0065-2601(08)60332-9

Rogers, S.L., Broadbent, R., Brown, J., Fraser, A., Speelman, C.P., 2022. Realistic Motion Avatars

are the Future for Social Interaction in Virtual Reality. Front Virtual Real 2, 1–12.

https://doi.org/10.3389/frvir.2021.750729

Salles, J.I., Silva, C., Wolff, A., Orwert, L., Ribeiro, P., Velasques, B., Morrissey, D., 2022.

Anxiety, insomnia, and depression during COVID-19 lockdown in elite individual and team

sport athletes. Psicologia: Reflexao e Critica 35. https://doi.org/10.1186/s41155-022-00233-z

Shima, T., Jesmin, S., Nakao, H., Tai, K., Shimofure, T., Arai, Y., Kiyama, K., Onizawa, Y., 2021a.

Association between self-reported empathy and level of physical activity in healthy young

adults. J Phys Fit Sports Med 10, 45–49. https://doi.org/10.7600/jpfsm.10.45

Shima, T., Jesmin, S., Nakao, H., Tai, K., Shimofure, T., Arai, Y., Kiyama, K., Onizawa, Y., 2021b.

Small Amounts of Physical Activity During the COVID-19 Pandemic May Contribute to

Improve Empathy in Young Adults: An Observational Study. Asia Pacific Journal of Public

Health 33, 635–637. https://doi.org/10.1177/10105395211016333

Shima, T., Jesmin, S., Onishi, H., Yoshikawa, T., Saitoh, R., 2022a. Physical activity associates

empathy in Japanese young adults with specific gene variations of oxytocin receptor and

399   vasopressin V1B receptor. Physiol Behav 255, 113930.

400   https://doi.org/10.1016/j.physbeh.2022.113930

401 Shima, T., Nakao, H., Tai, K., Shimofure, T., Jesmin, S., Arai, Y., Kiyama, K., Onizawa, Y., 2022b.

402   The Influences of Changes in Physical Activity Levels With Easing Restriction of Access to

403   the University Campus on Empathy and Social Supports in College Students During the

404   COVID-19 Pandemic. Asia Pacific Journal of Public Health 34, 406–410.

405   https://doi.org/10.1177/10105395221083381

406 Shima, T., Tai, K., Nakao, H., Shimofure, T., 2021c. Association between self-reported empathy

407   and sport experience in young adults. Journal of Physical Education and Sport 21, 66–72.

408   https://doi.org/10.7752/jpes.2021.01009

409 Sikich, L., Kolevzon, A., King, B.H., McDougle, C.J., Sanders, K.B., Kim, S.-J., Spanos, M.,

410   Chandrasekhar, T., Trelles, M.D.P., Rockhill, C.M., Palumbo, M.L., Witters Cundiff, A.,

411   Montgomery, A., Siper, P., Minjarez, M., Nowinski, L.A., Marler, S., Shuffrey, L.C.,

412   Alderman, C., Weissman, J., Zappone, B., Mullett, J.E., Crosson, H., Hong, N., Siecinski, S.K.,

413   Giamberardino, S.N., Luo, S., She, L., Bhapkar, M., Dean, R., Scheer, A., Johnson, J.L.,

414   Gregory, S.G., Veenstra-VanderWeele, J., 2021. Intranasal Oxytocin in Children and

415   Adolescents with Autism Spectrum Disorder. New England Journal of Medicine 385, 1462–

416   1473. https://doi.org/10.1056/NEJMoa2103583

ten Have, M., Tuithof, M., van Dorsselaer, S., Schouten, F., Luik, A.I., de Graaf, R., 2023.

Prevalence and trends of common mental disorders from 2007‐2009 to 2019‐2022: results

from the Netherlands Mental Health Survey and Incidence Studies (NEMESIS), including

comparison of prevalence rates before vs. during the COVID‐19 pandemic. World Psychiatry

22, 275–285. https://doi.org/10.1002/wps.21087

Viana, R.B., de Lira, C.A.B., 2020. Exergames as Coping Strategies for Anxiety Disorders During

the COVID-19 Quarantine Period. Games Health J 9, 147–149.

https://doi.org/10.1089/g4h.2020.0060

Yoon, S., Kim, Y.-K., 2022. Possible oxytocin-related biomarkers in anxiety and mood disorders.

Prog Neuropsychopharmacol Biol Psychiatry 116, 110531.

https://doi.org/10.1016/j.pnpbp.2022.110531

430   **Figures:**

431

432   **Fig. 1**

434   **Fig. 2**

436   **Fig. 3**

Fig. 4

Fig. 5

Figure legends:

Fig. 1. Experimental protocol in the current study.

Fig. 2. (A) The changes of heart rate in each condition. (B) The difference of mean cycling speed

between Ex and Ex+AR condition. Ex: standard solitary cycling exercise, Ex+AR: augmented-

reality-based multi-person cycling exercise. Data are expressed as mean $\pm$ SD. Plots represent

individual data, n = 14. $^{*}p < 0.05,$ $^{****}p < 0.0001.$

Fig. 3. The changes of (A) anger-hostility, (B) confusion-bewilderment, (C) depression-ejection,

(D) fatigue-inertia, (E) tension-anxiety, (F) vigor-activity, (G) friendliness and (H) total mood

disturbance in each condition. Ex: standard solitary cycling exercise, Ex+AR: augmented-reality-

based multi-person cycling exercise. Data are expressed as mean $\pm$ SD. Plots represent individual

data, n = 14. $^{*}p < 0.05.$

Fig. 4. (A) The changes of salivary oxytocin levels in each condition. Ex: standard solitary cycling

exercise, Ex+AR: augmented-reality-based multi-person cycling exercise. Data are expressed as

mean $\pm$ SD. Plots represent individual data, n = 14. $^{***}p < 0.001.$ (B) The correlations between

cycling speeds and baseline-normalized changes in salivary oxytocin levels. Blue circles, the Ex

condition; red circles, the Ex+AR condition.

Fig. 5. The correlations of baseline-normalized changes in salivary oxytocin levels with exercise-

induced changes in (A) anger-hostility, (B) confusion-bewilderment, (C) depression-ejection, (D)

fatigue-inertia, (E) tension-anxiety, (F) vigor-activity, (G) friendliness and (H) total mood

disturbance in each condition. Ex: standard solitary cycling exercise, Ex+AR: augmented-reality-

based multi-person cycling exercise. White circles, the Rest condition; blue circles, the Ex

condition; red circles, the Ex+AR condition.