1   **Global niche partitioning of purine and pyrimidine cross-feeding among ocean microbes**
2

3   R. Braakman[1],*, B. Satinsky[2,3], T.J. O'Keefe[2,4], K. Longnecker[5], S.L Hogle[2,6], J.W. Becker[2,7], K. Dooley[2,8],
4   A. Arellano[2,9], M.C. Kido Soule[5], E.B. Kujawinski[5], S.W. Chisholm[2,10]
5
[1] Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge,
MA, USA
[2] Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA
[3] current address: Dyno Therapeutics, Watertown, MA, USA
[4] current address: Institute of Marine Sciences, University of North Carolina at Chapel Hill, Morehead City, NC, USA
[5] Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA,
USA
[6] current address: Department of Biology, University of Turku, Turku, Finland
[7] current address: Department of Science, Alvernia University, Reading, PA, USA
[8] current address: Renewable Resources and Enabling Sciences Center, National Renewable Energy Laboratory,
Golden, CO, USA
[9] current address: Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA
[10] Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA
19
20   *Corresponding author: braakman@mit.edu
21

22   **Abstract**
23

Characterizing the diversity of cross-feeding pathways in ocean microbes illuminates forces shaping
co-evolution, ecosystem self-assembly and carbon cycling. Here we uncover a purine and pyrimidine
cross-feeding network in globally abundant groups. The cyanobacterium *Prochlorococcus* exudes
both compound classes, which metabolic reconstructions suggest follows synchronous daily genome
replication. Co-occurring heterotrophs differentiate into purine or pyrimidine specialists, or
generalists, and use compounds for different purposes. The most abundant heterotroph, SAR11, is a
specialist that uses purines as sources of energy, carbon and/or nitrogen, with subgroups
differentiating along ocean-scale gradients in the supply of energy and nitrogen, in turn leading to
putative cryptic nitrogen cycles that link many microbes. Finally, in a SAR11 subgroup that dominates
where *Prochlorococcus* is abundant, adenine additions to cultures inhibit DNA synthesis, poising cells
for replication. We argue this subgroup uses inferred daily pulses of adenine from *Prochlorococcus* to
metabolically synchronize to the daily supply of photosynthate from surrounding phytoplankton.
36

37   **Introduction**
38
Microbial processes and interactions lie at the heart of the oceanic carbon cycle (1-3). Carbon enters
the marine biosphere through photosynthesis (4,5), is transformed through metabolism (6) and viral
infections and lysis (7,8), is transferred to higher trophic levels through grazing (2,9), and is ultimately
broken back down and released as CO_2 through bacterial catabolism and respiration (1-3,6). An
increasingly recognized link within this global cycle of organic carbon is cross-feeding, in which cells
excrete specific compounds that are consumed by surrounding cells (10-13). Evidence that cross-

feeding is widespread in the ocean comes from the highly streamlined genomes and complex nutrient

dependencies of many highly abundant oceanic bacteria (14-16). However, partly due to the difficulty

of measuring extracellular metabolites in seawater (3), the diversity and distribution of cross-feeding

pathways has not been systematically characterized, obscuring both the ecological roles they play and

the forces shaping their evolution.

To explore this using a bounded model system, we have begun characterizing the production of

organic carbon by the highly abundant oceanic cyanobacterium *Prochlorococcus* (17) and assessing its

fate. By performing ~10-15% of oceanic CO_2-fixation (18,19), *Prochlorococcus* is a major source of

organic carbon to ocean ecosystems, and over geologic timescales is thought to have evolved

metabolic interdependencies with co-occurring heterotrophs (20). In recent experiments, we noticed

that pyrimidines and purines are both abundant exudates in *Prochlorococcus* (21). Here we further

examine the production and consumption of these compounds and uncover a globally distributed

cross-feeding network involving many abundant oceanic microbes. We begin to characterize the

forces driving large-scale niche partitioning within this network.

Results and Discussion

***Pyrimidine and purine exudation by* Prochlorococcus**

To better understand organic carbon production by *Prochlorococcus* we have measured the

extracellular abundance of a large suite of organic compounds in exponentially growing cultures of

several strains under a range of light and nutrient conditions (21). A signal standing out is that

thymidine is among the most abundant extracellular compounds across strains and culture conditions

(Fig. 1, Table S1). The exudation of high levels of this nitrogen-rich pyrimidine is unexpected, since it is

generally thought that the genome and proteome of *Prochlorococcus* are highly streamlined to

minimize the nitrogen requirements of growth in oligotrophic waters (22,23). We calculated (Methods)

that over the course of the experiment production of extracellular thymidine amounted to almost half

of what was incorporated into DNA (Fig. 1, Table S1). We also detected adenine and guanine in the

exudates of *Prochlorococcus*, and their levels were higher under phosphorus-limitation than in nutrient

replete conditions (Fig. 1, Table S1). While we could not precisely quantify these compounds due to an

extraction efficiency <1% (21), their uncorrected levels reached ~2-3% of that of thymidine, suggesting

their exudation was at a similar scale (Fig. 1, Table S1).

Fig. 1 Extra- and intracellular pyrimidines and purines in cultures of *Prochlorococcus*. Extracellular levels of the pyrimidine thymidine (orange), and the purines adenine, guanine and 5-methylthioadenosine (shades of blue are normalized to the total amounts of purines or pyrimidines incorporated into DNA and shown alongside cytosolic concentrations (gray shading) of these compounds, for three different strains of *Prochlorococcus* grown under different conditions. Error bars show the standard deviation between biological replicates. Light levels are in units of μmol photons $m^{-2} s^{-1}$, 'R' stands for batch culture growth in nutrient replete media and '–P' stands for growth under semi-continuous phosphorus limitation. Exudation levels of adenine and guanine (marked with asterisks) are uncorrected from the measured concentrations due to having an extraction efficiency of less than 1%. For a sense of scale relative to thymidine exudation levels, gray dashed bars are included at 100x the measured adenine and guanine levels (i.e. the theoretical correction factor for an extraction efficiency of 1%). Incidences in which metabolites were only detected in single replicates are denoted with 'detected', while incidences in which metabolites were not detected in any replicate are denoted with 'nd'.

To understand the excretion of thymidine, adenine and guanine, we searched for possible pathways producing and/or consuming them in *Prochlorococcus*. The only pathway we identified involving these compounds (Methods, Text S1) is a putative pathway for recycling deoxyribonucleotides, the direct precursors to DNA (Fig. 2A). There thymidine, deoxyadenosine and deoxyguanosine are generated from dTMP, dAMP and dGMP, respectively, through the action of 5'-nucleotidase (SurE) (24). Deoxyadenosine and deoxyguanosine are then further processed to adenine and guanine through the action of methylthioadenosine phosphorylase (MTAP), and then to AMP and GMP through the action of adenine phosphoribosyltransferase (apt), together forming a salvage pathway also observed in other systems (25). We could not identify other genes involved in processing thymidine, suggesting it forms an endpoint and is not further processed after the first (phosphate-salvaging) step of the pathway, perhaps because it is only involved in DNA synthesis, while deoxyadenosine and deoxyguanosine can be repurposed for ATP/GTP and RNA synthesis. This is consistent with the fact that we never measured intracellular thymidine, suggesting it is rapidly excreted after the phosphate is removed from dTMP. Original genome annotations assigned a narrower functionality to MTAP (i.e.

acting only on methylthioadenosine) than we hypothesize here, but this enzyme is known to act on

multiple substrates in other microbial systems, including as part of the pathway we identify here

(25,26). Enzyme multi-functionality is generally thought to be an important strategy in bacteria with

streamlined genomes like *Prochlorococcus* (27,28). A lack of other identifiable pathways thus leads us

to tentatively conclude that thymidine, adenine and guanine are all intermediates or endpoints of a

shared, broad specificity deoxyribonucleotide recycling pathway (Fig. 2A).

114
Fig. 2 Putative deoxyribonucleotide recycling pathway in *Prochlorococcus*. A) Structure of the deoxyribonucleotide

recycling pathway in relation to pathways for DNA and RNA synthesis and polyamine metabolism. Metabolites highlighted in

boxes are exuded by *Prochlorococcus* (Fig. 1). **B)** transcriptional dynamics of genes involved in the deoxyribonucleotide

recycling pathway, as well as DNA and RNA polymerase genes, in *Prochlorococcus* MED4 cells synchronized to a diurnal

light:dark cycle (29). In both panels deoxyribonucleotide recycling pathway reactions/genes are highlighted using the orange

spectrum, and DNA and RNA polymerase are highlighted in shades of blue. Abbreviations: NDP = nucleoside diphosphate,

dNDP = deoxynucleoside diphosphate, NTP = nucleoside triphosphate, dNTP = deoxynucleoside triphosphate, dTMP =

deoxythymidine monophosphate, dGMP = deoxyguanosine monophosphate, dAMP = deoxyadenosine monophosphate, dA =

deoxyadenosine, MTA = 5-methyl-thioadenosine, SAMA = S-adenosylmethioninamine, PUT = putrescine, SPE = spermidine,

MTRP = methyl-5-thioribose, PRPP = phosphoribose diphosphate, Pi = orthophosphate, SurE = survival protein E (5'-

nucleotidase), MTAP = methylthioadenosine phosphorylase, apt = adenine phosphoribosyltransferase, SpeE = spermidine

synthase.

To further explore the possible existence of a deoxyribonucleotide recycling pathway in

Prochlorococcus we examined existing data on the expression dynamics of the relevant genes in

transcriptional data obtained from a strain whose growth was synchronized to a diurnal light:dark cycle
(29). Under such conditions, metabolic processes are segregated in time, allowing us to examine the
potential role of the putative deoxyribonucleotide recycling pathway in the broader processes of
growth and cellular replication. The expression of all three genes (SurE, MTAP, apt) in the putative
pathway are synchronized and maximally expressed just after dusk, when genome replication is
complete and cells transition to RNA synthesis (Fig. 2B), consistent with their collective functioning in
deoxyribonucleotide recycling.
137
Properties of the deoxyribonucleotide recycling pathway are also consistent with observed metabolite
dynamics in *Prochlorococcus* cells under phosphorus-limited growth conditions. That is, regenerating
AMP and GMP from adenine and guanine requires phosphate (Fig. 2A), and adenine accumulates
intracellularly in phosphate-limited relative to nutrient-replete cells (Fig. 1). This suggests that
exudation of adenine and guanine results from the emergence of a bottleneck in the
deoxyribonucleotide recycling pathway due to a depletion of intracellular phosphate. This is consistent
with the observation that methylthioadenosine, a byproduct of polyamine metabolism that is recycled
via the same pathway (Fig. 2A), also accumulates and is excreted under phosphate limited relative to
replete conditions (Fig. 1). A recent study showed that some strains of the purple sulfur bacterium
Rhodopseudomonas palustris exude adenine when a bottleneck arises in its intracellular salvage due
to low levels of the enzyme *apt* (30), highlighting a similar principle. In contrast, the pyrimidine
thymidine, which unlike purines is not internally recycled (Fig. 2), never accumulates, and its
extracellular levels are not significantly different under phosphorus stress (Fig. 1). Taken together,
metabolite, genomic and transcriptional data thus suggest that *Prochlorococcus* cells generate excess
dNTP during genome replication and excrete whatever they do not reuse or recycle when transitioning
to ribosome synthesis. This in turn implies that *Prochlorococcus* provides daily pulses of pyrimidines
and purines to the surrounding ecosystem. Time-resolved quantification of metabolites in the field and
in diel L:D synchronized cultures of *Prochlorococcus* are a key area of future study that will help
constrain both the timing of release and the relative contributions of *Prochlorococcus* to rhythms of
these substrates in the ocean ecosystem.
158
Niche partitioning of purine and pyrimidine usage among heterotrophic bacterioplankton
We next sought to understand the ecological consequences of purine and pyrimidine excretion by
Prochlorococcus. To this end, we searched for purine and pyrimidine usage genes in 305 partial and
complete genomes of SAR11, SAR86 and SAR116, each among the most abundant of
Prochlorococcus' sympatric heterotrophs (16,31-33). In SAR11, the occurrence of purine usage genes
reaches frequencies similar to the average genome completeness of ~72%, suggesting purine usage
is a universal or nearly universal trait in this genus, whereas pyrimidine usage genes occur at
frequencies of only ~3% (Fig. 3, Data S1). In contrast, in SAR86, thymidine usage genes occur at
frequencies similar to the average genome completeness of ~71%, suggesting thymidine usage is a
universal or nearly universal trait in the genus, whereas purine usage genes are absent (Fig. 3, Data
S1). Finally, in SAR116, purine usage genes occur at frequencies of ~49%, whereas pyrimidine genes
occur at frequencies of ~43% (Fig. 3, Data S1), both below the average genome completeness of
~75%. These observations suggest that resource niche partitioning within the oceanic
bacterioplankton community has given rise to purine and pyrimidine specialists, as well as generalists

that use both classes of compounds, in turn driving differentiation of cross-feeding with
Prochlorococcus.

176
Fig. 3 Niche partitioning of purine and thymidine usage in oceanic bacterioplankton. A) Phylometabolic diagrams show
presence/absence (filled/empty symbols) of genes involved in purine and pyrimidine usage in the abundant often co-occurring
heterotrophic bacterioplankton groups SAR11, SAR86 and SAR116. Rows in the phylometabolic diagrams represent the
metabolic gene profiles within individual genomes in the tree. Transporter genes are indicated with pink asterisks. Colors
represent different functions of pathways: purine assimilation genes are in yellow, purine catabolism genes in purple, allantoin

catabolism genes in green, thymidine usage genes in dark red and urea assimilation genes in light blue. SAR116 purine
phosphoribosyltransferase (PPRT), which occurs within both purine and thymidine usage gene operons (Fig. S2), is colored in
dark gray. Genome completeness statistics (%) are shown as histogram next to gene profiles. **B)** Representative genome profiles
of SAR11 sub-groups specializing in either purine assimilation (IMCC9063 and HTCC7211), purine assimilation and allantoin
catabolism (HTCC1040), or full purine catabolism (HIMB083). Genes: a = purine NCS2 permease, b = purine deaminase (tadA), c
= purine ABC transporter permease, d = purine MFS permease, e = purine phosphoribosyltransferase (PPRT), f = urate oxidase
(PuuD), g = purine deaminase (tadA), h = Xanthine dehydrogenase Mo-subunit (XDH_Mo) , i = xanthine dehydrogenase FAD-
subunit (XDH_FAD) , j = 5-hydroxyisourate lyase (hiuH), k = 2-oxo-4-hydroxy-4-carboxy-5-ureidoimidazoline decarboxylase
(OHCU decarboxylase, uraD), l = allantoin NCS1 permease, m = allantoicase (alc), n = ureidoglycolate lyase (ugl) , o =
allantoinase (PuuE), p = uracil phosphoribosyltransferase, q = cytidine deaminase, r = thymidine kinase, s = pyrimidine ABC
transporter, t = urease, u = urea ABC transporter, v = purine ABC transporter, w = purine ABC transporter, x = XDH accessory
protein, y = purine NCS2 permease, z = purine deaminase, aa = allantoin racemase, ab = ureidoglycine aminohydrolase, ac =
purine TRAP transporter, ad = DMT transporter, ae = purine/pyrimidine phosphoribosyltransferase, af = CNT transporter.
195

Niche partitioning of purine and pyrimidine cross-feeding occurs across multiple taxonomic scales. For
example, while at the genus-level SAR116 is a purine and pyrimidine generalist, it also contains
subgroups that have genes for only purine or pyrimidine usage, but not both (Fig. 3, Fig. S2). In
addition, different parts of the SAR116 tree contain variations in the pathway of purine catabolism that
are characterized by differences in the presence or absence of allantoin racemase and the production
of either ammonia or urea in the final steps of the pathway (Fig. 3, Fig. S3), as well as an association
with primary, ATP-consuming transporters or with secondary, ion-gradient-driven transporters (Fig. 3).
203

Similarly, different SAR11 sub-groups have gene profiles suggesting they use purines for different
purposes. For example, most genes making up a large operon for purine catabolism (34) occur in ~18-
22% of genomes, but an uptake transporter and a purine phosphoribosyltransferase that together
perform the initial assimilatory portion of the pathway occur at frequencies similar to the average
genome completeness of ~72% (Fig. 3, Data S1), suggesting they are universal or nearly universal. This
suggests that some SAR11 cells break purines down to their constituent components, while others use
them intact. Further, an allantoin transporter occurs in ~12% of genomes (Data S1), including in at least
10 genomes that lack the full purine catabolism operon but retain genes involved in allantoin
catabolism (Fig. 3), leading to an overall higher frequency of allantoin catabolism genes (~22%) than
upstream purine catabolism genes (~18%) in SAR11 (Data S1). Allantoin is the product of the first
sequence of reactions of purine catabolism in which one of the rings is cleaved and energy is released,
while its further degradation releases carbon and nitrogen (Fig. 4). The occurrence of this allantoin-
specialist SAR11 subgroup raises the possibility that in some environments other SAR11 cells primarily
in search of energy incompletely degrade purines, releasing allantoin that is taken up and further
degraded by surrounding cells (Fig. 4).
219

Fig. 4. Purine catabolism in SAR11 and associated putative purine:urea-mediated cryptic nitrogen cycles.
Prochlorococcus, Thaumarcheota and potentially *Synechococcus* (see text) supply purines that are used by SAR11. Different
SAR11 cell types, labeled by representative strains (Fig. 3), possess variants of purine catabolism as shown using sequences of
reactions that perform different purine usage functions. Colors of purine usage functions are as defined in Fig. 3: purine
assimilation genes in yellow, purine catabolism genes in purple and allantoin catabolism genes in green. Many SAR11 cells lack
genomic capacity to use urea produced during purine catabolism, leading to its release and availability to surrounding groups,
including *Prochlorococcus, Thaumarcheota* and *Synechococcus*. Some SAR11 cells lack purine catabolism but have genomic
capacity to use allantoin, which is inferred to be released by other SAR11 cells that incompletely catabolize purines. OHCU = 2-
oxo-4-hydroxy-4-carboxy-5-ureidoimidazoline. Reactions marked with a red asterisk represent enzymes that in subsets of SAR11
genomes are collocated with transporter genes (Fig. 3). Black arrows represent inferred cross-feeding pathways.

In contrast to the diversity in pyrimidine/purine usage functions observed in SAR116 and SAR11, nearly
all SAR86 genomes contain just two, often-collocated thymidine usage genes: thymidine kinase and a
thymidine transporter (Fig. 3, Fig. S2). SAR116 genomes, as well as some deep-branching SAR11
genomes, contain other genes for interconverting among pyrimidines that are collocated with
thymidine kinase and the transporter, including uridine ribohydrolase, uracil
phosphoribosyltransferase and cytidine deaminase, but these are absent in SAR86 (Fig. 3, Fig. S2). This
suggests that while SAR116 may have a more general pyrimidine usage strategy, SAR86 focuses
specifically on thymidine assimilation.
239
Niche partitioning of purine usage strategies in SAR11
Since SAR11 is the most abundant of *Prochlorococcus*' sympatric heterotrophs (31,33), we decided to
examine its diversity of purine usage strategies more closely. As noted, purine catabolism in SAR11
leads to three products: energy in the form of NADH, carbon in the form of glyoxylate, and nitrogen in
the forms of ammonia and urea (Fig. 4). However, the bulk of nitrogen released during purine
catabolism is released as urea, whereas assimilation of nitrogen into biomolecules universally starts
from ammonia. To use nitrogen in urea, cells across the tree of life must first break it down to ammonia
using urease. We searched SAR11 genomes for urease genes, and found they occur at a much lower
frequency (~5%) than purine catabolism genes (~18-22%) (Fig. 3, Data S1). This suggests additional
niche partitioning of purine catabolism within SAR11, with some cells using it to obtain nitrogen, while
others use it primarily to obtain energy and/or carbon, releasing urea back to the environment.
Genomes in our sample that contain allantoin catabolism genes but lack the rest of the purine
catabolism operon also lack urease (Fig. 3), similarly suggesting that these cells use allantoin as a
carbon source and release urea back to the environment.
254
These observations suggest the existence of purine:urea-mediated cryptic nitrogen cycles involving
many abundant microbial groups (Fig. 4). That is, in addition to exuding purines (21) (Fig. 1),
Prochlorococcus is also a major urea user (35,36). This creates a putative cycle in which purines transfer
energy and carbon from *Prochlorococcus* to SAR11 and urea transfers nitrogen back to
Prochlorococcus. Incomplete purine catabolism by some SAR11 cells leading to potential transfer of
allantoin to other surrounding SAR11 cells (Fig. 3 and surrounding discussion) may add an additional
step to this cycle in some environments (Fig. 4). Similarly, marine *Synechococcus* also uses urea
(35,36), and while purine and thymidine exudation have not been documented in this genus, a marine
strain of *Synechococcus elongatus* was found to exude thymidine (37), suggesting the
deoxyribonucleotide recycling pathway we identified in *Prochlorococcus* (Fig. 2) could have a broader
distribution across other cyanobacteria. Further, ammonia-oxidizing archaea (Thaumarcheota), which
are highly abundant autotrophic (CO_2-fixing) microbes lying at the base of the food web below the
euphotic zone (38), also exude thymidine and purines (39) and use urea (40). Finally, SAR11 has a
broad geographic range that extends across all these groups, from coastal to open waters and from
the surface to the deep (41,42). Together these observations suggest that purine:urea-mediated
cryptic nitrogen cycles (Fig. 4) may operate throughout much of the world's oceans, linking many
abundant groups, warranting their further study in both laboratory and field settings.
272

To further disentangle the forces shaping SAR11 purine usage functions and associated putative
nitrogen cycles, we next examined the abundance of relevant genes in metagenomes collected from
diverse oceans. We estimated the fraction of SAR11 cells with the capacity for different purine usage
strategies in different environments by normalizing the abundance of purine usage genes to a set of
single-copy core genes (Methods). We found that genes involved in purine assimilation are always
abundant and have a relatively uniform distribution geographically, with depth, and as a function of
nutrient concentrations (Fig. 5, Fig. S4, Table S4), consistent with their near-universal presence in
SAR11 genomes (Fig. 3).
281

282
Figure 5. Distribution of SAR11 purine usage genes. Metagenomically-derived estimates for the gene/genome frequency of
different purine usage genes in SAR11 are shown as a function of **A)** surface (i.e. upper 100 meters) biogeography, **B)** depth in
the water column in the North Atlantic, and **C)** inorganic nitrogen concentration. Purine usage functions are color coded as
defined in Fig. 3. For surface biogeography panels each data point represents the mean across samples within the upper 100
meters of the water column at a given station. Spearman rank correlation coefficients (r_S) of the correlation between
gene/genome frequency and nitrogen concentration are shown as insets in panels on the right.
289

Genes mediating other purine usage functions in SAR11 have distinct, non-uniform distributions:
genes of the carbon- and energy-harvesting portions of the pathway increase in frequency in
environments where nutrients are elevated, including in surface regions in the Arabian Sea, the
upwelling regions of the Eastern Equatorial Pacific and the Banguelas system off the tip of South Africa,
and the Amazon River plume off the northern coast of South America (Fig. 5, Fig. S4). Genes involved

in harvesting nitrogen from purines follow the opposite pattern, reaching their highest abundances in
environments where nutrients are depleted (Fig. 5, Fig. S4). While energy and carbon harvesting can
proceed without the downstream process of nitrogen harvesting, nitrogen harvesting can only
proceed if the full pathway is present (Fig. 4). Hence, the similar abundance of genes for both functions
in the nutrient-poor open ocean – where the abundances of genes for carbon/energy-harvesting reach
their minimum – suggests that in those environments SAR11 primarily uses purine catabolism to obtain
nitrogen.
302
Further insight into the forces underlying niche partitioning come from the depth distribution of purine
catabolism functions in SAR11, as seen most clearly in the North Atlantic (Fig. 5, Fig. S4). That is, both
energy- and carbon-harvesting genes are most abundant at depth, where inorganic nutrient
concentrations increase but the energy supply to heterotrophs from primary producers decreases. In
contrast, nitrogen-harvesting genes are most abundant near the surface where energy is more
available, but nutrients are depleted. Taken together these observations suggest that niche
partitioning of SAR11 purine usage functions occurs along a central axis defined by the relative
availability of nitrogen and energy. This is consistent with the different transporters associated with the
different functions: nitrogen-harvesting genes are associated with an ABC transporter, a high-affinity
but energetically costly system requiring ATP hydrolysis (43), while both energy- and carbon-
harvesting genes are associated with more energetically efficient nucleotide:cation symporters (44)
(Fig. S4, Table S4).
315
While genes for the allantoin transporter characteristic of the allantoin-using SAR11 subgroup (Figs
3,4) follow a broadly similar trend as genes involved in energy-harvesting, they are nearly absent at the
surface where energy-harvesting genes remain present (Fig. 5, Fig. S4). The frequency of carbon-
harvesting genes also has a stronger positive correlation with nutrient concentrations than energy-
harvesting genes (Fig. 5, Fig. S4, Table S4). These observations are consistent with a decoupling of
energy- and carbon-harvesting portions of the purine catabolism pathway in some SAR11 cells as the
total flux of purine catabolism increases, thereby making allantoin available to other surrounding cells
(Fig. 4). This is further consistent with the use of symporter systems to acquire purines and allantoin at
depth, as it suggests that levels of both are higher in deeper waters than at the surface.
325
Some of the patterns of niche partitioning of SAR11 purine usage functions appear to be linked to the
niche partitioning of clades, adding further support to our conclusions. For example, clades IC and IIB,
which dominate in the mesopelagic (41,42) and upper mesopelagic (41,45), respectively, have higher
frequencies of genes involved in harvesting energy (~71% in clade IC, ~47% in clade IIB) and carbon
(~57% in clade IC, ~44% in clade IIB) from purines than other clades, while also having a high
frequency of allantoin transporter genes (~57% in clade IC, ~50% in clade IIB), but lacking urease
genes (Fig. S6, Data S1). This is consistent with our conclusions that the harvesting of energy and
carbon from purines and the cross-feeding of allantoin become more prominent at depth, whereas
nitrogen harvesting from purines is restricted to surface environments (Fig. 5 and surrounding
discussion). Further, genomes of clade IA.1, which is abundant in cold surface waters that are typically
richer in nutrients (46) have high frequencies of genes involved in allantoin transport (~57%) and
catabolism (~71%), but lack genes involved in harvesting energy from purines. This is consistent with

our conclusion that in some environments other SAR11 cells more focused on obtaining energy from
purines make allantoin available to surrounding cells (Figs 3 and 4 and surrounding discussion).
Finally, clades IV, IIIA and IIA, all of which are abundant in surface waters in the Northwestern Atlantic
from spring through early fall when waters are relatively more stratified (41), lack all genes for purine
catabolism but retain genes for purine assimilation (Fig. S6, Data S1), consistent with our conclusion
that this function is key in oligotrophic surface waters. Together these patterns suggest that
differentiation of purine usage strategies played a role in the broader metabolic differentiation of
SAR11 clades.
346
Inferred role of purines in the synchronization of ocean-surface populations of SAR11 to daily
rhythms in the supply of carbon from photosynthesis
Phylogenomic and metagenomic analyses of purine usage in SAR11 pose an interesting question:
what is the function of purine assimilation (yellow pathway in Fig. 4) in cells that lack the downstream
purine catabolism pathway? In surface waters of the open ocean, where *Prochlorococcus* dominates,
the frequency of purine catabolism genes in SAR11 genomes generally remains below ~30-35% (Fig.
5), suggesting that assimilation of intact purines is the major form of purine usage in these populations.
To examine possible benefits of purine assimilation, we cultured SAR11 HTCC7211, a strain
possessing assimilation genes but lacking purine catabolism genes (Fig. 3), in media amended with
adenine at a range of concentrations. Since purines are nitrogen-rich and our metagenomic analyses
suggest that SAR11 cells near the surface are often in search of nitrogen (Fig. 5), we performed these
experiments in both nutrient-balanced (pyruvate:glycine:methionine = 1:1:0.2) and glycine-depleted
(pyruvate:glycine:methionine = 50:1:10) media (see Methods). We expected that by lowering the need
for *de novo* purine synthesis, SAR11 cultures might receive a modest growth boost from the adenine
additions, reflected either as an increased growth rate and/or a higher final culture density. However,
we did not observe growth stimulation, but rather a decrease in growth rate at higher adenine
concentrations (Fig. 6). The decrease in growth rate occurred at lower adenine concentrations in
glycine-depleted than in nutrient-balanced cultures (Fig. 6).
365

366
Figure 6. Growth and cell physiology of SAR11 HTCC7211 grown with and without adenine. Top panels: growth curves of
SAR11 cells in nutrient-balanced (pyruvate:glycine:methionine = 1:1:0.2) and glycine-depleted (pyruvate:glycine:methionine =
50:1:10) media amended with different concentrations of adenine (shades of blue). Black bars at bottom reflect the time frame
used to determine growth rates, which are shown in the insets. Middle and bottom panels: histograms of the averages of bead-
normalized relative DNA fluorescence per cell and bead-normalized forward angle light scatter per cell (FALS), respectively, at
several time points between mid-exponential growth and stationary phase. For all panels, error bars reflect the standard
deviation across three biological replicates, while values significantly different (p<0.05, 2-tailed Student's t-test) between
adenine-treated and unamended controls are highlighted with a pink asterisk.
375

Puzzled by these results, we wondered if adenine could provide an indirect benefit to cells lacking
purine catabolism. A potentially relevant clue comes from the observation that in warm oligotrophic
waters where *Prochlorococcus* dominates, SAR11 metabolism is synchronized to the diurnal light:dark
cycle (47) despite not being a photoautotroph, and this coupling dissipates in nutrient-enhanced
upwelling regions with higher nutrients and fewer *Prochlorococcus* (48). Further, negative feedbacks
are often central in the synchronization of biological systems (49,50), and in eukaryotic systems purine
and pyrimidine additions have been used to synchronize cell cultures (51-53). These purine and
pyrimidine "block" techniques rely on the fact that the specificity of ribonucleotide reductase, which
generates all dNTP precursors to DNA (Fig. 2), is controlled through allosteric binding of dNTP
molecules (54). Consequently, imbalances in intracellular dNTP pools due to pyrimidine/purine
amendments can drive depletion in one or more dNTPs, thereby slowing down or fully inhibiting DNA
synthesis and genome replication (51). Subsequent removal of the pyrimidine/purine block
amendments by washing cultures results in cells undergoing genome replication and cell division in
synchrony (51,52). If similar mechanisms play out in bacteria, this raises the possibility that inferred
daily pulses of purines from *Prochlorococcus* (Fig. 2 and surrounding discussion) could act as a signal
that helps SAR11 synchronize its metabolism to the daily rhythms of primary production.
392
To explore this hypothesis, we first examined flow cytometric data from SAR11 cultures amended with
adenine to better understand how adenine inhibits SAR11 growth. Cells in SAR11 cultures amended
with adenine have a lower average DNA content, and DNA content decreases as adenine
concentrations increase (Fig. 6, Fig. S7). Further, cells from adenine-amended cultures are also larger
at the higher concentrations of adenine that inhibit growth (Fig. 6). This increase in size is particularly
noticeable in glycine-depleted cultures where growth inhibition sets in at lower adenine
concentrations (Fig. 6). Together these results suggest that adenine additions inhibit DNA synthesis in
SAR11, which, as the inhibition strengthens, leads to inhibition of cell division. This is similar to how
purine and pyrimidine "blocks" function in eukaryotic systems (51-53).
402
Next, we examined whether SAR11 cells whose DNA synthesis has been inhibited by adenine are
poised to replicate. We grew SAR11 cultures in media with and without adenine, washed cells in mid-
exponential growth, and compared their growth after resuspending in adenine-free media. Control
cultures grown without adenine often experienced significant growth retardation after washing, with
>60% of replicates experiencing major lags and/or never fully recovering (Fig. 7), suggesting that the
ultra-centrifugation-based washing protocol induces stress in SAR11 cells. The reasons for the
significant divergence of growth trajectories after washing among replicates are not entirely clear, but
it was a highly repeatable hallmark of cultures not exposed to adenine. In contrast, cultures treated
with adenine display fewer signs of stress after washing, with <20% of replicates experiencing lags or
failing to resume growth under most conditions tested (Fig. 7). In nutrient-balanced cultures the
benefit of adenine additions is clearest at higher adenine concentrations, while in glycine-depleted
cultures the benefit of adenine additions was clear across the full range of concentrations we
examined (Fig. 7). Higher adenine concentrations, as well as adenine amendments in glycine-depleted
relative to nutrient-balanced cultures, are both associated with greater inhibition of DNA synthesis and
cell division (Fig. 6). More consistent reignition of growth after washing of SAR11 cultures experiencing
greater inhibition of DNA synthesis is consistent with the inhibition poising cells for replication.

Figure 7. Post-washing growth of SAR11 HTCC7211 grown with and without adenine. Growth curves of SAR11 cultures grown with and without adenine, before and after washing and resuspension in adenine-free media. Three identical experiments were performed in both nutrient-balanced and glycine-depleted cultures, which are labeled using roman numerals. Each experiment included cultures grown with adenine at three different concentrations as well as an adenine-free control, with 6 biological replicates for each condition. Empty entries at given concentrations in different rows represent adenine concentrations not included in different experiments. Fractions of replicates experiencing a lag of 2 or more days after washing at a given adenine concentration and culture condition are shown as inset within each experimental panel, and as fractions of totals across experiments at bottom. Statistically significant differences between adenine amended and adenine-free cultures according to a 2-tailed Fisher's exact test are shown as single pink asterisks for $p<0.05$ and double pink asterisks for $p<0.01$.

From these observations, a conceptual model emerges in which SAR11 populations in surface waters of the open ocean use purines as a signal to achieve synchrony to daily rhythms in the supply of energy and carbon from photosynthesis (Fig. 8). In this model, inferred daily pulses of purines

originating from *Prochlorococcus* (and possibly others) at night (Fig. 2) inhibit SAR11 metabolism,
while the daily supply of photosynthate during daytime hours positively stimulates SAR11 metabolism.
Over long timescales these alternating signals act in concert to drive synchronization to Earth's
rotations, as reflected in *in situ* gene expression (47), allowing SAR11 to achieve greater coupling to
the daily supply of solar energy, thereby providing the selective force that maintains purine
assimilation genes despite loss of other purine catabolism genes (Fig. 3). In this context, it is worth
reconsidering SAR86's genomic profile for using thymidine, which our metabolic reconstruction
suggests is also released by *Prochlorococcus* in nightly pulses (Fig. 2). That is, SAR86 has a genomic
profile that suggests it focuses specifically on assimilating – and not interconverting, or breaking down
– thymidine (Fig. 2), and in that sense is like SAR11 populations that possess purine assimilation genes
but lack purine catabolism genes. Further, as in SAR11, *in situ* gene expression indicates that SAR86
metabolism is synchronized to the diurnal light:dark cycle in oligotrophic waters (47), but not in
regions of upwelling where *Prochlorococcus* is less abundant (48). This raises the possibility that
thymidine could play a similar metabolic role in SAR86 that adenine plays in SAR11, further
highlighting the potential value of field studies of the dynamics of dissolved purines and pyrimidines,
as well as of the coupled dynamics of *Prochlorococcus* and sympatric heterotrophs.

451
452   **Fig. 7. Hypothesized role of purines in driving synchronization of SAR11 metabolism to the diurnal light:dark cycle.**
453   Organic carbon produced by phytoplankton during the day (photosynthate, blue) stimulates SAR11 metabolism, while purines
454   (orange) produced by *Prochlorococcus* at night after genome replication (Fig. 2) inhibit DNA synthesis in SAR11 (Fig. 6). Daily
455   alternation of these positive and negative signals over long timescales drives synchronization to the 24-hour period (47).
456
457

Methods

Determining the scale of pyrimidine and purine production and exudation by Prochlorococcus

Data on the intra- and extracellular abundance of metabolites in *Prochlorococcus* were obtained in Ref
(21) (also available at MetaboLights (https://www.ebi.ac.uk/metabolights/MTBLS567). Briefly, in our
previous study (21), spent media was obtained in mid-exponential growth from nutrient replete batch
cultures of *Prochlorococcus* strains MIT9301, MIT0801 and MIT9313 grown at multiple light levels, as
well as from semi-continuous P-limited cultures of *Prochlorococcus* MIT9301. Intracellular metabolites
were directly extracted from cells using organic solvents and water. Metabolites from the spent media
were first extracted using solid phase extraction. Extracts were then analyzed using UPLC-MS/MS to

determine concentrations. For extracellular metabolites, concentrations in the extracts were corrected
for the extraction efficiency using the formula: $[m]_{media} = 100 \times [m]_{extract} / EF\%$, where $[m]_{media}$ is the
concentration of metabolite m in the spent media, $[m]_{extract}$ is the measured concentration in the
extract, and EF% (range = 0 – 100) is the efficiency with which the solid phase extraction resinextracts
metabolite m from seawater (55). This correction is only performed for metabolites with EF% > 1 due
to associated uncertainties in precisely determining extraction efficiencies when they are low (21).
While thymidine and methylthioadenosine have an EF%>1, adenine and guanine have an EF%<1, and
so their concentrations are left uncorrected.
477
478 Next, extracellular concentrations of purines and thymidine were normalized to the total amount of
479 these compounds incorporated into DNA over the course of the experiment using experimentally
480 determined cell counts and known GC% of the genomes for each strain, and assuming one
481 chromosome per *Prochlorococcus* cell (29). Thymidine, adenine and methylthioadenosine were
482 normalized to the AT% of genomes, while guanine was normalized to the GC% of genomes.
483
For intracellular purines and pyrimidines, concentrations were measured in units of ng/mL in the cell
extract and converted to fg/cell using the extract volume and the total number of cells captured on the
filter. Total cell numbers are obtained by multiplying culture volume and culture density as determined
by flow cytometry. Concentrations in units of fg/cell are then divided by the molar mass of metabolites
to obtain concentrations in units of moles/cell. Finally, concentrations in units of moles/cell are divided
by cellular volume to obtain molar concentrations. Cellular volumes for *Prochlorococcus* were
determined as follows. The volumes of *Prochlorococcus* strains MIT9313 (0.44 μm^3) and MED4 (0.2
μm^3) were measured in Ref (56), while the cellular dry masses of strains MIT9301 (60 femtograms),
MED4 (66 femtograms), NATL2A (91 femtograms) and MIT9313 (158 grams) were measured in Ref
(57). The volume of MIT9301 was estimated from that of MED4 by scaling by the relative dry weight
masses of the two strains (i.e. 60 fg/cell for MIT9301 and 66 fg/cell for MED4, giving a ratio of 60/66),
leading to 0.18 μm^3 for MIT9301. While there are no volume or mass measurements of MIT0801 it
belongs to the same "low light adapted I" (LLI) ecotype as NATL2A and so we estimated its volume
from that of MED4 by scaling by the relative dry weight masses of NATL2A and MED4 (i.e. 91 fg/cell for
NATL2A and 66 fg/cell for MED4, giving a ratio of 91/66), leading to 0.28 μm^3 for MIT0801.
499
500 *Reconstructing putative deoxyribonucleotide recycling pathway*
501
To identify possible pathways involved in the production and/or consumption of thymidine, adenine
and guanine, we searched the annotated genomes of all *Prochlorococcus* strains available on the
KEGG database (58) for all possible genes involved in the metabolism of these compounds. All strains
used in this study (MIT9301, MIT0801 and MIT9313) are included on KEGG. To place findings in a
broader context we also searched all *Prochlorococcus* genomes for all genes involved in the
metabolism of nucleobases, nucleosides, deoxynucleosides, ribonucleotide monophosphates and
deoxyribonucleotide monophosphates, all intermediates of a larger nucleotide metabolic network that
thymidine, adenine and guanine could be a part of. To account for the possibility of gene
misannotations or incorrect functional assignments in some *Prochlorococcus* strains, homologs
identified in some *Prochlorococcus* strains were used to query all *Prochlorococcus* genomes using the

BLASTp algorithm within the KEGG BLAST feature. Finally, to identify potentially relevant genes that
are present but not functionally assigned in any *Prochlorococcus* genome, we also blasted reference
gene sequences from *E. coli* (which has an extensive network of nucleotide metabolism) against
Prochlorococcus genomes. We included in our search FAMIN, a recently characterized enzyme that is
highly multi-functional and catalyzes multiple reactions within nucleotide metabolism (59). Together
this led to a list of 9 genes, 5 of which are universal in all strains used in this study (Table S2). Data on
the temporal transcriptional profiles of all 9 genes were obtained from a study in which the growth of
Prochlorococcus MED4 was synchronized to diurnal L:D conditions (29). To evaluate transcriptional
dynamics of these genes relative to DNA and RNA synthesis, the transcriptional profile of each gene
was plotted alongside the transcriptional profile of DNA polymerase and RNA polymerase (Fig. S1).
Pearson correlation coefficients of the expression of each gene relative to DNA and RNA polymerase
genes were calculated in Excel (Table S3).

Phylometabolic analysis of purine and pyrimidine strategies in heterotrophs

Genomes, curated genome phylogenies and associated metadata for SAR11, SAR86 and SAR116 were
obtained from the MARMICRODB database (60). The MARMICRODB database includes genomes from
a variety of sources, including isolate genomes, single-cell amplified genomes (SAGs) and
metagenome assembled genomes (MAGs), but genomes with %completeness – 5x %contamination <
30 were excluded in the final dataset, resulting in 186 SAR11 genomes, 57 SAR86 genomes and 59
SAR116 genomes. These genomes were first searched for the presence of purine and pyrimidine
usage genes using reference sequences in other organisms obtained from the KEGG database (58).
This led to the identification of homologs in SAR11, SAR86 and SAR116 that typically co-occur in
operons. Homologs from those operons were subsequently used for local searches using the BLASTp
algorithm obtained from NCBI (61), with final counting of gene frequencies relying on a combination
of sequence similarity and genomic context. For example, many SAR11 and SAR116 genomes that
contain a complete purine catabolism operon contain a second homolog also annotated as 5-
hydroxyisourate lyase, but which occurs in a different genomic location and has low sequence
similarity to the homolog occurring in the operon. Since this gene further occurs in many genomes that
do not contain any other purine catabolism genes, we conclude it has a different function and leave it
out of our final tally. All purine and pyrimidine usage genes thus identified are listed in Data S1, and
their presence/absence data for individual genomes was mapped onto the leaves of the SAR11,
SAR86 and SAR116 genome phylogenies to generate phylometabolic trees.

Metagenomic analyses

675 metagenomes were obtained from TARA and BioGeoTRACES (62,63). These metagenomes were
searched for SAR11 purine usage genes and SAR11 ribosomal proteins using custom Hidden Markov
Models (HMM's) that were created as follows. First, a set of ~20 representative sequences for the
genes of each individual step in the purine catabolism pathway (Figs. 3,4), as well as for a set of 14
single-copy ribosomal proteins (64) were obtained at Uniprot (65) and assembled into HMM's. For
phosphoribosyltransferase, which has many homologs operating on different nucleobases, the initial
set was very narrowly chosen to consist of primarily SAR11 variants occurring with purine catabolism

operons. Resulting HMM's were searched against the Uniprot database using the PHMMER algorithm
within the HMMER web portal (66), and results were manually inspected to determine e-value cut offs
(Data S2) that captured maximal diversity while eliminating homologous genes associated with other
functions. The original 20-gene HMM's were then searched against the MARMICRODB database
(33,60) and all sequences falling below the e-value cutoffs determined from HMMER results were
included our final HMM's to ensure they have broad coverage of diversity within the extant microbial
oceans. Next, our custom HMM's were searched against our full set of metagenomes using the GraftM
algorithm (64). The results of this search were again manually inspected to determine GraftM cutoff
scores (Data S2) that mostly eliminate homologous genes associated with other functions while mostly
capturing sequences of interest. To estimate gene/genome frequencies of individual purine usage
genes in SAR11, the counts for each purine usage gene were then normalized to the HMM amino acid
length and to the median count of the 14 single-copy ribosomal protein genes. Chemical and
hydrographic data associated with the BioGEOTRACES metagenomes (63) were obtained from the
GEOTRACES Intermediate Data Product IDP2017 version 2 (accessed January 2019), specifically from
sections GA02 (67,68) GA03, GA10 (69), and GP13. Environmental data from the Tara Oceans project
was obtained from https://doi.pangaea.de/10.1594/PANGAEA.875579. Metagenome sequence data
were quality controlled, environmental data matched to metagenomic samples, and missing
environmental data were imputed as described in detail in the supplementary material of Hogle et al.
2022 (70). Spearman-rank correlations between nutrient concentrations and gene frequencies were
determined in R.
575
576 *SAR11 adenine amendment experiments*
577
Cells of SAR11 strain HTCC7211 were cultured under constant light at 22 °C in ProMS medium (33).
ProMS is a seawater-based modified version of the artificial AMS1 medium (15), in which the
concentrations of the main sources of carbon (pyruvate), nitrogen (glycine) and sulfur (methionine)
have been lower 50-fold to promote detection of the impact of added organic substrates or
phytoplankton exudates (33). In addition, HTCC7211 cells were cultured in a glycine-depleted version
of ProMS in which the concentration of all nutrients other than glycine was increased 50-fold to bring
them back to the levels of the AMS1 medium. These cultures were then amended with adenine at
concentrations ranging seven orders of magnitude, from 10 pM to 100 mM, and their growth was
tracked alongside that of adenine-free control cultures for an entire growth curve. Culture density and
purity and cell size and DNA content were measured using flow cytometry. Because sub-populations in
different phases of the cell cycle are difficult to distinguish in SAR11 (15), we instead quantified the
averages and standard deviations over the total bulk fluorescence distributions (Fig. S7). To examine
whether adenine exposure poises SAR11 cells for replication, cultures grown with and without adenine
were harvested in mid-exponential growth and washed three times using adenine-free culture media
and ultracentrifugation at ~25,000 x g (15,000 rpm) for 45 minutes at 22 °C. After the final washing
step cell pellets were resuspended in adenine-free media and subsequent growth was monitored via
flow cytometry. Fisher's exact tests of the significance in the differences of the number of cultures
experiencing lags were determined in R.
596
597

Acknowledgements

We thank Eli Salcedo for help in generating relative DNA fluorescence/cell histograms from raw flow cytometry data for SAR11 adenine amendment experiments, Gretchen Swarr for help in the processing of samples for metabolomics analysis, and Steve Biller for discussions regarding genome replication in bacteria. This research was supported by grants from the Simons Foundation (Award ID 509034SCFY20 to R.B. and S.W.C., Award ID 509034FY20 to E.B.K., and SCOPE Award ID 329108 to M.J. Follows) and from the National Science Foundation (OCE-2019589 to R.B.). This is the NSF Center for Chemical Currencies of a Microbial Planet (C-CoMP) publication #33.

References

1. Pomeroy, L.R., 1974. The ocean's food web, a changing paradigm. *Bioscience*, *24*(9), pp.499-504.
2. Azam, F., Fenchel, T., Field, J.G., Gray, J.S., Meyer-Reil, L.A. and Thingstad, F., 1983. The ecological role of water-column microbes in the sea. *Marine ecology progress series. Oldendorf*, *10*(3), pp.257-263.
3. Moran, M.A., Kujawinski, E.B., Schroer, W.F., Amin, S.A., Bates, N.R., Bertrand, E.M., Braakman, R., Brown, C.T., Covert, M.W., Doney, S.C, Dyhrman, S.T., A.S. Edison, A.M. Eren, N.M. Levine, L. Li, A.C. Ross, M.A. Saito, A.E. Santoro, D. Segrè, A. Shade, M.B. Sullivan and A. Vardi, 2022. Microbial metabolites in the marine carbon cycle. *Nature microbiology*, *7*(4), pp.508-523.
4. Field, C.B., Behrenfeld, M.J., Randerson, J.T. and Falkowski, P., 1998. Primary production of the biosphere: integrating terrestrial and oceanic components. *science*, *281*(5374), pp.237-240.
5. Falkowski, P.G. and Raven, J.A., 2013. *Aquatic photosynthesis*. Princeton University Press.
6. Braakman, R. and Smith, E., 2012. The compositional and evolutionary logic of metabolism. *Physical biology*, *10*(1), p.011001.
7. Fuhrman, J.A., 1999. Marine viruses and their biogeochemical and ecological effects. *Nature*, *399*(6736), pp.541-548.
8. Wilhelm, S.W. and Suttle, C.A., 1999. Viruses and nutrient cycles in the sea: viruses play critical roles in the structure and function of aquatic food webs. *Bioscience*, *49*(10), pp.781-788.
9. Ward, B.A., Dutkiewicz, S., Jahn, O. and Follows, M.J., 2012. A size-structured food-web model for the global ocean. *Limnology and Oceanography*, *57*(6), pp.1877-1891.
10. Durham, B.P., Sharma, S., Luo, H., Smith, C.B., Amin, S.A., Bender, S.J., Dearth, S.P., Van Mooy, B.A., Campagna, S.R., Kujawinski, E.B. and Armbrust, E.V., 2015. Cryptic carbon and sulfur cycling between surface ocean plankton. *Proceedings of the National Academy of Sciences*, *112*(2), pp.453-457.
11. Segev, E., Wyche, T.P., Kim, K.H., Petersen, J., Ellebrandt, C., Vlamakis, H., Barteneva, N., Paulson, J.N., Chai, L., Clardy, J. and Kolter, R., 2016. Dynamic metabolic exchange governs a marine algal-bacterial interaction. *elife*, *5*, p.e17473.
12. Fritts, R.K., McCully, A.L. and McKinlay, J.B., 2021. Extracellular metabolism sets the table for microbial cross-feeding. *Microbiology and Molecular Biology Reviews*, *85*(1), pp.e00135-20.
13. Giri, S., Oña, L., Waschina, S., Shitut, S., Yousif, G., Kaleta, C. and Kost, C., 2021. Metabolic dissimilarity determines the establishment of cross-feeding interactions in bacteria. *Current Biology*, *31*(24), pp.5547-5557.
14. Giovannoni, S.J., Tripp, H.J., Givan, S., Podar, M., Vergin, K.L., Baptista, D., Bibbs, L., Eads, J., Richardson, T.H., Noordewier, M. and Rappé, M.S., 2005. Genome streamlining in a cosmopolitan oceanic bacterium. *Science*, *309*(5738), pp.1242-1245.

15. Carini, P., Steindler, L., Beszteri, S. and Giovannoni, S.J., 2013. Nutrient requirements for growth of the
extreme oligotroph 'Candidatus Pelagibacter ubique'HTCC1062 on a defined medium. *The ISME journal*,
7(3), pp.592-602.

16. Swan, B.K., Tupper, B., Sczyrba, A., Lauro, F.M., Martinez-Garcia, M., González, J.M., Luo, H., Wright, J.J.,
Landry, Z.C., Hanson, N.W. and Thompson, B.P., 2013. Prevalent genome streamlining and latitudinal
divergence of planktonic bacteria in the surface ocean. *Proceedings of the National Academy of Sciences*,
110(28), pp.11463-11468.

17. Chisholm, S.W., Olson, R.J., Zettler, E.R., Goericke, R., Waterbury, J.B. and Welschmeyer, N.A., 1988. A
novel free-living prochlorophyte abundant in the oceanic euphotic zone. *Nature*, *334*(6180), pp.340-343.

18. Flombaum, P., Gallegos, J.L., Gordillo, R.A., Rincón, J., Zabala, L.L., Jiao, N., Karl, D.M., Li, W.K., Lomas,
M.W., Veneziano, D. and Vera, C.S., 2013. Present and future global distributions of the marine
Cyanobacteria Prochlorococcus and Synechococcus. *Proceedings of the National Academy of Sciences*,
110(24), pp.9824-9829.

19. Letscher, R.T., Moore, J.K., Martiny, A.C. and Lomas, M.W., 2023. Biodiversity and Stoichiometric Plasticity
Increase Pico-Phytoplankton Contributions to Marine Net Primary Productivity and the Biological Pump.
Global Biogeochemical Cycles, *37*(8), p.e2023GB007756.

20. Braakman, R., Follows, M.J. and Chisholm, S.W., 2017. Metabolic evolution and the self-organization of
ecosystems. *Proceedings of the National Academy of Sciences*, *114*(15), pp.E3091-E3100.

21. Kujawinski, E.B., Braakman, R., Longnecker, K., Becker, J.W., Chisholm, S.W., Dooley, K., Kido Soule, M.C.,
Swarr, G.J. and Halloran, K., 2023. Metabolite diversity among Prochlorococcus strains belonging to
divergent ecotypes. *mSystems*, e01261-22.

22. Dufresne, A., Garczarek, L. and Partensky, F., 2005. Accelerated evolution associated with genome
reduction in a free-living prokaryote. *Genome biology*, *6*, pp.1-10

23. Grzymski, J.J. and Dussaq, A.M., 2012. The significance of nitrogen cost minimization in proteomes of
marine microorganisms. *The ISME journal*, *6*(1), pp.71-80.

24. Proudfoot, M., Kuznetsova, E., Brown, G., Rao, N.N., Kitagawa, M., Mori, H., Savchenko, A. and Yakunin,
A.F., 2004. General enzymatic screens identify three new nucleotidases in Escherichia coli: biochemical
characterization of SurE, YfbR, and YjjG. *Journal of Biological Chemistry*, *279*(52), pp.54687-54694.

25. Vodnala, M., Ranjbarian, F., Pavlova, A., de Koning, H.P. and Hofer, A., 2016. Trypanosoma brucei
methylthioadenosine phosphorylase protects the parasite from the antitrypanosomal effect of
deoxyadenosine: implications for the pharmacology of adenosine antimetabolites. *Journal of Biological*
Chemistry, *291*(22), pp.11717-11726.

26. Cacciapuoti, G., Bertoldo, C., Brio, A., Zappia, V. and Porcelli, M., 2003. Purification and characterization
of 5′-methylthioadenosine phosphorylase from the hyperthermophilic archaeon Pyrococcus furiosus:
substrate specificity and primary structure analysis. *Extremophiles*, *7*, pp.159-168.

27. Noell, S.E. and Giovannoni, S.J., 2019. SAR11 bacteria have a high affinity and multifunctional glycine
betaine transporter. *Environmental Microbiology*, *21*(7), pp.2559-2575.

28. Noell, S.E., Barrell, G.E., Suffridge, C., Morré, J., Gable, K.P., Graff, J.R., VerWey, B.J., Hellweger, F.L. and
Giovannoni, S.J., 2021. Sar11 cells rely on enzyme multifunctionality to metabolize a range of polyamine
compounds. *MBio*, *12*(4), pp.e01091-21.

29. Zinser, E.R., Lindell, D., Johnson, Z.I., Futschik, M.E., Steglich, C., Coleman, M.L., Wright, M.A., Rector, T.,
Steen, R., McNulty, N. and Thompson, L.R., 2009. Choreography of the transcriptome, photophysiology,
and cell cycle of a minimal photoautotroph, Prochlorococcus. *PloS one*, *4*(4), p.e5135.

30. Chuang, Y.-C. , Haas, N.W. , Pepin, R., Behringer, M., Oda, Y., LaSarre, B., Harwood, C.S., McKinlay, J.B.,
2023. A purine salvage bottleneck leads to bacterial adenine cross-feeding, bioRxiv 2023.10.17.562681

31. Morris, R.M., Rappé, M.S., Connon, S.A., Vergin, K.L., Siebold, W.A., Carlson, C.A. and Giovannoni, S.J.,
2002. SAR11 clade dominates ocean surface bacterioplankton communities. *Nature*, *420*(6917), pp.806-
810.

32. Treusch, A.H., Vergin, K.L., Finlay, L.A., Donatz, M.G., Burton, R.M., Carlson, C.A. and Giovannoni, S.J.,
2009. Seasonality and vertical structure of microbial communities in an ocean gyre. *The ISME journal*,
3(10), pp.1148-1163.

33. Becker, J.W., Hogle, S.L., Rosendo, K. and Chisholm, S.W., 2019. Co-culture and biogeography of
Prochlorococcus and SAR11. *The ISME journal*, *13*(6), pp.1506-1519.

34. Haro-Moreno, J.M., Rodriguez-Valera, F., Rosselli, R., Martinez-Hernandez, F., Roda-Garcia, J.J., Gomez,
M.L., Fornas, O., Martinez-Garcia, M. and López-Pérez, M., 2020. Ecogenomics of the SAR11 clade.
Environmental Microbiology, *22*(5), pp.1748-1763.

35. Moore, L.R., Post, A.F., Rocap, G. and Chisholm, S.W., 2002. Utilization of different nitrogen sources by
the marine cyanobacteria Prochlorococcus and Synechococcus. *Limnology and oceanography*, *47*(4),
pp.989-996.

36. Berthelot, H., Duhamel, S., L'helguen, S., Maguer, J.F., Wang, S., Cetinić, I. and Cassar, N., 2019.
NanoSIMS single cell analyses reveal the contrasting nitrogen sources for small phytoplankton. *The ISME*
Journal, *13*(3), pp.651-662.

37. Fiore, C.L., Longnecker, K., Kido Soule, M.C. and Kujawinski, E.B., 2015. Release of ecologically relevant
metabolites by the cyanobacterium *Synechococcus elongatus* CCMP 1631. *Environmental microbiology*,
17(10), pp.3949-3963.

38. Karner, M.B., DeLong, E.F. and Karl, D.M., 2001. Archaeal dominance in the mesopelagic zone of the
Pacific Ocean. *Nature*, *409*(6819), pp.507-510.

39. Bayer, B., Hansman, R.L., Bittner, M.J., Noriega-Ortega, B.E., Niggemann, J., Dittmar, T. and Herndl, G.J.,
2019. Ammonia-oxidizing archaea release a suite of organic compounds potentially fueling prokaryotic
heterotrophy in the ocean. *Environmental Microbiology*, *21*(11), pp.4062-4075.

40. Kitzinger, K., Padilla, C.C., Marchant, H.K., Hach, P.F., Herbold, C.W., Kidane, A.T., Könneke, M., Littmann,
S., Mooshammer, M., Niggemann, J. and Petrov, S., 2019. Cyanate and urea are substrates for nitrification
by Thaumarchaeota in the marine environment. *Nature microbiology*, *4*(2), pp.234-243.

41. Giovannoni, S.J., 2017. SAR11 bacteria: the most abundant plankton in the oceans. *Annual review of*
marine science, *9*, pp.231-255.

42. Cameron Thrash, J., Temperton, B., Swan, B.K., Landry, Z.C., Woyke, T., DeLong, E.F., Stepanauskas, R.
and Giovannoni, S.J., 2014. Single-cell enabled comparative genomics of a deep ocean SAR11
bathytype. *The ISME journal*, *8*(7), pp.1440-1451.

43. Rees, D.C., Johnson, E. and Lewinson, O., 2009. ABC transporters: the power to change. *Nature reviews*
Molecular cell biology, *10*(3), pp.218-227

44. Weyand, S., Shimamura, T., Yajima, S., Suzuki, S.I., Mirza, O., Krusong, K., Carpenter, E.P., Rutherford,
N.G., Hadden, J.M., O'Reilly, J. and Ma, P., 2008. Structure and molecular mechanism of a nucleobase–
cation–symport-1 family transporter. *Science*, *322*(5902), pp.709-713

45. Carlson, C.A., Morris, R., Parsons, R., Treusch, A.H., Giovannoni, S.J. and Vergin, K., 2009. Seasonal
dynamics of SAR11 populations in the euphotic and mesopelagic zones of the northwestern Sargasso
Sea. *The ISME journal*, *3*(3), pp.283-295.

46. Brown, M.V., Lauro, F.M., DeMaere, M.Z., Muir, L., Wilkins, D., Thomas, T., Riddle, M.J., Fuhrman, J.A.,
Andrews-Pfannkoch, C., Hoffman, J.M. and McQuaid, J.B., 2012. Global biogeography of SAR11 marine
bacteria. *Molecular systems biology*, *8*(1), p.595.

47. Ottesen, E.A., Young, C.R., Gifford, S.M., Eppley, J.M., Marin III, R., Schuster, S.C., Scholin, C.A. and
DeLong, E.F., 2014. Multispecies diel transcriptional oscillations in open ocean heterotrophic bacterial
assemblages. *Science*, *345*(6193), pp.207-212.

48. Ottesen, E.A., Young, C.R., Eppley, J.M., Ryan, J.P., Chavez, F.P., Scholin, C.A. and DeLong, E.F., 2013.
Pattern and synchrony of gene expression among sympatric marine microbial populations. *Proceedings*
of the National Academy of Sciences, *110*(6), pp.E488-E497.

49. Gonze, D., Halloy, J. and Goldbeter, A., 2002. Robustness of circadian rhythms with respect to molecular
noise. *Proceedings of the National Academy of Sciences*, *99*(2), pp.673-678.
50. Mondragón-Palomino, O., Danino, T., Selimkhanov, J., Tsimring, L. and Hasty, J., 2011. Entrainment of a
population of synthetic genetic oscillators. *Science*, *333*(6047), pp.1315-1319.
51. Xeros, N., 1962. Deoxyriboside control and synchronization of mitosis. *Nature*, *194*(4829), pp.682-683.
52. Mitchison, J.M. and Creanor, J., 1971. Induction synchrony in the fission yeast Schizosaccharomyces
pombe. *Experimental cell research*, *67*(2), pp.368-374.
53. Bjursell, G. and Reichard, P., 1973. Effects of thymidine on deoxyribonucleoside triphosphate pools and
deoxyribonucleic acid synthesis in Chinese hamster ovary cells. *Journal of Biological Chemistry*, *248*(11),
pp.3904-3909.
54. Jordan, A. and Reichard, P., 1998. Ribonucleotide reductases. *Annual review of biochemistry*, *67*(1),
pp.71-98.
55. Johnson, W.M., Kido Soule, M.C. and Kujawinski, E.B., 2017. Extraction efficiency and quantification of
dissolved metabolites in targeted marine metabolomics. *Limnology and Oceanography: Methods*, *15*(4),
pp.417-428.
56. Ting, C.S., Hsieh, C., Sundararaman, S., Mannella, C. and Marko, M., 2007. Cryo-electron tomography
reveals the comparative three-dimensional architecture of Prochlorococcus, a globally important marine
cyanobacterium. *Journal of bacteriology*, *189*(12), pp.4485-4493.
57. Cermak, N., Becker, J.W., Knudsen, S.M., Chisholm, S.W., Manalis, S.R. and Polz, M.F., 2017. Direct single-
cell biomass estimates for marine bacteria via Archimedes' principle. *The ISME journal*, *11*(3), pp.825-828.
58. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. and Morishima, K., 2017. KEGG: new perspectives on
genomes, pathways, diseases and drugs. *Nucleic acids research*, *45*(D1), pp.D353-D361.
59. Cader, M.Z., de Almeida Rodrigues, R.P., West, J.A., Sewell, G.W., Muhammad, N., Reikine, S., Sirago, G.,
Unger, L.W., Iglesias-Romero, A.B., Ramshorn, K. and Haag, L.M., 2020. FAMIN is a multifunctional purine
enzyme enabling the purine nucleotide cycle. Cell, 180(2), pp.278-295.
60. Hogle, S. L. (2019). MARMICRODB database for taxonomic classification of (marine) metagenomes (1.0.0)
[Data set]. Zenodo. https://doi.org/10.5281/zenodo.3520509
61. Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K. and Madden, T.L., 2009.
BLAST+: architecture and applications. *BMC bioinformatics*, *10*, pp.1-9.
62. Sunagawa, S., et al., 2015. Structure and function of the global ocean microbiome. *Science*,
348(6237), p.1261359.
63. Biller, S.J., Berube, P.M., Dooley, K., Williams, M., Satinsky, B.M., Hackl, T., Hogle, S.L., Coe, A., Bergauer,
K., Bouman, H.A. and Browning, T.J., 2018. Marine microbial metagenomes sampled across space and
time. *Scientific data*, *5*(1), pp.1-7.
64. Boyd, J.A., Woodcroft, B.J. and Tyson, G.W., 2018. GraftM: a tool for scalable, phylogenetically informed
classification of genes within metagenomes. *Nucleic Acids Research*, *46*(10), pp.e59-e59.
65. "UniProt: the universal protein knowledgebase in 2023." *Nucleic Acids Research* 51, no. D1 (2023): D523-
D531.
66. Potter, S.C., Luciani, A., Eddy, S.R., Park, Y., Lopez, R. and Finn, R.D., 2018. HMMER web server: 2018
update. *Nucleic acids research*, *46*(W1), pp.W200-W204.
67. Salt, L.A., van Heuven, S.M., Claus, M.E., Jones, E.M. and De Baar, H.J.W., 2015. Rapid acidification of
mode and intermediate waters in the southwestern Atlantic Ocean. *Biogeosciences*, *12*(5), pp.1387-1401.
68. Rijkenberg, M.J., Middag, R., Laan, P., Gerringa, L.J., van Aken, H.M., Schoemann, V., de Jong, J.T. and
De Baar, H.J., 2014. The distribution of dissolved iron in the West Atlantic Ocean. *PloS one*, *9*(6),
p.e101323.

69. Wyatt, N.J., Milne, A., Woodward, E.M.S., Rees, A.P., Browning, T.J., Bouman, H.A., Worsfold, P.J. and
Lohan, M.C., 2014. Biogeochemical cycling of dissolved zinc along the GEOTRACES South Atlantic
transect GA10 at 40 S. *Global Biogeochemical Cycles*, *28*(1), pp.44-56.
70. Hogle, S.L., Hackl, T., Bundy, R.M., Park, J., Satinsky, B., Hiltunen, T., Biller, S., Berube, P.M. and Chisholm,
S.W., 2022. Siderophores as an iron source for picocyanobacteria in deep chlorophyll maximum layers of
the oligotrophic ocean. *The ISME Journal*, *16*(6), pp.1636-1646.