Title:

An increasing Arctic-boreal CO$_2$ sink offset by wildfires and source regions

Coauthors:

Anna-Maria Virkkala[1], Brendan M. Rogers[1], Jennifer D. Watts[1], Kyle A. Arndt[1], Stefano Potter[1], Isabel Wargowsky[1], Edward A. G. Schuur[2,3], Craig See[2], Marguerite Mauritz[4], Julia Boike[5,6], Syndonia M. Bret-Harte[7], Eleanor J. Burke[8], Arden Burrell[1], Namyi Chae[9], Abhishek Chatterjee[10], Frederic Chevallier[11], Torben R. Christensen[12,13], Roisin Commane[14], Han Dolman[15,16], Bo Elberling[17], Craig A. Emmerton[18], Eugenie S. Euskirchen[7], Liang Feng[19], Mathias Goeckede[20], Achim Grelle[21], Manuel Helbig[32,22,39], David Holl[23], Järvi Järveoja[24], Hideki Kobayashi[25z], Lars Kutzbach[23], Junjie Liu[10], Ingrid Liujkx[44], Efrén López-Blanco[12,46], Kyle Lunneberg[26], Ivan Mammarella[27], Maija E. Marushchak[28], Mikhail Mastepanov[12,13], Yojiro Matsuura[29], Trofim Maximov[30], Lutz Merbold[31], Gesa Meyer[32,33], Mats B. Nilsson[24], Yosuke Niwa[34], Walter Oechel[26], Sang-Jong Park[35], Frans-Jan W. Parmentier[36], Matthias Peichl[24], Wouter Peters[44,37], Roman Petrov[30], William Quinton[38], Christian Rödenbeck[20], Torsten Sachs[39, 45], Christopher Schulze[32,40], Oliver Sonnentag[32], Vincent St.Louis[18], Eeva-Stiina Tuittila[41], Masahito Ueyama[42], Andrej Varlagin[43], Donatella Zona[26], and Susan M. Natali[1]

Correspondence: avirkkala@woodwellclimate.org

Affiliations:

Woodwell Climate Research Center, Falmouth, USA
Center for Ecosystem Science and Society, Northern Arizona University Flagstaff, USA
Department of Biological Sciences, Northern Arizona University Flagstaff, USA
Biological Sciences, University of Texas at El Paso, El Paso, USA
Permafrost Research Section, Alfred Wegener Institute Helmholtz Center for Polar and Marine Research, Potsdam, Germany
Department of Geography, Humboldt University, Berlin, Germany
Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, USA
Met Office Hadley Centre, Exeter, UK
Institute of Life Science and Natural Resources, Korea University, Seoul, South Korea
NASA Jet Propulsion Laboratory, California Institute of Technology, USA
Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette, France
Department of Ecoscience, Arctic Research Center, Aarhus University, Roskilde, Denmark
Water, energy and environmental engineering research unit, University of Oulu, Oulu, Finland
Department of Earth and Environmental Sciences, Columbia University, NY, USA
Royal Netherlands Institute for Sea Research, Texel, Netherlands
Vrije Universiteit, Amsterdam, the Netherlands
Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark
Department of Biological Sciences, University of Alberta, Edmonton, Canada
NCEO, School of GeoSciences, University of Edinburgh, UK
Max Planck Institute for Biogeochemistry, Jena, Germany
Linnaeus University, Växjö, Sweden

22 Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Canada
23 Institute of Soil Science, Center for Earth System Research and Sustainability (CEN), Universität
Hamburg, Hamburg, Germany
24 Department of Forest Ecology and Management, Swedish University of Agricultural Sciences, Umeå,
Sweden
25 Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology
26 Department Biology, San Diego State University, San Diego, USA
27 Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of
Helsinki, Finland
28 Department of Environmental and Biological Sciences, University of Eastern Finland, Finland
29 Forestry and Forest Products Research Institute, Tsukuba, Ibaraki, Japan
30 Institute for Biological Problems of Cryolithozone of the Siberian Branch of the RAS - Division of
Federal Research Centre "The Yakut Scientific Centre of the Siberian Branch of the Russian Academy of
Sciences", Yakutsk, Russia
31 Integrative Agroecology, Agroecology and Environment, Agroscope, Zurich, Switzerland
32 Département de géographie, Université de Montréal, Montréal, Québec, Canada
33 Environment and Climate Change Canada, Climate Research Division, Victoria, Canada
34 Earth System Division, National Institute for Environmental Studies
/Department of Climate and Geochemistry Research, Meteorological Research Institute, Japan
35 Division of Atmospheric Sciences, Korea Polar Research Institute, Incheon, Republic of Korea
36 Centre for Biogeochemistry in the Anthropocene, Department of Geosciences, University of Oslo,
Oslo, Norway
37 Centre for Isotope Research, Energy and Sustainability Research Institute Groningen, Groningen
University, The Netherlands
38 Cold Regions Research Centre, Wilfrid Laurier University, Waterloo, Canada
39 GFZ German Research Centre for Geosciences, Germany
40 Department of Renewable Resources, University of Alberta, Edmonton, Canada
41 School of Forest Sciences, University of Eastern Finland
42 Graduate School of Agriculture, Osaka Metropolitan University
43 A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia
44 Dept. of Meteorology and Air Quality, Wageningen University, The Netherlands
45 Institute of Geoecology, Technische Universität Braunschweig, Braunschweig, Germany
46 Department of Environment and Minerals, Greenland Institute of Natural Resources, Nuuk, Greenland
81

82   **Abstract (151 words)**
83

The Arctic-Boreal Zone (ABZ) is rapidly warming, impacting its large soil carbon stocks. We use
a new compilation of terrestrial ecosystem CO_2 fluxes, geospatial datasets and random forest
models to show that although the ABZ was an increasing terrestrial CO_2 sink from 2001 to 2020
(mean ± standard deviation in net ecosystem exchange: -548 ± 140 Tg C yr^{-1}; trend: -14 Tg C
yr^{-1}, p<0.001), more than 30% of the region was a net CO_2 source. Tundra regions may have
already started to function on average as CO_2 sources, demonstrating a critical shift in carbon
dynamics. After factoring in fire emissions, the increasing ABZ sink was no longer statistically
significant (budget: -319 ± 140 Tg C yr^{-1}; trend: -9 Tg C yr^{-1}), with the permafrost region
becoming CO_2 neutral (budget: -24 ± 123 Tg C yr^{-1}; trend: -3 Tg C yr^{-1}), underscoring the
importance of fire in this region.
94

Main text (3171 words)

Estimating terrestrial net ecosystem CO_2 exchange (NEE) of the Arctic-Boreal Zone (ABZ) poses a significant challenge[1–4] due to their complex functions [4–6] and a limited network of field measurements [7,8]. As a result, models show a wide range of CO_2 budgets, from substantial net atmospheric sinks (-1,800 Tg C yr^{-1}) to moderate atmospheric sources (600 Tg C yr^{-1}) [1,4,5,9,10], a concerning discrepancy as northern permafrost soils hold nearly half of global soil organic carbon stocks [11]. The release of this soil carbon to the atmosphere as CO_2 could significantly exacerbate climate change [12]. Thus, there is an urgent need to improve CO_2 budget estimates across the ABZ.

The rapid climate change of the ABZ makes this discrepancy even more critical [13]. Increasing air and soil temperatures in both summer and non-summer seasons are causing changes in the CO_2 budget that remain poorly understood [9]. Furthermore, it is not known how the widespread but spatially heterogeneous increase in vegetation productivity and greening [14,15] impacts the annual CO_2 balance although links to enhanced CO_2 sinks during the spring-summer period have been found [16,17]. Some of the enhanced uptake might be offset by CO_2 losses associated with vegetation dieback ('browning'), and the escalating frequency and intensity of disturbances such as abrupt permafrost thaw (e.g., thermokarst), drought and fires, further complicating the understanding of ABZ carbon dynamics and climate feedbacks [18–20].

Current evidence on recent ABZ CO_2 budget trends and their main drivers is limited to few in-situ data-driven synthesis and modeling studies without a regional perspective on where and why CO_2 budgets are changing [1,5,9,10]. These studies have focused primarily on ecosystem CO_2 fluxes (i.e., not incorporating fire emissions), coarse annual or seasonal CO_2 fluxes (i.e., overlooking the intra-annual dynamics), and spatial patterns in CO_2 fluxes with data from only one to two decades. Most importantly, earlier studies have not extended into the 2020s, a period of time where warming has further accelerated and more fires have occurred [21]. Thus, we lack a comprehensive understanding of the regional and seasonal patterns in recent ecosystem CO_2 fluxes, including fire emissions, and their multidecadal trends, and the links to changing environmental conditions across the ABZ.

Here, we address this knowledge gap using the most comprehensive site-level ABZ CO_2 flux dataset to-date —including monthly terrestrial photosynthesis (gross primary production; GPP), ecosystem respiration (R_{eco}), and NEE data from 200 terrestrial eddy covariance and flux chamber sites (4,897 site-months). This dataset is at least four times larger than in earlier upscaling efforts and covers a longer time period with data extending to 2020. The same dataset was previously used to analyze in-situ CO_2 flux trends in permafrost versus non-permafrost regions, with the conclusion that the annual net uptake is increasing in the non-permafrost region but not in the permafrost region [22]. Here we extend that study from the site level to the full ABZ region by combining flux observations with meteorological, remote sensing, and soil data, together with random forest models to estimate CO_2 budgets across the ABZ. We do this upscaling over two periods, 2001-2020 (1-km resolution) and 1990-2016 (8-km resolution); results in the main text are based on the 1-km models unless stated otherwise. We

139 then assess regional and seasonal patterns and trends in ABZ ecosystem CO_2 fluxes and their
140 environmental drivers. We also integrate annual fire emissions from 2002 to 2020 [23] to provide
141 near-complete terrestrial CO_2 budget estimates (referred as NEE + fire).
142
143 **Results**
144
145 **CO_2 budgets across the ABZ**
146
Using machine learning models that had a high predictive performance (up to two times higher
cross-validated R^2 compared to earlier efforts [5,9]), we find that from 2001-2020 circumpolar
tundra was on average CO_2 neutral without accounting for fire emissions (in-situ NEE: -4 ± 44 g
C m^{-2} yr^{-1}; upscaled NEE: 7 ± 3 g C m^{-2} yr^{-1}; upscaled budget 45 ± 53 Tg C yr^{-1}; mean ±
standard deviation; Table 1). In contrast, the boreal was a strong sink (in-situ NEE: -42 ± 82 g C
m^{-2} yr^{-1}; upscaled NEE: -43 ± 7 g C m^{-2} yr^{-1}; upscaled budget -593 ± 101 Tg C yr^{-1}). Including fire
emissions (on average 237 Tg C yr^{-1} [23], i.e., 2% of R_{eco} and 43% of the ABZ net CO_2 uptake
budget) changed the budget to -383 ± 101 Tg C yr^{-1} in the boreal and to 64 ± 53 Tg C yr^{-1} in the
tundra. With fire emissions included, the permafrost region turned into CO_2 neutral (NEE: -249 ±
123 Tg C yr^{-1}, NEE + fire: -24 ± 123 Tg C yr^{-1}).
157
Although the entire ABZ domain was a terrestrial CO_2 sink across all years during 2001-2020
with an average NEE of -548 ± 140 Tg C yr^{-1}, our upscaling of NEE revealed a large areal
fraction of annual ecosystem CO_2 sources across the domain (34% of the total region, Fig. 1).
For the permafrost domain, the fraction of annual CO_2 sources was even higher (41% of the
region). This large fraction is also seen in our in-situ CO_2 flux database, with 29% of sites being
CO_2 sources (NEE between 0-142 g C m^{-2} yr^{-1}). These CO_2 source sites were mostly in Alaska
(44%), but also in northern Europe (25%), Canada (19%), and Siberia (13%). One key factor
driving CO_2 sources is the long and persistent non-summer season (September-May) emissions
in the tundra that, on average, exceed the short summer (June-August) net CO_2 uptake (Table
1). In the boreal, longer summers with strong uptake still dominate over non-summer emissions.
168
169 **Model performance and comparison**
170
We observed moderate correlation of our upscaled NEE results with an ensemble of
atmospheric inversions [24] across space (Pearson's correlation 0.5, p<0.001), but the correlation
between the temporal trends was weaker (Pearson's correlation 0.2, p<0.001) (Fig. 1).
However, the ensemble net uptake budgets from the inversions, as well as from a global
machine-learning based upscaling product (FLUXCOM-X-BASE [25,26]) were 1.5 to 3 times larger
than our upscaled budgets (Supplementary Section 5). Moreover, the global Coupled Model
Intercomparison Project Phase 6 (CMIP6) process model ensemble [27] had barely any annual
CO_2 sources across the ABZ, indicating that the process models may not accurately simulate
CO_2 source situations (Fig. 1), especially given the prevalence of site-level sources. The cross-
validated predictive performances of our random forest models for GPP, R_{eco}, and NEE showed
high correlations between observed and predicted fluxes (R^2 varied from 0.5 to 0.78 and root
mean square error from 19.4 to 37.3 g C m^{-2} $month^{-1}$; Supplementary Fig. 1-3), but upscaling

uncertainties remain. For example, areas with the most extensive strong sink or source
estimates rarely had in-situ data and were thus largely extrapolated (e.g., sources in central
Siberia, or sinks in southern Siberia, Supplementary Fig. 4). These areas also had the highest
uncertainties in our analysis (approximately twice as large uncertainties as in the more densely
measured areas; Supplementary Fig. 5).

Table 1. Mean gross primary productivity (GPP), ecosystem respiration (R_{eco}), and net
ecosystem exchange (NEE) fluxes and budgets over 2001-2020, and NEE + fire budgets from
2002-2020. Uncertainties are standard deviations across sites or pixels (for the mean fluxes) or
across bootstrapped budget estimates. Positive numbers for NEE indicate net CO_2 loss to the
atmosphere and negative numbers indicate net CO_2 uptake by the ecosystem. Mismatches in
the site-level versus upscaled CO_2 fluxes are likely related to sites being biased to certain
regions and years while upscaled summaries should provide more representative regional
estimates but are influenced by model performance. Mismatches in the NEE vs. GPP-R_{eco}
estimates are related to different numbers of sites and observations being available for the
different fluxes. Supplementary Table 1 shows the budgets for different vegetation types and
regions.

Class	In-situ average			Upscaled per-area average			Average regional budget			The proportion of summer net uptake budget of non-summer net emissions	Average regional budget with fire	Area (x 10^6 km^2)
Flux and unit	NEE g C m^{-2} yr^{-1}	GPP g C m^{-2} yr^{-1}	R_{eco} g C m^{-2} yr^{-1}	NEE g C m^{-2} yr^{-1}	GPP g C m^{-2} yr^{-1}	R_{eco} g C m^{-2} yr^{-1}	NEE Tg C yr^{-1}	GPP Tg C yr^{-1}	R_{eco} Tg C yr^{-1}	%	NEE + fire Tg C yr^{-1}	
Arctic-Boreal Zone	-32 (± 76)	618 (± 396)	588 (± 385)	-26 (± 5)	482 (± 20)	460 (± 15)	-548 (± 140)	9970 (± 144)	9525 (± 90)	1.4	-319	20.79
Tundra	-4 (± 44)	302 (± 125)	312 (± 133)	7 (± 3)	300 (± 14)	306 (± 12)	45 (± 53)	2049 (± 49)	2090 (± 33)	0.9	64	6.8
Boreal	-42 (± 82)	705 (± 402)	664 (± 398)	-43 (± 7)	572 (± 24)	537 (± 17)	-593 (± 101)	7920 (± 106)	7435 (± 74)	1.6	-383	13.9
Permafrost region	-21 (± 62)	458 (± 197)	445 (± 171)	-15 (± 5)	416 (± 20)	405 (± 16)	-249 (± 123)	6918 (± 109)	6719 (± 69)	1.2	-24	16.6

201
202

Figure 1. Maps showing the mean annual terrestrial NEE (a) and its trends (b) based on site-level data, our upscaling, atmospheric inversion ensemble, and CMIP6 process model ensemble. In-situ trends in b are based on sites that have >7 years of data. Supplementary Fig. 5c shows the significance of the trends. While the average upscaled NEE values can go up to 116 g C m^{-2} yr^{-1}, most of the values are below 60 g C m^{-2} yr^{-1}. While the NEE values of the inversion ensemble can go down to -1636 g C m^{-2} yr^{-1}, most of the values are higher than -200 g C m^{-2} yr^{-1}, similar to upscaling and CMIP6 model outputs.

Temporal trends in upscaled ABZ CO_2 budgets

The ABZ has been an increasing terrestrial CO_2 sink based on NEE alone from 2001 to 2020 (temporal trend: -14 Tg C yr^{-1}, p<0.001) (Fig. 2). However, the increasing sink strength was no longer statistically significant when fire emissions were added to NEE (average NEE + fire budget trend -9 Tg C yr^{-1} over 2002-2020). In the permafrost region, the NEE + fire trend was only -3.3 Tg C yr^{-1}. Nevertheless, based on our NEE upscaling, 23% of the region increased (p<0.05) net CO_2 uptake from 2001 to 2020 (Fig. 1), with increasing net sink pixels occurring across all key regions (Supplementary Fig. 31). Most of the increasing net sink activity was driven by an increase in GPP, especially in Siberia (Fig. 2). Some of the trends were also related to a declining R_{eco}, likely associated with disturbed ecosystems (e.g., forest fires, harvesting) with high R_{eco} during the first post-disturbance years now recovering [28]. However, evidence for the increasing overall net uptake trend from the in-situ data is limited due to the low number of long-term sites (>7 years of year-round measurements; 9 sites) out of which only one site showed a statistically significant trend (increasing uptake at a boreal forest site in Finland). Some of the relationships in our model are likely thus influenced by spatial differences across

229   the sites rather than temporal and truly causal patterns, creating some uncertainty in upscaled
230   trends [29]. However, the model reproduces temporal patterns at individual sites well (see
231   Supplementary Fig. 6), and our upscaled trends are similar to a recent in-situ time-series
232   analysis [22] and somewhat similar to those estimated from the inversion ensemble (Fig. 1),
233   providing confidence in our trend results.
234

235
Figure 2. Terrestrial CO_2 budgets for 1-km (blue; 2001-2020) and 8-km (grey; 1990-2016) NEE,
and 1-km NEE + fire emissions (red; 2002-2020) across the ABZ (a) and permafrost region (b).
An overlay analysis of NEE, GPP, Reco trend maps identifying how trends in GPP and Reco
relate to trends in NEE over 2001-2020 (includes significant and non-significant trends; c), and a
map showing also pixels burned during 2002-2020 (d). In a and b, trends are shown for the

2002-2020 (NEE + fire) and 2001-2020 (NEE) periods. Stars in the trend values depict the
significance of the trend (*= p<0.05, **=p<0.01, ***=p<0.001).
243

Figure 3. Average upscaled monthly NEE, GPP, and R_{eco} in boreal and tundra biomes during
the past two decades. Negative NEE values represent net uptake and positive net release. Error
bars are only shown for the 2011-2020 period but are similar for the 2001-2010 period. Note
that NEE was 1.4 g C m^{-2} $month^{-1}$ lower in September 2011-2020 compared to 2001-2020 in the
boreal biome, but this is not shown in the figure. For a similar figure made based on the in-situ
data, see Supplementary Fig. 7.

Parts of the ABZ also show increasing annual net CO_2 emissions over time (Fig. 1). Such trends
have been observed at six long-term sites (2 to 17 g C m^{-2} yr^{-1}, $p>0.05$), and in 2% of the
upscaled region ($p<0.05$) from 2001 to 2020. Most of the increasing net emission trends were
driven by an increase in R_{eco} instead of a decline in GPP (Fig. 2). Regions experiencing
increased net CO_2 emissions in upscaling were found especially in (i) northern Europe and
Canada (dominated by evergreen needleleaf forests with mild and moderately wet climates), (ii)
parts of central Alaska and northern Siberia (sparse boreal ecosystems and graminoid tundra
with permafrost and high soil carbon stocks), and (iii) Hudson Bay and Siberian lowlands
(wetlands with some permafrost and high soil organic carbon stocks) (Supplementary Fig. 31).
Some sites in Alaska have increasing net emissions of CO_2 due to permafrost thaw [18,30], but it is
unclear if similar changes are occurring in other regions with increasing net CO_2 emissions.
263
We calculated an overall 25% increase in seasonal amplitude of CO_2 fluxes from the upscaled
NEE time series from 2001 to 2020 across the ABZ, on par with earlier atmospheric and
modeling studies [31,32]. Both increasing summer uptake and non-summer season emissions—the
key dynamics driving increasing annual sinks and sources—were evident in the tundra and
boreal biomes (Fig. 3). However, over the 2001-2020 period, the increasing uptake (GPP)
during summer months dominated over increasing net emissions (R_{eco}) during non-summer
months across most of the domain. On average across both biomes, net uptake increased the
most during July (an average upscaled increase of -5 g C m^{-2} $month^{-1}$ in the boreal and -3 g C
m^{-2} $month^{-1}$ in the tundra in 2011-2020 compared to 2001-2010), and increasing net emissions
were occurring throughout the entire non-summer season, with no clear peaks (0.1-0.9 g C m^{-2}
$month^{-1}$). Although increases in early growing season (May-June) uptake were evident, late
growing season (September) trends were absent or minimal (Fig. 3).
276
277   **Drivers of ABZ CO_2 fluxes**
278
There are several environmental conditions driving CO_2 budgets across the ABZ. Our variable
importance analysis showed CO_2 fluxes, and thus the overall increasing sink strength, are
explained by dynamic variables of air or surface temperatures, solar radiation, the Normalized
Difference Vegetation Index (NDVI), and partially also by soil temperature, snow cover, and the
vapor pressure deficit (Supplementary Fig. 8-10). Other less important dynamic variables were
vegetation cover and atmospheric CO_2 concentration. Volumetric soil water content was not
important in our models, likely due to the large uncertainties and coarse spatial resolution in the
gridded product, although in-situ studies have shown drier soils to be linked to larger net CO_2
emissions and wetter soils to enhanced plant growth due to the lack of water limitation [33]. Static
variables (primarily vegetation type, soil carbon stock, soil pH) were also important in explaining
spatial differences.
290
The most important dynamic variables had a positive overall effect on net uptake, GPP, and
R_{eco} (Supplementary Fig. 8-10), however, these relationships are more nuanced in reality. In
fact, the recent permafrost in-situ trend analysis of CO_2 fluxes using the same database
suggests that the CO_2 flux response to warmer temperatures ranges from positive to negative,
depending on the availability of water and nutrients at the site [22]. Consequently, strong warming

or greening trends did not always translate into increasing net CO_2 sinks in our upscaling
(Supplementary Fig. 11). For example, while 49% of the region experienced greening (June-
August average NDVI; based on MODIS NDVI, $p<0.05$), only 12% of those greening pixels
showed an annual increasing net CO_2 uptake trend, and 29% an increasing June-August net
uptake.
301
302   **Continental and regional patterns in CO_2 budgets and their trends**
303
Our upscaling showed clear continental patterns in NEE budgets and trends (Fig. 4), with the
boreal biome primarily driving the budget and trend differences between the continents [1,34]. The
increasing net uptake trend was more pronounced in Eurasia (-11 Tg C yr^{-1}, $p<0.001$) compared
to North America (-3 Tg C yr^{-1}, $p<0.05$), which corresponds with the smaller area and weaker
warming, declining snow cover and greening trends in North America (Supplementary Fig. 12-
14). We found statistically significant declining summer soil moisture trends in Siberian boreal
(Supplementary Fig. 15), but this did not translate into stronger net emissions. When fire
emissions were added, continental differences were less pronounced due to the much larger
and more rapidly increasing CO_2 emissions from Siberian fires (on average 160 compared to 76
Tg C yr^{-1} in North America; Supplementary Fig. 16). Fire emissions even reversed some NEE
trends: the strong increasing sink in Siberia became a source when fire emissions were
included (trend: +0.7 Tg C yr^{-1}; $p > 0.05$). However, Siberian ecosystems have the largest
uncertainty for both the upscaled fluxes and inversion-based estimates due to lack of in situ
observations, making it challenging to accurately determine the magnitude of continental
differences (Fig. 4, Supplementary Fig. 17).
319
Alaska is an important contributor in the weaker North American CO_2 sink. Based on our
analysis, Alaska as a whole was consistently CO_2 neutral or a source over 2002-2020 (NEE +
fire emissions), both in the boreal (budget +5 Tg C yr^{-1}) and tundra (budget +7 Tg C yr^{-1}). Alaska
has a relatively high density of observations, making this result more certain compared to other
regions. Alaska is therefore different from the other ABZ regions where boreal regions still
remain on average CO_2 sinks. Potential reasons for the Alaskan CO_2 source include Alaska
having the most rapidly warming autumns and declining autumn snow cover, which also have
high inter-annual variability (Supplementary Fig. 14 and 18). Further, field measurements
suggest that many of the observed changes in Alaskan ecosystems can be attributed to
permafrost thaw [18,30]—a phenomenon that has accelerated significantly in response to Alaska's
pronounced warming trend since the 1950s [35]. However, we were unable to incorporate
permafrost thaw into our models as high-resolution geospatial data from 1990 to 2020 were not
available. The question of whether analogous trends will manifest in other regions across the
northern permafrost region remains an important research priority.
334

Figure 4. Terrestrial CO_2 budgets for NEE and NEE + fire in key regions and biomes across the boreal (a) and tundra (b). Terrestrial CO_2 budgets are shown for 1-km (blue; 2001-2020) and 8-km (grey; 1990-2016) NEE, and 1-km NEE + fire emissions (red; 2002-2020). Trends are shown for the 2002-2020 (NEE + fire) and 2001-2020 (NEE) periods. The inset in Alaskan boreal in (a) shows the time series with a narrower y axis compared to the main figure to better detect interannual variability. Stars in the trend values depict the significance of the trend (*= p<0.05, **=p<0.01, ***=p<0.001). Fire emissions alone are shown in Supplementary Fig. 16.

Discussion

Our results show that the ABZ was on average an increasing terrestrial CO_2 sink (GPP is increasing more than R_{eco} + fire), indicating that the region still creates an important negative feedback to global warming. However, our study also suggests some positive feedbacks to climate change that have been more regional and of shorter duration in recent decades. We show that the presence of annual sources was large as indicated by several site-level and regional studies [36–38], and even larger with fire emissions included [39]. There were also extreme years when fire emissions exceeded annual net CO_2 uptake (e.g., 2003 in Siberian boreal, 2012

in Canadian boreal, and several years in the permafrost region; Fig. 2). We also observed
increasing shoulder season net emission trends, particularly in Alaska [40] (Supplementary Fig.
34). Moreover, while summer net uptake increase still dominates over non-summer CO_2
emissions, net CO_2 uptake is increasing only in the early and peak growing season (May-August
in the boreal and June-August in the tundra) and not in the late growing season (September),
because GPP does not increase later in the season due to plant physiological limitations, and
drier and warmer conditions cause enhanced R_{eco} instead [41–46]. A better understanding of how
soil moisture and hydrology have been and will be changing, and the impact of these changes
on CO_2 fluxes is critical for more accurate ABZ CO_2 budgets.
362
In the tundra, our findings reveal a noteworthy shift in carbon dynamics. While the tundra region
has been a carbon sink for millennia [47], our results suggest that many tundra regions may now
have started to function as CO_2 sources. This transition from an ecosystem CO_2 sink to a CO_2
source may have begun prior to 1990 [48], yet the precise timing of this transformation remains
uncertain. The main drivers of this pattern may be related to warming-induced permafrost thaw,
the drying of soils, or vegetation shifts [49–51] but remain unresolved. Tundra regions are also
progressing towards conditions where average annual soil (0-7 cm) temperatures are above
freezing, resulting in more soil organic material being susceptible to decomposition
(Supplementary Fig. 12). Overall, the primary reason behind the annual CO_2 emissions from
tundra ecosystems is the limited duration of the high net CO_2 uptake period, and the substantial
non-summer season net emissions. However, we observed lower in-situ and upscaled October-
April season NEE fluxes and budgets compared to Natali et al. (2019) throughout the entire
period (Supplementary Fig. 19; [52]).
376
Our results demonstrate the need to further study Siberian CO_2 flux trends. Our upscaling
indicated that some of the strongest net sources and sinks, and strongest increasing sink trends
occur in the Siberian boreal. Increasing sink trends in Siberian tundra were also the strongest
across tundra regions. The Siberian sink trend might be explained by strong greening trends [53],
earlier growing season starts and increasing carbon uptake due to declining spring snow cover
(Supplementary Fig. 14), increases in tree growth and distribution [54,55], rapid recovery of
ecosystems after fire [56], and high cover of larch forests that can rapidly take up CO_2
(Supplementary Table 1)[8,57]. However, the large inversion model spread, sparse measurement
network, and our upscaling uncertainties indicate that it remains challenging to conclude the
magnitude of the Siberian CO_2 balance [2]. This is a significant problem given that Siberia stores
more than half of the permafrost region's C stocks and is now warming more rapidly than other
ABZ regions.
389
In summary, our study reveals distinct spatial and temporal patterns in CO_2 budgets across the
ABZ and underscores the importance of three decades' worth of data. Relatively robust spatial
patterns can be seen, such as the Alaskan CO_2 sources and southern Eurasian boreal sinks
while the temporal trends remain more uncertain. While CO_2 fluxes can be relatively well
modeled using machine learning and existing gridded datasets, gaps persist, such as the
incomplete characterization of fire, thermokarst and harvesting disturbances and their links to
ecosystem CO_2 fluxes, the lack of accurate predictors describing soil moisture [1], and the need

to quantify landscape heterogeneity and carbon dynamics at even higher spatial (meters) and
temporal resolutions (days). Sustaining long-term sites is crucial to accurately track trends in
ABZ CO_2 balance, while establishing new year-round sites in data-poor areas like Siberia and
the Canadian Arctic is vital to fill knowledge gaps and enhance our understanding of carbon
dynamics [58].

Online Methods

In-situ data overview

We used a recently compiled dataset of in-situ Arctic-boreal terrestrial ecosystem CO_2 flux
measurements (ABCflux, led by Virkkala et al. 2021 [8]) within the ABZ (Supplementary Methods
Section 1). The synthesized data were cumulative flux densities of gross NEE, GPP, and R_{eco}
aggregated at the monthly timescale (3,968 to 4,897 site-months depending on the flux). In
addition to eddy covariance data, we included fluxes estimated with the chamber method to
increase data coverage especially during the growing season. The dataset included metadata
out of which we used the site coordinate, biome, flux measurement method, and disturbance
history information in the analysis. For further details on the dataset, see Virkkala et al. (2021) [8]
and a description of additional data processing and screening in the Supplementary Methods
Section 2. Note that our study does not include lateral transport of carbon, or vertical lake and
river CO_2 emissions which were recently summarized to be 93, 66, and 164 Tg C yr^{-1},
respectively, in the northern permafrost region (i.e., greater than the NEE+fire budget calculated
in this study) [59].

This dataset is more comprehensive than the ones used in earlier upscaling studies as it
represents monthly fluxes from the entire year if available, while Virkkala et al. (2021) focused
on coarse seasonal or annual fluxes [5], Natali et al. (2019) on monthly winter fluxes [9], and Mu et
al. (2023) a more limited temporal period (2002-2017) [60]. Furthermore, we included more data
from recent years (805 monthly observations from 2015-2020 compared to 32 and 95 fluxes in
Virkkala et al. 2021 and Natali et al. 2019, respectively), and the sample size in our models was
4 to 25 times larger here compared to the earlier upscaling efforts.

Geospatial data

We used data from geospatial products as predictor variables to upscale fluxes. Our models
had the following predictors: month, incident solar radiation, vapor pressure deficit, atmospheric
CO_2 concentration, vegetation type, snow cover (the fraction covered by snow), soil temperature
(0-7 cm), soil moisture (0-7 cm), NDVI (MODIS- or AVHRR-based), land surface temperature
(or air temperature; MODIS- or ERA5 Land-based), compound topographic index (i.e.,
topographic wetness index), continuous vegetation fields describing percent non-tree vegetation
and non-vegetated fraction and percent tree cover (MODIS- or AVHRR-based), soil pH (0-5
cm), soil organic carbon stock in 0-2 meters, and permafrost probability. In our analysis, NDVI
was the primary predictor describing disturbances, with declines in NDVI being related to
disturbances [5]. Data were in daily, weekly, monthly, annual, and static format (i.e., no temporal

changes such as in the compound topographic index). If data were of higher temporal resolution
than monthly, they were aggregated to monthly time steps. Gaps in MODIS and AVHRR NDVI
time series were filled to produce a continuous time series. Data were re-projected to North Pole
Lambert Azimuthal Equal Area Projection at 1 and 8 km spatial resolution and extracted at the
flux sites. See Supplementary Section 3 for further descriptions and data sources.
446
We used the Global Fire Emissions Database (GFED) 500-m fire product [23] to calculate fire
emissions. The product is based on a global fire emissions model with a spatial resolution of
500 m using MODIS data. The model was developed using an updated field measurement
synthesis database of fuel load and consumption which included improvements, for example, in
boreal soil carbon combustion. The higher resolution of the 500-m model compared to earlier
coarser models improved the detection of small-scale fires and understanding of landscape
heterogeneity, and reduced the scale mismatch in comparing field measurements to model grid
cells. However, some small fires might still be undetected by this model, leading to potential
underestimations in carbon emissions in this product.
456
457   **Machine learning modeling**
458
We used random forest models to upscale GPP, R_{eco}, and NEE to the ABZ from 1990 to 2020,
the period with in-situ flux measurements. Two sets of predictive models were developed: (i)
models using primarily predictors with a spatial resolution ≤1 km from 2001 to 2020 (i.e., the
MODIS era) at 1-km spatial resolution (hereafter 1-km models;), and (ii) models using coarser-
resolution predictors from the AVHRR GIMMS era (1990-2016;) from 1990 to 2016 at 8-km
spatial resolution (hereafter 8-km models) (Supplementary Table 3). Each model included all
available monthly fluxes from the entire year, i.e. no separate models for individual months or
seasons were developed, as this approach resulted in the best predictive performance. All
models included 17 predictors, but the sample sizes were variable depending on the amount of
data available for each flux and time period; NEE models had the highest amount of model
training data compared to GPP and R_{eco} models. For the 1-km model, coarsest predictors were
at 9-km resolution but most important predictors were at 1 to 4-km resolution. For the 8-km
model, the coarsest predictor resolution was 9 km, and the most important variables had a
resolution of 1 to 9 km.
473
Model parameter tuning was performed based on leave-one-site out cross validation (CV) to
achieve minimum predictive error. The models were run using the "caret" package in R version
4.2 [61]. We assessed the predictive performance of the final models using the (1) R^2, (2) root
mean square error (RMSE), 3) mean absolute error (MAE), and 4) mean bias error (MBE)
between predicted and observed values using the CV data. Larger RMSE and MAE values
indicate larger errors, and positive MBE values indicate overestimation. The predictive
performance of our models was good or high, with R^2 ranging from 0.55 to 0.78 and RMSE from
19.4 to 37.3 g C m^{-2} month^{-1}, but there were uncertainties in model performance at both ends of
the flux gradient. Specifically, the model had a tendency to slightly overestimate fluxes, which
was particularly evident with the model struggling with strong sink observations (observations of
ca. -180 to -80 g C m^{-2} month^{-1} were predicted to be -80 to -20 g C m^{-2} month^{-1} in cross

validation; Supplementary Fig. 1). Other uncertainties were mostly due to 1) our model not
being able to identify landscape heterogeneity with nearby sites showing large differences in
CO_2 fluxes (e.g., a forest and wetland site), and 2) our model not capturing inter-annual
variability at individual sites, both of which are likely attributed to the coarse, uncertain, and
missing predictors characterizing such conditions (e.g., soil moisture, disturbances)
(Supplementary Fig. 1-3).
491
We evaluated the uncertainty of predictions by creating 20 bootstrapped model training datasets
(with replacement; same sample size as in the original model training data) and using those to
develop 20 individual models and predictions. Out of the 20 predictions, we calculated the
standard deviation to represent prediction uncertainty. The uncertainty ranges in NEE + fire
budgets only represent NEE uncertainties. We further assessed the area of extrapolation of the
models, and the influence of the flux measurement method and disturbance history information
on flux predictions by training models with different subsets. Further details of the uncertainty
analyses can be found in the Supplementary Methods Section 4.
500
501 **Model performance in burned ecosystems**
502
In addition to direct fire emissions from combustion (i.e., burning) derived with GFED, fires have
a profound impact on carbon budgets by modulating post-fire ecosystem CO_2 fluxes [28,67]. Our
current database has 21 sites that reported fire disturbance. Only four of those were longer-term
sites (operating for >3 years) with recent (<10 years since burn) fire history. All four of these
sites in young recovering ecosystems were measured year-round and originally had an
evergreen (black spruce) forest cover which underwent a shift to a more deciduous shrub and
tree-dominated cover after a stand-replacing fire. These include (i) CA-NS7 with 4 years of data
starting 4 years after the fire, (ii) CA-SF3 with 6 years of data starting 3 years after the fire, (iii)
US-Rpf with a 6-year time series starting 4 years after the fire, and (iv) US-CR-Fire with a 4-year
time series starting the next year after the fire.
513
Across all the burned sites, the in-situ flux data and remotely-sensed NDVI time series show a
clear pattern of July NDVI values, GPP and net carbon uptake steadily increasing after the fire
(Supplementary Fig. 24). This post-fire recovery signal is captured by our upscaling, as we see
our upscaled GPP and net uptake drop after a fire, and then return to higher levels after the fire
(Supplementary Fig. 24 and 25). However, while our random forest model fits the time series of
the longer-term sites with recent fire history relatively well, the predictions based on cross
validation (i.e., model training data excluding each site) are variable (Supplementary Fig. 26),
indicating that our model might struggle in extrapolating post-fire ecosystem CO_2 fluxes in other
areas. The model performance at sites experiencing recent fire or other disturbances was also
lower than at sites without disturbance or disturbance information, as the model had a lower R_2
and a tendency to underestimate NEE values (i.e., underestimate net CO_2 emissions or
overestimate net CO_2 uptake) (Supplementary Fig. 27).
526
527
528

Spatial upscaling of fluxes

We upscaled fluxes across the Arctic-boreal terrestrial area ≥49° N [62], which comprises 20.69 × 10^6 km^2 of land (excluding glaciers and ice sheets; Fig. 1) with lake and glacier areas removed. The models were applied at a monthly time step from 2001 to 2020 for the 1-km models and from 1990 to 2016 for the 8-km models.

We analyzed the upscaled flux maps as well as fire emission and environmental predictor rasters for temporal trends using the nonparametric Mann–Kendall test using the "zyp" package [63,64] with pre-whitening (Zhang method [65]) to remove autocorrelation. We report the significance of Kendall's correlation coefficient (the strength of the time-series) and the Theil–Sen slope to describe trends over time. Finally, we calculated zonal statistics of average annual, seasonal, and monthly fluxes and trends across key regions (Siberia defined as all land east from the Ural mountains, including a small portion of Mongolia; the rest of Eurasia, including Greenland are grouped within the European classes), biomes (tundra and boreal) [62], permafrost region [66], and vegetation types [8].

546
547

Comparison to process models and atmospheric inversions

We compared our estimates with the CMIP6 process models [27], atmospheric inversions used in the Global Carbon Project's Global Carbon Budget 2022 [68], and a global upscaling product FLUXCOM-X-BASE [26]. We included a subset of CMIP6 process models (13 in total) that had soil thermal processes at several depths to assure they had some information about the freeze-thaw patterns in the permafrost region. We included inversions with data from the whole 2001-2020 period (i.e., included five inversions and excluded four). Fire CO_2 emissions [23] were subtracted from the inversions. CMIP6 process model outputs were only available for the 2001-2014 period. The final model outputs used here represent terrestrial NEE (GPP-R$_{eco}$) in a similar way across the models except for inversions that also include vertical CO_2 fluxes from water bodies. There is some heterogeneity between individual inversions and CMIP6 models within the ensembles, but overall the ensemble results can be considered robust [69,70].

561
562

Acknowledgements

This work was supported by funding from the Gordon and Betty Moore Foundation (grant #8414) and funding catalyzed by the TED Audacious Project (Permafrost Pathways). We additionally acknowledge the funding from the NASA Arctic–Boreal Vulnerability Experiment and Carbon Cycle Science programs (NNX17AE13G), NSF PLR Arctic System Science Research Networking Activities (RNA, Grant#1931333), Minderoo Foundation, NSF (grants DEB LTREB 1354370, 2011257, DEB-0425328, DEB-0724514, and DEB-0830997), the US Geological Survey Climate R&D Program, NSF Arctic Observatory Network (grants 1936752, 1503912, 1107892), KAKENHI(19H05668),The Danish National Research Foundation (CENPERM

DNRF100), EU HORIZON GreenFeedBack, grant agreement No. 101056921, Danish Arctic
Climate support through Greenland Ecosystem Monitoring and ICOS grants, Natural Sciences
and Engineering Research Council, the Deutsche Forschungsgemeinschaft (DFG, German
Research Foundation) under Germany's Excellence Strategy – EXC 2037 'CLICCS - Climate,
Climatic Change, and Society' – Project Number: 390683824, NASA Grant/Cooperative
Agreement Number: NNX17AD69A, The Research Council of Norway (BioGov, project nr.
323945), ERC synergy project Q-Arctic (grant agreement no. 951288), the Copernicus
Atmosphere Monitoring Service, implemented by ECMWF on behalf of the European
Commission (Grant: CAMS2 55), the Environment Research and Technology Development
Fund of the Environmental Restoration and Conservation Agency of Japan
(JPMEERF21S20810), ArCSII(JPMXD142031886), Financial support from the Swedish
Research Council (VR) and consortium partners to ICOS Sweden (grants 2015−06020 and
2019−00205) and SITES (grant 2017−00635), VR grant 2019-04676 and 2018-03966, ArcticNet
and NSERC,NOAA Cooperative Agreement NA16SEC4810008, Research Council of Finland
(NPERM project nrs 341349, 330840, 349503 ICOS-FIRI), ICOS-FI via University of Helsinki
funding, the EU Horizon Europe (GreenFeedback nr. 101056921 and LiweFor nr. 101079192),
NRF-2021M1A5A1065425 (KOPRI-PN23011), NRF-2021M1A5A1065679, NRF-
2021R1I1A1A01053870, the Dutch Research Council (NWO) (project number VI.Vidi.213.143),
and The Natural Environment Research Council through the National Centre for Earth
Observation (NE/R000115/1). Part of the research was carried out at the Jet Propulsion
Laboratory, California Institute of Technology, under a contract with the National Aeronautics
and Space Administration (80NM0018D0004). Part of the inverse analyses were performed on
the supercomputer systems at the National Institute for Environmental Studies and
Meteorological Research Institutes (NEC SX-Aurora TSUBASA and FUJITSU PRIMERGY
CX2550M5) and at the HPC cluster Aether at the University of Bremen, financed by DFG within
the scope of the Excellence Initiative (Germany).

599
600 **Data availability**
601
602 In-situ data used here can be accessed from ORNL DAAC [71] and geospatial data from the links
603 and references provided in the Supplementary Tables 3 and 6. The 1-km and 8-km upscaled
604 rasters of NEE, GPP, and R_{eco} together with their uncertainties will be published via ORNL
605 DAAC.
606
607 **Code availability**
608 The main analysis codes can be found in the Supplement.
609

References

1.   Watts, J. D. *et al.* Carbon uptake in Eurasian boreal forests dominates the high-latitude net

ecosystem carbon budget. *Glob. Chang. Biol.* **29**, 1870–1889 (2023).

2.   Fan, L. *et al.* Siberian carbon sink reduced by forest disturbances. *Nat. Geosci.* **16**, 56–62

(2022).

3.   McGuire, A. D. *et al.* Dependence of the evolution of carbon dynamics in the northern

permafrost region on the trajectory of climate change. *Proc. Natl. Acad. Sci. U. S. A.* **115**,

3882–3887 (2018).

4.   McGuire, A. D. *et al.* Variability in the sensitivity among model simulations of permafrost

and carbon dynamics in the permafrost region between 1960 and 2009. *Global*

Biogeochem. Cycles **30**, 1015–1037 (2016).

5.   Virkkala, A.-M. *et al.* Statistical upscaling of ecosystem CO2 fluxes across the terrestrial

tundra and boreal domain: regional patterns and uncertainties. *Glob. Chang. Biol.* (2021)

doi:10.1111/gcb.15659.

6.   Pongracz, A., Wårlind, D., Miller, P. A. & Parmentier, F.-J. W. Model simulations of arctic

biogeochemistry and permafrost extent are highly sensitive to the implemented snow

scheme in LPJ-GUESS. *Biogeosciences* **18**, 5767–5787 (2021).

7.   Pallandt, M. *et al.* Representativeness assessment of the pan-Arctic eddy-covariance site

network, and optimized future enhancements. (2021) doi:10.5194/bg-2021-133.

8.   Virkkala, A.-M. *et al.* The ABCflux database: Arctic-Boreal CO_2 flux observations and

ancillary information aggregated to monthly time steps across terrestrial ecosystems.

(2021) doi:10.5194/essd-2021-233.

9.   Natali, S. M. *et al.* Large loss of CO2 in winter observed across the northern permafrost

region. *Nat. Clim. Chang.* **9**, 852–857 (2019).

10.   Hugelius, G. *et al.* Two decades of permafrost region CO2, CH4, and N2O budgets suggest

a small net greenhouse gas source to the atmosphere. *Authorea Preprints* (2023)

doi:10.22541/essoar.169444320.01914726/v1.

11. Hugelius, G. *et al.* Estimated stocks of circumpolar permafrost carbon with quantified

uncertainty ranges and identified data gaps. *Biogeosciences* **11**, (2014).

12. Schuur, E. A. G. *et al.* Permafrost and Climate Change: Carbon Cycle Feedbacks From the

Warming Arctic. *Annu. Rev. Environ. Resour.* **47**, 343–371 (2022).

13. Rantanen, M. *et al.* The Arctic has warmed nearly four times faster than the globe since

1979. *Communications Earth & Environment* **3**, 1–10 (2022).

14. Berner, L. T. *et al.* Summer warming explains widespread but not uniform greening in the

Arctic tundra biome. *Nat. Commun.* **11**, 4621 (2020).

15. Bjorkman, A. D. *et al.* Plant functional trait change across a warming tundra biome. *Nature*

562, 57–62 (2018).

16. Lund, M. *et al.* Variability in exchange of CO2 across 12 northern peatland and tundra sites.

Glob. Chang. Biol. **16**, 2436–2448 (2010).

17. Happonen, K., Virkkala, A.-M., Kemppinen, J., Niittynen, P. & Luoto, M. Relationships

between above-ground plant traits and carbon cycling in tundra plant communities. *J. Ecol.*

110, 700–716 (2022).

18. Euskirchen, E. S., Bret-Harte, M. S., Shaver, G. R., Edgar, C. W. & Romanovsky, V. E.

Long-Term Release of Carbon Dioxide from Arctic Tundra Ecosystems in Alaska.

Ecosystems **20**, 960–974 (2017).

19. Varner, R. K. *et al.* Permafrost thaw driven changes in hydrology and vegetation cover

increase trace gas emissions and climate forcing in Stordalen Mire from 1970 to 2014.

Philos. Trans. A Math. Phys. Eng. Sci. **380**, 20210022 (2022).

20. Arndt, K. A., Hashemi, J., Natali, S. M., Schiferl, L. D. & Virkkala, A.-M. Recent Advances

and Challenges in Monitoring and Modeling Non-Growing Season Carbon Dioxide Fluxes

from the Arctic Boreal Zone. *Current Climate Change Reports* **9**, 27–40 (2023).

21. Watts, J. Regional hotspots of change in northern high latitudes informed by observations

from space. *Authorea Preprints* (2024) doi:10.22541/au.170497370.03373595/v1.

22. See, Craig R, Virkkala, Anna-Maria ,Natali, Susan M, Rogers, Brendan M , Marguerite

Mauritz, Mika Aurela, Christina Biasi, Stef Bokhorst, Julia Boike, Syndonia M Bret5 Harte,

Gerardo Celis, Namyi Chae, Torben R Christensen, Roisin Commane, Sara June Connon,

Sigrid Dengel, Han Dolman, Colin W Edgar, Bo Elberling, Craig A Emmerton, Eugenie

Euskirchen, Mathias Goeckede, Achim Grelle, Liam Heffernan, Manuel Helbig, David Holl,

Elyn Humphreys, Hiroki Iwata, Järvi Järveoja, Hideki Kobayashi, John Kochendorfer, Pasi

Kolari, Ayumi Kotani, Lars Kutzbach, Min Jung Kwon, Emma R Lathrop, Efrén López-

Blanco, Ivan Mammarella, Maija E Marushchak, Mikhail Mastepanov, Yojiro Matsuura, Lutz

Merbold, Gesa Meyer, Christina Minions, Mats B Nilsson, Julia Nojeim, Steven F

Oberbauer, David Olefeldt, Sang-Jong Park, Frans-Jan W Parmentier, Matthias Peichl,

Darcy Peter, Roman Petrov, Rafael Poyatos, Anatoly S Prokushkin, William Quinton, Heidi

Rodenhizer, Torsten Sachs, Kathleen Savage, Christopher Schulze, Sofie Sjögersten,

Oliver Sonnentag, Vincent LSt. Louis, Margaret Torn, Eeva-Stiina Tuittila, Masahito

Ueyama, Andrej Varlagin, Carolina Voigt, Jennifer D. Watts, Donatella Zona, Viacheslav I.

Zyryanov, and Edward A. G. Schuur. Decadal increases in carbon uptake offset by

respiratory losses across northern permafrost ecosystems. *Nature Climate Change 2024 in*

review.

23. van Wees, D. *et al.* Global biomass burning fuel consumption and emissions at 500 m

spatial resolution based on the Global Fire Emissions Database (GFED). *Geosci. Model*

Dev. **15**, 8411–8437 (2022).

24. Friedlingstein, P. *et al.* Global carbon budget 2021. *Earth Syst. Sci. Data* **14**, 1917–2005

(2022).

25. Jung, M. *et al.* Scaling carbon fluxes from eddy covariance sites to globe: synthesis and

evaluation of the FLUXCOM approach. *Biogeosciences* **17**, 1343–1365 (2020).

26. Nelson, J. A. *et al.* X-BASE: the first terrestrial carbon and water flux products from an

extended data-driven scaling framework, FLUXCOM-X. *EGUsphere* (2024)

doi:10.5194/egusphere-2024-165.

27. Eyring, V. *et al.* Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6)

experimental design and organization. *Geosci. Model Dev.* **9**, 1937–1958 (2016).

28. Ueyama, M. *et al.* Carbon dioxide balance in early-successional forests after forest fires in

interior Alaska. *Agric. For. Meteorol.* **275**, 196–207 (2019).

29. Damgaard, C. A Critique of the Space-for-Time Substitution Practice in Community

Ecology. *Trends Ecol. Evol.* **34**, 416–421 (2019).

30. Celis, G. *et al.* Tundra is a consistent source of CO_2 at a site with progressive permafrost

thaw during 6 years of chamber and eddy covariance measurements. *Journal of*

Geophysical Research: Biogeosciences vol. 122 1471–1485 Preprint at

https://doi.org/10.1002/2016jg003671 (2017).

31. Forkel, M. *et al.* Enhanced seasonal CO_2 exchange caused by amplified plant productivity

in northern ecosystems. *Science* **351**, 696–699 (2016).

32. Graven, H. D. *et al.* Enhanced seasonal exchange of CO_2 by northern ecosystems since

1960. *Science* **341**, 1085–1089 (2013).

33. McGuire, A. D. *et al.* An assessment of the carbon balance of arctic tundra: comparisons

among observations, process models, and atmospheric inversions. *Biogeosci. Discuss.* **9**,

4543 (2012).

34. Lin, X. *et al.* Siberian and temperate ecosystems shape Northern Hemisphere atmospheric

CO_2 seasonal amplification. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 21079–21087 (2020).

35. Jorgenson, M. T., Racine, C. H., Walters, J. C. & Osterkamp, T. E. Permafrost Degradation

and Ecological Changes Associated with a WarmingClimate in Central Alaska. *Clim.*

Change **48**, 551–579 (2001).

36. Euskirchen, E. S., Edgar, C. W., Turetsky, M. R., Waldrop, M. P. & Harden, J. W.

Differential response of carbon fluxes to climate in three peatland ecosystems that vary in

the presence and stability of permafrost. *J. Geophys. Res. Biogeosci.* **119**, 1576–1595

(2014).

37. Harris, L. I. *et al.* Permafrost thaw causes large carbon loss in boreal peatlands while

changes to peat quality are limited. *Glob. Chang. Biol.* **29**, 5720–5735 (2023).

38. Euskirchen, E. S. *et al.* Persistent net release of carbon dioxide and methane from an

Alaskan lowland boreal peatland complex. *Glob. Chang. Biol.* **30**, e17139 (2024).

39. Hayes, D. J. *et al.* Is the northern high-latitude land-based CO2sink weakening? *Global*

Biogeochemical Cycles vol. 25 Preprint at https://doi.org/10.1029/2010gb003813 (2011).

40. Commane, R. *et al.* Carbon dioxide sources from Alaska driven by increasing early winter

respiration from Arctic tundra. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 5361–5366 (2017).

41. Helbig, M. *et al.* Warming response of peatland CO2 sink is sensitive to seasonality in

warming trends. *Nat. Clim. Chang.* **12**, 743–749 (2022).

42. Zona, D. *et al.* Pan-Arctic soil moisture control on tundra carbon sequestration and plant

productivity. *Glob. Chang. Biol.* **29**, 1267–1281 (2023).

43. Zona, D. *et al.* Earlier snowmelt may lead to late season declines in plant productivity and

carbon sequestration in Arctic tundra ecosystems. *Sci. Rep.* **12**, 3986 (2022).

44. Byrne, B. *et al.* Multi-year observations reveal a larger than expected autumn respiration

signal across northeast Eurasia. *Biogeosciences* **19**, 4779–4799 (2022).

45. Pulliainen, J. *et al.* Early snowmelt significantly enhances boreal springtime carbon uptake.

Proc. Natl. Acad. Sci. U. S. A. **114**, 11081–11086 (2017).

46. Parker, T. C., Unger, S. L., Moody, M. L., Tang, J. & Fetcher, N. Intraspecific variation in

phenology offers resilience to climate change for Eriophorum vaginatum. *Arct. Sci.* 1–17

(2021).

47. Brovkin, V. *et al.* Comparative carbon cycle dynamics of the present and last interglacial.

Quat. Sci. Rev. **137**, 15–32 (2016).

48. Oechel, W. C. *et al.* Recent change of Arctic tundra ecosystems from a net carbon dioxide

sink to a source. *Nature* **361**, 520–523 (1993).

49. Lawrence, D. M., Koven, C. D., Swenson, S. C., Riley, W. J. & Slater, A. G. Permafrost

thaw and resulting soil moisture changes regulate projected high-latitude CO_2 and CH_4

emissions. *Environ. Res. Lett.* **10**, 094011 (2015).

50. Parker, T. C., Subke, J.-A. & Wookey, P. A. Rapid carbon turnover beneath shrub and tree

vegetation is associated with low soil carbon stocks at a subarctic treeline. *Glob. Chang.*

Biol. **21**, 2070–2081 (2015).

51. Voigt, C. *et al.* Warming of subarctic tundra increases emissions of all three important

greenhouse gases - carbon dioxide, methane, and nitrous oxide. *Glob. Chang. Biol.* **23**,

3121–3138 (2017).

52. Schiferl, L. D. *et al.* Using atmospheric observations to quantify annual biogenic carbon

dioxide fluxes on the Alaska North Slope. *Biogeosciences* **19**, 5953–5972 (2022).

53. Piao, S. *et al.* Characteristics, drivers and feedbacks of global greening. *Nature Reviews*

Earth & Environment **1**, 14–27 (2019).

54. Frost, G. V. & Epstein, H. E. Tall shrub and tree expansion in Siberian tundra ecotones

since the 1960s. *Glob. Chang. Biol.* **20**, 1264–1277 (2014).

55. Kharuk, V. I., Ranson, K. J., Im, S. T. & Petrov, I. A. Climate-induced larch growth response

within the central Siberian permafrost zone. *Environ. Res. Lett.* **10**, 125009 (2015).

56. Schulze, E.-D. *et al.* Factors promoting larch dominance in central Siberia: fire versus

growth performance and implications for carbon dynamics at the boundary of evergreen

and deciduous conifers. *Biogeosciences* **9**, 1405–1421 (2012).

57. Hiyama, T. *et al.* Lessons learned from more than a decade of greenhouse gas flux

measurements at boreal forests in eastern Siberia and interior Alaska. *Polar Sci.* **27**,

100607 (2021).

58. Pallandt, M. *et al.* High-latitude eddy covariance temporal network design and optimization.

Authorea Preprints (2023) doi:10.22541/essoar.169755225.54015522/v1.

59. Ramage, J. L. *et al.* The net GHG balance and budget of the permafrost region (2000-

2020) from ecosystem flux upscaling. *Authorea Preprints* (2023)

doi:10.22541/essoar.169462008.85493456/v1.

60. Mu, C. *et al.* Ecosystem CO_2 exchange and its economic implications in northern

permafrost regions in the 21st century. *Global Biogeochem. Cycles* **37**, (2023).

61. Kuhn, M. Building Predictive Models in R Using the caret Package. *J. Stat. Softw.* **28**, 1–26

(2008).

62. Dinerstein, E. *et al.* An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm.

Bioscience vol. 67 534–545 (2017).

63. Sen, P. K. Estimates of the Regression Coefficient Based on Kendall's Tau. *J. Am. Stat.*

Assoc. **63**, 1379–1389 (1968).

64. Zhang + Yue-Pilon Trends Package [R package zyp version 0.11-1]. (2023).

65. Zhang, X., Vincent, L. A., Hogg, W. D. & Niitsoo, A. Temperature and precipitation trends in

Canada during the 20th century. *Atmosphere-Ocean* **38**, 395–429 (2000).

66. Heginbottom, J., Brown, J., Ferrians, O. & Melnikov, E. S. Circum-arctic map of permafrost

and ground-ice conditions, version 2. NSIDC https://doi.org/10.7265/SKBG-KF16 (2002).

67. Amiro, B. D., Ian MacPherson, J., Desjardins, R. L., Chen, J. M. & Liu, J. Post-fire carbon

dioxide fluxes in the western Canadian boreal forest: evidence from towers, aircraft and

remote sensing. *Agric. For. Meteorol.* **115**, 91–107 (2003).

68. Friedlingstein, P. *et al.* Global carbon budget 2022. *Earth Syst. Sci. Data* **14**, 4811–4900

(2022).

69. Liu, Z. *et al.* Respiratory loss during late-growing season determines the net carbon dioxide

sink in northern permafrost regions. *Nat. Commun.* **13**, 5626 (2022).

70. Treat, C. C. *et al.* Permafrost carbon: Progress on understanding stocks and fluxes across

northern terrestrial ecosystems. *J. Geophys. Res. Biogeosci.* **129**, (2024).

71. Virkkala, A.-M. *et al.* The ABCflux Database: Arctic-boreal CO_2 flux and site environmental

data, 1989-2020. *ORNL DAAC* Preprint at https://doi.org/10.3334/ORNLDAAC/1934

(2021).