Title:

An increasing Arctic-Boreal CO$_2$ sink despite strong regional sources

Coauthors:

Anna-Maria Virkkala[1], Brendan M. Rogers[1], Jennifer D. Watts[1], Kyle A. Arndt[1], Stefano Potter[1], Isabel Wargowsky[1], Edward A. G. Schuur[2,3], Craig See[2], Marguerite Mauritz[4], Julia Boike[5,6], Syndonia M. Bret-Harte[7], Eleanor J. Burke[8], Arden Burrell[1], Namyi Chae[9], Abhishek Chatterjee[10], Frederic Chevallier[11], Torben R. Christensen[12,13], Roisin Commane[14], Han Dolman[15,16], Bo Elberling[17], Craig A. Emmerton[18], Eugenie S. Euskirchen[7], Liang Feng[19], Mathias Goeckede[20], Achim Grelle[21], Manuel Helbig[32,22], David Holl[23], Järvi Järveoja[24], Hideki Kobayashi[25], Lars Kutzbach[23], Junjie Liu[10], Ingrid Liujkx[44], Efrén López-Blanco[12,25], Kyle Lunneberg[26], Ivan Mammarella[27], Maija E. Marushchak[28], Mikhail Mastepanov[12,13], Yojiro Matsuura[29], Trofim Maximov[30], Lutz Merbold[31], Gesa Meyer[32,33], Mats B. Nilsson[24], Yosuke Niwa[34], Walter Oechel[26], Sang-Jong Park[35], Frans-Jan W. Parmentier[36], Matthias Peichl[24], Wouter Peters[44,37], Roman Petrov[30], William Quinton[38], Christian Rödenbeck[20], Torsten Sachs[39], Christopher Schulze[32,40], Oliver Sonnentag[32], Vincent St.Louis[18], Eeva-Stiina Tuittila[41], Masahito Ueyama[42], Andrej Varlagin[43], Donatella Zona[26], and Susan M. Natali[1]

Correspondence: avirkkala@woodwellclimate.org

Affiliations:

Woodwell Climate Research Center, Falmouth, USA
Center for Ecosystem Science and Society, Northern Arizona University Flagstaff, USA
Department of Biological Sciences, Northern Arizona University Flagstaff, USA
Biological Sciences, University of Texas at El Paso, El Paso, USA
Permafrost Research Section, Alfred Wegener Institute Helmholtz Center for Polar and Marine Research, Potsdam, Germany
Department of Geography, Humboldt University, Berlin, Germany
Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, USA
Met Office Hadley Centre, Exeter, UK
Institutes of Life Sciences and Natural Resources, Korea University, Seoul, South Korea
NASA Jet Propulsion Laboratory, California Institute of Technology, USA
Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette, France
Department of Ecoscience, Arctic Research Center, Aarhus University, Roskilde, Denmark
Oulanka research station, University of Oulu, Oulu, Finland
Department of Earth and Environmental Sciences, Columbia University, NY, USA
Royal Netherlands Institute for Sea Research, Texel, Netherlands
Vrije Universiteit, Amsterdam, the Netherlands
Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark
Department of Biological Sciences, University of Alberta, Edmonton, Canada
NCEO, School of GeoSciences, University of Edinburgh, UK
Max Planck Institute for Biogeochemistry, Jena, Germany
Linnaeus University, Växjö, Sweden

22 Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Canada
23 Institute of Soil Science, Center for Earth System Research and Sustainability (CEN), Universität
Hamburg, Hamburg, Germany
24 Department of Forest Ecology and Management, Swedish University of Agricultural Sciences, Umeå,
Sweden
25 Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology
26 Department Biology, San Diego State University, San Diego, USA
27 Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of
Helsinki, Finland
28 Department of Environmental and Biological Sciences, University of Eastern Finland, Finland
29 Forestry and Forest Products Research Institute, Tsukuba, Ibaraki, Japan
30 Institute for Biological Problems of Cryolithozone of the Siberian Branch of the RAS - Division of
Federal Research Centre "The Yakut Scientific Centre of the Siberian Branch of the Russian Academy of
Sciences", Yakutsk, Russia
31 Integrative Agroecology, Agroecology and Environment, Agroscope, Zurich, Switzerland
32 Département de géographie, Université de Montréal, Montréal, Québec, Canada
33 Environment and Climate Change Canada, Climate Research Division, Victoria, Canada
34 Earth System Division, National Institute for Environmental Studies
/Department of Climate and Geochemistry Research, Meteorological Research Institute, Japan
35 Division of Atmospheric Sciences, Korea Polar Research Institute, Incheon, Republic of Korea
36 Centre for Biogeochemistry in the Anthropocene, Department of Geosciences, University of Oslo,
Oslo, Norway
37 Centre for Isotope Research, Energy and Sustainability Research Institute Groningen, Groningen
University, The Netherlands
38 Cold Regions Research Centre, Wilfrid Laurier University, Waterloo, Canada
39 GFZ German Research Centre for Geosciences, Germany
40 Department of Renewable Resources, University of Alberta, Edmonton, Canada
41 School of Forest Sciences, University of Eastern Finland
42 Graduate School of Agriculture, Osaka Metropolitan University
43 A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia
44 Dept. of Meteorology and Air Quality, Wageningen University, The Netherlands

79
80
81

Abstract (151 words)

The Arctic-Boreal Zone (ABZ) is rapidly warming, impacting its large soil carbon stocks. We use
a new compilation of terrestrial ecosystem CO_2 fluxes, geospatial datasets and random forest
models to show that although the ABZ was an increasing terrestrial CO_2 sink from 2001 to 2020
(mean ± standard deviation in net ecosystem exchange: -548 ± 140 Tg C yr^{-1}; trend: -14 Tg C
yr^{-1}, p<0.001), more than 30% of the region was a net CO_2 source. Tundra regions may have
already started to function on average as CO_2 sources, demonstrating a critical shift in carbon
dynamics. After factoring in fire emissions, the increasing ABZ sink was no longer statistically
significant (budget: -319 ± 140 Tg C yr^{-1}; trend: -9 Tg C yr^{-1}), with the permafrost region
becoming CO_2 neutral (budget: -24 ± 123 Tg C yr^{-1}; trend: -3 Tg C yr^{-1}), underscoring the
importance of fire in this region.

Main text (3159 words)

Estimating terrestrial net ecosystem CO_2 exchange (NEE) of the Arctic-Boreal Zone (ABZ) poses a significant challenge[1–4] due to their complex functions [4–6] and a limited network of field measurements [7,8]. As a result, models show a wide range of CO_2 budgets, from substantial net atmospheric sinks (-1,800 Tg C yr^{-1}) to moderate atmospheric sources (600 Tg C yr^{-1}) [1,4,5,9,10], a concerning discrepancy as northern permafrost soils hold nearly half of global soil organic carbon stocks [11]. The release of this soil carbon to the atmosphere as CO_2 could significantly exacerbate climate change [12]. Thus, there is an urgent need to improve CO_2 budget estimates across the ABZ.

The rapid climate change of the ABZ makes this discrepancy even more critical [13]. Increasing air and soil temperatures in both summer and non-summer seasons are causing changes in the CO_2 budget that remain poorly understood [9]. Furthermore, it is not known how the widespread but spatially heterogeneous increase in vegetation productivity and greening [14,15] impacts the annual CO_2 balance although links to enhanced CO_2 sinks during the spring-summer period have been found [16,17]. Some of the enhanced uptake might be offset by CO_2 losses associated with vegetation dieback ('browning'), and the escalating frequency and intensity of disturbances such as abrupt permafrost thaw (e.g., thermokarst), drought and fires, further complicating the understanding of ABZ carbon dynamics and climate feedbacks [18–20].

Current evidence on recent ABZ CO_2 budget trends and their main drivers is limited to few in-situ data-driven synthesis and modeling studies without a regional perspective on where and why CO_2 budgets are changing [1,5,9,10]. These studies have focused primarily on ecosystem CO_2 fluxes (i.e., not incorporating fire emissions), coarse annual or seasonal CO_2 fluxes (i.e., overlooking the intra-annual dynamics), and spatial patterns in CO_2 fluxes with data from only one to two decades. Most importantly, earlier studies have not extended into the 2020s, a period of time where warming has further accelerated and more fires have occurred [21]. Thus, we lack a comprehensive understanding of the regional and seasonal patterns in recent ecosystem CO_2 fluxes, including fire emissions, and their multidecadal trends, and the links to changing environmental conditions across the ABZ.

Here, we address this knowledge gap using the most comprehensive site-level ABZ CO_2 flux dataset to-date —including monthly terrestrial photosynthesis (gross primary production; GPP), ecosystem respiration (R_{eco}), and NEE data from 200 terrestrial eddy covariance and flux chamber sites (4,897 site-months). This dataset is at least four times larger than in earlier upscaling efforts and covers a longer time period with data extending to 2020. The same dataset was previously used to analyze in-situ CO_2 flux trends in permafrost versus non-permafrost regions, with the conclusion that the annual net uptake is increasing in the non-permafrost region but not in the permafrost region [22]. Here we extend that study from the site level to the full ABZ region by combining flux observations with meteorological, remote sensing, and soil data, together with random forest models to estimate CO_2 budgets across the ABZ. We do this upscaling over two periods, 2001-2020 (1-km resolution) and 1990-2016 (8-km resolution); results in the main text are based on the 1-km models unless stated otherwise. We

139 then assess regional and seasonal patterns and trends in ABZ ecosystem CO_2 fluxes and their
140 environmental drivers. We also integrate annual fire emissions from 2002 to 2020 [23] to provide
141 near-complete terrestrial CO_2 budget estimates (referred as NEE + fire).
142
143 **Results**
144
145 **CO_2 budgets across the ABZ**
146
Using machine learning models that had a high predictive performance (up to two times higher
cross-validated R^2 compared to earlier efforts [5,9]), we find that from 2001-2020 circumpolar
tundra was on average CO_2 neutral without accounting for fire emissions (in-situ NEE: -4 ± 44 g
C m^{-2} yr^{-1}; upscaled NEE: 7 ± 3 g C m^{-2} yr^{-1}; upscaled budget 45 ± 53 Tg C yr^{-1}; mean ±
standard deviation; Table 1). In contrast, the boreal was a strong sink (in-situ NEE: -41 ± 82 g C
m^{-2} yr^{-1}; upscaled NEE: -43 ± 7 g C m^{-2} yr^{-1}; upscaled budget -593 ± 101 Tg C yr^{-1}). Including
fire emissions (on average 237 Tg C yr^{-1} [23], i.e., 2% of R_{eco} and 43% of the ABZ net CO_2 uptake
budget) changed the budget to -383 ± 101 Tg C yr^{-1} in the boreal and to 64 ± 53 Tg C yr^{-1} in the
tundra. With fire emissions included, the permafrost region turned into CO_2 neutral (NEE: -249 ±
123 Tg C yr^{-1}, NEE + fire: -24 ± 123 Tg C yr^{-1}).
157
Although the entire ABZ domain was a terrestrial CO_2 sink across all years during 2001-2020
with an average NEE of -548 ± 140 Tg C yr^{-1}, our upscaling of NEE revealed a large areal
fraction of annual ecosystem CO_2 sources across the domain (34% of the total region, Fig. 1).
For the permafrost domain, the fraction of annual CO_2 sources was even higher (41% of the
region). This large fraction is also seen in our in-situ CO_2 flux database, with 30% of sites being
CO_2 sources (NEE between 0-142 g C m^{-2} yr^{-1}). These CO_2 source sites were mostly in Alaska
(44%), but also in northern Europe (25%), Canada (19%), and Siberia (13%). One key factor
driving CO_2 sources is the long and persistent non-summer season (September-May) emissions
in the tundra that, on average, exceed the short summer (June-August) net CO_2 uptake (Table
1). In the boreal, longer summers with strong uptake still dominate over non-summer emissions.
168
169 **Model performance and comparison**
170
We observed moderate correlation of our upscaled NEE results with an ensemble of
atmospheric inversions [24] across space (Pearson's correlation 0.5, p<0.001), but the correlation
between the temporal trends was weaker (Pearson's correlation 0.2, p<0.001) (Fig. 1).
However, the ensemble net uptake budgets from the inversions, as well as from a global
machine-learning based upscaling product (FLUXCOM [25]) were 1.5 to 3 times higher than our
upscaled budgets (Supplementary Section 5). Moreover, the global Coupled Model
Intercomparison Project Phase 6 (CMIP6) process model ensemble [26] had barely any annual
CO_2 sources across the ABZ, indicating that the process models may not accurately simulate
CO_2 source situations (Fig. 1), especially given the prevalence of site-level sources. The cross-
validated predictive performances of our random forest models for GPP, R_{eco}, and NEE showed
high correlations between observed and predicted fluxes (R^2 varied from 0.5 to 0.78 and root
mean square error from 19.4 to 37.3 g C m^{-2} $month^{-1}$; Supplementary Fig. 1-3), but upscaling

 uncertainties remain. For example, areas with the most extensive strong sink or source
184 estimates rarely had in-situ data and were thus largely extrapolated (e.g., sources in central
185 Siberia, or sinks in southern Siberia, Supplementary Fig. 4). These areas also had the highest
186 uncertainties in our analysis (approximately twice as large uncertainties as in the more densely
187 measured areas; Supplementary Fig. 5).
188

Table 1. Mean gross primary productivity (GPP), ecosystem respiration (R_{eco}), and net
ecosystem exchange (NEE) fluxes and budgets over 2001-2020, and NEE + fire budgets from
2002-2020. Uncertainties are standard deviations across sites or pixels (for the mean fluxes) or
across bootstrapped budget estimates. Positive numbers for NEE indicate net CO_2 loss to the
atmosphere and negative numbers indicate net CO_2 uptake by the ecosystem. Mismatches in
the site-level versus upscaled CO_2 fluxes are likely related to sites being biased to certain
regions and years while upscaled summaries should provide more representative regional
estimates but are influenced by model performance. Mismatches in the NEE vs. GPP-R_{eco}
estimates are related to different numbers of sites and observations being available for the
different fluxes. Supplementary Table 1 shows the budgets for different vegetation types and
regions.
200

Class	In-situ average			Upscaled per-area average			Average regional budget			The proportion of summer net uptake budget of non-summer net emissions	Average regional budget with fire	Area (x 10^6 km^2)
Flux and unit	NEE g C m^{-2} yr^{-1}	GPP g C m^{-2} yr^{-1}	R_{eco} g C m^{-2} yr^{-1}	NEE g C m^{-2} yr^{-1}	GPP g C m^{-2} yr^{-1}	R_{eco} g C m^{-2} yr^{-1}	NEE Tg C yr^{-1}	GPP Tg C yr^{-1}	R_{eco} Tg C yr^{-1}	%	NEE + fire Tg C yr^{-1}	
Arctic-Boreal Zone	-32 (± 76)	617 (± 396)	587 (± 385)	-26 (± 5)	482 (± 20)	460 (± 15)	-548 (± 140)	9970 (± 144)	9525 (± 90)	1.4	-319	20.79
Tundra	-4 (± 44)	302 (± 124)	311 (± 133)	7 (± 3)	300 (± 14)	306 (± 12)	45 (± 53)	2049 (± 49)	2090 (± 33)	0.9	64	6.8
Boreal	-41 (± 82)	705 (± 402)	664 (± 398)	-43 (± 7)	572 (± 24)	537 (± 17)	-593 (± 101)	7920 (± 106)	7435 (± 74)	1.6	-383	13.9
Permafrost region	-21 (± 62)	459 (± 197)	447 (± 172)	-15 (± 5)	416 (± 20)	405 (± 16)	-249 (± 123)	6918 (± 109)	6719 (± 69)	1.2	-24	16.6

201
202

Figure 1. Maps showing the mean annual terrestrial NEE (a) and its trends (b) based on site-level data, our upscaling, atmospheric inversion ensemble, and CMIP6 process model ensemble. In-situ trends in b are based on sites that have >7 years of data. Supplementary Fig. 5c shows the significance of the trends. While the average upscaled NEE values can go up to 116 g C m^{-2} yr^{-1}, most of the values are below 60 g C m^{-2} yr^{-1}. While the NEE values of the inversions can go down to -1636 g C m^{-2} yr^{-1}, most of the values are higher than -200 g C m^{-2} yr^{-1}, similar to upscaling and CMIP6 model outputs.

Temporal trends in upscaled ABZ CO_2 budgets

The ABZ has been an increasing terrestrial CO_2 sink based on NEE alone from 2001 to 2020 (temporal trend: -14 Tg C yr^{-1}, p<0.001) (Fig. 2). However, the increasing sink strength was no longer statistically significant when fire emissions were added to NEE (average NEE + fire budget trend -9 Tg C yr^{-1} over 2002-2020). In the permafrost region, the NEE + fire trend was only 3.3 Tg C yr^{-1}. Nevertheless, based on our NEE upscaling, 23% of the region increased (p<0.05) net CO_2 uptake from 2001 to 2020 (Fig. 1). Most of the increasing net sink activity was driven by an increase in GPP, especially in Siberia (Fig. 2). Some of the trends were also related to a declining R_{eco}, likely associated with disturbed ecosystems (e.g., forest fires, harvesting) with high R_{eco} during the first post-disturbance years now recovering [27]. However, evidence for the increasing overall net uptake trend from the in-situ data is limited due to the low number of long-term sites (>7 years of year-round measurements; 9 sites) out of which only one

site showed a statistically significant trend (increasing uptake at a boreal forest site in Finland).
Some of the relationships in our model are likely thus influenced by spatial differences across
the sites rather than temporal and truly causal patterns, creating some uncertainty in upscaled
trends [28]. However, the model reproduces temporal patterns at individual sites well (see
Supplementary Fig. 6), and our upscaled trends are similar to a recent in-situ time-series
analysis [22] and somewhat similar to those estimated from the inversion ensemble (Fig. 1),
providing confidence in our trend results.

234
Figure 2. An overlay analysis of NEE, GPP, R_{eco} trend maps identifying how trends in GPP and
R_{eco} relate to trends in NEE over 2001-2020 (includes significant and non-significant trends),
showing also pixels that burned during 2002-2020 (a). Terrestrial CO_2 budgets for 1-km (blue;
2001-2020) and 8-km (grey; 1990-2016) NEE, and 1-km NEE + fire emissions (red; 2002-2020)
across the ABZ (b) and permafrost region (c). Trends are shown for the 2002-2020 (NEE + fire)
and 2001-2020 (NEE) periods. Uncertainties are standard deviations in bootstrapped estimates.
Stars in the trend values depict the significance of the trend (*= p<0.05, **=p<0.01,
***=p<0.001).

Parts of the ABZ also show increasing annual net CO_2 emissions over time (Fig. 1). Such trends
have been observed at six long-term sites (2 to 17 g C m^{-2} yr^{-1}, p>0.05), and in 2% of the
upscaled region (p<0.05) from 2001 to 2020. Most of the increasing net emission trends were

driven by an increase in R_{eco} instead of a decline in GPP (Fig. 2). Regions experiencing
increased net CO_2 emissions in upscaling were found especially in (i) northern Europe and
Canada (dominated by evergreen needleleaf forests with mild and moderately wet climates), (ii)
parts of central Alaska and northern Siberia (sparse boreal ecosystems and graminoid tundra
with permafrost and high soil carbon stocks), and (iii) Hudson Bay and Siberian lowlands
(wetlands with some permafrost and high soil organic carbon stocks). Some sites in Alaska
have increasing net emissions of CO_2 due to permafrost thaw [18,29], but it is unclear if similar
changes are occurring in other regions with increasing net CO_2 emissions.
255
We calculated an overall 25% increase in seasonal amplitude of CO_2 fluxes from the upscaled
NEE time series from 2001 to 2020 across the ABZ, on par with earlier atmospheric and
modeling studies [30,31]. Both increasing summer uptake and non-summer season emissions—the
key dynamics driving increasing annual sinks and sources—were evident in the tundra and
boreal biomes (Fig. 3). However, over the 2001-2020 period, the increasing uptake (GPP)
during summer months dominated over increasing net emissions (R_{eco}) during non-summer
months across most of the domain. On average across both biomes, net uptake increased the
most during July (an average upscaled increase of -5 g C m^{-2} $month^{-1}$ in the boreal and -3 g C
m^{-2} $month^{-1}$ in the tundra in 2011-2020 compared to 2001-2010), and increasing net emissions
were occurring throughout the entire non-summer season, with no clear peaks (0.1-0.9 g C m^{-2}
$month^{-1}$). Although increases in early growing season (May-June) uptake were evident, late
growing season (September) trends were absent or minimal (Fig. 3).
268
269 **Drivers of ABZ CO_2 fluxes**
270
There are several environmental conditions driving CO_2 budgets across the ABZ. Our variable
importance analysis showed CO_2 fluxes, and thus the overall increasing sink strength, are
explained by dynamic variables of air or surface temperatures, solar radiation, the Normalized
Difference Vegetation Index (NDVI), and partially also by soil temperature, snow cover, and the
vapor pressure deficit (Supplementary Fig. 8-10). Other less important dynamic variables were
vegetation cover and atmospheric CO_2 concentration. Volumetric soil water content was not
important in our models, likely due to the large uncertainties and coarse spatial resolution in the
gridded product, although in-situ studies have shown drier soils to be linked to larger net CO_2
emissions and wetter soils to enhanced plant growth due to the lack of water limitation [32]. Static
variables (primarily vegetation type, soil carbon stock, soil pH) were also important in explaining
spatial differences.
282
The most important dynamic variables had a positive overall effect on net uptake, GPP, and
R_{eco} (Supplementary Fig. 8-10), however, these relationships are more nuanced in reality. In
fact, the recent permafrost in-situ trend analysis of CO_2 fluxes using the same database
suggests that the CO_2 flux response to warmer temperatures ranges from positive to negative,
depending on the availability of water and nutrients at the site [22]. Consequently, strong warming
or greening trends did not always translate into increasing net CO_2 sinks in our upscaling
(Supplementary Fig. 11). For example, while 49% of the region experienced greening (June-
August average NDVI; based on MODIS NDVI, $p < 0.05$), only 12% of those greening pixels

showed an annual increasing net CO_2 uptake trend, and 29% an increasing June-August net
uptake.
293

Figure 3. Average upscaled monthly NEE, GPP, and R_{eco} in boreal and tundra biomes during the past two decades. Negative NEE values represent net uptake and positive net release. Error bars represent standard deviations across bootstrapped estimates and are only shown for the 2011-2020 period but are similar for the 2001-2010 period. Note that NEE was 1.4 g C m^{-2} month^{-1} lower in September 2011-2020 compared to 2001-2020 in the boreal biome, but this is not shown in the figure. For a similar figure made based on the in-situ data, see Supplementary Fig. 7.

Continental and regional patterns in CO$_2$ budgets and their trends

Our upscaling showed clear continental patterns in NEE budgets and trends (Fig. 4), with the boreal biome primarily driving the budget and trend differences between the continents [1,33]. The increasing net uptake trend was more pronounced in Eurasia (-11 Tg C yr^{-1}, p<0.001) compared to North America (-3 Tg C yr^{-1}, p<0.05), which corresponds with the smaller area, weaker warming, declining snow cover and greening trends in North America (Supplementary Fig. 12-14). We found statistically significant declining summer soil moisture trends in Siberian boreal (Supplementary Fig. 15), but this did not translate into stronger net emissions. When fire emissions were added, continental differences were less pronounced due to the much larger and more rapidly increasing CO$_2$ emissions from Siberian fires (on average 160 compared to 76 Tg C yr^{-1} in North America; Supplementary Fig. 16). Fire emissions even reversed some NEE trends: the strong increasing sink in Siberia became a source when fire emissions were included (trend: +0.7 Tg C yr^{-1}; p > 0.05). However, Siberian ecosystems have the largest uncertainty for both the upscaled fluxes and inversion-based estimates due to lack of in situ observations, making it challenging to accurately determine the magnitude of continental differences (Fig. 4, Supplementary Fig. 17).

Alaska is an important contributor in the weaker North American CO$_2$ sink. Based on our analysis, Alaska as a whole was consistently CO$_2$ neutral or a source over 2002-2020 (NEE + fire emissions), both in the boreal (budget +5 Tg C yr^{-1}) and tundra (budget +7 Tg C yr^{-1}). Alaska has a relatively high density of observations, making this result more certain compared to other regions. Alaska is therefore different from the other ABZ regions where boreal regions still remain on average CO$_2$ sinks. Potential reasons for the Alaskan CO$_2$ source include Alaska having the most rapidly warming autumns and declining autumn snow cover, which has high inter-annual variability (Supplementary Fig. 14 and 18). Further, field measurements suggest that many of the observed changes in Alaskan ecosystems can be attributed to permafrost thaw [18,29]—a phenomenon that has accelerated significantly in response to Alaska's pronounced warming trend since the 1950s [34]. However, we were unable to incorporate permafrost thaw into our models as high-resolution geospatial data from 1990 to 2020 were not available. The question of whether analogous trends will manifest in other regions across the northern permafrost region remains an important research priority.

Figure 4. Terrestrial CO₂ budgets for NEE and NEE + fire in key regions and biomes across the
boreal (a) and tundra (b). Terrestrial CO₂ budgets are shown for 1-km (blue; 2001-2020) and 8-
km (grey; 1990-2016) NEE, and 1-km NEE + fire emissions (red; 2002-2020). Trends are shown
for the 2002-2020 (NEE + fire) and 2001-2020 (NEE) periods. The inset in Alaskan boreal in (a)
shows the time series with a narrower y axis compared to the main figure to better detect
interannual variability. Uncertainties are standard deviations in bootstrapped estimates. Stars in
the trend values depict the significance of the trend (*= p<0.05, **=p<0.01, ***=p<0.001). Fire
emissions alone are shown in Supplementary Fig. 16.

Discussion

Our results show that the ABZ was on average an increasing terrestrial CO_2 sink (GPP is
increasing more than R_{eco} + fire), indicating that the region still creates an important negative
feedback to global warming. However, our study also suggests some positive feedbacks to
climate change that have been more regional and of shorter duration in recent decades. We

show that the presence of annual sources was large as indicated by several site-level and
regional studies [35,36], and even larger with fire emissions included [37]. There were also extreme
years when fire emissions exceeded annual net CO_2 uptake (e.g., 2003 in Siberian boreal, 2012
in Canadian boreal, and several years in the permafrost region; Fig. 2). Moreover, while
summer net uptake increase still dominates over non-summer CO_2 emissions, net CO_2 uptake
is increasing only in the early and peak growing season (May-August in the boreal and June-
August in the tundra) and not in the late growing season (September), because GPP does not
increase later in the season due to plant physiological limitations, and drier and warmer
conditions cause enhanced R_{eco} instead [38–43]. A better understanding of how soil moisture and
hydrology have been and will be changing, and the impact of these changes on CO_2 fluxes is
critical for more accurate ABZ CO_2 budgets.
363
In the tundra, our findings reveal a noteworthy shift in carbon dynamics. While the tundra region
has been a carbon sink for millennia [44], our results suggest that many tundra regions may now
have started to function as CO_2 sources. This transition from an ecosystem CO_2 sink to a CO_2
source may have begun prior to 1990 [45], yet the precise timing of this transformation remains
uncertain. The main drivers of this pattern may be related to warming-induced permafrost thaw,
the drying of soils, or vegetation shifts [46–48] but remain unresolved. Tundra regions are also
progressing towards conditions where average annual soil (0-7 cm) temperatures are above
freezing, resulting in more soil organic material being susceptible to decomposition
(Supplementary Fig. 12). Overall, the primary reason behind the annual CO_2 emissions from
tundra ecosystems is the limited duration of the high net CO_2 uptake period, and the substantial
non-summer season net emissions. However, we observed lower in-situ and upscaled October-
April season NEE fluxes and budgets compared to Natali et al. (2019) throughout the entire
period (Supplementary Fig. 19; [49]).
377
Our results demonstrate the need to further study Siberian CO_2 flux trends. Our upscaling
indicated that some of the strongest net sources and sinks, and strongest increasing sink trends
occur in the Siberian boreal. Increasing sink trends in Siberian tundra were also the strongest
across tundra regions. The Siberian sink trend might be explained by strong greening trends [50],
earlier growing season starts and increasing carbon uptake due to declining spring snow cover
(Supplementary Fig. 14), increases in tree growth and distribution [51,52], rapid recovery of
ecosystems after fire [53], and high cover of larch forests that can rapidly take up CO_2
(Supplementary Table 1)[8,54]. However, the large inversion model spread, sparse measurement
network, and our upscaling uncertainties indicate that it remains challenging to conclude the
magnitude of the Siberian CO_2 balance [2]. This is a significant problem given that Siberia stores
more than half of the permafrost region's C stocks and is now warming more rapidly than other
ABZ regions.
390
In summary, our study reveals distinct spatial and temporal patterns in CO_2 budgets across the
ABZ and underscores the importance of three decades' worth of data. Relatively robust spatial
patterns can be seen, such as the Alaskan CO_2 sources and southern Eurasian boreal sinks
while the temporal trends remain more uncertain. While CO_2 fluxes can be relatively well
modeled using machine learning and existing gridded datasets, gaps persist, such as the

incomplete characterization of thermokarst and harvesting disturbances and their links to
ecosystem CO_2 fluxes, and the lack of accurate predictors describing soil moisture [1]. Sustaining
long-term sites is crucial to accurately track trends in ABZ CO_2 balance, while establishing new
year-round sites in data-poor areas like Siberia and the Canadian Arctic is vital to fill knowledge
gaps and enhance our understanding of carbon dynamics [55].
401
402   **Online Methods**
403
404   **In-situ data overview**
405
We used a recently compiled dataset of in-situ Arctic-boreal terrestrial ecosystem CO_2 flux
measurements (ABCflux, led by Virkkala et al. 2021 [8]) within the ABZ (Supplementary Methods
Section 1). The synthesized data were cumulative fluxes of gross NEE, GPP, and R_{eco}
aggregated at the monthly timescale (3,968 to 4,897 site-months depending on the flux). In
addition to eddy covariance data, we included fluxes estimated with the chamber method to
increase data coverage especially during the growing season. The dataset included metadata
out of which we used the site coordinate, biome, flux measurement method, and disturbance
history information in the analysis. For further details on the dataset, see Virkkala et al. (2021) [8]
and a description of additional data processing and screening in the Supplementary Methods
Section 2. Note that our study does not include lateral transport of carbon which was recently
summarized to be 93 Tg C yr^{-1} in a roughly similar region (i.e., 17 % of the net uptake budget
calculated in this study) [56].
418
This dataset is more comprehensive than the ones used in earlier upscaling studies as it
represents monthly fluxes from the entire year if available, while Virkkala et al. (2021) focused
on coarse seasonal or annual fluxes [5], Natali et al. (2019) on monthly winter fluxes [9], and Mu et
al. (2023) a more limited temporal period (2013-2017) [57]. Furthermore, we included more data
from recent years (805 monthly observations from 2015-2020 compared to 32 and 95 fluxes in
Virkkala et al. 2021 and Natali et al. 2019, respectively), and the sample size in our models was
4 to 25 times larger here compared to the earlier upscaling efforts.
426
427   **Geospatial data**
428
We used data from geospatial products as predictor variables to upscale fluxes. Our models
had the following predictors: month, incident solar radiation, vapor pressure deficit, atmospheric
CO_2 concentration, vegetation type, snow cover (the fraction covered by snow), soil temperature
(0-7 cm), soil moisture (0-7 cm), NDVI (MODIS- or AVHRR-based), land surface temperature
(or air temperature; MODIS- or ERA5 Land-based), compound topographic index (i.e.,
topographic wetness index), continuous vegetation fields describing percent non-tree vegetation
and non-vegetated fraction and percent tree cover (MODIS- or AVHRR-based), soil pH (0-5
cm), soil organic carbon stock in 0-2 meters, and permafrost probability. In our analysis, NDVI
was the primary predictor describing disturbances, with declines in NDVI being related to
disturbances [5]. Data were in daily, weekly, monthly, annual, and static format (i.e., no temporal
changes such as in the compound topographic index). If data were of higher temporal resolution

than monthly, they were aggregated to monthly time steps. Gaps in MODIS and AVHRR NDVI
time series were filled to produce a continuous time series. Data were re-projected to North Pole
Lambert Azimuthal Equal Area Projection at 1 and 8 km spatial resolution and extracted at the
flux sites. See Supplementary Section 3 for further descriptions and data sources.
444
We used the Global Fire Emissions Database (GFED) 500-m fire product [23] to calculate fire
emissions. The product is based on a global fire emissions model with a spatial resolution of
500 m using MODIS data. The model was developed using an updated field measurement
synthesis database of fuel load and consumption which included improvements, for example, in
boreal soil carbon combustion. The higher resolution of the 500-m model compared to earlier
coarser models improved the detection of small-scale fires and understanding of landscape
heterogeneity, and reduced the scale mismatch in comparing field measurements to model grid
cells. However, some small fires might still be undetected by this model, leading to potential
underestimations in carbon emissions in this product.
454

Machine learning modeling

455
We used random forest models to upscale GPP, R_{eco}, and NEE to the ABZ from 1990 to 2020,
the period with in-situ flux measurements. Two sets of predictive models were developed: (i)
models using primarily predictors with a spatial resolution ≤1 km from 2001 to 2020 (i.e., the
MODIS era) at 1-km spatial resolution (hereafter 1-km models;), and (ii) models using coarser-
resolution predictors from the AVHRR GIMMS era (1990-2016;) from 1990 to 2016 at 8-km
spatial resolution (hereafter 8-km models) (Supplementary Table 3). Each model included all
available monthly fluxes from the entire year, i.e. no separate models for individual months or
seasons were developed, as this approach resulted in the best predictive performance. All
models included 17 predictors, but the sample sizes were variable depending on the amount of
data available for each flux and time period; NEE models had the highest amount of model
training data compared to GPP and R_{eco} models. For the 1-km model, coarsest predictors were
at 9-km resolution but most important predictors were at 1 to 4-km resolution. For the 8-km
model, the coarsest predictor resolution was 9 km, and the most important variables had a
resolution of 1 to 9 km.
470
Model parameter tuning was performed based on leave-one-site out cross validation (CV) to
achieve minimum predictive error. The models were run using the "caret" package in R version
4.2 [58]. We assessed the predictive performance of the final models using the (1) R^2, (2) root
mean square error (RMSE), 3) mean absolute error (MAE), and 4) mean bias error (MBE)
between predicted and observed values using the CV data. Larger RMSE and MAE values
indicate larger errors, and positive MBE values indicate overestimation. The predictive
performance of our models was good or high, ranging from 0.55 to 0.78 for R^2 and 19.4 to 37.3
g C m^{-2} month^{-1} for RMSE, but was occasionally limited (Supplementary Fig. 1-3), mostly due to
1) our model not being able to identify landscape heterogeneity with nearby sites showing large
differences in CO_2 fluxes (e.g., a forest and wetland site), and 2) our model not capturing inter-
annual variability at individual sites, both of which are likely attributed to the coarse, uncertain,
and missing predictors characterizing such conditions (e.g., soil moisture, disturbances).
483

We evaluated the uncertainty of predictions by creating 20 bootstrapped model training datasets
(with replacement; same sample size as in the original model training data) and using those to
develop 20 individual models and predictions. Out of the 20 predictions, we calculated the
standard deviation to represent prediction uncertainty. The uncertainty ranges in NEE + fire
budgets only represent NEE uncertainties. We further assessed the area of extrapolation of the
models, and the influence of the flux measurement method and disturbance history information
on flux predictions by training models with different subsets. Further details of the uncertainty
analyses can be found in the Supplementary Methods Section 4.
492
493 **Spatial upscaling of fluxes**
494
495 We upscaled fluxes across the Arctic-boreal terrestrial area ≥49° N [59], which comprises 20.69
496 × 10^6 km^2 of land (excluding glaciers and ice sheets; Fig. 1) with lake and glacier areas
497 removed. The models were applied at a monthly time step from 2001 to 2020 for the 1-km
498 models and from 1990 to 2016 for the 8-km models.
499
We analyzed the upscaled flux maps as well as environmental predictor rasters for temporal
trends using the nonparametric Mann–Kendall test using the "zyp" package [60,61] with pre-
whitening (Zhang method [62]) to remove autocorrelation. We report the significance of Kendall's
correlation coefficient (the strength of the time-series) and the Theil–Sen slope to describe
trends over time. Finally, we calculated zonal statistics of average annual, seasonal, and
monthly fluxes and trends across key regions (Siberia defined as all land east from the Ural
mountains, including a small portion of Mongolia; the rest of Eurasia, including Greenland are
grouped within the European classes), biomes (tundra and boreal) [59], permafrost region [63], and
vegetation types [8].
509
510 **Comparison to process models and atmospheric inversions**
511
We compared our estimates with the CMIP6 process models [26], atmospheric inversions used in
the Global Carbon Project's Global Carbon Budget 2022 [64], and a global upscaling product
FLUXCOM [25]. We included a subset of CMIP6 process models (13 in total) that had soil thermal
processes at several depths to assure they had some information about the freeze-thaw
patterns in the permafrost region. We included inversions with data from the whole 2001-2020
period (i.e., included five inversions and excluded four). Fire CO_2 emissions [23] were subtracted
from the inversions. CMIP6 process model and FLUXCOM outputs were only available for the
2001-2014 and 2000-2013 period, respectively. The final model outputs used here represent
terrestrial NEE (GPP-R_{eco}) in a similar way across the models except for inversions that also
include vertical CO_2 fluxes from water bodies.
522

523 **Acknowledgements**
524
This work was supported by funding from the Gordon and Betty Moore Foundation (grant
#8414) and funding catalyzed by the TED Audacious Project (Permafrost Pathways). We
additionally acknowledge the funding from the NASA Arctic-Boreal Vulnerability Experiment and

Carbon Cycle Science programs (NNX17AE13G), NSF PLR Arctic System Science Research
Networking Activities (RNA, Grant#1931333), Minderoo Foundation, KAKENHI(19H05668),The
Danish National Research Foundation (CENPERM DNRF100), EU HORIZON GreenFeedBack,
grant agreement No. 101056921, Danish Arctic Climate support through Greenland Ecosystem
Monitoring and ICOS grants, Natural Sciences and Engineering Research Council, the
Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's
Excellence Strategy – EXC 2037 'CLICCS - Climate, Climatic Change, and Society' – Project
Number: 390683824, NASA Grant/Cooperative Agreement Number: NNX17AD69A, The
Research Council of Norway (BioGov, project nr. 323945), ERC synergy project Q-Arctic (grant
agreement no. 951288), the Copernicus Atmosphere Monitoring Service, implemented by
ECMWF on behalf of the European Commission (Grant: CAMS2 55), the Environment
Research and Technology Development Fund of the Environmental Restoration and
Conservation Agency of Japan (JPMEERF21S20810), ArCSII(JPMXD142031886), Financial
support from the Swedish Research Council (VR) and consortium partners to ICOS Sweden
(grants 2015−06020 and 2019−00205) and SITES (grant 2017−00635), VR grant 2019-04676
and 2018-03966, ArcticNet and NSERC,NOAA Cooperative Agreement NA16SEC4810008,
Research Council of Finland (NPERM project nrs 341349, 330840, 349503 ICOS-FIRI), ICOS-
FI via University of Helsinki funding, the EU Horizon Europe (GreenFeedback nr. 101056921
and LiweFor nr. 101079192), NRF-2021M1A5A1065425 (KOPRI-PN23011), NRF-
2021M1A5A1065679, NRF-2021R1I1A1A01053870, the Dutch Research Council (NWO)
(project number VI.Vidi.213.143), and The Natural Environment Research Council through the
National Centre for Earth Observation (NE/R000115/1). Part of the research was carried out at
the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the
National Aeronautics and Space Administration (80NM0018D0004). Part of the inverse
analyses were performed on the supercomputer systems at the National Institute for
Environmental Studies and Meteorological Research Institutes (NEC SX-Aurora TSUBASA and
FUJITSU PRIMERGY CX2550M5) and at the HPC cluster Aether at the University of Bremen,
financed by DFG within the scope of the Excellence Initiative (Germany).
556

Data availability

In-situ data used here can be accessed from ORNL DAAC [65]. Environmental predictors and
links to the datasets are: TerraClimate (https://www.climatologylab.org/terraclimate.html), ERA5
Land (https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=overview),
MODIS land surface temperature MOD11A2v006
(https://lpdaac.usgs.gov/products/mod11a2v006/), Barrow CO_2 concentrations
(doi:10.15138/yaf1-bk21), ESA CCI land cover (https://www.esa-landcover-cci.org/), CAVM
vegetation type (https://geobotany.uaf.edu/), MODIS NDVI MOD13A1v006
(https://lpdaac.usgs.gov/products/mod13a1v006/), GIMMS3g NDVI
(https://climatedataguide.ucar.edu/climate-data/ndvi-normalized-difference-vegetation-index-
3rd-generation-nasagfsc-gimms), MOD44B continuous vegetation fields
(https://developers.google.com/earth-engine/datasets/catalog/MODIS_006_MOD44B),
MEaSUREs Vegetation Continuous Fields (https://lpdaac.usgs.gov/products/vcf5kyrv001/), ESA
CCI permafrost probability (https://doi.pangaea.de/10.1594/PANGAEA.888600), SoilGrids

(https://files.isric.org/soilgrids/former/2017-03-10/data/), MERIT DEM topographic indices
(https://doi.pangaea.de/10.1594/PANGAEA.899135) (for more details see Supplementary Table
3). CMIP6 process model outputs can be accessed at https://aims2.llnl.gov/search/cmip6/, and
Global Carbon Budget inversion outputs at https://meta.icos-
cp.eu/objects/GahdRITjT22GGmq_GCi4o_wy (for more details see Supplementary Table 6).
The 1-km and 8-km upscaled rasters of NEE, GPP, and R_{eco} together with their uncertainties will
be published via ORNL DAAC upon publication.
579

580   **Code availability**
581   The main analysis codes can be found in the Supplement.

593   References

594   1.   Watts, J. D. *et al.* Carbon uptake in Eurasian boreal forests dominates the high-latitude net

595   ecosystem carbon budget. *Glob. Chang. Biol.* **29**, 1870–1889 (2023).

596   2.   Fan, L. *et al.* Siberian carbon sink reduced by forest disturbances. *Nat. Geosci.* **16**, 56–62

597   (2022).

598   3.   McGuire, A. D. *et al.* Dependence of the evolution of carbon dynamics in the northern

599   permafrost region on the trajectory of climate change. *Proc. Natl. Acad. Sci. U. S. A.* **115**,

600   3882–3887 (2018).

601   4.   McGuire, A. D. *et al.* Variability in the sensitivity among model simulations of permafrost

602   and carbon dynamics in the permafrost region between 1960 and 2009. *Global*

603   *Biogeochem. Cycles* **30**, 1015–1037 (2016).

604   5.   Virkkala, A.-M. *et al.* Statistical upscaling of ecosystem CO2 fluxes across the terrestrial

605   tundra and boreal domain: regional patterns and uncertainties. *Glob. Chang. Biol.* (2021)

606   doi:10.1111/gcb.15659.

607   6.   Pongracz, A., Wårlind, D., Miller, P. A. & Parmentier, F.-J. W. Model simulations of arctic

608   biogeochemistry and permafrost extent are highly sensitive to the implemented snow

609   scheme in LPJ-GUESS. *Biogeosciences* **18**, 5767–5787 (2021).

610   7.   Pallandt, M. *et al.* Representativeness assessment of the pan-Arctic eddy-covariance site

611   network, and optimized future enhancements. (2021) doi:10.5194/bg-2021-133.

612   8.   Virkkala, A.-M. *et al.* The ABCflux database: Arctic-Boreal CO_2 flux observations and

613   ancillary information aggregated to monthly time steps across terrestrial ecosystems.

614   (2021) doi:10.5194/essd-2021-233.

615   9.   Natali, S. M. *et al.* Large loss of CO2 in winter observed across the northern permafrost

616   region. *Nat. Clim. Chang.* **9**, 852–857 (2019).

10.   Hugelius, G. *et al.* Two decades of permafrost region CO2, CH4, and N2O budgets suggest

a small net greenhouse gas source to the atmosphere. *Authorea Preprints* (2023)

doi:10.22541/essoar.169444320.01914726/v1.

11. Hugelius, G. *et al.* Estimated stocks of circumpolar permafrost carbon with quantified

uncertainty ranges and identified data gaps. *Biogeosciences* **11**, (2014).

12. Schuur, E. A. G. *et al.* Permafrost and Climate Change: Carbon Cycle Feedbacks From the

Warming Arctic. *Annu. Rev. Environ. Resour.* **47**, 343–371 (2022).

13. Rantanen, M. *et al.* The Arctic has warmed nearly four times faster than the globe since

1979. *Communications Earth & Environment* **3**, 1–10 (2022).

14. Berner, L. T. *et al.* Summer warming explains widespread but not uniform greening in the

Arctic tundra biome. *Nat. Commun.* **11**, 4621 (2020).

15. Bjorkman, A. D. *et al.* Plant functional trait change across a warming tundra biome. *Nature*

562, 57–62 (2018).

16. Lund, M. *et al.* Variability in exchange of CO2 across 12 northern peatland and tundra sites.

Glob. Chang. Biol. **16**, 2436–2448 (2010).

17. Happonen, K., Virkkala, A.-M., Kemppinen, J., Niittynen, P. & Luoto, M. Relationships

between above-ground plant traits and carbon cycling in tundra plant communities. *J. Ecol.*

110, 700–716 (2022).

18. Euskirchen, E. S., Bret-Harte, M. S., Shaver, G. R., Edgar, C. W. & Romanovsky, V. E.

Long-Term Release of Carbon Dioxide from Arctic Tundra Ecosystems in Alaska.

Ecosystems **20**, 960–974 (2017).

19. Varner, R. K. *et al.* Permafrost thaw driven changes in hydrology and vegetation cover

increase trace gas emissions and climate forcing in Stordalen Mire from 1970 to 2014.

Philos. Trans. A Math. Phys. Eng. Sci. **380**, 20210022 (2022).

20. Arndt, K. A., Hashemi, J., Natali, S. M., Schiferl, L. D. & Virkkala, A.-M. Recent Advances

and Challenges in Monitoring and Modeling Non-Growing Season Carbon Dioxide Fluxes

from the Arctic Boreal Zone. *Current Climate Change Reports* **9**, 27–40 (2023).

21. Watts, J. D. et al. Regional hotspots of change in northern high latitudes informed by

observations from space. *Authorea Preprints* (2024)

doi:10.22541/au.170497370.03373595/v1. In review at Geophysical Research Letters.

22. See, C. R. et al. Decadal increases in carbon uptake offset by respiratory losses across

northern permafrost ecosystems. In review at *Nature Climate Change* (2024).

23. van Wees, D. *et al.* Global biomass burning fuel consumption and emissions at 500 m

spatial resolution based on the Global Fire Emissions Database (GFED). *Geosci. Model*

Dev. **15**, 8411–8437 (2022).

24. Friedlingstein, P. *et al.* Global carbon budget 2021. *Earth Syst. Sci. Data* **14**, 1917–2005

(2022).

25. Jung, M. *et al.* Scaling carbon fluxes from eddy covariance sites to globe: synthesis and

evaluation of the FLUXCOM approach. *Biogeosciences* **17**, 1343–1365 (2020).

26. Eyring, V. *et al.* Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6)

experimental design and organization. *Geosci. Model Dev.* **9**, 1937–1958 (2016).

27. Ueyama, M. *et al.* Carbon dioxide balance in early-successional forests after forest fires in

interior Alaska. *Agric. For. Meteorol.* **275**, 196–207 (2019).

28. Damgaard, C. A Critique of the Space-for-Time Substitution Practice in Community

Ecology. *Trends Ecol. Evol.* **34**, 416–421 (2019).

29. Celis, G. *et al.* Tundra is a consistent source of CO2 at a site with progressive permafrost

thaw during 6 years of chamber and eddy covariance measurements. *Journal of*

Geophysical Research: Biogeosciences vol. 122 1471–1485 Preprint at

https://doi.org/10.1002/2016jg003671 (2017).

30. Forkel, M. *et al.* Enhanced seasonal CO2 exchange caused by amplified plant productivity

in northern ecosystems. *Science* **351**, 696–699 (2016).

31. Graven, H. D. *et al.* Enhanced seasonal exchange of CO2 by northern ecosystems since

1960. *Science* **341**, 1085–1089 (2013).

32. McGuire, A. D. *et al.* An assessment of the carbon balance of arctic tundra: comparisons

among observations, process models, and atmospheric inversions. *Biogeosci. Discuss.* **9**,

4543 (2012).

33. Lin, X. *et al.* Siberian and temperate ecosystems shape Northern Hemisphere atmospheric

CO2 seasonal amplification. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 21079–21087 (2020).

34. Jorgenson, M. T., Racine, C. H., Walters, J. C. & Osterkamp, T. E. Permafrost Degradation

and Ecological Changes Associated with a WarmingClimate in Central Alaska. *Clim.*

Change **48**, 551–579 (2001).

35. Commane, R. *et al.* Carbon dioxide sources from Alaska driven by increasing early winter

respiration from Arctic tundra. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 5361–5366 (2017).

36. Euskirchen, E. S., Edgar, C. W., Turetsky, M. R., Waldrop, M. P. & Harden, J. W.

Differential response of carbon fluxes to climate in three peatland ecosystems that vary in

the presence and stability of permafrost. *J. Geophys. Res. Biogeosci.* **119**, 1576–1595

(2014).

37. Hayes, D. J. *et al.* Is the northern high-latitude land-based CO2sink weakening? *Global*

Biogeochemical Cycles vol. 25 Preprint at https://doi.org/10.1029/2010gb003813 (2011).

38. Helbig, M. *et al.* Warming response of peatland CO2 sink is sensitive to seasonality in

warming trends. *Nat. Clim. Chang.* **12**, 743–749 (2022).

39. Zona, D. *et al.* Pan-Arctic soil moisture control on tundra carbon sequestration and plant

productivity. *Glob. Chang. Biol.* **29**, 1267–1281 (2023).

40. Zona, D. *et al.* Earlier snowmelt may lead to late season declines in plant productivity and

carbon sequestration in Arctic tundra ecosystems. *Sci. Rep.* **12**, 3986 (2022).

41. Byrne, B. *et al.* Multi-year observations reveal a larger than expected autumn respiration

signal across northeast Eurasia. *Biogeosciences* **19**, 4779–4799 (2022).

42. Pulliainen, J. *et al.* Early snowmelt significantly enhances boreal springtime carbon uptake.

Proc. Natl. Acad. Sci. U. S. A. **114**, 11081–11086 (2017).

43. Parker, T. C., Unger, S. L., Moody, M. L., Tang, J. & Fetcher, N. Intraspecific variation in

phenology offers resilience to climate change for Eriophorum vaginatum. *Arct. Sci.* 1–17

(2021).

44. Brovkin, V. *et al.* Comparative carbon cycle dynamics of the present and last interglacial.

Quat. Sci. Rev. **137**, 15–32 (2016).

45. Oechel, W. C. *et al.* Recent change of Arctic tundra ecosystems from a net carbon dioxide

sink to a source. *Nature* **361**, 520–523 (1993).

46. Lawrence, D. M., Koven, C. D., Swenson, S. C., Riley, W. J. & Slater, A. G. Permafrost

thaw and resulting soil moisture changes regulate projected high-latitude CO2 and CH4

emissions. *Environ. Res. Lett.* **10**, 094011 (2015).

47. Parker, T. C., Subke, J.-A. & Wookey, P. A. Rapid carbon turnover beneath shrub and tree

vegetation is associated with low soil carbon stocks at a subarctic treeline. *Glob. Chang.*

Biol. **21**, 2070–2081 (2015).

48. Voigt, C. *et al.* Warming of subarctic tundra increases emissions of all three important

greenhouse gases - carbon dioxide, methane, and nitrous oxide. *Glob. Chang. Biol.* **23**,

3121–3138 (2017).

49. Schiferl, L. D. *et al.* Using atmospheric observations to quantify annual biogenic carbon

dioxide fluxes on the Alaska North Slope. *Biogeosciences* **19**, 5953–5972 (2022).

50. Piao, S. *et al.* Characteristics, drivers and feedbacks of global greening. *Nature Reviews*

Earth & Environment **1**, 14–27 (2019).

51. Frost, G. V. & Epstein, H. E. Tall shrub and tree expansion in Siberian tundra ecotones

since the 1960s. *Glob. Chang. Biol.* **20**, 1264–1277 (2014).

52. Kharuk, V. I., Ranson, K. J., Im, S. T. & Petrov, I. A. Climate-induced larch growth response

within the central Siberian permafrost zone. *Environ. Res. Lett.* **10**, 125009 (2015).

53. Schulze, E.-D. *et al.* Factors promoting larch dominance in central Siberia: fire versus

growth performance and implications for carbon dynamics at the boundary of evergreen

722 and deciduous conifers. *Biogeosciences* **9**, 1405–1421 (2012).

723 54. Hiyama, T. *et al.* Lessons learned from more than a decade of greenhouse gas flux

724 measurements at boreal forests in eastern Siberia and interior Alaska. *Polar Sci.* **27**,

725 100607 (2021).

726 55. Pallandt, M. *et al.* High-latitude eddy covariance temporal network design and optimization.

727 *Authorea Preprints* (2023) doi:10.22541/essoar.169755225.54015522/v1.

728 56. Ramage, J. L. *et al.* The net GHG balance and budget of the permafrost region (2000-

729 2020) from ecosystem flux upscaling. *Authorea Preprints* (2023)

730 doi:10.22541/essoar.169462008.85493456/v1.

731 57. Mu, C. *et al.* Ecosystem CO_2 exchange and its economic implications in northern

732 permafrost regions in the 21st century. *Global Biogeochem. Cycles* **37**, (2023).

733 58. Kuhn, M. Building Predictive Models in R Using the caret Package. *J. Stat. Softw.* **28**, 1–26

734 (2008).

735 59. Dinerstein, E. *et al.* An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm.

736 *Bioscience* vol. 67 534–545 (2017).

737 60. Sen, P. K. Estimates of the Regression Coefficient Based on Kendall's Tau. *J. Am. Stat.*

738 *Assoc.* **63**, 1379–1389 (1968).

739 61. Zhang + Yue-Pilon Trends Package [R package zyp version 0.11-1]. (2023).

740 62. Zhang, X., Vincent, L. A., Hogg, W. D. & Niitsoo, A. Temperature and precipitation trends in

741 Canada during the 20th century. *Atmosphere-Ocean* **38**, 395–429 (2000).

742 63. Heginbottom, J., Brown, J., Ferrians, O. & Melnikov, E. S. Circum-arctic map of permafrost

743 and ground-ice conditions, version 2. (2002) doi:10.7265/SKBG-KF16.

744 64. Friedlingstein, P. *et al.* Global carbon budget 2022. *Earth Syst. Sci. Data* **14**, 4811–4900

745 (2022).

746 65. Virkkala, A.-M. *et al.* The ABCflux Database: Arctic-boreal CO2 flux and site environmental

747 data, 1989-2020. *ORNL DAAC* Preprint at https://doi.org/10.3334/ORNLDAAC/1934

748   (2021).