What "unexplored" means: Mapping undersampled regions in natural history collections

Laymon Ball[1], Ana M. Bedoya[1], Sheila Rodriguez Machado[1,2], Diego Paredes-Burneo[1,3], Samantha Rutledge[1,2], David Boyd[2], David Vander Pluym[1,2], Spenser Babb-Biernacki[1,2], Austin S. Chipps[1,2], Rafet C. Ozturk[2,4], Yahya Terzi[2,4], Prosanta Chakrabarty[1-2,5-6]*

[1] Department of Biological Sciences, Louisiana State University; Baton Rouge, Louisiana, United States of America
[2] Museum of Natural Science, Department of Biological Sciences, Louisiana State University; Baton Rouge, Louisiana 70803, United States of America
[3] Museo de Historia Natural; Jesús María, Lima, Perú
[4] Department of Fisheries Technology Engineering, Faculty of Marine Sciences; Karadeniz Technical University, Trabzon, Türkiye
[5] American Museum of Natural History; Division of Vertebrate Zoology, Central Park West at 79th Street, New York, New York, United States of America
[6] Smithsonian Institution, National Museum of Natural History; Department of Vertebrate Zoology, Washington, District of Columbia, United States of America

* Corresponding author

Email: prosanta@lsu.edu (PC)

Abstract

We examined global records of accessible natural history voucher collections (with publicly available data) for terrestrial and freshwater vascular plants, fungi, freshwater fishes, birds, mammals, and herpetofauna (amphibians and reptiles) and highlight areas of the world that would be considered undersampled and sometimes called "unexplored" (*i.e.*, have relatively low, or no evidence of, past sampling efforts) under typical Western-scientific descriptions. We also question what "unexplored" may actually mean in these contexts and explain how retiring the term in favor of more nuanced phrasing can mitigate future misunderstandings of natural history science.

Introduction

People have been collecting natural history specimens for centuries and these specimens have been used for gathering valuable data about our changing planet [1–3]. While the number of collections from some regions has steadily increased over time, many other regions have persistent gaps in their collection records. It may be that no naturalist history collectors have ventured into that space because of inaccessibility (*e.g.*, the tops of a tepui, the bottom of a tsingy), political reasons (*e.g.*, it is protected indigenous land, or in a war-torn/unsafe zone), or sometimes these areas may have simply been overlooked [4–5]— for example, some areas may be presumed to be biodiversity-poor (e.g., deserts) and overlooked in favor of well-known areas of high species richness (e.g., coral reefs, rainforests). Alternatively, it could be that specimens from these areas actually exist but have not been digitized— digitization being the "translating [of] metadata associated with a physical specimen object into flexible digital data formats" [6]; funding is a known barrier to digitization at institutions globally [7]. Or collections from these areas may exist, but they remain in a "blind spot" because they have not been aggregated into publicly accessible global databases such as the Global Biodiversity Information Facility (GBIF) which is the single largest biodiversity repository for such data. In this paper, we aim to shed light on what could be meant by "unexplored" by examining the different factors that explain the dearth of apparent collection activities in some parts of the globe.

Collections-based natural history research is the work of observing, securing, obtaining, and preserving wild organisms for future study. Data collection from these organisms can include photographing, measuring, and obtaining blood or tissue samples for future molecular and/or biochemical analyses. These organisms are preserved in a fixative (such as formalin), dried, skinned, and stuffed, or otherwise prepared for long-term comparative use as a reference "voucher" specimen [8–10]. Vouchers are evidence of the existence of these organisms in a particular place and time (that can be compared with specimens from other time periods and locations). These reference specimens document morphological, genetic, and phenological variation of a species or population and are used as proof of a species new to science (as type specimens). Scientific specimens can be used for countless research applications including as part of the "extended specimen" concept [11–14]. In particular, digitized collections facilitate large-scale studies on critical questions in global change biology [6,15], and GBIF-enabled research extends across all major science disciplines [16]. Publicly available natural history collections also facilitate opportunities for international and interdisciplinary collaboration [12].

Unfortunately, natural history collections are in serious decline [17]. Part of this decline is the
shifting in focus of traditional natural history museums away from collections research towards a
focus on public exhibits and sustainability [18], as well as cultural and political resistance to the
idea of collecting certain organisms [19, 20]. It is also related to a decline in basic science
research funding in natural history and the growth of more applied scientific research with
economic goals [18], leading to greater inequalities and mistrust between researchers from
resource-rich versus resource-poor countries or institutions (see "Institutions Holding
Collections" table from [5]). These inequalities lead to the pressing need for institutions
conducting collections-based research to connect and share opportunities more equitably.

Here, we aim to map the regions of the globe highlighting areas where there are relatively few or
no publicly available natural history records and consider the potential reasons behind these
deficiencies. We focus on terrestrial and freshwater habitats, and exclude the vast and still
relatively poorly explored and enormous marine realm [21–23]. Thus, we limited our searches to
terrestrial and freshwater vascular plants, and to terrestrial vertebrates (i.e., birds, herpetofauna,
and mammals) and freshwater fishes. Though oceans comprise the most abundant habitat on
Earth, most natural history research has focused on terrestrial environments [24, 25]. Therefore,
the lack of collections in these terrestrial areas may be more notable and informative than that of
the relatively overlooked and vast oceans.

Regional gaps in specimen collection may not be consistent across taxonomic groups [4]. For
example, an area depauperate in collections of mammals may not be so for freshwater fishes, and
vice versa. Myer [26] identified both taxonomic and geographic biases in plant collections,
particularly towards North America, Western Europe, and Australia, and found that outside of
these regions, high collection coverage tended to be associated with a specific botanical program,
such as the Missouri Botanical Garden's multi-decadal efforts in Madagascar [27]. Therefore, we
attempt here to address regional collection biases for each taxon examined separately and
identify "blind spot" regions that lack scientific collections for multiple taxa. We highlight areas
historically deemed "unexplored" in order to better recognize the reasons behind sampling biases
and raise awareness about habitats that might be worthy of future natural history research and
preservation. Future research may include aiding smaller, private, or non-digitized collections to
become part of the global collections infrastructure by helping to make their vouchers and
associated data accessible and public. We also aim to identify areas needing additional
conservation protection if the current status is insufficient. Our work may help recent advances
in predictive modeling that can be used to track regions with a high potential to hold hidden
biodiversity [28]. We hope this study will help identify the gaps in our understanding of the
world's biodiversity and draw attention to how we identify areas that have been potentially
overlooked by those interested in the exploration and conservation of the natural world.

Results and Discussion

We examined over 55 million (55,200,291) occurrence records from each continental region
(Table 1) as defined herein, and compared the patterns of sampling efforts for natural history
collections for all six major taxonomic groups. We focus on the shared poorly-collected regions
and those unique to each of the six taxonomic groups included. In some countries, an artifact of

older digitized data is that localities may be listed as simply "Brazil" or "Peru," causing the
centroid (relative midpoint) of that country to be the area highlighted as most "collected." We
avoid descriptions of such artifacts here. However, we note clear cases where the presence of a
local museum, herbarium or fungarium increases local specimen collections relative to other
regions.

Table 1: Specimen records by continental region and taxon

	South America	North and Middle America	Africa	Australia and Oceania	Eurasia	Antarctica
Birds	361,711	1,856,474	227,471	445,378	535,312	2,202
Fishes	360,685	1,484,529	188,976	179,446	443,824	1,128
Fungi	162,228	1,280,467	59,101	416,647	1,972,661	9,967
Herps	436,020	2,278,628	256,039	825,493	251,687	-
Mammals	279,689	2,033,383	317,256	278,725	555,010	-
Plants	6,723,374	10,271,258	2,089,744	6,342,071	12,269,675	4,032

125 Data, originally obtained from GBIF, were filtered to exclude observational data without
126 vouchers and to only include specimen records with geographic coordinates in terrestrial
127 systems, see "Materials and Methods" section for more information. Note: "Herps" is a
128 shorthand referring to "Herpetofauna" i.e., amphibians and reptiles.
129

South America
Collecting records across South America show that across taxa, publicly available collections are
more numerous in mountainous regions and well-known biodiversity hotspots (*i.e.* the Andes,
Atlantic forest, and the Guiana Highlands; Fig 1). This pattern is likely a result of collecting
efforts being centered in well-known areas of high species richness. Similarly, digitized
collections along the Andean mountains show a latitudinal gradient in sampling effort. This
pattern is consistent across taxonomic groups and is likely linked to a well-documented
latitudinal biodiversity gradient [29]. Every group examined appears to have a high number of
collections in the Atlantic Forest of Brazil and Uruguay relative to dryer adjacent regions
including the savannas of the Cerrado. However, estimates of undersampled terrestrial vertebrate
diversity made using publicly available data have identified hotspots like the Andean mountains
as among the most undersampled regions [4]. Assessing the extent of the scarcity of data
collection in highly biodiverse regions remains a challenge. There are relevant records stored in
private collections, as well as specimens at local institutions which have not, or have only
partially been aggregated into digital repositories (e.g., Museum of Zoology of the University of

São Paulo, Cartagena Botanical Garden "Guillermo Piñeres", the National Museum of Brazil, the
Museo de Historia Natural in Lima), which remain hidden to the international scientific
community.
148
149 **Fig 1. Maps of collecting efforts on South America.** Sample (voucher) quantity for each taxon
150 considered in this study is shown. Red colors represent 100K+ specimens, white and very light
151 colors represent the so-called 'unexplored' areas or areas with few collections.
152
The Amazon basin, one of the largest and most biodiverse regions of the world (holding 10% of
all named plant and vertebrate species; [30–35]), has lower sampling across taxa (particularly for
fungi and mammals; Fig 1). Large sections of the Amazon remain largely undersampled, with
botanical census data covering <0.0003% of the region [32]. Other lowland areas with fewer
publicly available records across taxa include the Orinoco basin, the Atacama Desert, the
southern portion of the Chaco, and the Patagonian grasslands. While fewer digitized collections
in the Atacama may be representative of the region's low diversity for the six taxonomic groups
examined here, accessibility and armed conflict may be relevant factors limiting exploration in
the other regions. Scarcity of roads or navigable water bodies are a challenge to transportation to
and from large sections of those regions, increasing the costs, risks, and logistics for field
exploration [36–37]. Political and civil unrest in Latin America have historically limited
collecting in these and other regions and may have kept many collections there from being part
of a global digitized database. This lack of digitization may be reflected in the dearth of
collections represented in what are very biodiverse countries. In fact, such biodiversity is related
in part to a positive relationship between forest cover and the intensity of armed conflict [38–
41].
169
Specimen data for plants are the most abundant compared to the other taxonomic groups
examined here. However, it has been estimated that there are zero publicly available botanical
records from about 10% of tropical South America [32]. A higher number of botanical
collections may be due to higher plant species richness compared to other groups. In addition,
the common practice of collecting multiple replicates per specimen to be shared across herbaria
could result in duplicated digitized records and in a higher probability of a given specimen being
digitized.
177
Freshwater fishes are the most speciose group of vertebrates that we examined, and their species
richness peaks in South America, specifically along the main tributaries of the Amazon River
basin (circa 7000 species; [42]). The Amazon Basin and major tributaries appear much better
collected (and digitized) than the Orinoco River Basin which covers much of Venezuela and
Colombia. In Venezuela, only the Universidad Central has published data in GBIF (for insects),
and many are privately held. The "La Plata" region (including the Parana and Paraguay) in east
central South America is also poorly sampled and/or digitized relative to the Amazon. Notably,
the Deseado River estuary, Lake Musters, and Nuevo Gulf in Argentina are well-sampled for
fishes. Compared to the other groups examined, there are comparatively fewer records of fungi.
Intensively "explored" regions for this group may be a consequence of the presence of
institutions where this taxonomic group has been well sampled and where collections have been
digitized. For instance, southeastern Brazil appears more intensively sampled because the
Herbarium at Federal University of Pernambuco is located in this area and has the largest

collection of fungi in Brazil [43]. Relative to northern South America, Brazil appears sparsely
sampled for specimens of mammals, herps, and birds, especially relative to plants and fishes.
193

North and Middle America [including landmasses in the Gulf of Mexico, Caribbean and West to Hawaii]

196
North America is one of the most consistently sampled continents with all taxonomic groups
exhibiting similar decreases in sampling effort along a south-to-north latitudinal gradient and few
areas devoid of specimen samples of any group south of Canada (Fig 2). Excluding Greenland,
interior Canada is the largest gap for all plant and animal groups examined, from Ontario
northwest to NW Territories and east to Newfoundland and Labrador. Canada may remain
relatively undersampled in part because of its enormous size; the lack of knowledge about its
abundant freshwater resources has been previously noted [44]. Relative to adjacent countries it
appears Nicaragua is poorly sampled for all vertebrate groups and fungi despite having an
abundance of biodiversity-rich regions including the Mosquito Coast, Lake Nicaragua, and the
largest tropical rainforest north of the Amazon [45]. Otherwise, most of Central America and the
Caribbean/Greater Antilles appear well sampled with a clear gap for Cuba, due more so to
conflicting politics and policies (at least with the United States) than to a lack of biodiversity
[46]. Conservation work in Cuba is among the best in the region [47] and the Museo Nacional de
Historia Natural de Cuba in Havana has digitized collections but these are not publicly available.
Perhaps not surprisingly the United States is very well sampled, and climatic differences between
the wetter, more humid east versus the more arid west can be seen when comparing fungi and
fishes from those regions; with the more temperate Pacific Northwest being an exception.
214
215 **Fig 2. Maps of collecting efforts on North America and adjacent regions.** Sample (voucher)
216 quantity for each taxon considered in this study is shown.
217
Mammals have weaker collection records from the southeast and midwest United States than
other vertebrate groups. Specimens of amphibians and reptiles (herpetofauna) display the
sharpest decline moving northward of all examined groups, with few specimens collected in and
east of the Rocky Mountains from Montana to Illinois. Fishes are, unsurprisingly, sparsely
collected from the northern Rocky Mountains south to central Mexico including adjacent desert
regions. Although most vertebrate groups display strong holdings from the southeastern United
States, non-coastal southern Appalachia is relatively poorly represented in Bird collections, as
are Montana, Wyoming, and Utah in the west. Plants are the most consistently sampled group
across North America with few, if any, undersampled regions, while Fungi have the highest
number of unsampled areas, especially in and around the Chihuahuan Desert.
228

Africa [including Madagascar and adjacent landmasses]

230 Perhaps more than any other region, Africa's "unexplored" areas likely reflect the sparsity of
231 digitization of regional collections as well as the general lack of collections (Fig 3). A wide
232 variation in climate such as the dry Sahara region versus sub-Saharan tropical regions are also
233 important explanations for variation in collecting effort within the continent. A striking
234 juxtaposition is seen between relatively well-sampled Madagascar and adjacent islands such as

the Seychelles of the southeastern coastline, particularly for plants, which is likely a result of sampling efforts from the Missouri Botanical Garden which has a permanent base situated on the large island, and a long history of botanical exploration there; today, the Garden accounts for an estimated 72% of vascular plant collections from Madagascar. Unfortunately, this means the largest collection of Malagasy plants exists well outside of the island which would hinder the growth of local knowledge on these organisms.

Fig 3. Maps of collecting efforts in Africa and adjacent regions. Sample (voucher) quantity for each taxon considered in this study is shown.

Herps show a similar pattern of natural history sampling effort to plants with a diagonal band across the tropical southern equatorial regions. Collections for most groups are generally tracking with humidity and rainfall, save for fungal and fish diversity which is likely higher than represented by the few collections that have been made across the continent, including Madagascar. Notably, Cape Verde is a bright spot for fungal collections on the West Coast of Africa almost entirely accounted for by collections from lichenologists at the Senckenberg Research Institute and Natural History Museum in Frankfurt, Germany, demonstrating the outsized impact a single research program can have on perceptions of whether an area has been 'explored' [48]. The Great Rift Valley region is particularly well sampled for mammals, as are the southern reaches of West Africa (from Liberia to Benin). Madagascar, given its high proportion of endemic mammal species, has few museum specimens; this is likely due to the conservation status of many of the species, including the endangered endemic lemurs.

Despite being largely tropical, Nigeria appears to be a surprising gap for several groups relative to adjacent countries. Similarly, Somalia, the Central African Republic, the Democratic Republic of the Congo, and Angola are gaps for several groups. The sparsity of collections in these regions can be attributed to a combination of complex factors including a history of colonization, lack of scientific infrastructure and institutional support, geographic accessibility, and historical and present-day political instability (particularly in central Africa; plant collections have been noted to decline during periods of war [49]). While not exclusive to these poorly collected regions, cultural factors are also an important consideration (e.g., in some areas there may be cultural beliefs that discourage the collection of natural history specimens; [34, 49]). We call for collections in Africa to join the GBIF network but also call on institutions outside of Africa with African collections to publish their records (some Western museums have collections from the region that are not yet digitized). Egyptian collections from Cairo, Alexandria, and along the Red Sea for several groups are notable and likely were a target for several important historical collections. Interior reaches of Africa, outside of the desert regions, including the Okavango Delta still have relatively few collections. However, given that regions celebrated conservation status, efforts to collect in this area should be well-regulated. Notably many collections from this region are in Europe and the United States, and efforts to digitize existing collections and create new regional collections should be considered as they would benefit local knowledge.

Australia and Oceania

Unsurprisingly, the large swath of arid land through central Australia is less densely sampled than tropical northern regions and the temperate south, due to the relative dearth of species

biodiversity in this arid climate, particularly for large vertebrates (Fig 4; [50]). This portion also remains logistically difficult to access for collecting, due to its desert climate and relative lack of travel infrastructure (i.e., roads). Tasmania appears to be a hotspot for specimens of all taxonomic groups, as does the region near Perth in the Southwest, due in part to their more tropical climates, easier accessibility, and the presence of natural history museums in each locale [43, 50].

Fig 4. Maps of collecting efforts on Australia and adjacent regions in Oceania (such as the Indonesian archipelago and New Zealand). Sample (voucher) quantity for each taxon considered in this study is shown.

A sharp divide exists in New Guinea between New Guinea (the western Indonesian side often called "Irian Jaya") and the Eastern side of the island, Papua New Guinea, which has been a target for many natural history expeditions [51-53]. Another noticeable division occurs between north and south of the New Guinea Highlands, with the south being poorly collected versus its northern counterpart. This is the case for both the Indonesian and Papuan halves of New Guinea, most likely due to the heavy monsoon flooding that seasonally submerges areas in that portion of the island [54].

Despite its tropical climate and abundance of biodiversity, there appear to be few collections from Indonesia, particularly for fungi. Contrastingly, the Philippines (central top of map) are densely collected relative to Indonesia, again showing how Indonesia appears poorly collected for natural history specimens. It is also possible that Indonesia only appears relatively "unexplored" because of a dearth of accessible digitized specimens (likely in collections in the West). Indonesia has been a target for museum-based collections since occupation by the Dutch and Wallace's early expeditions to the 'Malay Archipelago.' Few if any of the specimens from southeast Asia collected by Wallace or used as type material by Western zoologists are curated by local institutions. Specimens collected from this region more recently are curated in large institutions like the Museum Zoologicum Bogoriense in Java, Indonesia, which has yet to have its extensive collections digitized. Notably, no museum collection from Oceania is listed among the 73 of the world's largest natural history museums and herbaria from 28 countries with the nearest of these surveyed collections being in India and Australia [5]). Representation of people from this region and digitization of local collections should be a priority before labeling this entire archipelago "unexplored." The absence of a fungarium (despite there being herbaria) is an oversight and can be linked to why there are few fungal collections in an area that should be a hotspot for mycologists. Borneo appears remarkably underrepresented in natural history collections for herps and mammals, particularly relative to nearby Java; the lack of mammal and herp collections from New Zealand is less surprising because of the relatively few species of these groups known to exist there. Mammals and herps are particularly well collected in the Philippines and Papua New Guinea making adjacent areas even more glaringly under-sampled. Birds and plants appear to be the best sampled taxa from Indonesia, perhaps as a result of historic collecting efforts but large gaps remain in New Guinea, Borneo, Sulawesi, and Sumatra.

Eurasia

Much of Europe has been well collected and the specimens digitized from the region are among
the highest on the planet; in stark contrast to the rest of Eurasia except the far East (particularly
Japan and Taiwan which have substantial collections and active natural history researchers; Fig
5). The relative absence of digitized collections from India for plants, fungi and mammals is
unexpected, given the region's rich biodiversity. Sri Lanka appears well-sampled relative to
India for several groups which may reflect a greater accessibility to collections and digitization
of specimens from that island nation. South Korea, Nepal, and Taiwan appear to have a high
number of collections of birds, mammals, plants, and herps relative to much of "mainland"
China, although the warm temperate areas of Southern China appear well collected for some
groups such as plants and herps making the absence of mammals and birds from this region even
more striking. Much of the cold vastness of Russia too seems undersampled for natural history
specimens. Yet bright spots on the Russian map shed light on the importance of local museums
in the pursuit of cataloging biological diversity. For example, the high concentration of fungi
samples near the western central part of the country is not a centroid artifact but the Yugra State
University Biological Collection: an institution that has been collecting mycological specimens
since the early 20th century [55]. Lack of digitization, collecting efforts, climate, and politics
probably have a great deal to do with the lack of vouchered samples from the world's largest
country.

342
343 **Fig 4. Maps of collecting efforts on Eurasia**. Sample (voucher) quantity for each taxon
344 considered in this study is shown.
345

Antarctica

Antarctica is notable only in the absence of many collections for obvious reasons, there are no
collections of freshwater fishes, amphibians or reptiles because none survive on the continent.
Large mammal and bird collections do exist but are generally restricted to historic collections or
from areas near the coasts including some not yet recorded on GBIF, similarly, lichen and plants
are generally restricted to ice-free zones off of the continent (Fig 5). Observations of Antarctic
fauna have been made but many of these are of invertebrates and fishes below the sea ice [56].

353
354 **Fig 5. Maps of collecting efforts on Antarctica.** Sample (voucher) quantity for each taxon
355 considered in this study is shown.

Conclusions

The term "unexplored" can be used in many unrelated contexts, and it may not accurately reflect
the state of knowledge in a given area. To some, "unexplored" may mean an area that has never
been seen by human beings, while to others, it may mean an area that is not exploited by
Westerners or is uninhabited. In examining the so-called "unexplored" places that remain on
land, we find that a lack of digitized natural history collections may be a common cause for this
label. In some cases, there may be a lack of natural history samples from an area because there
are relatively few organisms there to collect (e.g., the Atacama Desert in Chile and Sahara Desert
of Africa which are some of the driest places on Earth, as well as many parts of Antarctica). In
some regions, political unrest has prevented access to naturalists (e.g., Libya, Venezuela,
Colombia), and a lack of infrastructure for obtaining permits and permissions hurt both the

global understanding of a region and local biodiversity education. Other areas appear simply to
have been overlooked by natural history research— the limited number of practicing naturalists
with even more limited funding may be an explanation for some of these regions remaining
poorly researched. The term "unexplored" in these contexts can be harmful, as such terminology
may inadvertently perpetuate misconceptions or undermine the value of existing research and the
efforts of local scientists and indigenous people or disregard the underlying resource-imbalances
impeding natural history research, collection, and digitization in many regions. For these reasons,
we advocate for retiring the word "unexplored" for more precise and inclusive language (e.g.,
"this region lacks digitized natural history collections") and suggest "biodiversity blindspots" as
an alternative.

In this paper, we highlight areas lacking or having few collections, not to encourage their
exploitation, but to call for an increased understanding of these areas in the context of global
biological diversity for the sake of conservation and recognition in a natural history context. A
multitude of factors have kept these places from the growing knowledge base of natural history
and biodiversity, and in some cases, particularly where indigenous stakeholders are the
protectors [57], traditional Western approaches to science may not be the most effective means
of including these regions [58]. Good faith partnerships and collaborations, a fair exchange of
knowledge and resources, and the acknowledgement that there are many different ways of
knowing that are equally contributable, will be necessary starting points to conversations related
to natural history in many of these areas (e.g., large swaths of the Amazon Rainforest that are
home to uncontacted tribes). Recent calls for equity-based fieldwork should be heeded, and
special care should be taken to avoid practicing parachute science, a term that refers to situations
in which researchers, often from more economically developed or privileged countries, visit
other marginalized countries to conduct research and then return to their home institutions
without genuinely engaging or collaborating with locals [59–61]. Some of the most biologically
diverse and well collected countries are also among the most exploited and experience high
incidences of parachute science. Any future work in undercollected regions should be performed
in ways that are considerate of and inclusive to local researchers and communities.

Accessibility to financial resources that make digitization (especially large-scale efforts) possible
are a significant barrier for many museums and institutions— even for some larger institutions in
the West. (e.g., based on GBIF data most natural history specimen collections from Costa Rica,
Mozambique, and India are found outside of those countries). There are large museums that are
not yet digitized and part of GBIF in some of the most biodiverse places, including but not
limited to the Museum of Zoology of the University of São Paulo, the Museu Paraense Emílio
Goeldi, Inpa Coleçoes Zoológicas, and the National Museum in Rio de Janeiro (which suffered a
devastating fire in 2018; [62]). Modernizing access to the reference vouchers housed at these
institutions would likely benefit research groups globally and facilitate natural history and
conservation research. However, international digitization efforts will require sharing resources
to create scientific infrastructure where it is lacking, especially in places where biodiversity is
richest and the people poorest. It should be noted that while there are many benefits to
digitization, care needs to be taken to not perpetuate colonialism. The benefits of digitization will
not be realized equitably without global collaborative efforts that are inclusive to regions with
fewer economic resources, and offer a fair exchange knowledge, technology, and support to
empower these areas in their scientific endeavors.

The goal of this study was to increase our knowledge of "biodiversity blindspots" by
highlighting these poorly-collected or undigitized places, and we hope these focus areas can be
topics of discussion by researchers, politicians, indigenous stakeholders, and others who are
interested in the discovery-based science of biodiversity research.

Materials and Methods

Occurrence records of preserved specimens with GPS coordinates were downloaded from GBIF
between July 1 and December 31, 2022 (Global Biodiversity Information Facility;
https://www.gbif.org/) for each of six taxonomic groups: fungi, terrestrial and freshwater
vascular plants, freshwater fishes, herpetofauna (amphibians and reptiles), birds, and mammals.
As we are interested in general trends in collection records across continents, only occurrence
records with associated voucher specimens were considered for each of the six taxonomic groups
included in this study (we did not include observational data where no specimens were
collected). To maximize the amount of georeferenced data retrieved from GBIF, we did not
apply any other filters (e.g., continent, year) as georeferenced records may be missing
information for such fields. Given the size of the vascular plant's dataset (>126 million records),
we split records by continent in R v.x.x (readr package; [63]) using the maximum and minimum
latitude and longitude for each continent.

Available records from GBIF for each dataset (taxonomic group and data subsets for vascular
plants) were cleaned in QGIS v.3.10.9 (QGIS.org, 2023. QGIS Geographic Information System.
QGIS Association. http://www.qgis.org), to include records that fall within continent limits
(including islands) using the "World Countries" shapefile as downloaded from ArcGIS. We used
QGIS to split the world map into separate shape files corresponding to six regions: South
America (with adjacent islands), North America (including landmasses in the Gulf of Mexico,
Caribbean and west to Hawaii), Africa (including Madagascar and adjacent landmasses),
Australia + Oceania (including islands in Oceania such as the Indonesian archipelago and New
Zealand), Eurasia (including all of continental Europe and Asia as well as adjacent islands such
as the British Isles, Iceland, Svalbard Archipelago, Japan and the Philippines), and Antarctica.
These continent shape files were used to define the extent of polygon grids, which were in turn
used to count the number of occurrence records per one-degree squared grid cell. Occurrence
records that did not intersect with continent layers were removed. This procedure also removed
data points exactly at 0°, 90°, or ≥180°, or in the ocean. Using the "count points in polygon" tool,
the number of collections per grid cell were calculated across each continent. The number of
collections in grids were log10 transformed using Field Calculator in QGIS. Collection gaps
were then visually identified. To assess whether global differences in collection intensity could
be due to climatic differences (e.g., lower species diversity in hot desert areas might lead to
fewer collections in these regions - see [64]), collections maps were also compared with Köppen-
Geiger climate maps (http://koeppen-geiger.vu-wien.ac.at/present.htm).

An interactive map was generated using QGIS2Web plugin (ver. 3.16.0) and can be found at
https://canon-network.github.io/ under the "Specimen Database" tab. This map shows all
collected specimen numbers divided into one-degree squared grid cells covering the globe and
can be toggled for individual taxonomic groups. In the static maps shown below (Figs 1-5)

lighter areas are poorly sampled regions based on GBIF records, while darkly colored (red) regions are well-collected (100K+ specimens) with many specimens from those areas. The specimen data was log-transformed to increase the color contrast of poorly sampled areas so they would be pronounced well on the maps. Non-log transformed data can be visualized in the interactive map described earlier in this paragraph.

References

1. Lane MA. Roles of natural history collections. Annals of the Missouri Botanical Garden. 1996 Jan 1:536-45.
2. Lister AM. Natural history collections as sources of long-term datasets. Trends in ecology & evolution. 2011 Apr 1;26(4):153-4.
3. Rocha LA, Aleixo AL, Allen G, Almeda F, Baldwin CC, Barclay MV, et al. Specimen collection: an essential tool. Science. 2014 May 23;344(6186):814-5.
4. Šmíd J. Geographic and taxonomic biases in the vertebrate tree of life. Journal of Biogeography. 2022 Dec;49(12):2120-9.
5. Johnson KR, Owens IF, Global Collection Group. A global approach for natural history museum collections. Science. 2023 Mar 24;379(6638):1192-4.
6. Hedrick BP, Heberling JM, Meineke EK, Turner KG, Grassa CJ, Park DS, et al. Digitization and the future of natural history collections. BioScience. 2020 Mar 1;70(3):243-51.
7. de Araujo ML, Quaresma AC, Ramos FN. GBIF information is not enough: national database improves the inventory completeness of Amazonian epiphytes. Biodiversity and Conservation. 2022 Sep;31(11):2797-815.
8. Remsen JV. The importance of continued collecting of bird specimens to ornithology and bird conservation. Bird Conservation International. 1995 Sep;5(2-3):146-80.
9. Buckner JC, Sanders RC, Faircloth BC, Chakrabarty P. The critical importance of vouchers in genomics. Elife. 2021 Jun 1;10:e68264.
10. Poo S, Whitfield SM, Shepack A, Watkins-Colwell GJ, Nelson G, Goodwin J, et al. Bridging the research gap between live collections in zoos and preserved collections in natural history museums. BioScience. 2022 May;72(5):449-60.
11. Webster MS, Cicero C, Bates J, Hackett S, Joseph L. Ornithological Collections in the 21st Century 1. InThe Extended Specimen 2017 Jul 20 (pp. 219-232). CRC Press.
12. Meineke EK, Davies TJ, Daru BH, Davis CC. Biological collections for understanding biodiversity in the Anthropocene. Philosophical Transactions of the Royal Society B. 2019 Jan 7;374(1763):20170386.
13. Lendemer J, Thiers B, Monfils AK, Zaspel J, Ellwood ER, Bentley A, et al. The extended specimen network: A strategy to enhance US biodiversity collections, promote research and education. BioScience. 2020 Jan 1;70(1):23-30.
14. Monfils AK, Krimmel ER, Linton DL, Marsico TD, Morris AB, Ruhfel BR. Collections Education: The Extended Specimen and Data Acumen. BioScience. 2022 Feb;72(2):177-88.
15. Ford KL, Albert JS, Summers AP, Hedrick BP, Schachner ER, Jones AS, et al. A New Era of Morphological Investigations: Reviewing methods for comparative anatomical studies. Integrative organismal biology. 2023;5(1):obad008.

16. Heberling JM, Miller JT, Noesgaard D, Weingart SB, Schigel D. Data integration enables
global biodiversity synthesis. Proceedings of the National Academy of Sciences. 2021
Feb 9;118(6):e2018093118.
17. Rohwer VG, Rohwer Y, Dillman CB. Declining growth of natural history collections
fails future generations. PLoS Biology. 2022 Apr 19;20(4):e3001613.
18. NAGGS F. The tragedy of the Natural History Museum, London. Megataxa. 2022 Jul
11;7(1):86-112.
19. Byrne AQ. Reimagining the future of natural history museums with compassionate
collection. PLoS biology. 2023 May 4;21(5):e3002101.
20. Nachman MW, Beckman EJ, Bowie RC, Cicero C, Conroy CJ, Dudley R, et al.
Specimen collection is essential for modern science. PLoS biology. 2023 Nov 22;21(11):
e3002318.
21. Costello MJ, Coll M, Danovaro R, Halpin P, Ojaveer H, Miloslavich P. A census of
marine biodiversity knowledge, resources, and future challenges. PloS one. 2010 Aug
2;5(8):e12110.
22. Albano PG, Azzarone M, Amati B, Bogi C, Sabelli B, Rilov G. Low diversity or poorly
explored? Mesophotic molluscs highlight undersampling in the Eastern Mediterranean.
Biodiversity and Conservation. 2020 Dec;29(14):4059-72.
23. Molony BW, Ford AT, Sequeira AM, Borja A, Zivian AM, Robinson C, et al.
Sustainable Development Goal 14-Life Below Water: Towards a Sustainable Ocean.
Frontiers in Marine Science. 2022 Feb 24;8:829610.
24. Dayton PK. The Importance of the Natural Sciences to Conservation: (An American
Society of Naturalists Symposium Paper). The American Naturalist. 2003 Jul;162(1):1-3.
25. Oestreich WK, Chapman MS, Crowder LB. A comparative analysis of dynamic
management in marine and terrestrial systems. Frontiers in Ecology and the Environment.
2020 Nov;18(9):496-504.
26. Meyer C, Weigelt P, Kreft H. Multidimensional biases, gaps and uncertainties in global
plant occurrence information. Ecology letters. 2016 Aug;19(8):992-1006.
27. Birkinshaw C, Lowry II PP, Raharimampionona J, Aronson J. Supporting Target 4 of the
Global Strategy for Plant Conservation by Integrating Ecological Restoration into the
Missouri Botanical Garden's Conservation Program in Madagascar1. Annals of the
Missouri Botanical Garden. 2013 Dec;99(2):139-46.
28. Parsons DJ, Pelletier TA, Wieringa JG, Duckett DJ, Carstens BC. Analysis of
biodiversity data suggests that mammal species are hidden in predictable places.
Proceedings of the National Academy of Sciences. 2022 Apr 5;119(14):e2103400119.
29. Fine PV. Ecological and evolutionary drivers of geographic variation in species diversity.
Annual Review of Ecology, Evolution, and Systematics. 2015 Dec 4;46:369-92.
30. Nelson BW, Ferreira CA, da Silva MF, Kawasaki ML. Endemism centres, refugia and
botanical collection density in Brazilian Amazonia. Nature. 1990 Jun 21;345(6277):714-
6.
31. Schulman L, Toivonen T, Ruokolainen K. Analysing botanical collecting effort in
Amazonia and correcting for it in species range estimation. Journal of Biogeography.
2007 Aug;34(8):1388-99.
32. Feeley K. Are we filling the data void? An assessment of the amount and extent of plant
collection records and census data available for tropical South America. PLoS One. 2015
Apr 30;10(4):e0125629.

33. Winemiller KO, McIntyre PB, Castello L, Fluet-Chouinard E, Giarrizzo T, Nam S, et al.
Balancing hydropower and biodiversity in the Amazon, Congo, and Mekong. Science.
2016 Jan 8;351(6269):128-9.
34. Stropp J, Umbelino B, Correia RA, Campos-Silva JV, Ladle RJ, Malhado AC. The
ghosts of forests past and future: deforestation and botanical sampling in the Brazilian
Amazon. Ecography. 2020 Jul;43(7):979-89.
35. Albert JS, Carnaval AC, Flantua SG, Lohmann LG, Ribas CC, Riff D, et al. Human
impacts outpace natural processes in the Amazon. Science. 2023 Jan
27;379(6630):eabo5003.
36. Daru BH, Park DS, Primack RB, Willis CG, Barrington DS, Whitfeld TJ, et al.
Widespread sampling biases in herbaria revealed from large-scale digitization. New
Phytologist. 2018 Jan;217(2):939-55.
37. Hijmans RJ, Garrett KA, Huaman Z, Zhang DP, Schreuder M, Bonierbale M. Assessing
the geographic representativeness of GenBank collections: the case of Bolivian wild
potatoes. Conservation Biology. 2000 Dec 18;14(6):1755-65.
38. Álvarez MD. Forests in the time of violence: conservation implications of the Colombian
war. InWar and Tropical Forests 2020 Sep 10 (pp. 49-70). CRC Press.
39. McNeely JA. Conserving forest biodiversity in times of violent conflict. Oryx. 2003
Apr;37(2):142-52.
40. Hanson T, Brooks TM, Da Fonseca GA, Hoffmann M, Lamoreux JF, Machlis G, et al.
Warfare in biodiversity hotspots. Conservation Biology. 2009 Jun. 23(3):578-87.
41. Negret PJ, Allan J, Braczkowski A, Maron M, Watson JE. Need for conservation
planning in postconflict Colombia. Conservation Biology. 2017 Feb.
42. Albert JS, Reis R, editors. Historical biogeography of Neotropical freshwater fishes. Univ
of California Press; 2011 Mar 8.
43. Major increase in Brazilian plant and fungus data shared through gbif [Internet]. 2015
[cited 2023 Feb 7]. Available from: https://www.gbif.org/news/82393/major-increase-in-
brazilian-plant-and-fungus-data-shared-through-gbif
44. Desforges JE, Clarke J, Harmsen EJ, Jardine AM, Robichaud JA, Serré S, et al. The
alarming state of freshwater biodiversity in Canada. Canadian Journal of Fisheries and
Aquatic Sciences. 2022;79(2):352-65.
45. Weaver PL, Lombardo DM, Martinez-Sanchez JC. Biodiversity and tropical forest
conservation, protection and management in Nicaragua: assessment and
recommendations. Evaluation performed for USAID/Nicaragua. 2003 Apr 15.
46. Denis D, Cruz-Flores DD, Testé E. Biodiversity in Cuba. In: Global Biodiversity 2018
Dec 7 (pp. 139-176). Apple Academic Press.
47. Goulart F, Galán ÁL, Nelson E, Soares-Filho B. Conservation lessons from Cuba:
Connecting science and policy. Biological Conservation. 2018 Jan 1;217:280-8.
48. Büdel B, Mies B. Notes on Cape Verde Islands lichen flora. II: Heppiaceae and
Peltulaceae (lichenized Ascomycotina). Nova Hedwigia. 1993;56(3-4):505-14.
49. Sosef MS, Dauby G, Blach-Overgaard A, van der Burgt X, Catarino L, Damen T, et al.
Exploring the floristic diversity of tropical Africa. BMC biology. 2017 Dec;15(1):1-23.
50. Dickman CR. Biodiversity in Australia: an overview. Global biodiversity: volume 4:
selected countries in the Americas and Australia. 2018 Dec 7:515-58.

51. Mayr E, Gilliard ET. Birds of central New Guinea: results of the American Museum of
Natural History expeditions to New Guinea in 1950 and 1952. Bulletin of the AMNH; v.
103, article 4.
52. Webb VL. Photographs of Papua New Guinea: American Expeditions 1928-29. Pacific
Arts. 1995 Jul 1(11/12):72-81.
53. Cookson M. From the archives: the Archbold expeditions to New Guinea: a preliminary
survey of archival materials held at the American Museum of Natural History, New York
City.
54. Tanaka M. The onset and retreat dates of the Austral summer monsoon over Indonesia,
Australia and New Guinea. Journal of the Meteorological Society of Japan. Ser. II. 1994
Apr 25;72(2):255-67.
55. Filippova N, Arefyev S, Zvyagina E, Kapitonov V, Makarova T, Mukhin V, et al. Fungal
literature records database of the Northern West Siberia (Russia). Biodiversity data
journal. 2020;8.
56. Chakrabarty P, Desai UR, Peck C, Mah C, Singer R, Downey R, et al. Preliminary
Checklist of the undersea fauna of the Ross Sea Antarctic Continental Shelf based on Yo-
yo camera surveys. Miscellaneous Publications of the Museum of Zoology, University of
Michigan. 2021, 1-11.
57. Watts J, Vidal J. Environmental defenders being killed in record numbers globally, new
research reveals. Chain Reaction. 2017 Sep(130):40-1.
58. Chapman M, Goldstein BR, Schell CJ, Brashares JS, Carter NH, Ellis-Soto D, et al.
Biodiversity monitoring for a just planetary future. Science. 2024 Jan 5;383(6678):34-6.
59. Ramírez-Castañeda V, Westeen EP, Frederick J, Amini S, Wait DR, Achmadi AS, et al.
A set of principles and practical suggestions for equitable fieldwork in biology.
Proceedings of the National Academy of Sciences. 2022 Aug 23;119(34):e2122667119.
60. Demery AJ, Pipkin M. Safe Fieldwork Strategies for At-Risk Individuals.
61. Paknia O, Rajaei Sh H, Koch A. Lack of well-maintained natural history collections and
taxonomists in megadiverse developing countries hampers global biodiversity
exploration. Organisms Diversity & Evolution. 2015 Sep;15:619-29.
62. Lucia Araujo A. The death of Brazil's national museum. The American Historical
Review. 2019 Apr 1;124(2):569-80.
63. Wickham H, Hester J, Francois R, Bryan J, Bearrows S, Jylänki J, et al. Package 'readr'.
Read Rectangular Text Data. Available online: https://cran. r-project.
org/web/packages/readr/readr. pdf (accessed on 23 August 2023). 2024 Jan 11.
64. Coelho MT, Barreto E, Rangel TF, Diniz-Filho JA, Wüest RO, Bach W, Skeels A,
McFadden IR, Roberts DW, Pellissier L, Zimmermann NE. The geography of climate
and the global patterns of species diversity. Nature. 2023 Oct 19;622(7983):537-44.
625

Acknowledgments

Greg Thom and Nathan Lord helped at various stages of manuscript preparations. We thank the
thousands of natural history researchers, explorers, and collectors who contributed to the samples
we examined here. We also thank the hundreds of museum researchers who prepared, digitized,
and curated these voucher samples. We are also grateful to the databasing experts, particularly
those at the Global Biodiversity Information Facility who have made this type of information
widely accessible.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6