1   Polygenic basis of Bt Cry resistance evolution in wild *Helicoverpa zea*
2
3

4   **Authors:**
5
6   Katherine L Taylor[1], Jane Quackenbush[1], Cara Lamberty[1], Kelly A Hamby[1] & Megan L Fritz[1*]
7
8   [1] Department of Entomology, University of Maryland, College Park MD 20742
9
10   *Corresponding author
11   Email: mfritz13@umd.edu

12
13   **Keywords:**
14   *Helicoverpa zea*, *Bacillus thuringiensi*s, Cry toxin, resistance evolution, gene amplification,
15   structural variant, trypsin, polygenic adaptation
16
17   **Short title:** Polygenic Bt resistance in *Helicoverpa zea*

Abstract

Strong and shifting selective pressures of the anthropocene are rapidly shaping phenomes and genomes of organisms worldwide. One major selective force on insect genomes is crops expressing pesticidal proteins from *Bacillus thuringiensis* (Bt). Here we characterize a rapid response to selection by Bt crops in a major crop pest, *Helicoverpa zea*. We reveal the polygenic architecture of Bt resistance evolution in *H. zea* and identify multiple genomic regions underlying this trait. In the genomic region of largest effect, we identified a gene cluster amplification, where resistant individuals showed variation in copy number for multiple genes. Signals of this amplification increased over time, consistent with the history of field-evolved Bt resistance evolution. Modern wild populations from disparate geographical regions are positive for this variant at high, but not fixed, allele frequencies. We also detected selection against single copy variants at this locus in wild *H. zea* collected from Bt expressing plants, further supporting its role in resistance. Seven trypsin genes were present in this genomic region and all appeared to be significantly upregulated in Bt resistant *H. zea*. Biochemically inhibiting trypsin activity decreased *H. zea*'s tolerance to Bt. These findings characterize rapid genome evolution in a major crop pest following anthropogenic selection and highlight the role that gene copy number variants can have in rapid evolutionary responses.

Introduction

The study of evolution in the Anthropocene has revealed that adaptive organismal responses can occur on timescales much shorter than previously thought possible (Carroll et al. 2007). Human activities impose strong and shifting selection pressure on communities of organisms, shaping their phenomes and genomes (Palumbi, 2001). Empirical evidence of rapid evolutionary change is mounting (*e.g.* Bergland et al., 2014; Bi et al., 2019; Campbell-Staton et al., 2017; Chaturvedi et al., 2021; Ergon et al., 2019; Mikheyev et al., 2015; Roberts Kingman et al., 2021; Rudman et al., 2022; Schiebelhut et al., 2018; Stahlke et al., 2021; Storz & Wheat, 2010), and has linked the strength and nature of selection to specific genomic variants that enhance the fitness of wild organisms. Rapid evolution can result from allele frequency changes at a single major effect locus, or at many loci across the genome (reviewed in Bay et al., 2017). Currently, a major objective of the field of evolutionary genomics is to move beyond documenting the phenomenon of rapid evolution, and instead, uncover the rules that govern how these responses occur on short timescales. This includes characterizing the types of genomic variants (*i.e.* single nucleotide polymorphisms, insertion/deletion polymorphisms, copy number variants, epigenetic modifications) that are most critical for and facilitate rapid evolutionary responses. Perhaps even more important is understanding how fitness-conferring variants came to be; whether they arose *de novo*, were introduced through migration, or were selected from standing genetic variation.

Agricultural ecosystems (agroecosystems) are useful for investigating mechanisms of rapid evolution, both due to the well-understood nature of selection in these ecosystems and their relatively simplified ecosystem structure (Chen & Schoville, 2018). One common feature of agroecosystems is the use of population suppression practices, which exert strong pressure on so-called agricultural pests to evade management through resistance evolution. High levels of

resistance can evolve rapidly (Brevik et al., 2018), and in arthropods alone, resistance to over

300 pesticidal compounds in more than 600 species has been documented (Fritz, 2022; Mota-

Sanchez & Wise, 2022). Rapid pesticide resistance evolution also has economic and

environmental consequences, which provides incentive for its study. Resistance evolution

causes billions of dollars in crop losses every year, and often results in increased environmental

pesticide inputs in an attempt to prevent these losses (Gould et al., 2018; Palumbi, 2001).

We have adopted the Bt cropping system as an experimental model for the study of rapid

organismal evolution. Since 1996, Bt crops have provided area-wide pest suppression, while

decreasing reliance on harmful broad-spectrum pesticides (Cattaneo et al., 2006; Dively et al.,

2018; Hutchison et al., 2010; Kathage & Qaim, 2012; Perry et al., 2016). Bt crops have been

engineered to express genes from the bacterium, *Bacillus thuringiensis*, which encode crystalline

(Cry) and vegetative insecticidal proteins (Vip). The proteins specifically target lepidopteran

and coleopteran species that damage crops, some of which have been historically challenging to

manage with sprayable, synthetic pesticides. They have been widely adopted for insect

management around the globe, and at present, 85% of corn and 89% of cotton planted in the

United States express Bt traits (USDA-ERS, 2023). Bt crops are, therefore, a major selective

force in agroecosystems (Gassmann & Reisig, 2023; Tabashnik & Carrière, 2017).

To improve the durability of these crops and slow resistance evolution, management

plans strongly rooted in evolutionary theory, modeling-based studies, and past empirical

observations were developed (*e.g.* Alstad & Andow, 1995; Frutos et al., 1999; Gould et al., 2018;

McGaughey et al., 1998). Bt crops were designed to produce a "high dose", or 25 times the

amount of pesticidal protein needed to cause mortality in susceptible target pests (US EPA-SRP,

1998). Such a high dose was predicted to favor resistance arising from a single mutation of

major effect, rather than from standing genetic variation across the genome (McKenzie &

Batterham, 1994). Multiple toxins were also pyramided into Bt crops to redundantly kill pests

that evolved resistance to a single protein. Finally, non-expressing plants, or a "refuge" planted

nearby, should produce homozygous susceptible pests that dilute resistance alleles in a local

landscape (Gould, 1998; Huang et al., 2011). Under these circumstances, if resistance was

recessive and resistance alleles incurred a fitness cost in the absence of these toxic proteins, the

emergence and spread of resistance may be preventable (Gould, 1998). Although there have

been cases of successful resistance management in some target pests of Bt crops, practical

resistance has emerged in several lepidopteran and coleopteran species, including the

polyphagous pest, *Helicoverpa zea* (Tabashnik & Carrière, 2017).

H. zea, a pest of corn and cotton, was successfully managed by Bt crops when they were

commercially released in 1996. Yet within 20 years, susceptibility to Cry proteins in *H. zea* had

decreased and crop damage increased (Dively et al., 2016; Dively et al., 2023; Kaur et al., 2019;

Yang et al., 2019; Yang et al., 2022). Resistance evolution in *H. zea* had been predicted because

Cry expressing Bt crops do not produce a "high" dose for *H. zea* (Horner et al., 2003), and refuge

implementation for corn and cotton was often insufficient (Reisig, 2017; Reisig & Kurtz, 2018).

Resistance may also have been facilitated by cross pollination between Bt and refuge corn,

which produces kernels expressing low Cry toxin doses (Pezzini et al., 2024; Dively et al., 2020;

Yang et al., 2015). The Cry resistance that emerged in wild *H. zea* has provided an opportunity

to empirically examine the mechanisms underlying rapid adaptation in this human-altered

system. If *H. zea*'s Cry toxin exposure was not high dose and the strength of selection allowed

for survivors within their natural viability distribution, adaptive responses should have a

polygenic trait architecture (McKenzie & Batterham, 1994). Our previous work supported this

prediction, using two single-family QTL analyses to characterize the molecular architecture of

resistance to two separate Bt crop cultivars (Taylor et al., 2021).

Cry toxicity is generally understood to begin with solubilization and activation of the Cry

protoxin by proteolytic enzymes in the insect digestive system, although this activation step is

not necessary for the activated toxins expressed by many Bt crops (Clark et al., 2005; Gould,

1998; Székács et al., 2010). Activated toxins interact with a series of midgut receptors,

eventually causing lysis of the midgut epithelial cells and leading to growth suppression or

mortality in targeted insects (Heckel, 2020; Jurat-Fuentes et al., 2021; L. Liu et al., 2021). The

multi-step and complex molecular mechanism of Cry toxicity provides many molecular and

physiological pathways for the evolution of resistance. In other lepidopteran species, resistance

has generally been connected to the disruption of the midgut toxin binding sites or reduced

expression of target genes, although there are also examples of resistance associated with altered

Bt protein processing, detoxification, and changes in immune function (Heckel, 2020; Jurat-

Fuentes et al., 2021; L. Liu et al., 2021). Our prior work with *H. zea* led us to reject most of the

common Cry resistance mechanisms described in other lepidopteran species, and instead pointed

to novel resistance-related genomic regions (Taylor et al., 2021), with one under particularly

strong selection by Cry toxins (Pezzini et al. 2024).

Here, we have expanded our analysis of the rapid adaptation to Bt crops observed in wild

H. zea, describing its underlying genetic architecture. We provide further evidence to support a

polygenic trait architecture of Cry resistance in wild *H. zea*. We also provide the first evidence

that a copy number variant (CNV) containing a cluster of 10 genes plays a major role in the Cry

resistance phenotype we observed in *H. zea*. Our results connect the introduction and adoption

of transgenic crops in the landscape, which resulted in a resistance phenotype, to genomic targets

of selection, and the nature of the genomic variants that facilitated *H. zea*'s rapid evolutionary

response.

Results

Cry resistance and general growth phenotypes

To link regions of the *H. zea* genome to Cry resistance phenotypes, we used a replicated

quantitative trait locus (QTL) analysis. This well-established approach relies on crosses between

individuals from phenotypically distinct inbred lines or outbred populations to produce F_2

offspring, whose chromosomes represent unique genomic combinations from cross founders

(Falconer, 1996; Lynch & Walsh, 1998). Correlated traits within phenotypically distinct cross-

founding populations are, therefore, uncorrelated in F_2 progeny due to genome-wide

recombination during meiosis. We crossed field-evolved resistant grandparents to one of very

few known susceptible *H. zea* populations in our replicated design. This allowed us to test for

genome-wide associations between field-relevant Cry resistance phenotypes and genotypes

across recombined F_2 chromosomes. While multiple genomic regions were associated with Cry

resistance in our previous work (Taylor et al. 2021), testing of additional replicate families

increased our power to detect loci of major effect that commonly contribute to field-evolved Cry

resistance in wild *H. zea*.

We initially measured Cry resistance phenotypes for a field resistant population and a

long-term laboratory-reared susceptible population, as well as the F_2 hybrid offspring from 10

intercross families. Resistant individuals should be able to grow while feeding on diets

containing Cry toxins, though at different rates depending on the number and effect size of their

Cry resistance alleles, while susceptible individuals should not. Thus, measuring weight

following 7 days exposure to Cry toxin-incorporated diet served as our metric of resistance. A

critical component of our experimental design included measurement of intra-family controls,

where half of the offspring from each cross were assayed on identical diets lacking Cry toxin.

These intra-family controls served as a way to measure and, if necessary, exclude general

growth-related QTL as candidates for resistance. If a QTL were truly related to Cry resistance,

we reasoned that QTL associated with growth on Cry toxin-incorporated diet in one half of the

family should not overlap with QTL for general growth in their siblings reared on a diet lacking

Cry toxins. Therefore, larvae from each family were exposed to diets containing either a

diagnostic dose of Cry1Ab or Cry1A.105 + Cry2Ab2 expressing corn leaf tissue and leaf tissue

from the non-expressing corn near isolines, as described in Dively et al. (2016) and Taylor et al.

(2021) (**Figure 1, Figure S1, Table S1, Table S2)**.

The larval population from which resistant cross-founding grandparents were drawn

grew significantly larger than laboratory-reared, susceptible larvae on both Cry1Ab and

Cry1A.105 + Cry2Ab2 expressing corn leaf tissue incorporated diets **(Figure 1, Table S1)**. Yet

even in the absence of Cry toxins, the susceptible population grew significantly less than did the

resistant population, suggesting that growth on toxin and general growth were correlated in the

grandparental populations. This was likely a result of long-term reproductive isolation between

these populations resulting in substantial phenotypic and genomic divergence between them.

Growth phenotypes in the F_2 offspring suggested that resistance to all the Cry toxins we tested,

as well as growth on the control treatments were quantitative genetic traits (**SI Results**). Based

on our experimental design, we predicted that recombination in the F1 generation should reduce

correlations between growth on control and toxin-incorporated diet within individual F_2

offspring, resulting in limited overlap between general growth and Cry resistance QTLs

discovered in our assays.

Marker generation and variant calling

Grandparents, parents, and progeny from 10 replicated intercross families were

sequenced on an Illumina NovaSeq 6000. Whole genome sequencing (WGS) of the 20 cross

parents resulted in a total of 4,716,009 high quality genome wide single nucleotide

polymorphisms (SNPs) following read filter-trimming and alignment to the *H. zea* genome (v.

1.0, PRJNA767434; Benowitz et al., 2022). Genome wide average sequencing coverage of

18.6$\times$ (*st. dev.* = 2.4) ensured accuracy of the genotyping calls. Divergence between the field

collected resistant and laboratory susceptible founding populations was high genome wide

(**Figures S2 & 3, SI Results**), as would be expected due to the long-term sexual isolation

between them.

Double digest restriction site associated DNA (ddRAD) sequencing of F_2 offspring

resulted in 78,580 - 79,408 high quality SNP markers for genotype-phenotype association in

each treatment. Final SNP marker numbers likely varied because of true differences in SNP

presence, different loci passing quality control filters, and genetic variation among individuals at

restriction cut sites used for marker development (Davey et al., 2013). The average sequencing

coverage per ddRAD locus across all F_2s was 52,225$\times$ (*st. dev.* = 45,011), and the average

coverage per locus per individual was 63$\times$. SNPs were further filtered to include only those

variants where the allele origin population (field resistant vs. lab susceptible) could be reliably

predicted to determine their directional effect on growth. This produced the smaller filtered sets

of 6,717 - 6,749 genome wide SNPs for each treatment that were used for visualization of effect

size and direction.

Polygenic Cry resistance architecture

To link ddRAD marker genotypes with Cry resistance phenotypes, we used gemma

(Zhou et al., 2013). This software estimates both the number of loci underlying complex traits

(including those of small effect size), as well as the impact of each large effect locus on a trait,

while accounting for the relatedness among individuals in a test population. A significant

proportion of phenotypic variation (PVE) in the F_2 offspring could be explained by the full set of

ddRAD markers. Mean PVE estimates were 69.7% [95% CI = 45.4 - 93.7] for the Cry1Ab

treatment and 71.1% [95% CI = 43.6 - 96.5] for the Cry1A.105 + Cry2Ab2 toxin blend, and

lower for the control growth treatments 44.7% [95% CI = 20.3 - 71.9] and 51.2% [95% CI = 27.1

- 74.1] (**Figure S4**).

The numbers of variants predicted to have large effect sizes on growth were similar

across treatments, with mean estimates ranging from 11.3 - 34.1, further supporting a polygenic

trait architecture. In all cases, there were wide overlapping credible intervals around these

means, suggesting that tens of genomic variants of major effect underlie phenotypic differences

in all growth-related traits (**Figure S4**). Large effect size variants contributed to 74.0% [95% CI

= 36.9 - 98.6] of the Cry1Ab growth-related variance explained by our full marker panel (PGE).

For the Cry1A.105 + Cry2Ab2 treatment, 77.8% [95% CI = 46.8 - 98.9] of the growth-related

variance could be explained by variants of large effect size. These values dropped to 53.4%

[95% CI = 0.5 - 97.3%] and 73.6% [95% CI = 24 - 99.2] for the control treatments (**Figure S4**).

Notably, the high PVE and PGE estimates for both Cry treatments suggest that our data captured

much of the genetic basis of field evolved resistance in *H. zea*.

Genomic regions underlying Bt Cry resistance

To identify the chromosomal regions underlying field evolved Cry resistance, we

estimated the additive effect (β) of a single resistant cross-founder allele on 7 day weight in mg.

Smoothed additive effects of genome-wide markers on growth are shown for Cry1Ab,

Cry1A.105 + Cry2Ab2, and control treatments in **Figure 2**. The major effect QTL revealed in

Figure 2 were also identified using multiple SNP filtering criteria and in the unsmoothed data set

(**Figures S5 & 6**), indicating that our detection of major effect QTL is robust to differences in

bioinformatic and analytical approaches. A resistant parent allele in F_2 offspring significantly

and strongly increased growth on Cry1Ab in regions of Chromosomes (Chr) 2, 3, 6, 9, and 30

with a significance threshold of Bonferroni corrected $p < 0.01$ (**Figure 2A**). With the less

stringent criteria of Bonferroni corrected $p < 0.05$ we also detected an association of parts of Chr

11 and 21 (**Figure 2A**). For Cry1A.105 + Cry2Ab2, we only detected a significant effect of Chr

9, though non-significant peaks suggest possible shared QTL with the Cry1Ab treatment on Chr

2 and 30 (**Figure 2C**). We detected genomic regions associated with general growth in the

laboratory assays on Chr 10, 14 and 27 (**Figure 2B**). As predicted for F_2 offspring, we observed

little overlap between QTL underlying growth on toxin-containing and control treatments. Chr

10 was the only Chr with a major effect on growth on both control and toxin-containing

(Cry1Ab) treatments (**Figures 2A & B**). Therefore, we did not consider this region to be Cry

resistance related. The negative direction of this effect also suggested that the growth associated

allele on Chr 10 is more common in the susceptible parent population. Notably, we did not

detect any fitness cost of Cry resistance-associated genomic regions, which would have appeared

as negative values for these variants in the subsets of F_2s on non-Cry expressing treatments

(**Figures 2B & D**).

The region of the genome with the largest additive effect on resistance was on Chr 9 for

both Cry-containing treatments **(Figures 2A & C)**. There was no signal of an effect of Chr 9 on

general growth, however **(Figures 2B & D)**. For individuals grown on the Cry1Ab treatment, we

identified a clear peak between 5 and 6 Mb on Chr 9. This aligned well with previously

identified signals of genomic divergence over time at 5.75 Mb on Chr 9 in wild *H. zea* (Taylor et

al. 2021). We also identified a directly overlapping divergence peak at 5.75 Mb between

resistant and susceptible cross founders (SI Results). Between 5 and 6 Mb on Chr 9, there were

45 genome annotations (**Table S3**), several of which we examined further (see below). The QTL

on Chr 30 had clear peaks at ~2.5 and ~3.3 Mb, making it possible to describe nearby gene

candidates for the Cry resistance observed in our study. Within 100 Kb of those Chr 30

association peaks (2.4 - 3.4 Mb) were a MAP kinase-activated protein kinase 2-like,

HzeaTryp129, and HzeaABCC11, toll-like receptors, a cluster of carboxylesterases, and

carboxylase-like genes (**Table S4**), suggesting that several known Bt candidate gene families and

insecticide resistance related genes could underlie this QTL. The other identified QTL on Chr 2,

3 and 6 had more diffuse genomic signals with less well-defined peaks, making it difficult to

identify narrow regions most linked to resistance. In the SI Results, we further describe the

limited evidence for any role of other major candidate genes.

Resistance-associated differential gene expression

We paired our QTL study with differential gene expression analyses to narrow in on

potential targets of selection. The mechanism of Cry toxin action takes place in the larval

midgut, and changes in gene expression in this tissue have strong potential to impact resistance.

Therefore, we initially focused on differential midgut gene expression between wild, resistant *H.*

zea larvae collected directly from Cry1A.105+Cry2Ab2-expressing corn and laboratory-reared,

susceptible larvae. Of the 14,600 annotated genes, 527 genes were significantly upregulated and

352 were significantly downregulated in the population collected from Cry expressing corn

relative to the susceptible population (p-adjusted < 0.01; **Table S5**). Of those significantly

differentially expressed genes, 33 were within 100 kb of at least one SNP significantly associated

with resistance to Cry1Ab, Cry1A.105 + Cry2Ab2, or both treatments (**Table S6**). This analysis

identified several significantly upregulated genes in resistant larvae that were found near QTL,

signaling their potential involvement in resistance. One was an aminopeptidase gene found on

Chr 9 (*apn1*) (**Table S6**). *apn1* is a known Cry resistance candidate gene, for which reduction of

expression increases Cry resistance (Herrero et al., 2005; X. Ma et al., 2022; Sun et al., 2022). In

our work, however, *apn1* expression increased in our resistant population, which was

inconsistent with its previously described role in resistance. The most striking gene expression

differences were found among seven significantly upregulated trypsin genes arranged in tandem

between 5 and 6 Mb in the Chr 9 QTL. Six of these trypsins in the tandem array on Chr 9 were

among the top 50 most differentially expressed genes genome wide **(Figure 3B, Table S5)**.

Trans-acting factors also impact regulation of gene expression, which would result in

separation of differentially expressed genes from QTL. When we considered only our

differential gene expression analysis, six other trypsins spread across multiple chromosomes

were also among the top 50 most differentially expressed genes (**Table S5**). Several other

insecticide resistance candidate gene families also had members among the top 50 most

differentially expressed genes, including, a cadherin-87A-like gene and a cytochrome P450

(CYP301B1), both of which were downregulated in resistant individuals (**Table S5**).

We also tested for Cry inducible gene expression changes between wild *H. zea* larvae

collected from Cry expressing and non-expressing corn. A comparison of midgut gene

expression for these groups revealed that expression of a small number of genes was modulated

by Cry toxin exposure. Thirteen genes were significantly upregulated and 20 genes significantly

downregulated in larvae collected from expressing corn (**Table S7**). Four trypsins and one

chymotrypsin were among the most upregulated genes in Cry exposed resistant individuals. Only

one of the trypsins found between 5 and 6 Mb on Chr 9, *tryp80,* was in this group showing Cry

inducible expression changes. Downregulated genes in Cry exposed individuals included a

cadherin-87A-like gene and the immune-related gene, Lysozyme1.

A potential genetic mechanism of large effect on Bt Cry resistance

Taken together, evidence from the QTL and differential expression analysis suggested

that a region of Chr 9 containing a cluster of differentially expressed genes, including 7 trypsins,

was strongly associated with Cry resistance **(Figures 3B & C)**. Closer investigation of this

region revealed a structural variant from 5.22 - 5.37 Mb on Chr 9 **(Figure 4A)**. In this region

encompassing the full cluster of differentially expressed genes, resistant cross founders had

increased whole genome sequencing depth of coverage relative to their genome wide average

(Figure 4A). Elevated coverage depth indicated this region of the genome was at least

duplicated in resistant cross founders. The observed upregulation of most genes in this gene

cluster in resistant individuals was consistent with a mutational event that duplicated the entire

gene cluster **(Figure 3B)**.

To confirm that the coverage and differential expression signals we detected were truly

related to gene cluster duplication, we quantified the copy number variation for one

representative gene in the cluster (*tryp77*) using droplet digital polymerase chain reaction

(ddPCR) **(Figure 4B)**. If *tryp77* existed as a single copy gene, our ddPCR results should reveal a

copy number equal to two in this diploid species. Instead, our analysis showed that all resistant

cross founders from our study had more than two copies of *tryp77*, while all susceptible founders

had the expected two copies. Variation in copy number existed among the field-collected

resistant individuals used in our analysis, indicating there is not a single allele, but multiple

alleles linked to resistance at this Chr 9 locus. There was a strong relationship between copy

number detected by ddPCR and relative coverage depth of the gene cluster ($r^2 = 0.95$), suggesting

that WGS coverage depth is a reasonable proxy measure for copy number.

We used publicly available WGS data from Taylor et al. (2021) and Pezzini et al. (2024)

to determine whether variation in copy number at this Chr 9 locus could also be found in other

wild North American *H. zea* populations. Increased depth of coverage was not found in WGS

data from the susceptible lab population or in field samples collected in 2002, decades before

Cry resistance was widespread **(Figures 4A-D)**. Absence of this variant in samples collected in

2002 suggested that: 1) it may have arisen after 2002, 2) it existed as a rare allele at our study

site prior to 2002, or 3) it arose in or before 2002, but at a location not sampled for our work.

Analysis of WGS data from these *H. zea* samples collected from Louisiana in 2017

(Figures 4C & D), and North Carolina in 2019 **(Figures 4E & F)** revealed signals of increased

coverage depth in this Chr 9 region, similar to what we observed in the resistant cross founding

individuals collected from Maryland in 2019 and 2020 **(Figure 4A)**. The presence of this

sequence duplication in samples collected across 3 separate regions of North America over

multiple years, suggests that this genetic variant is widespread. Notably, depth of coverage at

this locus increased over time, with increased coverage depth first appearing in 2012 and the

strength of that signal increasing in 2017 **(Figures 4C & D)**.

In 109 samples collected from experimental plots in North Carolina in 2019, we detected

increased coverage depth in this Cry resistance-associated Chr 9 region in individuals collected

from Cry expressing plants compared to individuals collected from non-Bt plants in structured

and blended refuge plots (Structured refuge: t = 2.61, adjusted p = 0.039, Blended refuge: t =

2.67, adjusted p = 0.033) **(Figure 4E & F)**. The consistent difference in coverage depth between

the population collected on Cry expressing plants and both refuge conditions and suggests that

selection by the Cry expressing plants limited the growth and survival of individuals with lower

sequence duplication in this region, providing another line of evidence for the role of this genetic

variant in Cry resistance evolution. In these samples, we also observed a continued increase in

sequencing depth over time, with samples from 2019 and 2020 (**Figure 4A & E**) showing

stronger signals of this amplification than those from 2012 and 2017 (**Figure 4C**). This could

reflect increased numbers of individuals homozygous for a duplication or additional duplication

events. Though none of the resistant samples collected in MD and NC in 2019 and 2020 appear

to be single copy in this region (**Figure 4A & E & F**), some individuals collected from non-

expressing plants are (**Figure 4E & F**), indicating that this variant is likely at high frequency but

not fixed in modern populations. Samples collected in 2019 and 2020 also have markedly higher

coverage between 5.32 and 5.35 Mb, region contains *tryp79* and *tryp80,* relative to other parts of

the duplicated cluster and historical samples. We speculate that this particular region may have

undergone multiple duplication events.

The strong association of *tryp77* copy number and depth of WGS coverage motivated us

to analyze the correlation between ddRAD sequencing coverage depth and weight for the F_2

offspring from our experimental crosses **(Figure 4G)**. We reasoned that F_2 individuals with

higher coverage at this locus should, on average, have higher weights than those with low

coverage, if copy number variation was involved in Cry resistance. Likewise, if increasing copy

number is only related to Cry resistance and not generally associated with weight gain, we

reasoned that any positive correlation between weight and coverage depth should be observed for

individuals fed on Cry-treated diet, but not for their siblings on untreated diet. Indeed, ddRAD-

seq coverage depth of the Chr 9 gene cluster was significantly higher for the F_2 offspring that

grew most on both toxin containing diets compared to the offspring which grew the least

(Cry1A.105 + Cry2Ab2: t = 5.2054, p adjusted < 0.001, Cry1Ab: t = 3.4574, p adjusted <0.001)

(Figure 4G). On both control diets, there was no relationship between relative coverage and

growth (Control 1: t = -1.5115, p = 0.1323, Control 2: t = 0.46397, p = 0.6432) **(Figure 4G)**. As

this is a polygenic trait and the Chr 9 locus was not the only one to confer Cry resistance **(Figure**

2A), we expected that F_2 genotype at this locus would not perfectly predict phenotypes.

However, the strong association of ddRAD-seq coverage with resistance phenotype indicates that

the region containing this amplified gene cluster likely explains most of the association between

Chr 9 and Cry resistance.

Synergistic effect of trypsin inhibition and a Cry toxin

Ten genes were found in the region of copy number variation on Chr 9, and seven of

these genes encoded trypsins. To assess whether trypsins, including those on Chr 9, were

involved in Cry resistance, we inhibited their activity with N-α-tosyl-L-lysine chloromethyl

ketone (TLCK) in resistant and susceptible populations of *H. zea*. If trypsin activity, including

the activity of those on Chr 9, were involved in Cry resistance, we reasoned that their inhibition

should interfere with growth on Cry treated diet. Trypsins are generally involved in lepidopteran

metabolism (Muhlia-Almazán et al., 2008), and their inhibition could also lead to stunted growth

on untreated diet. Yet if the midgut expressed trypsins on Chr 9 had activity unique to Cry

resistance, we reasoned that increasing dosages of a trypsin inhibitor should more strongly

impact the weights of larvae feeding on Cry-treated diet, compared to those fed trypsin inhibitor

on an untreated control diet. This trend should be true for both susceptible individuals bearing

single copy genes in the cluster, as well as resistant individuals with copy number variation. We

solubilized TLCK in a phosphate saline buffer (PBS) to make different concentrations of trypsin

inhibitor, and each concentration was mixed with our leaf tissue-incorporated diets. Early

second instar larvae were grown on either a diagnostic diet of Cry1Ab expressing corn leaf tissue

or leaf tissue from the non-expressing near isoline, each with increasing dosages of TLCK.

Larval weight was examined after a 7 day diet exposure. We confirmed that incorporation of

PBS did not impact larval growth relative to our standard non-expressing leaf tissue diet ($p >$

0.05, data not shown), and therefore larval growth on buffer alone was used as a control for all

further analyses and visualization.

408   To understand the extent to which trypsin inhibition influenced Cry resistance, we

calculated a growth ratio, which compared growth on the Cry-treated and untreated diets at each

TLCK dose. If trypsins, including those in our Chr 9 cluster, showed no special activity related

to Cry resistance, we reasoned that the ratio of larval weights on these diet treatments should

remain constant with increasing doses of a trypsin inhibitor. Instead the growth ratio was

negatively correlated with TLCK dose, suggesting that increased suppression of trypsin activity

had a proportionally greater impact on the weights of larvae grown on Cry-treated diets, than

those on untreated diet **(Figure 5)**. The interaction between TLCK dose and Cry1Ab exposure

on larval weight was statistically significant for both resistant and susceptible *H. zea* populations

($p < 0.001$) **(Figure 5, Table S8)**, and post hoc contrasts of the estimated marginal means

confirmed that a trypsin inhibitor paired with Cry1Ab toxin significantly decreased larval growth

(Table S9). Notably, the mean weight of larvae from the resistant population fed on the Cry1Ab

and 10X TLCK treatment was 17.4 mg, suggesting that strongly inhibiting trypsins in resistant

individuals could reduce their Cry tolerance to levels observed in the susceptible population

(mean = 20.7 mg; **Table S8**). These results failed to falsify a role for trypsins in Cry resistance,

including those from our Chr 9 copy number variant (CNV). Overall, our findings suggest that

trypsin activity serves as one protective mechanism against Cry1Ab toxin expressed in corn.

Discussion

Here, we confirm the polygenic nature of *H. zea*'s adaptation to transgenic crops in

agroecosystems, and we begin to shed light on the mechanisms underlying their rapid

evolutionary response. Using replicated, split-family QTL analyses, we identified multiple

genomic regions associated with Bt resistance, most of which are not shared by siblings that

grew well on untreated diets (**Figure 2**). This lack of overlap demonstrates that the multiple

genomic regions associated with growth on Cry-containing diets are Cry resistance loci.

Polygenic adaptation to Cry toxins was predicted for *H. zea* because selection by Bt crops was

expected to act within its broad phenotypic range of Cry tolerance values. Early studies of field-

collected *H. zea* described this range and documented that laboratory selection could rapidly

increase population-level Cry tolerance (Luttrell et al., 1999). Our data provide empirical support

for the prediction that polygenic adaptation is favored when selection acts on phenotypes within

the natural range of variation of a population (McKenzie and Batterham 1994).

Tens of loci contribute to the polygenic basis of Cry resistance in *H. zea*, but the genomic

region of largest effect on both Cry1Ab and Cry1A.105 + Cry2Ab2 resistance was found on Chr

9. All resistance associated genomic regions are likely to be important to the Cry tolerance

observed in wild *H. zea*, and other loci should be the focus of future investigations. It is likely

that genetic interactions among the Cry resistance loci revealed in our work are critical for

production of strong Cry resistance phenotypes, as is observed in other species (Ma et al. 2022,

Sun et al. 2022). However, much of our analyses focused on Chr 9, providing us with insight

into the nature of the genetic mutations that facilitated rapid evolution in this system. We used

multiple lines of evidence to document the nature of the Chr 9 locus and demonstrate its

importance to Cry1Ab resistance both in our lab-based QTL experiments, as well as in field-

collected populations.

For our first line of evidence, we showed that field-collected Cry resistant *H. zea*

overexpressed multiple genes in a 10 gene cluster situated within the Chr 9 genotype-phenotype

association peak (**Figure 3**). Changes in gene expression are well known to underlie adaptive

phenotypes, including for cases of rapid insecticide resistance evolution (Amezian et al., 2021;

Guo et al., 2021; Nauen et al., 2022; Wilding, 2018). An alternative explanation for increased

expression of an entire gene cluster is sequence duplication, however (Heckel, 2022;

Kondrashov, 2012). We used ddPCR of our cross-founding parents to demonstrate that field-

evolved Cry resistant *H. zea* currently carry multiple copies of at least one representative gene

(*tryp77*) in the cluster. From this, we concluded that a CNV found in the 5-6 Mb region of Chr 9

was strongly (but not exclusively) associated with field-evolved Cry resistance in *H. zea*.   It was

not possible to reconstruct the full sequence of this large CNV with the short read sequencing

approaches used in this study, and future studies should employ long read sequencing to describe

haplotypic variation in this genomic region.

Our second line of evidence for the role of this region in Cry resistance was the change in

copy number variation on Chr 9 over time in wild *H. zea*. Changes in Cry1Ab and

Cry1A.105+Cry2Ab2 tolerance in wild *H. zea* slowly increased following commercialization of

Bt crops in 1996 (Dively et al. 2016). Early warning signs of resistance emerged in 2008

(Brévault et al., 2013) and widespread practical resistance to both Cry1A and Cry2A toxins was

documented by 2016 (Dively et al., 2016; Reisig et al., 2018; US-EPA, 2018; Yang et al., 2019).

Using WGS read coverage depth as a proxy for copy number, we documented an increase in

copy number at this locus in wild *H. zea* over time. Consistent with previous resistance

observations, *H. zea* collected from LA in 2002 showed no evidence of a CNV, those from 2012

showed evidence of a rare CNV, and most 2017 individuals carried the variant. This indicated

that the Chr 9 CNV was widespread by the time *H. zea* could readily feed on Cry-expressing

corn ears in the field.

474   Our recent collections of the cross-founding parents from MD in 2019 and 2020 showed

that the Chr 9 CNV was not fixed for a specific copy number. While all resistant cross founders

carried a Chr 9 CNV, some carried different numbers of copies **(Figure 4B)**, which likely confer

different levels of fitness under selection pressure by Cry toxins. This idea motivated the

analysis which provides a third line of evidence for the involvement of the Chr 9 CNV in *H.*

zea's rapid adaptation to Bt crops. A re-analysis of WGS data from wild *H. zea* collected as part

of a field experiment in NC in 2019-2020 (Pezzini et al. 2024) documented that the Chr 9 locus

continues to be under strong selection by Cry toxins. Larvae collected directly from

Cry1Ab+Cry1F expressing field corn showed significantly greater WGS coverage depth of this

loci, our proxy for copy number variation, than did individuals collected from structured or

blended refuge ears **(Figures 4E & F)**. We previously showed that SNPs in this CNV-

containing region were under strong Cry selection (Pezzini et al. 2024), but here we

demonstrated that a generation of selection by Cry1Ab+Cry1F increased the copy number in

wild *H. zea*. Moreover, coverage signals suggested that no individuals collected from

Cry1Ab+Cry1F expressing corn had single copy genotypes, while their conspecifics collected

from non-expressing corn in the same small experimental plots did. These data suggest that the

CNV on Chr 9 continues to be under selection for increased copy number, even though copy

number variation is widespread in the North American landscape.

Interestingly, seven of the ten genes within the CNV were trypsins, all of which were

differentially expressed **(Figure 3B)**. The highest WGS coverage depth in wild *H. zea* collected

on Cry1Ab+Cry1F expressing corn in NC also spanned two of those trypsins. Moreover,

expression of one of the trypsin genes within this cluster, *tryp80*, appeared to be inducible upon

exposure to Cry-expressing corn **(Table S7)**. Trypsins are a gene family previously thought to

be involved in lepidopteran Cry resistance via toxin activation or degradation (Gong et al., 2020;

González-Cabrera et al., 2013; C. Liu et al., 2014; Y. Ma et al., 2013).   We demonstrated that

trypsin inhibition synergistically reduced growth in Cry containing treatments for both resistant

and susceptible populations. These findings demonstrated a significant protective effect of

trypsin activity in Cry exposed *H. zea*. When exposed to the highest TLCK dose and Cry toxin

together, growth in the resistant population was suppressed to the level of the susceptible

population on Cry toxin alone. The shared response of susceptible and resistant populations to

TLCK in this assay suggests that the increased expression and duplication of trypsins might

allow the resistant population to better utilize a mechanism already present in the susceptible

population.

Previous work has also linked trypsin activity to Cry resistance in *H. zea*, for example,

Lawrie et al. (2020) found strong signals of trypsin upregulation in resistant *H. zea* while Zhang

et al. (2019) found that trypsin down regulation was related to *H. zea* Cry resistance in a

laboratory selected line. The upregulation of trypsins in resistant individuals observed here

suggests degradation rather than activation as the potential mechanism for these genes'

involvement in resistance. This would be expected in the case of resistance to crops that express

activated toxin. Phenotypic resistance due to Cry toxin degradation by trypsins has been reported

in field populations of *Spodoptera exigua* (Y. Ma et al., 2013). Our findings along with results

from other Lepidoptera suggest a potential resistance mechanism: gene amplification causes

increased expression of these trypsins which, in turn, may degrade activated Cry toxins.

The CNV on Chr 9, likely plays an important role in resistance, but we emphasize that it

is not the only region involved in *H. zea*'s rapid adaptation to Bt crops. We also identified

resistance associated genomic regions on Chr 2, 3, 6, and 30. Multiple promising candidate genes

were found in the QTL peak on Chr 30 and near other QTL. There were also multiple candidate

Bt resistance genes among the top differentially expressed genes between resistant and

susceptible *H. zea,* including a cadherin gene. The genomic architecture of Bt resistance likely

involves interactions between some of these candidates of large to moderate effect size, as well

as undetected small effect size loci to produce the resistance observed in wild *H. zea*. Indeed,

our results showed that, on average, large Cry-exposed F_2s had significantly higher depth of

sequencing coverage, and therefore copy number in our Chr 9 CNV, than did smaller Cry-

exposed F_2s (**Figure 4G**). However, some small F_2s contained the Chr 9 CNV, while some large

F_2s did not. As expected based on the multiple QTL regions uncovered in our analysis, this

suggests one or more additional segregating resistance alleles are additively or synergistically

interacting with the Chr 9 CNV to produce resistance phenotypes. Consistent with our findings,

interactions between mutations in multiple genes have been described in laboratory studies of

other lepidopteran species (Ma et al. 2022, Sun et al. 2022). Further experimental work targeting

the Cry resistance associated regions on Chr 2, 3, 6 and 30 will be necessary to identify the

additional mutations underlying field-evolved Cry resistance.

In our non-expressing control growth treatments, we did not detect any fitness costs of

this Cry resistance associated gene amplification, or any of the other detected major effect loci

(**Figure 2**). The lack of fitness cost for resistance, the low dose of the toxins, incomplete refuge

implementation, and cross pollination, likely contributed together to the failure of Cry resistance

management strategies for *H. zea*. A degradation mechanism for toxin resistance could explain

the rapid evolution of resistance to multiple toxins that do not share binding sites. Our findings

link the same gene amplification to resistance to both Cry1Ab and blends of Cry1A.105 +

Cry2Ab2 and Cry1Ab+Cry1F. It may also be associated with resistance to other Cry toxins not

tested here. If, as our results suggest, trypsin activity underlies Bt resistance phenotypes in field

H. zea populations, then targeted synergist trypsin inhibitors could potentially be developed to

restore efficacy of Bt crops in controlling *H. zea* (Correy et al., 2019).

Gene duplications, an important source of genetic variation for phenotypic evolution,

have been linked to many cases of insecticide and herbicide resistance evolution (Bass & Field,

2011; Heckel, 2022). Duplications of detoxification related genes can immediately confer

resistance by increasing the quantity of already functional detoxification enzymes or can

additionally lead to new mechanisms of resistance by neofunctionalization. Multiple cases of

insecticide resistance have been functionally linked to new duplications of cytochrome P450s

and carboxylesterases, two detoxification related gene families (Bass et al., 2013; Cattel et al.,

2021; Mouches et al., 1990; Puinean et al., 2010; Schmidt et al., 2010). Our findings here

suggest a new example of the duplication of a detoxification related gene leading to insecticide

resistance, and the first case of this mechanism linked to Bt resistance. It is possible that gene

duplication may be more likely to underlie rapid phenotypic evolution than other variant types,

but the prevalence of structural variants is not well described as detection is difficult with current

technologies (Ho et al., 2020; Mahmoud et al., 2019). As gene amplifications have frequently

been linked to resistance evolution, genomic monitoring for resistance should be extended to

consider a range of genomic variant types beyond SNPs (Fritz 2022).

Methods

Insect samples and phenotypes

Field samples of Cry resistant *H. zea*, defined as surviving to late instar on a Bt corn plant

expressing Cry1A.105 + Cry2Ab2, were collected in 2019 and 2020 in Prince George's County

(MD) at the University of Maryland CMREC farm in Beltsville. Late instars were collected from

Cry1A.105 + Cry2Ab2 expressing sweet corn and reared in the lab on a 16:8 long day light cycle

at 25°C and 50% room humidity. Newly emerged adults from field collections were single pair

mated to produce a second generation of resistant field derived *H. zea*. Susceptible *H. zea* were

acquired from a population that has been maintained in the laboratory without exposure to Cry

toxins at Benzon Research (Cumberland County, PA). Cry tolerance in both the resistant field

and susceptible lab populations was measured with a diagnostic dose in a corn leaf tissue

incorporation assay as described in Taylor et al. (2021) to confirm resistant and susceptible

status. Briefly, early second instar caterpillars were fed for seven days on a laboratory diet mixed

with powdered lyophilized leaf tissue from Bt expressing corn and their non-Bt expressing near

isolines. Resistance was measured by weight gain after seven days on Bt expressing leaf diets.

Weight gain on the non-expressing near isoline control diets served as a measure of non-

resistance associated growth.

The field derived progeny that grew well in the Cry expressing resistance assay (not in

the bottom quartile for growth) were single pair mated to a susceptible individual from the lab

colony. The specific susceptible individuals used in crosses were not exposed to the Cry toxin

resistance assay due to the significant growth suppression that would be experienced by this

highly susceptible population. Offspring of the single pair matings between the resistant and

susceptible populations were sibling mated to produce an F_2 generation. Cry toxin resistance

phenotypes were assessed in the F_2 progeny also using the leaf tissue incorporation assay, as

described in Taylor et al. (2021). Five F_2 intercross families were assessed for resistance to

Cry1Ab by measuring growth on diet with incorporated powdered Cry1Ab expressing leaf

tissue. The other half of the offspring were assayed for growth related advantages on the diets

with powdered leaf tissue from the non-expressing near isoline of Cry1Ab expressing corn. Five

different intercross families were assessed for resistance to Cry1A.105 + Cry2Ab2 by measuring

growth on diet with leaf tissue expressing both toxins for half of the offspring. The other half of

the offspring were used to assess growth related advantages on diets containing leaf tissue from

the non-expressing near isoline. Between 146 and 282 F_2 offspring were split across treatments

and phenotyped from each family, resulting in 73 to 142 F_2 offspring phenotyped per treatment

from each family. F_2 numbers varied due to differences in larval availability for assays. After

phenotyping, the larvae were grown on a non-expressing lab diet to increase in size before DNA

isolation. All samples were flash frozen and stored at -80°C.

The impact of experimental treatment and population on weight phenotypes after seven

days of growth in the laboratory assays was assessed using a model reduction approach between

nested general linear models all with a gamma distribution, as is appropriate for the continuous

and positive weight phenotypes (Bolker, 2008). The treatments compared were 1) Cry1Ab

expressing corn leaf tissue, 2) control tissue from the non-expressing near isoline of Cry1Ab

corn, 3) Cry1A.105 + Cry2Ab2 expressing corn leaf tissue, and 4) control tissue from the non-

expressing near isoline for Cry1A.105 + Cry2Ab2 corn. The populations compared were 1) the

resistant field derived F_0, 2) susceptible lab colony F_0, and 3) the intercross F_2 offspring. The

models were compared using a likelihood ratio χ^2 test with an *a priori* **α** of 0.01. Post hoc

pairwise contrasts were calculated with the R package emmeans (v. 1.8.6; Lenth et al., 2023)

with a bonferroni correction to p-values.

DNA extraction and sequencing

Individuals in the top and bottom 20% of weight phenotypes from each treatment and

family were selected for DNA extraction and sequencing. When individuals in the top or bottom

20% did not survive to preservation, they were replaced by individuals in the top or bottom 30%.

Genomic DNA was extracted from 833 high and low weight intercross F_2 offspring and all F_1

and F_0 parents with a DNeasy (Qiagen, Hilden, Germany) extraction kit using the modified

mouse tail protocol of Fritz et al. (2020). Larval and pupal tissue from the rear ½ - ⅓ of the

insect was used for extraction, while for adults, ½ of the thorax was used. DdRAD libraries were

prepared for all F_2 samples as described in Fritz et al. (2016, 2018). Briefly, samples were

digested with the restriction enzymes EcoRI and MSPI (New England Biolabs, Ipswich, MA),

and unique 6-mer or 8-mer barcodes were annealed prior to pooling with 38 - 39 other samples.

Pools were size-selected for 450-650 bp fragments using a Pippin Prep (Sage Scientific, Beverly,

MA), then 12 replicated PCR reactions with 14 cycles each were used to amplify DNA and add a

standard Illumina TruSeq index as an identifier to each pool. Pools were sequenced at North

Carolina State Genomic Sciences Laboratory on an S4 flow cell of the Illumina NovaSeq 6000,

resulting in 173.9 - 97.5 million raw 150 bp paired-end reads per sample pool with an average of

128 million reads per pool. Additionally whole genome sequencing was performed for the 20

cross founding parents on part of an S4 flow cell of an Illumina NovaSeq 6000 at the University

of Maryland Baltimore Institute for Genome Science. This sequencing resulted in 30.9 - 48

million raw reads per sample with an average of 39.5 million raw reads.

Bioinformatic analysis

F_2 ddRAD sequencing reads were demultiplexed and quality controlled using the stacks

script process_radtags (v.2.61; Rochette et al., 2019). The reads were only retained if they had

exact matches to the correct barcode sequence, quality scores above 20 in all 15 bp length sliding

windows, and no evidence of adapter contamination allowing up to 2 bp mismatch in adapter

sequence. Quality controlled reads were aligned to a *H. zea* chromosome scale assembly (v. 1.0,

PRJNA767434; Benowitz et al., 2022) with Bowtie (v.2.2.5; Langmead & Salzberg, 2012) using

the "very sensitive" alignment option. Variants were called with bcftools mpileup (v. 1.9;

Danecek et al., 2016) and pruned to include biallelic SNPs with a minor allele frequency > 0.05,

with a quality score > 50, present in $> 50\%$ of samples. The average sequencing coverage per

individual at a called ddRAD locus in the final variant set was 63✕. The total number of ddRAD

sequencing markers remaining for genotype phenotype association ranged from 78,580 - 79,408

(Cry1Ab treatment group = 79,261, Cry1Ab control group = 79,182, Cry1A.105 + Cry2Ab2

treatment group = 79,408, Cry1A.105 + Cry2Ab2 control group = 78,580).

Whole genome sequencing reads were quality controlled with Trimmomatic (v. 0.39;

Bolgar et al 2014), to remove Illumina adapters, low quality regions at the beginning or end of

the read or where the average quality of a four base pair window fell below 30, and any reads

shorter than 36 bp. Trimmed reads were aligned and variants were called as described above for

the ddRADseq reads. In the filtered WGS data, a total of 4,716,009 high quality SNPs were

identified between resistant and susceptible populations, with a genome wide average sequencing

coverage depth of 18.6✕ (*st. dev.* = 2.4). For those analyses of the ddRAD sequencing data that

identified the direction of allele effect, SNPs were further filtered to include only those variants

where the allele origin (resistant or susceptible population) could be predicted with some

certainty. To identify SNPs where the likely origin population of the allele could be predicted,

only those SNPs with an allele count difference > 15 (out of a possible 20 alleles in each

population) between the resistant and susceptible parent populations were used. The population

where the allele was more common was used to identify the allele as resistance or susceptibility

associated. After filtering for effect direction, 6,717 - 6,749 SNPs remained (Cry1Ab treatment

group = 6,737, Cry1Ab control group = 6,749, Cry1A.105 + Cry2Ab2 treatment group = 6,733,

Cry1A.105 + Cry2Ab2 control group = 6,717).

Resistant and susceptible founding populations genome wide comparison

Genome-wide patterns of differentiation between the founding resistant and susceptible

populations were described using the variants identified by whole genome sequencing of the ten

field resistant and ten laboratory susceptible F_0 individuals founding the family crosses. A

principal components analysis was performed with Plink (v. 1.90b; Chang et al., 2015).

Divergence was measured by Weir and Cockerham's windowed F_{ST} across 40kb windows with a

10kb step using VCFtools (v. 0.1.17; Danecek et al., 2011). A genome wide zF_{ST} significance

threshold of 6 was used (Rubin et al., 2010). Downstream analysis of this data and all other data

presented here were performed in R (v.4.2.1; (R Development Core Team, 2023) with tidyverse

R package (v.2.0.0; Wickham et al., 2019), and visualizations were made with the R package

ggplot2 (v.3.4.1; Wickham, 2016)

Quantitative trait mapping for Cry resistance

Cry resistance genomic architecture was characterized by Bayesian sparse linear mixed

models (BSLMM) in gemma (v. 0.98.4; Zhou et al., 2013). For each treatment the model was run

five separate times, each time with a total of 5 million sampling steps with the first 500 thousand

discarded as burn-in. Hyperparameter estimates for each treatment were averaged across all five

runs. The association between SNPs across the genome and weight phenotype from the

resistance and control growth assays were detected with linear mixed models (LMM) also in

gemma (v. 0.98.4; Zhou & Stephens, 2012). Both the BSLMM and LMM account for relatedness

among the five replicated families for each treatment.

Differential gene expression analysis

In 2022, Cry1A.105 + Cry2Ab2 expressing corn ears and ears from the non-Bt

expressing near isoline were collected from the University of Maryland CMREC farm at

Beltsville. Ears that were infested with *H. zea* were identified by the loosened silk and brought to

the laboratory. In the lab, each corn ear was open and any 5th instar larvae present were

identified and removed from the ear; caterpillars at other developmental stages were not included

in the experiment. Immediately following removal, larvae were chilled and immobilized in a

70% solution of ice cold RNAlater (Invitrogen, Waltham, MA) and PBS. The head capsule on

each larva was removed and stored for later measurement to confirm instar. All larvae were later

confirmed as 5th instar by head measurements between 1.60 to 2.70 mm (Bilbo et al., 2019). The

midgut was cut away from the crop and intestine, and the malpighian tubules were removed.

Corn was cleared from midgut using forceps and a rinse of 70% RNAlater. Midguts were flash

frozen and stored at -80°C. The dissection process was repeated with individuals from the

laboratory susceptible colony, which had been reared on a standard laboratory diet not containing

any corn tissue or Bt toxins.

Midguts of 5th instars from Bt expressing corn, non-Bt corn and the susceptible

laboratory colony were organized into eight randomized pools per treatment with four samples in

each pool (n = 32 per treatment). RNA was extracted from each pool using am RNeasy® Mini

Kit (Qiagen, Hilden, Germany) according to the manufacturer protocol except for the initial

pooling of samples. Poly-A selection, library preparation, and sequencing were completed by the

University of Maryland Baltimore Genomics Core Facility. All RNA libraries had a RIN value

between 8.7 and 10, with 22/24 having RIN of 10, indicating very high quality. Libraries were

100 bp paired end sequenced on an Illumina NovaSeq 6000.

Raw reads were quality controlled with Trimmomatic (v. 0.39; Bolger et al., 2014) to

remove Illumina adapters, trim reads where 4 bp sliding window quality score fell below 20, and

drop any reads shorter than 36 bp after trimming. The paired quality controlled reads were

aligned to a chromosome scale *H. zea* assembly (Benowitz et al., 2022; GCA_022343045.1) with

HISAT2 (v. 2.2.1; Kim et al., 2019) using default parameters. Gene annotations were lifted from

the older reference genome Hzea_1.0 (Pearce et al., 2017; GCA_002150865.1) to the new

chromosome scale assembly using Liftoff (v. 1.6.1; Shumate & Salzberg, 2021). Gene

expression counts were generated with HTSeq-count (v. 0.13.5; Anders et al., 2015). Expression

patterns across all 24 pools were compared through differential expression analysis in DESeq2

(v. 1.38.3; Love et al., 2014). Two-way differential expression analysis was conducted between

resistant field individuals collected from Bt expressing corn and susceptible lab colony samples

to identify expression changes associated with resistance. An additional two-way differential

expression analysis between the field collected individuals from Bt expressing corn and non

expressing corn was conducted to identify Cry exposure inducible gene expression changes.

Statistical significance was indicated by a false discovery rate adjusted p-value of 0.01.

Structural variant detection

Sequencing coverage depth signals were extracted from aligned .bam files from the WGS

of the cross founders, the ddRAD sequencing of F_2, and two publicly available data sets. The

genome wide coverage depth and the average coverage depth for the trypsin cluster were

calculated with Qualimap (v. 2.2.1; Okonechnikov et al., 2016), while windowed coverage depth

calculations were from samtools depth (v. 1.10; Danecek et al., 2021) results. Calculations of

mean coverage of the amplified region for these analyses included all reads aligning between

5.22 - 5.37 Mb on Chr 9. Significant differences in coverage depth across samples were

determined by pairwise t-tests with a bonferroni correction to p. The presence of the structural

variant in 2002, 2012, and 2017 was assessed using the publicly available whole genome

sequencing data from Taylor et al. (2021) (PRJNA751583). The presence of the structural

variant in samples collected from Cry expressing and non-expressing corn was assessed using

sequencing data from the Pezzini et al. (2024) (PRJNA1055981).

The structural variant was validated with digital droplet PCR (ddPCR) for *tryp77*, one of

seven trypsins in the putative duplication. ATP dependent DNA helicase was used as a control

gene, as it is single copy in Lepidoptera in OrthoDB (Zdobnov et al., 2021). Twenty-five ng of

DNA per sample was mixed with HindIII and analyzed in multiplexed assay for *tryp77* and ATP

dependent DNA helicase at MOgene (St. Louis, MO). The genes were targeted with the primers

shown in **(Table S10)** and the following amplification conditions: activation at 95℃ for 10

minutes, followed by 40 cycles of denaturing at 94℃ for 30 seconds, annealing and extension at

58℃ for 1 minute, finally the enzyme was deactivated at 98℃ for 10 minutes and held at 4℃.

ddPCR was performed with QX200 Automated Droplet Generator and Reader and analyzed

using the QX Manager 1.2 Standard Edition software (Bio-Rad, Hercules CA). All samples were

run with a synthesized gBlock positive control and a no template negative control. Copy number

for samples was calculated as in Karlin-Neumann et al. (2012) with the following formula:

$$CN = \frac{ln\left(1 - \frac{target\ gene\ positive\ droplets}{total\ number\ of\ droplets}\right)}{ln\left(1 - \frac{control\ gene\ positive\ droplets}{total\ number\ of\ droplets}\right)} * 2$$

741
Field resistant colony collection and rearing

Cry resistant *H. zea* late instar caterpillars were collected from Cry1Ab expressing sweet

corn ('BC0805') ears grown at the Central Maryland Research and Education Center in

Beltsville, Maryland. Approximately 450 caterpillars were collected during each of two

collection dates (August 18 and September 7, 2023). Caterpillars were reared on *H. zea* diet

(Southland Products Inc., Arkansas, USA) until adulthood and bucket mated in a growth

chamber set at 25°C with 70% relative humidity and 16:8 light:dark cycle. Eggs from bucket

matings were placed onto diet and reared until the second instar under the same conditions.

Occasionally, larvae were held at 4°C once they reached the appropriate size to ensure sufficient

numbers at the same developmental stage for assays.

Trypsin inhibition assay

To determine whether trypsin activity impacts Cry toxicity in *H. zea*, we

compared 7-day larval weight after exposure to corn leaf tissue incorporated diet. Larvae were

exposed to one of 12 treatments: 1) 1X TLCK trypsin inhibitor and Cry1Ab expressing leaf

tissue, 2) 2X TLCK trypsin inhibitor and Cry1Ab expressing leaf tissue, 3) 5X TLCK trypsin

inhibitor and Cry1Ab expressing leaf tissue, 4) 10X TLCK trypsin inhibitor and Cry1Ab

expressing leaf tissue, 5) 1X TLCK trypsin inhibitor and non-expressing leaf tissue, 6) 2X

TLCK trypsin inhibitor and non-expressing leaf tissue, 7) 5X TLCK trypsin inhibitor and non-

expressing leaf tissue, 8) 10X TLCK trypsin inhibitor and non-expressing leaf tissue, 9) buffer

and Cry1Ab expressing leaf tissue, 10) buffer and non-expressing leaf tissue, 11) Cry1Ab

expressing leaf tissue alone, 12) non-expressing leaf tissue alone. These treatments enabled us to

separately determine the growth effect of Cry expressing tissue, the buffer, and the inhibitor at

different concentrations. Second instars from the laboratory Cry susceptible *H. zea* population

from Benzon research and field-collected Cry resistant *H. zea* were reared on these treatments

for seven days before they were individually weighed to assess growth. In each assay, 8 - 16

individuals were exposed to each treatment, due to availability of appropriately sized larvae.

Final sample sizes were $\geq$ 70 in each treatment, and are reported in Table S8.

Based upon Ma et al. (2013) and preliminary trials, we selected N-α-tosyl-ʟ-lysine

chloromethyl ketone hydrochloride (TLCK, Sigma-Aldrich®, St. Louis, MO) as our trypsin

inhibitor. To make the inhibitor solutions, 350 mg of TLCK was dissolved in 10 mL of

phosphate buffer solution (0.1M, pH 5.8, bioWORLD, Dublin, OH) to make a stock solution of

35 mg/mL (10X). This solution was further diluted to 3.5 mg/mL (1X), 7 mg/mL (2X), and 17.5

mg/mL (5X) as needed. The inhibitor solutions were freshly made for each assay date. A 10 mL

aliquot of phosphate buffer was used for the buffer control treatment. The bioassays were

prepared as in Dively (2016), Taylor et al. (2021), and as described above, except, in the

inhibitor and buffer treatments 1.6 mL of the TLCK solution or phosphate buffer were mixed

into 25 mL of experimental diet, resulting in the following final TLCK concentrations: 1X: 224.5

μg of TLCK per mL diet, 2X: 449 μg of TLCK per mL diet, 5X: 1.1 mg of TLCK per mL diet,

10X: 2.2 mg of TLCK per mL diet.

First we compared the two control groups (incorporated leaf tissue only and incorporated

leaf tissue with buffer) using t tests with an α of 0.01 to identify any potential effects of the

buffer alone. Then, we tested for a synergistic effect of trypsin inhibition and Cry exposure on

growth in *H. zea* with a general linear model comparison approach. A gamma distribution was

used for the weight response variable. The fit of the models was compared with a likelihood ratio

chi-squared test using an α of 0.01. We tested for combined effects by comparing a full model

with Cry treatment and trypsin inhibition main effects and an interaction between the two, to a

model with only the two main effects. Bonferroni corrected post hoc contrasts were calculated

with the R package emmeans (v. 1.8.6; Lenth et al., 2023). We also calculated a growth ratio at

each TLCK dose as:

791
$$GR = \frac{weight\ on\ Cry\ expressing\ leaf\ tissue\ at\ TLCK\ dose\ x}{weight\ on\ the\ non-expressing\ leaf\ tissue\ at\ TLCK\ dose\ x}$$

792   The growth ratio measured the combined effect of TLCK and Cry toxins, and would be expected

793   to stay consistent across doses if the effects of TLCK and Cry toxins on growth suppression are

794   not linked.

Acknowledgements

We thank Juan Luis Jurat-Fuentes, and David Heckel for suggestions that improved our manuscript. We thank Galen Dively for access to experimental plots for field collections. We thank Katherine Bell, Thea Bliss, Ben Burgunder, Lasair ní Chochlain, Maria Cramer, Kyree Day, Dominique Desmarattes, Shea Ill, Emma Kohanski, Ava Lamberty, Scott McCluen, Declan McHugh, Oliva Moy, Hiral Patel, Daniella Anconeira Sayco, Rachel Sanford, Torsten Schöneberg, Robert Starkenburg, Olivia Rosen, and Joshua Yeroshefsky for assistance with H. zea collections, crosses, rearing, bioassays, and/or DNA isolation. This work was funded by the US Department of Agriculture, National Institute of Food and Agriculture Biotechnology Risk Assessment Grants 2019-33522-29992 and 2022-33522-37744.

Figure Legends

Figure 1. Distribution of weights for *H. zea* individuals after seven days in a laboratory leaf tissue incorporation assay for Cry toxin and control treatments.

Figure 2. Association of genome wide markers and weight after seven days of feeding on Cry toxin containing and control treatments. The additive effect (beta from LMM) is the weight in mg associated with the presence of a single resistant population allele smoothed for visualization by averaging over 21 SNP windows with a 5 SNP step for markers on 31 *H. zea* chromosomes. The genotype-phenotype association is shown for **A.** weight after seven days of exposure to Cry1Ab, **B.** weight on the non-toxin near isoline control diet for Cry1Ab, **C.** weight after seven days of exposure to Cry1A.105 + Cry2Ab2, **D.** weight on the non-toxin near isoline control diet for Cry1A.105 + Cry2Ab2. Chromosomes are plotted in alternating dark and light gray with each point representing a 21 SNP window. Windows with at least one significantly associated SNP are highlighted in color, with orange indicating an adjusted p value < 0.05 and red indicating an adjusted p value < 0.01.

Figure 3. Genomic signals indicate that a region between 5.2 and 5.4 Mb on *H. zea* Chr 9 is duplicated in field resistant individuals and associated with resistance phenotype. The absolute value of the effect size (beta from the LMM) for ddRAD SNPs is plotted in (**A**). A differential expression plot for all genes on Chr 9 between resistant and susceptible populations is shown in (**B**). In (**C**) the gene annotations between 5.2 and 5.4 Mb on Chr 9 are shown above a plot of relative depth of sequencing coverage for field resistant *H. zea*. Gene abbreviations are as follows: Myr1 = myrosinase 1-like, Dlh = disks large homolog 4-like, Tryp = Trypsin, Npr = neuropeptide receptor A35. In all panels the region between 5.2 and 5.4 Mb on Chr 9 is highlighted in blue.

Figure 4. Resistance evolution and the gene cluster amplification on *H. zea* Chr 9. **(A)** Relative sequencing depth is plotted for susceptible and resistant cross founders. In A, C, and E each point represents the average sequencing depth in that 10 kb window for a single individual

relative to that individual's genome wide average sequencing coverage. **(B)** Coverage is
associated with ddPCR copy number variant analysis for *trypsin 77* ($r^2 = 0.95$) for the resistant
(dark gray) and susceptible (light gray) founders of mapping families. **(C)** Relative sequencing
depth is plotted for individuals collected in LA in 2002 before Cry resistance evolution, 2012 as
resistance was evolving, and 2017 after practical resistance was first detected. **(D)** The mean
coverage depth of the amplified region from 5.22 - 5.37 Mb is plotted for the same individuals
shown in C. **(E)** Relative sequencing depth is plotted for individuals collected in 2019 in NC
from structured refuge, blended refuge and Cry1Ab+Cry1F expressing corn. **(F)** The mean
coverage depth of the amplified region is plotted for the same individuals shown in E. Coverage
was significantly higher for the individuals taken from Cry expressing plants compared to both
refuge types ($t = 2.61$, $p = 0.039$; $t = 2.67$, $p = 0.033$), suggesting selection for higher copy
numbers. **(G)** Mean ddRAD sequencing coverage for F_2 offspring that were the top and bottom
for 7 day weight in each treatment. Relative coverage was significantly higher for the F_2
offspring that grew largest on both toxins ($t = 3.46$, $p < 0.001$; $t = 5.21$, $p < 0.001$). On both
control treatments there was no relationship between coverage and growth ($t = 0.46$, $p > 0.05$; $t =$
-1.51, $p > 0.05$).
860
861
862   **Figure 5.** Effect of trypsin inhibition. **(A)** Distribution of weights for laboratory susceptible and
863   field resistant *H. zea* individuals after seven days in a trypsin inhibition assay with Cry1Ab
864   containing corn leaf tissue and leaf tissue from the non-expressing near isoline. **(B)** The growth
865   ratio for each trypsin inhibitor dose, a measure of how much weight is suppressed by Cry toxins
866   and TLCK combined compared to the effects TLCK alone.

References

Alstad, D. N., & Andow, D. A. (1995). Managing the evolution of insect resistance to transgenic plants. *Science*, *268*(5219), 1894–1896. https://doi.org/10.1126/science.268.5219.1894

Amezian, D., Nauen, R., & Le Goff, G. (2021). Transcriptional regulation of xenobiotic detoxification genes in insects—An overview. *Pesticide Biochemistry and Physiology*, *174*, 104822. https://doi.org/10.1016/j.pestbp.2021.104822

Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq—A Python framework to work with high-throughput sequencing data. *Bioinformatics*, *31*(2), 166–169. https://doi.org/10.1093/bioinformatics/btu638

Bass, C., & Field, L. M. (2011). Gene amplification and insecticide resistance. *Pest Management Science*, *67*(8), 886–890. https://doi.org/10.1002/ps.2189

Bass, C., Zimmer, C. T., Riveron, J. M., Wilding, C. S., Wondji, C. S., Kaussmann, M., Field, L. M., Williamson, M. S., & Nauen, R. (2013). Gene amplification and microsatellite polymorphism underlie a recent insect host shift. *Proceedings of the National Academy of Sciences*, *110*(48), 19460–19465. https://doi.org/10.1073/pnas.1314122110

Bay, R. A., Rose, N., Barrett, R., Bernatchez, L., Ghalambor, C. K., Lasky, J. R., Brem, R. B., Palumbi, S. R., & Ralph, P. (2017). Predicting responses to contemporary environmental change using evolutionary response architectures. *The American Naturalist*, *189*(5), 463–473. https://doi.org/10.1086/691233

Benowitz, K. M., Allan, C. W., Degain, B. A., Li, X., Fabrick, J. A., Tabashnik, B. E., Carrière, Y., & Matzkin, L. M. (2022). Novel genetic basis of resistance to Bt toxin Cry1Ac in *Helicoverpa zea*. *Genetics*, *221*(1), iyac037. https://doi.org/10.1093/genetics/iyac037

Bergland, A. O., Behrman, E. L., O'Brien, K. R., Schmidt, P. S., & Petrov, D. A. (2014). Genomic evidence of rapid and stable adaptive oscillations over seasonal time scales in *Drosophila*. *PLOS Genetics*, *10*(11), e1004775. https://doi.org/10.1371/journal.pgen.1004775

Bi, K., Linderoth, T., Singhal, S., Vanderpool, D., Patton, J. L., Nielsen, R., Moritz, C., & Good, J. M. (2019). Temporal genomic contrasts reveal rapid evolutionary responses in an alpine mammal during recent climate change. *PLOS Genetics*, *15*(5), e1008119. https://doi.org/10.1371/journal.pgen.1008119

Bilbo, T. R., Reay-Jones, F. P. F., Reisig, D. D., Greene, J. K., & Turnbull, M. W. (2019). Development, survival, and feeding behavior of *Helicoverpa zea* (Lepidoptera: Noctuidae) relative to Bt protein concentrations in corn ear tissues. *PLoS ONE*, *14*(8), e0221343. https://doi.org/10.1371/journal.pone.0221343

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Bolker, B. M. (2008). Ecological models and data in R. In *Ecological Models and Data in R*.

Princeton University Press. https://doi.org/10.1515/9781400840908

Brévault, T., Heuberger, S., Zhang, M., Ellers-Kirk, C., Ni, X., Masson, L., Li, X., Tabashnik, B.
E., & Carrière, Y. (2013). Potential shortfall of pyramided transgenic cotton for insect
resistance management. *Proceedings of the National Academy of Sciences*, *110*(15),
5806–5811. https://doi.org/10.1073/pnas.1216719110

Brevik, K., Schoville, S. D., Mota-Sanchez, D., & Chen, Y. H. (2018). Pesticide durability and
the evolution of resistance: A novel application of survival analysis. *Pest Management*
Science, *74*(8), 1953–1963. https://doi.org/10.1002/ps.4899

Campbell-Staton, S. C., Cheviron, Z. A., Rochette, N., Catchen, J., Losos, J. B., & Edwards, S.
V. (2017). Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the
green anole lizard. *Science*, *357*(6350), 495–498.
https://doi.org/10.1126/science.aam5512

Cattaneo, M. G., Yafuso, C., Schmidt, C., Huang, C., Rahman, M., Olson, C., Ellers-Kirk, C.,
Orr, B. J., Marsh, S. E., Antilla, L., Dutilleul, P., & Carrière, Y. (2006). Farm-scale
evaluation of the impacts of transgenic cotton on biodiversity, pesticide use, and yield.
Proceedings of the National Academy of Sciences, *103*(20), 7571–7576.
https://doi.org/10.1073/pnas.0508312103

Cattel, J., Haberkorn, C., Laporte, F., Gaude, T., Cumer, T., Renaud, J., Sutherland, I. W., Hertz,
J. C., Bonneville, J.-M., Arnaud, V., Fustec, B., Boyer, S., Marcombe, S., & David, J.-P.
(2021). A genomic amplification affecting a carboxylesterase gene cluster confers
organophosphate resistance in the mosquito *Aedes aegypti*: From genomic
characterization to high-throughput field detection. *Evolutionary Applications*, *14*(4),
1009–1022. https://doi.org/10.1111/eva.13177

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015).
Second-generation PLINK: Rising to the challenge of larger and richer datasets.
GigaScience, *4*(1). https://doi.org/10.1186/s13742-015-0047-8

Chaturvedi, A., Zhou, J., Raeymaekers, J. A. M., Czypionka, T., Orsini, L., Jackson, C. E.,
Spanier, K. I., Shaw, J. R., Colbourne, J. K., & De Meester, L. (2021). Extensive standing
genetic variation from a small number of founders enables rapid adaptation in *Daphnia*.
Nature Communications, *12*(1), 4306. https://doi.org/10.1038/s41467-021-24581-z

Chen, Y., & Schoville, S. (2018). Editorial overview: Ecology: Ecological adaptation in
agroecosystems: novel opportunities to integrate evolutionary biology and agricultural
entomology. *Current Opinion in Insect Science*, *26*, iv–viii.
https://doi.org/10.1016/j.cois.2018.03.003

Clark, B. W., Phillips, T. A., & Coats, J. R. (2005). Environmental fate and effects of *bacillus*
thuringiensis (Bt) proteins from transgenic crops: A review. *Journal of Agricultural and*
Food Chemistry, *53*(12), 4643–4653. https://doi.org/10.1021/jf040442k

Correy, G. J., Zaidman, D., Harmelin, A., Carvalho, S., Mabbitt, P. D., Calaora, V., James, P. J.,
Kotze, A. C., Jackson, C. J., & London, N. (2019). Overcoming insecticide resistance

through computational inhibitor design. *Proceedings of the National Academy of*
Sciences, *116*(42), 21012–21021. https://doi.org/10.1073/pnas.1909130116

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R.
E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000 Genomes
Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*,
27(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

Danecek, P., Schiffels, S., & Durbin, R. (2016). *Multiallelic calling model in bcftools (-m),*
samtools.github.io/bcftools/call-m.pdf [Computer software].

Davey, J. W., Cezard, T., Fuentes-Utrilla, P., Eland, C., Gharbi, K., & Blaxter, M. L. (2013).
Special features of RAD sequencing data: Implications for genotyping. *Molecular*
Ecology, *22*(11), 3151–3164. https://doi.org/10.1111/mec.12084

Dively, G. P., Huang, F., Oyediran, I., Burd, T., & Morsello, S. (2020). Evaluation of gene flow
in structured and seed blend refuge systems of non-Bt and Bt corn. *Journal of Pest*
Science, *93*(1), 439–447. https://doi.org/10.1007/s10340-019-01126-4

Dively, G. P., Venugopal, P. D., Bean, D., Whalen, J., Holmstrom, K., Kuhar, T. P., Doughty, H.
B., Patton, T., Cissel, W., & Hutchison, W. D. (2018). Regional pest suppression
associated with widespread Bt maize adoption benefits vegetable growers. *Proceedings*
of the National Academy of Sciences, *115*(13), 3320–3325.
https://doi.org/10.1073/pnas.1720692115

Dively, G. P., Venugopal, P. D., & Finkenbinder, C. (2016). Field-evolved resistance in corn
earworm to Cry proteins expressed by transgenic sweet corn. *PLOS ONE*, *11*(12),
e0169115. https://doi.org/10.1371/journal.pone.0169115

Ergon, Å., Skøt, L., Sæther, V. E., & Rognli, O. A. (2019). Allele frequency changes provide
evidence for selection and identification of candidate loci for survival in red clover
(*Trifolium pratense L.*). *Frontiers in Plant Science*, *10*.
https://doi.org/10.3389/fpls.2019.00718

Falconer, D. S. (1996). *Introduction to quantitative genetics* (4th ed). Prentice Hall.

Fritz, M. L. (2022). Utility and challenges of using whole-genome resequencing to detect
emerging insect and mite resistance in agroecosystems. *Evolutionary Applications*,
15(10), 1505–1520. https://doi.org/10.1111/eva.13484

Fritz, M. L., DeYonke, A. M., Papanicolaou, A., Micinski, S., Westbrook, J., & Gould, F.
(2018). Contemporary evolution of a Lepidopteran species, *Heliothis virescens*, in
response to modern agricultural practices. *Molecular Ecology*, *27*(1), 167–181.
https://doi.org/10.1111/mec.14430

Fritz, M. L., Paa, S., Baltzegar, J., & Gould, F. (2016). Application of a dense genetic map for
assessment of genomic responses to selection and inbreeding in *Heliothis virescens*.
Insect Molecular Biology, *25*(4), 385–400. https://doi.org/10.1111/imb.12234

Frutos, R., Rang, C., & Royer, M. (1999). Managing insect resistance to plants producing

Bacillus thuringiensis toxins. *Critical Reviews in Biotechnology*, *19*(3), 227–276. https://doi.org/10.1080/0738-859991229251

Gassmann, A. J., & Reisig, D. D. (2023). Management of insect pests with Bt crops in the United States. *Annual Review of Entomology*, *68*(1). https://doi.org/10.1146/annurev-ento-120220-105502

Gong, L., Kang, S., Zhou, J., Sun, D., Guo, L., Qin, J., Zhu, L., Bai, Y., Ye, F., Akami, M., Wu, Q., Wang, S., Xu, B., Yang, Z., Bravo, A., Soberón, M., Guo, Z., Wen, L., & Zhang, Y. (2020). Reduced expression of a novel midgut trypsin gene involved in protoxin activation correlates with Cry1Ac resistance in a laboratory-selected strain of *Plutella xylostella* (L.). *Toxins*, *12*(2), Article 2. https://doi.org/10.3390/toxins12020076

González-Cabrera, J., García, M., Hernández-Crespo, P., Farinós, G. P., Ortego, F., & Castañera, P. (2013). Resistance to Bt maize in *Mythimna unipuncta* (Lepidoptera: Noctuidae) is mediated by alteration in Cry1Ab protein activation. *Insect Biochemistry and Molecular Biology*, *43*(8), 635–643. https://doi.org/10.1016/j.ibmb.2013.04.001

Gould, F. (1998). Sustainability of transgenic insecticidal cultivars: Integrating pest genetics and ecology. *Annual Review of Entomology*, *43*(1), 701–726. https://doi.org/10.1146/annurev.ento.43.1.701

Gould, F., Brown, Z. S., & Kuzma, J. (2018). Wicked evolution: Can we address the sociobiological dilemma of pesticide resistance? *Science*, *360*(6390), 728–732. https://doi.org/10.1126/science.aar3780

Guo, Z., Kang, S., Wu, Q., Wang, S., Crickmore, N., Zhou, X., Bravo, A., Soberón, M., & Zhang, Y. (2021). The regulation landscape of MAPK signaling cascade for thwarting *Bacillus thuringiensis* infection in an insect host. *PLOS Pathogens*, *17*(9), e1009917. https://doi.org/10.1371/journal.ppat.1009917

Heckel, D. G. (2020). How do toxins from *Bacillus thuringiensis* kill insects? An evolutionary perspective. *Archives of Insect Biochemistry and Physiology*, *104*(2), e21673. https://doi.org/10.1002/arch.21673

Heckel, D. G. (2022). Perspectives on gene copy number variation and pesticide resistance. *Pest Management Science*, *78*(1), 12–18. https://doi.org/10.1002/ps.6631

Herrero, S., Gechev, T., Bakker, P. L., Moar, W. J., & de Maagd, R. A. (2005). *Bacillus thuringiensis* Cry1Ca-resistant *Spodoptera exigua* lacks expression of one of four Aminopeptidase N genes. *BMC Genomics*, *6*(1), 96. https://doi.org/10.1186/1471-2164-6-96

Ho, S. S., Urban, A. E., & Mills, R. E. (2020). Structural variation in the sequencing era. *Nature Reviews Genetics*, *21*(3), Article 3. https://doi.org/10.1038/s41576-019-0180-9

Horner, T. A., Dively, G. P., & Herbert, D. A. (2003). Development, survival and fitness performance of *Helicoverpa zea* (Lepidoptera: Noctuidae) in MON810 Bt field corn. *Journal of Economic Entomology*, *96*(3), 914–924. https://doi.org/10.1093/jee/96.3.914

Hutchison, W. D., Burkness, E. C., Mitchell, P. D., Moon, R. D., Leslie, T. W., Fleischer, S. J., Abrahamson, M., Hamilton, K. L., Steffey, K. L., Gray, M. E., Hellmich, R. L., Kaster, L. V., Hunt, T. E., Wright, R. J., Pecinovsky, K., Rabaey, T. L., Flood, B. R., & Raun, E. S. (2010). Areawide suppression of European corn borer with Bt maize reaps savings to non-Bt maize growers. *Science*, *330*(6001), 222–225. https://doi.org/10.1126/science.1190242

Jurat-Fuentes, J. L., Heckel, D. G., & Ferré, J. (2021). Mechanisms of resistance to insecticidal proteins from *Bacillus thuringiensis*. *Annual Review of Entomology*, *66*(1), 121–140. https://doi.org/10.1146/annurev-ento-052620-073348

Karlin-Neumann, G., Montesclaros, L., Heredia, N., Hodges, S., Regan, J., Troup, C., Mukhopadhyay, D., & Litterst, C. (2012). *Probing copy number variations using Bio-Rad's QX100^{TM} Droplet Digital^{TM} PCR system*.

Kathage, J., & Qaim, M. (2012). Economic impacts and impact dynamics of Bt (*Bacillus thuringiensis*) cotton in India. *Proceedings of the National Academy of Sciences*, *109*(29), 11652–11656. https://doi.org/10.1073/pnas.1203647109

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, *37*(8), Article 8. https://doi.org/10.1038/s41587-019-0201-4

Kondrashov, F. A. (2012). Gene duplication as a mechanism of genomic adaptation to a changing environment. *Proceedings. Biological Sciences*, *279*(1749), 5048–5057. https://doi.org/10.1098/rspb.2012.1108

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, *9*(4), 357–359. https://doi.org/10.1038/nmeth.1923

Lawrie, R. D., Mitchell Iii, R. D., Deguenon, J. M., Ponnusamy, L., Reisig, D., Pozo-Valdivia, A. D., Kurtz, R. W., & Roe, R. M. (2020). Multiple known mechanisms and a possible role of an enhanced immune system in Bt-resistance in a field population of the Bollworm, *Helicoverpa zea* : Differences in gene expression with RNAseq. *International Journal of Molecular Sciences*, *21*(18), E6528. https://doi.org/10.3390/ijms21186528

Lenth, R. V., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love, J., Miguez, F., Riebl, H., & Singmann, H. (2023). *emmeans: Estimated Marginal Means, aka Least-Squares Means* (1.8.7) [Computer software]. https://cran.r-project.org/web/packages/emmeans/index.html

Liu, C., Xiao, Y., Li, X., Oppert, B., Tabashnik, B. E., & Wu, K. (2014). Cis-mediated down-regulation of a trypsin gene associated with Bt resistance in cotton bollworm. *Scientific Reports*, *4*(1), Article 1. https://doi.org/10.1038/srep07219

Liu, L., Li, Z., Luo, X., Zhang, X., Chou, S.-H., Wang, J., & He, J. (2021). Which is stronger? A continuing battle between Cry toxins and insects. *Frontiers in Microbiology*, *12*. https://www.frontiersin.org/article/10.3389/fmicb.2021.665101

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.
https://doi.org/10.1186/s13059-014-0550-8

Luttrell, R. G., Wan, L., & Knighten, K. (1999). Variation in susceptibility of Noctuid
(Lepidoptera) larvae attacking cotton and soybean to purified endotoxin proteins and
commercial formulations of *Bacillus thuringiensis*. *Journal of Economic Entomology*,
92(1), 21–32. https://doi.org/10.1093/jee/92.1.21

Lynch, M., & Walsh, B. (1998). *Genetics and analysis of quantitative traits*. Sinauer.

Ma, X., Shao, E., Chen, W., Cotto-Rivera, R. O., Yang, X., Kain, W., Fei, Z., & Wang, P.
(2022). Bt Cry1Ac resistance in *Trichoplusia ni* is conferred by multi-gene mutations.
Insect Biochemistry and Molecular Biology, *140*, 103678.
https://doi.org/10.1016/j.ibmb.2021.103678

Ma, Y., Zhang, Y., Chen, R.-R., Ren, X.-L., Wan, P.-J., Mu, L.-L., & Li, G.-Q. (2013).
Combined effects of three crystalline toxins from *Bacillus thuringiensis* with seven
proteinase inhibitors on beet armyworm, *Spodoptera exigua* Hübner (Lepidoptera:
Noctuidae). *Pesticide Biochemistry and Physiology*, *105*(3), 169–176.
https://doi.org/10.1016/j.pestbp.2013.01.007

Mahmoud, M., Gobet, N., Cruz-Dávalos, D. I., Mounier, N., Dessimoz, C., & Sedlazeck, F. J.
(2019). Structural variant calling: The long and the short of it. *Genome Biology*, *20*(1),
246. https://doi.org/10.1186/s13059-019-1828-7

McGaughey, W., Gould, F., & Gelernter, W. (1998). Bt resistance management. *Nature*
Biotechnology, *16*, 144–146. https://doi.org/10.1038/nbt0298-144

McKenzie, J. A., & Batterham, P. (1994). The genetic, molecular and phenotypic consequences
of selection for insecticide resistance. *Trends in Ecology & Evolution*, *9*(5), 166–169.
https://doi.org/10.1016/0169-5347(94)90079-5

Mikheyev, A. S., Tin, M. M. Y., Arora, J., & Seeley, T. D. (2015). Museum samples reveal rapid
evolution by wild honey bees exposed to a novel parasite. *Nature Communications*, *6*(1),
7991. https://doi.org/10.1038/ncomms8991

Mota-Sanchez, D., & Wise, J. C. (2022). *The arthropod pesticide resistance database*.
www.pesticideresistance.org/search.php

Mouches, C., Pauplin, Y., Agarwal, M., Lemieux, L., Herzog, M., Abadon, M., Beyssat-
Arnaouty, V., Hyrien, O., De Saint Vincent, B. R., & Georghiou, G. P. (1990).
Characterization of amplification core and esterase B1 gene responsible for insecticide
resistance in *Culex*. *Proceedings of the National Academy of Sciences*, *87*(7), 2574–2578.
https://doi.org/10.1073/pnas.87.7.2574

Muhlia-Almazán, A., Sánchez-Paz, A., & García-Carreño, F. L. (2008). Invertebrate trypsins: A
review. *Journal of Comparative Physiology B*, *178*(6), 655–672.
https://doi.org/10.1007/s00360-008-0263-y

Nauen, R., Bass, C., Feyereisen, R., & Vontas, J. (2022). The role of cytochrome P450s in insect toxicology and resistance. *Annual Review of Entomology*, *67*(1), 105–124. https://doi.org/10.1146/annurev-ento-070621-061328

Palumbi, S. R. (2001). Humans as the world's greatest evolutionary force. *Science*, *293*(5536), 1786–1790.

Pearce, S. L., Clarke, D. F., East, P. D., Elfekih, S., Gordon, K. H. J., Jermiin, L. S., McGaughran, A., Oakeshott, J. G., Papanikolaou, A., Perera, O. P., Rane, R. V., Richards, S., Tay, W. T., Walsh, T. K., Anderson, A., Anderson, C. J., Asgari, S., Board, P. G., Bretschneider, A., … Wu, Y. D. (2017). Genomic innovations, transcriptional plasticity and gene loss underlying the evolution and divergence of two highly polyphagous and invasive *Helicoverpa* pest species. *BMC Biology*, *15*(1), 63. https://doi.org/10.1186/s12915-017-0402-6

Perry, E. D., Ciliberto, F., Hennessy, D. A., & Moschini, G. (2016). Genetically engineered crops and pesticide use in U.S. maize and soybeans. *Science Advances*, *2*(8), e1600850. https://doi.org/10.1126/sciadv.1600850

Puinean, A. M., Foster, S. P., Oliphant, L., Denholm, I., Field, L. M., Millar, N. S., Williamson, M. S., & Bass, C. (2010). Amplification of a cytochrome P450 gene is associated with resistance to neonicotinoid insecticides in the aphid *Myzus persicae*. *PLOS Genetics*, *6*(6), e1000999. https://doi.org/10.1371/journal.pgen.1000999

R Development Core Team. (2023). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. http://www.R-project.org.

Reisig, D. D. (2017). Factors associated with willingness to plant non-Bt maize refuge and suggestions for increasing refuge compliance. *Journal of Integrated Pest Management*, *8*(1), 9. https://doi.org/10.1093/jipm/pmx002

Reisig, D. D., Huseth, A. S., Bacheler, J. S., Aghaee, M.-A., Braswell, L., Burrack, H. J., Flanders, K., Greene, J. K., Herbert, D. A., Jacobson, A., Paula-Moraes, S. V., Roberts, P., & Taylor, S. V. (2018). Long-term empirical and observational evidence of practical *Helicoverpa zea* resistance to cotton with pyramided Bt toxins. *Journal of Economic Entomology*, *111*(4), 1824–1833. https://doi.org/10.1093/jee/toy106

Reisig, D. D., & Kurtz, R. (2018). Bt resistance implications for *Helicoverpa zea* (Lepidoptera: Noctuidae) insecticide resistance management in the United States. *Environmental Entomology*, *47*(6), 1357–1364. https://doi.org/10.1093/ee/nvy142

Roberts Kingman, G. A., Vyas, D. N., Jones, F. C., Brady, S. D., Chen, H. I., Reid, K., Milhaven, M., Bertino, T. S., Aguirre, W. E., Heins, D. C., von Hippel, F. A., Park, P. J., Kirch, M., Absher, D. M., Myers, R. M., Di Palma, F., Bell, M. A., Kingsley, D. M., & Veeramah, K. R. (2021). Predicting future from past: The genomic basis of recurrent and rapid stickleback evolution. *Science Advances*, *7*(25), eabg5285. https://doi.org/10.1126/sciadv.abg5285

Rochette, N. C., Rivera-Colón, A. G., & Catchen, J. M. (2019). Stacks 2: Analytical methods for

paired-end sequencing improve RADseq-based population genomics. *Molecular Ecology*, *28*(21), 4737–4754. https://doi.org/10.1111/mec.15253

Rubin, C. J., Zody, M. C., Eriksson, J., Meadows, J. R. S., Sherwood, E., Webster, M. T., Jiang, L., Ingman, M., Sharpe, T., Ka, S., Hallböök, F., Besnier, F., Carlborg, Ö., Bed'hom, B., Tixier-Boichard, M., Jensen, P., Siegel, P., Lindblad-Toh, K., & Andersson, L. (2010). Whole-genome resequencing reveals loci under selection during chicken domestication. *Nature*, *464*(7288), Article 7288. https://doi.org/10.1038/nature08832

Rudman, S. M., Greenblum, S. I., Rajpurohit, S., Betancourt, N. J., Hanna, J., Tilk, S., Yokoyama, T., Petrov, D. A., & Schmidt, P. (2022). Direct observation of adaptive tracking on ecological time scales in Drosophila. *Science*, *375*(6586), eabj7484. https://doi.org/10.1126/science.abj7484

Schiebelhut, L. M., Puritz, J. B., & Dawson, M. N. (2018). Decimation by sea star wasting disease and rapid genetic change in a keystone species, *Pisaster ochraceus*. *Proceedings of the National Academy of Sciences*, *115*(27), 7069–7074. https://doi.org/10.1073/pnas.1800285115

Schmidt, J. M., Good, R. T., Appleton, B., Sherrard, J., Raymant, G. C., Bogwitz, M. R., Martin, J., Daborn, P. J., Goddard, M. E., Batterham, P., & Robin, C. (2010). Copy number variation and transposable elements feature in recent, ongoing adaptation at the Cyp6g1 locus. *PLOS Genetics*, *6*(6), e1000998. https://doi.org/10.1371/journal.pgen.1000998

Stahlke, A. R., Epstein, B., Barbosa, S., Margres, M. J., Patton, A. H., Hendricks, S. A., Veillet, A., Fraik, A. K., Schönfeld, B., McCallum, H. I., Hamede, R., Jones, M. E., Storfer, A., & Hohenlohe, P. A. (2021). Contemporary and historical selection in Tasmanian devils (*Sarcophilus harrisii*) support novel, polygenic response to transmissible cancer. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1951), 20210577. https://doi.org/10.1098/rspb.2021.0577

Storz, J. F., & Wheat, C. W. (2010). Integrating evolutionary and functional approaches to infer adaptation at specific loci. *Evolution*, *64*(9), 2489–2509. https://doi.org/10.1111/j.1558-5646.2010.01044.x

Sun, D., Zhu, L., Guo, L., Wang, S., Wu, Q., Crickmore, N., Zhou, X., Bravo, A., Soberón, M., Guo, Z., & Zhang, Y. (2022). A versatile contribution of both aminopeptidases N and ABC transporters to Bt Cry1Ac toxicity in the diamondback moth. *BMC Biology*, *20*(1), 33. https://doi.org/10.1186/s12915-022-01226-1

Székács, A., Lauber, É., Juracsek, J., & Darvas, B. (2010). Cry1Ab toxin production of MON 810 transgenic maize. *Environmental Toxicology and Chemistry*, *29*(1), 182–190. https://doi.org/10.1002/etc.5

Tabashnik, B. E., & Carrière, Y. (2017). Surge in insect resistance to transgenic crops and prospects for sustainability. *Nature Biotechnology*, *35*(10), 926–935. https://doi.org/10.1038/nbt.3974

Taylor, K. L., Hamby, K. A., DeYonke, A. M., Gould, F., & Fritz, M. L. (2021). Genome

evolution in an agricultural pest following adoption of transgenic crops. *Proceedings of*
the National Academy of Sciences, *118*(52). https://doi.org/10.1073/pnas.2020853118

US-EPA. (2018). *White Paper on Resistance in Lepidopteran Pests of Bacillus thuringiensis (Bt)*
Plant-incorporated Protectants in the United States.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund,
G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M.,
Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019).
Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686.
https://doi.org/10.21105/joss.01686

Wilding, C. S. (2018). Regulating resistance: CncC:Maf, antioxidant response elements and the
overexpression of detoxification genes in insecticide resistance. *Current Opinion in*
Insect Science, *27*, 89–96. https://doi.org/10.1016/j.cois.2018.04.006

Yang, F., González, J. C. S., Williams, J., Cook, D. C., Gilreath, R. T., & Kerns, D. L. (2019).
Occurrence and ear damage of *Helicoverpa zea* on transgenic *Bacillus thuringiensis*
maize in the field in Texas, U.S. and its susceptibility to Vip3A protein. *Toxins*, *11*(2),
Article 2. https://doi.org/10.3390/toxins11020102

Yang, F., Kerns, D., & Huang, F. (2015). Refuge-in-the-bag strategy for managing insect
resistance to Bt maize. *Outlooks on Pest Management*, *26*(5), 226–228.
https://doi.org/10.1564/v26_oct_10

Yang, F., Kerns, D. L., Little, N., Brown, S. A., Stewart, S. D., Catchot, A. L., Cook, D. R.,
Gore, J., Crow, W. D., Lorenz, G. M., Towles, T., & Tabashnik, B. E. (2022). Practical
resistance to Cry toxins and efficacy of Vip3Aa in Bt cotton against *Helicoverpa zea*.
Pest Management Science, *78*(12), 5234–5242. https://doi.org/10.1002/ps.7142

Zdobnov, E. M., Kuznetsov, D., Tegenfeldt, F., Manni, M., Berkeley, M., & Kriventseva, E. V.
(2021). OrthoDB in 2020: Evolutionary and functional annotations of orthologs. *Nucleic*
Acids Research, *49*(D1), D389–D393. https://doi.org/10.1093/nar/gkaa1009

Zhang, M., Wei, J., Ni, X., Zhang, J., Jurat-Fuentes, J. L., Fabrick, J. A., Carrière, Y., Tabashnik,
B. E., & Li, X. (2019). Decreased Cry1Ac activation by midgut proteases associated with
Cry1Ac resistance in *Helicoverpa zea*. *Pest Management Science*, *75*(4), 1099–1106.
https://doi.org/10.1002/ps.5224

Zhou, X., Carbonetto, P., & Stephens, M. (2013). Polygenic modeling with Bayesian sparse
linear mixed models. *PLOS Genetics*, *9*(2), e1003264.
https://doi.org/10.1371/journal.pgen.1003264

Zhou, X., & Stephens, M. (2012). Genome-wide efficient mixed-model analysis for association
studies. *Nature Genetics*, *44*(7), 821–824. https://doi.org/10.1038/ng.2310

Position on Chromosome 9 (Mb)