Effective lowering of α-synuclein expression by targeting G-quadruplex structures within the SNCA genome

Valentina Pirota,[1,2,†,] Federica Rey,[3,2,†] Valentina Fantini,[4] Letizia Esposito,[3] Cecilia Pandini,[5] Rosalinda Di Gerlando,[6,7] Filippo Doria,[1] Mariella Mella,[1] Orietta Pansarasa,[8] Paolo Gandellini,[5] Mauro Freccero,[1,‡] Stephana Carelli,[3,9,*,‡] Cristina Cereda,[9,‡]*

[1] Department of Chemistry, University of Pavia, Pavia, Italy

[2] G4-INTERACT, USERN, Pavia, Italy

[3] Pediatric Clinical Research Center "Fondazione Romeo ed Enrica Invernizzi", Department of Biomedical and Clinical Sciences "L. Sacco", University of Milan, Milan, Italy

[4] Laboratory of Neurobiology and Neurogenetic, Golgi-Cenci Foundation, Abbiategrasso, Italy

[5] Department of Biosciences, University of Milan, Milan, Italy

[6] Department of Biology and Biotechnology "Lazzaro Spallanzani", University of Pavia, Pavia, Italy

[7] Molecular Biology and Transcriptomic Unit, IRCCS Mondino Foundation, Pavia, Italy

[8] Cellular Models and Neuroepigenetics Units, IRCCS Mondino Foundation, Pavia, Italy

[9] Center of Functional Genomics and Rare diseases, Buzzi Children's Hospital, Milan, Italy

KEYWORDS: G-quadruplex, SNCA gene modulation, Alpha-synuclein, Parkinson's disease.

ABSTRACT

Alpha-synuclein, encoded by the SNCA gene, is a pivotal protein implicated in the pathogenesis of synucleinopathies, including Parkinson's disease. Current approaches for modulating alpha-synuclein levels involve antisense nucleotides, siRNAs, and small molecules targeting SNCA's 5'-UTR mRNA. Here, we propose a groundbreaking strategy targeting G-quadruplex structures to effectively modulate SNCA gene expression and lowering alpha-synuclein amount. Novel G-quadruplex sequences, identified on the SNCA gene's transcription starting site and 5'-UTR of SNCA mRNAs, were experimentally confirmed for their stability through biophysical assays and *in vitro* experiments on human genomic DNA. Biological validation in differentiated SH-SY5Y cells revealed that well-known G-quadruplex ligands remarkably stabilized these structures, inducing the modulation of SNCA mRNAs expression, and the effective decrease in alpha-synuclein amount. Besides, a novel peptide nucleic acid conjugate, designed to selectively disrupt of G-quadruplex within the SNCA gene promoter, caused a promising lowering of both SNCA mRNA and alpha-synuclein protein. Altogether our findings highlight G-quadruplexes' key role as intriguing biological targets in achieving a notable and successful reduction in alpha-synuclein expression, pointing to a novel approach against synucleinopathies.

INTRODUCTION

Alpha-synuclein (α-Syn) is a presynaptic protein of 140 residues, encoded by the SNCA gene, widely expressed throughout the brain, with a high concentration in the presynaptic nerve terminals.[1] Although its physiological role is not yet fully characterized, it is mainly related to membrane binding, synaptic vesicle recycling, neurotransmitter release, and dopamine

metabolism.[2] Under pathological conditions, impaired regulation of SNCA (including duplication or triplication of the SNCA gene locus) can induce an increase of α-Syn expression as high as 200% relative to its physiological levels. This leads to the accumulation of α-Syn monomers resulting in the formation of misfolded proteins that assemble into oligomers through the formation of β-sheet-rich fibrils.[3] These aggregates of α-Syn are the major components of Lewy Bodies, spherical filamentous protein inclusions that constitute the distinctive histopathological hallmarks in neurons of Parkinson's disease (PD) affected brains.[3] The accumulation and release of α-Syn aggregates from affected brain regions can trigger synaptic and mitochondrial dysfunctions, changes in the lysosomal autophagy system, increased oxidative stress, as well as disruption of axonal transport, leading to neurodegeneration and progressive spreading of PD.[1, 4]

In principle, these pieces of evidence suggest the potential of new therapeutic strategies aimed at lowering α-Syn production or increasing its clearance through the prevention and/or removal of its toxic aggregated deposits.[5] Many different approaches have been investigated in the last decade for preventing excessive buildup of α-Syn (a reduction of 25%–50% is expected to be sufficient to restore normal levels), and among them, the downregulation of SNCA gene expression attracted increasing interest in slowing or even stopping PD pathogenesis.[6] So far, promising results have been reached by epigenetic modification of SNCA through DNA methylation,[7] β2-adrenergic receptor agonists Clenbuterol and Salbutamol,[8] Gapmer-type antisense nucleotides,[9-11] short-interfering RNAs,[12] and small molecules acting on SNCA mRNAs.[13, 14]

In this context, a new and promising strategy for lowering α-Syn production could be obtained by targeting G-quadruplex (G4) nucleic acid secondary structural motives. These are non-canonical high-order structures that occur in specific regions of the genome rich in guanines, like telomeres, gene promoters, transcription start sites (TSS), introns, and both 5' and 3' untranslated regions

(UTRs).[15-18] Under physiological conditions, four guanine bases can self-associate via Hoogsteen-type hydrogen bonds base-pairing generating square-planar guanine quartets, known as G-tetrads. Two or more G-tetrads can stack on top of each other, through a cooperative stabilization due to monovalent cations (e.g. K^+), constituting the backbone of G4 structures.[15] It is well-verified that G4s act as key structural elements in the regulation of a multitude of biological processes, among which is the modulation of gene expression.[19] Indeed, there are strong pieces of evidence that the stabilization of G4 structures in gene promoter regions interplays with the control of the transcriptional activity either increasing or decreasing it.[20, 21] Moreover, G4s on 5'- and 3'-UTR mRNAs regions have been correlated with the regulation of the initial stage of the translation process, generally acting as structural silencing factors.[17] Therefore, the selective targeting of G4s, which has been reached successfully by small molecules,[22] is emerging as a novel therapeutic approach for a growing list of different pathologies, such as cancers,[16, 18, 23] bacterial and viral infections,[24, 25] and neurodegenerative diseases.[26] About the last topic, it is important to note that G4s are often recognized as pathogenic drivers in various neurological disorders, with protective or deleterious effects in the cascade neurotoxic processes.[26] Moreover, the well-defined folded nucleic acid structures of G4s make them advantageous therapeutic targets for upcoming drug development, mimicking the strategy of protein-targeted therapies.[21]

In this context, this work demonstrates, for the first time, the discovery and modulation of two new G4s, respectively located on the TSS of the SNCA gene (pSNCA) and on the 5'-UTR of the SNCA mRNA (mSNCA), in steering SNCA transcription process as well as in lowering α-Syn production.

Biophysical studies demonstrated the folding of these sequences into very stable G4s under physiological conditions, which were further stabilized by well-known G4 ligands (G4Ls). In

addition, the disruption of pSNCA G4 structure was achieved by a new conjugate based on a peptide nucleic acid sequence able to hybridize with pSNCA nucleo bases (NLS-PNA-C343). The treatment of differentiated SH-SY5Y cells with HPHAM[25, 27] and Pyridostatin[28] G4-ligands, leads to the G4s' stabilization, with an increase in the neosynthesis of SNCA mRNAs and a simultaneous decrease in α-Syn production. Conversely, the treatment with NLS-PNA-C343 leads to a decrease in SNCA mRNAs and α-Syn expression.

Altogether these results emphasize the promising role of these new G4s as very promising targets for understanding SNCA regulation processes and reaching a novel and effective decrease in α-Syn production.

RESULTS AND DISCUSSION

G-quadruplex folding prediction in SNCA promoter and transcripts.

The opportunity to exploit the targeting of G4 structures for the modulation of α-Syn imbalance was suggested by the bioinformatic identification of two promising putative quadruplex sequences (PQSs): one straddles the SNCA TSS (pSNCA) and one within the 5'-UTR of SNCA-transcripts in the internal ribosome entry site (mSNCA).

These PQSs were identified by using the online algorithm-based software "QGRS Mapper",[29] "G4 Hunter",[30] and "QuadBase2",[31] where the likelihood of forming a stable G4 is evaluated by a G-score ranking.

The SNCA gene is located on chromosome 4q22.1 (PARK1/4 loci), it contains 6 exons and spans about 114 kb. The main encoded transcript is SNCA140, translated by the last 5 exons of the gene. Other alternatively spliced transcripts encoding different isoforms have been identified for this

gene, and in particular three (SNCA98, SNCA112, and SNCA126) result from alternative splicing of exons 3 and 5.[32]

SNCA promoter region (Chr4: 89835200-89838801) is rich in G bases (22.4%), with a total %GC of 49.4% (Figure S1). Excluding the overlaps, 19 PQSs were identified, among which the only one with a QGRS Mapper G-score higher than 21 was pSNCA ($G_4ATG_4CAG_5CGCG_4$; location Chr4: 89837301-89837324; 24nt length) (Table S1). Moreover, this PQS was the only one also identified by "QuadBase2"[31] configuring medium stringency for G4 motifs. pSNCA sequence is composed of four consecutive G-tracts of four guanines (Gs), respectively divided by a double-nucleotide tract (AT) and by two three-base sequences (CAG and CGC). It is located in the CpG island region, in the non-coding strand, and it straddles the TSS of the SNCA gene (-14/+10), reinforcing its potential as a promising target involved in the regulation of SNCA expression. Interestingly, this sequence shows QGRS Mapper and G4 Hunter score values higher than those of well-known G4s (Table 1),[33] emphasizing its potential ability in G4 folding with extraordinary structural stability.

The 5′-UTR of the human SNCA mRNA is a 264 nt long G-rich (41.2%) region, with an overall 65.5% of GC content. Previous bioinformatics investigation of this region led to the discovery of three non-overlapping PQS motifs (G1, G2, and G3 in Table 1)[34] with rather subpar QGRS Mapper and G4 Hunter scores in comparison to verified mRNA G4 structures. Considering that their combined effect resulted in 45% translation inhibition,[34] the 5′-UTR of the human SNCA mRNA was further analyzed to uncover a very promising PQS that spans G1 and G2 (mSNCA in Table 1; $GA_2(G_2A)_3CUA(G_2A)_3G_2$; 23 nt length). mSNCA sequence is made up of two repetitions of the GGA triplet belonging to G1, a GGACUA exa-nucleotide stretch, and three repetitions of the GGA triplet with a terminal two-bases tract (GG) belonging to G2. Interestingly, it alone comprises

approximately 46% of guanines that have been previously mutated[34] to confirm the relevance of G4 structures within 5′-UTR SNCA mRNA in translation inhibition.

Table 1. G-score values of pSNCA and mSNCA compared with the ones of well-known DNA and RNA G4s.

	Name	Base sequences	QGRS Mapper G-score	G4Hunter score
DNA sequences	pSNCA	$G_4ATG_4CAG_5CGCG_4$	62	2.75
	Myc-23456	$GAG_3TG_4AG_3TG_4 A_2G_2$	41	2.39
	VEGF	$G_3AG_3T_2G_4TG_3$	41	2.53
	Bcl2	$G_3CG_3CGCG_3AG_2A_2G_5CG_3$	40	2.11
	KRAS	$AG_3CG_2TGTG_3A_2GAG_3A_2GAG_5AG_2$	40	1.78
	CKIT-1	$AG_3AG_3CGCTG_3AG_2AG_3$	39	1.77
	22AG	$AG_3T_2AG_3T_2AG_3T_2AG_3$	42	1.64
	CKIT-2	$C_3G_3CG_3CGCGAG_3AG_4AG_2$	38	1.42
	HRAS-1	$TCG_3T_2GCG_2CGCAG_2CACG_3CG$	41	1.19
RNA sequences	ADAM10	$G_3ACG_3UAG_4CG_3$	42	2.28
	MMP16	$G_3AG_3AG_3AGAG_3$	40	2.18
	CCND3	$G_3CG_2CG_3CG_3CUG_4$	39	2.15
	TERRA	$AG_3U_2AG_3U_2AG_3U_2AG_3$	42	1.64
	G2	$(G_2A)_3G_2$	21	1.45
	mSNCA	$(G_2A)_3CUA(G_2A)_3G_2$	21	1.17
	G1	$G_2AGA_2G_2AGA_2G_2AG_2$	18	1.06
	G3	$G_4ATG_4CAG_5CGCG_4$	12	1.61

Confirmation of G4s folding in SNCA promoter and transcripts.

CD, UV-vis, FRET, and NMR Studies on SNCA model oligonucleotides

The ability of the pSNCA DNA sequence ($G_4ATG_4CAG_5CGCG_4$, 24nt) to fold into G4 structures was first confirmed by recording circular dichroic (CD) spectra in the presence of different KCl concentrations (from 0 to 100 mM). In the presence of high potassium concentrations (100 mM K^+), pSNCA (2.5 µM) is mainly folded into an antiparallel topology (calculated equal to 64%),[35] featured by a positive peak around 295 nm and a negative one around 260 nm (Figure 1A).[36] A quantitative secondary structural analysis[35] estimated the presence of 29.4% *syn-anti* and 24.5% *anti-syn* glycosidic bond conformations, with 47.1% of diagonal and lateral loops.[35] As the potassium concentration decreases, the sequence rebalances itself towards a 3+1 hybrid topology, highlighted by the growth of a positive shoulder at around 270 nm (hybrid topology percentage equal to 68% with 0.5 mM K^+). To strengthen the biophysical model making it more consistent with the real biological SNCA G4-folding equilibria, an extended sequence, pSNCAext ($GATG_4ATG_4CAG_5CGCG_4TGA$, 30-nt), that includes the 3 previous nucleobases (GAT) and the 3 subsequent nucleobases (TGA) present at the genomic level, was also studied. pSNCAext allowed us to better explore how the involvement of the adjacent bases influences the folding of the G4 structure. In this case, the addition of 6 nucleobases, including two more guanines, shifts the equilibrium towards a 3+1 hybrid topology (92% with 100 mM K^+), characterized by a positive peak at around 295 nm with a shoulder at around 275 nm (Figure 1B).[36] In this case, the evaluation of secondary structural element fractions included many more combinations: 6.3% *anti-anti*, 19.0% *syn-anti*, and 11.8% *anti-syn* of glycosidic bond conformations, with 41.9% of diagonal and later loops and 21.0% of remaining base steps.[35] By decreasing potassium chloride, a transition to a mixture of hybrid and antiparallel topologies (equal to 62.1% and 37.9% respectively with 0.5 mM K^+) was obtained. A complete analysis of estimated secondary and tertiary structure fractions

of pSNCA and pSNCAext G4s as a function of potassium chloride concentration is reported as Supporting Information (Figure S2).

CD-melting experiments on both the model oligonucleotides described their folding into extremely stable G4s, showing mid-transition temperature (T_m) values equal to $61.7 \pm 0.1°C$ and $56.8 \pm 0.1°C$ for pSNCA and pSNCAext respectively in the presence of only 1 mM K^+. This potassium chloride concentration is over two orders of magnitude lower than the physiological value in the cell (around 150 mM),[37] but the high stability of these two new G4s makes it difficult to have a correct measurement of T_m values by biophysical methods already at 50 mM KCl. Indeed, by CD-melting, T_m values are very close to the detection limit set at 95°C with 50 mM K^+ ($94.3 \pm 0.7°C$ and 91.8 ± 0.3 °C respectively for pSNCA and pSNCAext) and over 95°C with 100 mM K^+. To further underline the remarkable stability of these pSNCA-G4s, we performed CD-melting experiments on well-known G4-models (Myc-23456, cMyc, 22ag, CKIT-1, and LTR-III) [33, 38] with different KCl concentrations to maintain T_m values in the 50-60 °C range. It should be noted that the model sequence of the most stable parallel G4 known, Myc-23456, showed a T_m of $61.2 \pm 0.4°C$ in the presence of 5 mM K^+, about 12 degrees lower than T_m obtained with pSNCA ($74.6 \pm 0.1°C$) and pSNCAext ($73.2 \pm 0.6°C$) in the same conditions (Table S2).

The ability of pSNCA and pSNCAext (2.5 µM) to fold into highly stable G4s was also confirmed by UV-vis melting experiments in the presence of 1 mM KCl. Following the absorbance decrease at 295 nm, mid-transition temperature values were calculated to be $61.5 \pm 0.4°C$ and 55.1 ± 0.1 °C for pSNCA and pSNCAext respectively,[39] excellent agreement with CD-melting data. From the melting curves, model-dependent thermodynamic parameters were extracted to better evaluate the thermodynamics of these novel G4s.[39] In particular, for pSNCA Van't Hoff enthalpy and entropy values corresponded to -55.5 ± 0.6 kcalmol^{-1} and -166 ± 2 calmol^{-1}K^{-1}, while for pSNCAext we

obtained $\Delta H° = -52.3 \pm 0.9$ kcalmol^{-1} and $\Delta S° = -159 \pm 3$ calmol^{-1}K^{-1}.[39] So, at the physiological

temperature of 37°C, free Gibbs enthalpy ($\Delta G°$) resulted to be -4.02 kcalmol^{-1} and -2.99 kcalmol^{-1} for pSNCA and pSNCAext respectively (Table S3), further underlining the high stability of these

G4s in the presence of only 1 mM potassium chloride.[40]

The folding of these two sequences into G4 structures was also confirmed by UV-vis thermal

difference spectra (TDS), obtained by the subtraction of the spectra of unfolded (temperature

above T_m) and folded (low temperature) G4s, as highlighted by the TDS features peaks: a negative

one at 295 nm, and positive ones at 273 nm, 257 nm, and 243 nm (Figures 1C-D).[41] Moreover, the

calculated TDS factors ($\Delta A240nm/\Delta A295nm$), equal to 1.06 ± 0.08 and 1.3 ± 0.1, confirmed their

folding into antiparallel (group III) and hybrid (group II) topologies.[41]

Fluorescence resonance energy transfer (FRET)-melting experiments, using 5'-FAM/3'-TAMRA

labeled oligonucleotides, were recorded to corroborate mid-transition temperature data. Following

the G4-unfolding of 0.25 µM model oligonucleotides through a temperature-dependent increase

in donor emission of FAM (25-95°C temperature range) in the presence of 1 mM KCl, we

measured pSNCA and pSNCAext T_m equal to 64.3 ± 0.4°C and 58.8 ± 0.8 °C, respectively.[42]

Supporting Information contains a summary of the computed stabilities as a function of potassium

chloride concentrations, determined by exploiting the thermal unfolding of pSNCA and pSNCAext

G4s by different techniques (Table S4).

G4 folding of pSNCA and pSNCAext was finally confirmed by preliminary one-dimensional ^{1}H-

NMR spectra recorded on a Bruker Avance 700 MHz at 25°C (Figures 1E-F). The signals within

the range of 10-12 ppm are characteristic of the Gs imino protons when a G-tetrad is formed, as

compared to 13-14 ppm for those involved in Watson-Crick base pairing.[43]

As no biophysical analysis was previously conducted on the PQSs identified in the 5'-UTR-RNA region of SNCA, a deep biophysical analysis was performed on mSNCA model oligonucleotide ((G_2A)$_3$CUA(G_2A)$_3G_2$, 23 nt) to confirm its propensity to fold into a stable mRNA G4.

Dichroic spectra of mSNCA (2.5 µM) recorded at different potassium concentrations (0, 10, 100 mM) in 10 mM lithium cacodylate buffer pH 7.4, confirmed its folding into a mainly parallel G4 structure (68.3% with 100 mM KCl, Figure S2E), characterized by a positive peak at 264 nm and a negative one at around 240 nm (Figure 1G). The positive signal between 290 nm and 310 nm conforms to sequence reorganization into an antiparallel G4 structure, as confirmed by tertiary structure analysis (around 25% with 100 mM KCl; Figure S2E). Estimation of secondary structural element fractions in 100 mM KCl (Figure S2F) highlighted around 30% of *anti-anti* glycosidic bond conformations, with around 18% of diagonal and later loops and 40.0% of other base steps.[35] CD and UV melting experiments in 10 mM lithium cacodylate buffer pH 7.4 and in the presence of 100 mM KCl, showed a melting temperature of 62.4 ± 0.1 °C, as well as FRET-melting analysis revealed a $T_m = 68.5 \pm 0.6$ in the same experimental conditions. These results emphasized that mSNCA is a pretty stable mRNA G4 when compared to other well-known mRNA G4s (TERRA, ADAM10, MMP16)[23, 44, 45] which melt at around 60°C in the presence of much less potassium chloride (Table S2), following the G-score values identified by bioinformatic analysis. A summary of melting data was reported in Table S4. Model-dependent thermodynamic parameters were extracted from the UV-melting curve, resulting in $\Delta H° = -62.5 \pm 0.2$ kcalmol^{-1} and $\Delta S° = -183.6 \pm 0.4$ calmol^{-1}K^{-1}.[39] As a consequence, $\Delta G°$ resulted to be -5.58 kcalmol^{-1} in the presence of 100 mM potassium chloride at 37 °C (calculated parameters estimated in the presence of 10 mM KCl were reported in Table S3).

mSNCA folding into a G4 structure was also confirmed by UV-vis thermal difference spectra, which revealed a negative peak at around 295 nm, and positive ones at around 273 nm, 257 nm, and 243 nm (Figures 1H).[41] Moreover, the calculated TDS factor (ΔA240nm/ΔA295nm) of 2.4 ± 0.2 highlighted its folding into a mainly parallel G4 (group I).[41]

Figure 1. *Biophysical data supporting the folding of SNCA model oligonucleotides into G4*. CD spectra of 2.5 µM of A) pSNCA and B) pSNCAext in 10 mM lithium cacodylate buffer (pH 7.4) with different potassium concentrations (ranging from 0 to 100 mM). The ionic strength is maintained constant at 100 mM by adding proper amounts of LiCl. TDS spectra of 2.5 µM C) pSNCA and D) pSNCAext in 10mM lithium cacodylate buffer pH 7.4, with 1 mM KCl and 99 mM LiCl. ^{1}H-NMR of 0.25 mM E) pSNCA and F) pSNCAext in 90% H_2O/10% D_2O solution with 100 mM KCl and 20 mM phosphate buffer pH 7.4, at 25°C. G) CD spectra of 2.5 µM of mSNCA in 10 mM lithium cacodylate buffer (pH 7.4) with different potassium concentrations (0-10-100 mM). The ionic strength is maintained constant at 100 mM by adding proper amounts of LiCl. H) TDS spectra of 2.5 µM mSNCA in 10 mM lithium cacodylate buffer pH 7.4, with 100 mM KCl.

Chromatin Immunoprecipitation and qPCR stop assay confirmed SNCA-G4 folding on genomic DNA

To confirm the aforementioned observations on pSNCA G4-folding to the cellular setting, Chromatin Immunoprecipitation (ChIP) and quantitative PCR (qPCR) stop assays were performed. Both ChIP and qPCR stop assays were accomplished on genomic DNA from SH-SY5Y cells, a renowned and widely used neuronal cell model for neurobiology research,[46] after 7 days of neural differentiation.

Primers covering a 182 base-pair region (SNCA_G4-FW and SNCA_G4-REV) centered on TSS were used to check the folding of pSNCA into a G4 structure by G4-ChIP. ChIP assay showed the enrichment of SNCA-G4s DNA upon chromatin immunoprecipitation with the anti-G4 antibody BG4[47] as compared to an anti-FLAG antibody (Figure 2A). In contrast, a non-G4 forming GAPDH

region, used as a negative control, was not enriched, emphasizing that the pSNCA sequence folds into a G4 structure *in vivo*.

qPCR stop assay was performed by applying different potassium chloride concentrations (from 0 to 50 mM), exploiting the same SNCA-G4 primers used for G4-ChIP. Stabilization of SNCA G4 structure by K^+ ions caused an obstacle for the DNA polymerase during DNA amplification, with a reduction of amplicons even with 0.1 mM KCl (Figure 2B). This trend parallels the results obtained for MYC, which was used as a positive control, highlighting the high stability of this new SNCA-G4 (Figure S3). Primers spanning a non-G4 sequence in the GAPDH gene were used as a negative control, and indeed its amplification products did not show a significant alteration in response to the increase in K^+ (Figure S3).

The presence of G4s in the 5'-UTR-RNA region of SNCA was demonstrated by G4-RIP analysis highlighting a specific enrichment for the mRNA SNCA-G4 (Figure 2C). Moreover, the qPCR stop assay performed on cDNA also showed a decreased target amplification at increasing concentrations of KCl (Figure 2D).

Figure 2. *Biological validation of the presence of G4 structures within SNCA promoter and SNCA 5'-UTR mRNA.* A) BG4 and anti-FLAG antibodies were used for chromatin immunoprecipitation and enrichment for the SNCA-G4 DNA region was determined by qPCR. A non-G4 forming GAPDH region was used as a negative control. B) SNCA-G4 abundance at promoter level was assayed via qPCR stop assay with increasing KCl concentrations, using a non-G4 forming GAPDH region as a housekeeping gene. Data are expressed as mean ± SEM of 5 independent samples (n = 5, *p<0.05, ***p<0.001 vs 0mM). C) BG4 and FLAG were used for RNA immunoprecipitation

and enrichment for the SNCA-G4 cDNA region was determined by qPCR. A non-G4 forming GAPDH region was used as a negative control. D) SNCA-G4 abundance at 5'-UTR mRNA level was assayed via qPCR stop assay with increasing potassium chloride concentrations, using a non-G4 forming GAPDH region as a housekeeping gene. Data are expressed as mean ± SEM of 5 independent samples (n = 6). All assays were performed on SH-SY5Y cells after 7 days of neural differentiation.

SNCA-G4 folding modulation by well-known G4-ligands and PNA conjugate

FRET- and CD-melting experiments to select the best-stabilizing G4 ligands

To define the involvement of G4s in the modulation of SNCA gene expression, pSNCA and pSNCAext stabilization were measured in the presence of four well-known G4 ligands. Among the countless known G4 binders, we selected the well-known HPHAM (core-extended naphthalene diimide), PDS (Pyridostatin, a bis-quinoline derivative), BRACO19 (acridine-based ligand), and TMPyP4 (porphyrin-based ligand).[22]

From FRET- and CD-melting results, the three ligands HPHAM, PDS, and BRACO19 stabilize to different extents both model sequences, with the best-performing stabilizer being HPHAM, which is able to stabilize both model oligonucleotides of around 13-16°C (Table 2). On the contrary, TMPyP4 destabilized both the model structures, from 2 to 11°C, consistent with its sporadically described propensity to unfold some G4 structures.[48]

In addition, when analyzing the effect of the binding on the original topology of SNCA-G4s, we realized that among the analyzed binders, only HPHAM can stabilize both pSNCA and pSNCAext G4 structures preserving their original topology unaltered (Figure S4). This is confirmed at both 1 mM and 100 mM KCl concentrations, corresponding respectively to the working concentration of

FRET-/CD-melting and an experimental environment with a physiological-like potassium ions concentration.

Considering the excellent results obtained with HPHAM, its stabilizing power was also evaluated on the mSNCA model oligonucleotide. PDS was also tested as the best-known binder reference. In the presence of 100 mM KCl, both the ligands are able to strongly stabilize the mSNCA G4 structure so much so that it was not possible to identify the melting temperature within 95°C (Table 2).

Table 2. – Melting data summary describing the G4-stabilization/destabilization induced by G4-ligands. pSNCA (or pSNCAext, or mSNCA) 2.5µM and 0.25µM for CD and FRET melting experiments respectively, in the presence of 4 equiv. of G4-ligands, and 10 mM lithium cacodylate buffer pH 7.4. Data for pSNCA and pSNCAext were recorded with 1 mM KCl and 99 mM LiCl. Data for mSNCA were recorded with 100 mM KCl.

	ΔT_m values (°C)					
	pSNCA		pSNCAext		mSNCA	
	1 mM KCl		1 mM KCl		100 mM KCl	
G4 Ligand	**CD-melting**	**FRET-melting**	**CD-melting**	**FRET-melting**	**CD-melting**	**FRET-melting**
HPHAM	13.1 ± 0.5	15.2 ± 0.3	14.4 ± 0.3	16.1 ± 0.3	> 32.6	> 26.5
PDS	8.9 ± 0.3	9.1 ± 0.5	4.3 ± 0.2	6.4 ± 0.1	> 32.6	> 26.5
BRACO19	7.0 ± 0.3	8.2 ± 0.8	9.5 ± 0.2	11.5 ± 0.7		
TMPyP4	-7.1 ± 0.8	-11 ± 1	-2.2 ± 0.6	-3.5 ± 0.5		

CD study to test the ability of a new PNA conjugate to disrupt pSNCA G4 structure

Although the literature has demonstrated that the presence of G4 structures in the SNCA 5'-UTR mRNA region, and consequently their stabilization, mediates a negative response on SNCA

mRNA translation,[34] it was to be defined how the newly discovered G4 on the promoter interferes with transcription processes. Therefore, in parallel with the identification of good binders capable of strongly stabilizing its structure, a short peptide nucleic acids (PNA)[49] sequence (C_4GTC_3) capable of hybridizing with the central bases of the pSNCA (or pSNCAext) was synthesized. PNA sequence was chosen to effectively force the unfolding of the highly stable G4 structure within the TSS of the SNCA gene, in favour of the PNA-DNA duplex formation.[50, 51] PNAs, compared to other antisense tools, allow for a tighter bond with natural nucleic acids due to the lack of polyanionic electrostatic interactions (the backbone of PNAs is an achiral aminoethyl glycyl spacer) and greater stability as they are not recognized by nucleases or proteases.[50] In addition, PNAs can efficiently discriminate a one-base mutation with remarkable specificity,[52] making this tool highly selective.

A nuclear localization sequence (NLS) H_2N-ProLysLysLysArgLysVal-OH was conjugated to the PNA C-terminus, with the inclusion of a glycine residue as a spacer, to accomplish PNAs delivery into the nuclei. In addition, a Coumarin 343 (C343) residue was coupled to the N-terminus of PNA by separating the dye through a lysine residue, which is positively charged and increases the electrostatic contact between the two strands, implementing water-solubility and cellular uptake.

The power of the new NLS-PNA-C343 conjugate (H_2NCO-NLS-Gly-C_4GTC_3-Lys-C343) was tested in the interaction with the model sequence pSNCAext to better mimic its effect on an extended region. CD spectrum of 2.5 μM pSNCAext with 100 mM KCl in 10 mM lithium cacodylate buffer, pH 7.4, was recorded after the addition of 4 equivalents of NLS-PNA-C343 left to equilibrate in solution for 4 hours at 25°C. A drastic change in dichroic signal with a positive band around 278 nm and a negative one around 252 nm was obtained, confirming the PNA-DNA duplex formation, following G4 unfolding (Figure S5). The melting temperature of this new hetero

duplex resulted to be 54.3 ± 0.5 °C, therefore, emphasizing a stable interaction at the physiological temperature of 37°C.

To underline the enormous difference in the dichroic spectra features, due to the interaction with the new PNA conjugate, a Principal Component Analysis (PCA) was conducted. PCA was run on a reference library of 23 known G4s CD spectra.[35] The scores plot of the first two components PC1 (explained variance 62.82%) and PC2 (explained variance 18.87%) showed the separation of three clusters, corresponding to parallel, hybrid, and antiparallel G4 topologies (Figure S6). While the CD spectrum of pSNCAext fits perfectly into the set relating to the hybrid topology G4s (as well as the CD spectrum of pSNCA fits with antiparallel G4s group), the spectrum recorded after the addition of NLS-PNA-C343 does not have the appropriate characteristics to fall into any of the three PCA clusters (Figure S6). This analysis confirms that the dichroic characteristics of the new structure do not correlate with those of known G4s.

Evaluation of SNCA mRNA and α-Syn expression following G4 stabilization

To assess the impact of SNCA-G4 stabilization on SNCA mRNA and α-Syn expression, *in vitro* experimental assays on the SH-SY5Y cell line following neural differentiation in the presence of Retinoic Acid were performed [53, 54]. Considering the biophysical results, we selected HPHAM and PDS as the best stabilizing G4-ligands to test, firstly identifying their non-cytotoxic concentration in the SH-SY5Y cell line by MTT assay. We found that 625 nM was the highest non-toxic dosage for HPHAM, even after 48 hours of treatment (Figure S7). At the same time, a non-toxic concentration of 5 μM was chosen for PDS, in complete agreement with already reported studies on the SH-SY5Y cell line (Figure S8).[55, 56] In addition, taking advantage of the intrinsic red

fluorescence emission of HPHAM, we highlighted that HPHAM efficiently enters cells at 625 nM, already after 4 hours (Figure S9).

By Real-Time PCR we identified an increase over time in SNCA mRNA expression following HPHAM treatment (Figure 3A), which, however, remained stable for up to 48 hours when cells were treated with PDS (Figure 3B).

In both cases, the G4-ligand activity was validated by assessing the expression of MYC as a positive control, which, as expected, was reduced following both HPHAM and PDS treatments (Figures 3A and 3B). TATA Binding Protein (TBP) was used as a housekeeping gene as we found this to be the most stable housekeeping not influenced by G4 stabilization (as opposed to GAPDH and 18S; Figure S10).

Considering the high stability of pSNCA G4, along with its unique localization straddling the TSS, these results highlighted that in the presence of SNCA-G4 stabilizing ligands an increase in SNCA expression is induced. Despite the concept that G4s behave as gene-silencing structural elements was supported over the years in several studies [57], these results are in agreement with the more recent knowledge that G4 formation and stabilization can also actively contribute to enhancing, rather than repressing, gene expression.[21]

To evaluate whether SNCA was newly synthesized, a decay assay was performed treating cells with HPHAM or PDS and the transcriptional inhibitor Actinomycin D was performed (Figures 3C and 3D).[58] In both cases, following treatment with the G4-stabilizer, a faster decay of RNA expression was emphasized, most significant after 8 hours of treatment (around 50% with both the ligands), suggesting that the increase we note with HPHAM stabilization is a result of novel RNA synthesis.

To assess the impact of G4-stabilizing ligands on α-Syn protein expression, Western Blotting analyses were performed on protein extracts obtained from SH-SY5Y cells treated with HPHAM or PDS at non-cytotoxic concentrations (625 nM and 5 μM, respectively) for 0, 4, 8, 24, and 48 hours. In this case, β-actin was used as a housekeeping protein as its expression is not impacted by treatment with the two G4-binders. A drastic reduction in the protein expression was highlighted following treatment with both compounds, with a dropped α-Syn protein production higher than 50% after 24h (Figures 3E-3F).

This promising protein reduction could be due to the action of these G4-ligands on SNCA gene transcription, indeed, both SNCA mRNA increase and impacted stability can affect α-Syn protein production. On the other hand, there is also the possibility that α-Syn protein reduction derived from the impairment of the translation process due to HPHAM- or PDS-induced stabilization of G4s on the 5'-UTR of SNCA mRNAs.[34]

Figure 3. Assessment of SNCA mRNA and α-Syn expression as a consequence of G4 stabilization. SNCA and MYC mRNA expression levels were assessed via Real-Time PCR following treatment with 625 nM HPHAM (A) and 5 µM PDS (B) for 0, 4, 8, 24, and 48 hours.

TBP was used as a housekeeping gene and data are expressed as mean ± SEM of 3 replicate values in 3 independent experiments (n=9; *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 vs T0). SNCA expression levels were also assessed via Real-Time PCR following treatment with 1.5 µg/mL Actinomycin D and either 625 nM HPHAM (C) or 5 µM PDS (D) for 0, 4, 8, 24, and 48 hours. α-Syn expression levels were assessed via Western Blotting following treatment with 625 nM HPHAM (E) and 5 µM PDS (F) for 0, 4, 8, 24, and 48 hours. β-Actin was used as a housekeeping protein and data are expressed as mean ± SEM of 3 independent experiments (n=3; *p<0.05 vs T0).

Selective disruption of TSS G4 leads to the reduction in SNCA mRNA and α-Syn expression.

To better understand the specific role of the G4 structure within the promoter region of the SNCA gene, NLS-PNA-C343 was tested for its ability to disrupt the pSNCA G4 folding. NLS-PNA-C343 resulted in being non-toxic at 5 µM dosage after 24 hours (Figure S11), thus selecting this dosage for further experiments. Following 7 days of neural differentiation, cells were treated with NLS-PNA-C343 for 0, 4, 8, and 24 hours.

The effective disruption of pSNCA G4 drove a decrease in SNCA mRNA expression of around 50% after only 4 hours (Figure 4A), conversely to results obtained with G4s' stabilizing ligands. Moreover, this selective targeting led to a decrease in α-Syn expression of around 70% after 24 hours, suggesting that targeting and disruption of the G4 structure at the promoter level can be successfully used to down-regulate both SNCA mRNAs and α-Syn protein expression (Figure 4B).

Figure 4. Assessment of SNCA mRNA and α-Syn expression as a consequence of G4 destabilization. (A) SNCA mRNA expression levels were assessed via Real-Time PCR following treatment with 5 µM SNCA-G4 destabilizer for 0, 4, 8, and 24 hours. TBP was used as a housekeeping gene and data are expressed as mean ± SEM of 3 replicate values in 3 independent experiments (n=9; *p<0.05, **p<0.01, ****p<0.0001 vs T0). (B) α-Syn expression levels were assessed via Western Blotting following treatment with 5 µM SNCA-G4 destabilizer for 0, 4, 8, 24, and 48 hours. β-Actin was used as a housekeeping protein and data are expressed as mean ± SEM of 2 independent experiments (n=2; *p<0.05 vs T0).

CONCLUSIONS

In this work, we present the discovery and modulation of two novel G-quadruplex structures on the TSS of the SNCA gene and within the 5' UTR of SNCA mRNAs, highlighting that they are promising biological targets in reaching an effective lowering of α-Syn expression. The high propensity to fold into G4 structures was demonstrated on model oligonucleotides (pSNCA, pSNCAext, and mSNCA) by different biophysical studies as well as in genomic DNA and cDNA. Our results demonstrate that both pSNCA and pSNCAext are very stable in comparison with other highly structured DNA-G4 models. Moreover, the novel mSNCA G4 turned out to be moderately

stable as RNA-G4 with promising bioinformatic and biophysical results although less effective than those of the known RNA-G4s. The use of well-known G4-ligands (HPHAM and PDS) emphasized the possibility of strengthening SNCA-G4s stability leading to a modulation of SNCA transcription *in vitro* on the SH-SY5Y cell line. The use of both HPHAM and PDS induced a robust decrease in α-Syn protein production (higher than 50%), data which drastically reinforces the possibility of using this new strategy to restore α-Syn healthy levels in neurons of neurodegenerative disease-affected brains. Moreover, selectively targeting the TSS G4 in the SNCA gene promoter led to a decrease in both mRNA and protein levels (around 70%), suggesting that the targeting of this interesting and novel G4 is enough to successfully modulate both gene and protein expression.

Although the development of new hyper-selective ligands for these innovative SNCA-G4s is of pivotal importance, these data, with unprecedented nucleic acid targets and their innovative mechanism of action, offer a novel mechanistic perspective for the possible future development of personalized therapeutic strategies against synucleinopathies.

EXPERIMENTAL SECTION

Solvents, salts, BRACO19 hydrochloride, and TMPyP4 were purchased from Merck and TCI Chemicals and were used as they were without further purification. HPHAM,[59] Pyridostatin,[60] and C343[61] were synthesized according to the previously reported protocols, and their identity was assessed by [1]H-NMR. Analytic HPLC profiles were recorded to confirm HPHAM and Pyridostatin purity major of 98% (Figures S12 and S13). For their chemical structure, please see Figure S14. Fmoc-protected amino acids and resin for SPPS were bought from Novabiochem® (Merck, Milan, Italy), while Fmoc/Bhoc-protected PNA monomers were from Panagene (Daejeon, South Korea).

Oligonucleotides

All the model oligonucleotides were purchased from Eurogentec (Seraing, Belgium) and Biomers.net GmbH (Ulm, Germany) with an HPLC grade of purity (See Table S5 for sequence details).

They were dissolved in nuclease-free water and stored at -20°C. Exact oligonucleotide strand concentrations were determined by absorption measurements at 260 nm, using the extinction coefficient values provided by the manufacturer.

NLS-PNA-C343 synthesis

NLS-PNA-C343 was sequentially assembled using microwave-assisted Fmoc solid phase peptide synthesis on a semi-automatic synthesizer (Biotage® Initiator+). The synthesis was conducted on a Rink Amide resin (100-200 mesh; loading 0.37 mmol/g) operating in a 0.06 mmol scale. It was pre-swelled in DCM and DMF for 1 h each and Fmoc-deprotected using 25% (v/v) piperidine in DMF.

Coupling reactions for NLS synthesis were conducted at 50°C for 15 minutes, using 3 equivalents of Fmoc-protected amino acids, 3 equivalents of HOBt, 3 equivalents of PyBOP, and 6 equivalents of DIPEA in a 1:1 DMF and DCM mixture. In order to assemble the PNA sequence, coupling steps were carried out for 35 min at 75°C with 3 equivalents of Fmoc/Bhoc protected PNA monomers, 3.5 equivalents of Oxyma Pure®, and 3.5 equivalents of N,N′-diisopropylcarbodiimide in 1:1 NMP and DCM mixture.

All the Fmoc-deprotection steps were done by treating the resin twice (3 and 10 min) with a 25% piperidine solution in DMF at room temperature.

Coumarine 343 was directly conjugated on the solid phase at room temperature using 5 equivalents of HATU and 10 equivalents of DIPEA in DMF, by stirring the mixture for 2h.

Resin cleavage was performed using a TFA:TIS:MilliQ H_2O (95:2.5:2.5, v/v) solution. Following three hours of stirring at room temperature, the resin was filtrated under low pressure, and cold diethyl ether was added to induce the precipitation of PNA-conjugate. The crude was resuspended in acidic water (0.1% TFA) and purified by preparative HPLC (Agilent Technologies 1260 Infinity HPLC fitted with a diode array detector) using a Waters XSelectHSS C18 (2.5 µm, 50 x 4.6 mm) column. Eluents: acidic water (0.1% TFA) and acetonitrile. Purification procedure: 2 min of 95% acidic water, 14 min of gradient step to reach 60% aqueous phase; 30 mL/min flow rate; λ = 210 nm. The purity of NLS-PNA-C343 conjugate was evaluated by analytic HPLC, resulting equal to 97.6% (Figure S15).

PNA conjugates were characterized using a surveyor UHPLC system (Thermo Finnigan, San Jose, CA, USA) equipped with a BEH Acquity UPLC column (1.7 µm) 2.1 x 50 mm, and an LCQ ADV MAX ion-trap mass spectrometer, with an ESI ion source (Figures S16).

NLS-PNA-C343. MW: 3651.9 u.m.a.; UHPLC-MS (positive mode): 913.68 [NLS-PNA-C343+4H]$^{4+}$; 731.22 [NLS-PNA-C343+5H]$^{5+}$; 609.75 [NLS-PNA-C343+6H]$^{6+}$; 522.85 [NLS-PNA-C343+7H]$^{7+}$; 457.45 [NLS-PNA-C343+8H]$^{8+}$ m/z + signals related to conjugates with TFA.

Bioinformatic prediction

SNCA promoter region (Chr4: 89835200-89838801; ENSR00000170730) and human SNCA mRNA (NM_000345.3) were analyzed by online available algorithm-based software *QGRS Mapper,*[29] *G4 Hunter,*[30] and *QuadBase2* [31] for predicting putative G4 sequences in both coding and non-coding strands. For *QGRS Mapper* analysis, were considered the following restrictions:

maximum length of 30 nt; minimum G-Group size of 2 nt; loop size of 0−15 nt. For *G4 Hunter* calculations have been set at a threshold of 1.5 with a window size of 20 nt. For *QuadBase2* evaluation has been configured with a medium stringency (G_3L_{1-7}), considering stem length of 3 nt, loop size 1-7 nt, and a Greedy algorithm type.

Circular Dichroism studies

Circular Dichroism analyses were recorded by JASCO J-1500 spectropolarimeter (JASCO corporation, Easton, MD, USA), equipped with a Peltier temperature controller, in a quartz cuvette with a 10 mm optical length.

All oligonucleotides were diluted from stocks to the final concentration (2.5 µM) in lithium cacodylate buffer (10 mM, pH 7.4) containing different concentrations of KCl (0, 0.5, 1, 5, 10, 50, 100 mM). Ionic strength was kept constant by adding suitable quantities of LiCl (to 100 mM chloride salts). The solutions were annealed by heating at 95°C for 5 min and gradually cooling to room temperature over 4h.

CD spectra were recorded at 293 K, an instrument scanning speed of 200 nm/min with a response of 2 s over a wavelength range of 210-340 nm with a 1 nm sampling interval. The reported spectra of each sample represent the average of 4 scans and are baseline-corrected for signal contributions due to the buffer mixture. CD spectra were normalized to $\Delta\varepsilon$ ($M^{-1}cm^{-1}$) according to the relationship $\Delta\varepsilon = \vartheta / (32980 \times C \times l)$, where ϑ corresponded to observed ellipticities (in millidegrees), C is DNA molar concentration, and l is the path length in centimeters. Secondary and tertiary structure fractions predictions were calculated by exploiting the algorithm developed by Del Villar-Guerra et al.[35]

For the CD-melting experiments of G4 structures (2.5 µM), the ellipticity was recorded at 295 nm for pSNCA and pSNCAext and at 264 nm for mSNCA every 0.5°C, with a temperature scan rate of 1 °C/min in the range of 15-95 °C, after a pre-equilibrium of the solution at 15°C for 5 minutes. Data were the average of three independent experiments. G4-ligands were added in 10 µM concentration (4 equivalents) and left to equilibrate for 16 h at room temperature before recording the melting curve. NLS-PNA-C343 was added in 10 µM concentration (4 equivalents) and left to equilibrate for 4 h at room temperature before recording the CD spectrum and the melting curve. Melting temperature values were calculated according to the Van't Hoff equation, applied for a two-state transition from a folded to unfolded state, assuming that the heat capacity of the folded and unfolded states are equal. ΔT_m was calculated as the difference between T_m in the presence and absence of the compounds.

UV-vis studies

UV-visible spectra were run on an Agilent Cary 60 spectrophotometer (Agilent Technologies, Santa Clara, CA, USA) equipped with a Single-Cell Peltier temperature controller using a quartz cuvette of 10 mm path length. pSNCA and pSNCAext model oligonucleotides were diluted from stocks to final concentration (2.5 µM) in lithium cacodylate buffer (10 mM, pH 7.4) with 1 mM or 5 mM KCl. Ionic strength was kept constant by adding suitable quantities of LiCl (to 100 mM chloride salts). mSNCA was suspended in the same conditions but in the presence of 10 mM or 100 mM KCl. The solutions were annealed by heating at 95°C for 5 min and gradually cooling to room temperature over 4h. The absorbance was recorded as a function of temperature (20-95°C) from 220 to 340 nm wavelength intervals at a scan rate of 50 nm/min and a slit width of 0.5 nm. Mid-transition temperatures were determined following the absorbance decreasing at 295 nm in

the function of temperature (20-95°C). Data were the average of three independent experiments. Model-dependent thermodynamic parameters were estimated by following the protocol published by J.L. Mergny and collaborator [39]. In particular, the association constants (Ka) were calculated in function of the folded fractions ($\emptyset_T$), accordingly the equation $Ka = \emptyset_T/(1- \emptyset_T)$. Van't Hoff enthalpy and entropy values were estimated accordingly the equation $LogKa = (−\Delta H°/R) \times ((1/T) + \Delta S°/R)$, by restricting the temperature range for which folded fraction is $0.15 <\emptyset< 0.85$. R, the perfect gas constant, was considered equal to 1.987 calmol^{-1}K^{-1}. UV-vis thermal difference spectra have been determined by the mathematical subtraction of the spectra of fully unfolded oligonucleotides (above Tm, 95°C) and the spectra of fully folded oligonucleotides (low T, 20°C) [39]. TDS factors correspond to the average of $\Delta A_{240nm}/\Delta A_{295nm}$ absolute values obtained at 20, 30, 40, and 50 °C.

FRET-melting

FRET-melting measurements were recorded by an Agilent AriaMx Real-time PCR System (Agilent Technologies, Santa Clara, CA, USA) using 5'-FAM (6-carboxyfluorescein) and 3'-TAMRA (6-carboxy-tetramethylrhodamine) labeled oligonucleotides.[62] In a total volume of 60 µL, 0.25 µM of tagged oligonucleotides were dissolved in 10 mM lithium cacodylate buffer pH 7.4 in the presence of proper KCl/LiCl concentrations (suitable quantities of LiCl were added to maintain the total concentration of chloride salts equal to 100 mM). The mixtures were annealed by heating at 95 °C for 3 min followed by an equilibration step in ice for 30 min. The folded samples were stored at -4°C. Melting experiments were performed on 96 wells plates, by analyzing 20 µL of samples. After the first equilibration step at 25 °C for 5 minutes, a stepwise increase of 1°C/min was performed to reach 95°C, measuring the fluorescence emission at 516 nm (excitation at 462 nm) according to SYBR/FAM optical cartridge (Agilent). Data were the average of three

independent experiments, each conducted in triplicate conditions. G4-ligands were added in 1 µM concentration (4 equivalents) and left to equilibrate for 16 h at room temperature before recording the FRET-melting curve. The fluorescence emission curves were normalized between 0 and 1, and T_m was defined as the temperature corresponding to 0.5 normalized emissions. ΔT_m was calculated as the difference between T_m in the presence and absence of the compounds.

NMR studies

[1]H-NMR spectra were recorded on Bruker Avance NEO 700MHz (BRUKER, Billerica, MA, USA), equipped with a triple resonance helium-cooled cryoprobe 5 mm. In a total volume of 800 µL, 0.17 mM of oligonucleotides were dissolved in 90% H_2O/10% D_2O solution with 100 mM KCl and 20 mM potassium phosphate buffer at pH 7.0. The mixtures were annealed by heating at 95 °C for 5 min followed by an equilibration step in ice for 30 min. The folded samples were stored at -4°C. [1]H-NMR spectra were recorded at 25°C using a zgesgp technique to suppress the water signal. All NMR spectra were processed and analyzed using Bruker TopSpin 2.1.

Chemometric analysis on CD spectra

Principal Component Analysis (PCA) was performed using R software in which the Chemometric Agile Tool was loaded. PCA was initially applied to a reference dataset of 23 G4s CD spectra (7 parallels, 8 hybrids, and 8 antiparallels) reported in the literature,[35] by considering only the first two components PC1 and PC2. The variables were centered and scaled. Calculated scores from pSNCA and pSNCAext spectra (recorded at 2.5 µM oligonucleotides, 100 mM KCl, and 10 mM lithium cacodylate buffer, pH 7.4) were projected on the training set. The dataset was then utilized

to examine NLS-PNA-C343's impact on the pSNCAext structure (spectra recorded after the addition of 4 equivalents of PNA conjugate and an equilibration time of 4h).

SH-SY5Y culture and maintenance

SH-SY5Y cells were maintained in DMEM/F12 Complete Medium supplemented with 10% of Fetal Bovine Serum (FBS; Microtech), 1% glutamine (L-GLUT; GIBCO), and 1% Penicillin/Streptomycin (P/S, GIBCO) at 37 °C in a 5% CO_2. For all performed experiments, cells were differentiated with retinoic acid (RA) at a concentration of 10 μM for 7 days in a 1% FBS DMEM High Glucose medium to induce differentiation.

Chromatin Immunoprecipitation (ChIP)

ChIP G4-immunoprecipitation was performed on SH-SY5Y cells following a previously published protocol.[16] Cells were fixed in 1% formaldehyde for 10 minutes and quenched with 0.125 M glycine. Chromatin was sonicated using Bioruptor® plus sonicator (Diagenode) to shear to an average size of 1,000–1,500 bp and incubated in the presence of BG4 (Merck, Milan, Italy, MABE817, Lot: 3534080). Anti-FLAG antibody (Sigma-Aldrich, Milan, Italy, cat. F1804) coated Protein-G magnetic beads (Pierce™ ThermoFisher Scientific) were used to capture BG4-G4 complexes. After de-crosslinking, samples were purified with the ExpIn Combo GP mini kit (GeneAll). G4 enrichment was quantified via qPCR, using the Optimum QMastermix with SYBER GREEN (Genespin) and with the CFX Connect Real-Time PCR System (Bio-Rad). GAPDH was used as a negative control, primer sequences are reported in Table S6. Relative enrichments were calculated with respect to their inputs.

RNA Immunoprecipitation

RNA G4-immunoprecipitation was performed on SH-SY5Y cells. Cells were lysed with lysis buffer (10 mM Tris HCl, 10 mM NaCl, 2 mM EDTA, 0.5 Triton X100) and incubated on ice for 10 minutes. 120mM NaCl was then added and cells were incubated for a further 5 minutes. Cells were then pelleted at 16000 g for 15 minutes and the surnated was retained for pre-clearing. Following pre-clearing, the lysate was incubated in the presence of BG4 (Merck, Milan, Italy, MABE817, Lot: 3534080). Anti-FLAG antibody (Sigma-Aldrich, Milan, Italy, cat. F1804) coated Protein-G magnetic beads (Pierce™ ThermoFisher Scientific) were used to capture BG4-G4 complexes. After elution, samples were purified with TRIZOL® reagent (Thermo Fisher Scientific) following the manufacturer's instructions. Samples were retrotranscribed and analyzed via Real-Time PCR (see below). GAPDH was used as a negative control, primer sequences are reported in Table S6. Relative enrichments were calculated with respect to their inputs.

qPCR-Stop assay

qPCR stop assay was used to determine whether the putative genomic pSNCA region folds into a G4 structure.[63] Primers were designed to surround the G4 and increasing KCl concentrations (0, 0.1 mM, 1 mM, 10 mM, and 50 mM) were added to stabilize the G4 and thus assess Taq activity. Real-Time PCR was performed with the CFX Connect Real-Time PCR System (Bio-Rad) using the Optimum QMastermix with SYBER GREEN (Genespin). Primer sequences are reported in Table S6.

MTT assays

SH-SY5Y cells were seeded in a 96-well plate at 2×10^5 cells/well density and they were treated with increasing concentrations of HPHAM stabilizer. 10 μl MTT assay kit reagent (Merck) was then added to each well, and cells were incubated for 3 h. MTT crystals were eluted with 100 μl of elution solution, composed of 4 mM HCl, 0.1% (v/v) NP40 all in isopropanol for 30 min. The relative absorbance was measured with a Multiskan GO spectrophotometer (Thermo Fisher Scientific) at $\lambda = 570$ nm.

Immunocytochemistry

SH-SY5Y cells were seeded on ethanol-washed glass coverslips and cultured in a standard medium. Following treatment with HPHAM, cells were fixed with 4% paraformaldehyde in 0.1 M PBS (Thermo Fisher Scientific), pH 7.4, for 20 min at room temperature, and then washed with PBS. Nuclei were stained with DAPI at the final concentration of 1 μg/ml for 30 min. Glass coverslips were mounted using the FluorSave Reagent (Calbiochem) and analyzed by confocal microscopy (Confocal laser scanning microscopy platform Leica TCS SP8, Leica Microsystems). Co-localization rate was analyzed with Leica Analysis Software.

RNA extraction and real-time PCR

Total RNA was extracted using TRIZOL® reagent (Thermo Fisher Scientific) following the manufacturer's instructions and then quantified using the Multiskan GO spectrophotometer (Thermo Fisher Scientific). For stability assays, cells were co-treated with 1.5 μg/ml Actinomycin D. 1500 ng of RNA were retro-transcribed using the Super Mix 5X Full (Genespin) kit following the manufacturer's instructions. Real-Time PCR was performed with the CFX Connect Real-Time PCR System (Bio-Rad) using the Optimum QMastermix with SYBER GREEN (Genespin). The

NCBI's Primer-BLAST tool was used to design primers. Primer sequences are reported in Table S6. Gene expression was calculated using the $2^{-\Delta\Delta Ct}$ method. TBP was used as endogenous control.

Western blotting analysis

Protein extracts were obtained by means of RadioImmunoPrecipitation Assay (RIPA) lysis buffer, proteins were quantified with the Bradford Assay following a standard protocol (Coomassie Plus—The Better Bradford Assay™ Reagent, Thermo Scientific), and equal amounts of solubilized proteins were heated in Laemmli sample buffer (Bio-Rad) containing 70 mM 2-β-mercaptoethanol (Sigma Aldrich). Samples were separated in a 12% SDS-PAGE gel and electroblotted onto a nitrocellulose membrane (Trans-blot, Bio-Rad). Membranes were then fixed with 4% paraformaldehyde and 0.01% Glutaraldehyde in PBS (Thermo Fisher Scientific), for 30 min at room temperature, and then blocked in 5% slim milk (diluted in TBS with 0.05% Tween-20). After that, membranes were probed with the appropriate primary antibody: Recombinant Anti-Alpha-synuclein antibody [MJFR1] (1:2000 Abcam, ab138501) or β-Actin (1:5000 Sigma-Aldrich, A1978), overnight at 4°C. Lastly, membranes were incubated with the appropriate secondary antibody Peroxidase AffiniPure Goat Anti-Rabbit/Mouse IgG (1:5000 dilution; Jackson Immuno Research). Proteins were visualized by means of an enhanced chemiluminescence detection solution (Thermo Fisher Scientific). After acquisition by a GelDoc™ image capture system (Uvitec, Eppendorf), densitometric analysis of the bands was performed using the ImageJ software.

SUPPLEMENTARY INFORMATION

The online version contains supplementary material. %GC Content distribution; putative G-quadruplex sequences identified; estimation of the fractions of tertiary and secondary structural elements; melting data; thermodynamic parameters; qPCR stop assay for the negative control; CD spectra of pSNCAext/NLS-PNA-C343 hetero duplex; PCA score plot; MTT assay; immunofluorescence assay; real-time PCR of different housekeeping genes; analytic HPLC profiles; chemical structures of HPHAM, PDS, BRACO19, TMPyP4, and NLS-PNA-C343; name and nucleobases sequences of the DNA model oligonucleotides analyzed; LC-MS analysis of NLS-PNA-C343; List of primers used in this study.

AVAILABILITY OF DATA AND MATERIALS The datasets supporting the conclusions of this article are included within the article (and its additional file). Additional material or information can be provided upon request by contacting the corresponding author.

AUTHOR INFORMATION

* To whom correspondence should be addressed. Tel: +39 0382987668; Email: valentina.pirota@unipv.it; Correspondence may also be addressed to Stephana Carelli. Email: stephana.carelli@unimi.it

Author Contributions

† The authors wish it to be known that, in their opinion, the first two authors should be regarded as Joint First Authors.

‡ These authors contributed equally

Concept and design: V.P., S.C.; Acquisition and analysis: V.P., F.R., V.F., L.E., C.P., R.D.G.; Interpretation of data, V.P., F.R., V.F., M.M., P.G., M.F., S.C., C.C.; Drafting of the manuscript,

V.P., F.R.; Critical revision of the manuscript for important intellectual content: V.P., F.R., P.G., M.F., S.C.; Administrative, technical, or material support: V.P., F.D., M.F., S.C.; C.C.. All authors have given approval to the final version of the manuscript.

Funding Sources

Postdoctoral research fellowship (V.P.) DCHIMICA2020-A01 from University of Pavia**.**

ABBREVIATIONS

α-Syn : Alpha-synuclein; PD : Parkinson's disease; G4(s) : G-quadruplex(es); TSS : transcription start sites; UTR : untranslated region; NLS: nuclear localization sequence; PNA: peptide nucleic acid; TDS: thermal difference spectra; Tm: mid-transition temperature; C343: Coumarin 343; PCA: Principal Component Analysis.

REFERENCES

(1) Morato Torres, C. A.; Wassouf, Z.; Zafar, F.; Sastre, D.; Outeiro, T. F.; Schüle, B. The Role of Alpha-Synuclein and Other Parkinson's Genes in Neurodevelopmental and Neurodegenerative Disorders. *International Journal of Molecular Sciences* **2020**, *21* (16). DOI: 10.3390/ijms21165724.

(2) Burre, J.; Sharma, M.; Tsetsenis, T.; Buchman, V.; Etherton, M. R.; Sudhof, T. C. Alpha-synuclein promotes SNARE-complex assembly in vivo and in vitro. *Science* **2010**, *329* (5999), 1663-1667. DOI: 10.1126/science.1195227.

(3) Schaser, A. J.; Osterberg, V. R.; Dent, S. E.; Stackhouse, T. L.; Wakeham, C. M.; Boutros, S. W.; Weston, L. J.; Owen, N.; Weissman, T. A.; Luna, E.; et al. Alpha-synuclein is a DNA binding protein that modulates DNA repair with implications for Lewy body disorders. *Scientific Reports* **2019**, *9* (1), 10919. DOI: 10.1038/s41598-019-47227-z.

(4) Irwin, D. J.; Lee, V. M. Y.; Trojanowski, J. Q. Parkinson's disease dementia: convergence of α-synuclein, tau and amyloid-β pathologies. *Nature Reviews Neuroscience* **2013**, *14* (9), 626-636. DOI: 10.1038/nrn3549.

(5) Colombo, D.; Pnevmatikou, P.; Melloni, E.; Keywood, C. Therapeutic innovation in Parkinson's disease: a 2020 update on disease-modifying approaches. *Expert Rev Neurother* **2020**, 1-18. DOI: 10.1080/14737175.2020.1800454 .

(6) Fields, C. R.; Bengoa-Vergniory, N.; Wade-Martins, R. Targeting Alpha-Synuclein as a Therapy for Parkinson's Disease. *Frontiers in Molecular Neuroscience* **2019**, *12*, Review.

(7) Kantor, B.; Tagliafierro, L.; Gu, J.; Zamora, M. E.; Ilich, E.; Grenier, C.; Huang, Z. Y.; Murphy, S.; Chiba-Falek, O. Downregulation of SNCA Expression by Targeted Editing of DNA Methylation: A Potential Strategy for Precision Therapy in PD. *Mol Ther* **2018**, *26* (11), 2638-2649. DOI: 10.1016/j.ymthe.2018.08.019.

(8) Mittal, S.; Bjørnevik, K.; Im Doo, S.; Flierl, A.; Dong, X.; Locascio Joseph, J.; Abo Kristine, M.; Long, E.; Jin, M.; Xu, B.; et al. β2-Adrenoreceptor is a regulator of the α-synuclein gene driving risk of Parkinson's disease. *Science* **2017**, *357* (6354), 891-898. DOI: 10.1126/science.aaf3934.

(9) Nakamori, M.; Junn, E.; Mochizuki, H.; Mouradian, M. M. Nucleic Acid-Based Therapeutics for Parkinson's Disease. *Neurotherapeutics* **2019**, *16* (2), 287-298. DOI: 10.1007/s13311-019-00714-7 .

(10) Uehara, T.; Choong, C. J.; Nakamori, M.; Hayakawa, H.; Nishiyama, K.; Kasahara, Y.; Baba, K.; Nagata, T.; Yokota, T.; Tsuda, H.; et al. Amido-bridged nucleic acid (AmNA)-modified antisense oligonucleotides targeting α-synuclein as a novel therapy for Parkinson's disease. *Sci Rep* **2019**, *9* (1), 7567. DOI: 10.1038/s41598-019-43772-9.

(11) Cole, T. A.; Zhao, H.; Collier, T. J.; Sandoval, I.; Sortwell, C. E.; Steece-Collier, K.; Daley, B. F.; Booms, A.; Lipton, J.; Welch, M.; et al. α-Synuclein antisense oligonucleotides as a disease-modifying therapy for Parkinson's disease. *JCI Insight* **2021**, *6* (5). DOI: 10.1172/jci.insight.135633

(12) Alarcón-Arís, D.; Recasens, A.; Galofré, M.; Carballo-Carbajal, I.; Zacchi, N.; Ruiz-Bronchal, E.; Pavia-Collado, R.; Chica, R.; Ferrés-Coy, A.; Santos, M.; et al. Selective α-Synuclein Knockdown in Monoamine Neurons by Intranasal Oligonucleotide Delivery: Potential Therapy for Parkinson's Disease. *Mol Ther* **2018**, *26* (2), 550-567. DOI: 10.1016/j.ymthe.2017.11.015

(13) Zhang, P.; Park, H. J.; Zhang, J.; Junn, E.; Andrews, R. J.; Velagapudi, S. P.; Abegg, D.; Vishnu, K.; Costales, M. G.; Childs-Disney, J. L.; et al. Translation of the intrinsically disordered protein α-synuclein is inhibited by a small molecule targeting its structured mRNA. *Proc Natl Acad Sci U S A* **2020**, *117* (3), 1457-1467. DOI: 10.1073/pnas.1905057117

(14) Kuo, Y. M.; Nwankwo, E. I.; Nussbaum, R. L.; Rogers, J.; Maccecchini, M. L. Translational inhibition of α-synuclein by Posiphen normalizes distal colon motility in transgenic Parkinson mice. *Am J Neurodegener Dis* **2019**, *8* (1), 1-15.

(15) Spiegel, J.; Adhikari, S.; Balasubramanian, S. The Structure and Function of DNA G-Quadruplexes. *Trends Chem* **2020**, *2* (2), 123-136. DOI: 10.1016/j.trechm.2019.07.002

(16) Lago, S.; Nadai, M.; Ruggiero, E.; Tassinari, M.; Marušič, M.; Tosoni, B.; Frasson, I.; Cernilogar, F. M.; Pirota, V.; Doria, F.; et al. The MDM2 inducible promoter folds into four-tetrad antiparallel G-quadruplexes targetable to fight malignant liposarcoma. *Nucleic Acids Res* **2021**, *49* (2), 847-863. DOI: 10.1093/nar/gkaa1273

(17) Kharel, P.; Becker, G.; Tsvetkov, V.; Ivanov, P. Properties and biological impact of RNA G-quadruplexes: from order to turmoil and back. *Nucleic Acids Research* **2020**, *48* (22), 12534-12555. DOI: 10.1093/nar/gkaa1126

(18) Frasson, I.; Pirota, V.; Richter, S. N.; Doria, F. Multimeric G-quadruplexes: A review on their biological roles and targeting. *International Journal of Biological Macromolecules* **2022**, *204*, 89-102. DOI: https://doi.org/10.1016/j.ijbiomac.2022.01.197.

(19) Varshney, D.; Spiegel, J.; Zyner, K.; Tannahill, D.; Balasubramanian, S. The regulation and functions of DNA and RNA G-quadruplexes. *Nature Reviews Molecular Cell Biology* **2020**, *21* (8), 459-474. DOI: 10.1038/s41580-020-0236-x.

(20) Tian, T.; Chen, Y.-Q.; Wang, S.-R.; Zhou, X. G-Quadruplex: A Regulator of Gene Expression and Its Chemical Targeting. *Chem* **2018**, *4* (6), 1314-1344. DOI: https://doi.org/10.1016/j.chempr.2018.02.014.

(21) Robinson, J.; Raguseo, F.; Nuccio, S. P.; Liano, D.; Di Antonio, M. DNA G-quadruplex structures: more than simple roadblocks to transcription? *Nucleic Acids Research* **2021**, *49* (15), 8419-8431. DOI: 10.1093/nar/gkab609.

(22) Pirota, V.; Stasi, M.; Benassi, A.; Doria, F. Chapter Six - An overview of quadruplex ligands: Their common features and chemotype diversity. In *Annual Reports in Medicinal Chemistry*, Neidle, S. Ed.; Vol. 54; Academic Press, 2020; pp 163-196.

(23) Pirota, V.; Lunghi, E.; Benassi, A.; Crespan, E.; Freccero, M.; Doria, F. Selective Binding and Redox-Activity on Parallel G-Quadruplexes by Pegylated Naphthalene Diimide-Copper Complexes. *Molecules* **2021**, *26* (16). DOI: 10.3390/molecules26165025.

(24) Tassinari, M.; Zuffo, M.; Nadai, M.; Pirota, V.; Sevilla Montalvo, A. C.; Doria, F.; Freccero, M.; Richter, S. N. Selective targeting of mutually exclusive DNA G-quadruplexes: HIV-1 LTR as paradigmatic model. *Nucleic Acids Res* **2020**, *48* (9), 4627-4642. DOI: 10.1093/nar/gkaa186

(25) Cebrián, R.; Belmonte-Reche, E.; Pirota, V.; de Jong, A.; Morales, J. C.; Freccero, M.; Doria, F.; Kuipers, O. P. G-Quadruplex DNA as a Target in Pathogenic Bacteria: Efficacy of an Extended Naphthalene Diimide Ligand and Its Mode of Action. *Journal of Medicinal Chemistry* **2021**. DOI: 10.1021/acs.jmedchem.1c01905.

(26) Wang, E.; Thombre, R.; Shah, Y.; Latanich, R.; Wang, J. G-Quadruplexes as pathogenic drivers in neurodegenerative disorders. *Nucleic Acids Research* **2021**, *49* (9), 4816-4830. DOI: 10.1093/nar/gkab164

(27) Pirota, V.; Nadai, M.; Doria, F.; Richter, S. N. Naphthalene Diimides as Multimodal G-Quadruplex-Selective Ligands. *Molecules* **2019**, *24* (3). DOI: 10.3390/molecules24030426

(28) Liu, L.-Y.; Ma, T.-Z.; Zeng, Y.-L.; Liu, W.; Mao, Z.-W. Structural Basis of Pyridostatin and Its Derivatives Specifically Binding to G-Quadruplexes. *Journal of the American Chemical Society* **2022**, *144* (26), 11878-11887. DOI: 10.1021/jacs.2c04775.

(29) Kikin, O.; D'Antonio, L.; Bagga, P. S. QGRS Mapper: a web-based server for predicting G-quadruplexes in nucleotide sequences. *Nucleic Acids Res* **2006**, *34* (Web Server issue), W676-682. DOI: 10.1093/nar/gkl253

(30) Bedrat, A.; Lacroix, L.; Mergny, J. L. Re-evaluation of G-quadruplex propensity with G4Hunter. *Nucleic Acids Res* **2016**, *44* (4), 1746-1759. DOI: 10.1093/nar/gkw006

(31) Dhapola, P.; Chowdhury, S. QuadBase2: web server for multiplexed guanine quadruplex mining and visualization. *Nucleic acids research* **2016**, *44* (W1), W277-W283. DOI: 10.1093/nar/gkw425

(32) Ozansoy, M.; Başak, A. N. The Central Theme of Parkinson's Disease: α-Synuclein. *Molecular Neurobiology* **2013**, *47* (2), 460-465. DOI: 10.1007/s12035-012-8369-3.

(33) Patel, D. J.; Phan, A. T.; Kuryavyi, V. Human telomere, oncogenic promoter and 5'-UTR G-quadruplexes: diverse higher order DNA and RNA targets for cancer therapeutics. *Nucleic Acids Res* **2007**, *35* (22), 7429-7455. DOI: 10.1093/nar/gkm711

(34) Koukouraki, P.; Doxakis, E. Constitutive translation of human α-synuclein is mediated by the 5'-untranslated region. *Open Biol* **2016**, *6* (4), 160022. DOI: 10.1098/rsob.160022

(35) Del Villar-Guerra, R.; Trent, J. O.; Chaires, J. B. G-Quadruplex Secondary Structure Obtained from Circular Dichroism Spectroscopy. *Angew Chem Int Ed Engl* **2018**, *57* (24), 7171-7175. DOI: 10.1002/anie.201709184

(36) Kejnovská, I.; Renčiuk, D.; Palacký, J.; Vorlíčková, M. CD Study of the G-Quadruplex Conformation. In *G-Quadruplex Nucleic Acids: Methods and Protocols*, Yang, D., Lin, C. Eds.; Springer New York, 2019; pp 25-44.

(37) Zacchia, M.; Abategiovanni, M. L.; Stratigis, S.; Capasso, G. Potassium: From Physiology to Clinical Implications. *Kidney Dis (Basel)* **2016**, *2* (2), 72-79. DOI: 10.1159/000446268

(38) Nadai, M.; Doria, F.; Scalabrin, M.; Pirota, V.; Grande, V.; Bergamaschi, G.; Amendola, V.; Winnerdy, F. R.; Phan, A. T.; Richter, S. N.; et al. A Catalytic and Selective Scissoring Molecular Tool for Quadruplex Nucleic Acids. *Journal of the American Chemical Society* **2018**, *140* (44), 14528-14532. DOI: 10.1021/jacs.8b05337.

(39) Mergny, J. L.; Lacroix, L. UV Melting of G-Quadruplexes. *Curr Protoc Nucleic Acid Chem* **2009**, *Chapter 17*, Unit 17.11. DOI: 10.1002/0471142700.nc1701s37

(40) Lane, A. N.; Chaires, J. B.; Gray, R. D.; Trent, J. O. Stability and kinetics of G-quadruplex structures. *Nucleic Acids Res* **2008**, *36* (17), 5482-5515. DOI: 10.1093/nar/gkn517

(41) Karsisiotis, A. I.; Hessari, N. M. a.; Novellino, E.; Spada, G. P.; Randazzo, A.; Webba da Silva, M. Topological Characterization of Nucleic Acid G-Quadruplexes by UV Absorption and Circular Dichroism. *Angewandte Chemie International Edition* **2011**, *50* (45), 10645-10648, https://doi.org/10.1002/anie.201105193.

(42) De Cian, A.; Guittat, L.; Kaiser, M.; Sacca, B.; Amrane, S.; Bourdoncle, A.; Alberti, P.; Teulade-Fichou, M. P.; Lacroix, L.; Mergny, J. L. Fluorescence-based melting assays for studying quadruplex ligands. *Methods* **2007**, *42* (2), 183-195. DOI: 10.1016/j.ymeth.2006.10.004

(43) Adrian, M.; Heddi, B.; Phan, A. T. NMR spectroscopy of G-quadruplexes. *Methods* **2012**, *57* (1), 11-24. DOI: https://doi.org/10.1016/j.ymeth.2012.05.003.

(44) Lammich, S.; Kamp, F.; Wagner, J.; Nuscher, B.; Zilow, S.; Ludwig, A.-K.; Willem, M.; Haass, C. Translational Repression of the Disintegrin and Metalloprotease ADAM10 by a Stable G-quadruplex Secondary Structure in Its 5′-Untranslated Region*. *Journal of Biological Chemistry* **2011**, *286* (52), 45063-45072. DOI: https://doi.org/10.1074/jbc.M111.296921.

(45) von Hacht, A.; Seifert, O.; Menger, M.; Schütze, T.; Arora, A.; Konthur, Z.; Neubauer, P.; Wagner, A.; Weise, C.; Kurreck, J. Identification and characterization of RNA guanine-quadruplex binding proteins. *Nucleic Acids Res* **2014**, *42* (10), 6630-6644. DOI: 10.1093/nar/gku290

(46) Xicoy, H.; Wieringa, B.; Martens, G. J. M. The SH-SY5Y cell line in Parkinson's disease research: a systematic review. *Molecular Neurodegeneration* **2017**, *12* (1), 10. DOI: 10.1186/s13024-017-0149-0.

(47) Biffi, G.; Tannahill, D.; McCafferty, J.; Balasubramanian, S. Quantitative visualization of DNA G-quadruplex structures in human cells. *Nat Chem* **2013**, *5* (3), 182-186. DOI: 10.1038/nchem.1548

(48) Joshi, S.; Singh, A.; Kukreti, S. Porphyrin induced structural destabilization of a parallel DNA G-quadruplex in human MRP1 gene promoter. *Journal of Molecular Recognition* **2022**, *35* (3), e2950, https://doi.org/10.1002/jmr.2950.

(49) Pellestor, F.; Paulasova, P. The peptide nucleic acids (PNAs), powerful tools for molecular genetics and cytogenetics. *European Journal of Human Genetics* **2004**, *12* (9), 694-700. DOI: 10.1038/sj.ejhg.5201226.

(50) Swenson, C. S.; Lackey, H. H.; Reece, E. J.; Harris, J. M.; Heemstra, J. M.; Peterson, E. M. Evaluating the effect of ionic strength on PNA:DNA duplex formation kinetics. *RSC Chemical Biology* **2021**, *2* (4), 1249-1256, 10.1039/D1CB00025J. DOI: 10.1039/D1CB00025J.

(51) Green, J. J.; Ying, L.; Klenerman, D.; Balasubramanian, S. Kinetics of Unfolding the Human Telomeric DNA Quadruplex Using a PNA Trap. *Journal of the American Chemical Society* **2003**, *125* (13), 3763-3767. DOI: 10.1021/ja029149w.

(52) Komiyama, M.; Ye, S.; Liang; Yamamoto, Y.; Tomita, T.; Zhou, J.-M.; Aburatani, H. PNA for One-Base Differentiating Protection of DNA from Nuclease and Its Use for SNPs Detection. *Journal of the American Chemical Society* **2003**, *125* (13), 3758-3762. DOI: 10.1021/ja0295220.

(53) Rey, F.; Pandini, C.; Messa, L.; Launi, R.; Barzaghini, B.; Zangaglia, R.; Raimondi, M. T.; Gagliardi, S.; Cereda, C.; Zuccotti, G. V.; et al. α-Synuclein antisense transcript SNCA-AS1 regulates synapses- and aging-related genes suggesting its implication in Parkinson's disease. *Aging Cell* **2021**, *20* (12), e13504. DOI: 10.1111/acel.13504

(54) Kalinovskii, A. P.; Osmakov, D. I.; Koshelev, S. G.; Lubova, K. I.; Korolkova, Y. V.; Kozlov, S. A.; Andreev, Y. A. Retinoic Acid-Differentiated Neuroblastoma SH-SY5Y Is an Accessible In Vitro Model to Study Native Human Acid-Sensing Ion Channels 1a (ASIC1a). *Biology (Basel)* **2022**, *11* (2). DOI: 10.3390/biology11020167

(55) Moruno-Manchon, J. F.; Koellhoffer, E. C.; Gopakumar, J.; Hambarde, S.; Kim, N.; McCullough, L. D.; Tsvetkov, A. S. The G-quadruplex DNA stabilizing drug pyridostatin promotes DNA damage and downregulates transcription of Brca1 in neurons. *Aging (Albany NY)* **2017**, *9* (9), 1957-1970. DOI: 10.18632/aging.101282

(56) Yang, J.; Qin, G.; Niu, J.; Wei, Y.; Li, X.; Zhao, C.; Wang, C.; Ren, J.; Qu, X. Targeting G-quadruplexes in an ageing epigenetic regulator promoter for rescuing mitochondrial dysfunction in Alzheimer's disease. *Chemical Communications* **2023**, *59* (8), 1078-1081, DOI: 10.1039/D2CC05957F.

(57) Balasubramanian, S.; Hurley, L. H.; Neidle, S. Targeting G-quadruplexes in gene promoters: a novel anticancer strategy? *Nat Rev Drug Discov* **2011**, *10* (4), 261-275. DOI: 10.1038/nrd3428

(58) Zhang, A.; Zhao, J. C.; Kim, J.; Fong, K. W.; Yang, Y. A.; Chakravarti, D.; Mo, Y. Y.; Yu, J. LncRNA HOTAIR Enhances the Androgen-Receptor-Mediated Transcriptional Program and Drives Castration-Resistant Prostate Cancer. *Cell Rep* **2015**, *13* (1), 209-221. DOI: 10.1016/j.celrep.2015.08.069

(59) Doria, F.; Nadai, M.; Sattin, G.; Pasotti, L.; Richter, S. N.; Freccero, M. Water soluble extended naphthalene diimides as pH fluorescent sensors and G-quadruplex ligands. *Organic & Biomolecular Chemistry* **2012**, *10* (19), 3830-3840, DOI: 10.1039/C2OB07006E.

(60) Rodriguez, R.; Miller, K. M.; Forment, J. V.; Bradshaw, C. R.; Nikan, M.; Britton, S.; Oelschlaegel, T.; Xhemalce, B.; Balasubramanian, S.; Jackson, S. P. Small-molecule-induced DNA damage identifies alternative DNA structures in human genes. *Nat Chem Biol* **2012**, *8* (3), 301-310. DOI: 10.1038/nchembio.780

(61) Liu, S.; Wang, W.-H.; Dang, Y.-L.; Fu, Y.; Sang, R. Rational design and efficient synthesis of a fluorescent-labeled jasmonate. *Tetrahedron Letters* **2012**, *53* (32), 4235-4239. DOI: https://doi.org/10.1016/j.tetlet.2012.06.006.

(62) Wang, K.; Flaherty, D. P.; Chen, L.; Yang, D. High-Throughput Screening of G-Quadruplex Ligands by FRET Assay. In *G-Quadruplex Nucleic Acids: Methods and Protocols*, Yang, D., Lin, C. Eds.; Springer New York, 2019; pp 323-331.

(63) Jamroskovic, J.; Obi, I.; Movahedi, A.; Chand, K.; Chorell, E.; Sabouri, N. Identification of putative G-quadruplex DNA structures in S. pombe genome by quantitative PCR stop assay. *DNA Repair* **2019**, *82*, 102678. DOI: https://doi.org/10.1016/j.dnarep.2019.102678.