1   A screen of mutants generated and imaged by the International Mouse Phenotyping Consortium
2   identifies 39 novel genes regulating the development of the secondary palate.

10   Ashwin Bhaskar[1] and Sophie Astrof[2, *]
11
12
13
14   1. Rutgers University, School of Arts and Sciences Honors Program, New Brunswick, NJ,
15   08901, USA
16

17   2. Department of Cell Biology and Molecular Medicine, Cardiovascular Research Institute,

18   Rutgers Biomedical and Health Sciences, 185 South Orange Ave, Newark, NJ, 07103, USA.

21   *Corresponding author:

22   E-mail: sophie.astrof@rutgers.edu

Abstract

The International Mouse Phenotyping Consortium (IMPC) has generated thousands of knockout mouse lines, of which a large proportion is embryonic or early neonatal lethal. The IMPC has generated and imaged embryos from lethal lines and made the three-dimensional image data sets publicly available. In this study, we used this resource to screen homozygous null mutants for defects in the development of the secondary palate. Altogether, we analyzed optical sections from 3216 embryos isolated at embryonic day (E) 15.5 and E18.5 from 478 homozygous mutant lines. Through this analysis, we discovered 39 novel genes important for palatal development. These studies provide new insights into the molecular regulation of palatogenesis and craniofacial disease and offer a useful resource for future exploration.

Introduction

Cleft palate with or without a cleft lip (known as orofacial clefts) is the most prevalent craniofacial birth defect, affecting approximately 1 in 800 babies each year (Rahimov et al., 2012). There is a significant difference in the incidence of orofacial clefting among different ethnic groups: the greatest incidence was found among Native Americans at 3.6 per 1000 births, followed by Japanese and Chinese populations, with a prevalence of 2.1 and 1.7 per 1000 births, respectively, while the lowest incidence is among African-Americans, at approximately 0.3 per 1000 births (Wojcicki et al., 2016). Children born with this disorder have difficulties with feeding and speech when untreated. In developed countries, orofacial clefts are surgically treated early in childhood to close the clefts (Wojcicki et al., 2016). Despite the treatment, some affected individuals may still experience negative psychological, psychosocial, and financial hardships (Wehby and Cassell, 2010).

Palate formation proceeds in multiple phases (Hammond and Dixon, 2022; Som and Naidich, 2013). During the early stages of palate development, multi-potent neural crest (NC) cells migrate to populate the first branchial arch, which subsequently gives rise to the maxillary and mandibular prominences. The secondary palate, the tissue that separates the nasal and oral cavities, forms when the paired outgrowths from the maxillary processes fuse at the midline and then with the primary palate, which is derived from the frontonasal prominence. In humans, palatal development takes place between weeks 4 and 10 of gestation (Mossey et al., 2009; Som and Naidich, 2013). Due to the distinct developmental processes underlying the formation of the primary and secondary palate, several types of clefting are observed in patients and vertebrate model organisms: isolated cleft lip, isolated cleft palate, or clefting can be present both in the lip and palate (Mossey et al., 2009; Selleri and Rijli, 2023).

Mice are an established model for studying palate formation and clefting due to similarities in palate development between mice and humans (Selleri and Rijli, 2023). In mice, the growth of paired palatal shelves from the maxillary processes can be observed at embryonic day (E) 11.5. These processes undergo a dramatic increase in size through the proliferation of NC-derived mesenchyme and initially move downwards to be positioned on either side of the tongue by E13.5. The paired palatal shelves then elevate to become positioned horizontally above the tongue by E14.5. The fusion of palatal shelves at the midline begins soon afterward and results in the formation of a continuous shelf of the secondary palate. The fusion of the

secondary palatal shelves is completed by E15.5, and the fusion of the primary and secondary

palatal shelves is typically completed by E17.5 (Bush and Jiang, 2012).

The causes of cleft lip and palate in the human population are not completely understood.

Epidemiological studies, genetic analyses, and modeling of lip and palate clefting in mice

indicated that both environmental factors and genetic mutations are at play (Hammond and

Dixon, 2022; Mossey et al., 2009; Selleri and Rijli, 2023). Among common mechanisms

mediating palatal development in humans and mice are those regulated by Sonic Hedgehog

(Shh) signaling (Cohen, 2010; Mansilla et al., 2006). Modeling studies in mice indicated that

precise levels of Shh gene expression in the NC-derived cells are essential for proper palate

development, and both downregulation and upregulation of Shh in the NC-derived cells cause

clefting and other defects in facial development (Jeong et al., 2004; Li et al., 2017). Additional

players important for palate development include genes encoding secreted growth factors such as

BMPs, Wnts, and Fgfs, and transcriptional regulators such as the Pbx family of transcription

factors and *Snail1* (Ferretti et al., 2011; Losa et al., 2018; Welsh et al., 2018). However, much

still needs to be learned. Therefore, it is critical to identify additional genes and their roles in

palate development to broaden our understanding of mechanisms leading to the clefting of the lip

and palate and to create more effective prognostic, diagnostic, and treatment options. To discover

additional genes regulating the formation of the secondary palate, we examined images of mouse

mutants generated and imaged by the International Mouse Phenotyping Consortium (IMPC)

using micro-computed tomography (micro-CT) and looked for the presence of clefting in the

secondary palate at E15.5 and E18.5 in coronal optical sections.

Materials and Methods

Image Analysis: To identify novel genes regulating the development of the secondary palate, we

screened E15.5 and E18.5 homozygous-null mutant embryos generated and imaged by the

IMPC: https://www.mousephenotype.org). Coronal optical sections of micro-CT images were

analyzed using the tools embedded within the IMPC website in order to identify embryos with

defective palates. Embryos at E18.5 were preferred for analysis due to the palate being fully

fused at this stage in wild-type mouse embryos. E15.5 embryos were also analyzed. In E15.5

controls, the anterior-most and the posterior palate was often not yet fused. Therefore, cleft

palate was only marked in those E15.5 mutants in which the palatal shelves were not fused

throughout the entire length of the palate.

Statistical Analysis: According to the Center for Disease Control, the incidence of cleft palate in

the human population is 1 in 1700 (Mai et al., 2019). Of 100 E18.5 wild-type embryos examined

in this study, only one had a cleft palate (wild-type embryo ID: X63574-04). Using the binomial

probability formula, we obtained a value of p=0.0554948102 (rounded up to 0.06 or 6%), which

is the incidence of cleft palate expected solely by chance. Among the identified novel mutants

with cleft palate, homozygous-null mutants in the *Spop* gene exhibited the smallest penetrance of

cleft palate; only 1 of 11 *Spop*-null embryos had cleft palate. The binomial probability of

observing 1 cleft palate out of 11 individuals is 0.007, indicating that the presence of a cleft

palate was unlikely by chance alone, even at such low penetrance. In fact, the cutoff would be 1

cleft palate in 80 embryos (p=0.047).

Identification of novel cleft palate genes: To find which of the cleft palate mutations were

novel, we determined Mammalian Phenotype ontology terms associated with each gene on our

list (Supplemental Table 1). To do so, we queried the list of genes in Supplemental Table 1

against mammalian phenotype terms (https://www.informatics.jax.org/batch). In addition, we

compared the genes on our list with those in a cleft gene database (https://bioinfo.uth.edu).

Genes not previously identified as playing a role in palatal development were considered novel

(Supplemental Table 1, red ink).

Pathway enrichment analyses: We used Metascape (Zhou et al., 2019) to identify enriched

biological processes and pathways (Supplemental Table 2).

Results and discussion

123   We examined the palatal morphology using coronal optical sections in 3216 embryos

from 478 homozygous mutant lines that were generated by the IMPC and imaged using micro-

CT by the IMPC (Table 3). Out of 478 lines, 61 lines displayed cleft palate (Table 1). To

determine which lines represented novel cleft palate genes, we used gene ontology terms and the

Mouse Genome Informatics (MGI) along with the cleft palate database

(https://bioinfo.uth.edu/CleftGeneDB/). These analyses indicated that of 61 genes, mutations in

39 were not previously associated with cleft palate; therefore, these 39 genes were marked as

novel cleft palate genes (Supplemental Table 1, marked in red ink, and Figures 1-2). 38 of the 39

genes were not previously associated with cleft palate phenotypes according to both the MGI and

the CleftGeneDB databases. The *Col27a1* gene was associated with the cleft palate MP term in

the MGI database but not in the CleftGeneDB database. Mutant *Col27a1* alleles were generated

independently of the IMPC; however, palate development was not examined in these mice

(Plumb et al., 2011). And even though mutations in *Col27a1* were found in some patients with

Steel syndrome whose symptoms included cleft palate (Girisha et al., 2022), no studies directly

linked mutations in *Col27a1* to cleft palate. Therefore, we considered *Col27a1* a novel cleft

palate gene. A gene that was initially considered a novel cleft palate gene but ultimately not

considered novel in our analysis was Bmi1. Bmi1 was not associated with cleft palate MP terms

in the MGI database, but it was present in the CleftGeneDB. Upon further literature review, we

found a statement, albeit without supporting data, that 2 of 6 Bmi1-null embryos had cleft palate

(Akasaka et al., 2001), and therefore, we excluded Bmi1 from our list of novel cleft palate genes.

Using the list of novel cleft palate genes as an input, we performed pathway enrichment analysis

using Metascape (Zhou et al., 2019), which utilizes ontology resources including *Kegg Pathway,*

Reactome Gene Sets, GO Biological Processes, Canonical Pathways, CORUM, and

Wikipathways in order to discover enriched pathways regulated by cleft palate genes identified in

our screen.

Homozygous mutants in *Arhgef18, Memo1,* and *Tctn3* exhibited a nearly 100% incidence

of cleft palate at E18.5 (Fig. 1, Supplemental Table 1). Arhgef18 is RhoA guanine nucleotide

exchange factor (GEF), also known as p114RhoGEF, that regulates RhoA activation and

actomyosin contractility in multiple contexts (Beal et al., 2021; Nakajima and Tanoue, 2011;

Tsuji et al., 2010; Xu et al., 2013). Arhgef18, thus, may be important for the fusion of the palatal

shelves. As the two palatal shelves contact each other, the ectodermal epithelial cells, a.k.a.

medial edge epithelial layers (MEE) that border the palatal mesenchyme, intercalate to form the

midline epithelial seam (MES) (Lan et al., 2015). Activation of Rho and actomyosin contractility

are essential for this intercalation and cell extrusion to form a continuous palatal tissue (Kim et

al., 2015; Rosenblatt et al., 2001), and Arhgef18 could be an essential activator of RhoA in this

process. Future conditional mutagenesis studies are necessary to test this hypothesis and to

determine the precise tissues in which Arhgef18 functions during palatal development. This is

especially important since recent studies found mutations in Arhgef18 in patients with sporadic

cleft palate (El-Sibai et al., 2021).

162   Interestingly, Memo1 also regulates RhoA signaling, cytoskeletal dynamics, and cell

migration (Marone et al., 2004; Zaoui et al., 2008), and thus, could regulate palatal fusion by

mediating cell intercalation and extrusion behaviors leading to MES formation and dissolution.

The identification of *Arhgef18* and *Memo1* as important regulators of palatal development

provides novel molecular players to explore in the future and gain additional insights into the

regulation of MES formation and dissolution, which are relatively understudied but essential

processes for palatal morphogenesis.

169   A homozygous null mutation in *Tctn3* (tectonic family member 3) also resulted in

penetrant cleft palate (Fig. 1 and Supplemental Table 1). In humans, mutations in *Tctn*3 are

associated with the Mohr-Majewski Syndrome, and cleft palate was found in one case (Thomas

et al., 2012). However, palatal development was not examined in Tctn3-null mutants generated

prior to the IMPC screen (Wang et al., 2017); therefore, it was not clear whether Tctn3 was

required for the development of the palate. Tctn3 is a membrane protein that localizes to the

primary cilium and is essential for the activation of the Shh signaling (Reiter and Skarnes, 2006;

Wang et al., 2017). Ablation of Tctn3 resulted in defective ciliogenesis (Wang et al., 2017).

Interestingly, ciliopathies (defects resulting in dysfunctional primary cilia) are common causes of

craniofacial defects and underly a number of craniofacial syndromes, such as Joubert syndrome

[OMIM 213300] and Meckel-Gruber syndrome [OMIM 249000], Ellis-van Creveld syndrome

[OMIM 225500], Oral–facial–digital syndromes types IX [OMIM 258865] and XIV (Adly et al.,

2014; Thauvin-Robinet et al., 2014), and Weyers acrofacial dysostosis [OMIM 193530] (Moore,

2022; Reynolds et al., 2020). Consistent with this, our gene set enrichment analysis (GSEA)

showed that "Genes related to primary cilium" category was a statistically enriched category in

our dataset (q=0.04624) and included the following genes: *Tctn3, Cep97, Inpp5e, Fuz*, and

Clasp1 (Supplemental Table 2), among which *Tctn3, Cep97*, and *Clasp1* are novel cleft palate

genes identified in our screen (Figs. 1-2). Cep97 localizes at the centrioles and regulates cilia

assembly (Spektor et al., 2007). Clasp1 is a plus-end microtubule-binding protein regulating the

microtubule dynamics (Akhmanova et al., 2001; Maiato et al., 2005), an essential process for the

biogenesis of cilia (Malicki and Johnson, 2017). Thus, our screen identified two additional cilia
genes (*Cep97* and *Clasp1*) with roles in microtubule dynamics and palate development.

Perhaps not surprisingly, "Embryonic organ development" and "Skeletal system
development" were the top two enriched categories (q=2.477E-4) of our GSEA. Among the 13
genes that fell into these 2 categories, 2 genes, *Celf1 and Col27a1*, were not previously known to
regulate palate development. Our phenotyping analyses indicated that 1 of 6 homozygous *Celf1*-
null mutants imaged by the IMPC exhibited a cleft palate (Fig. 1). The binomial probability of
cleft palate in this strain is p<0.004, which indicates that cleft palate did not occur simply by
chance. *Celf1* (CUGBP Elav-like family member 1) is an RNA-binding protein regulating
diverse aspects of mRNA biology, including mRNA alternative splicing, stability, and miRNA
processing (Vlasova-St Louis et al., 2013). *Celf1* knockout animals were previously generated on
a mixed C57Bl6-129Sv background, and although ~50% of the knockout animals did not survive
past day 10 after birth, palatal development was not examined (Kress et al., 2007). Celf1 protein
binds GU-rich sequences in introns as well as in 5'- and 3'-UTRs of mRNAs. The role of Celf1
in palate development could be related to its ability to promote epithelial-to-mesenchymal
transition (EMT) due to its ability to bind to and increase translation of mRNAs encoding EMT
regulators (Chaudhury et al., 2016). Among these regulators is *Snail1*, which is important for
palate development (Losa et al., 2018; Murray et al., 2007). Thus, *Celf1* could be another novel
player in regulating EMT during the dissolution of the MES.

Col27a1 (a.k.a. type XXVII collagen) is an evolutionarily conserved member of extracellular
matrix proteins that belongs to a family of fibrillar collagens (Boot-Handford et al., 2003). The
high conservation in vertebrates, the unusual domain composition relative to other fibrillar
collagens, and the developmentally restricted expression of Col27a1 suggested that this protein
plays important developmental roles (Plumb et al., 2007). Mutations in the *Col27a1* genes have
been associated with Steel syndrome, a rare autosomal recessive disease of the skeleton (Girisha
et al., 2022; Gonzaga-Jauregui et al., 2020; Kotabagi et al., 2017). Among these patients, a pair
of twin brothers was also diagnosed with cleft palate (Girisha et al., 2022). However, no other
patients with *Col27a1* mutations or mouse mutants in *Col27a1* had palatal clefting (Plumb et al.,
2011). Although only one of eight *Col27a1*-null mutants exhibited cleft palate (**Fig. 2**), this
incidence is significant, p<0.005, excluding that cleft palate occurred purely by chance in this

strain. However, the low incidence of cleft palate in *Col27a1*-null mutants may complicate

further studies to understand the role of this protein in palatal shelf elevation or fusion.

"Melanoma" pathway (*CDK4, FGF9, FGF10, FGF19*) was also a statistically significant

enriched category of genes (q=0.02398) in our dataset. Melanocytes are derived from the neural

crest and melanomas are highly proliferative metastatic tumors (Mort et al., 2015). This is

consistent with the literature indicating that control of cell proliferation is essential during palate

development (Hammond and Dixon, 2022). Among genes in this category, *Cdk4* is a novel

regulator of palatal development, with two of eight *Cdk4*-null mutants displaying cleft palate in

our screen (Fig. 2).

Since the neural crest gives rise to the palatal mesenchyme and craniofacial bones and

cartilage, it was consistent that pathways related to neural crest development were significantly

enriched in our dataset. These pathways were: "Ear development (q = 0.03899)", "sensory organ

development (q = 0.005662)", "digestive system development (q=0.021777)", and "roof of

mouth development (q = 0.03882)". The enrichment in the "Ear Development" pathway was

especially intriguing as it contained the greatest number of novel cleft palate genes identified in

our screen (Supplemental Table 2). 22 genes from our screen comprised this category, and 8 of

them had not been associated with palate development before our work. These genes are *MAF,*

CDK4, RUNX1T1, PODXL, COL27A1, CELF1, TMEM115, UGT1A6. *Maf* encodes a group of

transcription factors that can act as transcriptional activators or repressors. Of the four MAFs

(*MAFA, MAFB, c-MAF, NRL*), *MAFB* is essential for the formation of inner ears, among its other

functions, such as the development of pancreatic endocrine cells (Takahashi, 2021). *Runx1t1*

encodes a transcriptional corepressor implicated in the positive regulation of angiogenesis (Liao

et al., 2017). In addition, Runx1t1 is commonly mutated in acute myeloid leukemia through its

fusion with Runx1 (Grinev et al., 2021). Curiously, there was a case study reporting a mutation

in Runx1t1 in a patient with a protruding ear (Huynh et al., 2012). The enrichment in the "Ear

development" category in our dataset suggests that screening for ear anomalies and hearing loss

may be reasonable in patients with cleft palate.

The statistically significant enrichment in the "Kidney development" category

(q=0.03548) was surprising mainly because the set of genes in this category did not include cilia

genes (*AP2B1, CDKN1C, EDNRA, FGF10, PODXL, PYGO2*), since defects in cilia development

contribute both to craniofacial and kidney disorders (McConnachie et al., 2021; Moore, 2022).

Published studies indicated that *Pygo2,* a component of canonical Wnt signaling required for the

development of both the palate and the kidney (Schwab et al., 2007), although the mechanism by

which palate development is regulated by *Pygo2* mutants is still unknown. The enrichment in

this category suggests the existence of common regulatory mechanisms mediating the

development of the kidney and the palate in addition to those regulated by cilia and suggests that

patients presenting with cleft palate may need to be followed up for anomalies in other organ

systems, such as the sensory nervous system, the digestive system, and the ear, as well as in a

seemingly unexpected organ, such as the kidney. Analyses of additional IMPC mutants in the

future will provide important insights into the mechanisms of palatal development and identify

genes and additional pathways important in this process and disease. Our screen of IMPC

mutants identified 39 novel cleft palate genes, providing fertile grounds for more mechanistic

studies in the future. Another value of our analyses lies in identifying additional and

unanticipated organs/pathways that could be affected alongside the cleft palate in patients,

allowing a more thorough clinical follow-up and treatment of all associated symptoms.

Figure Legends:

Figure 1. E18.5 mutants with cleft palate. Snapshots of coronal optical sections from micro-

CT images were taken at eye level. Eyes were marked by arrowheads. Long arrows point at

palatal shelves. Short arrows point at the tongue. Supplemental Table 4 contains additional

information about the displayed embryos. All scale bars are 1 mm.

Figure 2. E15.5 mutants with cleft palate. Snapshots of coronal optical sections from micro-

CT images were taken at eye level. Eyes were marked by arrowheads. Long arrows point at

palatal shelves. Short arrows point at the tongue. Supplemental Table 5 contains additional

information about the displayed embryos. All scale bars are 1 mm.

Acknowledgments

These studies were supported by the National Heart, Lung, and Blood Institute of the National

Institutes of Health (R01 HL103920, R01 HL134935, and R01 HL158049) to SA.

References:

Adly, N., Alhashem, A., Ammari, A., Alkuraya, F.S., 2014. Ciliary genes TBC1D32/C6orf170
and SCLT1 are mutated in patients with OFD type IX. Hum Mutat 35, 36-40.

Akasaka, T., van Lohuizen, M., van der Lugt, N., Mizutani-Koseki, Y., Kanno, M., Taniguchi,
M., Vidal, M., Alkema, M., Berns, A., Koseki, H., 2001. Mice doubly deficient for the Polycomb
Group genes Mel18 and Bmi1 reveal synergy and requirement for maintenance but not initiation
of Hox gene expression. Development 128, 1587-1597.

Akhmanova, A., Hoogenraad, C.C., Drabek, K., Stepanova, T., Dortland, B., Verkerk, T.,
Vermeulen, W., Burgering, B.M., De Zeeuw, C.I., Grosveld, F., Galjart, N., 2001. Clasps are
CLIP-115 and -170 associating proteins involved in the regional regulation of microtubule
dynamics in motile fibroblasts. Cell 104, 923-935.

Beal, R., Alonso-Carriazo Fernandez, A., Grammatopoulos, D.K., Matter, K., Balda, M.S., 2021.
ARHGEF18/p114RhoGEF Coordinates PKA/CREB Signaling and Actomyosin Remodeling to
Promote Trophoblast Cell-Cell Fusion During Placenta Morphogenesis. Front Cell Dev Biol 9,
658006.

Boot-Handford, R.P., Tuckwell, D.S., Plumb, D.A., Rock, C.F., Poulsom, R., 2003. A novel and
highly conserved collagen (pro(alpha)1(XXVII)) with a unique expression pattern and unusual
molecular characteristics establishes a new clade within the vertebrate fibrillar collagen family. J
Biol Chem 278, 31067-31077.

Bush, J.O., Jiang, R., 2012. Palatogenesis: morphogenetic and molecular mechanisms of
secondary palate development. Development 139, 231-243.

Chaudhury, A., Cheema, S., Fachini, J.M., Kongchan, N., Lu, G., Simon, L.M., Wang, T., Mao,
S., Rosen, D.G., Ittmann, M.M., Hilsenbeck, S.G., Shaw, C.A., Neilson, J.R., 2016. CELF1 is a
central node in post-transcriptional regulatory programmes underlying EMT. Nat Commun 7,
13362.

Cohen, M.M., Jr., 2010. Hedgehog signaling update. Am J Med Genet A 152A, 1875-1914.

El-Sibai, M., El Hajj, J., Al Haddad, M., El Baba, N., Al Saneh, M., Daoud Khatoun, W.,
Helaers, R., Vikkula, M., El Atat, O., Sabbagh, J., Abou Chebel, N., Ghassibe-Sabbagh, M.,
2021. Dysregulation of Rho GTPases in orofacial cleft patients-derived primary cells leads to
impaired cell migration, a potential cause of cleft/lip palate development. Cells Dev 165,
203656.

Ferretti, E., Li, B., Zewdu, R., Wells, V., Hebert, J.M., Karner, C., Anderson, M.J., Williams, T.,
Dixon, J., Dixon, M.J., Depew, M.J., Selleri, L., 2011. A conserved Pbx-Wnt-p63-Irf6 regulatory
module controls face morphogenesis by promoting epithelial apoptosis. Dev Cell 21, 627-641.

Girisha, K.M., Jacob, P., SriLakshmi Bhavani, G., Shah, H., Mortier, G.R., 2022. Steel
syndrome: Report of three patients, including monozygotic twins and review of clinical and
mutation profiles. Eur J Med Genet 65, 104521.

Gonzaga-Jauregui, C., Yesil, G., Nistala, H., Gezdirici, A., Bayram, Y., Nannuru, K.C.,
Pehlivan, D., Yuan, B., Jimenez, J., Sahin, Y., Paine, I.S., Akdemir, Z.C., Rajamani, S., Staples,
J., Dronzek, J., Howell, K., Fatih, J.M., Smaldone, S., Schlesinger, A.E., Ramirez, N., Cornier,
A.S., Kelly, M.A., Haber, R., Chim, S.M., Nieman, K., Wu, N., Walls, J., Baylor-Hopkins
Center for Mendelian Genetics, R.G.C., Poueymirou, W., Siao, C.J., Sutton, V.R., Williams,
M.S., Posey, J.E., Gibbs, R.A., Carlo, S., Tegay, D.H., Economides, A.N., Lupski, J.R., 2020.
Functional biology of the Steel syndrome founder allele and evidence for clan genomics
derivation of COL27A1 pathogenic alleles worldwide. Eur J Hum Genet 28, 1243-1264.

Grinev, V.V., Barneh, F., Ilyushonak, I.M., Nakjang, S., Smink, J., van Oort, A., Clough, R.,
Seyani, M., McNeill, H., Reza, M., Martinez-Soria, N., Assi, S.A., Ramanouskaya, T.V.,
Bonifer, C., Heidenreich, O., 2021. RUNX1/RUNX1T1 mediates alternative splicing and
reorganises the transcriptional landscape in leukemia. Nat Commun 12, 520.

Hammond, N.L., Dixon, M.J., 2022. Revisiting the embryogenesis of lip and palate
development. Oral Dis 28, 1306-1326.

Huynh, M.T., Beri-Dexheimer, M., Bonnet, C., Bronner, M., Khan, A.A., Allou, L., Philippe, C.,
Vigneron, J., Jonveaux, P., 2012. RUNX1T1, a chromatin repression protein, is a candidate gene
for autosomal dominant intellectual disability. Am J Med Genet A 158A, 1782-1784.

Jeong, J., Mao, J., Tenzen, T., Kottmann, A.H., McMahon, A.P., 2004. Hedgehog signaling in
the neural crest cells regulates the patterning and growth of facial primordia. Genes Dev 18, 937-
951.

Kim, S., Lewis, A.E., Singh, V., Ma, X., Adelstein, R., Bush, J.O., 2015. Convergence and
extrusion are required for normal fusion of the mammalian secondary palate. PLoS Biol 13,
e1002122.

Kotabagi, S., Shah, H., Shukla, A., Girisha, K.M., 2017. Second family provides further
evidence for causation of Steel syndrome by biallelic mutations in COL27A1. Clin Genet 92,
323-326.

Kress, C., Gautier-Courteille, C., Osborne, H.B., Babinet, C., Paillard, L., 2007. Inactivation of
CUG-BP1/CELF1 causes growth, viability, and spermatogenesis defects in mice. Mol Cell Biol
27, 1146-1157.

Lan, Y., Xu, J., Jiang, R., 2015. Cellular and Molecular Mechanisms of Palatogenesis. Curr Top
Dev Biol 115, 59-84.

Li, C., Lan, Y., Jiang, R., 2017. Molecular and Cellular Mechanisms of Palate Development. J
Dent Res 96, 1184-1191.

Liao, K.H., Chang, S.J., Chang, H.C., Chien, C.L., Huang, T.S., Feng, T.C., Lin, W.W., Shih,
C.C., Yang, M.H., Yang, S.H., Lin, C.H., Hwang, W.L., Lee, O.K., 2017. Endothelial
angiogenesis is directed by RUNX1T1-regulated VEGFA, BMP4 and TGF-beta2 expression.
PLoS One 12, e0179758.

Losa, M., Risolino, M., Li, B., Hart, J., Quintana, L., Grishina, I., Yang, H., Choi, I.F., Lewicki,
P., Khan, S., Aho, R., Feenstra, J., Vincent, C.T., Brown, A.M.C., Ferretti, E., Williams, T.,
Selleri, L., 2018. Face morphogenesis is promoted by Pbx-dependent EMT via regulation of
Snail1 during frontonasal prominence fusion. Development 145.

Mai, C.T., Isenburg, J.L., Canfield, M.A., Meyer, R.E., Correa, A., Alverson, C.J., Lupo, P.J.,
Riehle-Colarusso, T., Cho, S.J., Aggarwal, D., Kirby, R.S., National Birth Defects Prevention,
N., 2019. National population-based estimates for major birth defects, 2010-2014. Birth Defects
Res 111, 1420-1435.

Maiato, H., Khodjakov, A., Rieder, C.L., 2005. Drosophila CLASP is required for the
incorporation of microtubule subunits into fluxing kinetochore fibres. Nat Cell Biol 7, 42-47.

Malicki, J.J., Johnson, C.A., 2017. The Cilium: Cellular Antenna and Central Processing Unit.
Trends Cell Biol 27, 126-140.

Mansilla, M.A., Cooper, M.E., Goldstein, T., Castilla, E.E., Lopez Camelo, J.S., Marazita, M.L.,
Murray, J.C., 2006. Contributions of PTCH gene variants to isolated cleft lip and palate. Cleft
Palate Craniofac J 43, 21-29.

Marone, R., Hess, D., Dankort, D., Muller, W.J., Hynes, N.E., Badache, A., 2004. Memo
mediates ErbB2-driven cell motility. Nat Cell Biol 6, 515-522.

McConnachie, D.J., Stow, J.L., Mallett, A.J., 2021. Ciliopathies and the Kidney: A Review. Am
J Kidney Dis 77, 410-419.

Moore, E.R., 2022. Primary Cilia: The New Face of Craniofacial Research. Biomolecules 12.

Mort, R.L., Jackson, I.J., Patton, E.E., 2015. The melanocyte lineage in development and
disease. Development 142, 620-632.

Mossey, P.A., Little, J., Munger, R.G., Dixon, M.J., Shaw, W.C., 2009. Cleft lip and palate.
Lancet 374, 1773-1785.

Murray, S.A., Oram, K.F., Gridley, T., 2007. Multiple functions of Snail family genes during
palate development in mice. Development 134, 1789-1797.

Nakajima, H., Tanoue, T., 2011. Lulu2 regulates the circumferential actomyosin tensile system
in epithelial cells through p114RhoGEF. J Cell Biol 195, 245-261.

Plumb, D.A., Dhir, V., Mironov, A., Ferrara, L., Poulsom, R., Kadler, K.E., Thornton, D.J.,
Briggs, M.D., Boot-Handford, R.P., 2007. Collagen XXVII is developmentally regulated and

forms thin fibrillar structures distinct from those of classical vertebrate fibrillar collagens. J Biol
Chem 282, 12791-12795.

Plumb, D.A., Ferrara, L., Torbica, T., Knowles, L., Mironov, A., Jr., Kadler, K.E., Briggs, M.D.,
Boot-Handford, R.P., 2011. Collagen XXVII organises the pericellular matrix in the growth
plate. PLoS One 6, e29422.

Rahimov, F., Jugessur, A., Murray, J.C., 2012. Genetics of nonsyndromic orofacial clefts. Cleft
Palate Craniofac J 49, 73-91.

Reiter, J.F., Skarnes, W.C., 2006. Tectonic, a novel regulator of the Hedgehog pathway required
for both activation and inhibition. Genes Dev 20, 22-27.

Reynolds, K., Zhang, S., Sun, B., Garland, M.A., Ji, Y., Zhou, C.J., 2020. Genetics and signaling
mechanisms of orofacial clefts. Birth Defects Res 112, 1588-1634.

Rosenblatt, J., Raff, M.C., Cramer, L.P., 2001. An epithelial cell destined for apoptosis signals
its neighbors to extrude it by an actin- and myosin-dependent mechanism. Curr Biol 11, 1847-
1857.

Schwab, K.R., Patterson, L.T., Hartman, H.A., Song, N., Lang, R.A., Lin, X., Potter, S.S., 2007.
Pygo1 and Pygo2 roles in Wnt signaling in mammalian kidney development. BMC Biol 5, 15.

Selleri, L., Rijli, F.M., 2023. Shaping faces: genetic and epigenetic control of craniofacial
morphogenesis. Nat Rev Genet 24, 610-626.

Som, P.M., Naidich, T.P., 2013. Illustrated review of the embryology and development of the
facial region, part 1: Early face and lateral nasal cavities. AJNR Am J Neuroradiol 34, 2233-
2240.

Spektor, A., Tsang, W.Y., Khoo, D., Dynlacht, B.D., 2007. Cep97 and CP110 suppress a cilia
assembly program. Cell 130, 678-690.

Takahashi, S., 2021. Functional analysis of large MAF transcription factors and elucidation of
their relationships with human diseases. Exp Anim 70, 264-271.

Thauvin-Robinet, C., Lee, J.S., Lopez, E., Herranz-Perez, V., Shida, T., Franco, B., Jego, L., Ye,
F., Pasquier, L., Loget, P., Gigot, N., Aral, B., Lopes, C.A., St-Onge, J., Bruel, A.L., Thevenon,
J., Gonzalez-Granero, S., Alby, C., Munnich, A., Vekemans, M., Huet, F., Fry, A.M., Saunier,
S., Riviere, J.B., Attie-Bitach, T., Garcia-Verdugo, J.M., Faivre, L., Megarbane, A., Nachury,
M.V., 2014. The oral-facial-digital syndrome gene C2CD3 encodes a positive regulator of
centriole elongation. Nat Genet 46, 905-911.

Thomas, S., Legendre, M., Saunier, S., Bessieres, B., Alby, C., Bonniere, M., Toutain, A.,
Loeuillet, L., Szymanska, K., Jossic, F., Gaillard, D., Yacoubi, M.T., Mougou-Zerelli, S., David,
A., Barthez, M.A., Ville, Y., Bole-Feysot, C., Nitschke, P., Lyonnet, S., Munnich, A., Johnson,
C.A., Encha-Razavi, F., Cormier-Daire, V., Thauvin-Robinet, C., Vekemans, M., Attie-Bitach,
T., 2012. TCTN3 mutations cause Mohr-Majewski syndrome. Am J Hum Genet 91, 372-378.

Tsuji, T., Ohta, Y., Kanno, Y., Hirose, K., Ohashi, K., Mizuno, K., 2010. Involvement of p114-
RhoGEF and Lfc in Wnt-3a- and dishevelled-induced RhoA activation and neurite retraction in
N1E-115 mouse neuroblastoma cells. Mol Biol Cell 21, 3590-3600.

Vlasova-St Louis, I., Dickson, A.M., Bohjanen, P.R., Wilusz, C.J., 2013. CELFish ways to
modulate mRNA decay. Biochim Biophys Acta 1829, 695-707.

Wang, C., Li, J., Meng, Q., Wang, B., 2017. Three Tctn proteins are functionally conserved in
the regulation of neural tube patterning and Gli3 processing but not ciliogenesis and Hedgehog
signaling in the mouse. Dev Biol 430, 156-165.

Wehby, G.L., Cassell, C.H., 2010. The impact of orofacial clefts on quality of life and healthcare
use and costs. Oral Dis 16, 3-10.

Welsh, I.C., Hart, J., Brown, J.M., Hansen, K., Rocha Marques, M., Aho, R.J., Grishina, I.,
Hurtado, R., Herzlinger, D., Ferretti, E., Garcia-Garcia, M.J., Selleri, L., 2018. Pbx loss in
cranial neural crest, unlike in epithelium, results in cleft palate only and a broader midface. J
Anat 233, 222-242.

Wojcicki, P., Kozlik, M.J., Wojcicka, K., 2016. Genetic Factors in Selected Complex Congenital
Malformations with Cleft Defect. Adv Clin Exp Med 25, 977-987.

Xu, X., Jin, D., Durgan, J., Hall, A., 2013. LKB1 controls human bronchial epithelial
morphogenesis through p114RhoGEF-dependent RhoA activation. Mol Cell Biol 33, 2671-2682.

Zaoui, K., Honore, S., Isnardon, D., Braguer, D., Badache, A., 2008. Memo-RhoA-mDia1
signaling controls microtubules, the actin network, and adhesion site formation in migrating
cells. J Cell Biol 183, 401-408.

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O., Benner, C.,
Chanda, S.K., 2019. Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. Nat Commun 10, 1523.
447

Figure 1

Figure 2