Evolutionary and biological mechanisms underpinning chitin degradation in aquatic

fungi

Nathan Chrismas[1], Kimberley Bird[1], Davis Laundon[1,2], Poppy Hesketh-Best[1,3], Chloe Lieng[1,3]

& Michael Cunliffe[1,3]

[1] Marine Biological Association, The Laboratory, Citadel Hill, Plymouth, PL1 2PB.

[2] School of Environmental Sciences, University of East Anglia, Norwich Research Park,

Norwich, NR4 7TJ.

[3] School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth,

PL4 8AA.

Corresponding author: Michael Cunliffe

14   Marine Biological Association, The Laboratory, Citadel Hill,

15   Plymouth, PL1 2PB, UK.

16   E: micnli@mba.ac.uk   T: +44 (0)1752 426328

Keywords: Chytridiomycota, chytrids, saprotrophy, gene expansion, gene transfer, evolution.

Abstract

Fungal biology underpins major processes in ecosystems. The Chytridiomycota (chytrids) is

a group of early-diverging fungi, many of which function in ecosystems as saprotrophs

processing high molecular weight biopolymers, however the mechanisms underpinning

chytrid saprotrophy are poorly understood. Genome sequences from representatives across

the group and the use of model chytrids offers the potential to determine new insights into

their evolution. In this study, we focused on the biology underpinning chitin saprotrophy, a

common ecosystem function of aquatic chytrids. The genomes of chitinophilic chytrids have

expanded inventories of glycoside hydrolase genes responsible for chitin processing,

complemented with bacteria-like chitin-binding modules (CBMs) that are absent in other

chytrids. In the model chitinophilic saprotroph *Rhizoclosmatium globosum* JEL800, the

expanded repertoire of chitinase genes is diverse and almost half were detected as proteins

in the secretome when grown with chitin. Predicted models of the secreted chitinases

indicate a range of active site sizes and domain configurations. We propose that increased

diversity of secreted chitinases is an adaptive strategy that facilitates chitin degradation in

the complex heterologous organic matrix of the arthropod exoskeleton. Free swimming *R.*

globosum JEL800 zoospores are chemotactic to the chitin monomer N-acetylglucosamine

and accelerate zoospore development when grown with chitin. Our study sheds light on the

underpinning biology and evolutionary mechanisms that have supported the saprotrophic

niche expansion of some chytrids to utilise lucrative chitin-rich particles in aquatic

ecosystems and is a demonstration of the adaptive capability of this successful fungal group.

Introduction

Fungi are leading drivers of ecosystem processes, including the degradation of high-

molecular weight, often recalcitrant, biogenic organic polymers. As high-molecular weight

material is too large to be directly absorbed across fungal cell walls, substrates must first be

degraded in the extracellular environment. This process is carried out by enzymes secreted

by the fungus into the environment before subsequent breakdown products are absorbed via

osmotrophy (Richards and Talbot 2018). Our understanding of the biological mechanisms

underpinning fungal saprotrophy is largely based on plant cell wall decomposition by

terrestrial Ascomycota and Basidiomycota, the evolution of which is synonymous with the

successful expansion of the kingdom through exploitation of diverse ecological niches

(Berbee et al. 2017, Nagy et al. 2017, Naranjo-Ortiz and Gabaldón 2019).

The Chytridiomycota (chytrids) is a group of early-diverging fungi that are typically

unicellular, developing into a sporangium with substrate-attaching and feeding rhizoids (Fig.

1a). Within mature sporangia, multiple uniflagellate zoospores form that lack a cell wall and

are usually motile when released into the environment. Once a suitable substrate is found by

swimming zoospores they attach and encyst, developing into a new sporangium with

rhizoids to repeat the life cycle (Medina and Buchler 2020). Chytrids broadly occupy two

major functional roles in ecosystems, either as (i) parasites, mainly of algae with the

exceptions of a few amphibian pathogens, or (ii) saprotrophs that degrade a range of

recalcitrant organic biopolymers. A recent phylogenomic study has proposed that

saprotrophy is a derived trait in chytrids and that the ancestral trait is algal parasitism

(Thomé et al. 2023). The biological mechanisms underpinning aquatic chytrid saprotrophy

are currently poorly understood including in an evolutionary context (Laundon and Cunliffe

2021).

Chitin-based particles such as arthropod exoskeleton remains (exuviae) are

prominent aquatic chytrid habitats (Sparrow 1960) (Fig. 1b-d). Chitin is a polymer of N-

acetylglucosamine (GlcNAc) and is the second most abundant polysaccharide on Earth after

cellulose. Chitin degradation is carried out by chitinases that either remodel cellular chitin

(e.g. in the fungal cell wall), or externally process chitin substrate (Duo-Chuan 2006). Fungal

chitinases belong to the glycoside hydrolase family 18 (GH18) of carbohydrate-active

enzymes (CAZymes) (Funkhouser and Aronson 2007) and act by introducing random breaks

into crystal chitin (endochitinases) or cleaving terminal ends of GlcNAc polymers

(exochitinases). GH18 chitinases are classified into two main related groups (Goughenour et

al. 2021). Group AC are typically exochitinases with a deep, tunnel shaped substrate-

interacting groove whereas Group B are typically endochitinases with a shallow, open

groove. GH18 domain-containing proteins can also include carbohydrate binding modules

(CBMs) belonging to ChtBD3/ChiC_BD domain families, which increase substrate binding

affinity. Following primary chitinolytic activity by GH18s, chitin-derived GlcNAc is further

processed by GH20 family β-*N*-acetyl-D-hexosaminidases.

The aims of this study were to determine the biological mechanisms underpinning

chytrid chitin saprotrophy by focusing on the following specific questions: (i) what is the

underlying molecular machinery (i.e. genes, mRNA and proteins/enzymes) involved in chitin

saprotrophy, and (ii) what chitin-associated behaviour is shown by swimming zoospores. Our

approach was to compare the reference genome sequences from chitinophilic chytrids with

representatives across the Chytridiomycota that occupy a range of different ecological

niches, combined with the use of the model saprotroph *Rhizoclosmatium globosum* JEL800

to determine new insights into chytrid biology. *Rhizoclosmatium globosum* is an aquatic

chytrid found prevalent in nature attached to chitin-rich arthropod exuviae (Sparrow 1960).

The strain *R. globosum* JEL800 is an emerging model for biological and evolutionary studies

of early diverging fungi (Laundon et al. 2020, Venard et al. 2020, Prostak et al. 2021,

Galindo et al. 2022, Laundon et al. 2022) that was isolated by chitin baiting (Joyce Longcore,

pers. comm.) and grows readily on chitin as a sole carbon source (Laundon et al. 2020).

Results

Expansion of genes encoding chitin degradation enzymes in chitinophilic chytrids

The abundance of GH18 chitinase encoding genes in the reference genomes of 36 chytrids

across the Chytridiomycota was compared, including the Chytridiomycetes Chytridiales

(n=7), Caulochtriales (n=1), Cladochytriales (n=2), Polychytriales (n=1), Rhizophydiales

(n=5), Spizellomycetales (n=6), *Blyttiomyces helicus incertae sedis* (n=1), as well as

Monoblepharidomycetes (n=2) and Neocallimastigomycetes (n=11) that occupy a diverse

range of ecological niches (Fig 2ai, Supplementary File, Supplementary Figure 1).

All chytrids associated with arthropod exuviae in nature and isolated by chitin-baiting

belong to the Chytridiales except *Polychytrium aggregatum* JEL109 from the sister order

Polychytriales. All the chitinophilic chytrids i.e. *R. globosum* JEL800, *Obelidium mucronatum*

JEL802, *Chytriomyces hyalinus* JEL632, *Chytriomyces* sp. MP 7 and *P. aggregatum* JEL109

showed the highest enrichment of GH18 encoding genes (Fig 2ai, Supplementary Fig 1). All

other chytrids, i.e. occupying non-chitin associated niches, had relatively lower GH18 gene

abundances with the plant-processing Neocallimastigomycetes found in herbivore guts and

plant-baited *Cladochytrium replicatum* JEL0714 lowest (Fig 2ai, Supplementary Fig 1). A

similar pattern of increased gene frequency was observed for GH20 encoding genes,

particularly for the chitinophilic *R. globosum* JEL800, *O. mucronatum* JEL802 and *P.*

aggregatum JEL109 (Fig. 2aii, Supplementary Fig 1).

Bacteria-like chitin binding module genes in chitinophilic chytrids

The abundance of genes containing carbohydrate binding module 5 (CBM5) domains in the

genomes of the chitinophilic chytrids was distinctive, with *R. globosum* JEL800 and *O.*

mucronatum JEL802 particularly high (Fig. 2aiii, Supplementary Fig 1). Although GH18 and

GH20 encoding genes were present but at lower abundances in the genomes of non-chitin

associated chytrids (Fig. 2 ai-aii), genes containing CBM5s were only detected in the

genomes of chitinophilic chytrids except for the Monoblepharidomycete *Hyaloraphidium*

curvatum SAG235-1. In most cases, BLAST searches of the CBM5 domains from

chitinophilic chytrids revealed similarity with equivalent bacterial ChtBD3/ChiC_BD domains

(Supplementary File), many of which were found in bacteria with known chitin-degrading

capacity (e.g. *Chitinilyticum aquatile*, *Paenibacillus chitinolyticus*, *Chitinolyticbacter albus*).

Only three genes, one in *Chytriomyces* sp. MP71 (KAI8611252) and two in *Polychytrium*

aggregatum JEL109 (XP_052967452 and XP_052967585), contained domains more similar

to those found in non-chytrid fungi with matches to Agaricomycetes (Basidiomycota). In

addition, our CBM5 domain searches matched to a further three Chytridiales genomes not

included with the reference Mycocosm genomes (*Podochytrium* sp. JEL0797, *Siphonaria sp.*

JEL0065, and *Physocladia obscura* JEL0513), all of which were isolated from mayfly

(Ephemeroptera) exuviae (Joyce Longcore, pers. comm.).

Rhizoclosmatium globosum GH18 and GH20 gene diversity and domain architecture

Forty-nine *R. globosum* JEL800 genes contained GH18 domains. Maximum-Likelihood

phylogenetic analysis of the GH18 domains revealed two major clades that corresponded

with either Group AC (predominantly exochitinases) or Group B (predominantly

endochitinases) (Fig. 2b). Group AC contained domains for 24 proteins, with 21 forming a

large sub-clade, with two domains appearing in ORY47749. Although ORY48020 fell within

this clade, it was separated by a long branch with relatively low bootstrap support.

ORY40370 and ORY40369 fell outside the main clade. Group B contained domains from 25

proteins. One clade contained domains from 16 proteins and a second clade contained a

variety of GH18 domains from nine proteins with low bootstrap support for several branches

within this clade.

Chitin binding domains, all of which were CBM5-type, were found on eight genes

(Fig. 2b). Seven were associated with Group AC (ORY40369, ORY41001, ORY53850,

ORY32764, ORY35945, ORY37254 and ORY34525) and one with Group B (ORY39595).

Signal peptides associated with protein secretion were found in 17 proteins. Eleven of these

were found in Group A proteins (45.8% of Group A) and six with Group B (24% of Group B).

Fourteen *R. globosum* JEL800 genes were found to contain GH20 domains (Fig. 2c),

Maximum-Likelihood phylogenetic analysis of which revealed two well supported clades,

each containing seven genes, nine of which had associated signal peptides.

GH18 and GH20 in the Rhizoclosmatium globosum secretome and transcriptome

A total of 23 proteins from GH18 domain-containing genes were detected in the *R. globosum*

JEL800 secretome when grown on chitin as the sole source of carbon: 10 from Group AC

(exochitinases), 12 from Group B (endochitinases) and one other (Fig. 3a), accounting for

almost half (47%) of the GH18 genes detected in the genome. Proteins for six of the GH20

domain-containing proteins were also detected in the secretome (Fig. 3a). All GH18 and

GH20 proteins detected in the secretome matched reciprocal mRNA sequences detected in

the cellular transcriptome, with a positive correlation of secretome GH18 and GH20 protein

abundance with mRNA fpkm (Fig. 3b). Predicted structures of the proteins encoded from

GH18 domain-containing genes that were detected as secretome proteins suggests that *R.*

globosum JEL800 chitinases are variable both in terms of total size and substrate binding

site size (Fig. 3c and Supplementary Figures 2 and 3). Modelling of the abundant GH18

protein ORY35945 predicted a C-terminal GH18 domain and N-terminal CBM5 domain

joined by a linker domain (Fig. 3d).

Rhizoclosmatium globosum zoospore behaviour indicates adaptations to chitinophilic

lifestyle

Swimming *R. globosum* JEL800 zoospores are chemotactic towards the chitin monomer

GlcNAc. Targeted swimming occurred with zoospores from cultures grown with GlcNAc as

the sole carbon source (Fig. 4ai) and when grown without GlcNAc in the complex medium

PmTG (Fig. 4aii). Even though *R. globosum* grows with glucose as an alternative sole

carbon source (Supplementary Figure 4), chemotaxis did not occur towards glucose with

either GlcNAc or PmTG grown cultures (Fig. 4ai, Fig. 4aii).

We made several attempts to determine chemotaxis in zoospores from cultures

grown with chitin but were unable to conduct the assay due to apparent reduction in

zoospore motility. To explore this further, we compared general motility (i.e. swimming

versus stationary) of zoospores harvested from cultures grown with chitin to the motility of

zoospores from cultures grown with PmTG, showing that chitin-grown zoospores rapidly lose

motility after release compared to zoospores grown with PmTG which remained more motile

for longer (Fig. 4b).

Discussion

Gene expansion and gene transfer underpin chitinophilic chytrid evolution

Here we provide genome-based evidence for two complementary mechanisms through which chitin saprotrophy has evolved in chytrids. The abundance of genes encoding GH18 chitinases and GH20 β-N-acetyl-d-hexosaminidases responsible for the breakdown of chitin and subsequent processing of chitin-derived GlcNAc respectively were increased in chitinophilic chytrids relative to non-chitinolytic chytrids. Alongside GH18 gene expansion, bacteria-like chitin-binding CBM5 domains were in the genomes of the chitinophilic chytrids, indicating that horizontal gene transfer (HGT) has taken place, potentially from chitinophilic bacteria to chitinophilic chytrids. The acquisition of chitin-binding capability likely improved the functional efficiency of chytrid chitinases to degrade extracellular chitin-based substrates and therefore has clear fitness advantages.

Lineage specific gene family expansion is a mechanism of adaptive evolution in eukaryotes and including in fungi (Lespinet et al. 2002). Expansion of hydrolase gene families related to host/substrate specificity has been reported in other fungi. In non-chytrid fungi, necrotrophic Ascomycota and Basidiomycota exhibit expansion of GH28 family pectinases (Sprockett et al. 2011), and white rot fungi including *Armillaria* spp. (Basidiomycota) have an increased complement of genes encoding cellulose- and xylan-degrading enzymes (Sipos et al. 2017). Increased copy numbers of proteases have been associated with the evolution of amphibian pathogenicity in *Batrachochytrium dendrobatidis* by facilitating attachment and penetration of host cells (Joneson et al. 2011, Farrer et al. 2017). Specifically in terms of GH18 gene expansion, as we show here in early-diverging chitinophilic chytrids, similar has been observed across other fungi including various Ascomycota (Karlsson and Stenlid 2008) and ectomycorrhizal fungi (Maillard et al. 2023).

A key feature of gene family expansion in fungi is that it facilitates exploitation of specific niches by allowing increased rates of enzyme production and functional novelty (Gladieux et al. 2014). The increased GH18 and GH20 inventory in chitinophilic chytrids may confer the ability to react quickly to substrate availability by transcribing multiple genes. Although cellular mRNA was detected for almost all of the GH18 and GH20 genes encoded within the *R. globosum* JEL800 genome, not all transcribed genes were detected in the secretome. As well as accounting for cell wall processing cellular chitinases (which are yet to be characterised in chytrids), some of this untranslated mRNA may play a role maintaining a bank of mRNA for a variety of chitinases, *R. globosom* JEL800 may be able to remain 'primed' for rapid protein synthesis.

Increased chitinase gene copy number in chitinophilic chytrids contrasts with chytrids occupying other niches which have reduced abundances of chitinase encoding genes. It is important to reiterate that GH18 chitinases are likely used for remodelling chitin in the chytrid

cell wall based on analogous mechanisms in other fungi (Gow et al. 2017) and therefore

necessary in all chytrids. Our study suggests that chitinase gene family expansion is varied

within lineages, with an apparent increase in chitinase gene copy number associated with

the base of the Chytridiales encompassing the closely related chitinophilic *Polychytrium*

aggregatum (Polychytriales) followed by subsequent gene reduction in the plant saprotroph

Entophlyctis luteolus (Chytridiales).

The presence of chitin-binding CBM5 domains is a distinguishing feature of

chitinophilic chytrid genomes. In *R. globosum* JEL800, chitin-binding domains are present on

multiple chitinases and are potentially derived from xenologs originating from chitinophilic

bacteria. Horizontal gene transfer is well established in prokaryotes but has only relatively

recently been determined in fungi including early-diverging fungi. Comparative assessment

of herbivore gut anaerobic Neocallimastigomycetes has shown the HGT played a role in

their evolution (Murphy et al. 2019). As we propose here for aquatic chitinophilic chytrids,

anaerobic gut Neocallimastigomycetes HGT occurred from functionally similar bacteria that

likely co-occur in the same niche i.e. anaerobes in herbivore guts that degrade plant-derived

polysaccharides and is especially associated with CAZymes and extracellular

polysaccharide degradation. Murphy *et al.* (2019) proposed that HGT, including the acquired

ability to degrade plant-derived polysaccharides, was a key feature of the expansion of the

Neocallimastigomycetes, as we propose here with the evolution of the Chytridiales. Wider

comparison across fungal divisions including Mucoromycota indicates that the early

diverging groups of aquatic fungi are more likely to undergo HGT compared to terrestrial

taxa (Ciach et al. 2024). Further work is needed using more sophisticated bioinformatic

approaches to explore HGT in chitinophilic chytrids and other saprotrophic guilds.

Diversity in chitinase gene architecture and modelled secretome chitinase structure

Relatively few crystal structures of fungal chitinases have been characterised (Jiang et al.

2021) and prediction of specific chitinase functions is therefore difficult (Goughenour et al.

2021) but variation in binding site volume relative to overall protein volume indicates

structural diversity which may be related to variation in substrate, with different enzyme

activities able to account for substrate complexity (discussed further below). The high levels

of secretion of Group AC exochitinases compared to Group B endochitinases suggests that

cleavage of terminal GlcNAc dimers from chitin is a particularly important characteristic of

the chitinophilic lifestyle.

Alongside size complexity, we also show that some chitinases are multimodular with

additional domains that potentially expand the chytrids chitin degradation capability.

Considering the major secreted chitinase ORY35945 (Fig. 3d), this has broadly analogous

domain configurations with the cellulose degrading endoglucanase (EG11) in *Rhizopus*

stolonifer (Mucoromycota) with a C-terminal catalytic domain linked with N-terminal CBM

(Tang et al. 2015). The function of linkers between catalytic and binding domains has not

been considered with chitinases but has been studied with cellulose and lignocellulose

degrading enzymes and suggest that linkers may have different functions including general

improvements in catalytic efficiency (Payne et al. 2013, Tang et al. 2015, Courtade et al.

2018), promotion of localised and repeated substrate processing (Courtade et al. 2018) and

also have an additional substrate binding role (Payne et al. 2013). In terms of the

chitinophilic chytrids hosting bacteria-like CBM5 domains, homologous CBM5 from the

chitinophilic bacterium *Serratia marcescens* exhibits affinity specifically for α-chitin and

promotes chitin degradation in the presence of other polysaccharides (Liu et al. 2023).

In nature, chitinophilic chytrids are observed growing on arthropod exoskeleton

remains (Sparrow 1960) in which chitin is not available as a distinct molecule but instead

packaged within a complex heterologous organic matrix. Even though exoskeleton matrices

likely vary between source taxa, the general structure is based around crystalline α-chitin

fibres in chitin-protein planes that are stacked in a twisted 'plywood' structure (Raabe et al.

2006). Taken together, we propose that the configuration diversity from varied chitinase

sizes, associated bacteria-like CBM5 domains with likely α-chitin affinity and further

capability caused the presence (or absence) of linkers between catalytic sites and binding

domains has adaptive advantages for saprotrophy within the naturally heterologous organic

matrix in which the primarily α-chitin substrate is intertwined.

Zoospore behaviours as adaptive traits for the chitinophilic lifestyle

Rhizoclosmatium globosum JEL800 zoospores are chemotactic to the chitin monomer

GlcNAc independent of the growth conditions of the previous generation, a trait that has

obvious advantages for targeting chitin-rich substrates such as arthropod exuviae in complex

aquatic environments. Chemotaxis has been shown in non-saprotrophic chytrid zoospores,

particularly species that are parasites. *Batrachochytrium dendrobatidis* zoospores are

chemotactic towards amphibian-produced compounds (Moss et al. 2008) and algal parasites

target cell extracts of stressed hosts (Scholz et al. 2017). The biological mechanisms

underpinning chemotaxis in aquatic chytrids are poorly understood (Laundon and Cunliffe

2021). It is possible that response to environmental signals could be linked to the

rumposome, an enigmatic structure connecting the zoospore cell surface to the flagellar

machinery that is present in *R. globosum* JEL800 zoospores (Laundon et al. 2022) and is

likely a site of calcium-mediated signalling (Powell 1983). Phototaxis has also been shown in

chytrid zoospores, including with *R. globosum* JEL800, facilitated by a CyclOp-fusion protein

based sensory system (Galindo et al. 2022). Together, this suggests that via chemotaxis and

phototaxis, swimming chytrid zoospores utilise multi-signal mechanisms to navigate the
aquatic environment.
The chemotactic response to GlcNAc shown here suggests that, in nature,
chitinophilic chytrid zoospores respond to signals of 'on-going' chitin saprotophy that is
releasing GlcNAc via degradation by other organisms. Chitin degradation in aquatic
ecosystems is also performed by chitinophilic bacteria that share the same habitat and
potential niche as chitinophilic chytrids such as *R. globosum* (Roberts et al. 2020). It is likely
that a range of chytrid-bacteria interactions are associated with chitin-rich particles, including
particle colonisation, that involve the release of chemotactic signals from bacteria to chytrids
and *vice versa* depending on which are the primary or secondary colonisers (Roberts et al.
2020).
When grown with chitin, we observed faster loss of next generation zoospore motility
compared to non-chitin grown *R. globosum* JEL800 which remained motile for longer.
Accelerated transition from a non-feeding swimming zoospore to the substrate attaching
stage of the life cycle (i.e. encystment) is a possible trait that could facilitate the rapid
colonisation and exploitation of ephemeral chitin-rich substrates such as mayfly exuviae in a
dynamic resource-complex and competitive environment. Chytrid zoospore biology is poorly
understood, including controls on the cell cycle (Medina and Buchler 2020, Laundon et al.
2022), however based on frog mucus exposure experiments with *B. dendrobatidis*, likely
involve critical host/substrate chemical cues that initiate encystment via calcium signalling
and actin cytoskeleton remodelling (Robinson et al. 2022). Chytrid zoospores are
metabolically active, yet are transcriptionally and translationally inactive, with ribosomes held
dormant (Medina and Buchler 2020). As with other chytrids, in *R. globosum* JEL800
zoospores, ribosomes are maintained dormant in a distinct cluster that only dissipates once
the cell transitions to the germling stage when transcription and translation are initiated
(Laundon et al. 2022). Therefore, we propose the molecular mechanisms underpinning the
zoospore behavioural traits that we report here that likely support chitin utilisation are
maternally derived during zoosporogenesis and combine independent (i.e. GlcNAc
chemotaxis) and dependent (i.e. accelerated zoospore-germling transition) of maternal
growth conditions (i.e. with or without chitin).
326
Conclusion
Our study has shown mechanisms that underpin and have likely supported the evolution of
chitin-based saprotrophy in chytrids. We propose that the complementary expansion of
existing GH18 and GH20 genes, probably from cell wall remodelling functions, alongside
niche-specific novel gene transfer of GH18-associated chitin-binding CBM5 domains,
potentially from functionally similar bacteria, facilitated chitin saptrophy evolution. Expansion

of GH18 genes, acquisition of CBM5 domains and linkers, allowed the diversification of
secreted chitinases that may facilitate utilisation of chitin within the complex arthropod
exoskeleton matrix and, along with substrate-dependent zoospore behavioural traits,
determined the widespread and successful exploitation of the chitin-based particle niche in
aquatic ecosystems.
338
Methods
Comparison of GH18, GH20 and CBM5 encoding genes across chytrid genomes
Counts of predicted GH18, GH20 and CBMs genes were retrieved from Chytridiomycota
reference genomes from Mycocosm (https://mycocosm.jgi.doe.gov/, accessed 10/2023)
(Supplementary File, Supplementary Figure 5). Conserved domains in *R. globosum* JEL800
GH18 and GH20 genes were identified using CD-search (Yang et al. 2020) and aligned
separately in MAFFT (Katoh and Standley 2013) using E-INS-i before maximum likelihood
phylogenetic analysis was carried out using the CIPRES (Miller et al. 2010) implementation
of RaxML 8.2.10 (Stamatakis 2014) using the CAT model and the BLOSUM62 matrix for
divergent protein sequences with automatic bootstrapping. Domain trees were visualised
using FigTree v.1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/). GH18 domains were further
subclassified by comparison with a phylum-wide sampling of fungal GH18 domains (Junges
et al. 2014) (Supplementary Figure 5). Presence of signal peptides in each gene were
predicted using the SignalP 4.1 webserver (Petersen et al. 2011). Complete protein domains
were plotted using genoplotR (Guy et al. 2010). Chitin binding domains (ChtBD3/ChiC_BD)
in genes annotated as CBM5 in Mycocosm for all CBM5-containing lineages were identified
using CD-search. Where domain homology was identified, domains were searched against
GenBank using BLASTp (e-value cutoff = 0.00001) excluding hits for the chitinophilic
chytrids in this study.
358
Rhizoclosmatium globosum JEL800 maintenance, growth and microscopy
Rhizoclosmatium globosum JEL800 was routinely maintained on peptidised milk, tryptone,
and glucose (PmTG) agar plates (Barr 1986) at 22°C in the dark. For experimental analysis,
R. globosum JEL800 was also grown with modified Bold's Basal Medium (BBM) (Laundon et
al. 2020) diluted 10-fold and with the addition of 0.25 g ammonium sulfate and 0.5 mL f/2
vitamin stock in 1 L (Guillard 1975). To produce samples for transcriptome and secretome
analysis, *R. globosum* was grown in 50 mL BBM with 10mM chitin in 250 mL Erlenmeyer
flasks inoculated with ~14,000 zoospores and shaking at 70 rpm. Cultures were harvested
after 7 days growth by transferring contents to a 50 mL centrifuge tube and centrifuging at
4,000 G for 10 min. Cell pellets were snap frozen with liquid nitrogen. The supernatant was
removed and Halt™ Protease & Phosphate Inhibitor Cocktail (Thermo) added at 5 μL / 50

mL with mixing by inversion. Protease treated supernatant (~ 50 mL) was concentrated to ~

250 µL in 15 mL batches at 4,000 G for 20 min each (total ~ 1 hour) using 10 kDa Amicon®

Ultra 15 mL Centrifugal Filters (Millipore). Concentrated supernatant was transferred to a 3

kDa Pierce™ Protein Concentrator PES 0.5 mL (Thermo) and centrifuged for 20 min at

15,000 G to a final volume of ~ 100 µL. All centrifuge steps were carried out at 4 °C and

samples stored at -80 °C.

To screen a range of alternative non-chitin carbon sources used by *R. globosum*

JEL800 for growth, mature cultures on PmTG agar plates were sporulated using 4 mL dH_2O

for ~ 2 hours at room temperature. Zoospores were then harvested and passed through a 10

µm sieved before being washed 3 times in carbon free BBM by centrifuging at 4000 G for 10

min at 4 °C and re-suspended in carbon free BBM. Biolog PM 1 and PM 2A Carbon

Utilisation Assay plates were inoculated with 150 µL per well at a final density of 1×10^3

zoospores per well. Assay plates were incubated at 22 °C and absorbance at 405 nm

(OD_{405}) was measured periodically using a plate reader (CLARIOstar).

To demonstrate *R. globosum* JEL800 growth on an arthropod exoskeleton, *Daphnia*

were acquired from a local aquarium supplier and cleaned by incubation in 10% sodium

dodecyl sulphate solution for 30 mins, followed by 100% ethanol for 1 hour and then

autoclaved in ddH_2O. Carapaces were washed twice in sterile ddH_2O between each step.

Culture dishes containing 3mL BBM and one carapace were inoculated with 500 µl zoospore

suspension (1.03×10^6 zoospores.ml^{-1}) and incubated at 22°C. Chytrids growing on

carapaces were imaged with DIC using a Leica Dmi8 microscope (Leica, Germany).

Transcriptome and secretome analysis

RNA was extracted from the cell pellets using the RNeasy Micro kit (Qiagen) according to

the manufacturer's instructions. Pellets were thawed in 2 mL RLT lysis buffer (Qiagen)

containing 10 µL mL^{-1} of 2-mercaptoethanol, incubated at room temperature for 5 minutes

and mixed periodically using a vortex. Cell debris was removed by centrifuging at 4000 G for

1 min, the lysate was recovered through a QIA shredder (Qiagen). An equal volume of 100

% ethanol was added to the homogenised lysate before being transferred in batches to the

RNeasy extraction column including on column DNase digestion step using the RNase-Free

DNase set (Qiagen). RNA was quantified using a NanoDrop 1000 spectrophotometer

(Thermo) and the RNA BR assay kit (Invitrogen) on the Qubit 4 fluorometer (Invitrogen).

RNA quality was assessed using the RNA 6000 Nano kit total RNA assay (Agilent) run on

the 2100 Bioanalyzer instrument (Agilent). Sequencing was carried out using Illumina

NovaSeq 6000 technology and assembled with the Novogene WTS pipeline. Raw reads are

deposited in SRA under BioProject PRNJA1076503. GH18 and GH20 domain containing

genes were identified in the assembled transcriptome by performing blastn searches for

each gene identified in the *R. globosum* JEL800 genome (e-value cutoff = 0.00001).

Concentrated supernatant samples (i.e. secretome) were sent to University of Bristol

Proteomics Facility for comparative proteomics using tandem mass tagging (TMT). Identified

peptides were cross referenced with predicted proteins from mRNA identified within the

transcriptome.

Protein modelling from gene sequences

GH18 and GH20 models were retrieved from the AlphaFold Protein Structure Database

(Jumper et al. 2021, Varadi et al. 2022). Total model volumes were predicted using VADAR

1.8 (Willard et al. 2003) and binding site volumes were predicted using CASTp 3.0 (Tian et

al. 2018) with default settings. The complete model for ORY35945 including the GH18

domain, linker, and CBM5 domain was visualised in UCSF Chimera (Pettersen et al. 2004).

Graphics were edited using Inkscape (http://inkscape.org).

Zoospore chemotaxis and motility

Prior to chemotaxis and motility experiments, *R. globosum* JEL800 was maintained on agar

plates made with PmTG, or BBM with 5mM GlcNAc or chitin for at least 3 generations.

Mature plates were sporulated as described above and 1 mL zoospores spread onto fresh

media and dried to create a lawn. Lawns were incubated at 22°C for 48 hours (PmTG and

GlcNAc) and 24 hours (chitin) to allow sporangia to develop and mature. For chemotaxis

experiments, mature lawns were sporulated with 5mL sterile dH_2O, and after 30 min

zoospores were harvested and passed through a 10 µm sieve, enumerated using a LUNA™

Automated Cell Counter and motility checked using a haemocytometer. Only zoospore

suspensions with motility >80% and ~$1x10^7$ cells/ mL were used for experiments. 2mL

zoospore suspension was placed in a 50mL centrifuge tube and the top sealed with parafilm.

1 mL syringes with 21g needles containing either 150 µL 10mM GlcNAc, 10mM Glucose or

10mM H_2O (control) were held in the zoospore suspension and incubated at 22°C in the

dark. After 1 hour, the contents of the syringe was emptied into a 1.5mL tube and fixed with

glutaraldehyde (final concentration of 0.5%) and zoospores enumerated using a

haemocytometer. For motility experiments, zoospore lawns were sporulated with 3mL sterile

dH_2O, a 1 mL sample was taken immediately, a subsample of this was glutaraldehyde fixed

to determine the starting concentration. Non-motile zoospores were enumerated using a

haemocytometer every 5 min for 1 hour and % motility was calculated from the starting

concentration.

Acknowledgements

We thank Joyce E. Longcore (University of Maine) for providing *R. globosum* JEL800, now

held at the Collection of Zoosporic Eufungi at the University of Michigan (CZEUM), and

information on the isolation of *Podochytrium* sp. JEL0797, *Siphonaria* sp. JEL0065 and

Physocladia obscura JEL0513. We also thank Antonia Feilden (University of York) for useful

suggestions on protein modelling. NC, KB and MC were supported by the European

Research Council (ERC) (MYCO-CARB project; ERC grant agreement no. 772584). DL was

supported by an EnvEast Doctoral Training Partnership (DTP) PhD studentship funded from

the UKRI Natural Environment Research Council (NERC grant no. NE/L002582/1).

References

Barr, D. J. S. 1986. *Allochytridium expandens* rediscovered: Morphology, physiology and
zoospore ultrastructure. Mycologia **78**:439-448.

Berbee, M. L., T. Y. James, and C. Strullu-Derrien. 2017. Early Diverging Fungi: Diversity
and Impact at the Dawn of Terrestrial Life. Annual Review of Microbiology **71**:41-60.

Ciach, M. A., J. Pawłowska, P. Górecki, and A. Muszewska. 2024. The interkingdom
horizontal gene transfer in 44 early diverging fungi boosted their metabolic, adaptive,
and immune capabilities. Evolution Letters.

Courtade, G., Z. Forsberg, E. B. Heggset, V. G. H. Eijsink, and F. L. Aachmann. 2018. The
carbohydrate-binding module and linker of a modular lytic polysaccharide
monooxygenase promote localized cellulose oxidation. The Journal of biological
chemistry **293**:13006-13015.

Duo-Chuan, L. 2006. Review of fungal chitinases. Mycopathologia **161**:345-360.

Farrer, R. A., A. Martel, E. Verbrugghe, A. Abouelleil, R. Ducatelle, J. E. Longcore, T. Y.
James, F. Pasmans, M. C. Fisher, and C. A. Cuomo. 2017. Genomic innovations
linked to infection strategies across emerging pathogenic chytrid fungi. Nat Commun
8:14742.

Funkhouser, J. D., and N. N. Aronson. 2007. Chitinase family GH18: evolutionary insights
from the genomic history of a diverse protein family. BMC Evolutionary Biology **7**:96.

Galindo, L. J., D. S. Milner, S. L. Gomes, and T. A. Richards. 2022. A light-sensing system in
the common ancestor of the fungi. Current Biology **32**:3146-3153.e3143.

Gladieux, P., J. Ropars, H. Badouin, A. Branca, G. Aguileta, D. M. de Vienne, R. C.
Rodríguez de la Vega, S. Branco, and T. Giraud. 2014. Fungal evolutionary
genomics provides insight into the mechanisms of adaptive divergence in
eukaryotes. Mol Ecol **23**:753-773.

Goughenour, K. D., J. Whalin, J. C. Slot, and C. A. Rappleye. 2021. Diversification of Fungal
Chitinases and Their Functional Differentiation in Histoplasma capsulatum. Mol Biol
Evol **38**:1339-1355.

Gow, N. A. R., J.-P. Latge, and C. A. Munro. 2017. The Fungal Cell Wall: Structure,
Biosynthesis, and Function. Microbiology Spectrum **5**:10.1128/microbiolspec.funk-
0035-2016.

Guillard, R. R. L. 1975. Culture of phytoplankton for feeding marine invertebrates. Pages 26-
60 *in* W. L. Smith and M. H. Chanley, editors. Culture of Marine Invertebrate Animals
Plenum Press, New York.

Guy, L., J. R. Kultima, and S. G. E. Andersson. 2010. genoPlotR: comparative gene and
genome visualization in R. Bioinformatics (Oxford, England) **26**:2334-2335.

Jiang, Z., S. Hu, J. Ma, Y. Liu, Z. Qiao, Q. Yan, Y. Gao, and S. Yang. 2021. Crystal structure
of a chitinase (RmChiA) from the thermophilic fungus Rhizomucor miehei with a real
active site tunnel. Biochim Biophys Acta Proteins Proteom **1869**:140709.

Joneson, S., J. E. Stajich, S. H. Shiu, and E. B. Rosenblum. 2011. Genomic transition to
pathogenicity in chytrid fungi. PLoS Pathog **7**:e1002338.

Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K.
Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A.
Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T.
Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska,
T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu,
P. Kohli, and D. Hassabis. 2021. Highly accurate protein structure prediction with
AlphaFold. Nature **596**:583-589.

Junges, Â., J. T. Boldo, B. K. Souza, R. L. Guedes, N. Sbaraini, L. Kmetzsch, C. E.
Thompson, C. C. Staats, L. G. de Almeida, A. T. de Vasconcelos, M. H. Vainstein,
and A. Schrank. 2014. Genomic analyses and transcriptional profiles of the glycoside
hydrolase family 18 genes of the entomopathogenic fungus Metarhizium anisopliae.
PLOS One **9**:e107864.

Karlsson, M., and J. Stenlid. 2008. Comparative evolutionary histories of the fungal chitinase
gene family reveal non-random size expansions and contractions due to adaptive
natural selection. Evol Bioinform Online **4**:47-60.

Katoh, K., and D. M. Standley. 2013. MAFFT Multiple Sequence Alignment Software Version
7: Improvements in Performance and Usability. Molecular Biology and Evolution
30:772-780.

Laundon, D., N. Chrismas, K. Bird, S. Thomas, T. Mock, and M. Cunliffe. 2022. A cellular
and molecular atlas reveals the basis of chytrid development. eLIFE **11**:e73933.

Laundon, D., N. Chrismas, G. Wheeler, and M. Cunliffe. 2020. Chytrid rhizoid
morphogenesis resembles hyphal development in multicellular fungi and is adaptive
to resource availability. Proc. R. Soc. B **287**.

Laundon, D., and M. Cunliffe. 2021. A call for a better understanding of aquatic chytrid
biology. Frontiers in Fungal Biology **2**.

Lespinet, O., Y. I. Wolf, E. V. Koonin, and L. Aravind. 2002. The Role of Lineage-Specific
Gene Family Expansion in the Evolution of Eukaryotes. Genome Research **12**:1048-
1059.

Liu, J., Q. Xu, Y. Wu, D. Sun, J. Zhu, C. Liu, and W. Liu. 2023. Carbohydrate-binding
modules of ChiB and ChiC promote the chitinolytic system of Serratia marcescens
BWL1001. Enzyme Microb Technol **162**:110118.

Maillard, F., A. Kohler, E. Morin, C. Hossann, S. Miyauchi, I. Ziegler-Devin, D. Gérant, N.
Angeli, A. Lipzen, K. Keymanesh, J. Johnson, K. Barry, I. V. Grigoriev, F. M. Martin,
and M. Buée. 2023. Functional genomics gives new insights into the ectomycorrhizal
degradation of chitin. New Phytol **238**:845-858.

Medina, E. M., and N. E. Buchler. 2020. Chytrid fungi. Current Biology **30**:R516-R520.

Miller, M. A., W. Pfeiffer, and T. Schwartz. 2010. Creating the CIPRES Science Gateway for
inference of large phylogenetic trees. 2010 Gateway Computing Environments
Workshop (GCE):1-8.

Moss, A. S., N. S. Reddy, I. M. Dortaj, and M. J. San Francisco. 2008. Chemotaxis of the
amphibian pathogen Batrachochytrium dendrobatidis and its response to a variety of
attractants. Mycologia **100**:1-5.

Murphy, C. L., N. H. Youssef, R. A. Hanafy, M. B. Couger, J. E. Stajich, Y. Wang, K. Baker,
S. S. Dagar, G. W. Griffith, I. F. Farag, T. M. Callaghan, and M. S. Elshahed. 2019.
Horizontal Gene Transfer as an Indispensable Driver for Evolution of
Neocallimastigomycota into a Distinct Gut-Dwelling Fungal Lineage. Appl Environ
Microbiol **85**.

Nagy, L. G., R. Tóth, E. Kiss, J. Slot, A. Gácser, and G. M. Kovács. 2017. Six Key Traits of
Fungi: Their Evolutionary Origins and Genetic Bases. Microbiology Spectrum
5:10.1128/microbiolspec.funk-0036-2016.

Naranjo-Ortiz, M. A., and T. Gabaldón. 2019. Fungal evolution: major ecological adaptations
and evolutionary transitions. Biol Rev Camb Philos Soc **94**:1443-1476.

Payne, C. M., M. G. Resch, L. Chen, M. F. Crowley, M. E. Himmel, L. E. Taylor, 2nd, M.
Sandgren, J. Ståhlberg, I. Stals, Z. Tan, and G. T. Beckham. 2013. Glycosylated
linkers in multimodular lignocellulose-degrading enzymes dynamically bind to
cellulose. Proc Natl Acad Sci U S A **110**:14646-14651.

Petersen, T. N., S. Brunak, G. von Heijne, and H. Nielsen. 2011. SignalP 4.0: discriminating
signal peptides from transmembrane regions. Nat Methods **8**:785-786.

Pettersen, E. F., T. D. Goddard, C. C. Huang, G. S. Couch, D. M. Greenblatt, E. C. Meng,
and T. E. Ferrin. 2004. UCSF Chimera--a visualization system for exploratory
research and analysis. J Comput Chem **25**:1605-1612.

Powell, M. J. 1983. Localization of antimonate-mediated precipitates of cations in zoospores
ofChytriomyces hyalinus. Experimental Mycology **7**:266-277.

Prostak, S. M., K. A. Robinson, M. A. Titus, and L. K. Fritz-Laylin. 2021. The actin networks
of chytrid fungi reveal evolutionary loss of cytoskeletal complexity in the fungal
kingdom. Current Biology **31**:1192-1205.e1196.

Raabe, D., P. Romano, C. Sachs, H. Fabritius, A. Al-Sawalmih, S. B. Yi, G. Servos, and H.
G. Hartwig. 2006. Microstructure and crystallographic texture of the chitin–protein
network in the biological composite material of the exoskeleton of the lobster
Homarus americanus. Materials Science and Engineering: A **421**:143-153.

Richards, T. A., and N. J. Talbot. 2018. Osmotrophy. Curr Biol **28**:R1179-r1180.

Roberts, C., R. Allen, K. E. Bird, and M. Cunliffe. 2020. Chytrid fungi shape bacterial
communities on model particulate organic matter. Biol Lett **16**:20200368.

Scholz, B., F. C. Küpper, W. Vyverman, H. G. Ólafsson, and U. Karsten. 2017.
Chytridiomycosis of Marine Diatoms-The Role of Stress Physiology and Resistance
in Parasite-Host Recognition and Accumulation of Defense Molecules. Mar Drugs **15**.

Sipos, G., A. N. Prasanna, M. C. Walter, E. O'Connor, B. Bálint, K. Krizsán, B. Kiss, J. Hess,
T. Varga, J. Slot, R. Riley, B. Bóka, D. Rigling, K. Barry, J. Lee, S. Mihaltcheva, K.
LaButti, A. Lipzen, R. Waldron, N. M. Moloney, C. Sperisen, L. Kredics, C. Vágvölgyi,
A. Patrignani, D. Fitzpatrick, I. Nagy, S. Doyle, J. B. Anderson, I. V. Grigoriev, U.
Güldener, M. Münsterkötter, and L. G. Nagy. 2017. Genome expansion and lineage-
specific genetic innovations in the forest pathogenic fungi Armillaria. Nat Ecol Evol
1:1931-1941.

Sparrow, F. K. 1960. Aquatic Phycomycetes. The University of Michigan Press, Ann Arbour.

Sprockett, D. D., H. Piontkivska, and C. B. Blackwood. 2011. Evolutionary analysis of
glycosyl hydrolase family 28 (GH28) suggests lineage-specific expansions in
necrotrophic fungal pathogens. Gene **479**:29-36.

Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics **30**:1312-1313.

Tang, B., L. Wei, W. Tang, S. Li, and R. Zhou. 2015. Effect of Linker Flexibility on the
Catalytic Features of a Glycoside Hydrolase Family 45 Endoglucanase from
Rhizopus stolonifer. Applied Biochemistry and Biotechnology **176**:2242-2252.

Thomé, P. C., J. Wolinska, S. V. D. Wyngaert, A. Reñé, D. Ilicic, R. Agha, H.-P. Grossart, E.
Garcés, M. T. Monaghan, and J. F. H. Strassert. 2023. Phylogenomics including new

sequence data of phytoplankton-infecting chytrids reveals multiple independent

lifestyle transitions across the phylum. bioRxiv:2023.2006.2028.546836.

Tian, W., C. Chen, X. Lei, J. Zhao, and J. Liang. 2018. CASTp 3.0: computed atlas of

surface topography of proteins. Nucleic Acids Res **46**:W363-w367.

Varadi, M., S. Anyango, M. Deshpande, S. Nair, C. Natassia, G. Yordanova, D. Yuan, O.

Stroe, G. Wood, A. Laydon, A. Žídek, T. Green, K. Tunyasuvunakool, S. Petersen, J.

Jumper, E. Clancy, R. Green, A. Vora, M. Lutfi, M. Figurnov, A. Cowie, N. Hobbs, P.

Kohli, G. Kleywegt, E. Birney, D. Hassabis, and S. Velankar. 2022. AlphaFold Protein

Structure Database: massively expanding the structural coverage of protein-

sequence space with high-accuracy models. Nucleic Acids Res **50**:D439-d444.

Venard, C. M., K. K. Vasudevan, and T. Stearns. 2020. Cilium axoneme internalization and

degradation in chytrid fungi. Cytoskeleton (Hoboken) **77**:365-378.

Willard, L., A. Ranjan, H. Zhang, H. Monzavi, R. F. Boyko, B. D. Sykes, and D. S. Wishart.

2003. VADAR: a web server for quantitative evaluation of protein structure quality.

Nucleic Acids Res **31**:3316-3319.

Yang, M., M. K. Derbyshire, R. A. Yamashita, and A. Marchler-Bauer. 2020. NCBI's

Conserved Domain Database and Tools for Protein Domain Analysis. Current

Protocols in Bioinformatics **69**:e90.

Figure 1 a) A *Rhizoclosmatium globosum* JEL800 cell with branching rhizoids (r) used for attaching to substrates and sporangium (s) inside which develop multiple zoospores. b-d) *Rhizoclosmatium globosum* JEL800 attached to the *Daphnia* exoskeleton across various levels of magnification. Note rhizoids (r) penetrating the exoskeleton surface.

Figure 2 ai) Comparison of ratios of glycoside hydrolases GH18 chitinases and aii) GH20 β-
N-acetyl-D-hexosaminidases with total GH encoding genes, and aiii) carbohydrate binding
module CBM5 genes with total CBM encoding genes and genome sizes of 36 members of
the Chytridiomycota that occupy diverse ecological niches and utilise a range of substrates
(full labels in Supplementary Figure 1). Point colours indicate the substrate strains were
isolated from, convex hulls link strains isolated from the same substrate; point shapes
indicate order. Maximum likelihood phylogenies of *Rhizoclosmatium globosum* JEL800 b)
GH18 and c) GH20 domains showing architecture of GH domain-containing genes. Genes
encode for proteins with either d) endo- (GH18) or exochitinase (GH18, GH20) activity.

Figure 3 a) Secretome abundance and mRNA (fpkm) for all *Rhizoclosmatium globosum*
JEL800 glycoside hydrolases present in the secretome. b) Relationship of abundance of
proteins in the secretome to equivalent mRNA in the transcriptome c) Relationship of
substrate binding site volume to total protein volume d) AlphaFold predicted model of
ORY35945. Colours are based on conserved domains: the carbohydrate binding module
domain (CBM) (amino acid residues 22 - 36) is shown in yellow and the GH18 domain

641 (amino acid residues 203 - 596) is shown in red. Both are joined by a linker region. Signal

642 peptide region not shown.

644
Figure 4 ai-ii) Chemotactic response of *Rhizoclosmatium globosum* JEL800 zoospores to

the chitin monomer N-acetylglucosamine (GlcNAc) compared to glucose and no-attractant

control from cultures grown with GlcNAc as the sole carbon source and in complex GlcNAc-

free medium PmTG. Significance values (pairwise Wilcoxon test) are indicated as *** = p <

0.001, * = p < 0.05, NS. = not significant. Points are binned at 1/30 range of the data. b)

Loss of motility in *R. globosum* JEL800 zoospores from cultures grown with chitin as the sole

carbon source compared to the complex medium PmTG.