# Clearance of protein aggregates during cell division

Shoukang Du[1,3], Yuhan Wang[1], Bowen Chen[1], Shuangshuang Xie[1], Kuan Yoow Chan[1,3], David C. Hay[4], Ting Gang Chew[1,2,3,5,#]

**Affiliations**

[1]The Zhejiang University-University of Edinburgh Institute, Zhejiang University School of Medicine, Zhejiang University, Haining 314400, China

[2]Department of Cardiology of the Second Affiliated Hospital, Zhejiang University School of Medicine, Zhejiang University, Hangzhou 310058, China

[3]College of Medicine and Veterinary Medicine, The University of Edinburgh, Edinburgh, EH4 2XR, UK

[4]Centre for Regenerative Medicine, Institute for Regeneration and Repair, University of Edinburgh, 5 Little France Drive, Edinburgh EH16 4UU, UK

[5]State Key Laboratory of Biobased Transportation Fuel Technology, Zhejiang University, Hangzhou 310027, China

#Correspondending author: Ting Gang Chew (tinggchew@intl.zju.edu.cn)

**Abstract**

Protein aggregates are spatially organized and regulated in cells to prevent deleterious effects of proteostatic stress. Misfolding of proteins in the endoplasmic reticulum (ER) result in aggregate formation, but how the aggregates are processed especially during cell division is not well understood. Here, we induced proteostatic stress and protein aggregation using a proteostasis reporter, which is prone to misfolding and aggregation in the ER. Unexpectedly, we detected solid-like protein aggregates deposited mainly in the nucleus and surrounded by the ER membrane. The membrane-bound aggregates were then cleared as cells progressed through mitosis and cytokinesis. Aggregate clearance depended on Hsp70 family chaperones in the ER, particularly BiP, and proteasomal activity. The clearance culminated at mitotic exit and required cyclin-dependent kinase 1 (Cdk1) inactivation and ER reorganization but was independent of the anaphase-promoting complex (APC/C). Thus, dividing cells have the capacity to clear protein aggregates to maintain proteostasis in the newly divided cells, which could have implications for human disease development and aging.

**Introduction**

Nascent polypeptides fold into three-dimensional structures to perform their biological functions (Hipp et al., 2019). Misfolded proteins tend to have their hydrophobic residues exposed and result in aggregation of proteins (Hartl et al., 2011). Intramolecular beta-sheets in some proteins are also structural elements prone for protein aggregation (Tyedmers et al., 2010). Protein misfolding and aggregation not only disrupt the native function of the protein but also could interfere other proteins' functions by co-aggregation (Olzscha et al., 2011; Woerner et al., 2016). This challenges proteome

homeostasis (proteostasis) and causes proteostatic stress in cells, which is a main driver of cellular

aging and neurodegenerative disorders (Balch et al., 2008; Labbadia and Morimoto, 2015).

Cells evolve several protein quality control (PQC) systems to maintain their proteostasis in which

components of PQC systems facilitate refolding, degradation, and spatial deposition of misfolded

protein aggregates (Jayaraj et al., 2020). Molecular chaperones are key players in the PQC system.

They are involved to fold nascent polypeptides, target misfolded proteins for degradation by

ubiquitin-proteasome system or autophagy, and promote solubilization of protein aggregates

(Tyedmers et al., 2010). Eukaryotic cells confine misfolded proteins and smaller aggregates into

distinct cellular deposition sites, which reduce the reactivity of these harmful species to the proteome

(Kaganovich et al., 2008). These deposition sites include aggresomes or aggresome-like induced

structures locating near centrosomes, INQ (intranuclear quality control compartment) residing in the

nucleus, cytosolic CytoQ, and the perivacuolar IPOD (insoluble protein deposit) (Kaganovich et al.,

2008). Confinement of protein aggregates in a deposition site facilitates retention of aggregates in

short-lived cells (mother cells) during asymmetric cell division (Aguilaniu et al., 2003; Fuentealba et

al., 2008; Rujano et al., 2006; Singhvi and Garriga, 2009; Zhou et al., 2014).

The endoplasmic reticulum (ER) processes one-third of cellular proteome and possesses stronger

ability to maintain aggregation-prone proteins in a non-toxic state than the cytosol owing to its

specific molecular chaperone environment (Vincenz-Donnelly et al., 2018). Chaperones in the ER

such as HSPA5/BiP regulates the unfolded protein response (UPR), which blocks instant protein

synthesis and activates transcription of genes involved in PQC (Preissler and Ron, 2019; Wiseman

et al., 2022). In addition, the proteostatic stress in the ER triggers ER-associated degradation (ERAD) and ER-phagy to remove misfolded proteins (Mochida and Nakatogawa, 2022; Olzmann et al., 2013). Despite the high capacity of proteostatic stress response in the ER (Rousseau et al., 2004; Vincenz-Donnelly et al., 2018), excessive misfolded proteins still lead to formation of protein aggregates in the ER (Melo et al., 2022; Miyata et al., 2020). In response to protein aggregation in the ER, BiP functions as a disaggregase to promote solubilization of these aggregates. In cells undergoing asymmetric cell division such as in the budding yeast, the ER diffusion barrier and the ER stress surveillance (ERSU) pathway ensure ER protein aggregates are retained preferentially in the short-lived mother cells during cell division (Clay et al., 2014; Pina and Niwa, 2015). How ER protein aggregates are regulated in dividing cells, especially those that do not undergo asymmetric cell division, is less well studied. We addressed the fate of ER protein aggregates in human cells undergoing cell division symmetrically and identified a clearance mechanism of protein aggregates when cells are exiting mitosis.

**Results**

**Targeting a proteostasis reporter to the ER results in protein aggregate formation in the nucleus.**

To investigate how cells maintain their proteostasis during cell division, we employed a proteostasis reporter consists of a firefly luciferase mutant prone for protein mis-folding and protein aggregation fused to a green fluorescent protein (FlucDM-eGFP) (Gupta et al., 2011). Next, we targeted the reporter to the ER by fusing a ER-targeting sequence at its N-terminus (ER-FlucDM-eGFP). Interestingly, we found that ER-FlucDM-eGFP assembled into visible protein aggregates in the

nucleus when stably expressed in mammary epithelial MCF10A cells in the absence of any

perturbation of the proteome stability (Figure 1A). The protein aggregates were below detection limit

of our microscopy in first 2 days after lentivirus transduction but progressively accumulated in cells

and stabilized in the population several days after lentivirus transduction. This contrasted with a

previous study in which the ER-FlucDM-eGFP was transiently expressed in cells and hence did not

allow long term tracking of the protein (Sharma et al., 2018). ER-FlucDM-mCherry localized to the

ER network indicated by the ER membrane protein Sec61β (Figure 1B). Since we detected

aggregate formation in the nucleus in cells expressing ER-FlucDM-eGFP, we tested if targeting

FlucDM-eGFP to the nucleus (NLS-FlucDM-eGFP) would result in the formation of visible protein

aggregates. Our data showed that NLS-FlucDM-eGFP did not form protein aggregates in the nucleus

and was not co-localized with the ER-FlucDM-mCherry, and accumulated lower protein abundance

than ER-FlucDM-eGFP (Figure S1A-D). Luciferase activity measurement showed that ER-FlucDM-

eGFP has significantly reduced enzymatic activity and was not functional possibly due to mis-folding

and protein aggregation (Figure 1C). ER-FlucDM-eGFP was also detected in the insoluble fraction

after heat stress (Figure S1E). Moreover, we observed the colocalization of ER-FlucDM-eGFP

aggregates and thioflavin T (ThT), a universal dye to detect amyloid fibril, further supporting these

aggregates were misfolded protein aggregates (Figure S1F). Thus, expression of ER-FlucDM-eGFP

results in proteostatic stress and allows us to study how protein aggregates are regulated in human

cells.

Formation of protein aggregates in the nucleus by ER-FlucDM-eGFP was not limited to MCF10A

cells. The protein aggregates were also detected in the nucleus when ER-FlucDM-eGFP was

expressed in other cell lines such as A549, MDA-MB-231 and U2OS cells (Figure 1A and S1G).

Similar observations were found when a different N-terminal signal peptide sequence was used

(Figure 1D and S1H). In addition, when ER-FlucDM was replaced by ER-HaloDM, which is a

modified haloalkane dehalogenase prone to mis-folding and aggregation (Melo et al., 2022), we

observed similar protein aggregates in the nucleus (Figures 1D and S1H). However, only very low

number of cells formed aggregates in the nucleus when intact GFP was targeted to the ER (Figure

S1I). Thus, ER-FlucDM-eGFP formed protein aggregates in the nucleus independent of cell line,

fluorescent protein types, specific ER targeting sequence and FlucDM.

Despite localizing in the nucleus, we found that the protein aggregates were surrounded by the ER

membrane as stained by the ER-tracker (Figure 1E). To further confirm this, we sectioned the cells

and used electron microscopy to observe the protein aggregates. We found that the aggregates

were surrounded by a single layer membrane closed to the inner membrane of the nucleus (Figure

1F). Thus, the aggregates are intra-nuclear membranous structure surrounded by the ER membrane.

**Differential and slow turnover of protein aggregates in cells at interphase and mitosis.**

Pathological proteins aggregates such as Z-α1-antitrypsin accumulated in the ER exhibited minimal

turnover (Dickens et al., 2016). To test whether ER-FlucDM-eGFP aggregates has a reduced

dynamics as well, we performed fluorescence recovery after photobleaching (FRAP) in cells

expressing both ER-FlucDM-eGFP and ER-FlucDM-mCherry. The ER-FlucDM-eGFP signal served

as a tracker for the location of aggregates and the ER-FlucDM-mCherry was used to probe the

aggregate dynamics. We photobleached ER-FlucDM-mCherry with 561 nm laser while ensuring ER-

FlucDM-eGFP signal was not affected. In interphase cells, ER-FlucDM-mCherry aggregates

intensity barely recovered after 4.5 min post-bleaching, indicating that ER-FlucDM-mCherry

aggregates possessed minimal turnover similar to that of in other protein aggregates such as Z-α1-

antitrypsin (Figure 2A). Similarly, ER-HaloDM-eGFP aggregates in the nucleus also displayed low

recovery after photobleaching (Figure S2A). When ER-FlucDM-mCherry aggregates were

photobleached in mitotic cells, the recovery of intensity was also much lesser than the ER-FlucDM-

mCherry signal present in the ER network (Figures 2B and 2C). However, ER-FlucDM-mCherry

aggregates in mitotic cells showed a higher recovery than that of in interphase cells (Figure 2D;

aggregates of interphase cells recovered from 0.131 ± 0.093 to 0.157 ± 0.095; aggregates of mitotic

cells recovered from 0.196 ± 0.137 to 0.356 ± 0.151).

Consistently, the single-cell analysis revealed that in mitotic cells, there was an increase of the

recovery intensity of ER-FlucDM-mCherry in late time-point as compared to that of in early time-

points after photobleaching. The increased recovery intensity was not observed in ER-FlucDM-

mCherry aggregates of interphase cells (Figure S2B). Taken together, ER-FlucDM-mCherry

aggregates in the nucleus have low turnover and the aggregate displays increased turnover in mitotic

cells.

**Protein aggregates are cleared during cell division.**

Since aggregates displayed some levels of turnover during mitosis (Figures 2B, 2C and 2D), we next

investigated the aggregate behaviour throughout cell division. To this end, we labelled cells

expressing ER-FlucDM-eGFP with SiR-Tubulin to distinguish cells at various stages based on the

microtubule organization. We categorized cells into interphase and prophase, prometaphase, metaphase, anaphase, telophase and early G1 cells based on their microtubule structures (Figure 3A). Next, we analyzed the number and area of aggregates in cells of each category. Interestingly, we observed that the number and area of aggregates in telophase and early G1 cells were significantly lower than in interphase cells, suggesting that aggregates are cleared gradually when cells progress through mitosis and cytokinesis (Figure 3B).

To test whether this was the case, we employed time-lapse imaging to profile aggregates in cells undergoing division. To increase the number of dividing cells during imaging, we synchronized cells at G2/M and released them into mitosis prior to imaging. We observed that upon entry into mitosis, the ER-FlucDM-eGFP aggregates were released from the nucleus during nuclear envelope breakdown and gradually decreased in numbers as cells progress through mitosis and cytokinesis (Figure 3C and Video 1). In majority of early divided G1 cells, there was essentially no detectable aggregates (Figure 3D). Interestingly, the aggregates released from the nucleus during mitosis were still surrounded by the ER membrane as indicated by the ER-tracker (Figures S3A and S3B). The decrease of aggregates appears to happen specifically in dividing cells as the number and area of aggregates remained largely unchanged in interphase cells (Figures 3C and 3D). Furthermore, aggregates formed in cells expressing ER-HaloDM-eGFP decreased in numbers when cells progress through mitosis and cytokinesis (Figure S3C and Video 2), suggesting that the clearance of protein aggregates during cell division was not specific to ER-FlucDM-eGFP.

**Effects of ER stress inducers on aggregate clearance in mitotic cells**.

Expression of ER-FlucDM-eGFP results in proteostatic stress in cells. Consistently, we observed that gene expression for chaperones and co-chaperones in the ER were upregulated in these cells (Figure 4A), indicating cells involved ER proteostasis control during adaptation. Proteomic analysis of cells expressing ER-FlucDM-GFP showed up-regulation of proteins involved in ER stress response (Figure S4A). We speculated that acute perturbation to ER homeostasis could exacerbate the proteostatic stress in dividing cells expressing ER-FlucDM-eGFP and affect aggregate clearance during mitosis. To test if this was the case, synchronized cells expressing ER-FlucDM-eGFP were released into mitosis in the presence of either DMSO (control) or 1 μM thapsigargin (Thaps), which blocks the ER calcium ion pump and causes ER stress. In cells at prometaphase and metaphase (35 min after release), DMSO and Thaps treatments showed comparable number and area of aggregates in cells (Figures 4B and 4C). However, in cells at the late stage of cell division (65 min after release), there were significantly higher number of aggregates retained in the cytosol of Thaps-treated cells than DMSO (Figures 4B and 4C). Consistently, time-lapse microscopy revealed that ER-FlucDM-eGFP aggregates retained in the cytosol after cytokinesis in Thaps-treated cells, whereas the aggregates were largely cleared in control cells completing cytokinesis (Figures 4D, 4E, S4B and Video 4). Thus, acute treatment of ER stress inducer Thaps in dividing cells prevents clearance of ER-FlucDM-eGFP aggregates.

A previous study showed that cells with protein aggregates treated for hours with Thaps or Tunicamycin (Tuni), which inhibits protein glycosylation in the ER, promotes aggregate clearance in cells (Melo et al., 2022). To test if prolonged treatment of cells expressing ER-FlucDM-eGFP with Tuni could promote aggregate clearance, we pretreated cells blocked at G2/M with varying

concentrations of Tuni for 3 hours and released cells into mitosis. Interestingly, prolonged

pretreatment of Tuni prior to mitosis promoted aggregate clearance in dividing cells (Figure S4C).

Thus, acute treatment of cells expressing ER-FlucDM-eGFP with ER stress inducers prevents

aggregate clearance in dividing cells while prolonged treatment with ER stress inducers promotes

aggregate clearance.

**Hsp70 family protein BiP is required for aggregate clearance during cell division.**

Since we observed that the expression of a chaperone Hsp70 (HSPA5/BiP) was upregulated in cells

expressing ER-FlucDM-eGFP (Figure 4A) and Hsp70 is involved in elimination of protein aggregates

(Melo et al., 2022; Nillegoda et al., 2015), we next tested whether the clearance of ER-FlucDM-eGFP

aggregates in dividing cells was mediated by Hsp70 family proteins. To this end, cells entering

mitosis were treated with either DMSO (control) or VER-155008 (VER), an inhibitor of HSP70's

ATPase activity (Samanta et al., 2021). We found that VER-treated dividing cells fixed after 65

minutes released from the G2/M boundary retained high number of ER-FlucDM-eGFP aggregates

in the cytosol compared to control treatment (Figures 5A and 5B). Time-lapse microscopy showed

that when cells were treated with VER, ER-FlucDM-eGFP aggregates released from the nucleus

upon entry into mitosis were not cleared as to the same extent as control cells (Figure 5C and Video

5).

HSPA5/BiP is a major Hsp70 family member in regulating ER proteostasis. We next tested whether

inhibition of BiP function could affect clearance of aggregates in dividing cells. Similar to VER

treatment, inhibition of BiP's ATPase activity by YUM-70 (Samanta et al., 2021) led to a dose-

dependent accumulation of ER-FlucDM-eGFP aggregates in cells (Figures 5D and 5E). Moreover,

by using immunofluorescence staining, we found a strong colocalization of BiP and ER-FlucDM-

eGFP, consistent with a role of BiP in regulating ER-FlucDM-eGFP aggregates (Figures S5A to S5C).

Collectively, our data demonstrated that the clearance of ER-FlucDM-eGFP aggregates was

regulated by the HSP70 family proteins, particularly BiP.

**Aggregate clearance is a mitotic exit event but is independent of the anaphase promoting**

**complex (APC/C)**.

Protein aggregates are subjected to proteasomal degradation in cells (Tyedmers et al., 2010). To

test if aggregate clearance during cell division involves proteasomes, we inhibited the proteasome

activity with MG132 in dividing cells expressing ER-FlucDM-eGFP. After treating cells released from

the G2/M boundary with low or high concentrations of MG132 for 55 minutes, majority of mitotic cells

were arrested at metaphase and accumulated high number of aggregates as compared to control

cells treated with DMSO (Figures S6A and S6B). Next, we inhibited the proteasome with MG132 in

cells at late anaphase or telophase and examined the aggregates 35 minutes after MG132 treatment

(Figures S6C and S6F). In cells completing cytokinesis, there were significantly high number of

aggregates in the early divided cells treated with MG132 versus the control (Figures S6D, 6E, S6G

and S6H).

Since protein aggregates are cleared when cells progress through mitosis and cytokinesis, which

are the cell cycle stage with low Cyclin B/CDK1 activity, we speculated that inactivation of CDK1

activity in MG132-treated and metaphase-arrested cells could lead to clearance of ER-FlucDM-

eGFP aggregates in the cytosol. To test if this was the case, we released cells into mitosis in the

presence of MG132 for 55 minutes and treated metaphase-arrested cells with RO-3306 to inhibit

Cdk1 activity (Figure 6A). Interestingly, ER-FlucDM-eGFP aggregates were cleared in cells treated

with RO-3306 while the aggregates were retained in cells treated with DMSO (Figures 6B and S6I).

Consistently, live cell imaging demonstrated similar accelerated clearance in Cdk1 inhibited cells

(Figures 6C and 6D). Thus, clearance of the aggregates happens when cells exit from mitosis.

We established that clearance of aggregates during cell division requires proteasomes and CDK

inactivation, it is possible that the anaphase promoting complex (APC/C), which is a E3 ubiquitin

ligase regulating mitotic exit (Watson et al., 2019), is involved in clearing the aggregates. To test if

this was the case, we inhibited APC/C activity using a cocktail of APC/C inhibitors (APC/Ci) consists

of Apcin and ProTAME in dividing cells expressing ER-FlucDM-eGFP (Sackton et al., 2014).

Interestingly, inhibition of APC/C did not affect aggregate clearance as APC/Ci-treated cells that have

arrested at metaphase have comparable number of aggregates as in control DMSO-treated cells

(Figures 6E and 6F). When cells at late anaphase or telophase were treated with APC/Ci, the newly

divided cells have similar number of aggregates as in control cells (Figures 6G, 6H and 6I). Lastly,

we verified the efficacy of APC/C inhibition using cells expressing the APC/C reporter derived from

Geminin (Figure S6J). Consistently, APC/Ci treatment has prevented the removal of APC/C reporter

in newly divided cells indicating that the APC/C was inhibited by the dosage of APC/Ci used in our

experiments (Figure S6K). Taken together, our data showed that clearance of ER-FlucDM-eGFP

aggregates happens when cells exit from mitosis, is proteasome dependent but does not involve

APC/C.

To examine if ER-FlucDM-eGFP decreased in abundance while cells progressing through mitosis, we inhibited protein translation using cycloheximide (CHX) in cells released from G2/M boundary. Short half-life proteins like p53 showed a rapid decrease of abundance upon CHX treatment, indicating CHX prevented new protein synthesis in our treatment (Figure S7A and S7B). ER-FlucDM-eGFP did not show significant change of its abundance after CHX treatment, suggesting that ER-FlucDM-eGFP aggregates were not cleared by the protein degradation mechanism directly (Figure S7A and S7C). Consistently, when the total GFP intensity of ER-FlucDM-eGFP was quantitated in dividing cells 20-30 min (early) or 85-95 min (late) released from G2/M boundary, there was no significant change of GFP fluorescence intensity between early and late dividing cells (Figure S7D), further supported that the protein degradation mechanism was not responsible for the ER-FlucDM-eGFP aggregate clearance.

**Aggregate clearance depends on the ER reorganization during cell division**

Since we showed that ER-FlucDM-eGFP aggregates are not primarily cleared by the protein degradation mechanism, we considered the possibility that aggregate clearance requires reorganization of the ER that happens in cells exiting mitosis. To this end, we used CRISPR interference (CRISPRi) technique to knock down ER membrane fusion proteins Atlastin 2 (ATL2) and Atlastin 3 (ATL3) in cells expressing ER-FlucDM-eGFP and quantitated the aggregate numbers in cells exiting mitosis (65 min after release). Interestingly, dividing cells depleted of ATL2 and ATL3 accumulated more ER-FlucDM-eGFP aggregates than in control cells (Figure 7A and B).

The mitotic kinase Aurora A has been recently shown to remodel the ER network during mitosis (Zhang et al., 2024). We next tested if blocking Aurora A activity could affect aggregate clearance. When mitotic cells expressing ER-FlucDM-eGFP were treated with Aurora A inhibitor, MLN-8237, aggregate clearance was prevented compared to cells treated with DMSO (Figure 7C and D). Since the ER reorganization involves microtubule dynamics, we also perturbed microtubule networks by treating cells expressing ER-FlucDM-eGFP by nocodazole (Figure 7E and F). Our data showed that mitotic cells treated with nocodazole accumulated higher number of aggregates. Taken together, we found that clearance of ER-FlucDM-eGFP aggregates involves reorganization of the ER during mitotic exit (Lee et al., 1989).

**Discussion**

We showed that stably expression of ER-FlucDM-eGFP leads to proteostatic stress and formation of protein aggregates in human cells without the use of additional proteostatic stressors. ER-FlucDM-eGFP aggregates may derive from the invagination of the inner nuclear membrane. Morris et al reported a similar aggregate upon BiP overexpression (Morris et al., 1997). This fluorescent based live-cell imaging reporter enables us to study how protein aggregates are regulated in dividing cells, which are sensitive to genetic or environmental perturbations. Conventional methods that use proteasomal inhibitors to cause proteome imbalance are not ideal to investigate proteostasis in dividing cells given an essential role of protesomes in driving mitotic exit (Ghislain et al., 1993; Glotzer et al., 1991). Additionally, methods such as heat shock also perturb mitotic progression, especially in human cells (Kakihana et al., 2019). By using ER-FlucDM-eGFP that does not perturb cell cycle progression significantly, we reveal an unexpected process of clearing protein aggregates

in cells progressing through mitosis and cytokinesis.

Previous studies showed that the proteome of mitotic cells have higher structural stability and less

aggregation prone compared to that of in interphase cells (Becher et al., 2018; Wirth et al., 2013).

Post-translational modifications such as phosphorylation of mitotic proteins are suggested to be

responsible for the protein stability during cell division (Becher et al., 2018). The aggregate clearance

we identified in this study could represent an active mechanism that increases proteome stability in

dividing cells. We found that aggregate clearance requires Cdk1 inactivation, which is a hallmark of

mitotic exit, however, does not involve APC/C that is important in driving mitotic exit through protein

degradation. It is possible that APC/C targets mostly cytosolic proteins whereas ER-FlucDM-eGFP

aggregates are confined in a membrane during mitosis. Furthermore, the clearance of protein

aggregates coincides with ER reorganization that happens throughout mitosis and cytokinesis, which

may contribute to the aggregate clearance (Bergman et al., 2015; Schlaitz et al., 2013). A previous

study showed that the size of protein aggregates is limited by the ER tubules (Parashar et al., 2021).

In fact, when the ER reorganization is perturbed by depleting ER membrane fusion proteins ATL2

and 3, clearance of ER-FlucDM-eGFP aggregates is affected. Consistently, ER-FlucDM-eGFP

proteins do not turn over significantly in cells progressing through mitosis and cytokinesis. Thus, it is

less likely that ER-FlucDM-eGFP is ubiquitinated by APC/C and targeted for protein degradation.

Misfolded or damaged proteins form protein aggregates and are spatially organized into specific

subcellular sites to sequester them from other cellular processes (Hill et al., 2017). These aggregate

deposition sites are localized in the cytosol or in the nucleus to confine misfolded proteins and

aggregates for elimination from cells (Kaganovich et al., 2008). Similarly, we identified confinement

of ER-FlucDM-eGFP aggregates in the nucleus and are surrounded by a single layer membrane.

These intranuclear membranous structures contain ER membranes and BiP, which is a key Hsp70

family chaperone in the ER and translocates into the nucleus under stress (Liu et al., 2023). How

protein aggregates are targeted and assembled into the intranuclear membranous structure awaits

future investigation. Interestingly, upon entry into mitosis, concomitant with the nuclear membrane

breakdown, the membranous structure is released from the nucleus and the aggregates are cleared

when cells progress though mitosis till early G1 phase of the daughter cells. The confinement and

clearance of protein aggregates during cell division presumably protect dividing cells from harmful

effects of protein misfolding and aggregation and to ensure reliable inheritance of genetic materials.

It has been reported that aggregates are asymmetrically retained in one of the daughter cells that

are usually short-lived (Rujano et al., 2006). The aggregate clearance mechanism we report here

may provide an additional cellular defense strategy to overcome proteostatic stress and to maintain

proteome integrity during cell division. Proteomic analysis of cells expressing ER-FlucDM-eGFP

identified up-regulation of multiple proteins involved in ER stress response, indicating that cells

experience significant proteostatic stress upon expression of ER-FlucDM-eGFP. Moreover, several

proteins regulating cell cycle displayed higher abundance in the proteome, which together with

proteostatic stress indicate a perturbed cellular health.

Hsp70 chaperones regulate proteostasis by preventing protein aggregation, promoting

disaggregation and subjecting solubilized aggregates to degradation or refolding (Rosenzweig et al.,

2019). Our proteomic analysis revealed significant expression of BiP (HSPA5) in cells expressing

ER-FlucDM-eGFP but not other Hsp70 family members. A recent study showed that BiP solubilizes aggregates and drives aggregate clearance in the ER (Melo et al., 2022). Our study of mitotic aggregate clearance in dividing cells further extend the role of BiP in disaggregation. Consistently, we found that Cdk1 inactivation in the presence of the proteasomal inhibitor could still lead to aggregate clearance, suggesting a partial role of the proteosome in clearing aggregates and the disaggregation breaks down the aggregates into smaller and possibly soluble species. The role of proteasome in regulating ER-FlucDM-eGFP aggregate clearance appears indirect as ER-FlucDM-eGFP proteins are stable throughout cell division. It is possible that players involved in ER-FlucDM-eGFP aggregate clearance are regulated by proteasomal degradation, which are affected when proteasomes are inhibited. Interestingly, Melo et al. showed that aggregate clearance is promoted by treating ER-cells with ER stressors as the treatment triggers UPR, which in turn leads to higher activity of BiP to cope with the ER stress. We observed differential responses of aggregate clearance to ER stressors depending on the duration of the treatments. It is possible that acute treatment of the ER stressor on dividing cells does not provide sufficient time for cells to accumulate abundant amount of BiP given that mitotic cells are not very transcriptionally active (Palozola et al., 2017). Also, degradation of BiP could be accelerated upon the treatment of ER stressors, which in turn undermines the abundance of BiP in disaggregation (Shim et al., 2018). Thus, short term treatment of the ER stressor in cells exacerbates the ER stress and affects aggregate clearance during cell division, whereas longer treatment of the ER stressor promotes aggregate clearance. Mechanistic understanding of aggregate clearance during cell division could offer new insights to understand disaggregation in proteostasis control.

**Materials and methods**

**Cell lines and culture**

MCF10A cells were cultured in DMEM F-12 (Sigma) supplemented with 1% Glutamax (Gibco), 5% horse serum (Biological Industries), 20 ng/ml EGF (Gibco), 0.5 mg/ml Hydrocortisone (MedChemExpress), 100 ng/ml Cholera toxin (Sigma), 10 µg/ml Insulin (Biological Industries), and 1% Pen/Strep (Biological Industries). U2OS, A549, and HEK293Ta cells were maintained in DMEM (Sigma) with Glutamax (Gibco), 6% fetal bovine serum (Biological Industries), and 1% Pen/Strep (Biological Industries). MDA-MB-231 cells were cultured in DMEM F-12 (Sigma) with 1% Glutamax (Gibco) and 1% Pen/Strep (Biological Industries). All cell cultures were maintained at 37°C with 5% $CO_2$ in a humidified incubator. Cells expressing ER-FlucDM-eGFP, ER-eGFP, ER-HaloDM-eGFP, ER-FlucWT-eGFP, and ER$^{FSHR}$-FlucDM-eGFP, ER-FlucDM-mCherry, NLS-FlucDM-eGFP were generated through lentivirus transduction and selected with 5 µg/mL puromycin (Sangon Biotech). MCF10A cells expressing mStayGold-Sec61β were prepared by lentivirus transduction, selected with 50 µg/mL hygromycin B and only used in three days after transduction (Ando et al., 2024). MCF10A cells expressing CRISPRi plasmids to knock down ATL2 and ATL3 were selected by 6 µg/mL Blasticidin (Solarbio, 703X023) and only used in five days after transduction. MCF10A cells co-expressing ER-FlucDM-eGFP and ER-FlucDM-mCherry for FRAP or co-expressing ER-FlucDM-mCherry and mGreenLantern-hGeminin (degron) or co-expressing ER-FlucDM-mCherry and NLS-FlucDM-eGFP were generated by lentivirus transduction and selected by 5 ug/mL puromycin and followed by fluorescence-activated cell sorting (FACS) to enrich double-positive cells.

**Lentivirus packaging**

Lentiviral particles were generated using the 3rd generation lentiviral packaging system. Packaging plasmids pRSV-Rev, pMDLg/pRRE, pMD2.G (kindly provided by Didier Trono) and the transfer plasmid were chemically transfected using GeneTwin transfection reagent (Biomed, TG101) into the HEK293Ta packaging cell line (Genecopoeia, LT008). The cell culture medium was harvested after 2 days and filtered through a 0.45 μm filter. The filtrate was concentrated using a 5X precipitation solution (250 g/L PEG 8,000 and 43.83 g/L NaCl in ddH$_2$O) for overnight, and precipitated by centrifuging at 4000 × g for 25 min at 4°C. Lentiviral particles were resuspended in phosphate buffered saline (PBS) and added to the culture medium, which were supplemented with 1 μg/mL polybrene (HanBio).

**Lentivirus expression constructs**

Lentiviral transfer plasmids used in this study were listed below: pTGL0563 containing FlucDM-eGFP in which FlucDM was mutated from FlucWT (a gift from Dr. Mikael Bjorklund); pTGL0645 containing ER-FlucDM-eGFP-KDEL; pTGL0662 containing NLS-FlucDM-eGFP; pTGL0673 containing ER-FlucDM-mCherry-KDEL; pTGL0698 containing ER-FlucWT-eGFP-KDEL; pTGL0703 containing ER-HaloDM-eGFP-KDEL in which HaloDM was mutated from HaloWT; pTGL0707 containing ER-HaloWT-eGFP-KDEL; pTGL0733 containing ER-eGFP-KDEL; pTGL0743 containing ER[FSHR]-FlucDM-eGFP; pTGL0828 containing mGreenLantern-hGeminin (degron); pTGL0891 was the empty plasmid used in constructing pTGL0645 and used as a control for proteomics; pTGL0914 containing mStayGold-Sec61β; pTGL0386 was the empty plasmid in constructing CRISPRi plasmids; pCRISPRi0249 to knockdown ATL2 containing gRNA 5' GAGGGCAGCAACCGCACCAG 3'; pCRISPRi0252 to knockdown ATL3 containing gRNA 5' GAGCAGGGGTGCAGAGGAGA 3'.

**Quantitative real-time PCR (qPCR)**

Total RNA was extracted from cells using the FastPure Cell/Tissue Total RNA Isolation Kit V2 (Vazyme, RC112-01). Subsequently, the RNA was reversely transcribed into complementary DNA (cDNA) using the HiScript II Q RT SuperMix for qPCR (Vazyme, R223-01). The levels of the cDNAs were quantified using real-time PCR with the ChamQ Universal SYBR qPCR Master Mix (Vazyme, Q711-02). The PCR reaction mix was prepared on a hard-shell PCR plate (Bio-Rad, HSP9655), sealed with Microseal 'B' seals (Bio-Rad, MSB1001), and conducted in the CFX96 Touch Real-Time PCR Detection System (Bio-RAD, C1000). Primers used in the qPCR: GAPDH, 5' CAGGAGGCATTGCTGATGAT 3' and 5' GAAGGCTGGGGCTCATTT 3'; HSPA5, 5' CACAGTGGTGCCTACCAAGA 3' and 5' TGTCTTTTGTCAGGGGTCTTT 3'; Calreticulin, 5' ATAAAGGTTTGCAGACAAGC 3' and 5' CCACAGTCGATGTTCTGCTC 3'; HSP90B1, 5' TCCAGCAGAAAAGAGGCTGA 3' and 5' CAAATTCGGGAAGGGCCTGA 3'; CANX, 5' GCACCTATTCTGGAGGCGAG 3' and 5' ACAGCAACCACTTCCCTTCC 3'; P4HB, 5' TTCAGGAATGGAGACACGGC 3' and 5' TCCACGTCCTTGAAGAAGCC 3'; DNAJB9, 5' GTGGAGGAGCAGCAGTAGTC 3' and 5' CGCTCTGATGCCGATTTTGG 3'; ATL2, 5' CTGGTTCCATTGCTGCTTGC 3' and 5' CTTCAGCTGTTGCCTGAAGC 3'; ATL3, 5' TCACCCCAAGTCCATGCTTC 3' and 5' CTCCCCCACAAACCTCTTCC 3'.. The $C_T$ value for GAPDH was used for normalization to obtain the relative expression level.

Protein immunoblotting

Adherent cells were lysed with 1x LDS sample buffer (Beyotime) and boiled at 95°C for 10 min. The

1x LDS sample buffer was prepared by diluting 4 X LDS sample buffer (Beyotime) in 100 mM Tris-HCl (pH 7.4) with 2.5% v/v β-mercaptoethanol. Protein samples were separated by pre-cast SDS-PAGE gels (GenScript) and blotted to a PVDF membrane (EMD Millipore). Blots were blocked with Quick Block buffer (Beyotime) for at least 20 min at room temperature and were incubated at 4°C overnight with the primary antibody. After rinsing blots with TBS containing 0.05% Tween-20 for four times (5 min each time), blots were incubated at room temperature with the secondary antibody for 1h, followed by washing with TBS containing 0.05% Tween-20. Odyssey CLX was employed to image the blots and LI-COR Image Studio software was used to quantify the band intensity. To incubate different primary antibodies in the same blot, the blot was vigorously rinsed with the stripping buffer (CWBIO, 01427/12724) and washed 3 times with TBS containing 0.05% Tween-20 and processed as described before.

To examine the solubility of ER-FlucDM-eGFP in Figure S1E, the lysis buffer containing 100 mM Tris-HCl (pH 7.4), 1% protease inhibitor cocktail (Lablead, C0101) and different concentrations of NP-40 (Sigma) was used. To prepare supernatant and pellet fractions of ER-FlucDM-eGFP, cells were lysed with the lysis buffer containing 0.2%, 0.5% or 1.0% v/v NP-40. The lysate was then centrifuged at 14,000 ×g for 10 min at 4°C and the supernatant was collected. The cell pellet was further lysed with 1x LDS buffer on ice for 40 min and vortexed vigorously for 10 seconds every 10 min.

Primary antibodies used were: 1:3000 anti-GFP (Huabio, ET1601-13); 1:5000 anti-β-actin (Genscript, A00702); 1:1000 anti-Cyclin B1 (Proteintech, 55004-1-AP); 1:1000 anti-p53 (Huabio, ET1601-13).

Secondary antibodies were goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody conjugated to Alexa Fluor™ Plus 800 (Invitrogen, A32735) and goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody conjugated to Alexa Fluor™ Plus 680 (Invitrogen, A32729)

Proteomics

MCF10A cells transduced with pTGL0891 (empty vector) or pTGL0645 (ER-FlucDM-eGFP) for six days were trypsinized, centrifuged and washed twice with ice-cold PBS. The pellets were quickly frozen by liquid nitrogen and then sent to Tsingke (Wuhan, China) for further processing for proteomics. In brief, cell pellets were lysed with buffer containing 8 M urea, 1 mM PMSF and 2 mM EDTA. Samples were subjected to ultrasound sonication on ice for five minutes and followed by centrifugation at 15,000 xg for 10 minutes. The supernatants were then collected, and their concentrations were measured using BCA method (Beyotime). Solutions containing 100 $\mu$g was topped up to 200 µl with 8M urea and DTT (final concentration 5 mM) was added before incubation at 37°C for 45 min. Protein samples were alkylated using iodoacetamide (final concentration 11 mM) in a dark room at room temperature for 15 min. The solutions were then subjected to trypsin digestion and further analyzed using LC-MS/MS analysis. The MS/MS data were processed using DIA-NN (v1.8.1). The protein quantification of DIA-NN software is done by MaxLFQ algorithm.

**Luciferase activity test**

Each well of 96-well plates (Biosharp, BS-MP-96W for luminescence, BS-MP-96B for fluorescence) was seeded with 10,000 cells in 100 µL of medium. After incubation for 18 hours, 5 µM MG132 was added to the MG132+FlucDM-eGFP group, while other untreated wells remained unchanged.

Following an additional 6 hours, cells were washed once with PBS.

For luminescence detection, cell lysis was achieved by adding 50 μL of Steady-Glo Luciferase Assay

System buffer (Promega, E2510) to the wells, followed by a 15-minute incubation at room

temperature. Luminescence in each well was then recorded three times over 5 minutes using the

Spark® Multimode Microplate Reader (Tecan, 1000 ms exposure). Luminescence intensity in each

well was calculated by normalizing these three measurements and subtracting the background signal

from empty wells.

To determine abundance of luciferase in each treatments, $1 \times 10^5$ MCF10A cells were seeded in 12-

well for 24 h. The cells were lysed with 1x LDS sample buffer (Beyotime) and protein immunoblotting

using anti-β-actin and anti-GFP, respectively was performed. β-actin was first used to normalized the

GFP intensity, and the value was used to normalize the luminescence to calculate the luciferase

activity.

**Mitotic cell preparation**

To increase the quantity of dividing cells, cells were first seeded on imaging chambers (ibidi, 80826)

and were synchronized to the G2/M boundary by treating cells with 7.5 μm RO-3306 (Selleckchem,

S7747) for 16 to 20 hours. Cells were released to enter into mitosis synchronously by washing once

with pre-warmed fresh medium.

**Drug treatments**

507 The following drugs were used in this study: Thapsigargin (Abcam, ab120286), Tunicamycin (GlpBio,

508 GC16738), MG132 (CSN Pharm, CSN11436), YUM70 (Aladdin, Y413413), VER-155008 (CSN

509 Pharm, CSN13116), Apcin (Sigma, SML1503), proTAME (Cayman Chemical, 25835),

510 Cycloheximide (AbMole, M4879), Nocodazole (Beyotime, S1765), MLN-8237 (TargetMol, T2241).

512 **Immunofluorescence staining**

513 For HSPA5/BiP staining, MCF10A cells expressing ER-FlucDM-eGFP were seeded in 8-well ibidi

514 plates, fixed with 100% methanol for 10 minutes and permeabilized with 0.5% Triton X-100 for 15

515 minutes. The fixed cells were blocked with 5% FBS in PBS for 30 minutes. Subsequently, cells were

516 incubated with the mouse anti-BiP antibody (HuaBio, HA601076) at 1:200 dilution in 1% FBS

517 overnight at 4°C. The secondary antibody used was Donkey Anti-Mouse IgG H&L (Alexa Fluor® 568,

518 ab175472, Abcam) at 1:2000 dilution for one hour. For Tubulin staining, cells were fixed using 4%

519 PFA for 20 minutes and permeabilized with 0.5% Triton X-100 for 15 minutes. After blocking the fixed

520 cells with 5% FBS in PBS for 30 minutes, cells were stained with the mouse anti-$\alpha$-tubulin

521 monoclonal antibody (Proteintech, 66031-1-IG) at 1:200 dilution. The secondary antibody used was

522 Donkey Anti-Mouse IgG H&L (Alexa Fluor® 568, ab175472, Abcam) at 1:2000 dilution for one hour.

523 To stain DNA, DAPI Staining Solution (Sangon Biotech, E607303, 5 mg/L) was diluted in PBS to 50

524 ng/mL and applied to cells for 10 minutes. After the DAPI solution was removed, fixed cells were

525 washed with PBS twice before imaging.

527 **Transmission Electron Microscopy (TEM)**

528 To prepare samples for TEM, MCF10A cells expressing ER-FlucDM-eGFP were synchronized to the

G2/M boundary by treating cells with RO-3306 for 16 to 20 h and were released into mitosis by

removing RO-3306. After 45 min released from the G2/M boundary, cells were trypsinized for 10 min

before centrifugation. The cell pellet was fixed by 2.5% paraformaldehyde (in PBS) for 30 min at

room temperature before overnight at 4°C. Fixed pellet was further fixed by 1% osmium tetroxide

and then 2% uranium acetate. Ethanol and acetone were used to dehydrate samples before

embedding them into epoxy resin. Sectioned samples were observed by Tecnai G2 spirit 120kV

transmission electron microscopy (Thermo FEI).

**Live-cell staining**

The live-cell staining dyes used are as follow: 100 nM SiR-Tubulin (Cytoskeleton, CY-SC002), 1 µM

ER-tracker Red (Beyotime, C1041S) or 10 µM Thioflavin T (ThT) in the medium to label cells for

about 30 min in the incubator before imaging. For ThT staining, about ten percent aggregates could

be stained by ThT in ER-FlucDM-mCherry cells.

**Spinning-disk confocal microscopy imaging**

The hardware configuration of the spinning-disk confocal microscopy system was as described

previously (Wang et al., 2023). Imaging utilized a z-step size of 0.5 µm, with a x-y plane resolution

of 183.3 nm/pixel for the 60× lens and 110 nm/pixel for the 100× lens. Fluorophores were excited

using laser lines at wavelengths of 405, 488, 561, or 640 nm.

For FRAP analysis, images were acquired with 488 nm and 561 nm laser lines through the 100×

lens. The procedure involved acquiring three pre-bleach images every 30 seconds, followed by

bleaching the mCherry signal using the 561 nm laser. Subsequently, one image was acquired

immediately post-bleaching, followed by ten images acquired every 30 seconds. In total, fourteen

time-point images were acquired, and the total duration of each FRAP experiment was 6.5 minutes.

**Image analysis and processing**

Fiji was used to process images. Z-stack images for quantification were first projected using max-

intensity projection (Fiji/image/stacks/Z project). To quantify aggregates, projected images were

classified by Trainable Weka Segmentation (Fiji/Plugins/Segmentation/Trainable Weka

Segmentation). The segmented images were processed by using the ImageJ Macro commands:

run("8-bit"); setOption("BlackBackground", true); run("Convert to Mask"); run("Watershed") before

using Analyze Particles (Fiji/Analyze/Analyze Particles). For time-lapse images except images from

FRAP, the segmented images were processed by using the ImageJ Macro commands: run("8-bit");

setThreshold(0, 120); run("Make Binary", "method=Default background=Light black"); run("Invert

LUT"); run("Watershed", "stack") before using Analyze Particles (Fiji/Analyze/Analyze Particles).

To quantify total GFP intensities in live cells, Z-stack images (24 $\mu$m) were acquired and projected

using sum slices projection. Cell outlines were segmented using polygon selection in Fiji. To

quantitate the background intensity, the mean intensity of a small dark region of the field-of-view was

measured and multiplied by the area of the segmented cell. The background was subtracted from

the GFP intensity of segmented cells.

For FRAP analysis of ER-eGFP and ER-HaloDM-eGFP, region-of-interests (ROI) in projected

images were manually selected and the average fluorescence intensity of ROIs was measured in Fiji. For FRAP analysis of aggregates in cells co-expressing ER-FlucDM-eGFP and ER-FlucDM-mCherry (both interphase and mitosis), the aggregates were first segmented by Weka Segmentation. The unsegmented images from the mCherry channel were processed by Analyze Particles (Fiji/Analyze/Analyze Particles) based on the corresponding segmented images from the GFP channel. For fluorescence intensity quantification, background in all FRAP experiments were subtracted by one hundreds.

**Quantification and statistical analysis**

Datasets were analysed by Student's t test or Mann-Whitney test. Prism 8 (GraphPad) was used for the statistical analysis and data plotting. Statistical details are indicated in the figure legends.

**Author contributions**

S.D conceived the study, designed and performed experiments, and analyzed data. Y.W, B.C, S.X, K.Y.C provided some reagents used in the study. D.H participated in the discussion during project development. T.G.C. conceived the study, designed experiments and supervised the study. D.S and T.G.C prepared the manuscript with inputs from D.H.

**Acknowledgements**

This study was funded by National Natural Science Foundation of China (NSFC) RFIS-II grant (32350610247) and NFSC grant (32270770) given to T.G.C. We thank Xiaoxia Wan and Chenyu Yang in the Center of Cryo-Electron Microscopy (CCEM), Zhejiang University for their technical

assistance on Transmission Electron Microscopy. We thank Xuqi Chen, Xukai Gao and Zihan Chu

for participating in the early stage of the project. We thank Dr. Mike Shipston for participating in the

discussion. We thank all lab members for discussion.

**Conflict of interest**

The authors declare that they have no conflict of interest.

**Figure legends**

**Figure 1. ER-FlucDM-eGFP forms protein aggregates in the nucleus.**

(A) Representative images of ER-FlucDM-eGFP expressed in different cell lines. Maximum

projected images were shown.

(B) Microscopy images of cells expressing ER-FlucDM-mCherry and mStayGold-Sec61β. Single Z-

slice images were shown.

(C) Quantification of the luciferase activity in MCF10A cells expressing different constructs. Unpaired

t-test. Data were collected from three independent experiments. Abundance of GFP fusion

proteins was determined using protein blotting and was used in normalization.

(D) Representative images to show MCF10A cells expressing ER-FlucDM-mCherry, ER[FSHR]-

FlucDM-eGFP (signal peptide sequence from the N-terminus of FSHR), ER-HaloDM-eGFP or

ER-eGFP. Maximum projected images were shown.

(E) Images of ER-FlucDM-eGFP aggregates surrounded by the ER membrane in a MCF10A cell.

Single Z-slice images were shown.

(F) TEM images of sectioned MCF10A interphase cells expressing ER-FlucDM-eGFP.

Data represent mean + SD. Scale bar, 10 µm (A to E).

**Figure 2. Differential aggregate dynamics in cells at interphase and mitosis.**

(A) Representative images of interphase cells co-expressing ER-FlucDM-eGFP and ER-FlucDM-

mCherry before and after photobleaching. Yellow rectangles indicate bleached aggregates.

Maximum projected images were shown.

(B) Representative images of mitotic cells co-expressing ER-FlucDM-eGFP and ER-FlucDM-

mCherry before and after photobleaching. Yellow circles indicate bleached aggregates.

Maximum projected images were shown.

(C) Recovery curves of ER-FlucDM-mCherry intensity after photobleaching. 14 ROIs from the ER of

interphase cells, 14 aggregates from mitotic cells, 26 aggregates from interphase cells. Data

represent mean + SD. A small region of ER-FlucDM-mCherry in the ER network was chosen as

a control for photobleaching.

(D) Quantification of the ER-FlucDM-mCherry intensity of bleached aggregates right after bleaching

and after 5 min of recovery. 0 min and 5 min refer to the time points in (C). Paired t-test (two-

tailed).

Data were collected from four independent experiments. Scale bar, 10 µm.

**Figure 3. Aggregates are cleared during cell division.**

(A) Representative images of dividing ER-FlucDM-eGFP cells at different cell cycle phases. Grey

scale bar: telophase and early G1 cells; black scale bar: interphase and prophase, prometaphase,

metaphase, anaphase. Maximum projected images were shown. Scale bar, 10 µm.

(B) Quantification of ER-FlucDM-eGFP aggregates from dividing cells. Count and area are individual cell data. Each grey ring represents aggregates from one cell. Interphase and prophase, 91 cells; prometaphase, 45 cells; metaphase, 60 cells; anaphase, 20 cells; telophase and early G1, 73 cells. From three independent experiments. Unpaired two side Mann-Whitney U-test. Data represent mean ± SD

(C) Representative time-lapse images of cells expressing ER-FlucDM-eGFP released from the G2/M boundary. NEB (nuclear envelop breakdown) is set to be t = 0 min. Maximum projected images were shown. Scale bar, 10 $\mu$m.

(D) Quantification of aggregate number and area of dividing cells expressing ER-FlucDM-eGFP. Left panel: data of individual cells. Right panel: mean values + SD of single-cell data (left panel). Time point of NEB in each cell was set to 0 min. Aggregate count and total area of each cell at each time point is normalized to its own value at t = 0 min. Aggregate count data are from 61 interphase and 73 dividing cells of four independent experiments. Aggregate area of each cell is calculated from 32 interphase and 34 dividing cells of two independent experiments.

**Figure 4. Effects of ER stressors on aggregate clearance.**

(A) Relative gene expression levels of chaperones and co-chaperones in cells expressing ER-FlucDM-eGFP. Four independent experiments.

(B) Images of cells treated with DMSO or 1 µM Thaps and fixed at 35 min or 65 min after released from the G2/M boundary. Maximum projected images were shown.

(C) Quantification of cells treated with DMSO or 1 µM Thaps and fixed at 35 min or 65 min after

released from the G2/M boundary. 35 min_DMSO, 207 cells; 35 min_Thaps, 173 cells;

65min_DMSO, 244 cells; 65 min_Thaps, 209 cells. Unpaired two side Mann-Whitney U-test. The

solid black line in violin plot indicates median of data in the column. Gray dotted lines are quartiles.

Each light pink circle indicates measurements of individual cells.

(D) Representative time-lapse images of cells expressing ER-FlucDM-eGFP treated with DMSO or

1 µM Thaps after released from the G2/M boundary. Maximum projected images were shown.

(E) Quantification of cells expressing ER-FlucDM-eGFP treated with DMSO or 1 µM Thaps. Time

point of NEB in each cell was set to 0 min. Aggregates count and total area of each cell is

normalized to its own value at t = 0 min. Black dotted lines are 90 min since NEB. DMSO, 89

cells; Thaps, 65 cells.

Scale bar, 10 µm. Three (C and E) or four (A) independent experiments.

**Figure 5. Inhibition of HSP70 and BiP prevents aggregate clearance.**

(A) Representative images of cells expressing ER-FlucDM-eGFP treated with DMSO or 50 µM VER

and fixed at 65 min after released from the G2/M boundary. Maximum projected images were

shown.

(B) Quantification of cells expressing ER-FlucDM-eGFP treated with DMSO or 50 µM VER and fixed

at 65 min after released from the G2/M boundary. DMSO, 357 cells; VER, 465 cells.

(C) Time-lapse images of dividing cells expressing ER-FlucDM-eGFP treated with DMSO or 50 µM

VER. Maximum projected images were shown.

(D) Representative images of cells expressing ER-FlucDM-eGFP treated with DMSO or different

concentrations of YUM70. Maximum projected images were shown.

(E) Quantification of cells expressing ER-FlucDM-eGFP treated with DMSO or YUM70 and fixed at 65 min after released from the G2/M boundary. DMSO, 310 cells; 5 µM YUM70, 258 cells; 20 µM YUM70, 215 cells; 50 µM YUM70, 209 cells; 100 µM YUM70, 287 cells.

Scale bar, 10 µm. Unpaired two side Mann-Whitney U-test. Each solid line in violin plot indicates median of data in the column. Gray dotted lines are quartiles. A, B, D and E are from three independent experiments. C is from two independent experiments

**Figure 6. Effects of MG132, APCi and RO-3306 (Cdk1i) on aggregate clearance.**

(A) Representative images of cells expressing ER-FlucDM-eGFP arrested at mitosis by MG132 and further treated with Cdk1i to induce mitotic exit. Cells were first treated with 25 µM MG132 for 55 min and mitotic arrested cells were further treated with 25 µM MG132 plus 10 µM RO-3306 or 25 µM MG132 only. Cells were fixed 40 min after the second treatment. Maximum projected images were shown.

(B) Quantification of (A). MG132 only, 442 cells; MG132 + RO-3306, 374 cells.

(C) Time-lapse images of cells expressing ER-FlucDM-eGFP arrested at mitosis by MG132 and further treated with RO-3306 to induce mitotic exit. Cells were treated with 25 µM MG132 for 55 min, and further treated with 25 µM MG132 plus 10 µM RO-3306 or 25 µM MG132 before imaging. Maximum projected images were shown.

(D) Quantification of (C). Remaining aggregates at 40 min were calculated for their aggregates count or area and compared to that of in 0 min. MG132 only, 125 cells; MG132 + RO-3306, 134 cells.

(E) Images of cells expressing ER-FlucDM-eGFP treated with DMSO or APCi (20 µM proTAME and 300 µM Apcin) and fixed at 55 min after released from the G2/M boundary. Maximum projected

images were shown.

(F) Quantifications of cells expressing ER-FlucDM-eGFP treated with DMSO or APCi (20 µM
proTAME and 300 µM Apcin). DMSO, 450 cells; APCi, 549 cells.

(G) Images of cells at telophase or cells of newly divided upon treatment of DMSO or APCi for 35 to
40 min. Maximum projected images were shown.

(H) Cells at telophase or newly divided cells that have aggregates upon treatment of DMSO or APCi
were counted.

(I) Aggregates number and area were counted in cells at telophase or in newly divided cells after
treated with DMSO or APCi. Only cells with aggregates were included in the quantification.
DMSO, 297 cells; APCi, 319 cells.

Scale bar, 10 µm. Data are from at least three independent experiments and analysed by unpaired
two side Mann-Whitney U-test (B, D, F and I) or paired t-test (H).

**Figure 7. Perturbation of proteins that regulate ER reorganization in aggregate clearance**

(A) Images of cells expressing ER-FlucDM-eGFP after depletion of Atlastin 2 and Atlastin 3 and fixed
at 65 min after released from the G2/M boundary. Maximum projected images were shown.

(B) Quantification of cells expressing ER-FlucDM-eGFP after depletion of Atlastin 2 and Atlastin 3
and fixed at 65 min after released from the G2/M boundary. Control, 396 cells; ATL2/ATL3 KD,
362 cells.

(C) Images of cells expressing ER-FlucDM-eGFP treated with DMSO or 100 nM MLN-8237 and fixed
at 65 min after released from the G2/M boundary. Maximum projected images were shown.

(D) Quantification of cells expressing ER-FlucDM-eGFP treated with DMSO or 100 nM MLN-8237

and fixed at 65 min after released from the G2/M boundary. Only round mitotic cells were counted. DMSO, 102 cells; MLN-8237, 288 cells.

(E) Images of cells expressing ER-FlucDM-eGFP treated with DMSO or 331nM nocodazole and fixed at 65 min after released from the G2/M boundary. Maximum projected images were shown.

(F) Quantification of cells expressing ER-FlucDM-eGFP treated with DMSO or 331nM nocodazole and fixed at 65 min after released from the G2/M boundary. Only round mitotic cells were counted. DMSO, 122 cells; nocodazole, 320 cells.

Scale bar, 10 µm. Data are from three independent experiments and analysed by unpaired two side Mann-Whitney U-test.

**Supplementary figures**

**Figure S1. ER-FlucDM-eGFP forms protein aggregates, related to Figure 1**

(A) Microscopy images of cells expressing NLS-FlucDM-eGFP. Two cell lines were used to express NLS-FlucDM-eGFP. Single Z-slice images were shown.

(B) Microscopy images of cells expressing NLS-FlucDM-eGFP and ER-FlucDM-mCherry in MCF10A cells. Single Z-slice images were shown.

(C) Protein blots for ER-FlucDM-eGFP and NLS-FlucDM-eGFP using anti-GFP or anti-β-actin antibodies.

(D) Relative GFP protein intensity in S1(C) was quantitated and plotted. The intensity of β-actin was used in normalization.

(E) Protein blotting to examine solubility of ER-FlucDM-eGFP under different concentrations of detergents, NP-40. Cells expressing ER-FlucDM-eGFP were subjected to 45 °C for 1 hour before

harvesting cell lysates in heat shock experiments. Insoluble fractions were quantitated and

plotted.

(F) Colocalization of ER-FlucDM-eGFP and Thioflavin T (ThT), a marker for protein aggregates.

Single Z-slice images were shown.

(G) Quantification of ER-FlucDM-eGFP aggregate number (left) and total aggregate area (right) of

individual cells in different cell lines. MCF10A, 516 cells; A549, 395 cells; MDA-MB-231, 369

cells; U2OS, 528 cells.

(H) Schematic representation of different constructs used in this study.

(I) Quantification of aggregate number in cells expressing ER-eGFP, ER-HaloWT-eGFP, ER-

HaloDM-eGFP. ER-eGFP, 612 cells; ER-HaloWT-eGFP, 405 cells; 546 ER-HaloDM-eGFP, 546

cells; ER-FlucDM-eGFP, 516 cells.

Data are from three independent experiments, analysed by unpaired t-test and represented by mean

+ SD. Scale bar, 10 μm.

**Figure S2. FRAP analysis of protein aggregates, related to Figure 2.**

(A) Recovery curve of ER-HaloDM-eGFP before and after photobleaching. Seven interphase cells

from three independent experiments. Data represent mean + SD.

(B) Recovery curve of individual cell data in ER-FlucDM-eGFP before and after photobleaching. 14

ROIs from the ER of interphase cell, 14 aggregates from cells at mitosis, 26 aggregates from

cells at interphase. Data were collected from four independent experiments.

(C) Recovery curve of ER-eGFP before and after photobleaching. Seven interphase cells from three

independent experiments. Data represent mean + SD.

(D) Quantification of protein fluorescence intensity of different constructs right after photobleaching and 5 min after recovery. 0 min is the protein intensity right after photobleaching and 5 min is the protein intensity 5 min after recovery from photobleaching. At least three independent experiments. Data represent mean $\pm$ SD.

**Figure S3. Aggregates in dividing cells, related to Figure 3.**

(A) Images to show ER-FlucDM-eGFP aggregates of dividing cells were surrounded by the membrane stained by the ER-tracker. Scale bar, 10 μm. Single Z-slice images were shown.

(B) TEM images of sectioned MCF10A mitotic cells expressing ER-FlucDM-eGFP.

(C) Time-lapse images of cells expressing ER-HaloDM-eGFP released from the G2/M boundary. NEB (nuclear envelop breakdown) is set to be t = 0 min. Maximum projected images were shown. Scale bar, 10 μm.

(D) Linear regression analysis to show that cell division process is not linearly dependent on aggregates quantity. 74 dividing cells are from four independent experiments.

**Figure S4. Effect of Thapsigargin or Tunicamycin on aggregates clearance, related to Figure 4.**

(A) Proteomic analysis of cells expressing ER-FlucDM-eGFP. Three independent experiments. Proteins involved in ER stress and unfolded protein response (blue) or cell cycle and cell division (pink) were highlighted.

(B) Changes of aggregate number and total area as a function of time after cells expressing ER-FlucDM-eGFP were treated with DMSO or 1 μM Thaps.

(C) Quantification of aggregate number and total area of cells expressing ER-FlucDM-eGFP treated

with DMSO or Tunicamycin (Tuni) during mitosis. DMSO, 542 cells; 200 ng/mL Tuni, 385 cells;

2 µg/mL Tuni, 471 cells; 10 µg/mL Tuni, 439 cells. Four independent experiments. Unpaired two

side Mann-Whitney U-test. Each solid thick line in violin plot indicates median of data in the

column. Gray lines are quartiles.

**Figure S5. Colocalization of BiP and ER-FlucDM-eGFP, related to Figure 5.**

(A) Immunofluorescence staining of BiP and ER-FlucDM-eGFP in dividing cells. Maximum projected

images were shown.

(B) Colocalization analysis of BiP and ER-FlucDM-eGFP. R represents Pearson correlation

coefficient (r).

(C) Immunofluorescence staining of BiP and ER-FlucDM-eGFP in interphase cells. Single Z-slice

images were shown.

Scale bar, 10 µm. Three independent experiments.

**Figure S6. MG132, APCi and RO-3306 treatments on aggregate clearance, related to Figure 6.**

(A) Quantification of the aggregate number in cells expressing ER-FlucDM-eGFP treated with DMSO

or 25 µM MG132 and fixed at 55 min after released from the G2/M boundary. DMSO, 293 cells;

MG132, 350 cells. Maximum projected images were shown.

(B) Quantification of the aggregate number in cells expressing ER-FlucDM-eGFP treated with DMSO

or 5 µM MG132 and fixed at 55 min since release. DMSO, 295 cells; MG132, 436 cells. Maximum

projected images were shown.

(C) Images of cells at telophase or newly divided cells after treated with DMSO or 25 µM MG132 for 35 min. Maximum projected images were shown.

(D) Cells at telophase or newly divided cells that have aggregates upon treatment of 25 µM MG132 were counted. DMSO, 455 cells; MG132, 393 cells.

(E) Aggregates number and area were counted in cells at telophase or in newly divided cells after treated with DMSO or 25 µM MG132. Only cells with aggregates were included in the quantification. DMSO, 294 cells; MG132, 339 cells.

(F) Images of cells at telophase and newly divided cells after treated with DMSO or 5 µM MG132 for 35 min. Maximum projected images were shown.

(G) Cells at telophase or newly divided cells that have aggregates upon treatment of DMSO or 5 µM MG132. DMSO, 676 cells; APCi, 646 cells.

(H) Aggregates number and area were counted in cells at telophase or in newly divided cells after treated with DMSO or 5 µM MG132. Only cells with aggregates were included in the quantification. DMSO, 227 cells; MG132, 297 cells.

(I) Quantification of the aggregate number in cells expressing ER-FlucDM-eGFP treated with 5 µM MG132 or 5 µM MG132 + RO-3306.

(J) Images of cells co-expressing ER-FlucDM-mCherry and mGreenLantern-hGeminin (degron). Left panel: cells were treated with DMSO or APC/Ci after 65 min released from the G2/M. Right panel: after 65 min released from the G2/M, cells were treated with DMSO or APC/Ci for 40 minutes using live-cell imaging. Maximum projected images were shown.

(K) Quantification of mGreenLantern-hGeminin (degron) intensity of cells treated with DMSO or APC/Ci for 40 min.

Scale bar, 10 µm. Data are from at least three independent experiments and analysed by unpaired

two side Mann-Whitney U-test (A, B, E, H and I), unpaired t-test (K) and paired t-test (D and G).

**Figure S7. Cycloheximide chase experiment and microscopy quantification of cells**

**expressing ER-FlucDM-eGFP, related to Figure 6.**

(A) Protein blots of ER-FlucDM-eGFP, Cyclin B1, p53 and $\beta$-actin under DMSO or 200 $\mu$M CHX

treatments. Cells were synchronized to the G2/M boundary and released into medium containing

DMSO or CHX and lysates were harvested.

(B) Protein levels of p53 and Cyclin B1 in different time points after DMSO or CHX treatments were

measured and plotted. Levels of $\beta$-actin were used in normalization.

(C) Protein levels of ER-FlucDM-eGFP in different time points after DMSO or CHX treatments were

measured and plotted. Levels of $\beta$-actin were used in normalization.

(D) Relative intensity of ER-FlucDM-eGFP in early and late dividing cells.

Scale bar, 10 µm. Data are from three independent experiments and analysed by unpaired t-test.

**References**

Aguilaniu, H., L. Gustafsson, M. Rigoulet, and T. Nystrom. 2003. Asymmetric inheritance of oxidatively damaged
proteins during cytokinesis. *Science*. 299:1751-1753.
Ando, R., S. Shimozono, H. Ago, M. Takagi, M. Sugiyama, H. Kurokawa, M. Hirano, Y. Niino, G. Ueno, F. Ishidate, T.
Fujiwara, Y. Okada, M. Yamamoto, and A. Miyawaki. 2024. StayGold variants for molecular fusion and
membrane-targeting applications. *Nat Methods*. 21:648-656.
Balch, W.E., R.I. Morimoto, A. Dillin, and J.W. Kelly. 2008. Adapting proteostasis for disease intervention. *Science*.
319:916-919.
Becher, I., A. Andres-Pons, N. Romanov, F. Stein, M. Schramm, F. Baudin, D. Helm, N. Kurzawa, A. Mateus, M.T.
Mackmull, A. Typas, C.W. Muller, P. Bork, M. Beck, and M.M. Savitski. 2018. Pervasive Protein Thermal
Stability Variation during the Cell Cycle. *Cell*. 173:1495-1507 e1418.
Bergman, Z.J., J.D. McLaurin, A.S. Eritano, B.M. Johnson, A.Q. Sims, and B. Riggs. 2015. Spatial reorganization of
the endoplasmic reticulum during mitosis relies on mitotic kinase cyclin A in the early Drosophila embryo.
*PLoS One*. 10:e0117859.

Clay, L., F. Caudron, A. Denoth-Lippuner, B. Boettcher, S. Buvelot Frei, E.L. Snapp, and Y. Barral. 2014. A
867    sphingolipid-dependent diffusion barrier confines ER stress to the yeast mother cell. *Elife*. 3:e01883.

868 Dickens, J.A., A. Ordonez, J.E. Chambers, A.J. Beckett, V. Patel, E. Malzer, C.S. Dominicus, J. Bradley, A.A. Peden, I.A.
869    Prior, D.A. Lomas, and S.J. Marciniak. 2016. The endoplasmic reticulum remains functionally connected by
870    vesicular transport after its fragmentation in cells expressing Z-alpha1-antitrypsin. *FASEB J*. 30:4083-4097.

871 Fuentealba, L.C., E. Eivers, D. Geissert, V. Taelman, and E.M. De Robertis. 2008. Asymmetric mitosis: Unequal
872    segregation of proteins destined for degradation. *Proc Natl Acad Sci U S A*. 105:7732-7737.

873 Ghislain, M., A. Udvardy, and C. Mann. 1993. S. cerevisiae 26S protease mutants arrest cell division in
874    G2/metaphase. *Nature*. 366:358-362.

875 Glotzer, M., A.W. Murray, and M.W. Kirschner. 1991. Cyclin is degraded by the ubiquitin pathway. *Nature*. 349:132-
876    138.

877 Gupta, R., P. Kasturi, A. Bracher, C. Loew, M. Zheng, A. Villella, D. Garza, F.U. Hartl, and S. Raychaudhuri. 2011.
878    Firefly luciferase mutants as sensors of proteome stress. *Nat Methods*. 8:879-884.

879 Hartl, F.U., A. Bracher, and M. Hayer-Hartl. 2011. Molecular chaperones in protein folding and proteostasis. *Nature*.
880    475:324-332.

881 Hill, S.M., S. Hanzen, and T. Nystrom. 2017. Restricted access: spatial sequestration of damaged proteins during
882    stress and aging. *EMBO Rep*. 18:377-391.

883 Hipp, M.S., P. Kasturi, and F.U. Hartl. 2019. The proteostasis network and its decline in ageing. *Nat Rev Mol Cell*
884    *Biol*. 20:421-435.

885 Jayaraj, G.G., M.S. Hipp, and F.U. Hartl. 2020. Functional Modules of the Proteostasis Network. *Cold Spring Harb*
886    *Perspect Biol*. 12.

887 Kaganovich, D., R. Kopito, and J. Frydman. 2008. Misfolded proteins partition between two distinct quality control
888    compartments. *Nature*. 454:1088-1095.

889 Kakihana, A., Y. Oto, Y. Saito, and Y. Nakayama. 2019. Heat shock-induced mitotic arrest requires heat shock
890    protein 105 for the activation of spindle assembly checkpoint. *FASEB J*. 33:3936-3953.

891 Labbadia, J., and R.I. Morimoto. 2015. The biology of proteostasis in aging and disease. *Annu Rev Biochem*.
892    84:435-464.

893 Lee, C., M. Ferguson, and L.B. Chen. 1989. Construction of the endoplasmic reticulum. *J Cell Biol*. 109:2045-2055.

894 Liu, Z., G. Liu, D.P. Ha, J. Wang, M. Xiong, and A.S. Lee. 2023. ER chaperone GRP78/BiP translocates to the nucleus
895    under stress and acts as a transcriptional regulator. *Proc Natl Acad Sci U S A*. 120:e2303448120.

896 Melo, E.P., T. Konno, I. Farace, M.A. Awadelkareem, L.R. Skov, F. Teodoro, T.P. Sancho, A.W. Paton, J.C. Paton, M.
897    Fares, P.M.R. Paulo, X. Zhang, and E. Avezov. 2022. Stress-induced protein disaggregation in the
898    endoplasmic reticulum catalysed by BiP. *Nat Commun*. 13:2501.

899 Miyata, T., D. Hagiwara, Y. Hodai, T. Miwata, Y. Kawaguchi, J. Kurimoto, H. Ozaki, K. Mitsumoto, H. Takagi, H. Suga,
900    T. Kobayashi, M. Sugiyama, T. Onoue, Y. Ito, S. Iwama, R. Banno, M. Matsumoto, N. Kawakami, N. Ohno,
901    H. Sakamoto, and H. Arima. 2020. Degradation of Mutant Protein Aggregates within the Endoplasmic
902    Reticulum of Vasopressin Neurons. *iScience*. 23:101648.

903 Mochida, K., and H. Nakatogawa. 2022. ER-phagy: selective autophagy of the endoplasmic reticulum. *EMBO Rep*.
904    23:e55192.

905 Morris, J.A., A.J. Dorner, C.A. Edwards, L.M. Hendershot, and R.J. Kaufman. 1997. Immunoglobulin binding protein
906    (BiP) function is required to protect cells from endoplasmic reticulum stress but is not required for the
907    secretion of selective proteins. *J Biol Chem*. 272:4327-4334.

908 Nillegoda, N.B., J. Kirstein, A. Szlachcic, M. Berynskyy, A. Stank, F. Stengel, K. Arnsburg, X. Gao, A. Scior, R. Aebersold,
909    D.L. Guilbride, R.C. Wade, R.I. Morimoto, M.P. Mayer, and B. Bukau. 2015. Crucial HSP70 co-chaperone
910    complex unlocks metazoan protein disaggregation. *Nature*. 524:247-251.

Olzmann, J.A., R.R. Kopito, and J.C. Christianson. 2013. The mammalian endoplasmic reticulum-associated
degradation system. *Cold Spring Harb Perspect Biol*. 5.

Olzscha, H., S.M. Schermann, A.C. Woerner, S. Pinkert, M.H. Hecht, G.G. Tartaglia, M. Vendruscolo, M. Hayer-Hartl,
F.U. Hartl, and R.M. Vabulas. 2011. Amyloid-like aggregates sequester numerous metastable proteins
with essential cellular functions. *Cell*. 144:67-78.

Palozola, K.C., G. Donahue, H. Liu, G.R. Grant, J.S. Becker, A. Cote, H. Yu, A. Raj, and K.S. Zaret. 2017. Mitotic
transcription and waves of gene reactivation during mitotic exit. *Science*. 358:119-122.

Parashar, S., R. Chidambaram, S. Chen, C.R. Liem, E. Griffis, G.G. Lambert, N.C. Shaner, M. Wortham, J.C. Hay, and
S. Ferro-Novick. 2021. Endoplasmic reticulum tubules limit the size of misfolded protein condensates.
*Elife*. 10.

Pina, F.J., and M. Niwa. 2015. The ER Stress Surveillance (ERSU) pathway regulates daughter cell ER protein
aggregate inheritance. *Elife*. 4.

Preissler, S., and D. Ron. 2019. Early Events in the Endoplasmic Reticulum Unfolded Protein Response. *Cold Spring
Harb Perspect Biol*. 11.

Rosenzweig, R., N.B. Nillegoda, M.P. Mayer, and B. Bukau. 2019. The Hsp70 chaperone network. *Nat Rev Mol Cell
Biol*. 20:665-680.

Rousseau, E., B. Dehay, L. Ben-Haiem, Y. Trottier, M. Morange, and A. Bertolotti. 2004. Targeting expression of
expanded polyglutamine proteins to the endoplasmic reticulum or mitochondria prevents their
aggregation. *Proc Natl Acad Sci U S A*. 101:9648-9653.

Rujano, M.A., F. Bosveld, F.A. Salomons, F. Dijk, M.A. van Waarde, J.J. van der Want, R.A. de Vos, E.R. Brunt, O.C.
Sibon, and H.H. Kampinga. 2006. Polarised asymmetric inheritance of accumulated protein damage in
higher eukaryotes. *PLoS Biol*. 4:e417.

Sackton, K.L., N. Dimova, X. Zeng, W. Tian, M. Zhang, T.B. Sackton, J. Meaders, K.L. Pfaff, F. Sigoillot, H. Yu, X. Luo,
and R.W. King. 2014. Synergistic blockade of mitotic exit by two chemical inhibitors of the APC/C. *Nature*.
514:646-649.

Samanta, S., S. Yang, B. Debnath, D. Xue, Y. Kuang, K. Ramkumar, A.S. Lee, M. Ljungman, and N. Neamati. 2021.
The Hydroxyquinoline Analogue YUM70 Inhibits GRP78 to Induce ER Stress-Mediated Apoptosis in
Pancreatic Cancer. *Cancer Res*. 81:1883-1895.

Schlaitz, A.L., J. Thompson, C.C. Wong, J.R. Yates, 3rd, and R. Heald. 2013. REEP3/4 ensure endoplasmic reticulum
clearance from metaphase chromatin and proper nuclear envelope architecture. *Dev Cell*. 26:315-323.

Sharma, R., M.M.D. Pramanik, B. Chandramouli, N. Rastogi, and N. Kumar. 2018. Understanding organellar protein
folding capacities and assessing their pharmacological modulation by small molecules. *Eur J Cell Biol*.
97:114-125.

Shim, S.M., H.R. Choi, K.W. Sung, Y.J. Lee, S.T. Kim, D. Kim, S.R. Mun, J. Hwang, H. Cha-Molstad, A. Ciechanover,
B.Y. Kim, and Y.T. Kwon. 2018. The endoplasmic reticulum-residing chaperone BiP is short-lived and
metabolized through N-terminal arginylation. *Sci Signal*. 11.

Singhvi, A., and G. Garriga. 2009. Asymmetric divisions, aggresomes and apoptosis. *Trends Cell Biol*. 19:1-7.

Tyedmers, J., A. Mogk, and B. Bukau. 2010. Cellular strategies for controlling protein aggregation. *Nat Rev Mol
Cell Biol*. 11:777-788.

Vincenz-Donnelly, L., H. Holthusen, R. Korner, E.C. Hansen, J. Presto, J. Johansson, R. Sawarkar, F.U. Hartl, and M.S.
Hipp. 2018. High capacity of the endoplasmic reticulum to prevent secretion and aggregation of
amyloidogenic proteins. *EMBO J*. 37:337-350.

Wang, C., J. Ding, Q. Wei, S. Du, X. Gong, and T.G. Chew. 2023. Mechanosensitive accumulation of non-muscle
myosin IIB during mitosis requires its translocation activity. *iScience*. 26:107773.

Watson, E.R., N.G. Brown, J.M. Peters, H. Stark, and B.A. Schulman. 2019. Posing the APC/C E3 Ubiquitin Ligase to

Orchestrate Cell Division. *Trends Cell Biol*. 29:117-134.

Wirth, A.J., M. Platkov, and M. Gruebele. 2013. Temporal variation of a protein folding energy landscape in the
cell. *J Am Chem Soc*. 135:19215-19221.

Wiseman, R.L., J.S. Mesgarzadeh, and L.M. Hendershot. 2022. Reshaping endoplasmic reticulum quality control
through the unfolded protein response. *Mol Cell*. 82:1477-1491.

Woerner, A.C., F. Frottin, D. Hornburg, L.R. Feng, F. Meissner, M. Patra, J. Tatzelt, M. Mann, K.F. Winklhofer, F.U.
Hartl, and M.S. Hipp. 2016. Cytoplasmic protein aggregates interfere with nucleocytoplasmic transport of
protein and RNA. *Science*. 351:173-176.

Zhang, W., Z. Zhang, Y. Xiang, D.D. Gu, J. Chen, Y. Chen, S. Zhai, Y. Liu, T. Jiang, C. Liu, B. He, M. Yan, Z. Wang, J.
Xu, Y.L. Cao, B. Deng, D. Zeng, J. Lei, J. Zhuo, X. Lei, Z. Long, B. Jin, T. Chen, D. Li, Y. Shen, J. Hu, S. Gao,
and Q. Liu. 2024. Aurora kinase A-mediated phosphorylation triggers structural alteration of Rab1A to
enhance ER complexity during mitosis. *Nat Struct Mol Biol*. 31:219-231.

Zhou, C., B.D. Slaughter, J.R. Unruh, F. Guo, Z. Yu, K. Mickey, A. Narkar, R.T. Ross, M. McClain, and R. Li. 2014.
Organelle-based aggregation and retention of damaged proteins in asymmetrically dividing cells. *Cell*.
159:530-542.

# Figure 1

**A**

MCF10A | A549
ER-FlucDM-eGFP
MDA-MB-231 | U2OS

**B**

ER-FlucDM-mCherry | mStayGold-Sec61β | Merge

**C**

Luciferase activity (%)

a: MG132+FlucDM-eGFP
b: FlucDM-eGFP
c: ER-FlucDM-eGFP
d: ER-FlucWT-eGFP

$p < 0.015$   $p < 0.005$

a   b   c   d

**D**

ER-FlucDM-mCherry | ER$^{FSHR}$-FlucDM-eGFP | ER-HaloDM-eGFP | ER-eGFP

**E**

ER-FlucDM-eGFP | ER-Tracker | Merge

Intensity (normalized to max)

Distance (μm)

**F**

10 μm   1 μm   200 nm

# Figure 2

**A** Interphase

ER-FlucDM-eGFP          ER-FlucDM-mCherry

**B** Mitosis

ER-FlucDM-eGFP          ER-FlucDM-mCherry

# Figure 3

# Figure 4

# Figure 5

# Figure 6

# Figure 7

# Figure S1

**A** Only NLS-FlucDM-eGFP

MCF10A | U2OS

**B** NLS-FlucDM-eGFP + ER-FlucDM-mCherry

NLS-FlucDM-eGFP | ER-FlucDM-mCherry | Merge

**C**

GFP
β-actin
Lanes  1  2  3  4  5  6

A: ER-FlucDM-eGFP  B: NLS-FlucDM-eGFP + ER-FlucDM-mCherry

**D**

$p < 0.001$
Relative GFP intensity
A  B

**E**

0.2% NP-40 | 0.5% NP-40 | 1.0% NP-40
S  P  S  P  S  P  S  P  S  P  S  P
GFP
β-actin
Heat shock  −  −  +  +  −  −  +  +  −  −  +  +
Lanes  1  2  3  4  5  6  7  8  9  10  11  12

Insoluable fraction (GFP signal of insoluable/soluable)
● 0.2% NP-40
■ 0.5% NP-40
▲ 1.0% NP-40
Supernatant  Pellet

**F**

ER-FlucDM-eGFP | Thioflavin T (ThT) | Merge

**G**

Count / cell
MCF10A  A549  MDA-MB-231  U2OS

Area / cell (μm²)
MCF10A  A549  MDA-MB-231  U2OS

**H**

| Construct | N-terminal | Core | Fluorescent | C-terminal |
|---|---|---|---|---|
| ER-FlucDM-eGFP | N-terminal ER targeting sequence MGWSCIILFFVATAT | FlucDM (R188Q, R261Q) | eGFP | KDEL |
| ER-FlucDM-mCherry | N-terminal ER targeting sequence MGWSCIILFFVATAT | FlucDM (R188Q, R261Q) | mCherry | KDEL |
| ER$^{FSHR}$-FlucDM-eGFP | N-terminal sequence from FSHR MALLLVSLLAFLSLGSG | FlucDM (R188Q, R261Q) | eGFP | |
| ER-FlucWT-eGFP | N-terminal ER targeting sequence MGWSCIILFFVATAT | FlucWT | eGFP | KDEL |
| FlucDM-eGFP | | FlucDM (R188Q, R261Q) | eGFP | KDEL |
| ER-HaloWT-eGFP | N-terminal ER targeting sequence MGWSCIILFFVATAT | HaloWT | eGFP | KDEL |
| ER-HaloDM-eGFP | N-terminal ER targeting sequence MGWSCIILFFVATAT | HaloDM (M21K, F86L) | eGFP | KDEL |
| ER-eGFP | N-terminal ER targeting sequence MGWSCIILFFVATAT | eGFP | | KDEL |
| mGreenLantern-hGeminin (degron) | | mGreenLantern | hGeminin (degron) | |
| NLS-FlucDM-eGFP | nuclear targeting sequence | FlucDM (R188Q, R261Q) | eGFP | |
| mStayGold-Sec61β | mStayGold | Sec61β | | |

**I**

Count / cell
0.52 ± 0.58    0.39 ± 0.33    2.93 ± 1.57    9.73 ± 1.64
ER-eGFP  ER-HaloWT-eGFP  ER-HaloDM-eGFP  ER-FlucDM-eGFP

**Figure S2**

# Figure S3

**A**

ER-FlucDM-eGFP | ER-Tracker | Merge

Cell 1

Cell 2

Cell 3

**B**

10 μm

1 μm

200 nm

**C** ER-HaloDM-eGFP

| | Before NEB | NEB (t=0 min) | 30 min | 60 min | 90 min | 120 min |

Mitosis

Interphase

**D**

R² < 0.005

Aggregates count before NEB

NEB to ingression end (min)

# Figure S4

# Figure S5

**A**

ER-FlucDM-eGFP  BiP  Merge

**B**

Single layer
of Z-stack

Pearson's R= 0.933

BiP fluorescence intensity

ER-FlucDM-eGFP fluorescence intensity

**C**

ER-FlucDM-eGFP  BiP  Merge