Title page (Original Research Article)

Characterisation of RNA editing and gene therapy with a compact CRISPR-Cas13 in the retina

Satheesh Kumar[1,2,3], Yi-Wen Hsiao[1], Vickie H Y Wong[4], Deborah Aubin[5,6], Jiang-Hui Wang[1],

Leszek Lisowski[7,8,9], Elizabeth P Rakoczy[10], Fan Li[11], Luis Alarcon-Martinez[1,2], Anai

Gonzalez-Cordero[5,6], Bang V Bui[4], Guei-Sheung Liu[1,2,3*]

[1]Centre for Eye Research Australia, Royal Victorian Eye and Ear Hospital, East Melbourne,
VIC, Australia

[2]Ophthalmology, Department of Surgery, University of Melbourne, East Melbourne, VIC,
Australia

[3]Menzies Institute for Medical Research, University of Tasmania, Hobart, TAS, Australia

[4]Department of Optometry and Vision Sciences, University of Melbourne, Parkville, VIC,
Australia

[5]School of Medical Sciences, Faculty of Medicine and Health, The University of Sydney,
Camperdown, NSW, Australia

[6]Stem Cell Medicine Group, Children's Medical Research Institute, Westmead, NSW,
Australia

[7]Translational Vectorology Research Unit, Children's Medical Research Institute, Faculty of
Medicine and Health, The University of Sydney, Westmead, NSW, Australia

[8]Military Institute of Medicine, Laboratory of Molecular Oncology and Innovative Therapies,
Warsaw, Poland

[9]Australian Genome Therapeutics Centre, Children's Medical Research Institute and Sydney
Children's Hospitals Network, Westmead, NSW, Australia

[10]Centre for Ophthalmology and Visual Sciences, University of Western Australia, Perth, WA,
Australia

11Eye Center, Zhongshan City People's Hospital, Zhongshan, Guangdong Province, China

*Correspondence and requests for materials should be addressed to

Dr Guei-Sheung Liu (rickliu0817@gmail.com). Centre for Eye Research Australia. Address: Level 7, 32 Gisborne Street, East Melbourne, VIC 3002, Australia. Tel: +61399298360.

Key words: RNA silencing, CRISPR-Cas13, angiogenesis, ocular neovascularization, diabetic retinopathy, gene therapy, retinal organoids

ABSTRACT

CRISPR-Cas gene editing technology has revolutionised the development of novel genetic

therapies. CRISPR-Cas13 nucleases are programmable RNA-targeting effectors that can

silence gene expression in a reversible manner. Recent iterations of Cas13 nucleases are

compact and adequate for adeno-associated virus (AAV) delivery to achieve strong and

persistent transduction of various organs in a safe manner. Here, we report significant

transcriptomic signatures of Cas13bt3 expression and demonstrate that all-in-one AAV gene

therapy with Cas13bt3 can effectively silence *VEGFA* in human retinal organoids and

humanised mouse models. Specifically, human embryonic stem cells (hESC)-derived retinal

pigment epithelium cells show high expression of Cas13bt3 after virus delivery corresponding

to significant reduction of *VEGFA* mRNA. We further show that intravitreal delivery of

Cas13bt3 can transduce mouse retinal cells efficiently, reaching the photoreceptor cells for

specific knockdown of human *VEGFA*, without impacting mouse *VEGFA* in the trVEGF029

(Kimba) mouse. Our results reveal important considerations for assessing Cas13 activity and

establish Cas13bt3 as a potential anti-VEGF agent that can achieve long-term control of VEGF

for treatment of diabetic retinopathy and wet age-related macular degeneration.

INTRODUCTION

The emergence of Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) and CRISPR-associated (Cas) enzymes, known as CRISPR-Cas, has fuelled several gene editing efforts for novel therapeutics in the past decade[1]. Developed by repurposing the bacterial immune defence against phage viruses, CRISPR-Cas harnesses the ability to induce targeted changes to DNA or RNA sequences. Among CRISPR-Cas proteins, CRISPR-Cas13 is of particular interest for its exclusive specificity for RNA. Targeting RNA is especially attractive as it enables control of gene expression without perturbing the genome and introducing permanent changes and is therefore an emerging avenue for therapeutic development[2].

Eleven variants of CRISPR-Cas13 have been described to date[3]. Among them, Cas13d (commonly referred to as CasRx) described in 2018 garnered special attention for its compact size of 930aa, making it the first Cas13 enzyme amenable to adeno-associated virus (AAV) delivery. Studies investigating the development of RNA-targeting therapeutics have widely used CasRx to demonstrate preclinical efficacy against disorders affecting vision, hearing and brain function[4, 5, 6, 7], however significant unintended editing have been reported, hampering its clinical progress[8, 9], and the exact specificity profile of CasRx remains contested[10]. More recently, hypercompact Cas13bt3 (also reported as Cas13X) enzymes were described, ranging even smaller at 775aa[11, 12]. The newfound enzyme has been reported to possess high specificity, and its small size enables additional applications such as RNA base editing using a single AAV[13, 14, 15]. The therapeutic potential of Cas13bt3 is yet to be demonstrated, however initial *in vitro* studies show promising efficiency and safety[13, 16].

As a leading cause of blindness in the elderly population affecting 200 million people globally, ocular neovascularisation refers to a group of conditions characterised by the abnormal growth of leaky blood vessels in the eye. Vascular endothelial growth factor (VEGF)

has been described as a key mediator of this process, and the overexpression of *VEGFA*

underlies pathological angiogenesis and neovascularisation in common ocular diseases like

diabetic retinopathy (DR) and neovascular age-related macular degeneration (nAMD). Regular

(monthly/ bimonthly) administration of anti-VEGF drugs is the approved front-line treatment

modality for these patients[17]. Delivered through intravitreal injections, frequent administration

of anti-VEGF drugs poses a significant clinical burden on healthcare institutions and patients

due to risk of infection, patient anxiety, clinic capacity and availability of trained healthcare

personnel among others[18].

We have previously described a potential alternative therapy to current anti-VEGF

injections using an all-in-one AAV CRISPR-CasRx gene therapy for the treatment of ocular

neovascularisation. This approach was validated *in vitro* in HEK293FT and MIO-M1 cells,

demonstrating more than 90% and 50% *VEGFA* mRNA knockdown, respectively[19]. Here, we

utilize the novel Cas13bt3 to demonstrate potent *VEGFA* knockdown *in vitro* in human cell

lines, in 3D human embryonic stem cells (hESC)-derived retinal organoids and *in vivo* in

transgenic humanised mouse retinas. Promisingly, our results show significant knockdown of

VEGFA mRNA in human cell lines without impact on predicted off-target genes. Single cell

RNA sequencing (scRNA-seq) was used to characterise Cas13bt3-mediated *VEGFA* mRNA

knockdown at single cell resolution in human retinal organoids identifying, for the first time,

the overall transcriptomic impact and cell-specific activity of the enzyme. Lastly, *in vivo* gene

therapy showed robust *VEGFA* mRNA silencing in mouse models of proliferative diabetic

retinopathy along with control of neovascularisation symptoms. These findings support the

development of Cas13-mediated ocular gene therapy establishing the clinical potential of

Cas13bt3 and offering an alternative treatment to current anti-VEGFA treatments.

RESULTS

CRISPR-Cas13bt3 can knock down *VEGFA* mRNA in mammalian cells with minimal unintended targeting activity.

We used our previously described single guide RNAs (sgRNA) targeting Exon 4 of *VEGFA* [19], to develop our new Cas13bt3 constructs and assess RNA cutting efficiency as well as collateral and off target cleavages (**Figure 1A-D**). Collateral activity is the trans-cleavage by Cas13 enzymes of adjacent mRNA transcripts as the result of an activated nuclease domain (**Figure 1B**). Off-target activity, on the other hand, is the targeting of other non-target mRNA transcripts due to unspecific binding of the sgRNA (**Figure 1C**).

We sought to compare and validate efficiency of the new Cas13bt3 enzyme by testing RNA cutting against two previously established RNA-targeting methods, RNA interference (short hairpin RNA, shRNA) and CasRx (a variant of Cas13d). These are widely used for therapeutic and diagnostic applications. The final all-in-one AAV plasmids carrying either *VEGFA*-targeting shRNA or precursor single guide RNA (sgRNA hereafter) with either CasRx or Cas13bt3 (**Figure 1D**) were transfected into HEK293FT cells. Here, to further enhance the expression of Cas13bt3 we incorporated the stronger cytomegalovirus (CMV) promoter instead of the previously used elongation factor 1α short (EFS) promoter.

Significant *VEGFA* mRNA knockdown was observed from all three RNA-targeting methods, with up to 70% shRNA *VEGFA* mRNA knockdown ($P < 0.001$), while CasRx and Cas13bt3 achieved up to 90% knockdown ($P < 0.0001$) compared to vector controls (AAV plasmid carrying a fluorescent marker) (**Figure 1F**, n = 4 independent experiments). Next, we sought to determine if the reduction in mRNA would also lead to a decrease in protein levels. Interestingly, ELISAs for VEGFA revealed that only the Cas13bt3 treatment led to significant reduction of VEGFA protein in the cell supernatant (**Figure 1G**, $P < 0.05$; n = 4 independent

experiments). As both Cas13 variants proved potent in silencing *VEGFA* mRNA in HEK293FT cells, we next transfected immortalised human retinal pigment epithelium cells (ARPE19 cell line) with the *VEGFA*-targeting CasRx and Cas13bt3 plasmids and again observed up to 90% knockdown ($P < 0.05$, n = 3 independent experiments) with Cas13bt3 against vector controls (**Figure S1**), suggesting that Cas13bt3 is a promising candidate for RNA silencing of retinal cells.

To further understand transcriptome-wide effects, we performed bulk RNA sequencing (RNA-seq) on HEK293FT cells transfected with shRNA, CasRx and Cas13bt3. RNA-seq was also performed on cells treated with either control, Cas13bt3 only, Cas13bt3NT and Cas13bt3VEGFA constructs to identify collateral and off-target activity of Cas13. Differentially expressed genes (DEGs) were similar for all treated groups, shRNA, CasRx and Cas13bt3 (**Figure S2A-C**), indicating that collateral activity from the new Cas13bt3 is comparable to previously established RNA-targeting methods. Interestingly, the non-targeting (NT) sgRNA group exhibited the highest number of DEGs, a finding that corresponds with recent studies (**Figure 1G, H and S2A-D**) [16, 20]. Volcano plots of Cas13bt3 samples against control demonstrated and confirmed a significant downregulation of *VEGFA* mRNA only on cells that received the VEGFA targeting treatment (**Figure 1I).**

We identified potential off-target genes through NCBI BLAST of our VEGFA sgRNA sequence and studied the expression levels of the top 10 candidate off-target genes (**Figure S3A**). None of the off-target genes were significantly downregulated, indicating lack of off-target activity from Cas13bt3 (**Figure 1J**), as was the case for shRNA and CasRx (**Figure S2F**).

To further examine the varying DEG levels between the NT and targeting sgRNAs, we identified the top significantly enriched gene ontology (GO) terms for the Cas13bt3-NT

sgRNA and Cas13bt3-VEGFA sgRNA groups. Top 5 significant GO terms for the Cas13bt3-

NT sgRNA group were all positive enrichments and specifically showed upregulation of RNA

binding processes, such as "*pre-mRNA binding*" and "*mRNA 3' UTR binding*", possibly

explaining the elevated levels of DEGs in this group. On the other hand, with a VEGFA

sgRNA, the top 5 GO terms were all negative enrichments and particularly showed

downregulation of "*platelet-derived growth factor (PDGF) binding*" and "*PDGF receptor*

binding", both angiogenesis promoting factors (**Figure 1K**). A recent study showed CasRx to

have intrinsic targets that affected essential processes like RNA metabolism and mitotic cell

cycle [20] and opined that this could underlie the lethality seen upon CasRx expression in a mouse

model [21]. Here, we found laminin-related processes (e.g., "*laminin binding*", "*laminin*

complex") were strongly downregulated across all groups (Cas13bt3 only, Cas13bt3-NT

sgRNA and Cas13bt3-VEGFA sgRNA) suggesting intrinsic targets could exist for Cas13bt3

as well (**Extended data** 1).

While Cas13bt3 may undergo activation after maturation of a NT sgRNA; without

adequate homologous pairing, the catalytic pocket cannot be formed for efficient nuclease

activity [22]. On the other hand, with a targeting gRNA, Cas13bt3 would be activated to form the

Cas13bt3-gRNA-target complex for targeted knockdown. Our results suggest that with a NT

sgRNA, Cas13bt3 initiates an unspecific target-search process involving significant RNA

binding that greatly impacts the expression levels of several genes, while a targeting sgRNA

narrows the target-search process to reduce overall transcriptomic impact. Notably, despite

high levels of overall DEGs, the number of downregulated genes remained similar between the

NT sgRNA and VEGFA sgRNA groups suggesting that the NT sgRNA does not induce

formation of the catalytic pocket for RNA knockdown, and altered gene expression is

predominantly a result of increased RNA binding.

Figure 1. Comparison of RNA-targeting tools for in vitro testing of VEGFA mRNA silencing. (A) Schematic of CRISPR-Cas13 mechanism of action. **(B)** VEGFA sgRNA sequence and targeting region on VEGFA mRNA sequence. **(C)** Schematic of collateral and **(D)** off-target cleavage. **(E)** Schematic for domain structure of VEGFA targeting shRNA and Cas13 (CasRx and Cas13bt3) vectors. **(F)** VEGFA mRNA knockdown from shRNA, CasRx and Cas13bt3 in HEK293FT cells (n = 7). **(H)** VEGFA protein expression from ELISA after transfection of shRNA, CasRx and Cas13bt3 plasmids (n = 4). **(F)** Schematic for CRISPR-Cas13 collateral cleavage. **(G)** Venn diagram of total number of differentially expressed genes (DEGs) from Cas13bt3 only, Cas13bt3-NT sgRNA and Cas13bt3-VEGFA sgRNA plasmids. **(I)** Volcano plots showing DEGs from Cas13bt3 only, Cas13bt3-NT sgRNA and Cas13bt3-VEGFA sgRNA compared to transfection control in HEK293FT cells. VEGFA gene is denoted by a black arrow. **(J)** TPM plot for VEGFA and candidate off-target genes. **(K)** Top 5 enriched GO molecular function (MF) terms from Cas13bt3 only, Cas13bt3-NT sgRNA and Cas13bt3 VEGFA sgRNA. NES above zero represent significant activation of pathway, while NES below zero represents significant suppression of pathway. Orange text denotes key factors that are upregulated; blue text denotes key factors downregulated. Data presented as mean ± SEM. Statistical analysis was conducted by one-way ANOVA and Tukey's multiple comparison test **(F, G)** and Welch's t-test **(J)**. ns: not significant, $*P \leq 0.05$, $**P \leq 0.001$, $***P \leq 0.0001$, $****P \leq 0.00001$.

CRISPR-Cas13bt3 sgRNAs multiplexing for knockdown of target gene using a single-AAV vector.

The small size of Cas13bt3 and the significant downregulation of *VEGFA* mRNA observed prompted us to hypothesize that multiple sgRNAs can be delivered using a single AAV vector to increase overall targeting and RNA knockdown efficiency. In theory, multiple sgRNAs would increase rate of Cas13 activation and levels of intracellular Cas13-sgRNA complexes for RNA silencing.

To understand Cas13bt3 activation and demonstrate multiplex targeting, we transfected Cas13bt3-VEGFA sgRNA and Cas13bt3 with a VEGFA sgRNA array (carrying three VEGFA-targeting sgRNAs) (**Figure 2A-C**). Surprisingly, cells transfected with Cas13bt3-VEGFA sgRNA or Cas13bt3-VEGFA sgRNA array showed similar levels of significant

VEGFA mRNA knockdown (<90%) (**Figure 2D,** n = 3 independent experiments). We then

sought to study collateral and off-target cleavages from the VEGFA sgRNA array system using

RNA-seq. Overall DEGs for both single and multiple sgRNA groups when compared to NT

control were comparable and suggested minimal collateral activity from the sgRNA array

system (**Figure 2E**). Again, we identified top candidate off-target genes for the other *VEGFA*

sgRNAs (**Figure S4B and C**) and did not detect any significant off-target activity in these

genes by the Cas13bt3-VEGFA sgRNA array system as well (**Figure 2F**). Active Cas13

nucleases competing for target mRNA or low levels of *VEGFA* mRNA transcripts may have

resulted in the lack of improvement in targeting efficiency, however our results suggest that

multiple sgRNAs can be used simultaneously with Cas13bt3 while still avoiding significant

collateral activity.

Apart from competition for substrate mRNA, it was recently reported that the presence

of "UC" sites on the target mRNA may also affect knockdown efficiency, with "UC" sites

within 10-20 bases of spacer binding to allow for optimal cleavage (**Figure S3A**)[16]. We found

that while sgRNA1 and sgRNA2 had "UC" sites in these optimal regions, sgRNA3 had only

one distal "UC" site 40 nucleotides away from spacer binding (**Figure S3B**). While we did not

independently test the knockdown efficiency of our sgRNAs, this finding suggests sgRNA 3

alone would achieve low levels of RNA knockdown. As no improved knockdown was

observed from the VEGFA sgRNA array, we proceeded to investigate Cas13bt3 with a single

sgRNA further.

Figure 2 Guide-dependent and guide-independent off-targets from Cas13bt3-mediated VEGFA knockdown. **(A)** Schematic of single and multiplex targeting with Cas13bt3 and **(B)** U6-sgRNA domain in single sgRNA and multiple sgRNA vectors. **(C)** Guide RNA targeting regions in *VEGFA* mRNA. **(D)** *VEGFA* mRNA knockdown from Cas13bt3, non-targeting and targeting sgRNAs. **(E)** Volcano plots showing differentially expressed genes between Cas13bt3 with non-targeting and **(F)** TPM plots for VEGFA and candidate off-target genes. Target gene VEGFA is highlighted in red. Data presented as mean ± SEM. Statistical analysis

232   was conducted by one-way ANOVA and Tukey's multiple comparison test (**C**) and Welch's t-
233   test. ns: not significant, *P≤0.05, ***P≤0.0001, ****P≤0.00001.

Retinal organoids can be efficiently transduced by AAV2.7m8 for expression of Cas13bt3.

Human organoids are excellent models for gaining clinically relevant information on new

therapeutic agents, as they are derived from human pluripotent stem cells (PSCs) and reproduce

morphological, functional and transcriptomic features *in vitro*[23]. Recently, hESC-derived

retinal organoids have been used to investigate eye development and model inherited retinal

diseases as well as the testing of novel gene therapies[24, 25]. While CRISPR-Cas9 has entered

the clinic, Cas13-based therapies largely remain at the preclinical stage, and demonstrating

utility of Cas13 in retinal organoids will be a major advance for ocular gene therapy.

Here we utilised our well-established 2D/3D differentiation protocol to generate hESC-

derived retinal organoid containing all the major cell types in the retina[26]. First, to ensure

maximum expression of the Cas13 nuclease in the retinal cells, we subjected the Cas13bt3

protein sequence to codon optimisation to obtain two other variants of Cas13bt3 (Cas13bt3$_{COP1}$

and Cas13bt3$_{COP2}$) (**Figure S5A**). We compared the expression levels of the original Cas13bt3

with Cas13bt3$_{COP1}$ and Cas13bt3$_{COP2}$ by transfection in HEK293FT and ARPE19 cells and

found both variants to exhibit increased expression and knockdown efficiency (**Figure S5B, C**

and D). Among them, Cas13bt3$_{COP1}$ showed the highest expression and was used for

subsequent investigation. To elucidate the activity of Cas13bt3 and efficacy for AAV gene

therapy RNA editing in *bona fide* human retinal cells we transduced 19 weeks old organoids

with AAV2.7m8.U6.NTsgRNA.Cas13bt3 and AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 viral

vectors. Retinal organoids were then harvested for single cell RNA sequencing (scRNA-Seq)

6 weeks after treatment (~25 weeks) for (**Figure 3A**). First, we confirmed that the major retinal

cell types were clustered similarly across experimental conditions (S1: Untreated; S2:

AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3; and S3: AAV2.7m8.U6.NTsgRNA.Cas13bt3)

(**Figure 3B**, n = 3 pooled organoids per condition). We used our recently described retinal atlas

dataset to determine fidelity of retinal organoids and identified expression of the different

retinal cell markers[27], confirming exclusive cell-specific expression of the markers (**Figure 3C**

and S6, S7).

Next, to examine Cas13bt3 expression in the retinal organoids, we evaluated read

numbers and classified cells containing more than 10 reads of Cas13bt3 to be positive for viral

infection. Using this threshold, we determined an average transduction efficiency of 18.78%

(1403 out of 7467 cells) for the VEGFA sgRNA group and 25.95% (2260 out of 8706 cells)

for the NT sgRNA group. Next, we analysed transduction and presence of Cas13bt3 in specific

cell types. While all cell types showed noticeable levels of Cas13bt3, RPE cells, in particular,

showed high levels of expression with up to 50% of cells containing Cas13bt3 in the VEGFA

sgRNA (S2) treatment group. Interestingly, in the NT sgRNA group, only ~15% of RPE cells

expressed Cas13bt3, which could owe to varying culture conditions. Bipolar cells showed the

least expression of Cas13bt3 ($\leq$10%), while all other cell types showed moderate expression

($\leq$25%) (**Figure 3D**). Overall, our results indicated AAV2.7m8 as a useful delivery vector for

cells across retinal cell types.

To investigate the impact of the NT sgRNA in retinal cells, we first compared DEGs

between cells positive and negative for Cas13bt3 in the retinal organoid that received the

AAV2.7m8.U6.NTsgRNA.Cas13bt3. Here, we observed a significant number DEGs across all

cell types, and in similar trend to the HEK293FT cells where most DEGs were upregulated

genes (**Figure 3E**). Albeit at a lower extent the same pattern was observed in the retinal

organoid transduced with AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 (**Figure 3F**), suggesting

that the NT sgRNA exhibited greater impact on the transcriptome. To determine the specific

cell types that were more sensitive to Cas13bt3 expression, we identified unique DEGs from

each cell type (to remove common effects from AAV delivery and culturing conditions) across

the retinal cells and found macroglial and bipolar cells to exhibit the highest DEGs, which was

especially higher in the NT sgRNA group (Macroglial - NT: 5571 vs VEGFA: 2928, Bipolar

– NT: 3725 vs VEGFA: 1633) (**Figure 3G**).

To understand this effect from the NT sgRNA, we further identified DEGs unique to

the NT sgRNA and VEGFA sgRNA group and performed GO analysis to identify responses

exclusive to the delivered sgRNA. However, ubiquitin protein-like (UBL) associations (e.g.,

"*UBL ligase binding*", "*UBL transferase activity*") were top hits for both groups across

macroglial, bipolar, amacrine and RPE cells (**Figure 3H and Extended data 2**). Ubiquitin

processes are often activated in response to immune stress to facilitate the clearing of proteins

inside cells [28], and our results suggest the delivery of Cas13bt3 with an activating sgRNA (non-

targeting or targeting) to significantly elevate the immune response in retinal cells. This stress

response is likely higher with the NT sgRNA group due to unspecific RNA binding, causing

the increased transcriptomic impact. The strong stress response from all cells to Cas13bt3

delivery is also likely to have made sgRNA-specific responses challenging to distinguish.

Figure 3. Cas13bt3 treatment and single-cell RNA sequencing of 3D retinal organoids.
(A) Schematic of 3D retinal organoid scRNA-seq experimental procedure. **(B)** UMAP plots showing retinal cell type clusters for the untreated, NT sgRNA and VEGFA sgRNA treatment groups. **(C)** Expression of cell-type specific markers in cone cells. **(D)** Infection rate of retinal organoids by cell type and treatment group. **(E)** Multi-volcano plots showing differentially expressed genes (DEGs) between Cas13bt3 positive and negative cells from the NT sgRNA and **(F)** VEGFA sgRNA treatment groups. **(G)** Number of unique DEGs across retinal cell types. **(H)** GO molecular function terms related to unique DEGs from Cas13bt3-NT sgRNA treatment for macroglial and bipolar cells. UBL = Ubiquitin-like protein.

Cas13bt3 significantly knockdown *VEGFA* mRNA in retinal organoids.

Next, we examined Cas13bt3 targeting efficiency and effect on *VEGFA* expression across the retinal cell types. First, we determined *VEGFA* expression in control untreated organoids. *VEGFA* mRNA is present across all retinal organoids cell types. Expression of VEGFA varied significantly between retinal cell types with macroglial, rod, cone photoreceptors, and bipolar cells expressing significantly less *VEGFA* than RPE cells (**Figure S8A, B**).

Furthermore, *VEGFA* expression was also detected in all treated organoids, albeit at different levels (**Figure S8C-E**). DEGs were detected in all cell types between retinal organoids transduced with Cas13bt3-VEGFA sgRNA and Cas13bt3-NT sgRNA. However, a significant downregulation of *VEGFA* was only detected in RPE cells (**Figure S9**). While Cas13bt3 positive cells have been identified, the abundance of Cas13bt3 reads within cells may still vary (**Figure 4A**) and affect determination of targeting efficiency. We hypothesised that increased intracellular Cas13bt3 transcripts allows for higher targeting efficiency and thereby increased silencing of *VEGFA* mRNA (**Figure 4B**). We first examined the number of Cas13bt3 reads in cells that received AAV2.7m8.U6.NTsgRNA.Cas13bt3 and AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 treatments (**Figure 4C and S10**). To accurately determine Cas13bt3 effect and to understand abundance of the targeting complex within each

cell we examined AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 treated cells. Here, we found RPE

and macroglial cells expressed high levels of Cas13bt3 expression while most cones, rods and

horizontal and bipolar cells only showed low levels of Cas13bt3 expression (**Figure 4C**). We

selected a threshold to classify high ($\geq$ 50 reads) and low (<50 reads) expression of Cas13bt3

that enables evaluation of targeting efficiency. Next, we quantified *VEGFA* mRNA expression

in cells expressing high and low levels of Cas13bt3. To this end, we pooled cells within the

high and low Cas13bt3 thresholds and compared average *VEGFA* expression against Cas13bt3

negative cells to determine if loss of *VEGFA* is from is dependent on high levels of Cas13bt3.

Significant decrease of *VEGFA* in high Cas13bt3-expressing cells compared to low Cas13bt3-

expressing cells would indicate *VEGFA* silencing from increased targeting activity, while

significant reduction compared to negative cells would indicate *VEGFA* loss from viral

delivery. Here, we observed that RPE cells were the only cells to show significant reduction of

VEGFA expression in the high Cas13bt3 when compared to low Cas13bt3 cells ($P < 0.05$)

(**Figure 4D**, red square). Viral delivery appeared to upregulate *VEGFA* expression in

macroglial cells with both low and high Cas13bt3-expressing cells exhibiting significant

increase in VEGFA expression (**Figure 4D**). Interestingly, the increase of *VEGFA* was not

significant between low and high Cas13bt3 expressing macroglial cells, suggesting a

considerable loss of *VEGFA* (**Figure 4D**). We therefore further investigated retinal cells that

showed moderate transduction (~25%) of Cas13bt3-VEGFA sgRNA (i.e. amacrine, horizontal

and macroglial) and compared *VEGFA* expression to cells that received Cas13bt3-NTsgRNA.

Promisingly, we observed near significant downregulation of *VEGFA* in amacrine and

horizontal cells ($P = 0.075$ and $P = 0.051$ respectively), suggesting higher transduction rates

may lead to significant downregulation in the other cells as well (**Figure S11**). We then studied

DEGs in the RPE cells to determine if the off-target genes previously identified were affected

from the significant Cas13bt3-mediated *VEGFA* silencing. Importantly, none of the off-target

genes we previously identified were significantly affected (**Figure 4E and S12**). Finally, GO

analysis revealed that "*PDGF receptor binding*" was among the top five downregulated

processes (**Figure 4F and Extended data 3**), akin to our observations in HEK293FT cells.

While we only observed significant *VEGFA* mRNA knockdown in RPE cells, whether

this is a result of higher transduction or RPE cells carry cell-specific factors that are conducive

for Cas13 activity needs further investigation. Regardless, with RPE cells being major

contributors of *VEGFA* expression, our results show that Cas13bt3 can be a potent tool against

diseases such as DR and nAMD.

Figure 4. VEGFA knockdown efficiency in human retinal organoids. **(A)** Schematic of
variable Cas13bt3 expression in retinal organoid cells and **(B)** relationship between CRISPR-
Cas13bt3 expression and targeting efficiency. **(C)** Histograms showing level of Cas13bt3
expression by number of reads across retinal cell types in the VEGFA sgRNA group. **(D)** Violin
plots of *VEGFA* expression against level of Cas13bt3 expression across retinal cell types
compared to Cas13bt3 negative cells. Red box indicates cells of interest for high VEGFA
targeting activity. **(E)** Volcano plot of DEGs between RPE cells with high and low Cas13bt3
expression. Significant knockdown of *VEGFA* is highlighted in red. **(F)** Top five significant
GO molecular function terms between RPE cells expressing low and high Cas13bt3. Statistical
analyses were conducted using Student's t-test **(E)**. PPi: Pyrophosphatase. TM:
Transmembrane.
373

VEGFA knockdown impedes disease progression in humanised mouse model of

proliferative diabetic retinopathy.

Lastly, we sought to investigate if *VEGFA* knockdown can control neovascularisation *in vivo*

in humanised models of DR. This offers a potential new tool against eye diseases causing

neovascularisation. To this end, we utilised the trVEGF029 (Kimba) mouse, a humanised

transgenic mouse model where the neuroretina overexpresses h*VEGFA* leading to

neovascularisation like human proliferative DR[29]. To ensure that optimal transduction of the

retina is being achieved, we delivered a mCherry reporter gene under the control of CMV

promoter using the AAV2.7m8 serotype, known to have excellent transduction of the retina,

when delivered intravitreally into C57BL/6 mice (**Figure S13A and B**).

Immunohistochemistry analysis of AAV2.7m8.CMV.mCherry treated retinal flatmounts

showed high levels of fluorescence through all retinal layers (RGCs, bipolar and photoreceptor

cells) 4 weeks following injections, confirming quick onset and deep penetration of the retina

from intravitreal delivery (**Figure S13C, n = 6 eyes from 3 animals**).

Next we treated Kimba mice with AAV2.7m8 viruses driving our codon optimised

Cas13bt3 carrying NT sgRNA (AAV2.7m8.U6.NTsgRNA.Cas13bt3) or VEGFA sgRNA

(AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3). Due to severe disease progression in homozygous

Kimba (*VEGFA*$^{+/+}$), we opted for the heterozygous (*VEGFA*$^{+/-}$) breed. Six-week-old Kimba

mice received bilateral, intravitreal injections of AAV2.7m8.U6.NTsgRNA.Cas13bt3 or

AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3. Eight weeks following treatment eyes were

analysed for *VEGFA* mRNA expression, neovascular growth, and retinal degeneration (**Figure**

5A). Similar to the control mCherry treatments, high transduction of photoreceptor cells was

observed as demonstrated by co-localisation of mCherry and HA-tag (Cas13bt3) with

Rhodopsin positive cells in the ONL (**Figure 5B, C**). In addition qPCR analysis showed high

expression of Cas13bt3 in Kimba retinas injected with AAV.2.7m8.U6.NTsgRNA.Cas13bt3

and AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 (**Figure 5D,** n = 3-12 retinas). This

corresponded with ~50% reduction (*P* < 0.001) of human VEGFA (*hVEGFA)* mRNA (**Figure**

5E), but not mouse VEGFA (mVEGFA) (**Figure 5F**) in the

AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 group, while no reduction was observed with

AAV2.7m8.U6.NTsgRNA.Cas13bt3 or no treatment (**Figure 5E, F,** n = 3-12 retinas). These

results confirmed specific knockdown of *hVEGFA* from Cas13bt3, despite high

complementarity of *VEGFA* sequences (29 out of 30 nucleotides) in the targeting region

(**Figure 5G**). RNA sequencing further confirmed that mouse VEGFA expression and predicted

off-target genes were not significantly altered, indicating safety of Cas13bt3-mediated RNA

silencing (**Figure S13**).

To confirm if *VEGFA* mRNA knockdown influenced disease progression, we analysed

fundus fluorescein angiography (FFA) images from Kimba eyes for vessel leakage and

quantified vessel density and number of branch points, which are signs indicative for the

neovascular process. We used fluorescein to label vessels of the

AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3, AAV2.7m8.U6.NTsgRNA.Cas13bt3 and untreated

eyes. We observed large spots of vessel leakage, indicating neovascularisation had progressed

in both untreated and NT treated eyes (**Figure 5H,** green expression and white arrows).

However, in the AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 group, no leakage was observed

(**Figure 5H,** VEGFA targeting). Immunohistochemistry against laminin was used to image

through the retinal vascular layers (superficial, intermediate and deep vascular plexuses)

(**Figure S16A and B**). Blood vessels appeared diminished in the intermediate and deep

vascular layers for the VEGFA sgRNA group. In the NT sgRNA group, the neovascular

process could be observed through leaky vessel structure in the same layers (**Figure S16C,** n

= 4 retinas). The AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 treated eyes also showed

significantly lower vessel density and branch points compared to baseline at 6 weeks post-

treatment, unlike the untreated and the AAV27m8.U6.NTsgRNA.Cas13bt3 eyes (**Figure 5I**

and J, n = 4-13 retinas, P<0.05). Next, to determine whether *VEGFA* knockdown leads to

rescue of overall retinal structure. we performed optical coherence tomography (OCT) to

analyse total retinal thickness (TRT)   before and after treatments. We observed no significant

changes between the untreated and treated groups, and a general decline in TRT across all

animals (**Figure S17**). We postulate that the overexpression of *hVEGFA* promotes rapid retinal

degeneration that could not be controlled or reversed through *VEGFA* silencing.

Overall, our findings showed Cas13bt3-mediated RNA silencing to significantly reduce

VEGFA mRNA expression, specifically target *hVEGFA* and control hallmark signs of

neovascularisation in a humanised mouse model, promising a useful alternative to current anti-

VEGF drugs.

Figure 5. In vivo VEGFA mRNA knockdown in humanised Kimba mice. **(A)** Schematic of experimental procedure for intravitreal treatment of Kimba mice. **(B and C)** Immunohistochemistry of retinal flatmount and cryosection showing transduction of AAV2.7m8 and expression of Cas13bt3 (Magnification: 20X and 40X, respectively). **(D)** Expression of Cas13bt3 from Kimba mice retinas. **(E)** Expression of mouse and human *VEGFA* mRNA in Kimba retinas. **(F)** *VEGFA* mRNA targeting regions in human and mouse transcriptome. **(G)** Plot of vessel density and number of branch points from untreated and treated Kimba eyes at baseline (6 weeks) and endpoint (14 weeks) analysed from *in vivo* central fluorescein fundus angiography (FFA). **(H)** Central fluorescein fundus angiography (FFA) images obtained from Kimba mice at baseline (6 weeks) and endpoint (14 weeks) after AngioTool analysis and quantification. Images were edited using Adobe Photoshop to remove central retinal arteries and veins. Green spots and white arrows indicate leakage from blood vessels. Data presented as mean ± SEM. Statistical analysis was conducted by one-way

450 ANOVA (**D and E**) with Tukey's multiple comparison test and Multiple paired t-tests (**G**).
451 *P≤0.05, **P ≤0.001, ****P≤0.00001.
452

DISCUSSION

Our comprehensive *in vitro*, *ex vivo* and *in vivo* characterisation of Cas13bt3 activity and performance in the retina provides novel insights for therapeutic use of the enzyme. Cas13bt3 was first described only two years ago but has garnered significant attention for its compact size and reported specificity[11]. Structural characterisation and activation mechanisms have recently been described spearing immense interest in the molecule for therapeutic development[16, 30]. Due to ongoing controversy over the exact specificity of Cas13 nucleases[10], we aimed to characterise Cas13bt3 efficiency and specificity in mouse and human cells. We chose to target the *VEGFA* transcripts, as this growth factor is essential for both physiological and pathological angiogenesis. In the retina, physiological functions include its role as a survival factor for retinal ganglion cells[31], photoreceptors, Müller cells[32], and RPE[33]. In disease, overexpression of VEGFA leads to neovascularisation and vision loss. Therefore, clinical applicability of VEGFA downregulation in specific cell types of the retina is of great importance.

Previous studies have demonstrated transcript knockdown using CasRx proteins[11, 12]. Notably, CasRx-mediated *VEGFA* mRNA knockdown has also been shown to reduce choroidal neovascularisation after silencing VEGFA in RPE cells *in vivo* in laser-induced choroidal neovascularisation model[4]. Our results confirmed that Cas13bt3 mediates significant knockdown of *VEGFA* in human cell lines, hESC-derived retinal cells, and the mouse retina. We further investigated the transcriptomic impact of expressing active Cas13bt3 in the cells and found that initiating Cas13bt3's target search process with a non-targeting sgRNA can lead

to significant RNA binding, upregulating the expression of several genes. A targeting sgRNA,

such as with VEGFA, appears to narrow the target-search process and thereby reduce

transcriptomic impact.

In vitro, and specifically in HEK293FT cells, several studies have shown Cas13

nucleases to be highly precise[10]. Most of these studies, however, compared differentially

expressed genes to a NT control, not accounting for transcriptomic perturbations from a Cas13-

NT sgRNA complex. A recent study showed high collateral activity from CasRx when

compared to an empty vector[34]. Here, we compared targeting Cas13bt3 to vector controls

showing minimal, but positive collateral activity. More importantly, we show that Cas13 with

a NT sgRNA has significant transcriptomic impact resulting in high levels of DEGs. This

demonstrates that NT sgRNAs are not appropriate controls for determining specificity of Cas13

nucleases raising awareness against their use.

We also show that multiple sgRNAs can be incorporated within a single AAV vector

with Cas13bt3 driven by a strong ubiquitous promoter (CMV) for targeted knockdown of

VEGFA mRNA. Even though this appears to be an attractive approach to enhance transcript

knockdown our study established that multiple sgRNAs failed to improve transcript

degradation. The rate of mRNA knockdown was comparable to single sgRNA treatments.

Previously, we similarly observed with multiplexed RNA editing with CasRx carrying three

VEGFA-targeting sgRNAs, the multiple sgRNA system unexpectedly led to decreased

knockdown efficiency[19], raising the suspicion that substrate competition or inadequate Cas13

activation plays an inhibitory role in RNA targeting efficiency. Interestingly, the comparable

levels of DEGs in both single and multiple sgRNA vectors suggest that Cas13bt3 activation is

not drastically elevated with the presence of additional sgRNAs. While we did not investigate

the delivery of sgRNAs targeting multiple genes, our results show multiple sgRNAs do not

increase overall collateral activity, which might prove useful when several genes must be

targeted simultaneously to control disease progression. This approach could potentially be

applied even to ocular neovascularisation by simultaneously targeting the *VEGFA* and *PDGF*

genes for enhanced treatment[35].

Furthermore, we investigated for the first time AAV-mediated Cas13 RNA expression

and editing in retinal organoids at single cell resolution. While several studies have

demonstrated *VEGFA* silencing for control of neovascularisation, previous studies have

focused on specific cell types that are believed to be major contributors to *VEGFA* expression,

like RPE and Müller cells[4, 36]. While this proves useful to demonstrate proof-of-concept, it does

not provide a comprehensive understanding relevant for therapeutic development as *VEGFA*

silencing in one cell type may have collateral effects on other retinal cells. Retinal organoid

models enable the testing of therapies in an environment that mimics the human retina

containing lamination and major retinal cell types of the developing retina. Here we sought to

understand retina-wide effect following delivery of Cas13bt3 as well as targeted silencing of

VEGFA. Our findings not only support the use of Cas13bt3 for ocular gene therapy, but also

brings to light several considerations for its use. First, we showed transduction across all retinal

cell types studied with the clinically relevant AAV2.7m8 vector (NCT04418427) confirming

its effectiveness as a vector for retinal transduction and supporting its clinical utility[37,38].

Among the retinal cell types studied, bipolar cells showed the least transduction by AAV2.7m8,

while RPE cells seemed more susceptible. It is likely that these differences are due to cell-

specific transduction susceptibility, as RPE cells are easily transduced with a variety of AAV

capsids[39]. It should also be noted that overall transduction was determined using an arbitrary

cut-off (Cas13bt3 read counts $\geq$ 10) based on our previous experience with the retinal

organoids[26]. NT sgRNA activity following gene therapy led to numerous DEGs when

compared to VEGFA targeted sgRNA. This emphasises the need for a well-designed targeting

sgRNA for reduced collateral activity. Retinal organoids showed significantly more DEGs than

HEK293FT cells, which is expected partly due to cellular response to virus delivery. Still,

notwithstanding response to AAV, the higher DEG levels also indicate that retinal cells are

more sensitive to Cas13, demonstrating the importance of studying nuclease activity and

efficiency in physiological relevant models, an important consideration for Cas-mediated gene

therapies. As preclinical studies with Cas13 nucleases are commonly performed in HEK293FT

cells, its specificity has often been exaggerated in the literature. Our results strongly suggest

that Cas13 specificity must be evaluated in relation to untreated or empty vector controls in

organ-specific cells to accurately determine safety. On the front of therapeutic potential, we

could only confirm significant targeted *VEGFA* mRNA knockdown in RPE cells, which is

likely due to strong expression of Cas13bt3 in these cells. Lower copies of Cas13bt3 in the

other cells may have resulted in the lack of significant *VEGFA* knockdown, and further studies

are required to understand the levels of Cas13 expression necessary in other retinal cells for

effective activity. Testing of cell type specific promoters could be beneficial as CMV promoter

is known for its susceptibility to silencing in certain cell types, for example stage specific PSC-

derived photoreceptor cells (REF). Importantly, with RPE cells being the primary producers of

VEGFA in the retina, the significant downregulation of *VEGFA* mRNA in RPE strongly

suggests the benefits of Cas13bt3 gene therapy as a potential anti-VEGF tool for neovascular

eye disease.

Lastly, we tested the anti-VEGF potential of Cas13bt3 in the retina of a humanised

mouse model of diabetic retinopathy, the Kimba mouse. We opted to examine anti-VEGF

potential of Cas13bt3 in Kimba mice as it shows chronic disease and all major signs of

proliferative retinopathy [29]. Importantly, as Kimba mice expresses human *VEGFA*, the same

vectors tested in retinal organoids could also be used *in vivo*. Others have shown the control of

ocular neovascularisation using mouse models of laser-induced injury[4], streptozotocin (STZ)-

induced diabetes[40], or oxygen-induced retinopathy (OIR)[41]. However, these models have some

drawbacks. The laser-induced choroidal neovascularisation model often used in anti-VEGF

preclinical studies does not accurate model the chronic retinal degeneration observed in DR

and nAMD patients and presents *VEGFA* overexpression only as a response to injury[42].

Diabetic mouse models do not recapitulate proliferative retinopathy which is often the cause

of severe vision impairment. Vascular pathogenesis is also delayed and varied based on

response to STZ[43]. OIR mouse models rapidly present proliferative retinopathy, however

neovascularisation regresses within a week[44].

Following RNA editing gene therapy in Kimba retinas, we demonstrated the specific

knockdown of human *VEGFA*, with no significant impact on endogenous mouse *VEGFA*. This

shows strong evidence for the specificity of targeted Cas13bt3 activity. Silencing of Cas13bt3

VEGFA also rescued vascular phenotypes showing an improvement in controlling vascular

growth and leakiness. The major concern with current anti-VEGF treatment is the need for

frequent intravitreal injections. Recent efforts to address this problem have employed gene

editing strategies like targeting *VEGFA* using CRISPR-Cas9 [36, 45]. However, efficacy of such

treatments are limited by efficient and safe delivery of vectors due to the size of Cas9[46, 47], poor

efficiency of compact Cas9 nucleases[48] or have reported the risk of genomic alterations from

long-term expression of Cas9[49]. Our results indicate that a single intravitreal injection could

achieve durable and potent while reversible silencing of *VEGFA*, addressing drawbacks with

current gene editing strategies. While RNA editing is thought to be safer than DNA-targeting

options, it is prudent to consider effects of long-term Cas13 expression, and Cas13 systems

amenable to regulation are of consideration for therapeutic development.

There are a few limitations to our study. First, we did not extensively study Cas13bt3

expression in ARPE19 cells, where we also observed potent knockdown. While HEK293FT

appears resistant to high levels of collateral activity, ARPE19 cells may exhibit significantly

higher levels of transcriptomic effects. This is supported by the high levels of DEGs we

observed in various cells from the retinal organoids, suggesting significant inflammatory

responses as well as cell-specific collateral activity. Specifically, the upregulation of UBL

processes across cell types regardless of VEGFA silencing, strongly indicate increased cellular

stress from delivery of Cas13. As RNA targeting molecules like Cas13 require persistent

expression for therapeutic effect, this could pose a significant safety concern. Second, while

vessel density and branching were noticeably reduced, we could not extensively study recovery

of phenotype in our mouse model due to severity of the disease. OCT readings showed

significant retinal degeneration at treatment timepoint, making rescue of phenotype

challenging to assess. Lastly, we were also unable to confirm reduction of VEGFA protein

from the mice retinas as not enough protein could be obtained for quantitative analysis.

CRISPR-Cas technology has advanced at breakneck speed in the last decade

revolutionising medicine and offering potential treatment opportunities to numerous previous

uncurable diseases. The recent FDA approval of an *ex vivo* CRISPR therapy for sickle-cell

disease (branded Casgevy) is evidence of this progress. However, a comprehensive

understanding of the CRISPR nucleases is crucial. In this study we offer novel insights into

Cas13 targeting mechanisms and its utility in targeting retinal and RPE cells providing

evidence for the potential of Cas13-based therapies for ocular disease. This study demonstrated

that the hypercompact Cas13bt3 can be efficiently delivered and expressed in various retinal

cell types effectively controlling *VEGFA* expression and neovascularisation. In-depth

evaluation of collateral activity in retinal cells would facilitate development of a safe and

effective Cas13-mediated anti-VEGF AAV gene therapy.

MATERIALS AND METHODS

Plasmid construction

The protein sequences of CasRx and Cas13bt3 were gifts from P. Hsu[50] and H. Yang[11], respectively. Cas13bt3 was codon optimised using online tools from Integrated DNA Technologies (IDT), GenScript and Benchling. The all-in-one AAV plasmids for shRNA, CasRx and Cas13bt3 were designed as previously described[19], and synthesised by GenScript. All plasmids used in this study are available from Addgene.

Cell culture and transfections

HEK293FT cells (American Type Culture Collection, ATCC) were maintained in Dulbecco's Modified Eagle's Medium (DMEM; Thermo Fisher Scientific) supplemented with 10% (v/v) fetal bovine serum (FBS, Bovogen Biologicals) and 1% (v/v) antibiotic-antimycotic (Thermo Fisher Scientific) in a humidified 5% CO_2 incubator at 37°C. ARPE19 (ATCC) cells were maintained in DMEM/F-12 (Thermo Fisher Scientific) supplemented with 10% (v/v) FBS and 1% (v/v) antibiotic-antimycotic.

RT-qPCR

At 48 hours after transfection, RNA was extracted from cells using the Monarch® Total RNA Miniprep Kit (New England Biolabs and 200ng of RNA was converted to complementary DNA (cDNA) using High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific) according to the manufacturers' instructions. cDNA was diluted with water at the ratio of 1:10. qPCR was performed with 2uL of cDNA using the TaqMan™ Fast Advanced Master Mix (Thermo Fisher Scientific) according to manufacturers' instructions with the following TaqMan probes human VEGFA (Hs00900055_m1; Thermo Fisher Scientific), mouse VEGFA

(Mm00437306_m1; Thermo Fisher Scientific), human GAPDH (Hs02786624_g1; Thermo

Fisher Scientific), mouse GAPDH (Mm99999915_g1, Thermo Fisher Scientific). Mouse

eyeballs were enucleated and stored in 4°C in RNALater® until dissection. Retina was dissected

from injected animals and placed in lysis buffer and sonicated for 2 seconds prior to RNA

extraction and qPCR as previously described.

Western blot

At 48 hours after transfection, cells were harvested and collected in Pierce™ RIPA buffer

(Thermo Fisher Scientific) containing Pierce™ Phosphatase Inhibitor Mini Tablets (Thermo

Fisher Scientific). Cell lysate was sonicated for 5 seconds and centrifuged for 15 minutes at

13300rpm at 4°C. The supernatant was collected, and protein levels were quantified using the

Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific) as per manufacturers' instructions.

For Western blot, 10μg of protein mixed with NuPAGE™ LDS Sample Buffer (4X) (Thermo

Fisher Scientific) and loaded in NuPAGE™ 4 to 12% Mini Protein Gel (Thermo Fisher

Scientific) for gel electrophoresis at 150V for 30 minutes. Transfer to Amersham™ Hybond™

PVDF membrane (Cytiva) was performed at 30V for 1 hour. Protein blot was blocked with 5%

skim milk for 1 hour, followed by incubation with HA-tag (C29F4) Rabbit mAb (Cell Signaling

Technology®) or Anti-Actin Antibody, clone C4 (Merck) overnight at 4°C. Blots were then

stained with Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, HRP

(Invitrogen™) or Goat anti-Rabbit IgG (H+L) Secondary Antibody, HRP (Invitrogen™) for 2

hours at room temperature. Blot was incubated with Pierce™ ECL Western Blotting Substrate

for 1 minute before imaging using Bio-Rad ChemiDoc. All washes were performed with 1X

PBST for 5 minutes, repeated 3 times.

ELISA

At 48 hours after transfection, 100μL of supernatant was collected and ELISA was performed using the Human VEGF DuoSet ELISA (R & D Systems) as per manufacturers' instructions. Briefly, 96-well plates were incubated with Capture Antibody in ELISA Plate coating buffer overnight at room temperature. Plates were then blocked with Reagent Diluent for 1 hour at room temperature, and 100uL of supernatant was incubated for 2 hours at room temperature. Then, Detection antibody was added and incubated for 2 hours at room temperature. After incubation with Streptavidin-HRP and Substrate reagent for 20 minutes each, the provided stop solution was added, and plate was read using the Tecan Spark® Micro plate reader at 450nm with reference wavelength 570nm.

Immunohistochemistry

Retinal flatmounts were washed with 1X PBS containing 0.5% Triton-X-100 (PBST) and permeabilised in -80°C for 15 minutes. After washing with PBST, flatmounts were incubated in anti-rhodopsin antibody (1:500; Abcam) with 2% donkey serum and 2% PBST in 4°C for 72 hours. This was followed by incubation in Alexa Fluor 488 AffiniPure Donkey Anti-Mouse IgG (H+L) (1:500; Jackson ImmunoResearch) at room temperature for 2 hours. After washing, the flatmounts were stained with Anti-Laminin-2 (α-2 Chain) antibody, Rat monoclonal (1:250; Merck) and incubated in 4°C for 5 nights. This was followed by incubation in Alexa Fluor® 647 AffiniPure Donkey Anti-Rat IgG (H+L) (1:250; Jackson ImmunoResearch) at room temperature for 2 hours. Then, the retinae were washed and incubated in mCherry Monoclonal Antibody (16D7), Alexa Fluor 594 (1:500; Thermo Fisher Scientific) at 4°C for 72 hours. Retinal cryosections were blocked using serum-free protein block (Agilent Dako) at room

668 temperature for 1 hour, followed by incubation at 4°C overnight with primary antibody as

669 previously described. Secondary antibody incubation was performed at room temperature for

670 1 hour as previously described. Slides were stained with NucBlue™ Live ReadyProbes™

671 Reagent (Hoechst 33342; Thermo Fisher Scientific) for 10 minutes before mounting as

672 previously described and imaged on the Stellaris 5 confocal microscope.

hESC Maintenance and Retinal Differentiation Culture

675 hESCs were maintained on feeder-free conditions on E8 (Thermo Fisher) and Geltrex-coated

676 six-well plates. In brief, for retinal differentiation hESCs were maintained until 90%–95%

677 confluent, then FGF-free medium was added to the cultures for 2 days followed by a neural

678 induction period where RPE and retinal vesicles appeared. NRVs were excised and kept in low

679 binding 96-well plates for maturation.

AAV transduction of retinal organoids

Retinal organoids were maintained in ALT90 medium as previously described (West et al.,

2022) and AAV transduction was performed during the 15-19 weeks of differentiation. On the

day of transduction, AAV vectors were diluted in 300μL of fresh ALT90 medium at titer of

1x10E11 vg/organoids. The following day treated wells were supplemented with 700μL of

fresh ALT90 medium. Transduced organoids were incubated for 72 hours in the AAV vector-

containing medium. After the 72-hour incubation, the vector-containing medium was aspirated,

and 1mL of fresh ALT90 medium was added to each well. Transduced organoids underwent 6

weeks culture until harvesting for experimental procedures.

Immunohistochemistry of retinal organoids

Retinal organoids were fixed for 45 minutes in 4% paraformaldehyde and incubated overnight in 20% sucrose, prior to embedding in OCT. Cryosections (14 μm thick) were collected for analysis and preserved at - 20°C. Cryosections were blocked in 5% goat serum and 1% bovine serum albumin in PBS for 2 hours. Primary antibodies (ARR3: 1:200, Rhodopsin: 1:1000, VSX2: 1:200, CRALBP: 1:200, PROX1:1:200, and HuCHuD:1:500) were incubated overnight at 4°C in blocking solution with 5% goat or donkey serum. After washing of primary antibody with PBS sections were incubated with secondary antibody in blocking solution for 2 hours at RT, washed and counter-stained with DAPI (Sigma-Aldrich). Alexa fluor 488 and 546 secondary antibodies (Thermo Fisher Scientific) were used at a 1:500 dilution.

Intravitreal injections

All animal procedures were approved by the Animal Ethics Committees of University of Melbourne and conducted in accordance with The Association for Research in Vision and Ophthalmology (ARVO) Statement for the Use of Animal in Ophthalmic and Vision Research. Ethics approval was obtained from The Florey Institute of Neuroscience and Mental Health Animal Ethics Committee (22-036-UM). The Akimba mice developed by Rakoczy *et al.* [51] and the heterozygous Kimba mice ($hVEGFA^{+/-}$) were used for this study. Mice were bred in the Melbourne Brain Centre animal facility (Kenneth Myer Building, Parkville, VIC, Australia), and were housed in a well-ventilated environment that was kept at a constant room temperature of 21°C, with a 12-hour diurnal light-dark cycle (lights on: 7am, off: 7pm; room illumination < 50 lux). Mouse chow (Barastoc, Melbourne, VIC, Australia) and water were provided ad libitum. 6-week-old Kimba mice were anesthetized with ketamine (80mg/kg) and xylazine (10mg/kg) solution mix (Troy Laboratory, Smithfield, NSW, Australia; i.p. with 30G needle)

and given a drop of 0.5% proxymethacaine hydrochloride (Alcaine™, Alcon Laboratories,

Frenchs Forest, NSW, Australia) and 0.5% tropicamide (Mydriacyl™, Alcon Laboratories).

Baseline fundus fluoresceine angiography (FFA) were taken (as described below), and animals

intravitreally injected with 1uL of AAV2.7m8 viruses using Hamilton syringe attached to a 33-

gauge dental needle. Post injection, mice were subcutaneously administered 0.3mL of

atipamezole hydrochloride (0.25mg/mL) and allowed to recover on heating pads with food and

water *ad libitum*.

Fundus fluorescein angiography and optical coherence tomography

Retinal vasculature was assayed at baseline prior to intravitreal injection of AAVs and 8 weeks

after AAV-treatment using fundus fluorescein angiography (FFA) (Spectralis, Heidelberg

Engineering, Heidelberg, Germany) immediately after intraperitoneal injection of sodium

fluorescein (1%, 100 µL/kg, fluorescite 10%; Alcon Laboratories). Five images (central, nasal

superior, temporal superior, nasal inferior, and temporal inferior) were obtained for each eye

and analysed using AngioTool[52] for vessel density and branch points using default settings.

In vivo retinal structure was assayed with spectral domain optical coherence

tomography (SD-OCT) (Spectralis®, Heidelberg Engineering, Heidelberg, Germany) with

general anaesthesia, mydriasis and corneal hydration (Systane, Alcon Laboratories)

maintained. During image acquisition, retinal volumes ($8.1 \times 8.1 \times 1.9$ mm) centred on the

optic nerve head were acquired each consisting of 121 evenly spaced horizontal B-scans. Each

B-scan comprised of 768 A-scans with 3.87 µm axial depth resolution, 9.8 µm lateral

resolution. OCT scans were captured with an automated real-time tracking average of 6 frames,

at an average speed of 85,000 A-scans per second. The Heidelberg Eye Explorer 2 OCT reader

plugin (Heidelberg Engineering) was used for analysis. All retinal layers (retinal nerve fibre

layer, RNFL; ganglion cell inner plexiform layer, GCIPL; inner nuclear layer, INL; outer

plexiform layer, OPL; outer nuclear layer, ONL; photoreceptor segments, PR segment) as well

as total retinal thickness (TRT) were automatically segmented (Heidelberg Eye Explorer 2

software). A circular ring around the optic nerve head (Early Treatment Diabetic Retinopathy

Study [ETDRS] outer ring) was employed for analysis.

Determination of candidate off-target genes

Off-targets of the gRNAs were predicted using a sequence-based approach. gRNAs in length

of 30 bps were first aligned to the human transcriptome (Grch38 cdna from emsembl release

100) using blastn (megablast). Flexible options were set to capture the potential offtargets

which could tolerate up to 7 mismatches in this step, i.e., max_target_seqs = 10000, evalue =

10000, word_size = 5, perc_identity = 0.6. Secondly, the candidates were further filtered with

the threshold that at least 13 bp perfect match should exist in each alignment.

Bulk RNA-seq data analysis

Raw data processing was performed using FASTQC [53] to check the reads quality and

Trimmomatic [54] tool to further remove the bases and reads with low quality, adaptors and

barcodes. The clean reads were aligned to the reference human genome (version GRCh38),

utilizing STAR v2.7 [55] as the aligner. For RNA quantification, RSEM [56] was employed to

normalize raw read counts, transforming them into TPM (transcripts per million) values,

providing a relative expression level that is theoretically comparable across samples. The

magnitude (log2-transformed fold change) and significance (P-value) of differential expression

between groups were then calculated. Differentially expressed genes (DEGs) were selected

based on the criteria of a p-value < 0.05 and |log2 fold change| ≥ 1. Gene set enrichment analysis

(GSEA) [57], focused on Gene Ontology[58], was conducted using the R 4.3.1 Bioconductor

package clusterProfiler[59] to elucidate whole transcriptomic patterns between groups. To

account for multiple testing, P-values underwent adjustment through the Benjamini-Hochberg

false discovery rate (BH-FDR).

Single-cell RNA-seq data analysis

After checking the reads quality, the sequencing files were aligned to reference human genome

(version GRCh38) using 10X Genomics CellRanger software (v7.1.0) count pipeline to remove

non-cellular barcodes and keep the cellular barcodes and gene matrix. To further distinguish

the Cas13bt3-valid and -invalid cells, FASTX-toolkit [60] was used to keep the reads with cellular

barcodes and convert FASTAQ file into FASTA file. Then, Blastn [61] was used to compute the

similarity between cellular reads and Cas13bt3 reference sequence.

Based on the quality of the sequencing reads, appropriate cutoff values for both the percentage

of identical positions and the alignment length were employed to delineate cells as either

Cas13bt3-valid or Cas13bt3-invalid. For either Cas13bt3-valid or -invalid cells, we further

used Seurat v4.3 for data pre-processing, data normalization, feature selection, data scaling,

dimensional reduction and clustering. To refine the characterization of cell types within each

cluster, ScType [62] was employed, leveraging genesets specific to eight retinal cell types—

amacrine, bipolar, cones, horizontal, macroglial, microglial, rods, and RPE—validated by Kim

et al. [27]. The magnitude (log2-transformed fold change) and significance (P-value) of

differential expression between cell types were then calculated. Differentially expressed genes

(DEGs) were selected based on the criteria of a p-value < 0.05 and |log2 fold change| ≥ 1. Gene

set enrichment analysis (GSEA) [57], focused on Gene Ontology [58], was conducted using the R

4.3.1 Bioconductor package clusterProfiler [59] to elucidate whole transcriptomic patterns

between groups. To account for multiple testing, P-values underwent adjustment through the

Benjamini-Hochberg false discovery rate (BH-FDR).

Statistical analysis

To assess the strength of the relationship between gene expression values and the number of

Cas13bt3-valid read counts following log2 transformation in each cell type, Spearman's rank

correlation coefficient was employed. The associated *p-value* serves as an indicator of the

likelihood or probability that any observed correlation is merely a result of chance.

Acknowledgements

The authors would like to thank Liuhui Huang, Jesse Gardner-Russell and Yujie Wang for their assistance with retinal dissections, Prof Alex Hewitt and Dr Leilei Tu for providing resources for vector construction. The authors also extend their gratitude to the Vector and Genome Engineering Facility (VGEF) and the Single Cell Analytics Facility at the Children's Medical Research Institute (CMRI) for producing AAV vectors and the scRNA-seq experiments respectively and Azenta Life Sciences for conducting RNA sequencing. Schematic in figures were designed using Biorender.com.

Funding

This work was supported by grants from the National Health and Medical Research Council of Australia (NHMRC; GNT1185600), CERA Innovation Fund, National Natural Science Foundation of China (No. 82201227), and Natural Science Foundation of Guangdong Province, China (No. 2023A1515011225). Anai Gonzalez Cordero is an Al & Val Rosenstrauss Fellow (Rebecca Cooper Foundation). Funds supporting this work included the Medical Research Future Fund Stem Cell Mission (MRFF; MRF2008912), and the National Stem Cell Foundation - Metcalf Award Prize (No 00339). Satheesh Kumar was supported by the Melbourne Research Scholarship and Graeme Clark Institute Top-Up Scholarship. The Centre for Eye Research Australia receives Operational Infrastructure Support from the Victorian Government.

Author contributions

Satheesh Kumar: Conceptualization, Methodology, Formal analysis, Investigation, Data curation, Writing – original draft, Writing – review and editing. **Yi-Wen Hsiao**: Methodology,

Formal analysis, Investigation, Writing – review and editing. **Vickie H Y Wong**: Methodology,

Investigation. **Deborah Aubin**: Methodology, Investigation. **Jiang-Hui Wang**: Formal

analysis, Investigation. **Leszek Lisowski:** Resources. **Elizabeth Rakoczy:** Resources. **Fan Li**:

Resources. **Luis Alarcon-Martinez**: Methodology. **Anai Gonzalez-Cordero**: Methodology,

Investigation, Resources, Writing – review and editing, Funding acquisition. **Bang V. Bui**:

Methodology, Formal analysis, Investigation, Writing – review & editing. **Guei-Sheung Liu**:

Conceptualization, Methodology, Formal analysis, Resources, Data curation, Writing –

original draft, Writing – review and editing, Project administration, Funding acquisition,

Supervision.

Competing interests

The authors declare that they have no known competing financial interests.

Data and materials availability

RNA sequencing data generated from this study will be deposited in GEO and SRA. Other data

generated during and/or analysed during the current study will be available from the

corresponding author on reasonable request. Supporting data and source data can also be found

in the supplementary files.

REFERENCES

1. Wang JY, Doudna JA. CRISPR technology: A decade of genome editing is only the
beginning. *Science* **379**, eadd8643 (2023).

842
843 2. Kumar S*, et al.* RNA-targeting strategies as a platform for ocular gene therapy. *Prog*
844 *Retin Eye Res* **92**, 101110 (2023).

845
846 3. Hu Y*, et al.* Metagenomic discovery of novel CRISPR-Cas13 systems. *Cell Discovery*
847 **8**, 107 (2022).

848
849 4. Zhou C*, et al.* CasRx-mediated RNA targeting prevents choroidal neovascularization
850 in a mouse model of age-related macular degeneration. *Natl Sci Rev* **7**, 835-837 (2020).

851
852 5. Zhou H*, et al.* Glia-to-Neuron Conversion by CRISPR-CasRx Alleviates Symptoms of
853 Neurological Disease in Mice. *Cell* **181**, 590-603.e516 (2020).

854
855 6. Zheng Z*, et al.* Preventing autosomal-dominant hearing loss in Bth mice with
856 CRISPR/CasRx-based RNA editing. *Signal Transduction and Targeted Therapy* **7**, 79
857 (2022).

858
859 7. Powell JE*, et al.* Targeted gene silencing in the nervous system with CRISPR-Cas13.
860 *Science Advances* **8**, eabk2485 (2022).

861
862 8. Li Y*, et al.* Collateral cleavage of 28s rRNA by RfxCas13d causes death of mice.
863 *bioRxiv*, 2022.2001.2017.476700 (2022).

864
865 9. Shi P*, et al.* RNA-guided cell targeting with CRISPR/RfxCas13d collateral activity in
866 human cells. *bioRxiv*, 2021.2011.2030.470032 (2021).

867
868 10. Bot JF, van der Oost J, Geijsen N. The double life of CRISPR–Cas13. *Current Opinion*
869 *in Biotechnology* **78**, 102789 (2022).

870
871 11. Xu C*, et al.* Programmable RNA editing with compact CRISPR–Cas13 systems from
872 uncultivated microbes. *Nat Methods* **18**, 499–506 (2021).

873
874 12. Kannan S*, et al.* Compact RNA editors with small Cas13 proteins. *Nature*
875 *Biotechnology*, 1-4 (2021).

877   13.   Tong H, *et al.* High-fidelity Cas13 variants for targeted RNA degradation with minimal
878   collateral effects. *Nature Biotechnology*, (2022).

879
880   14.   Li G, *et al.* Mini-dCas13X–mediated RNA editing restores dystrophin expression in a
881   humanized mouse model of Duchenne muscular dystrophy. *The Journal of Clinical*
882   *Investigation* **133**, (2023).

883
884   15.   Xiao Q, *et al.* Rescue of autosomal dominant hearing loss by in vivo delivery of mini
885   dCas13X-derived RNA base editor. *Sci Transl Med* **14**, eabn0449 (2022).

886
887   16.   Deng X, *et al.* Structural basis for the activation of a compact CRISPR-Cas13 nuclease.
888   *Nature Communications* **14**, 5845 (2023).

889
890   17.   Lin F-L, *et al.* Gene Therapy Intervention in Neovascular Eye Disease: A Recent
891   Update. *Mol Ther* **28**, 2120-2138 (2020).

892
893   18.   Xu M, Fan R, Fan X, Shao Y, Li X. Progress and Challenges of Anti-VEGF Agents
894   and Their Sustained-Release Strategies for Retinal Angiogenesis. *Drug Des Devel Ther*
895   **16**, 3241-3262 (2022).

896
897   19.   Chuang Y-F, Wang P-Y, Kumar S, Lama S, Lin F-L, Liu G-S. Methods for in vitro
898   CRISPR/CasRx-Mediated RNA Editing. *Frontiers in Cell and Developmental Biology*
899   **9**, (2021).

900
901   20.   Li Z, *et al.* Intrinsic targeting of host RNA by Cas13 constrains its utility. *Nature*
902   *Biomedical Engineering*, (2023).

903
904   21.   Li Y, *et al.* The collateral activity of RfxCas13d can induce lethality in a RfxCas13d
905   knock-in mouse model. *Genome Biol* **24**, 20 (2023).

906
907   22.   Ofer K, Benjamin BL, Owen RSD, Aartjan JWtV, Cameron M. RNA structure
908   modulates   Cas13   activity   and   enables   mismatch   detection.   *bioRxiv*,
909   2023.2010.2005.560533 (2023).

910
911   23.   Lancaster MA, Knoblich JA. Organogenesis in a dish: Modeling development and
912   disease using organoid technologies. *Science* **345**, 1247125 (2014).

913
914   24.   Wahle P, *et al.* Multimodal spatiotemporal phenotyping of human retinal organoid
915   development. *Nature Biotechnology*, (2023).

917 25. Maryna VI, *et al.* PCDH15 Dual-AAV Gene Therapy for Deafness and Blindness in
918   Usher Syndrome Type 1F. *bioRxiv*, 2023.2011.2009.566447 (2023).

919
920 26. West EL*, et al.* Antioxidant and lipid supplementation improve the development of
921   photoreceptor outer segments in pluripotent stem cell-derived retinal organoids. *Stem*
922   *Cell Reports* **17**, 775-788 (2022).

923
924 27. Kim HJ*, et al.* Comprehensive characterization of fetal and mature retinal cell identity
925   to assess the fidelity of retinal organoids. *Stem Cell Reports* **18**, 175-189 (2023).

926
927 28. Hu H, Sun S-C. Ubiquitin signaling in immune responses. *Cell Research* **26**, 457-483
928   (2016).

929
930 29. Rakoczy EP*, et al.* Characterization of a mouse model of hyperglycemia and retinal
931   neovascularization. *Am J Pathol* **177**, 2659-2670 (2010).

932
933 30. Nakagawa R*, et al.* Structure and engineering of the minimal type VI CRISPR-
934   Cas13bt3. *Molecular Cell* **82**, 3178-3192.e3175 (2022).

935
936 31. Nishijima K*, et al.* Vascular Endothelial Growth Factor-A Is a Survival Factor for
937   Retinal Neurons and a Critical Neuroprotectant during the Adaptive Response to
938   Ischemic Injury. *Am J Pathol* **171**, 53-67 (2007).

939
940 32. Le Y-Z. VEGF production and signaling in Müller glia are critical to modulating
941   vascular function and neuronal integrity in diabetic retinopathy and hypoxic retinal
942   vascular diseases. *Vision Research* **139**, 108-114 (2017).

943
944 33. Ford KM, Saint-Geniez M, Walshe T, Zahr A, D'Amore PA. Expression and Role of
945   VEGF in the Adult Retinal Pigment Epithelium. *Investigative Ophthalmology & Visual*
946   *Science* **52**, 9478-9487 (2011).

947
948 34. Colognori D, Trinidad M, Doudna JA. Precise transcript targeting by CRISPR-Csm
949   complexes. *Nature Biotechnology* **41**, 1256-1264 (2023).

950
951 35. Jo N*, et al.* Inhibition of Platelet-Derived Growth Factor B Signaling Enhances the
952   Efficacy of Anti-Vascular Endothelial Growth Factor Therapy in Multiple Models of
953   Ocular Neovascularization. *Am J Pathol* **168**, 2036-2053 (2006).

954
955 36. Park J*, et al.* CRISPR/Cas9 mediated specific ablation of vegfa in retinal pigment
956   epithelium efficiently regresses choroidal neovascularization. *Sci Rep* **13**, 3715 (2023).

958 37.   Garita-Hernandez M, *et al.* AAV-Mediated Gene Delivery to 3D Retinal Organoids
959   Derived from Human Induced Pluripotent Stem Cells. *Int J Mol Sci* **21**, (2020).

960
961 38.   McClements ME, *et al.* Tropism of AAV Vectors in Photoreceptor-Like Cells of
962   Human iPSC-Derived Retinal Organoids. *Translational Vision Science & Technology*
963   **11**, 3-3 (2022).

964
965 39.   Duong TT, *et al.* Comparative AAV-eGFP Transgene Expression Using Vector
966   Serotypes 1–9, 7m8, and 8b in Human Pluripotent Stem Cells, RPEs, and Human and
967   Rat Cortical Neurons. *Stem Cells International* **2019**, 7281912 (2019).

968
969 40.   Eoi Jong S, Jeong AC, Jae-Young K, Young Hee Y. Aflibercept ameliorates retinal
970   pericyte loss and restores perfusion in streptozotocin-induced diabetic mice. *BMJ Open*
971   *Diabetes Research & Care* **8**, e001278 (2020).

972
973 41.   Tokunaga CC, *et al.* Effects of Anti-VEGF Treatment on the Recovery of the
974   Developing Retina Following Oxygen-Induced Retinopathy. *Investigative*
975   *Ophthalmology & Visual Science* **55**, 1884-1892 (2014).

976
977 42.   Shah CP, Garg SJ, Vander JF, Brown GC, Kaiser RS, Haller JA. Outcomes and Risk
978   Factors Associated with Endophthalmitis after Intravitreal Injection of Anti-Vascular
979   Endothelial Growth Factor Agents. *Ophthalmology* **118**, 2028-2034 (2011).

980
981 43.   Lai AKW, Lo ACY. Animal Models of Diabetic Retinopathy: Summary and
982   Comparison. *Journal of Diabetes Research* **2013**, 106594 (2013).

983
984 44.   Smith LEH, *et al.* Regulation of vascular endothelial growth factor-dependent retinal
985   neovascularization by insulin-like growth factor-1 receptor. *Nat Med* **5**, 1390-1395
986   (1999).

987
988 45.   Huang X, *et al.* Genome editing abrogates angiogenesis in vivo. *Nature*
989   *Communications* **8**, 112 (2017).

990
991 46.   Holmgaard A, *et al.* In Vivo Knockout of the Vegfa Gene by Lentiviral Delivery of
992   CRISPR/Cas9 in Mouse Retinal Pigment Epithelium Cells. *Molecular Therapy -*
993   *Nucleic Acids* **9**, 89-99 (2017).

994
995 47.   Holmgaard AB, *et al.* Targeted Knockout of the Vegfa Gene in the Retina
996   by Subretinal Injection of RNP Complexes Containing Cas9 Protein and Modified
997   sgRNAs. *Mol Ther* **29**, 191-207 (2021).

48. Wang Y*, et al.* Efficient Human Genome Editing Using SaCas9 Ribonucleoprotein Complexes. *Biotechnology journal* **14**, e1800689 (2019).

49. Chung SH*, et al.* CRISPR-based VEGF suppression using paired guide RNAs for treatment of choroidal neovascularization. *Molecular Therapy - Nucleic Acids* **28**, 613-622 (2022).

50. Konermann S, Lotfy P, Brideau NJ, Oki J, Shokhirev MN, Hsu PD. Transcriptome Engineering with RNA-Targeting Type VI-D CRISPR Effectors. *Cell* **173**, 665-676.e614 (2018).

51. Lai C-M. Generation of transgenic mice with mild and severe retinal neovascularisation. *Br J Ophthalmol* **89**, 911-916 (2005).

52. Zudaire E, Gambardella L, Kurcz C, Vermeren S. A Computational Tool for Quantitative Analysis of Vascular Networks. *PLoS One* **6**, e27385 (2011).

53. Andrews S. FastQC: a quality control tool for high throughput sequence data.). Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom (2010).

54. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

55. Dobin A*, et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

56. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC bioinformatics* **12**, 1-16 (2011).

57. Hung J-H, Yang T-H, Hu Z, Weng Z, DeLisi C. Gene set enrichment analysis: performance evaluation and usage guidelines. *Briefings in bioinformatics* **13**, 281-291 (2012).

58. Consortium GO. The Gene Ontology (GO) database and informatics resource. *Nucleic acids research* **32**, D258-D261 (2004).

59. Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics: a journal of integrative biology* **16**, 284-287 (2012).

1039 60.   Gordon A, Hannon GJ. Fastx-toolkit. *FASTQ/A short-reads preprocessing tools*
1040   *(unpublished) http://hannonlab cshl edu/fastx_toolkit* **5**, (2010).

1041

1042 61.   Ye J, McGinnis S, Madden TL. BLAST: improvements for better sequence analysis.
1043   *Nucleic acids research* **34**, W6-W9 (2006).

1044

1045 62.   Ianevski A, Giri AK, Aittokallio T. Fully-automated and ultra-fast cell-type
1046   identification using specific marker combinations from single-cell transcriptomic data.
1047   *Nature communications* **13**, 1246 (2022).

1048

1049 63.   Limb GA, Salt TE, Munro PM, Moss SE, Khaw PT. In vitro characterization of a
1050   spontaneously immortalized human Müller cell line (MIO-M1). *Invest Ophthalmol Vis*
1051   *Sci* **43**, 864-869 (2002).

1052

1053 64.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
1054   RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1055

1056 65.   Hao Y*, et al.* Dictionary learning for integrative, multimodal and scalable single-cell
1057   analysis. *Nature Biotechnology*, (2023).

1058

1059

1060

 Key Resources Table

Reagent or Resource	Source	Identifier
Chemicals		
DMEM	Thermo Fisher Scientific	Cat# 11965092
DMEM/F12	Thermo Fisher Scientific	Cat# 11320033
Opti-MEM	Thermo Fisher Scientific	Cat# 31985070
TrypLE™ Express	Thermo Fisher Scientific	Cat# 12605093
GlutaMAX™ Supplement	Thermo Fisher Scientific	Cat# 35050061
FBS	Bovogen Biologicals	SFBS-AU
Antibiotic-Antimycotic	Thermo Fisher Scientific	Cat# 15240062
DPBS	Thermo Fisher Scientific	Cat# 14190250
Lipofectamine™ 3000 Transfection Reagent	Thermo Fisher Scientific	Cat# L3000001
ViaFect™ Transfection Reagent	Promega Corporation	Cat# E4982
NEB® Stable Competent E. coli, High Efficiency	New England Biolabs	Cat# C3040I
rCutSmart™ Buffer	New England Biolabs	Cat# B6004S
NEB® 10-beta/Stable Outgrowth Medium	New England Biolabs	Cat# B9035S
NucBlue™ Live ReadyProbes™ Reagent (Hoechst 33342)	Thermo Fisher Scientific	Cat# R37605
LB (Lennox L) agar	Thermo Fisher Scientific	Cat# 22700025
2-YT Broth	Thermo Fisher Scientific	Cat# 22712020

Carbenicillin Disodium Salt	Thermo Fisher Scientific	Cat# 10177012
GelRed® nucleic acid gel stain	Biotium	Cat# 41003
Phosphate buffered saline	Merck	Cat# P3813
Tween-20	Merck	Cat# P1379
Triton-X-100	Amresco	Cat# 0694
RIPA Lysis and Extraction Buffer	Thermo Fisher Scientific	Cat# 89900
Pierce™ Phosphatase Inhibitor Mini Tablets	Thermo Fisher Scientific	Cat# A32957
NuPAGE™ MES SDS Running Buffer (20X)	Thermo Fisher Scientific	Cat# NP0002
NuPAGE™ LDS Sample Buffer (4X)	Thermo Fisher Scientific	Cat# NP0007
0.5% Proxymethacaine hydrochloride (Alcaine™)	Alcon	N/A
0.5% Tropicamide (Mydriacyl)	Alcon	N/A
GenTeal™ Gel	Alcon	N/A
Ketamine, 100 mg/mL	Troy Laboratories	Ketamil
Xylazine, 100 mg/mL	Troy Laboratories	Ilium Xylazil-100
Experimental models: Cell lines		
Human: HEK293FT	Thermo Fisher Scientific	R70007
Human: ARPE-19	ATCC	CRL-2302™
Human: Müller 1 (MIO-M1)	UCL Institute of Ophthalmology, London	[63]
Experimental models: Organisms/ strains		

Human: 3D retinal organoid	Children's Medical Research Institute	[27]
Kimba Mouse	The University of Western Australia	[29]
Antibodies		
Mouse anti-actin, clone C4	Merck	MAB1501
Rabbit anti-HA, C29F4	Cell Signaling Technology	#3724
Mouse anti-rhodopsin, 1D4	Abcam	ab5417
Rat mCherry monoclonal 16D7, Alexa Fluor™ 594	Thermo Fisher Scientific	Cat# M11240
Rat anti-laminin monoclonal, α-2 chain	Merck	L0663
Donkey anti-rat AffiniPure Fragment IgG (H + L), Alexa Fluor™ 647	Jackson ImmunoResearch	712-606-150
Donkey mouse AffiniPure Fragment IgG (H + L), Alexa Fluor™	Jackson ImmunoResearch	715-546-151
Goat anti-rabbit IgG (H+L) Secondary Antibody, HRP	Thermo Fisher Scientific	Cat# 31460
Goat anti-Mouse IgG (H+L) Secondary Antibody, HRP	Thermo Fisher Scientific	Cat# 62-6520
Anti-Arrestin C antibody, C-terminal	Abcam	189437
Monoclonal Anti-Opsin antibody produced in mouse	Merck	O4886

Sheep anti Human chx 10 (Visual system homeobox 2) (NT)	ExAlpha	XII80P
CRALBP Monoclonal Antibody (B2)	Thermo Fisher Scientific	MA1-813
Anti-Prox 1 Antibody	Merck	AB5475
HuC/HuD Monoclonal Antibody (16A11)	Thermo Fisher Scientific	A21271
Commercial kits		
RNeasy Mini kit	Qiagen	Cat# 74106
Monarch® Total RNA Miniprep Kit	New England Biolabs	Cat# T2010S
High-Capacity cDNA Reverse Transcription Kit	Thermo Fisher Scientific	Cat# 4368814
Q5® Hot Start High-Fidelity Master Mix	New England Biolabs	Cat# M0494S
TaqMan™ Fast Advanced Master Mix	Thermo Fisher Scientific	Cat# 4444557
SYBR™ Green PCR Master Mix	Thermo Fisher Scientific	Cat# 4309155
Human VEGF DuoSet ELISA	R & D systems	Cat# DY293B
Recombinant DNA		
pAAV.CMV-CasRx mCh	Addgene	203442
pAAV.CMV-CasRx-VEGFA sgRNA	Addgene	203443
pAAV.CMV-CasRx-VEGFA sgRNA array	Addgene	203444
pAAV.CMV-Cas13bt3-NT sgRNA	Addgene	203446

pAAV.CMV-Cas13bt3-VEGFA sgRNA	Addgene	203447
pAAV.CMV-Cas13bt3-VEGFA sgRNA array	Addgene	203448
pAAV.CMV-Cas13bt3 (GenScript CO)-NT sgRNA	Addgene	203451
pAAV.CMV-Cas13bt3 (GenScript CO)-VEGFA sgRNA	Addgene	203452

Software and algorithms

GraphPad Prism 10	GraphPad	https://www.graphpad.com/scientific-software/prism/E/
Benchling	Benchling	https://www.benchling.com/
Fiji-ImageJ	National Institutes of Health	https://imagej.net/software/fiji/
GenSmart™ Codon Optimisation	GenScript	https://www.genscript.com/gensmart-free-gene-codon-optimization.html
IDT Codon Optimisation Tool	Integrated DNA Technologies	https://sg.idtdna.com/pages/tools/codon-optimization-tool
STAR (2.7.11a)	[55]	https://github.com/alexdobin/STAR
DESeq2 (v3.18)	[64]	https://bioconductor.org/packages/release

		/bioc/html/DESeq2.html
Seurat (v5)	[65]	https://satijalab.org/seurat/
Cell Ranger (v7.2)	10X Genomics	https://www.10xgenomics.com/support/software/cell-ranger
ImageLab (v6.1)	Bio-Rad	https://www.bio-rad.com/en-au/product/image-lab-software
R (v4.3.2)	R Core Team	https://cran.r-project.org/
RStudio (v2023.09.1)	Posit PBC	https://posit.co/download/rstudio-desktop/
Adobe Photoshop (v25)	Adobe Inc.	https://www.adobe.com/au/products/photoshop.html
AngioTool (v0.5)	National Cancer Institute	https://ccrod.cancer.gov/confluence/display/ROB2/Downloads
Equipment		
HRA + OCT Spectralis	Heidelberg Engineering	N/A
Leica Stellaris 5 confocal microscope	Leica Microsystems	N/A

1062

1063 **Primers used in this study.**

Quantitative PCR

Name	Source	Identifier
Human VEGFA (Hs00900055_m1)	Thermo Fisher Scientific	Cat# 4331182
Mouse VEGFA (Mm00437306_m1)	Thermo Fisher Scientific	Cat# 4331182
Human GAPDH (Hs02786624_g1)	Thermo Fisher Scientific	Cat# 4448489
Mouse GAPDH (Mm99999915_g1)	Thermo Fisher Scientific	Cat# 4448489

Standard PCR

Name	Sequence (5' - 3')	Source
Cas13bt3_FWD	CGGCGAACAGGGTGATAAA	Integrated DNA Technologies
Cas13bt3_REV	CTGTGCTGGGCTTCAAGATA	Integrated DNA Technologies

1064

1065

1066

1067

1068

1069

1070 **SUPPLEMENTARY DATA**

1071

1072 **Supplementary figure 1. Evaluation of Cas13bt3 targeting in retinal cell lines.** VEGFA
1073 mRNA expression in ARPE19 cells following transfection of control, Cas13bt3-NT and
1074 Cas13bt3-VEGFA. A significant reduction in Cas13bt3-VEGFA was observed. Data presented
1075 as mean ± SEM. ns: not significant. *P≤0.05.

1076

1077

Supplementary figure 2. HEK293FT DEG and off target analysis. (A) Number of differentially expressed genes from the VEGFA-targeting shRNA, CasRx and Cas13bt3 transfections in HEK293FT cells compared to vector control. **(B)** Volcano plots of differentially expressed genes from VEGFA shRNA, **(C)** CasRx VEGFA sgRNA, **(D)** Cas13bt3 NT sgRNA and **(E)** Cas13bt3 VEGFA sgRNA compared to vector control. **(F)** Plot of TPM values for VEGFA and candidate off-target genes from Cas13bt3-NT sgRNA and Cas13bt3 VEGFA sgRNA groups compared to shRNA and CasRx targeting vectors. Statistical analysis was conducted by Welch's t test **(F)**. Data presented as mean ± SEM. ns: not significant. *P≤0.05, ****P≤0.00001.

1087

1088 **Supplementary figure 3. Prediction of Cas13bt3 cut sites**. **(A)** Schematic of Cas13bt3-gRNA and Cas13bt3-gRNA-target complex. **(B)** Potential
1089 Cas13bt3 cut sites on VEGFA mRNA target for VEGFA sgRNA1, 2 and 3. **(C)** Potential Cas13bt3 cut sites on top 10 candidate off-target genes
1090 for VEFGA sgRNA 1.

A

```
VEGFA      T A T G T G C T G G C C T T G G T G A G G T T T G A T C C G
PRKAR2B    T A T G T G C T G G C C T T G G - - - - - - - - - - - - - -
EHD4       - - - - - - - - - - - - - G G T G A G G T T T G A T C C -
USP22      - - - - T G C T G G C C T T G G T G A - - - - - - - - - - -
RHBDL1     - - - G T G C T G G C C T T G G T G - - - - - - - - - - -
POMT2      - - - - T G C T G G C C T T G G T G A - - - - - - - - - - -
SNX33      - - - - - - - - - - - T G G T G A G G T T T G A T - - -
RER1       - - - G T G C T G G C C T T G G T - - - - - - - - - - - -
POLA1      - - - - - - - - - - - - - - - G T G A G G T T T G A T C C -
SLC9B2     - - - G T G C T G G C C T T G G T - - - - - - - - - - - -
SYT10      - - - - - - - - - - C T T G G T G A G G T T T G - - - - -
```

B

```
VEGFA presgRNA 2   C A C T T C G T G A T G A T T C T G C C C T C C T C C T T C
BARX1              - - - - - - - - - - - - - T C T G C C C T C C T C C T T C
SLX4               - - - - - - - - - - - - - T C T G C C C T C C T C C T T C
LOC100192426       - - - - - - - - - - - - - T C T G C C C T C C T C C T T C
NRG1               - - - - - - - - G A T G A T T C T G C C C T C - - - - - - -
AGRN               - - - - - - - - - - - A T T C T G C C C T C C T C C - - -
TRIM56             - - - - - - - - - - - - - - C T G C C C T C C T C C T T C
CDH20              - - - - - - - - - - G A T T C T G C C C T C C T C C T T C
MORC3              - - - - - - - - - T G A T T C T G C C C T C C T - - - - -
ZNF32-AS3          - - - - - - - - - - - - - - C T G C C C T C C T C C T T C
SUGT1P4-STRA6LP    C A C T T C G T G A T G A T - - - - - - - - - - - - - - - -
```

C

```
VEGFA presgRNA 3   T C T G C A T T C A C A T T T G T T G T G C T G T A G G A A
NUP210L            - - - - - - - - - C A T T T G T T G T G C T G T - - - - -
LOC105373744       T C T G C A T T C A C A T T T - - - - - - - - - - - - - -
LYSMD3             - - - - - - - - - - C A T T T G T T G T G C T G T - - - - -
SKP1               - C T G C A T T C A C A T T T G - - - - - - - - - - - - -
CPA4               - - T G C A T T C A C A T T T G - - - - - - - - - - - - -
SENP5              - C T G C A T T C A C A T T T G T T - - - - - - - - - - -
```

Supplementary figure 4. Multiplexed VEGFA knockdown using Cas13bt3. (A) Top off-target genes for VEGFA sgRNA 1, **(B)** VEGFA sgRNA 2 and **(C)** VEGFA sgRNA 3.

1094

Supplementary Figure 5. Codon optimisation of Cas13bt3. (A) Schematic of codon optimisation procedure from Cas13e protein sequence. **(B)** VEGFA mRNA knockdown from original and codon optimised Cas13e compared to control transfection. **(C and D)** Expression of original and codon optimised determined from Western blot. Statistical analyses was conducted using One-way ANOVA with Tukey's multiple comparisons test (**B and C**). Data presented as mean ± SEM. *P≤0.05, **P≤0.001, ****P≤0.00001.

1101
1102
1103
Supplementary figure 6. Characterization of 30 weeks retinal organoids. (A) Retinal
organoids showed expression of mature photoreceptor markers Arrestin-3 (magenta) specific
to the cone cells and Rhodopsin (cyan) localized to the outer segments (OS) of rod
photoreceptor cells. **(B, C)** In the inner nuclear layer (INL) bipolar cells stained with VSX2
(cyan) and Müller glia stained with CRALBP (magenta) **(B)**, Amacrine and horizontal cells
stained with PROX1 (magenta) and ganglion cells stained with HuCHuD (cyan) **(C)**. **(D)**
Brightfield image of retinal organoid section showing the outer nuclear layer (ONL) and a
pigmented RPE island next to it. Scale bar: 116μm. Abbreviations: INL: inner nuclear layer
ONL: outer Nuclear Layer, OS: outer segment, RPE: retinal pigment epithelium.

1113

Supplementary figure 7. Dot plots of retinal cell marker genes for respective cell types in 3D retinal organoids. Complete list of cell marker genes can obtained from Kim et al., 2023 [25].

1116

Supplementary figure 8. *Expression of VEGFA *on* retinal cell clusters. UMAP plots of

showing expression on **(A)** Untreated retinal organoid. **(B)** Violin plots showing expression of

VEGFA in individual retinal cell types. **(C)** UMAP plots of retinal cell clusters with expression

of VEGFA from VEGFA-treated and NT-treated retinal organoids, **(C)** all cells, **(D)** Cas13bt3

positive cells and **(E)** Cas13bt3 negative cells. Data analyses were conducted using One-way

ANOVA with Tukey's multiple comparisons test (B). **P$\leq$0.001, ****P$\leq$0.00001.

1123

Supplementary figure 9. Differentially expressed genes in Cas13bt3-VEGFA against Cas13bt3-NT. Volcano plots from scRNA-seq for each retinal cell type showing differentially expressed genes between retinal organoid infected with Cas13bt3-VEGFA sgRNA and

1127 Cas13bt3-NT sgRNA. Significant downregulation of VEGFA in RPE cells is highlighted in
1128 red.

1129

1130 **Supplementary figure 10. Cas13bt3-NT sgRNA treatment in retinal organoids. (A)**
1131 Expression of Cas13bt3 by number of reads across retinal cell types in NT sgRNA treatment
1132 group. **(B)** Violin plots of VEGFA expression in cells with low and high expression of
1133 Cas13bt3 against Cas13bt3 negative cells showed no significant differences.

1134

Supplementary figure 11. Expression of Cas13bt3 plotted against expression of *VEGFA* mRNA across cell types in NT sgRNA and VEGFA sgRNA groups.

1139 **Supplementary figure 12. Correlation plots of Cas13bt3 with VEGFA expression in the various retinal cell types for the predicted off-**
1140 **target genes of VEGFA sgRNA.**

Supplementary figure 13. Intravitreal delivery of AAV2.7m8 achieves strong transduction of mice photoreceptors. (A) Schematic of AAV2.7m8 virus carrying mCherry reporter. **(B)** Schematic of experimental procedure for intravitreal injection of C57J/BL6 mice eyes. **(C)** Confocal microscope images of retinal flatmount 4 weeks post intravitreal injection, showing transduction of ganglion, bipolar, cone and rod cells with reference image of retinal cell layers.

Supplementary Figure 14. Cas13bt3 off-target effects in Kimba mice. **(A)** Schematic of candidate off-target genes to VEGFA sgRNA in mouse transcriptome. Volcano plots of differentially expressed genes from Kimba mice retinas treated with AAV2.7m8 carrying **(B)**

Cas13bt3-NTsgRNA and **(C)** Cas13bt3-VEGFAsgRNA compared to vector controls. **(D)** Expression in TPM of mouse VEGFA and candidate off-target genes from RNA-seq.

Supplementary figure 15. Original fundus fluorescein angiography images from Kimba mice retinae.

Supplementary figure 16. Imaging of retinal vascular plexuses Kimba mice treated with Cas13bt3-NTsgRNA and Cas13bt3-VEGFAsgRNA. (A) Schematic of experimental procedure for imaging retinal vascular plexuses. **(B)** Schematic of the different retinal plexuses. **(C)** Confocal imaging of the retinal plexuses, 8 weeks after Cas13bt3 treatment, immunolabelled with laminin. Magnification 40X, scale bar: 50µm. White arrows indicate leaky vessel structure.

Supplementary figure 17. Retinal thickness readings from optical coherence tomography (OCT) in Kimba mice. (A) Early Treatment of Diabetic Retinopathy Study (ETDRS) grid showing segments of the retina analysed using OCT. C:Central (optic nerve); N1: Inner nasal; N2: Outer Nasal; T1: Inner temporal; T2: Outer temporal; S1: inner superior; S2: Outer superior; I1: Inner inferior; I2: Outer inferior. **(B)** Vessel density and **(C)** number of junctions readings from FFA images of Kimba mice at baseline compared to wildtype (C57BL/6) mice. **(D)** Plot of difference in total retinal thickness in untreated and treated Kimba mice between baseline and endpoint. **(E)** Representative OCT images from untreated and treated Kimba mice at endpoint.