1   Title page (Original Research Article)

3   Characterisation of RNA editing and gene therapy with a compact CRISPR-Cas13 in the retina

Satheesh Kumar[1,2,3], Yi-Wen Hsiao[1], Vickie H Y Wong[4], Deborah Aubin[5,6], Jiang-Hui Wang[1],

Leszek Lisowski[7,8,9], Elizabeth P Rakoczy[10], Fan Li[11], Luis Alarcon-Martinez[1,2], Anai

Gonzalez-Cordero[5,6], Bang V Bui[4], Guei-Sheung Liu[1,2,3*]

[1]Centre for Eye Research Australia, Royal Victorian Eye and Ear Hospital, East Melbourne,

VIC, Australia.

[2]Ophthalmology, Department of Surgery, University of Melbourne, East Melbourne, VIC,

Australia.

[3]Menzies Institute for Medical Research, University of Tasmania, Hobart, TAS, Australia.

[4]Department of Optometry and Vision Sciences, University of Melbourne, Parkville, VIC,

Australia.

[5]School of Medical Sciences, Faculty of Medicine and Health, The University of Sydney,

Camperdown, NSW, Australia.

[6]Stem Cell Medicine Group, Children's Medical Research Institute, Westmead, NSW,

Australia.

[7]Translational Vectorology Research Unit, Children's Medical Research Institute, Faculty of

Medicine and Health, The University of Sydney, Westmead, NSW, Australia.

[8]Military Institute of Medicine, Laboratory of Molecular Oncology and Innovative Therapies,

Warsaw, Poland.

[9]Australian Genome Therapeutics Centre, Children's Medical Research Institute and Sydney

Children's Hospitals Network, Westmead, NSW, Australia.

[10]Centre for Ophthalmology and Visual Sciences, University of Western Australia, Perth, WA,

Australia.

[11]Eye Center, Zhongshan City People's Hospital, Zhongshan, Guangdong Province, China.

*Correspondence and requests for materials should be addressed to

Dr Guei-Sheung Liu (rickliu0817@gmail.com). Centre for Eye Research Australia. Address:

Level 7, 32 Gisborne Street, East Melbourne, VIC 3002, Australia. Tel: +61399298360.

Abstract

CRISPR-Cas13 nucleases are programmable RNA-targeting effectors that can silence gene
expression in a reversible manner. Recent iterations of Cas13 nucleases are compact for adeno-
associated virus (AAV) delivery to achieve strong and persistent expression in various organs
in a safe manner. Here, we report significant transcriptomic signatures of Cas13bt3 expression
and show all-in-one AAV gene therapy with Cas13bt3 can effectively silence *VEGFA* in human
retinal organoids and humanised mouse models. Specifically, human embryonic stem cells
(hESC)-derived retinal pigment epithelium cells show high expression of Cas13bt3 from virus
delivery corresponding to significant reduction of *VEGFA* mRNA. We further show that
intravitreal delivery of Cas13bt3 can transduce mouse retinal cells efficiently, reaching the
photoreceptors for specific knockdown of human *VEGFA* in the trVEGF029 (Kimba) mouse.
Our results reveal important considerations for assessing Cas13 activity and establishing
Cas13bt3 as a potential anti-VEGF agent that can achieve long-term control of VEGFA for
treatment of diabetic retinopathy and wet age-related macular degeneration.

Introduction

The emergence of Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) and CRISPR-associated (Cas) enzymes, known as CRISPR-Cas, has fuelled several gene editing efforts for novel therapeutics in the past decade[1]. Developed by repurposing the bacterial immune defence against phage viruses, CRISPR-Cas harnesses the ability to induce targeted changes to DNA or RNA sequences. Among CRISPR-Cas proteins, CRISPR-Cas13 is of particular interest for its exclusive specificity for RNA. Targeting RNA is especially attractive as it enables control of gene expression without perturbing the genome and introducing permanent changes and is therefore an emerging avenue for therapeutic development[2].

Eleven variants of CRISPR-Cas13 have been described to date[3]. Among them, Cas13d (commonly referred to as CasRx) described in 2018 garnered special attention for its compact size of 930aa, making it the first Cas13 enzyme amenable to adeno-associated virus (AAV) delivery. Studies investigating the development of RNA-targeting therapeutics have widely used CasRx to demonstrate preclinical efficacy against disorders affecting vision, hearing and brain function[4-7], however significant unintended editing have been reported, hampering its clinical progress[8, 9], and the exact specificity profile of CasRx remains contested[10]. More recently, hypercompact Cas13bt3 (also reported as Cas13X) enzymes were described, ranging even smaller at 775aa[11, 12]. The newfound enzyme has been reported to possess high specificity, and its small size enables additional applications such as RNA base editing using a single AAV[13-15]. The therapeutic potential of Cas13bt3 is yet to be demonstrated, however initial *in vitro* studies show promising efficiency and safety[13, 16].

As a leading cause of blindness in the elderly population affecting 200 million people globally, ocular neovascularisation refers to a group of conditions characterised by the abnormal growth of leaky blood vessels in the eye. Vascular endothelial growth factor (VEGF) has been described as a key mediator of this process, and the overexpression of *VEGFA* underlies pathological angiogenesis and neovascularisation in common ocular diseases like diabetic retinopathy (DR) and neovascular age-related macular degeneration (nAMD). Regular (monthly/ bimonthly) administration of anti-VEGF drugs is the approved front-line treatment modality for these patients[17]. Delivered through intravitreal injections, frequent administration of anti-VEGF drugs poses a significant clinical burden on healthcare institutions

and patients due to risk of infection, patient anxiety, clinic capacity and availability of trained healthcare personnel among others(*18*).

We have previously described a potential alternative therapy to current anti-VEGF injections using an all-in-one AAV CRISPR-CasRx gene therapy for the treatment of ocular neovascularisation. This approach was validated *in vitro* in HEK293FT and MIO-M1 cells, demonstrating more than 90% and 50% *VEGFA* mRNA knockdown, respectively(*19*). Here, we utilize the novel Cas13bt3 to demonstrate potent *VEGFA* knockdown *in vitro* in human cell lines, in 3D human embryonic stem cells (hESC)-derived retinal organoids and *in vivo* in transgenic humanised mouse retinas. Promisingly, our results show significant knockdown of *VEGFA* mRNA in human cell lines without impact on predicted off-target genes. Single cell RNA sequencing (scRNA-seq) was used to characterise Cas13bt3-mediated *VEGFA* mRNA knockdown at single cell resolution in human retinal organoids identifying, for the first time, the overall transcriptomic impact and cell-specific activity of the enzyme. Lastly, *in vivo* gene therapy showed robust *VEGFA* mRNA silencing in mouse models of proliferative diabetic retinopathy along with control of neovascularisation symptoms. These findings support the development of Cas13-mediated ocular gene therapy establishing the clinical potential of Cas13bt3 and offering an alternative treatment to current anti-VEGF treatments.

Results

CRISPR-Cas13bt3 can knock down VEGFA mRNA in mammalian cells with minimal unintended targeting activity

We used our previously described single guide RNAs (sgRNA) targeting Exon 4 of *VEGFA* (*19*), to develop our new Cas13bt3 constructs and assess RNA cutting efficiency as well as collateral and off target cleavages (**Figure 1A-D**). Collateral activity is the trans-cleavage by Cas13 enzymes of adjacent mRNA transcripts as the result of an activated nuclease domain (**Figure 1B**). Off-target activity, on the other hand, is the targeting of other non-target mRNA transcripts due to unspecific binding of the sgRNA (**Figure 1C**).

We sought to compare and validate efficiency of the new Cas13bt3 enzyme by testing RNA cutting against two previously established RNA-targeting methods, RNA interference (short hairpin RNA, shRNA) and CasRx (a variant of Cas13d). These are widely used for therapeutic

and diagnostic applications. The final all-in-one AAV plasmids carrying either *VEGFA*-targeting shRNA or precursor single guide RNA (sgRNA hereafter) with either CasRx or Cas13bt3 (**Figure 1E**) were transfected into HEK293FT cells. Here, to further enhance the expression of Cas13bt3 we incorporated the stronger cytomegalovirus (CMV) promoter instead of the previously used elongation factor 1α short (EFS) promoter.

Significant *VEGFA* mRNA knockdown was observed from all three RNA-targeting methods, with up to 70% shRNA *VEGFA* mRNA knockdown ($P < 0.001$), while CasRx and Cas13bt3 achieved up to 90% knockdown ($P < 0.0001$) compared to vector controls (AAV plasmid carrying a fluorescent marker) (**Figure 1F**, n = 4 independent experiments). Next, we sought to determine if the reduction in mRNA would also lead to a decrease in protein levels. Interestingly, ELISAs for VEGFA revealed that only the Cas13bt3 treatment led to significant reduction of VEGFA protein in the cell supernatant (**Figure 1G**, P< 0.05; n = 4 independent experiments). As both Cas13 variants proved potent in silencing *VEGFA* mRNA in HEK293FT cells, we next transfected immortalised human retinal pigment epithelium cells (ARPE19 cell line) with the *VEGFA*-targeting CasRx and Cas13bt3 plasmids and again observed up to 90% knockdown ($P < 0.05$, n = 3 independent experiments) with Cas13bt3 against vector controls (**Figure S1**), suggesting that Cas13bt3 is a promising candidate for RNA silencing of retinal cells.

To further understand transcriptome-wide effects, we performed bulk RNA sequencing (RNA-seq) on HEK293FT cells transfected with shRNA, CasRx and Cas13bt3. RNA-seq was also performed on cells treated with either control, Cas13bt3 only, Cas13bt3NT and Cas13bt3VEGFA constructs to identify collateral and off-target activity of Cas13. Differentially expressed genes (DEGs) were similar for all treated groups, shRNA, CasRx and Cas13bt3 (**Figure S2A-C**), indicating that collateral activity from the new Cas13bt3 is comparable to previously established RNA-targeting methods. Interestingly, the non-targeting (NT) sgRNA group exhibited the highest number of DEGs, a finding that corresponds with recent studies (**Figures 1G, H and S2A-D**) (*16, 20*). Volcano plots of Cas13bt3 samples against control demonstrated and confirmed a significant downregulation of *VEGFA* mRNA only on cells that received the VEGFA targeting treatment (**Figure 1I**).

We identified potential off-target genes through NCBI BLAST of our VEGFA sgRNA sequence and studied the expression levels of the top 10 candidate off-target genes (**Figure

S3A). None of the off-target genes were significantly downregulated, indicating lack of off-target activity from Cas13bt3 (**Figure 1J**), as was the case for shRNA and CasRx (**Figure S2F**).

To further examine the varying DEG levels between the NT and targeting sgRNAs, we identified the top significantly enriched gene ontology (GO) terms for the Cas13bt3-NT sgRNA and Cas13bt3-VEGFA sgRNA groups. Top 5 significant GO terms for the Cas13bt3-NT sgRNA group were all positive enrichments and specifically showed upregulation of RNA binding processes, such as "*pre-mRNA binding*" and "*mRNA 3' UTR binding*", possibly explaining the elevated levels of DEGs in this group. On the other hand, with a VEGFA sgRNA, the top 5 GO terms were all negative enrichments and particularly showed downregulation of "*platelet-derived growth factor (PDGF) binding*" and "*PDGF receptor binding*", both angiogenesis promoting factors (**Figure 1K**). A recent study showed CasRx to have intrinsic targets that affected essential processes like RNA metabolism and mitotic cell cycle (*20*) and opined that this could underlie the lethality seen upon CasRx expression in a mouse model (*21*). Here, we found laminin-related processes (e.g., "*laminin binding*", "*laminin complex*") were strongly downregulated across all groups (Cas13bt3 only, Cas13bt3-NT sgRNA and Cas13bt3-VEGFA sgRNA) suggesting intrinsic targets could exist for Cas13bt3 as well (**Data S1**).

While Cas13bt3 may undergo activation after maturation of a NT sgRNA; without adequate homologous pairing, the catalytic pocket cannot be formed for efficient nuclease activity (*22*). On the other hand, with a targeting gRNA, Cas13bt3 would be activated to form the Cas13bt3-gRNA-target complex for targeted knockdown. Our results suggest that with a NT sgRNA, Cas13bt3 initiates an unspecific target-search process involving significant RNA binding that greatly impacts the expression levels of several genes, while a targeting sgRNA narrows the target-search process to reduce overall transcriptomic impact. Notably, despite high levels of overall DEGs, the number of downregulated genes remained similar between the NT sgRNA and VEGFA sgRNA groups suggesting that the NT sgRNA does not induce formation of the catalytic pocket for RNA knockdown, and altered gene expression is predominantly a result of increased RNA binding.

Fig. 1. Comparison of RNA-targeting tools for in vitro testing of VEGFA mRNA
silencing. (A) Schematic of CRISPR-Cas13 mechanism of action. **(B)** VEGFA sgRNA
sequence and targeting region on VEGFA mRNA sequence. **(C)** Schematic of collateral and
(D) off-target cleavage. **(E)** Schematic for domain structure of VEGFA targeting shRNA and
Cas13 (CasRx and Cas13bt3) vectors. **(F)** VEGFA mRNA knockdown from shRNA, CasRx
and Cas13bt3 in HEK293FT cells (n = 7). **(G)** VEGFA protein expression from ELISA after
transfection of shRNA, CasRx and Cas13bt3 plasmids (n = 4). **(H)** Venn diagram of a total
number of differentially expressed genes (DEGs) from Cas13bt3 only, Cas13bt3-NT sgRNA
and Cas13bt3-VEGFA sgRNA plasmids. **(I)** Volcano plots showing DEGs from Cas13bt3
only, Cas13bt3-NT sgRNA and Cas13bt3-VEGFA sgRNA compared to transfection control in
HEK293FT cells. VEGFA gene is denoted by a black arrow. **(J)** TPM plot for VEGFA and
candidate off-target genes. **(K)** Top 5 enriched GO molecular function (MF) terms from
Cas13bt3 only, Cas13bt3-NT sgRNA and Cas13bt3 VEGFA sgRNA. NES above zero
represents significant activation of the pathway, while NES below zero represents significant
suppression of the pathway. Orange text denotes key factors that are upregulated; blue text
denotes key factors that are downregulated. Data presented as mean ± SEM. Statistical analysis
was conducted by one-way ANOVA and Tukey's multiple comparison test **(F, G)** and Welch's
t-test **(J)**. ns: not significant, *$P \leq 0.05$, **$P \leq 0.001$, ***$P \leq 0.0001$, ****$P \leq 0.00001$.

Multiplexing sgRNAs for CRISPR-Cas13bt3 knockdown of target gene from a single-AAV
vector

The small size of Cas13bt3 and the significant downregulation of *VEGFA* mRNA observed
prompted us to hypothesize that multiple sgRNAs can be delivered using a single AAV vector
to increase overall targeting and RNA knockdown efficiency. In theory, multiple sgRNAs
would increase rate of Cas13 activation and levels of intracellular Cas13-sgRNA complexes
for RNA silencing.

To understand Cas13bt3 activation and demonstrate multiplex targeting, we transfected
Cas13bt3-VEGFA sgRNA and Cas13bt3 with a VEGFA sgRNA array (carrying three
VEGFA-targeting sgRNAs) (**Figure 2A-C**). Surprisingly, cells transfected with Cas13bt3-
VEGFA sgRNA or Cas13bt3-VEGFA sgRNA array showed similar levels of significant

VEGFA mRNA knockdown (<90%) (**Figure 2D,** n = 3 independent experiments). We then

sought to study collateral and off-target cleavages from the VEGFA sgRNA array system using

RNA-seq. Overall DEGs for both single and multiple sgRNA groups when compared to NT

control were comparable and suggested minimal collateral activity from the sgRNA array

system (**Figure 2E**). Again, we identified top candidate off-target genes for the other *VEGFA*

sgRNAs (**Figure S4B and C**) and did not detect any significant off-target activity in these

genes by the Cas13bt3-VEGFA sgRNA array system as well (**Figure 2F**). Active Cas13

nucleases competing for target mRNA or low levels of *VEGFA* mRNA transcripts may have

resulted in the lack of improvement in targeting efficiency, however our results suggest that

multiple sgRNAs can be used simultaneously with Cas13bt3 while still avoiding significant

collateral activity.

Apart from competition for substrate mRNA, it was recently reported that the presence of "UC"

sites on the target mRNA may also affect knockdown efficiency, with "UC" sites within 10-20

bases of spacer binding to allow for optimal cleavage (**Figure S3A**)(*16*). We found that while

sgRNA1 and sgRNA2 had "UC" sites in these optimal regions, sgRNA3 had only one distal

"UC" site 40 nucleotides away from spacer binding (**Figure S3B**). While we did not

independently test the knockdown efficiency of our sgRNAs, this finding suggests sgRNA 3

alone would achieve low levels of RNA knockdown. As no improved knockdown was

observed from the VEGFA sgRNA array, we proceeded to investigate Cas13bt3 with a single

sgRNA further.

A

B pAAV.U6.presgRNA.CMV.Cas13bt3.HA

ITR — U6 — gRNA — CMV — Cas13bt3 — HA ITR — 3981bp
                                          pA

U6.presgRNA.pA

U6 — DR — gRNA — DR — pA

U6.presgRNAarray.pA

U6 — DR — gRNA1 — DR — gRNA2 — DR — gRNA3 — DR — pA
          Cassette of 3 gRNAs

C

VEGFA sgRNA 3
TCTGCATTCACATTTGTTGTGCTGTAGGAA

VEGFA sgRNA 2
CACTTCGTGATGATTCTGCCCTCCTCCTTC

VEGFA mRNA

Ex1 — Ex2 — Ex3 — Ex4 — // — Ex8

VEGFA sgRNA 1
TATGTGCTGGCCTTGGTGAGGTTTGATCCG

D

VEGFA expression

*** **** ****

Cas13bt3 : − + − − − + + +
NT : − − + − − + − −
VEGFA : − − − + − − + −
Array : − − − − + − − +

E

VEGFA sgRNA vs NT sgRNA

−Log10(p-value)

VEGFA

Log2FC

VEGFAarray sgRNA vs NT sgRNA

VEGFA

Log2FC

F

*   ns

TPM

VEGFA PRKAR2B EHD4 USP22 RHBDL1 POMT2 SNX33 RER1 POLA1 SLC9B2 SYT10 BARX1 LOC100192426 CDH20 MORC3 ZNF32-AS3 SUGT1P4-STRA6LP FAM172A CLRN2 MUC16 KCNQ5 PNMA8A NUP210L LOC105373744 LYSMD3 SKP1 CPA4 SENP5

nd   nd   nd   nd nd   nd   nd

■ Non-targeting ■ Cas13bt3-VEGFA ■ Cas13bt3 VEGFA array

Fig. 2. Guide-dependent and guide-independent off-targets from Cas13bt3-mediated VEGFA knockdown. (A) Schematic of single and multiplex targeting with Cas13bt3 and **(B)** U6-sgRNA domain in single sgRNA and multiple sgRNA vectors. **(C)** Guide RNA targeting regions in VEGFA mRNA. **(D)** VEGFA mRNA knockdown from Cas13bt3, non-targeting and targeting sgRNAs. **(E)** Volcano plots showing differentially expressed genes between Cas13bt3 with non-targeting and **(F)** TPM plots for VEGFA and candidate off-target genes. Target gene VEGFA is highlighted in red. Data presented as mean ± SEM. Statistical analysis was conducted by one-way ANOVA and Tukey's multiple comparison test **(D)** and Welch's t-test **(F)**. ns: not significant, *P≤0.05, ***P≤0.0001, ****P≤0.00001.

AAV2.7m8 delivery efficiently expresses Cas13bt3 in various retinal cells

Human organoids are excellent models for gaining clinically relevant information on new therapeutic agents, as they are derived from human pluripotent stem cells (PSCs) and reproduce morphological, functional and transcriptomic features *in vitro*(*23*). Recently, hESC-derived retinal organoids have been used to investigate eye development and model inherited retinal diseases as well as the testing of novel gene therapies(*24, 25*). While CRISPR-Cas9 has entered the clinic, Cas13-based therapies largely remain at the preclinical stage, and demonstrating utility of Cas13 in retinal organoids will be a major advance for ocular gene therapy.

Here we utilised our well-established 2D/3D differentiation protocol to generate hESC-derived retinal organoid containing all the major cell types in the retina(*26*). First, to ensure maximum expression of the Cas13 nuclease in the retinal cells, we subjected the Cas13bt3 protein sequence to codon optimisation to obtain two other variants of Cas13bt3 (Cas13bt3$_{COP1}$ and Cas13bt3$_{COP2}$) (**Figure S5A**). We compared the expression levels of the original Cas13bt3 with Cas13bt3$_{COP1}$ and Cas13bt3$_{COP2}$ by transfection in HEK293FT and ARPE19 cells and found both variants to exhibit increased expression and knockdown efficiency (**Figure S5B, C and D**). Among them, Cas13bt3$_{COP1}$ showed the highest expression and was used for subsequent investigation. To elucidate the activity of Cas13bt3 and efficacy for AAV gene therapy RNA editing in *bona fide* human retinal cells we transduced 19 weeks old organoids with AAV2.7m8.U6.NTsgRNA.Cas13bt3 and AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 viral vectors. Retinal organoids were then harvested for single cell RNA sequencing (scRNA-Seq)

6 weeks after treatment (~25 weeks) for (**Figure 3A**). First, we confirmed that the major retinal
cell types were clustered similarly across experimental conditions (S1: Untreated; S2:
AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3; and S3: AAV2.7m8.U6.NTsgRNA.Cas13bt3)
(**Figure 3B**, n = 3 pooled organoids per condition). We used our recently described retinal atlas
dataset to determine fidelity of retinal organoids and identified expression of the different
retinal cell markers(*27*), confirming exclusive cell-specific expression of the markers (**Figures**
3C and S6, S7).

Next, to examine Cas13bt3 expression in the retinal organoids, we evaluated read numbers and
classified cells containing more than 10 reads of Cas13bt3 to be positive for viral infection.
Using this threshold, we determined an average transduction efficiency of 18.78% (1403 out
of 7467 cells) for the VEGFA sgRNA group and 25.95% (2260 out of 8706 cells) for the NT
sgRNA group. Next, we analysed transduction and presence of Cas13bt3 in specific cell types.
While all cell types showed noticeable levels of Cas13bt3, RPE cells, in particular, showed
high levels of expression with up to 50% of cells containing Cas13bt3 in the VEGFA sgRNA
(S2) treatment group. Interestingly, in the NT sgRNA group, only ~15% of RPE cells expressed
Cas13bt3, which could owe to varying culture conditions. Bipolar cells showed the least
expression of Cas13bt3 (≤10%), while all other cell types showed moderate expression (≤25%)
(**Figure 3D**). Overall, our results indicated AAV2.7m8 as a useful delivery vector for cells
across retinal cell types.

To investigate the impact of the NT sgRNA in retinal cells, we first compared DEGs between
cells positive and negative for Cas13bt3 in the retinal organoid that received the
AAV2.7m8.U6.NTsgRNA.Cas13bt3. Here, we observed a significant number DEGs across all
cell types, and in similar trend to the HEK293FT cells where most DEGs were upregulated
genes (**Figure 3E**). Albeit at a lower extent the same pattern was observed in the retinal
organoid transduced with AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 (**Figure 3F**), suggesting
that the NT sgRNA exhibited greater impact on the transcriptome. To determine the specific
cell types that were more sensitive to Cas13bt3 expression, we identified unique DEGs from
each cell type (to remove common effects from AAV delivery and culturing conditions) across
the retinal cells and found macroglial and bipolar cells to exhibit the highest DEGs, which was
especially higher in the NT sgRNA group (Macroglial - NT: 5571 vs VEGFA: 2928, Bipolar
– NT: 3725 vs VEGFA: 1633) (**Figure 3G**).

To understand this effect from the NT sgRNA, we further identified DEGs unique to the NT

sgRNA and VEGFA sgRNA group and performed GO analysis to identify responses exclusive

to the delivered sgRNA. However, ubiquitin protein-like (UBL) associations (e.g., "*UBL ligase*

binding", "*UBL transferase activity*") were top hits for both groups across macroglial, bipolar,

amacrine and RPE cells (**Figure 3H and Data S2**). Ubiquitin processes are often activated in

response to immune stress to facilitate the clearing of proteins inside cells (*28*), and our results

suggest the delivery of Cas13bt3 with an activating sgRNA (non-targeting or targeting) to

significantly elevate the immune response in retinal cells. This stress response is likely higher

with the NT sgRNA group due to unspecific RNA binding, causing the increased

transcriptomic impact. The strong stress response from all cells to Cas13bt3 delivery is also

likely to have made sgRNA-specific responses challenging to distinguish.

Fig. 3. Cas13bt3 treatment and single-cell RNA sequencing of 3D retinal organoids. (A) Schematic of 3D retinal organoid scRNA-seq experimental procedure. **(B)** UMAP plots show retinal cell type clusters for the untreated, NT sgRNA and VEGFA sgRNA treatment groups. **(C)** Expression of cell-type specific markers in cone cells. **(D)** Infection rate of retinal organoids by cell type and treatment group. **(E)** Multi-volcano plots showing differentially expressed genes (DEGs) between Cas13bt3 positive and negative cells from the NT sgRNA and **(F)** VEGFA sgRNA treatment groups. **(G)** Number of unique DEGs across retinal cell types. **(H)** GO molecular function terms related to unique DEGs from Cas13bt3-NT sgRNA treatment for macroglial and bipolar cells. UBL = Ubiquitin-like protein.

Cas13bt3 can significantly knock down VEGFA mRNA in retinal organoids

Next, we examined Cas13bt3 targeting efficiency and effect on *VEGFA* expression across the retinal cell types. First, we determined *VEGFA* expression in control untreated organoids. *VEGFA* mRNA is present across all retinal organoids cell types. Expression of VEGFA varied significantly between retinal cell types with macroglial, rod, cone photoreceptors, and bipolar cells expressing significantly less *VEGFA* than RPE cells (**Figure S8A, B**).

Furthermore, *VEGFA* expression was also detected in all treated organoids, albeit at different levels (**Figure S8C-E**). DEGs were detected in all cell types between retinal organoids transduced with Cas13bt3-VEGFA sgRNA and Cas13bt3-NT sgRNA. However, a significant downregulation of *VEGFA* was only detected in RPE cells (**Figure S9**). While Cas13bt3 positive cells have been identified, the abundance of Cas13bt3 reads within cells may still vary (**Figure 4A**) and affect determination of targeting efficiency. We hypothesised that increased intracellular Cas13bt3 transcripts allows for higher targeting efficiency and thereby increased silencing of *VEGFA* mRNA (**Figure 4B**). We first examined the number of Cas13bt3 reads in cells that received AAV2.7m8.U6.NTsgRNA.Cas13bt3 and AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 treatments (**Figures 4C and S10**). To accurately determine Cas13bt3 effect and to understand abundance of the targeting complex within each cell we examined AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 treated cells. Here, we found RPE and macroglial cells expressed high levels of Cas13bt3 expression while most cones, rods and horizontal and bipolar cells only showed low levels of Cas13bt3 expression (**Figure 4C**). We

selected a threshold to classify high ($\geq$ 50 reads) and low (<50 reads) expression of Cas13bt3

that enables evaluation of targeting efficiency. Next, we quantified *VEGFA* mRNA expression

in cells expressing high and low levels of Cas13bt3. To this end, we pooled cells within the

high and low Cas13bt3 thresholds and compared average *VEGFA* expression against Cas13bt3

negative cells to determine if loss of *VEGFA* is from is dependent on high levels of Cas13bt3.

Significant decrease of *VEGFA* in high Cas13bt3-expressing cells compared to low Cas13bt3-

expressing cells would indicate *VEGFA* silencing from increased targeting activity, while

significant reduction compared to negative cells would indicate *VEGFA* loss from viral

delivery. Here, we observed that RPE cells were the only cells to show significant reduction of

VEGFA expression in the high Cas13bt3 when compared to low Cas13bt3 cells ($P < 0.05$)

(**Figure 4D**, red square). Viral delivery appeared to upregulate *VEGFA* expression in

macroglial cells with both low and high Cas13bt3-expressing cells exhibiting significant

increase in VEGFA expression (**Figure 4D**). Interestingly, the increase of *VEGFA* was not

significant between low and high Cas13bt3 expressing macroglial cells, suggesting a

considerable loss of *VEGFA* (**Figure 4D**). We therefore further investigated retinal cells that

showed moderate transduction (~25%) of Cas13bt3-VEGFA sgRNA (i.e. amacrine, horizontal

and macroglial) and compared *VEGFA* expression to cells that received Cas13bt3-NTsgRNA.

Promisingly, we observed near significant downregulation of *VEGFA* in amacrine and

horizontal cells ($P = 0.075$ and $P = 0.051$ respectively), suggesting higher transduction rates

may lead to significant downregulation in the other cells as well (**Figure S11**). We then studied

DEGs in the RPE cells to determine if the off-target genes previously identified were affected

from the significant Cas13bt3-mediated *VEGFA* silencing. Importantly, none of the off-target

genes we previously identified were significantly affected (**Figures 4E and S12**). Finally, GO

analysis revealed that "*PDGF receptor binding*" was among the top five downregulated

processes (**Figure 4F and Data S3**), akin to our observations in HEK293FT cells.

While we only observed significant *VEGFA* mRNA knockdown in RPE cells, whether this is

a result of higher transduction or RPE cells carry cell-specific factors that are conducive for

Cas13 activity needs further investigation. Regardless, with RPE cells being major contributors

of *VEGFA* expression, our results show that Cas13bt3 can be a potent tool against diseases

such as DR and nAMD.

Fig 4. VEGFA knockdown efficiency in human retinal organoids. (A) Schematic of variable Cas13bt3 expression in retinal organoid cells and **(B)** relationship between CRISPR-Cas13bt3 expression and targeting efficiency. **(C)** Histograms showing the level of Cas13bt3 expression by number of reads across retinal cell types in the VEGFA sgRNA group. **(D)** Violin plots of VEGFA expression against the level of Cas13bt3 expression across retinal cell types compared to Cas13bt3 negative cells. Red box indicates cells of interest for high VEGFA targeting activity. **(E)** Volcano plot of DEGs between RPE cells with high and low Cas13bt3 expression. Significant knockdown of VEGFA is highlighted in red. **(F)** Top five significant GO molecular function terms between RPE cells expressing low and high Cas13bt3. Statistical analyses were conducted using Student's t-test **(D)**. PPi: Pyrophosphatase. TM: Transmembrane.

VEGFA knockdown impedes disease progression in humanised mouse model of proliferative diabetic retinopathy

Lastly, we sought to investigate if *VEGFA* knockdown can control neovascularisation *in vivo* in humanised models of DR. This offers a potential new tool against eye diseases causing neovascularisation. To this end, we utilised the trVEGF029 (Kimba) mouse, a humanised transgenic mouse model where the neuroretina overexpresses h*VEGFA* leading to neovascularisation like human proliferative DR(*29*). To ensure that optimal transduction of the retina is being achieved, we delivered a mCherry reporter gene under the control of CMV promoter using the AAV2.7m8 serotype, known to have excellent transduction of the retina, when delivered intravitreally into C57BL/6 mice (**Figure S13A and B**). Immunohistochemistry analysis of AAV2.7m8.CMV.mCherry treated retinal flatmounts showed high levels of fluorescence through all retinal layers (RGCs, bipolar and photoreceptor cells) 4 weeks following injections, confirming quick onset and deep penetration of the retina from intravitreal delivery (**Figure S13C, n = 6 eyes from 3 animals**).

Next we treated Kimba mice with AAV2.7m8 viruses driving our codon optimised Cas13bt3 carrying NT sgRNA (AAV2.7m8.U6.NTsgRNA.Cas13bt3) or VEGFA sgRNA (AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3). Due to severe disease progression in homozygous Kimba (*VEGFA*$^{+/+}$), we opted for the heterozygous (*VEGFA*$^{+/-}$) breed. Six-week-old Kimba

mice received bilateral, intravitreal injections of AAV2.7m8.U6.NTsgRNA.Cas13bt3 or
AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3. Eight weeks following treatment eyes were
analysed for *VEGFA* mRNA expression, neovascular growth, and retinal degeneration (**Figure**
5A). Similar to the control mCherry treatments, high transduction of photoreceptor cells was
observed as demonstrated by co-localisation of mCherry and HA-tag (Cas13bt3) with
Rhodopsin positive cells in the ONL (**Figure 5B, C**). In addition qPCR analysis showed high
expression of Cas13bt3 in Kimba retinas injected with AAV.2.7m8.U6.NTsgRNA.Cas13bt3
and AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 (**Figure 5D,** n = 3-12 retinas). This
corresponded with ~50% reduction ($P < 0.001$) of human VEGFA (*hVEGFA)* mRNA (**Figure**
5E), but not mouse VEGFA (mVEGFA) (**Figure 5F**) in the
AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 group, while no reduction was observed with
AAV2.7m8.U6.NTsgRNA.Cas13bt3 or no treatment (**Figure 5E, F,** n = 3-12 retinas). These
results confirmed specific knockdown of *hVEGFA* from Cas13bt3, despite high
complementarity of *VEGFA* sequences (29 out of 30 nucleotides) in the targeting region
(**Figure 5G**). RNA sequencing further confirmed that mouse VEGFA expression and predicted
off-target genes were not significantly altered, indicating safety of Cas13bt3-mediated RNA
silencing (**Figure S13**).

To confirm if *VEGFA* mRNA knockdown influenced disease progression, we analysed fundus
fluorescein angiography (FFA) images from Kimba eyes for vessel leakage and quantified
vessel density and number of branch points, which are signs indicative for the neovascular
process. We used fluorescein to label vessels of the AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3,
AAV2.7m8.U6.NTsgRNA.Cas13bt3 and untreated eyes. We observed large spots of vessel
leakage, indicating neovascularisation had progressed in both untreated and NT treated eyes
(**Figure 5H,** green expression and white arrows). However, in the
AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 group, no leakage was observed (**Figure 5H,**
VEGFA targeting). Immunohistochemistry against laminin was used to image through the
retinal vascular layers (superficial, intermediate and deep vascular plexuses) (**Figure S16A**
and B). Blood vessels appeared diminished in the intermediate and deep vascular layers for
the VEGFA sgRNA group. In the NT sgRNA group, the neovascular process could be observed
through leaky vessel structure in the same layers (**Figure S16C,** n = 4 retinas). The
AAV2.7m8.U6.VEGFAsgRNA.Cas13bt3 treated eyes also showed significantly lower vessel

density and branch points compared to baseline at 6 weeks post-treatment, unlike the untreated and the AAV27m8.U6.NTsgRNA.Cas13bt3 eyes (**Figure 5I and J,** n = 4-13 retinas, $P < 0.05$). Next, to determine whether *VEGFA* knockdown leads to rescue of overall retinal structure. we performed optical coherence tomography (OCT) to analyse total retinal thickness (TRT) before and after treatments. We observed no significant changes between the untreated and treated groups, and a general decline in TRT across all animals (**Figure S17**). We postulate that the overexpression of *hVEGFA* promotes rapid retinal degeneration that could not be controlled or reversed through *VEGFA* silencing.

Overall, our findings showed Cas13bt3-mediated RNA silencing to significantly reduce *VEGFA* mRNA expression, specifically target *hVEGFA* and control hallmark signs of neovascularisation in a humanised mouse model, promising a useful alternative to current anti-VEGF drugs.

A Week 0 — Kimba hVEGFA+/-; Week 6 — Intravitreal injection; Week 14 — OCT, FFA, qPCR, RNAseq, immunostaining

B mCherry Rhodopsin 50μm

C HA-tag = Cas13bt3, mCherry = AAV2.7m8; ONL OS mCherry HA 25μm; Merge ONL OS DAPI mCherry HA 25μm

D Cas13bt3 expression (ΔΔCt) — Untreated, NT, VEGFA; **** ****

E hVEGFA expression — Untreated, NT, VEGFA; * **

F mVEGFA expression — Untreated, NT, VEGFA; ns ns

G VEGFA sgRNA target region
HUMAN CGGATCAAACCTCACCAAGGCCAGCACATA
MOUSE CGGATCAAACCTCACCAAAGCCAGCACATA

H Baseline / Endpoint — Untreated, Non-targeting, VEGFA-targeting; 2mm

I Vessel density (%) — Untreated, NT, VEGFA; ns ns *

J No. of branch points — Untreated, NT, VEGFA; Baseline Endpoint; ns ns *

Fig 5. *In vivo* VEGFA mRNA knockdown in humanised Kimba mice. (A) Schematic of experimental procedure for intravitreal treatment of Kimba mice. **(B and C)** Immunohistochemistry of retinal flatmount and cryosection showing transduction of AAV2.7m8 and expression of Cas13bt3 (Magnification: 20X and 40X, respectively). **(D)** Expression of Cas13bt3 from Kimba mice retinas. (E) Expression of mouse and human VEGFA mRNA in Kimba retinas. **(F)** VEGFA mRNA targeting regions in human and mouse transcriptome. **(G)** Plot of vessel density and number of branch points from untreated and treated Kimba eyes at baseline (6 weeks) and endpoint (14 weeks) analysed from in vivo central fluorescein fundus angiography (FFA). **(H)** Central fluorescein fundus angiography (FFA) images were obtained from Kimba mice at baseline (6 weeks) and endpoint (14 weeks) after AngioTool analysis and quantification **(I and J)**. Images were edited using Adobe Photoshop to remove central retinal arteries and veins. Green spots and white arrows indicate leakage from blood vessels. Data presented as mean ± SEM. Statistical analysis was conducted by one-way ANOVA with Tukey's multiple comparison test **(D-F)** and Multiple paired t-tests **(I-J)**. *$P \leq 0.05$, **$P \leq 0.001$, ****$P \leq 0.00001$.

Discussion

Our comprehensive *in vitro*, *ex vivo* and *in vivo* characterisation of Cas13bt3 activity and performance in the retina provides novel insights for therapeutic use of the enzyme. Cas13bt3 was first described only two years ago but has garnered significant attention for its compact size and reported specificity(*11*). Structural characterisation and activation mechanisms have recently been described spearing immense interest in the molecule for therapeutic development(*16, 30*). Due to ongoing controversy over the exact specificity of Cas13 nucleases(*10*), we aimed to characterise Cas13bt3 efficiency and specificity in mouse and human cells. We chose to target the *VEGFA* transcripts, as this growth factor is essential for both physiological and pathological angiogenesis. In the retina, physiological functions include its role as a survival factor for retinal ganglion cells(*31*), photoreceptors, Müller cells(*32*), and RPE(*33*). In disease, overexpression of VEGFA leads to neovascularisation and vision loss. Therefore, clinical applicability of VEGFA downregulation in specific cell types of the retina is of great importance.

Previous studies have demonstrated transcript knockdown using CasRx proteins(*11, 12*).

Notably, CasRx-mediated *VEGFA* mRNA knockdown has also been shown to reduce choroidal

neovascularisation after silencing VEGFA in RPE cells *in vivo* in laser-induced choroidal

neovascularisation model(*4*). Our results confirmed that Cas13bt3 mediates significant

knockdown of *VEGFA* in human cell lines, hESC-derived retinal cells, and the mouse retina.

We further investigated the transcriptomic impact of expressing active Cas13bt3 in the cells

and found that initiating Cas13bt3's target search process with a non-targeting sgRNA can lead

to significant RNA binding, upregulating the expression of several genes. A targeting sgRNA,

such as with VEGFA, appears to narrow the target-search process and thereby reduce

transcriptomic impact.

In vitro, and specifically in HEK293FT cells, several studies have shown Cas13 nucleases to

be highly precise(*10*). Most of these studies, however, compared differentially expressed genes

to a NT control, not accounting for transcriptomic perturbations from a Cas13-NT sgRNA

complex. A recent study showed high collateral activity from CasRx when compared to an

empty vector(*34*). Here, we compared targeting Cas13bt3 to vector controls showing minimal,

but positive collateral activity. More importantly, we show that Cas13 with a NT sgRNA has

significant transcriptomic impact resulting in high levels of DEGs. This demonstrates that NT

sgRNAs are not appropriate controls for determining specificity of Cas13 nucleases raising

awareness against their use.

We also show that multiple sgRNAs can be incorporated within a single AAV vector with

Cas13bt3 driven by a strong ubiquitous promoter (CMV) for targeted knockdown of *VEGFA*

mRNA. Even though this appears to be an attractive approach to enhance transcript knockdown

our study established that multiple sgRNAs failed to improve transcript degradation. The rate

of mRNA knockdown was comparable to single sgRNA treatments. Previously, we similarly

observed with multiplexed RNA editing with CasRx carrying three *VEGFA*-targeting sgRNAs,

the multiple sgRNA system unexpectedly led to decreased knockdown efficiency(*19*), raising

the suspicion that substrate competition or inadequate Cas13 activation plays an inhibitory role

in RNA targeting efficiency. Interestingly, the comparable levels of DEGs in both single and

multiple sgRNA vectors suggest that Cas13bt3 activation is not drastically elevated with the

presence of additional sgRNAs. While we did not investigate the delivery of sgRNAs targeting

multiple genes, our results show multiple sgRNAs do not increase overall collateral activity,

which might prove useful when several genes must be targeted simultaneously to control
disease progression. This approach could potentially be applied even to ocular
neovascularisation by simultaneously targeting the *VEGFA* and *PDGF* genes for enhanced
treatment(*35*).

Furthermore, we investigated for the first time AAV-mediated Cas13 RNA expression and
editing in retinal organoids at single cell resolution. While several studies have demonstrated
VEGFA silencing for control of neovascularisation, previous studies have focused on specific
cell types that are believed to be major contributors to *VEGFA* expression, like RPE and Müller
cells(*4, 36*). While this proves useful to demonstrate proof-of-concept, it does not provide a
comprehensive understanding relevant for therapeutic development as *VEGFA* silencing in one
cell type may have collateral effects on other retinal cells. Retinal organoid models enable the
testing of therapies in an environment that mimics the human retina containing lamination and
major retinal cell types of the developing retina. Here we sought to understand retina-wide
effect following delivery of Cas13bt3 as well as targeted silencing of *VEGFA*. Our findings not
only support the use of Cas13bt3 for ocular gene therapy, but also brings to light several
considerations for its use. First, we showed transduction across all retinal cell types studied
with the clinically relevant AAV2.7m8 vector (NCT04418427) confirming its effectiveness as
a vector for retinal transduction and supporting its clinical utility(*37, 38*). Among the retinal
cell types studied, bipolar cells showed the least transduction by AAV2.7m8, while RPE cells
seemed more susceptible. It is likely that these differences are due to cell-specific transduction
susceptibility, as RPE cells are easily transduced with a variety of AAV capsids(*39*). It should
also be noted that overall transduction was determined using an arbitrary cut-off (Cas13bt3
read counts $\geq$ 10) based on our previous experience with the retinal organoids(*26*). NT sgRNA
activity following gene therapy led to numerous DEGs when compared to VEGFA targeted
sgRNA. This emphasises the need for a well-designed targeting sgRNA for reduced collateral
activity. Retinal organoids showed significantly more DEGs than HEK293FT cells, which is
expected partly due to cellular response to virus delivery. Still, notwithstanding response to
AAV, the higher DEG levels also indicate that retinal cells are more sensitive to Cas13,
demonstrating the importance of studying nuclease activity and efficiency in physiological
relevant models, an important consideration for Cas-mediated gene therapies. As preclinical
studies with Cas13 nucleases are commonly performed in HEK293FT cells, its specificity has

often been exaggerated in the literature. Our results strongly suggest that Cas13 specificity
must be evaluated in relation to untreated or empty vector controls in organ-specific cells to
accurately determine safety. On the front of therapeutic potential, we could only confirm
significant targeted *VEGFA* mRNA knockdown in RPE cells, which is likely due to strong
expression of Cas13bt3 in these cells. Lower copies of Cas13bt3 in the other cells may have
resulted in the lack of significant *VEGFA* knockdown, and further studies are required to
understand the levels of Cas13 expression necessary in other retinal cells for effective activity.
Testing of cell type specific promoters could be beneficial as CMV promoter is known for its
susceptibility to silencing in certain cell types, for example stage specific PSC-derived
photoreceptor cells (REF). Importantly, with RPE cells being the primary producers of *VEGFA*
in the retina, the significant downregulation of *VEGFA* mRNA in RPE strongly suggests the
benefits of Cas13bt3 gene therapy as a potential anti-VEGF tool for neovascular eye disease.

Lastly, we tested the anti-VEGF potential of Cas13bt3 in the retina of a humanised mouse
model of diabetic retinopathy, the Kimba mouse. We opted to examine anti-VEGF potential of
Cas13bt3 in Kimba mice as it shows chronic disease and all major signs of proliferative
retinopathy (*29*). Importantly, as Kimba mice expresses human *VEGFA*, the same vectors
tested in retinal organoids could also be used *in vivo*. Others have shown the control of ocular
neovascularisation using mouse models of laser-induced injury(*4*), streptozotocin (STZ)-
induced diabetes(*40*), or oxygen-induced retinopathy (OIR)(*41*). However, these models have
some drawbacks. The laser-induced choroidal neovascularisation model often used in anti-
VEGF preclinical studies does not accurate model the chronic retinal degeneration observed in
DR and nAMD patients and presents *VEGFA* overexpression only as a response to injury(*42*).
Diabetic mouse models do not recapitulate proliferative retinopathy which is often the cause
of severe vision impairment. Vascular pathogenesis is also delayed and varied based on
response to STZ(*43*). OIR mouse models rapidly present proliferative retinopathy, however
neovascularisation regresses within a week(*44*).

Following RNA editing gene therapy in Kimba retinas, we demonstrated the specific
knockdown of human *VEGFA*, with no significant impact on endogenous mouse *VEGFA*. This
shows strong evidence for the specificity of targeted Cas13bt3 activity. Silencing of Cas13bt3
VEGFA also rescued vascular phenotypes showing an improvement in controlling vascular
growth and leakiness. The major concern with current anti-VEGF treatment is the need for

frequent intravitreal injections. Recent efforts to address this problem have employed gene

editing strategies like targeting *VEGFA* using CRISPR-Cas9 (*36, 45*). However, efficacy of

such treatments are limited by efficient and safe delivery of vectors due to the size of Cas9(*46,*

47), poor efficiency of compact Cas9 nucleases(*48*) or have reported the risk of genomic

alterations from long-term expression of Cas9(*49*). Our results indicate that a single intravitreal

injection could achieve durable and potent while reversible silencing of *VEGFA*, addressing

drawbacks with current gene editing strategies. While RNA editing is thought to be safer than

DNA-targeting options, it is prudent to consider effects of long-term Cas13 expression, and

Cas13 systems amenable to regulation are of consideration for therapeutic development.

There are a few limitations to our study. First, we did not extensively study Cas13bt3

expression in ARPE19 cells, where we also observed potent knockdown. While HEK293FT

appears resistant to high levels of collateral activity, ARPE19 cells may exhibit significantly

higher levels of transcriptomic effects. This is supported by the high levels of DEGs we

observed in various cells from the retinal organoids, suggesting significant inflammatory

responses as well as cell-specific collateral activity. Specifically, the upregulation of UBL

processes across cell types regardless of VEGFA silencing, strongly indicate increased cellular

stress from delivery of Cas13. As RNA targeting molecules like Cas13 require persistent

expression for therapeutic effect, this could pose a significant safety concern. Second, while

vessel density and branching were noticeably reduced, we could not extensively study recovery

of phenotype in our mouse model due to severity of the disease. OCT readings showed

significant retinal degeneration at treatment timepoint, making rescue of phenotype

challenging to assess. Lastly, we were also unable to confirm reduction of VEGFA protein

from the mice retinas as not enough protein could be obtained for quantitative analysis.

CRISPR-Cas technology has advanced at breakneck speed in the last decade revolutionising

medicine and offering potential treatment opportunities to numerous previously incurable

diseases. The recent FDA approval of an *ex vivo* CRISPR therapy for sickle-cell disease

(branded Casgevy) is evidence of this progress. However, a comprehensive understanding of

the CRISPR nucleases is crucial. In this study we offer novel insights into Cas13 targeting

mechanisms and its utility in targeting retinal and RPE cells providing evidence for the

potential of Cas13-based therapies for ocular disease. This study demonstrated that the

hypercompact Cas13bt3 can be efficiently delivered and expressed in various retinal cell types

effectively controlling *VEGFA* expression and neovascularisation. In-depth evaluation of
collateral activity in retinal cells would facilitate development of a safe and effective Cas13-
mediated anti-VEGF AAV gene therapy.

Materials and Methods

Plasmid construction

The protein sequences of CasRx and Cas13bt3 were gifts from P. Hsu(*50*) and H. Yang(*11*),
respectively. Cas13bt3 was codon optimised using online tools from Integrated DNA
Technologies (IDT), GenScript and Benchling. The all-in-one AAV plasmids for shRNA,
CasRx and Cas13bt3 were designed as previously described(*19*), and synthesised by
GenScript. All plasmids used in this study are available from Addgene.

Cell culture and transfections

HEK293FT cells (American Type Culture Collection, ATCC) were maintained in Dulbecco's
Modified Eagle's Medium (DMEM; Thermo Fisher Scientific) supplemented with 10% (v/v)
fetal bovine serum (FBS, Bovogen Biologicals) and 1% (v/v) antibiotic-antimycotic (Thermo
Fisher Scientific) in a humidified 5% CO_2 incubator at 37°C. ARPE19 (ATCC) cells were
maintained in DMEM/F-12 (Thermo Fisher Scientific) supplemented with 10% (v/v) FBS and
1% (v/v) antibiotic-antimycotic.

RT-qPCR

At 48 hours after transfection, RNA was extracted from cells using the Monarch® Total RNA
Miniprep Kit (New England Biolabs and 200ng of RNA was converted to complementary DNA
(cDNA) using High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific)
according to the manufacturers' instructions. cDNA was diluted with water at the ratio of 1:10.
qPCR was performed with 2uL of cDNA using the TaqMan™ Fast Advanced Master Mix
(Thermo Fisher Scientific) according to manufacturers' instructions with the following

TaqMan probes human VEGFA (Hs00900055_m1; Thermo Fisher Scientific), mouse VEGFA

(Mm00437306_m1; Thermo Fisher Scientific), human GAPDH (Hs02786624_g1; Thermo

Fisher Scientific), mouse GAPDH (Mm99999915_g1, Thermo Fisher Scientific). Mouse

eyeballs were enucleated and stored in 4°C in RNALater® until dissection. Retina was dissected

from injected animals and placed in lysis buffer and sonicated for 2 seconds prior to RNA

extraction and qPCR as previously described.

Western blot

At 48 hours after transfection, cells were harvested and collected in Pierce™ RIPA buffer

(Thermo Fisher Scientific) containing Pierce™ Phosphatase Inhibitor Mini Tablets (Thermo

Fisher Scientific). Cell lysate was sonicated for 5 seconds and centrifuged for 15 minutes at

13300rpm at 4°C. The supernatant was collected, and protein levels were quantified using the

Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific) as per manufacturers' instructions.

For Western blot, 10µg of protein mixed with NuPAGE™ LDS Sample Buffer (4X) (Thermo

Fisher Scientific) and loaded in NuPAGE™ 4 to 12% Mini Protein Gel (Thermo Fisher

Scientific) for gel electrophoresis at 150V for 30 minutes. Transfer to Amersham™ Hybond™

PVDF membrane (Cytiva) was performed at 30V for 1 hour. Protein blot was blocked with 5%

skim milk for 1 hour, followed by incubation with HA-tag (C29F4) Rabbit mAb (Cell Signaling

Technology®) or Anti-Actin Antibody, clone C4 (Merck) overnight at 4°C. Blots were then

stained with Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, HRP

(Invitrogen™) or Goat anti-Rabbit IgG (H+L) Secondary Antibody, HRP (Invitrogen™) for 2

hours at room temperature. Blot was incubated with Pierce™ ECL Western Blotting Substrate

for 1 minute before imaging using Bio-Rad ChemiDoc. All washes were performed with 1X

PBST for 5 minutes, repeated 3 times.

ELISA

At 48 hours after transfection, 100µL of supernatant was collected and ELISA was performed

using the Human VEGF DuoSet ELISA (R & D Systems) as per manufacturers' instructions.

Briefly, 96-well plates were incubated with Capture Antibody in ELISA Plate coating buffer

overnight at room temperature. Plates were then blocked with Reagent Diluent for 1 hour at

room temperature, and 100uL of supernatant was incubated for 2 hours at room temperature.

Then, Detection antibody was added and incubated for 2 hours at room temperature. After

incubation with Streptavidin-HRP and Substrate reagent for 20 minutes each, the provided stop

solution was added, and plate was read using the Tecan Spark® Micro plate reader at 450nm

with reference wavelength 570nm.

Immunohistochemistry

Retinal flatmounts were washed with 1X PBS containing 0.5% Triton-X-100 (PBST) and

permeabilised in -80°C for 15 minutes. After washing with PBST, flatmounts were incubated

in anti-rhodopsin antibody (1:500; Abcam) with 2% donkey serum and 2% PBST in 4°C for

72 hours. This was followed by incubation in Alexa Fluor 488 AffiniPure Donkey Anti-Mouse

IgG (H+L) (1:500; Jackson ImmunoResearch) at room temperature for 2 hours. After washing,

the flatmounts were stained with Anti-Laminin-2 (α-2 Chain) antibody, Rat monoclonal (1:250;

Merck) and incubated in 4°C for 5 nights. This was followed by incubation in Alexa Fluor®

647 AffiniPure Donkey Anti-Rat IgG (H+L) (1:250; Jackson ImmunoResearch) at room

temperature for 2 hours. Then, the retinae were washed and incubated in mCherry Monoclonal

Antibody (16D7), Alexa Fluor 594 (1:500; Thermo Fisher Scientific) at 4°C for 72 hours.

Retinal cryosections were blocked using serum-free protein block (Agilent Dako) at room

temperature for 1 hour, followed by incubation at 4°C overnight with primary antibody as

previously described. Secondary antibody incubation was performed at room temperature for

1 hour as previously described. Slides were stained with NucBlue™ Live ReadyProbes™

Reagent (Hoechst 33342; Thermo Fisher Scientific) for 10 minutes before mounting as

previously described and imaged on the Stellaris 5 confocal microscope.

hESC Maintenance and Retinal Differentiation Culture

hESCs were maintained on feeder-free conditions on E8 (Thermo Fisher) and Geltrex-coated

six-well plates. In brief, for retinal differentiation hESCs were maintained until 90%–95%

confluent, then FGF-free medium was added to the cultures for 2 days followed by a neural

induction period where RPE and retinal vesicles appeared. NRVs were excised and kept in low

binding 96-well plates for maturation.

AAV transduction of retinal organoids

Retinal organoids were maintained in ALT90 medium as previously described (West et al.,

2022) and AAV transduction was performed during the 15-19 weeks of differentiation. On the

day of transduction, AAV vectors were diluted in 300µL of fresh ALT90 medium at titer of

1x10E11 vg/organoids. The following day treated wells were supplemented with 700µL of

fresh ALT90 medium. Transduced organoids were incubated for 72 hours in the AAV vector-

containing medium. After the 72-hour incubation, the vector-containing medium was aspirated,

and 1mL of fresh ALT90 medium was added to each well. Transduced organoids underwent 6

weeks culture until harvesting for experimental procedures.

Immunohistochemistry of retinal organoids

Retinal organoids were fixed for 45 minutes in 4% paraformaldehyde and incubated overnight

in 20% sucrose, prior to embedding in OCT. Cryosections (14 µm thick) were collected for

analysis and preserved at - 20°C. Cryosections were blocked in 5% goat serum and 1% bovine

serum albumin in PBS for 2 hours. Primary antibodies (ARR3: 1:200, Rhodopsin: 1:1000,

VSX2: 1:200, CRALBP: 1:200, PROX1:1:200, and HuCHuD:1:500) were incubated overnight

at 4°C in blocking solution with 5% goat or donkey serum. After washing of primary antibody

with PBS sections were incubated with secondary antibody in blocking solution for 2 hours at

RT, washed and counter-stained with DAPI (Sigma-Aldrich). Alexa fluor 488 and 546

secondary antibodies (Thermo Fisher Scientific) were used at a 1:500 dilution.

Intravitreal injections

All animal procedures were approved by the Animal Ethics Committees of University of

Melbourne and conducted in accordance with The Association for Research in Vision and

Ophthalmology (ARVO) Statement for the Use of Animal in Ophthalmic and Vision Research. Ethics approval was obtained from The Florey Institute of Neuroscience and Mental Health Animal Ethics Committee (22-036-UM). The Akimba mice developed by Rakoczy *et al*. (*51*) and the heterozygous Kimba mice (*hVEGFA$^{+/-}$*) were used for this study. Mice were bred in the Melbourne Brain Centre animal facility (Kenneth Myer Building, Parkville, VIC, Australia), and were housed in a well-ventilated environment that was kept at a constant room temperature of 21°C, with a 12-hour diurnal light-dark cycle (lights on: 7am, off: 7pm; room illumination < 50 lux). Mouse chow (Barastoc, Melbourne, VIC, Australia) and water were provided ad libitum. 6-week-old Kimba mice were anesthetized with ketamine (80mg/kg) and xylazine (10mg/kg) solution mix (Troy Laboratory, Smithfield, NSW, Australia; i.p. with 30G needle) and given a drop of 0.5% proxymethacaine hydrochloride (Alcaine™, Alcon Laboratories, Frenchs Forest, NSW, Australia) and 0.5% tropicamide (Mydriacyl™, Alcon Laboratories). Baseline fundus fluoresceine angiography (FFA) were taken (as described below), and animals intravitreally injected with 1uL of AAV2.7m8 viruses using Hamilton syringe attached to a 33-gauge dental needle. Post injection, mice were subcutaneously administered 0.3mL of atipamezole hydrochloride (0.25mg/mL) and allowed to recover on heating pads with food and water *ad libitum*.

Fundus fluorescein angiography and optical coherence tomography

Retinal vasculature was assayed at baseline prior to intravitreal injection of AAVs and 8 weeks after AAV-treatment using fundus fluorescein angiography (FFA) (Spectralis, Heidelberg Engineering, Heidelberg, Germany) immediately after intraperitoneal injection of sodium fluorescein (1%, 100 µL/kg, fluorescite 10%; Alcon Laboratories). Five images (central, nasal superior, temporal superior, nasal inferior, and temporal inferior) were obtained for each eye and analysed using AngioTool(*52*) for vessel density and branch points using default settings.

In vivo retinal structure was assayed with spectral domain optical coherence tomography (SD-OCT) (Spectralis®, Heidelberg Engineering, Heidelberg, Germany) with general anaesthesia, mydriasis and corneal hydration (Systane, Alcon Laboratories) maintained. During image acquisition, retinal volumes (8.1 × 8.1 × 1.9 mm) centred on the optic nerve head were acquired each consisting of 121 evenly spaced horizontal B-scans. Each B-scan comprised of 768 A-

scans with 3.87 μm axial depth resolution, 9.8 μm lateral resolution. OCT scans were captured

with an automated real-time tracking average of 6 frames, at an average speed of 85,000 A-

scans per second. The Heidelberg Eye Explorer 2 OCT reader plugin (Heidelberg Engineering)

was used for analysis. All retinal layers (retinal nerve fibre layer, RNFL; ganglion cell inner

plexiform layer, GCIPL; inner nuclear layer, INL; outer plexiform layer, OPL; outer nuclear

layer, ONL; photoreceptor segments, PR segment) as well as total retinal thickness (TRT) were

automatically segmented (Heidelberg Eye Explorer 2 software). A circular ring around the

optic nerve head (Early Treatment Diabetic Retinopathy Study [ETDRS] outer ring) was

employed for analysis.

Determination of candidate off-target genes

Off-targets of the gRNAs were predicted using a sequence-based approach. gRNAs in length

of 30 bps were first aligned to the human transcriptome (Grch38 cdna from emsembl release

100) using blastn (megablast). Flexible options were set to capture the potential offtargets

which could tolerate up to 7 mismatches in this step, i.e., max_target_seqs = 10000, evalue =

10000, word_size = 5, perc_identity = 0.6. Secondly, the candidates were further filtered with

the threshold that at least 13 bp perfect match should exist in each alignment.

Bulk RNA-seq data analysis

Raw data processing was performed using FASTQC (*53*) to check the reads quality and

Trimmomatic (*54*) tool to further remove the bases and reads with low quality, adaptors and

barcodes. The clean reads were aligned to the reference human genome (version GRCh38),

utilizing STAR v2.7 (*55*) as the aligner. For RNA quantification, RSEM (*56*) was employed to

normalize raw read counts, transforming them into TPM (transcripts per million) values,

providing a relative expression level that is theoretically comparable across samples. The

magnitude (log2-transformed fold change) and significance (P-value) of differential expression

between groups were then calculated. Differentially expressed genes (DEGs) were selected

based on the criteria of a p-value < 0.05 and |log2 fold change| ≥ 1. Gene set enrichment analysis

(GSEA) (*57*), focused on Gene Ontology(*58*), was conducted using the R 4.3.1 Bioconductor

package clusterProfiler(*59*) to elucidate whole transcriptomic patterns between groups. To

account for multiple testing, P-values underwent adjustment through the Benjamini-Hochberg

false discovery rate (BH-FDR).

Single-cell RNA-seq data analysis

After checking the reads quality, the sequencing files were aligned to reference human genome

(version GRCh38) using 10X Genomics CellRanger software (v7.1.0) count pipeline to remove

non-cellular barcodes and keep the cellular barcodes and gene matrix. To further distinguish

the Cas13bt3-valid and -invalid cells, FASTX-toolkit (*60*) was used to keep the reads with

cellular barcodes and convert FASTAQ file into FASTA file. Then, Blastn (*61*) was used to

compute the similarity between cellular reads and Cas13bt3 reference sequence.

Based on the quality of the sequencing reads, appropriate cutoff values for both the percentage

of identical positions and the alignment length were employed to delineate cells as either

Cas13bt3-valid or Cas13bt3-invalid. For either Cas13bt3-valid or -invalid cells, we further

used Seurat v4.3 for data pre-processing, data normalization, feature selection, data scaling,

dimensional reduction and clustering. To refine the characterization of cell types within each

cluster, ScType (*62*) was employed, leveraging genesets specific to eight retinal cell types—

amacrine, bipolar, cones, horizontal, macroglial, microglial, rods, and RPE—validated by Kim

et al. (*27*). The magnitude (log2-transformed fold change) and significance (P-value) of

differential expression between cell types were then calculated. Differentially expressed genes

(DEGs) were selected based on the criteria of a p-value < 0.05 and |log2 fold change| ≥ 1. Gene

set enrichment analysis (GSEA) (*57*), focused on Gene Ontology (*58*), was conducted using

the R 4.3.1 Bioconductor package clusterProfiler (*59*) to elucidate whole transcriptomic

patterns between groups. To account for multiple testing, P-values underwent adjustment

through the Benjamini-Hochberg false discovery rate (BH-FDR).

Statistical analysis

Statistical analyses were performed using GraphPad Prism 10. Analytical methods are

described in respective figure legends with *p*-value indicators. For *ex vivo* studies, to assess the

strength of the relationship between gene expression values and the number of Cas13bt3-valid

read counts following log2 transformation in each cell type, Spearman's rank correlation

coefficient was employed. The associated *p-value* serves as an indicator of the likelihood or

probability that any observed correlation is merely a result of chance.

References

1. J. Y. Wang, J. A. Doudna, CRISPR technology: A decade of genome editing is only
the beginning. *Science* **379**, eadd8643 (2023).
2. S. Kumar, L. E. Fry, J.-H. Wang, K. R. Martin, A. W. Hewitt, F. K. Chen, G.-S. Liu,
RNA-targeting strategies as a platform for ocular gene therapy. *Prog. Retin. Eye Res.*
92, 101110 (2023).
3. Y. Hu, Y. Chen, J. Xu, X. Wang, S. Luo, B. Mao, Q. Zhou, W. Li, Metagenomic
discovery of novel CRISPR-Cas13 systems. *Cell Discovery* **8**, 107 (2022).
4. C. Zhou, X. Hu, C. Tang, W. Liu, S. Wang, Y. Zhou, Q. Zhao, Q. Bo, L. Shi, X. Sun,
H. Zhou, H. Yang, CasRx-mediated RNA targeting prevents choroidal
neovascularization in a mouse model of age-related macular degeneration. *Natl. Sci.*
Rev. **7**, 835-837 (2020).
5. H. Zhou, J. Su, X. Hu, C. Zhou, H. Li, Z. Chen, Q. Xiao, B. Wang, W. Wu, Y. Sun,
Y. Zhou, C. Tang, F. Liu, L. Wang, C. Feng, M. Liu, S. Li, Y. Zhang, H. Xu, H. Yao,
L. Shi, H. Yang, Glia-to-Neuron Conversion by CRISPR-CasRx Alleviates
Symptoms of Neurological Disease in Mice. *Cell* **181**, 590-603.e516 (2020).
6. Z. Zheng, G. Li, C. Cui, F. Wang, X. Wang, Z. Xu, H. Guo, Y. Chen, H. Tang, D.
Wang, M. Huang, Z.-Y. Chen, X. Huang, H. Li, G.-L. Li, X. Hu, Y. Shu, Preventing
autosomal-dominant hearing loss in Bth mice with CRISPR/CasRx-based RNA
editing. *Signal Transduction and Targeted Therapy* **7**, 79 (2022).
7. J. E. Powell, C. K. W. Lim, R. Krishnan, T. X. McCallister, C. Saporito-Magriña, M.
A. Zeballos, G. D. McPheron, T. Gaj, Targeted gene silencing in the nervous system
with CRISPR-Cas13. *Science Advances* **8**, eabk2485 (2022).
8. Y. Li, J. Xu, X. Guo, Z. Li, L. Cao, S. Liu, Y. Guo, G. Wang, Y. Luo, Z. Zhang, X.
Wei, Y. Zhao, T. Liu, X. Wang, H. Xia, M. Kuang, Q. Guo, J. Li, L. Chen, Y. Wang,
Q. Li, F. Wang, Q. Liu, F. You, Collateral cleavage of 28s rRNA by RfxCas13d
causes death of mice. *bioRxiv*, 2022.2001.2017.476700 (2022).
9. P. Shi, M. R. Murphy, A. O. Aparicio, J. S. Kesner, Z. Fang, Z. Chen, A. Trehan, X.
Wu, RNA-guided cell targeting with CRISPR/RfxCas13d collateral activity in human
cells. *bioRxiv*, 2021.2011.2030.470032 (2021).
10. J. F. Bot, J. van der Oost, N. Geijsen, The double life of CRISPR–Cas13. *Current*
Opinion in Biotechnology **78**, 102789 (2022).
11. C. Xu, Y. Zhou, Q. Xiao, B. He, G. Geng, Z. Wang, B. Cao, X. Dong, W. Bai, Y.
Wang, X. Wang, D. Zhou, T. Yuan, X. Huo, J. Lai, H. Yang, Programmable RNA
editing with compact CRISPR–Cas13 systems from uncultivated microbes. *Nat.*
Methods **18**, 499–506 (2021).

12. S. Kannan, H. Altae-Tran, X. Jin, V. J. Madigan, R. Oshiro, K. S. Makarova, E. V.
Koonin, F. Zhang, Compact RNA editors with small Cas13 proteins. *Nature*
Biotechnology, 1-4 (2021).
13. H. Tong, J. Huang, Q. Xiao, B. He, X. Dong, Y. Liu, X. Yang, D. Han, Z. Wang, X.
Wang, W. Ying, R. Zhang, Y. Wei, C. Xu, Y. Zhou, Y. Li, M. Cai, Q. Wang, M. Xue,
G. Li, K. Fang, H. Zhang, H. Yang, High-fidelity Cas13 variants for targeted RNA
degradation with minimal collateral effects. *Nature Biotechnology*, (2022).
14. G. Li, M. Jin, Z. Li, Q. Xiao, J. Lin, D. Yang, Y. Liu, X. Wang, L. Xie, W. Ying, H.
Wang, E. Zuo, L. Shi, N. Wang, W. Chen, C. Xu, H. Yang, Mini-dCas13X–mediated
RNA editing restores dystrophin expression in a humanized mouse model of
Duchenne muscular dystrophy. *The Journal of Clinical Investigation* **133**, (2023).
15. Q. Xiao, Z. Xu, Y. Xue, C. Xu, L. Han, Y. Liu, F. Wang, R. Zhang, S. Han, X. Wang,
G.-L. Li, H. Li, H. Yang, Y. Shu, Rescue of autosomal dominant hearing loss by in
vivo delivery of mini dCas13X-derived RNA base editor. *Sci. Transl. Med.* **14**,
eabn0449 (2022).
16. X. Deng, E. Osikpa, J. Yang, S. J. Oladeji, J. Smith, X. Gao, Y. Gao, Structural basis
for the activation of a compact CRISPR-Cas13 nuclease. *Nature Communications* **14**,
5845 (2023).
17. F.-L. Lin, P.-Y. Wang, Y.-F. Chuang, J.-H. Wang, V. H. Y. Wong, B. V. Bui, G.-S.
Liu, Gene Therapy Intervention in Neovascular Eye Disease: A Recent Update. *Mol.*
Ther. **28**, 2120-2138 (2020).
18. M. Xu, R. Fan, X. Fan, Y. Shao, X. Li, Progress and Challenges of Anti-VEGF
Agents and Their Sustained-Release Strategies for Retinal Angiogenesis. *Drug Des*
Devel Ther **16**, 3241-3262 (2022).
19. Y.-F. Chuang, P.-Y. Wang, S. Kumar, S. Lama, F.-L. Lin, G.-S. Liu, Methods for in
vitro CRISPR/CasRx-Mediated RNA Editing. *Frontiers in Cell and Developmental*
Biology **9**, (2021).
20. Z. Li, Z. Li, X. Cheng, S. Wang, X. Wang, S. Ma, Z. Lu, H. Zhang, W. Zhao, Z.
Chen, Y. Yao, C. Zhang, L. Chao, W. Li, T. Fei, Intrinsic targeting of host RNA by
Cas13 constrains its utility. *Nature Biomedical Engineering*, (2023).
21. Y. Li, J. Xu, X. Guo, Z. Li, L. Cao, S. Liu, Y. Guo, G. Wang, Y. Luo, Z. Zhang, X.
Wei, Y. Zhao, T. Liu, X. Wang, H. Xia, M. Kuang, Q. Guo, J. Li, L. Chen, Y. Wang,
Q. Li, F. Wang, Q. Liu, F. You, The collateral activity of RfxCas13d can induce
lethality in a RfxCas13d knock-in mouse model. *Genome Biol.* **24**, 20 (2023).
22. K. Ofer, B. L. Benjamin, R. S. D. Owen, J. W. t. V. Aartjan, M. Cameron, RNA
structure modulates Cas13 activity and enables mismatch detection. *bioRxiv*,
2023.2010.2005.560533 (2023).
23. M. A. Lancaster, J. A. Knoblich, Organogenesis in a dish: Modeling development and
disease using organoid technologies. *Science* **345**, 1247125 (2014).
24. P. Wahle, G. Brancati, C. Harmel, Z. He, G. Gut, J. S. del Castillo, A. Xavier da
Silveira dos Santos, Q. Yu, P. Noser, J. S. Fleck, B. Gjeta, D. Pavlinić, S. Picelli, M.
Hess, G. W. Schmidt, T. T. A. Lummen, Y. Hou, P. Galliker, D. Goldblum, M.
Balogh, C. S. Cowan, H. P. N. Scholl, B. Roska, M. Renner, L. Pelkmans, B.
Treutlein, J. G. Camp, Multimodal spatiotemporal phenotyping of human retinal
organoid development. *Nature Biotechnology*, (2023).
25. V. I. Maryna, M. H. Daniel, M. M. Eric, T. B. Kevin, W. Mantian, W. P. Cole, J. K.
Alex, C. Xinlan, L. Yaqiao, G. Bence, P. C. David, PCDH15 Dual-AAV Gene

Therapy for Deafness and Blindness in Usher Syndrome Type 1F. *bioRxiv*,
2023.2011.2009.566447 (2023).

26.　E. L. West, P. Majumder, A. Naeem, M. Fernando, M. O'Hara-Wright, E. Lanning,
M. Kloc, J. Ribeiro, P. Ovando-Roche, I. O. Shum, N. Jumbu, R. Sampson, M. Hayes,
J. W. B. Bainbridge, A. Georgiadis, A. J. Smith, A. Gonzalez-Cordero, R. R. Ali,
Antioxidant and lipid supplementation improve the development of photoreceptor
outer segments in pluripotent stem cell-derived retinal organoids. *Stem Cell Reports*
17, 775-788 (2022).

27.　H. J. Kim, M. O'Hara-Wright, D. Kim, T. H. Loi, B. Y. Lim, R. V. Jamieson, A.
Gonzalez-Cordero, P. Yang, Comprehensive characterization of fetal and mature
retinal cell identity to assess the fidelity of retinal organoids. *Stem Cell Reports* **18**,
175-189 (2023).

28.　H. Hu, S.-C. Sun, Ubiquitin signaling in immune responses. *Cell Research* **26**, 457-
483 (2016).

29.　E. P. Rakoczy, I. S. Ali Rahman, N. Binz, C.-R. Li, N. N. Vagaja, M. de Pinho, C.-M.
Lai, Characterization of a mouse model of hyperglycemia and retinal
neovascularization. *Am J Pathol* **177**, 2659-2670 (2010).

30.　R. Nakagawa, S. Kannan, H. Altae-Tran, S. N. Takeda, A. Tomita, H. Hirano, T.
Kusakizako, T. Nishizawa, K. Yamashita, F. Zhang, H. Nishimasu, O. Nureki,
Structure and engineering of the minimal type VI CRISPR-Cas13bt3. *Molecular Cell*
82, 3178-3192.e3175 (2022).

31.　K. Nishijima, Y.-S. Ng, L. Zhong, J. Bradley, W. Schubert, N. Jo, J. Akita, S. J.
Samuelsson, G. S. Robinson, A. P. Adamis, D. T. Shima, Vascular Endothelial
Growth Factor-A Is a Survival Factor for Retinal Neurons and a Critical
Neuroprotectant during the Adaptive Response to Ischemic Injury. *Am J Pathol* **171**,
53-67 (2007).

32.　Y.-Z. Le, VEGF production and signaling in Müller glia are critical to modulating
vascular function and neuronal integrity in diabetic retinopathy and hypoxic retinal
vascular diseases. *Vision Research* **139**, 108-114 (2017).

33.　K. M. Ford, M. Saint-Geniez, T. Walshe, A. Zahr, P. A. D'Amore, Expression and
Role of VEGF in the Adult Retinal Pigment Epithelium. *Investigative Ophthalmology*
& Visual Science **52**, 9478-9487 (2011).

34.　D. Colognori, M. Trinidad, J. A. Doudna, Precise transcript targeting by CRISPR-
Csm complexes. *Nature Biotechnology* **41**, 1256-1264 (2023).

35.　N. Jo, C. Mailhos, M. Ju, E. Cheung, J. Bradley, K. Nishijima, G. S. Robinson, A. P.
Adamis, D. T. Shima, Inhibition of Platelet-Derived Growth Factor B Signaling
Enhances the Efficacy of Anti-Vascular Endothelial Growth Factor Therapy in
Multiple Models of Ocular Neovascularization. *Am J Pathol* **168**, 2036-2053 (2006).

36.　J. Park, G. Cui, H. Lee, H. Jeong, J. J. Kwak, J. Lee, S. H. Byeon, CRISPR/Cas9
mediated specific ablation of vegfa in retinal pigment epithelium efficiently regresses
choroidal neovascularization. *Sci. Rep.* **13**, 3715 (2023).

37.　M. Garita-Hernandez, F. Routet, L. Guibbal, H. Khabou, L. Toualbi, L. Riancho, S.
Reichman, J. Duebel, J. A. Sahel, O. Goureau, D. Dalkara, AAV-Mediated Gene
Delivery to 3D Retinal Organoids Derived from Human Induced Pluripotent Stem
Cells. *Int J Mol Sci* **21**, (2020).

38.　M. E. McClements, H. Steward, W. Atkin, E. A. Goode, C. Gándara, V. Chichagova,
R. E. MacLaren, Tropism of AAV Vectors in Photoreceptor-Like Cells of Human

iPSC-Derived Retinal Organoids. *Translational Vision Science & Technology* **11**, 3-3
(2022).

39. T. T. Duong, J. Lim, V. Vasireddy, T. Papp, H. Nguyen, L. Leo, J. Pan, S. Zhou, H. I.
Chen, J. Bennett, J. A. Mills, Comparative AAV-eGFP Transgene Expression Using
Vector Serotypes 1–9, 7m8, and 8b in Human Pluripotent Stem Cells, RPEs, and
Human and Rat Cortical Neurons. *Stem Cells International* **2019**, 7281912 (2019).

40. S. Eoi Jong, A. C. Jeong, K. Jae-Young, Y. Young Hee, Aflibercept ameliorates
retinal pericyte loss and restores perfusion in streptozotocin-induced diabetic mice.
BMJ Open Diabetes Research & Care **8**, e001278 (2020).

41. C. C. Tokunaga, K. P. Mitton, W. Dailey, C. Massoll, K. Roumayah, E. Guzman, N.
Tarabishy, M. Cheng, K. A. Drenser, Effects of Anti-VEGF Treatment on the
Recovery of the Developing Retina Following Oxygen-Induced Retinopathy.
Investigative Ophthalmology & Visual Science **55**, 1884-1892 (2014).

42. C. P. Shah, S. J. Garg, J. F. Vander, G. C. Brown, R. S. Kaiser, J. A. Haller,
Outcomes and Risk Factors Associated with Endophthalmitis after Intravitreal
Injection of Anti-Vascular Endothelial Growth Factor Agents. *Ophthalmology* **118**,
2028-2034 (2011).

43. A. K. W. Lai, A. C. Y. Lo, Animal Models of Diabetic Retinopathy: Summary and
Comparison. *Journal of Diabetes Research* **2013**, 106594 (2013).

44. L. E. H. Smith, W. Shen, C. Perruzzi, S. Soker, F. Kinose, X. Xu, G. Robinson, S.
Driver, J. Bischoff, B. Zhang, J. M. Schaeffer, D. R. Senger, Regulation of vascular
endothelial growth factor-dependent retinal neovascularization by insulin-like growth
factor-1 receptor. *Nat. Med.* **5**, 1390-1395 (1999).

45. X. Huang, G. Zhou, W. Wu, Y. Duan, G. Ma, J. Song, R. Xiao, L. Vandenberghe, F.
Zhang, P. A. D'Amore, H. Lei, Genome editing abrogates angiogenesis in vivo.
Nature Communications **8**, 112 (2017).

46. A. Holmgaard, A. L. Askou, J. N. E. Benckendorff, E. A. Thomsen, Y. Cai, T. Bek, J.
G. Mikkelsen, T. J. Corydon, In Vivo Knockout of the Vegfa Gene by Lentiviral
Delivery of CRISPR/Cas9 in Mouse Retinal Pigment Epithelium Cells. *Molecular
Therapy - Nucleic Acids* **9**, 89-99 (2017).

47. A. B. Holmgaard, A. L. Askou, E. G. Jensen, S. Alsing, R. O. Bak, J. G. Mikkelsen,
T. J. Corydon, Targeted Knockout of the Vegfa Gene in the Retina by
Subretinal Injection of RNP Complexes Containing Cas9 Protein and Modified
sgRNAs. *Mol. Ther.* **29**, 191-207 (2021).

48. Y. Wang, B. Wang, H. Xie, Q. Ren, X. Liu, F. Li, X. Lv, X. He, C. Cheng, R. Deng,
J. Li, J. Zhao, Z. Song, F. Gu, Efficient Human Genome Editing Using SaCas9
Ribonucleoprotein Complexes. *Biotechnology journal* **14**, e1800689 (2019).

49. S. H. Chung, T.-N. Sin, B. Dang, T. Ngo, T. Lo, D. Lent-Schochet, R. K. Meleppat,
R. J. Zawadzki, G. Yiu, CRISPR-based VEGF suppression using paired guide RNAs
for treatment of choroidal neovascularization. *Molecular Therapy - Nucleic Acids* **28**,
613-622 (2022).

50. S. Konermann, P. Lotfy, N. J. Brideau, J. Oki, M. N. Shokhirev, P. D. Hsu,
Transcriptome Engineering with RNA-Targeting Type VI-D CRISPR Effectors. *Cell*
173, 665-676.e614 (2018).

51. C.-M. Lai, Generation of transgenic mice with mild and severe retinal
neovascularisation. *Br J Ophthalmol* **89**, 911-916 (2005).

52. E. Zudaire, L. Gambardella, C. Kurcz, S. Vermeren, A Computational Tool for
Quantitative Analysis of Vascular Networks. *PLoS One* **6**, e27385 (2011).

53. S. Andrews. (Babraham Bioinformatics, Babraham Institute, Cambridge, United
Kingdom, 2010).

54. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

55. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M.
Chaisson, T. R. Gingeras, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
29, 15-21 (2013).

56. B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC bioinformatics* **12**, 1-16 (2011).

57. J.-H. Hung, T.-H. Yang, Z. Hu, Z. Weng, C. DeLisi, Gene set enrichment analysis:
performance evaluation and usage guidelines. *Briefings in bioinformatics* **13**, 281-291
(2012).

58. G. O. Consortium, The Gene Ontology (GO) database and informatics resource.
Nucleic acids research **32**, D258-D261 (2004).

59. G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R package for comparing
biological themes among gene clusters. *Omics: a journal of integrative biology* **16**,
284-287 (2012).

60. A. Gordon, G. J. Hannon, Fastx-toolkit. *FASTQ/A short-reads preprocessing tools
(unpublished) http://hannonlab. cshl. edu/fastx_toolkit* **5**, (2010).

61. J. Ye, S. McGinnis, T. L. Madden, BLAST: improvements for better sequence
analysis. *Nucleic acids research* **34**, W6-W9 (2006).

62. A. Ianevski, A. K. Giri, T. Aittokallio, Fully-automated and ultra-fast cell-type
identification using specific marker combinations from single-cell transcriptomic
data. *Nature communications* **13**, 1246 (2022).

63. G. A. Limb, T. E. Salt, P. M. Munro, S. E. Moss, P. T. Khaw, In vitro characterization
of a spontaneously immortalized human Müller cell line (MIO-M1). *Invest
Ophthalmol Vis Sci* **43**, 864-869 (2002).

64. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

65. Y. Hao, T. Stuart, M. H. Kowalski, S. Choudhary, P. Hoffman, A. Hartman, A.
Srivastava, G. Molla, S. Madad, C. Fernandez-Granda, R. Satija, Dictionary learning
for integrative, multimodal and scalable single-cell analysis. *Nature Biotechnology*,
(2023).

1011

1012

Acknowledgments

1014 The authors would like to thank MAnusha H Shah, Liuhui Huang, Jesse Gardner-Russell and

1015 Yujie Wang for their assistance with retinal dissections, Prof Alex Hewitt and Dr Leilei Tu for

1016 providing resources for vector construction. The authors also extend their gratitude to the

1017 Vector and Genome Engineering Facility (VGEF) and the Single Cell Analytics Facility at the

1018 Children's Medical Research Institute (CMRI) for producing AAV vectors and the scRNA-seq

experiments respectively and Azenta Life Sciences for conducting RNA sequencing. Schematic in figures were designed using Biorender.com.

Funding: This work was supported by grants from the National Health and Medical Research Council of Australia (NHMRC; GNT1185600), CERA Innovation Fund, National Natural Science Foundation of China (No. 82201227), and Natural Science Foundation of Guangdong Province, China (No. 2023A1515011225). Anai Gonzalez Cordero is an Al & Val Rosenstrauss Fellow (Rebecca Cooper Foundation). Funds supporting this work included the Medical Research Future Fund Stem Cell Mission (MRFF; MRF2008912), and the National Stem Cell Foundation - Metcalf Award Prize (No 00339). Satheesh Kumar was supported by the Melbourne Research Scholarship and Graeme Clark Institute Top-Up Scholarship. The Centre for Eye Research Australia receives Operational Infrastructure Support from the Victorian Government.

Author contributions:
Conceptualization: SK, GSL. Methodology: SK, YWH, VHYW, DA, LAM, AGC, BVB, GSL. Investigation: SK, YWH, VHYW, DA, AGC, BVB, GSL. Validation: SK, YWH, GSL. Formal analysis: SK, YWH, GSL. Data curation: SK, YWH, GSL. Visualization: SK, YWH, DA, AGC, GSL. Resources: LAM, AGC, BVB, GSL. Supervision: GSL. Writing—original draft: SK, GSL. Writing—review & editing: SK, YWH, AGC, BVB, GSL. Project administration: GSL. Funding acquisition: AGC, GSL

Competing interests: The authors declare that they have no known competing financial interests.

Data and materials availability: RNA sequencing data generated from this study is available under the GEO accession numbers: GSE259219 and GSE259220. All data are available in the main text or the supplementary materials. Other data generated and/or analysed in this current study will be available from the corresponding author upon reasonable request.

1048   **Supplementary Materials**

1049

1050

1051   **Figure S1. Evaluation of Cas13bt3 targeting in retinal cell lines.** VEGFA mRNA expression

1052   in ARPE19 cells following transfection of control, Cas13bt3-NT and Cas13bt3-VEGFA. A

1053   significant reduction in Cas13bt3-VEGFA was observed. Data presented as mean ± SEM. ns:

1054   not significant. *$P \leq 0.05$.

1055

1056

1057

Figure S2. HEK293FT DEG and off target analysis. (A) Number of differentially expressed

genes from the VEGFA-targeting shRNA, CasRx and Cas13bt3 transfections in HEK293FT

cells compared to vector control. **(B)** Volcano plots of differentially expressed genes from

VEGFA shRNA, **(C)** CasRx VEGFA sgRNA, **(D)** Cas13bt3 NT sgRNA and **(E)** Cas13bt3

VEGFA sgRNA compared to vector control. **(F)** Plot of TPM values for VEGFA and candidate

off-target genes from Cas13bt3-NT sgRNA and Cas13bt3 VEGFA sgRNA groups compared

to shRNA and CasRx targeting vectors. Statistical analysis was conducted by Welch's t test

(F). Data presented as mean ± SEM. ns: not significant. *P≤0.05, ****P≤0.00001.

A

Cas13bt3-crRNA complex   Cas13bt3-crRNA-target complex

HEPN1

IDL1

IDL2   HEPN2

REC lobe

Activation   Cleavage

B

Target   GAAGGAGGAGGGCAGAAUCAUCACGAAGUGGUGAAGU**UC**AUGGAUG**UC**UA
VEGFA presgRNA 2   CUUCCUCCUCCCGUCUUAGUAGUGCUUCAC

Target   CGGAUCAAACCUCACCAAGGCCAGCACAUAGGAGAGAUGAGC**UC**CUACA
VEGFA presgRNA1   GCCUAGUUUGGAGUGGUUCCGGUCGUGUAU

VEGFA mRNA

| Exon 1 | Exon 2 | Exon 3 | Exon 4 | Exon 5 | Exon 6 | Exon 7 | Exon 8 |

VEGFA presgRNA 3   AAGGAUGUCGUGUUGUUUACAUUACGUCU
Target   UUCCUACAGCACAACAAAUGUGAAUGCAGACCAAAGAAAGAUAGAGCAAGACAAGAAAAAAAAUCAGU**UC**

C

PRKAR2B AUUUCACCAAUAGGCCAAGGCCAGCACAUAACUUA**UC**AAGCUAGAAAACUUUAAAACAAAAAUGAUUAAA
         --------U-------UUCCGGUCGUGUAU

EHD4 UGGAUCAAACCUCACCUGAAGUCCUUCCUAU**UC**CUACUAGAUAUUUUUU**UC**CACUGUGUGGGCCAAUGCA
      -CCUAGUUUGGAGUG------------AU

USP22 CGAUGCCAUCAUCACCAAGGCCAGCAUCAAGGACG**UC**CUGGACAGCGAAGGGUACUUGCUGU**UC**UA**UC**AC
       GC---G-U-G-AGUGGUUCCGGUCGU---U

RHBDL1 GGAGCUCAUGCACACCAAGGCCAGCACCAUC**UC**AGCAACUUGUAGGGACA**UC**UCA**UC**CCAGCCCAGU**UC**
        -C-----U--G-GUGGUUCCGGUCGUG---

POMT2 GGACAGCAGCGUCACCAAGGCCAGCAGGGCCCACCAGCCGACCGCC**UC**GAAGCGCCGUGAGCCCCAAGCA
       -C-------G-AGUGGUUCCGGUCGU----

SNX33 AAAAUCAAACCUCACCACCCCCCGACAUAGC**UC**AGAAGGGCUGAUUGUGAAACCCAGC**UC**UGU**UC**AACAU
       ---UAGUUUGGAGUGGU---GG-C------

RER1 CAGAGAGAAACCAACCAAGGCCAGCACCUCC**UC**GGAAGCAUUUGGCACACACACCUU**UC**AAUACACG**UC**A
      G-CU---UU-G--UGGUUCCGGUCGUG-A-

POLA1 UGGAUCAAACCUCACUGCAAGUCAGAGGGCCUAUGCGCCUGAGCAGCUGCAGAAACAGGAUAA**UC**UAACC
       -CCUAGUUUGGAGUG----------U----

SLC9B2 CCUUUGAAUCCAGACCAAGGCCAGCACGAACCAGAAUGAUAGACAGGGCUAUGCU**UC**UC**AAAGAGGAAGA
        G---A-UU-GG--UGGUUCCGGUCGUG--U

SYT10 AGAGCCAAACCUCACCAAGAAAGUAAUUUUCUGAGAAUGAAAGAAAAGUAUUA**UC**AGAUAAGUAACAUUA
       -C---GUUUGGAGUGGUUC------U--A-

UC: Cas13bt3 cut site

1066

1067 **Figure S3. Prediction of Cas13bt3 cut sites**. **(A)** Schematic of Cas13bt3-gRNA and Cas13bt3-gRNA-target complex. **(B)** Potential Cas13bt3 cut

1068 sites on VEGFA mRNA target for VEGFA sgRNA1, 2 and 3. **(C)** Potential Cas13bt3 cut sites on top 10 candidate off-target genes for VEFGA

1069 sgRNA 1.

1070

A VEGFA T A T G T G C T G G C C T T G G T G A G G T T T G A T C C G

PRKAR2B T A T G T G C T G G C C T T G G – – – – – – – – – – – – –

EHD4 – – – – – – – – – – – – G G T G A G G T T T G A T C C –

USP22 – – – – T G C T G G C C T T G G T G A – – – – – – – – – –

RHBDL1 – – – G T G C T G G C C T T G G T G – – – – – – – – – – –

POMT2 – – – – T G C T G G C C T T G G T G A – – – – – – – – – –

SNX33 – – – – – – – – – – – T G G T G A G G T T T G A T – – –

RER1 – – – G T G C T G G C C T T G G T – – – – – – – – – – –

POLA1 – – – – – – – – – – – – – G T G A G G T T T G A T C C –

SLC9B2 – – – G T G C T G G C C T T G G T – – – – – – – – – – –

SYT10 – – – – – – – – – – C T T G G T G A G G T T T G – – – – –

B VEGFA presgRNA 2 C A C T T C G T G A T G A T T C T G C C C T C C T C C T T C

BARX1 – – – – – – – – – – – – – – T C T G C C C T C C T C C T T C

SLX4 – – – – – – – – – – – – – – T C T G C C C T C C T C C T T C

LOC 100192426 – – – – – – – – – – – – – – T C T G C C C T C C T C C T T C

NRG1 – – – – – – – G A T G A T T C T G C C C T C – – – – – – –

AGRN – – – – – – – – – – A T T C T G C C C T C C T C C – – –

TRIM56 – – – – – – – – – – – – – C T G C C C T C C T C C T T C

CDH20 – – – – – – – – – – G A T T C T G C C C T C C T C C T T C

MORC3 – – – – – – – – T G A T T C T G C C C T C C T – – – – –

ZNF32-AS3 – – – – – – – – – – – – – C T G C C C T C C T C C T T C

SUGT1P4-STRA6LP C A C T T C G T G A T G A T – – – – – – – – – – – – – – –

C VEGFA presgRNA 3 T C T G C A T T C A C A T T T G T T G T G C T G T A G G A A

NUP210L – – – – – – – – – C A T T T G T T G T G C T G T – – – – –

LOC 105373744 T C T G C A T T C A C A T T T – – – – – – – – – – – – – –

LYSMD3 – – – – – – – – – C A T T T G T T G T G C T G T – – – – –

SKP1 – C T G C A T T C A C A T T T G – – – – – – – – – – – – –

CPA4 – – T G C A T T C A C A T T T G – – – – – – – – – – – – –

SENP5 – C T G C A T T C A C A T T T G T T – – – – – – – – – – –

1071

1072

1073 **Figure S4. Multiplexed VEGFA knockdown using Cas13bt3. (A)** Top off-target genes for

1074 VEGFA sgRNA 1, **(B)** VEGFA sgRNA 2 and **(C)** VEGFA sgRNA 3.

1075

1076

Figure S5. Codon optimisation of Cas13bt3. (A) Schematic of codon optimisation procedure
from Cas13e protein sequence. **(B)** VEGFA mRNA knockdown from original and codon
optimised Cas13e compared to control transfection. **(C and D)** Expression of original and
codon optimised determined from Western blot. Statistical analyses was conducted using One-
way ANOVA with Tukey's multiple comparisons test (**B and C**). Data presented as mean ±
SEM. *P≤0.05, **P≤0.001, ****P≤0.00001.

1084

1085

1086

Figure S6. Characterization of 30 weeks retinal organoids. (A) Retinal organoids showed

expression of mature photoreceptor markers Arrestin-3 (magenta) specific to the cone cells and

Rhodopsin (cyan) localized to the outer segments (OS) of rod photoreceptor cells. **(B, C)** In

the inner nuclear layer (INL) bipolar cells stained with VSX2 (cyan) and Müller glia stained

with CRALBP (magenta) **(B)**, Amacrine and horizontal cells stained with PROX1 (magenta)

and ganglion cells stained with HuCHuD (cyan) **(C)**. **(D)** Brightfield image of retinal organoid

section showing the outer nuclear layer (ONL) and a pigmented RPE island next to it. Scale

bar: 116μm. Abbreviations: INL: inner nuclear layer ONL: outer Nuclear Layer, OS: outer

segment, RPE: retinal pigment epithelium.

1096

1097

Figure S7. Dot plots of retinal cell marker genes for respective cell types in 3D retinal organoids. Complete list of cell marker genes can

obtained from Kim et al., 2023 (*27*).

Figure S8. *E*xpression of VEGFA *on* retinal cell clusters. UMAP plots of showing

expression on **(A)** Untreated retinal organoid. **(B)** Violin plots showing expression of VEGFA

in individual retinal cell types. **(C)** UMAP plots of retinal cell clusters with expression of

VEGFA from VEGFA-treated and NT-treated retinal organoids, all cells, **(D)** Cas13bt3

positive cells and **(E)** Cas13bt3 negative cells. Data analyses were conducted using One-way

ANOVA with Tukey's multiple comparisons test (B). **P≤0.001, ****P≤0.00001.

Cas13bt3 VEGFA sgRNA vs Cas13bt3 NT sgRNA

1109 **Figure S9. Differentially expressed genes in Cas13bt3-VEGFA against Cas13bt3-NT.**
1110 Volcano plots from scRNA-seq for each retinal cell type showing differentially expressed
1111 genes between retinal organoid infected with Cas13bt3-VEGFA sgRNA and Cas13bt3-NT
1112 sgRNA. Significant downregulation of VEGFA in RPE cells is highlighted in red.

1113

1114

1115 **Figure S10. Cas13bt3-NT sgRNA treatment in retinal organoids. (A)** Expression of
1116 Cas13bt3 by number of reads across retinal cell types in NT sgRNA treatment group. **(B)** Violin
1117 plots of VEGFA expression in cells with low and high expression of Cas13bt3 against
1118 Cas13bt3 negative cells showed no significant differences.

1119

1120

1121 **Figure S11. Expression of Cas13bt3 plotted against expression of VEGFA mRNA across**

1122 **cell types in NT sgRNA and VEGFA sgRNA groups.**

1125   **Figure S12. Correlation plots of Cas13bt3 with VEGFA expression in the various retinal cell types for the predicted off-target genes of**

1126   **VEGFA sgRNA.**

1127

1128 **Figure S13. Intravitreal delivery of AAV2.7m8 achieves strong transduction of mice**
1129 **photoreceptors. (A)** Schematic of AAV2.7m8 virus carrying mCherry reporter. **(B)** Schematic
1130 of experimental procedure for intravitreal injection of C57J/BL6 mice eyes. **(C)** Confocal
1131 microscope images of retinal flatmount 4 weeks post intravitreal injection, showing
1132 transduction of ganglion, bipolar, cone and rod cells with reference image of retinal cell layers.

1133

1134

1135

1136

A

VEGFA presgRNA	T A T G T G C T G G C C T T G G T G A G G T T T G A T C C G
Mok	– – – – – – – – – – – – GGTGAGGTTTGAT – – –
Olfml2b	– – – – – – – – – – CCTTGGTGAGGTT – – – – – –
Ehmt1	– – – – – – – TGGCCTTGGTGAG – – – – – – – –
Tbl2	– – – TGCTGGCCTTGGTG – – – – – – – – – –
Slc4a8	– – – GTGCTGGCCTTGGT – – – – – – – – –
Dcp1b	– – – – – – TGGCCTTGGTGAG – – – – – – –
Usp22	– – – – – TGCTGGCCTTGGTGA – – – – – – – –
Thbs2	– – – TGTGCTGGCCTTGG – – – – – – – – – –
Pdzd2	– – – – – GCTGGCCTTGGTG – – – – – – – – –
Lrba	– – – – – – – – GGCCTTGGTGAGG – – – – – – –

B NT vs Control

−Log10(p−value) vs Log2FC

Lrba, Tbl2, Dcp1b, Ehmt1, Usp22, Slc4a8, mVegfa, Pdzd2, Mok, Thbs2, Olfml2b

C VEGFA vs Control

−Log10(p−value) vs Log2FC

Usp22, Slc4a8, Lrba, Tbl2, Dcp1b, Ehmt1, Thbs2, Pdzd2, Olfml2b, Mok, mVegfa

D

ns

TPM

Control | NT sgRNA | VEGFA sgRNA

mVEGFA, Mok, Olfml2b, Ehmt1, Tbl2, Slc4a8, Dcp1b, Usp22, Thbs2, Pdzd2, Lrba

1137

1138 **Figure S14. Cas13bt3 off-target effects in Kimba mice. (A)** Schematic of candidate off-
1139 target genes to VEGFA sgRNA in mouse transcriptome. Volcano plots of differentially
1140 expressed genes from Kimba mice retinas treated with AAV2.7m8 carrying **(B)** Cas13bt3-
1141 NTsgRNA and **(C)** Cas13bt3-VEGFAsgRNA compared to vector controls. **(D)** Expression in
1142 TPM of mouse VEGFA and candidate off-target genes from RNA-seq.

1143

1144

1145

1146 **Figure S15. Original fundus fluorescein angiography images from Kimba mice retinae.**

1147

1148

1149

1150

1151 **Figure S16. Imaging of retinal vascular plexuses Kimba mice treated with Cas13bt3-**
1152 **NTsgRNA and Cas13bt3-VEGFAsgRNA. (A)** Schematic of experimental procedure for
1153 imaging retinal vascular plexuses. **(B)** Schematic of the different retinal plexuses. **(C)** Confocal
1154 imaging of the retinal plexuses, 8 weeks after Cas13bt3 treatment, immunolabelled with
1155 laminin. Magnification 40X, scale bar: 50μm. White arrows indicate leaky vessel structure.

1156

1157

Figure S17. Retinal thickness readings from optical coherence tomography (OCT) in
Kimba mice. (A) Early Treatment of Diabetic Retinopathy Study (ETDRS) grid showing
segments of the retina analysed using OCT. C:Central (optic nerve); N1: Inner nasal; N2: Outer
Nasal; T1: Inner temporal; T2: Outer temporal; S1: inner superior; S2: Outer superior; I1: Inner
inferior; I2: Outer inferior. **(B)** Vessel density and **(C)** number of junctions readings from FFA
images of Kimba mice at baseline compared to wildtype (C57BL/6) mice. **(D)** Plot of
difference in total retinal thickness in untreated and treated Kimba mice between baseline and
endpoint. **(E)** Representative OCT images from untreated and treated Kimba mice at endpoint.

1168

1169

Reagent or Resource	Source	Identifier
Chemicals		
DMEM	Thermo Fisher Scientific	Cat# 11965092
DMEM/F12	Thermo Fisher Scientific	Cat# 11320033
Opti-MEM	Thermo Fisher Scientific	Cat# 31985070
TrypLE™ Express	Thermo Fisher Scientific	Cat# 12605093
GlutaMAX™ Supplement	Thermo Fisher Scientific	Cat# 35050061
FBS	Bovogen Biologicals	SFBS-AU
Antibiotic-Antimycotic	Thermo Fisher Scientific	Cat# 15240062
DPBS	Thermo Fisher Scientific	Cat# 14190250
Lipofectamine™ 3000 Transfection Reagent	Thermo Fisher Scientific	Cat# L3000001
ViaFect™ Transfection Reagent	Promega Corporation	Cat# E4982
NEB® Stable Competent E. coli, High Efficiency	New England Biolabs	Cat# C3040I
rCutSmart™ Buffer	New England Biolabs	Cat# B6004S
NEB® 10-beta/Stable Outgrowth Medium	New England Biolabs	Cat# B9035S
NucBlue™ Live ReadyProbes™ Reagent (Hoechst 33342)	Thermo Fisher Scientific	Cat# R37605
LB (Lennox L) agar	Thermo Fisher Scientific	Cat# 22700025
2-YT Broth	Thermo Fisher Scientific	Cat# 22712020
Carbenicillin Disodium Salt	Thermo Fisher Scientific	Cat# 10177012
GelRed® nucleic acid gel stain	Biotium	Cat# 41003
Phosphate buffered saline	Merck	Cat# P3813
Tween-20	Merck	Cat# P1379
Triton-X-100	Amresco	Cat# 0694
RIPA Lysis and Extraction Buffer	Thermo Fisher Scientific	Cat# 89900

Pierce™ Phosphatase Inhibitor Mini Tablets	Thermo Fisher Scientific	Cat# A32957
NuPAGE™ MES SDS Running Buffer (20X)	Thermo Fisher Scientific	Cat# NP0002
NuPAGE™ LDS Sample Buffer (4X)	Thermo Fisher Scientific	Cat# NP0007
0.5% Proxymethacaine hydrochloride (Alcaine™)	Alcon	N/A
0.5% Tropicamide (Mydriacyl)	Alcon	N/A
GenTeal™ Gel	Alcon	N/A
Ketamine, 100 mg/mL	Troy Laboratories	Ketamil
Xylazine, 100 mg/mL	Troy Laboratories	Ilium Xylazil-100
Experimental models: Cell lines		
Human: HEK293FT	Thermo Fisher Scientific	R70007
Human: ARPE-19	ATCC	CRL-2302™
Human: Müller 1 (MIO-M1)	UCL Institute of Ophthalmology, London	(63)
Experimental models: Organisms/ strains		
Human: 3D retinal organoid	Children's Medical Research Institute	(27)
Kimba Mouse	The University of Western Australia	(29)
Antibodies		
Mouse anti-actin, clone C4	Merck	MAB1501
Rabbit anti-HA, C29F4	Cell Signaling Technology	#3724
Mouse anti-rhodopsin, 1D4	Abcam	ab5417
Rat mCherry monoclonal 16D7, Alexa Fluor™ 594	Thermo Fisher Scientific	Cat# M11240
Rat anti-laminin monoclonal, α-2 chain	Merck	L0663
Donkey anti-rat AffiniPure Fragment IgG (H + L), Alexa Fluor™ 647	Jackson ImmunoResearch	712-606-150

Donkey mouse AffiniPure Fragment IgG (H + L), Alexa Fluor™	Jackson ImmunoResearch	715-546-151
Goat anti-rabbit IgG (H+L) Secondary Antibody, HRP	Thermo Fisher Scientific	Cat# 31460
Goat anti-Mouse IgG (H+L) Secondary Antibody, HRP	Thermo Fisher Scientific	Cat# 62-6520
Anti-Arrestin C antibody, C-terminal	Abcam	189437
Monoclonal Anti-Opsin antibody produced in mouse	Merck	O4886
Sheep anti Human chx 10 (Visual system homeobox 2) (NT)	ExAlpha	XII80P
CRALBP Monoclonal Antibody (B2)	Thermo Fisher Scientific	MA1-813
Anti-Prox 1 Antibody	Merck	AB5475
HuC/HuD Monoclonal Antibody (16A11)	Thermo Fisher Scientific	A21271
Commercial kits		
RNeasy Mini kit	Qiagen	Cat# 74106
Monarch® Total RNA Miniprep Kit	New England Biolabs	Cat# T2010S
High-Capacity cDNA Reverse Transcription Kit	Thermo Fisher Scientific	Cat# 4368814
Q5® Hot Start High-Fidelity Master Mix	New England Biolabs	Cat# M0494S
TaqMan™ Fast Advanced Master Mix	Thermo Fisher Scientific	Cat# 4444557
SYBR™ Green PCR Master Mix	Thermo Fisher Scientific	Cat# 4309155
Human VEGF DuoSet ELISA	R & D systems	Cat# DY293B
Recombinant DNA		
pAAV.CMV-CasRx mCh	Addgene	203442
pAAV.CMV-CasRx-VEGFA sgRNA	Addgene	203443
pAAV.CMV-CasRx-VEGFA sgRNA array	Addgene	203444
pAAV.CMV-Cas13bt3-NT sgRNA	Addgene	203446

pAAV.CMV-Cas13bt3-VEGFA sgRNA	Addgene	203447
pAAV.CMV-Cas13bt3-VEGFA sgRNA array	Addgene	203448
pAAV.CMV-Cas13bt3 (GenScript CO)-NT sgRNA	Addgene	203451
pAAV.CMV-Cas13bt3 (GenScript CO)-VEGFA sgRNA	Addgene	203452
Software and algorithms		
GraphPad Prism 10	GraphPad	https://www.graphpad.com/scientific-software/prism/E/
Benchling	Benchling	https://www.benchling.com/
Fiji-ImageJ	National Institutes of Health	https://imagej.net/software/fiji/
GenSmart™ Codon Optimisation	GenScript	https://www.genscript.com/gensmart-free-gene-codon-optimization.html
IDT Codon Optimisation Tool	Integrated DNA Technologies	https://sg.idtdna.com/pages/tools/codon-optimization-tool
STAR (2.7.11a)	(*55*)	https://github.com/alexdobin/STAR
DESeq2 (v3.18)	(*64*)	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
Seurat (v5)	(*65*)	https://satijalab.org/seurat/
Cell Ranger (v7.2)	10X Genomics	https://www.10xgenomics.com/support/software/cell-ranger
ImageLab (v6.1)	Bio-Rad	https://www.bio-rad.com/en-au/product/image-lab-software
R (v4.3.2)	R Core Team	https://cran.r-project.org/
RStudio (v2023.09.1)	Posit PBC	https://posit.co/download/rstudio-desktop/

Adobe Photoshop (v25)	Adobe Inc.	https://www.adobe.com/au/products/photoshop.html
AngioTool (v0.5)	National Cancer Institute	https://ccrod.cancer.gov/confluence/display/ROB2/Downloads
Equipment		
HRA + OCT Spectralis	Heidelberg Engineering	N/A
Leica Stellaris 5 confocal microscope	Leica Microsystems	N/A

1170

1171   **Table S1.**

1172   Key resources.

1173

Quantitative PCR

Name		Source	Identifier
Human	**VEGFA**	Thermo Fisher Scientific	Cat# 4331182
(Hs00900055_m1)			
Mouse	**VEGFA**	Thermo Fisher Scientific	Cat# 4331182
(Mm00437306_m1)			
Human	**GAPDH**	Thermo Fisher Scientific	Cat# 4448489
(Hs02786624_g1)			
Mouse	**GAPDH**	Thermo Fisher Scientific	Cat# 4448489
(Mm99999915_g1)			

Standard PCR

Name	Sequence (5' - 3')	Source
Cas13bt3_FWD	CGGCGAACAGGGTGATAAA	Integrated DNA Technologies
Cas13bt3_REV	CTGTGCTGGGCTTCAAGATA	Integrated DNA Technologies

1174

1175 **Table S2.**

1176 Primers used in this study.

1177

1178

1179