Enhancing Bone Regeneration and Osseointegration using rhPTH(1-34) and Dimeric [R25C]PTH(1-34) in an Osteoporotic Beagle Model

Jeong-Oh Shin[1*], Jong-Bin Lee[2*], Sihoon Lee[3], Jin-Woo Kim[4]

[1] Department of Anatomy, Soon Chun Hyang University College of Medicine, Cheonan, Korea

[2] Department of Periodontology and Research Institute of Oral Sciences, College of Dentistry, Gangneung-Wonju National University, Gangneung, Republic of Korea

[3] Department of Internal Medicine and Laboratory of Genomics and Translational Medicine, Gachon University College of Medicine, Incheon, Korea

[4] Department of Oral and Maxillofacial Surgery, Research Institute for Intractable Osteonecrosis of the Jaw, College of Medicine, Ewha Womans University, Seoul, Korea

* These authors contributed equally.

Corresponding author

Sihoon Lee, *MD, PhD*
Professor, Department of Internal Medicine and Laboratory of Genomics and Translational Medicine,
Gachon University College of Medicine, Incheon, Korea
Tel: +82-32-458-2646, Fax: 82-32-460-2381
Email: shleemd@gachon.ac.kr

Jin-Woo Kim, *DDS, MSD, PhD, FIBCSOMS*
Professor, Department of Oral and Maxillofacial Surgery
School of Medicine, Ewha Womans University
Anyangcheon-ro 1071, Yangcheon-gu, Seoul, 158-710, Korea
Tel: 82-2-2650-2720, Fax: 82-2-2650-2754
E-mail: jinu600@gmail.com, jwkim84@ewha.ac.kr

Abstract

Introduction: Despite continuous parathyroid hormone (PTH) exposure potentially accelerating bone resorption, intermittent PTH administration has shown anabolic effects on bone microarchitecture. This study investigates the therapeutic impacts of two PTH analogs, rhPTH(1-34) and dimeric ^{R25C}PTH(1-34) on bone regeneration and osseointegration in a postmenopausal osteoporosis animal model.

Methods: Twelve female beagles, osteoporotic post-ovariectomy, underwent implant surgeries. Animals were divided into three groups: control, daily rhPTH(1-34) injection, and daily dimeric ^{R25C}PTH(1-34) injection. After 10 weeks, bone regeneration and implant osseointegration were evaluated using micro-CT, histological/histomorphometric analyses, and serum biochemical analysis.

Results: While the rhPTH(1-34) group demonstrated improved microarchitectural characteristics, such as BMD, BV, trabecular metrics, and osseointegration degree, the dimeric ^{R25C}PTH(1-34) group showed similarly enhanced anabolic effects around the titanium implants, albeit these were less pronounced than the rhPTH(1-34), yet significantly better than the control group. Histologic and TRAP assays revealed that both PTH analogs significantly promoted bone regeneration and remodeling, especially in artificially created bone defects.

Conclusion: This study demonstrated the therapeutic effects of rhPTH(1-34) and dimeric ^{R25C}PTH(1-34) on bone regeneration and titanium osseointegration in a beagle model with osteoporosis.

Keywords: parathyroid hormone (PTH); mutation; teriparatide; anabolic; bone regeneration

Introduction

Bone regeneration is a complex physiological process that is essential for the successful integration of dental implants into the jawbone. Osseointegration, the direct structural and functional connection between living bone and an implant's surface, is a critical determinant of implant stability and long-term viability in dental reconstructive therapy. Since excessive levels of circulating parathyroid hormone (PTH) increase osteoclastic activity and accelerate bone resorption, it might seem paradoxical that PTH can also be used as a treatment modality for diseases with bone loss, such as osteoporosis (*Jilka, 2007*). Administration of prathyroid hormone (PTH) analogs can be categorized into two distinct protocols: intermittent and continuous. Intermittent rhPTH(1-34) therapy, typically characterized by daily injections, is clinically used to enhance bone formation and strength. This method leverages the anabolic effects of rhPTH(1-34) without significant bone resorption, which can occur with more frequent or continuous exposure. On the other hand, continuous rhPTH(1-34) exposure, often modeled in research as constant infusion, tends to accelerate bone resorption activities, potentially leading to bone loss (Silva and Bilezikian, 2015; Jilka, 2007). Understanding these differences is crucial for interpreting the therapeutic implications of rhPTH(1-34) in bone health. Intermittent administration of PTH unlikely continuous exposure showed anabolic effects, indicating different responses relating to bone microarchitecture depending on the dose and frequency (*Silva and Bilezikian, 2015*). However, the underlying mechanisms remain largely unknown, although it was shown that PTH binds through PTH type 1 receptor (PTH1R) and that G-protein coupled receptors (GPCR) are associated with the protein kinase A (PKA)-dependent pathway, thereby demonstrating primary anabolic action on bone (*Cheloha et al., 2015; Jilka, 2007*). The anabolic effect of intermittent PTH administration is mediated by the downregulation of the Wnt/beta-catenin signaling pathway, which upregulates the transcriptional expression of growth factors, such as IGF1, FGF2, and Runx2, which are essential for the proliferation and differentiation of osteoblasts, leading to an increased number of osteoblasts and survival (*Krishnan et al., 2006; Lee et al., 2009*).

A PTH analog, teriparatide [recombinant human PTH(1-34){rhPTH(1-34)}], demonstrated its promising anabolic effects in a fracture prevention trial (FPT) (*Neer et al., 2001*), leading to its

approval by the United States Food and Drugs Administration, making it the first anabolic agent for

postmenopausal osteoporotic women. Additionally, its unique anabolic features, which are contrasted

with antiresorptive, lead to an increased application of rhPTH(1-34), whereby it was used in both

metabolic and pathologic bone diseases alongside various other conditions where bone formation

occurred (*Uusi-Rasi et al., 2005*). Conversely, recent concerns regarding the development of

osteonecrosis of the jaw (ONJ) have appeared in association with the use of antiresorptive, meaning

rhPTH(1-34) has gained attention for its potential to reduce the risk of ONJ and its therapeutic effects

in treating ONJ (*Jung et al., 2017; Kakehashi et al., 2015*). Although in vivo studies on fracture

healing, bone augmentation, and titanium osseointegration effects of rhPTH(1-34) have been

attempted, they have generally been limited to pilot studies in rodents (*Gomes-Ferreira et al., 2020;*

Jung et al., 2021; Yu and Su, 2020).

Interestingly, only nine mutations have been discovered since the PTH amino acid and nucleotide

sequences were confirmed (*Lee et al., 2020; Schipani et al., 1999*). Among them, the PTH R25C

mutation was discovered in three siblings with familial idiopathic hypoparathyroidism, which

consisted of a homozygous arginine to cysteine mutation at residue 25 (R25C) in the mature PTH(1-

84) polypeptide and exhibited distinct characteristics from the others (*Lee and Lee, 2022*). The other

mutations located in the prepro-leader region of the hormone resulted in defective synthesis and

secretion; however, the PTH R25C mutation is located within the mature bioactive domain of PTH

and does not affect synthesis or secretion (*Lee and Lee, 2022*). Although the capacity of [R25C]PTH(1-

34) to bind to the PTH1R and stimulate cAMP production was slightly lower in the human osteoblast-

derived SaOS-2 cell line (*Lee and Lee, 2022*), yet it showed comparable anabolic activity in a mouse

model (*Bae et al., 2016*). Furthermore, the dimeric formation of the [R25C]PTH(1-34) peptide,

presumably through disulfide bonding of the cysteine residues, has implied that dimeric [R25C]PTH(1-34)

might partake in unique biological actions, which potentiate its clinical applications (*Park et al.,*

2021).

In addition to the limited effectiveness toward certain types of fractures, such as non-vertebral

fractures, and the disadvantage of its limited duration of use, the administration of rhPTH(1-34) for

osteoporosis has raised concerns about its potential to induce cortical porosity, despite showing

favorable results in the treatment of trabecular microarchitecture (*Lindsay et al., 2016*). This has led to

concerns regarding the widespread use of PTH. However, recent notable studies have indicated that

cortical porosity is not induced when PTH is administered weekly (*Mosekilde et al., 1995*), which is

in contrast to daily administrations. (*Yamamoto et al., 2016; Zebaze et al., 2017*) These studies

suggest that the frequency and dosage of PTH administration can significantly affect the bone

response (*Hock et al., 1992*), while different forms of PTH might also induce different biological

responses, which means deeper investigations into the therapeutic effects of PTH are

required (*Bellido et al., 2005*).

Therefore, in this study, the authors used a large animal model that mimics postmenopausal

osteoporosis to investigate the therapeutic effects of two PTH analogs, rhPTH(1-34) and dimeric

^{R25C}PTH(1-34), on bone regeneration and osseointegration.

Results

Microarchitectural and histological analysis of titanium osseointegration

To investigate evaluating and comparing the efficacy of rhPTH(1-34) and the dimeric ^{R25C}PTH(1-

34) in promoting bone regeneration and healing in a clinically relevant animal model. In our study,

beagle dogs were selected as the model due to their anatomical similarity to human oral structures,

suitable size for surgeries, human-like bone turnover rates, and established oral health profiles,

ensuring comparable and ethically sound research outcomes. The normal saline injected-control group,

injected with 40ug/day PTH (Forsteo, Eli Lilly) group, and 40ug/day dimeric ^{R25C}PTH(1-34) -injected

group. Animals in each group were injected subcutaneously for 10 weeks.

Various characteristics of the right mandible were evaluated in this analysis, including bone

mineral density (BMD), bone volume (BV), trabecular number (Tb.N), trabecular thickness (Tb.Th),

and trabecular separation (Tb.S) (Figure 1). The group administered rhPTH(1-34) presented

consistently higher values of BMD, BV, Tb.N, and Tb.Th,, and lower value of Tb.S compared to the

control and dimeric [R25C]PTH(1-34) groups, indicating that rhPTH(1-34) administration enhanced

titanium osseointegration (Figure 1E). Moreover, this was consistent for all three titanium implants,

irrespective of artificial bone defects (second implant) or bone grafting (third implant). Interestingly,

the dimeric [R25C]PTH(1-34) group showed similar trends for the anabolic effects related to the titanium

implants. Morphometric analysis indicated that dimeric [R25C]PTH(1-34) administration enhanced the

BMD, BV, and other trabecular indices, resulting in a higher degree of osseointegration than in the

control group (P < 0.05), although lower than in the rhPTH(1-34) group (P > 0.05) (Figure 1E).

Histological analysis further clarified the micro-CT results. The implants in the control group

showed physiological bone osseointegration around the titanium implant; however, insufficient bone–

implant contact and exposure of implant threads were observed (Figure 2). This was especially

evident on the buccal side of the implant, which was more vulnerable due to the bundle bone structure.

Alternatively, the titanium implants in the rhPTH(1-34) group were in full contact with the green-

stained mineralized bone. The dimeric [R25C]PTH(1-34)group showed a better pattern of

osseointegration compared to the control group, although the bone–implant contact was lower than in

the rhPTH(1-34) group (Figure 2A-C). Notably, both the rhPTH(1-34) and dimeric [R25C]PTH(1-34)

groups presented evidence of bone regeneration for the second implant, whereby a bone defect was

created prior to placing the implant. However, both were insufficient compared to the control group

and did not illustrate any bone filling. The measured bone–implant contact ratio was $18.32 \pm 16.19\%$

for the control group, $48.13 \pm 29.81\%$ for the group, and $39.53 \pm 26.17\%$ (P < 0.05).

Histological and TRAP analyses of bone regeneration

Artificial bilateral bone defects were created in the left mandible and the effects of the two PTH

analogs were evaluated on the bone regeneration, with one left unfilled and the other filled with a

bone graft. Figure 3 demonstrates the effects of rhPTH(1-34) and dimeric [R25C]PTH(1-34) on bone

regeneration compared to the control group (Figure 3A-L). Following the formation of the bone

defects, the rhPTH(1-34) and dimeric [R25C]PTH(1-34)groups achieved sufficient morphological bone

regeneration over a period of 10 weeks, while the control group exhibited morphological

incompleteness over the same period. The rhPTH(1-34) group exhibited a mature trabecularcfghnc

architecture, while the dimeric ^{R25C}PTH(1-34)group showed a similar morphology, although some

immature bone formation remained stained blue in the Masson trichrome staining analysis (Figure

3A-L). While there was no clear difference in the bone defect between the sites with and without bone

grafting (Figure 3E,F,K,L), the site where bone grafting occurred exhibited a more mature bone

morphology, indicating that a xenograft-maintained space for new bone formation with

osteoconductive effects.

The capability of rhPTH(1-34) and dimeric ^{R25C}PTH(1-34) in bone remodeling was evaluated by

tartrate-resistant acid phosphatase (TRAP) immunohistochemical staining. (Figure 4) Both the

rhPTH(1-34) and dimeric ^{R25C}PTH(1-34) groups showed a significantly higher number of TRAP+

cells at both bone defects, with and without a xenograft, compared to the control group (Figure 4M,N).

($P < 0.05$) In addition, the number of TRAP+ cells in the dimeric ^{R25C}PTH(1-34)group was

significantly higher than in the vehicle, yet lower than in the rhPTH(1-34) group (Figure 4M,N).

Serum biochemical analysis

The levels of calcium, phosphorus, CTX, and P1NP were analyzed over time using RM-ANOVA

(Figure 5). There were no significant differences between the groups for calcium and phosphorus at

time points T0 and T1 (Figure 5A). However, after the PTH analog was administered at T2 (Figure

5A), the levels were highest in the rhPTH(1-34) group, followed by the dimeric ^{R25C}PTH(1-34) group,

and then, lowest in the control group, which was statistically significant (Figure 5B,C). ($P < 0.05$) The

differences between the groups over time for CTX and P1NP were not statistically significant (Figure

5D, E).

Discussion

This study investigated the therapeutic effects of rhPTH(1-34) and dimeric ^{R25C}PTH(1-34) on bone

regeneration and osseointegration in a large animal model with postmenopausal osteoporosis.

rhPTH(1-34) and dimeric ^{R25C}PTH(1-34) have shown significant clinical efficacy, and although there

have been a few studies investigating their effects on bone regeneration in rodents (*Garcia et al.,*

2013), the authors in this study aimed to investigate the effects using a large animal model that more

accurately mimics osteoporotic humans (*Cortet, 2011*). In the evaluation of titanium osseointegration,

the rhPTH(1-34) group consistently exhibited enhanced bone mineral density (BMD), bone volume

(BV), and other key parameters, thereby indicating superior titanium osseointegration compared to the

control and dimeric ^{R25C}PTH(1-34) groups (Figure 1E). Histological analyses confirmed these results,

emphasizing the stronger bone–implant contact observed in the rhPTH(1-34) group. Furthermore,

both PTH analogs significantly promoted bone regeneration in artificially created defects, with the

rhPTH(1-34) group displaying a more mature trabecular architecture, as evidenced by a notable

increase in the TRAP+ cell count during the bone remodeling assessments (Figure 2M,N).

Furthermore, by demonstrating that dimeric ^{R25C}PTH(1-34) exhibits a distinct pharmacological

profile different from rhPTH(1-34) but still provides a clear anabolic effect in the localized jaw region,

the authors have shown that it may possess different potential therapeutic indications from rhPTH(1-

34). This biological difference is thought to be due to dimeric ^{R25C}PTH(1-34) exhibiting a more

preferential binding affinity for the RG versus R0 PTH1R conformation, despite having a diminished

affinity for either conformation. (*Lee and Lee, 2022; Park et al., 2021*) Additionally, the potency of

cAMP production in cells was lower for dimeric R25CPTH compared to monomeric R25CPTH,

consistent with its lower PTH1R-binding affinity. (*Noh et al., 2024*) One of the potential clinical

advantages of dimeric R25CPTH(1-34) is its partial agonistic effect in pharmacodynamics. This

property may allow for a more fine-tuned regulation of bone metabolism, potentially reducing the risk

of adverse effects associated with full agonism, such as hypercalcemia and bone resorption by

osteolcast activity. Moreover, the dimeric form may offer a more sustained anabolic response, which

could be beneficial in the context of long-term treatment strategies. (*Noh et al., 2024*) Also, the effects

of dimer were prominent, as we mentioned better bone formation than the control group.

In addition to the effect of systemic bone mineral gain, the unique anabolic feature of PTH has received clinical attention as an emerging strategy due to the increased risk of ONJ following the use of antiresorptive and the rising need for implantation and bone augmentation in the field of orthopedics and maxillofacial surgery (*Ruggiero et al., 2022*). Moreover, site-specific differential effects of teriparatide have not been clarified, although it represents an issue often associated with selective concentrations of teriparatide that cause anabolic effects on the central skeleton and possible bone mineral decrease on the peripheral skeleton, including the skull (*McClung et al., 2005*). Moreover, it can be inferred that facial and jaw bones, which have the same developmental origin as the skull through membranous ossification, will show the same bone response as the skull (*Setiawati and Rahardjo, 2019*). Previous studies have demonstrated that the central skeleton, including the lumbar and thoracic spine and pelvis regions, showed an increase in areal BMD, while the arms, legs, and skull showed a decrease in bone minerals, suggesting an effect of rhPTH(1-34) on the redistribution of bone minerals from the peripheral to the central skeleton (*Paggiosi et al., 2018*).

However, the results of this study demonstrated that rhPTH(1-34) and dimeric [R25C]PTH(1-34) significantly improved the osseointegration of bone and titanium, as well as jawbone regeneration (Figure 3). The authors have attributed this phenomenon to the unique anatomical characteristics observed in the jawbone. The jawbone in the human body undergoes the most rapid bone remodeling and has excellent blood flow (*Huja et al., 2006*). since is continuously exposed to mechanical stress due to mastication and swallowing, thereby suggesting that the net anabolic effect of rhPTH(1-34) in the jawbone, which is not part of the central skeleton, is achieved through mechanical loading. A recent study by Robinson et al. demonstrated that rhPTH(1-34) and mechanical loading additively stimulate anabolic modeling and synergistically stimulate remodeling in trabecular bone, findings that further support this notion (*Robinson et al., 2021*). However, further investigation is needed to fully understand this relationship.

The anabolic effects of rhPTH(1-34) have been demonstrated through several large randomized controlled trials (*Neer et al., 2001; Tsai et al., 2013*), although the drawbacks arising from the unique

characteristics of rhPTH(1-34) have hindered its broad clinical application. In addition to the

disadvantages of continuous self-injection, high cost, and a two-year treatment limit, the greatest

concern with prolonged use of teriparatide might be accelerated bone remodeling, subsequent

hypercalcemic condition, and increased bone resorption (*Burr et al., 2001; Fox, Miller, Newman, et*

al., 2007; Fox, Miller, Recker, et al., 2007; Neer et al., 2001; Sato et al., 2004; Tsai et al., 2013).

However, recent studies have shown that the frequency and dosing of rhPTH(1-34) administration can

lead to different bone responses. (*Yamamoto et al., 2016; Yamane et al., 2017*) An in vivo study by

Yamamoto et al. reported that lower doses of rhPTH(1-34) with high-frequency administration

resulted in the formation of thin trabeculae, osteoclastogenesis, and accelerated bone remodeling,

while low-frequency rhPTH(1-34) administration showed a phenomenon of modeling-based

formation through thicker trabeculae, mature osteoblasts, and new bone formation (*Yamane et al.,*

2017). This supports the notion that modeling-based bone gain can be an important axis in PTH

anabolism and the primary principle of PTH as bone remodeling-driven bone anabolism.

The use of daily injections in this study was intended to simulate intermittent PTH therapy, a well-

established clinical approach for managing osteoporosis and enhancing bone regeneration.

Intermittent administration of PTH, as opposed to continuous exposure, is critical for maximizing the

anabolic response while minimizing the catabolic effects that are associated with higher frequency or

continuous hormone levels. Our findings support the notion that even with daily administration, both

rhPTH(1-34) and dimeric dimeric ^{R25C}PTH(1-34) promote bone formation and osseointegration,

consistent with the outcomes expected from intermittent therapy. It's important for future research to

consider the dosage and timing of administration to further optimize the therapeutic benefits of PTH

analogs (Dempster et al., 2001; Hodsman et al., 2005).

The limitation of this study is that the therapeutic responses of rhPTH(1-34) and dimeric

^{R25C}PTH(1-34) were focused on local surgical interventions, meaning that we could not investigate the

central skeletal responses, such as in the femur and lumbar. Therefore, further research is needed to

investigate these different responses by administering the PTH analogs at various frequencies and

doses.

Overall, the study demonstrated the therapeutic effects of rhPTH(1-34) and dimeric ^{R25C}PTH(1-34)

on bone regeneration and titanium osseointegration using a beagle model with osteoporosis.

Validation of the anabolic effects of rhPTH(1-34) and dimeric ^{R25C}PTH(1-34) in large animals has

resulted in a broader understanding of their physiological and therapeutic functions and further

expands their potential applications.

Material and Methods
271

Animal preparation
272

This study was conducted in compliance with the ARRIVE guidelines and approved by the Animal

Research Committee of Cronex Co., Ltd., Hwaseong, South Korea. (CRONEX-IACUC 201801002).

All animal experiments, including animal selection, management, preparation, and surgical protocols

were conducted in compliance with the Ewha Womans University rules for animal experiments. The

animals were housed in a standard laboratory environment (21 ± 1 ∘C with 40–70% humidity;

12 hours light/dark cycle) with a standardized food/water supply.

Beagles were chosen for this study because the bone size and dentition could accommodate human

dental implants and the application of mechanical force to implants. To induce osteoporosis, 12

female beagles underwent bilateral ovariectomy (OVX) at 12 weeks of age, followed by osteoporosis

development for 12 weeks before being used for the next experiments at the age of 24 weeks. Health

and oral hygiene were checked and maintained daily (Figure 1A).

Experiment protocol
285

Anesthesia was induced using Zolazepam/Tiletamine (10 mg/kg body weight, Zoletil; Virbac

Laboratories, Carros, France) and xylazine hydrochloride (Rumpun$^{®}$, Bayer, Leverkusen, Germany),

by intramuscular injection. The beagles were anesthetized using inhalation anesthesia for the implant

surgery, and antibiotics (Ceftriaxone, Kyungdong Pharm, Seoul, South Korea) were administered for

three days.

Both mandibular premolars 1–4 were extracted at 12 weeks after OVX. Then, 12 weeks later, three

dental titanium implants (TS III® 3.0 x 10 mm, Osstem, Seoul, Korea) were inserted into the right

lower jaw of each animal via a conventional implant surgical procedure under saline irrigation. Each

implant was placed over at least a 3.0 mm distance. In detail, the first implant was placed over 3.0 mm

behind the canine. The second and third implants were inserted to create a 3 mm circumferential bony

defect using a Ø 6 mm trephine bur. Then, an additional bone graft (Bio-oss® small particle 1.0 g and

Bio-gide®, Geistlich, Switzerland) was placed on the bone defect around the 3rd implant, and the bone

gap remained with nothing at the 2nd implant. On the left lower jaw, two artificial bone defects of 5 x

10 mm were made using trephine bur. The anterior hole remained defective without a graft, while a

bone graft (Bio-oss® small particle 1.0 g and Bio-gide®, Geistlich, Switzerland) was applied to the

posterior hole.

After 2 weeks of healing, twelve female OVX beagles were randomly designated into three groups

as follows: 1) control group with normal saline injection, 2) PTH(1-34) group with daily 40 µg/day

rhPTH(1-34) injection (Forsteo®, Eli Lilly and Company, Indianapolis, IN, USA), and 3) dimeric

^{R25C}PTH(1-34) group with 40 µg/day injection (dimeric ^{R25C}PTH(1-34), chemically synthesized by the

by Anygen, Gwangju, Republic of Korea). Each animal received one injection per day, aimed at

replicating the intermittent rhPTH(1-34) exposure proven beneficial for bone regeneration and overall

skeletal health in clinical settings (Neer et al., 2001; Kendler et al., 2018). This regimen was chosen to

investigate the potential anabolic effects of these specific PTH analogs under conditions closely

resembling therapeutic use. Animals were injected subcutaneously for 10 weeks, after which, they

were euthanized and the bone regeneration and implant osseointegration were evaluated (Figure 1A,

B, and C). To analyze the dynamic bone formation, 10 mg/kg of calcein green (Sigma, St, Louis, USA)

and 30 mg/kg of oxytetracycline yellow (Fluka, Shanghai, China) were intramuscularly administered

at 2 weeks and 10 weeks after operation.

Micro-computed tomographic (µCT) analysis

To examine the microarchitectural effects of PTH(1-34) and dimeric [R25C]PTH(1-34) on bone regeneration and osseointegration, radiographic analysis was performed using µCT on the right mandible (Figure 1D). The specimens were fixed in 4% paraformaldehyde for 48 hours before being assessed by micro-computed tomography (µCT, SkyScan1173 ver. 1.6, Bruker-CT, Kontich, Belgium). The specimens were imaged with a pixel size of 29.83 µm. The voltage and current intensities of the images were 130 kV and 60 µA, respectively. The regions of interest (ROI) were determined as the 10 x 10 mm square area, located 3 mm from the bottom of the implant. Bone mineral density (BMD), bone volume (BV; mm^3), trabecular number (Tb.N; 1/mm), trabecular thickness (Tb.Th; µm), and trabecular separation (Tb.S; µm) were analyzed.

Histological and histomorphometric analysis

Bone histomorphometric parameters were computed and shown in accordance with recommendations by the ASBMR histomorphometric nomenclature committee. (*Dempster et al., 2013*) Goldner's trichrome and Masson trichrome staining were performed on both the right implantation and left bone defect sites, respectively. Specimens were dehydrated in increasing concentrations of ethanol and embedded in a mixture of ethanol and Technovit 7200 resin (Heraeus Kulzer, Wehrheimm, Germany), with an increasing ratio of resin. Following resin infiltration, the specimens were hardened in a UV embedding system (KULZER EXAKT 520, Norderstedt, Germany) for a day. The undecalcified specimens were cut using an EXAKT diamond cutting system (EXAKT 300 CP, Norderstedt, Germany), and the soft tissue and bone were attached to an acryl slide by an adhesive system. The section width of the specimen was adjusted to 40 ± 5µm using a grinding system (EXAKT 400CS, KULZER, Norderstedt, Germany). The specimens on the right implantation site were stained with Goldner's trichrome and photographed by a Panoramic 250 Flash III system (3DHISTECH Ltd., Budapest, Hungary). The bone–implant contact ratio (BIC, %) was assessed as the linear percentage of the interface with direct contact between the bone and implant to the total

interface of the implant using CaseViewer program software (3DHISTECH Ltd.).

Tartrate-resistant acid phosphatase (TRAP) assay was performed on the left bone defect sites. The

bone specimens were fixed in 4% paraformaldehyde overnight and decalcified in 10%

ethylenediaminetetraacetic acid (EDTA) for 7 days. The decalcifying solution was changed every

other day. The specimens were embedded in paraffin and cut into sections. According to the

manufacturer's instructions, the sections were deparaffinized and stained using a TRAP staining kit

(Sigma, St. Louis. MO, USA). The number of TRAP-positive cells in the sections was counted under

a microscope (DM2500, Leica Microsystems, Wetzlar, Germany).

Serum biochemical analysis

Fasting blood samples were drawn in the morning at baseline (T0; 12 weeks), the start of the

injection at 26th weeks(T1), and euthanasia at 36th weeks (T2) for P1NP (procollagen type I N-

terminal propeptide), PTH (parathyroid hormone), CTx (C-terminal telopeptide), and calcium, and

phosphorus (Figure 5A). Calcium and phosphorus were analyzed using the Beckman AU480

Chemistry Analyzer (Beckman Coulter AU480). P1NP (procollagen type I N-terminal propeptide

ELISA kit, Mybiosource), PTH (parathyroid hormone ELISA kit, Aviva Systems Biology), and CTX

(C-terminal telopeptide (CTx-I) ELISA kit, Mybiosource) were analyzed by the ELISA method,

according to the ELISA kit manufacturer's instructions. In all analyses, the measured values were

below the limit of quantification for the standard curve.

Statistical analysis

Data for microarchitectural, histomorphometric, and serum biochemical analyses were expressed

as mean and standard deviation (SD). Non-parametric tests were performed, including the Mann–

Whitney and Kruskal–Wallis tests. Group differences in serum markers over time were compared by

repeatedly measuring the analysis of variance. Statistical analysis was performed using SPSS 26 (IBM

Corp., USA) and Prism 10 (GraphPad, San Diego, CA, USA). P values of < 0.05 were set as

statistically significant.

Acknowledgments and Conflicts of Interest: The authors declare no conflict of interest. This work

was supported by a grant from the Korea Health Technology R&D Project through the Korea Health

Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of

Korea (HI22C1377) and the National Research Foundation of Korea (NRF) grant funded by the

Korean government (MSIT) (No. 2022R1A2C3006002, RS-2024-00339519 and

2018R1D1A1B07041400). This work was supported by the Gachon University Gil Medical Center

(FRD2023-12) and Soonchunhyang University Fund.

Figure legends

Figure 1. Micro-CT analysis (A-D) Experimental design for the controlled delivery of rhPTH(1-34)

and dimeric ^{R25C}PTH(1-34) in ovariectomized beagle model. Representative images for injection and

placement of titanium implant. (E) Micro-CT analysis. bone mineral density (BMD), bone volume

(TV; mm3), trabecular number (Tb.N; 1/mm), trabecular thickness (Tb. Th; um), trabecular separation

(Tb.sp; μm). Error bars indicate standard deviation. Data are shown as mean ± s.d. *p<0.05, **p<0.01,

***p<0.001, n.s., not significant.　P, posterior. R, right

Figure 2. (A-I) Histological analysis of the different groups stained in Goldner's trichrome. The

presence of bone is marked by the green color and soft tissue in red. Red arrows indicate the position

with soft tissues without bone around the implant threads. The area of bone formed was the widest in

the rhPTH(1-34)-treated group. In the dimeric ^{R25C}PTH(1-34)treated group, there is a greater amount

of bone than vehicle-treated group. Green arrows represent the bone formed over the implant. blue

dotted line, margin of bone and soft tissue; Scale bars: 1mm

Figure 3. Histological analysis using Masson trichrome staining results in the rhPTH(1-34) and

dimeric ^{R25C}PTH(1-34)-treated group (A-L) Masson trichrome-stained sections of cancellous bone in

the mandibular bone. The formed bone is marked by the color red. Collagen is stained blue. Black

dotted box magnification region of trabecular bone in the mandible. Scale bars, A-C, G-I: 1mm; D-F,

J-L: 200 μm

Figure 4. Immunohistochemical analysis using TRAP staining for bone remodeling activity (A-L)

TRAP staining is used to evaluate bone remodeling by staining osteoclasts. Osteoclasts is presented

by the purple color. Black dotted box magnification region of trabecular bone in the mandible. (M, N)

The number of TRAP-positive cells in the mandible of the rhPTH(1-34) and dimeric ^{R25C}PTH(1-34)-

treated beagles. Scale bars, A-C, G-I: 1mm; D-F, J-L: 200 μm. Error bars indicate standard deviation.

Data are shown as mean ± s.d. *p<0.05, **p<0.01, n.s., not significant

Figure 5. Measurement of biochemical Marker Dynamics in serum. The serum levels of calcium,

phosphorus, P1NP, and CTX across three time points (T0, T1, T2) following treatment with dimeric

dimeric ^{R25C}PTH(1-34), rhPTH(1-34), or control. (A-B) Calcium and phosphorus levels exhibit an

upward trend in response to both PTH treatments compared to control, suggesting enhanced bone

mineralization. (C) P1NP levels, indicative of bone formation, remain relatively unchanged across

time and treatments. (D) CTX levels, associated with bone resorption, show no significant differences

between groups. Data points for the dimeric ^{R25C}PTH(1-34), rhPTH(1-34), and control are marked by

squares, circles, and triangles, respectively, with error bars representing confidence intervals.

Supplementary Figure. Three-dimensional reconstructed image of the bone surrounding the implants.

Three-dimensional reconstructed images of the peri-implant bone depicting the osseointegration after

different therapeutic interventions. (A) Represents the bone response to recombinant human

parathyroid hormone fragment (rhPTH 1-34) treatment, showing the most robust degree of bone

formation around the implant in the three groups. (B) Shows the bone response to a modified PTH

fragment (dimeric ^{R25C}PTH(1-34)), indicating a similar level of bone growth and integration as seen

with rhPTH(1-34), although to a slightly lesser extent. (C) Serves as the control group, demonstrating

the least amount of bone formation and osseointegration. The upper panel provides a top view of the

bone-implant interface, while the lower panel offers a cross-sectional view highlighting the extent of

bony ingrowth and integration with the implant surface.

Reference

Bae CH, Kang M, Park CY, Park BM, Zhang D, Nam HJ, Yang YM, Shin DM, Choi JY, Lim SK. 2016. A Novel Human PTH Analog [Cys25]hPTH(1-34) Restores Bone Mass in Ovariectomized Mice. *J Clin Endocrinol Metab* **101**(10):3700-3708. DOI:https://doi.org/10.1210/jc.2016-1640.

Bellido T, Ali A, Gubrij I, Plotkin L, Fu Q, O'brien C, Manolagas S, Jilka RJE. 2005. Chronic elevation of parathyroid hormone in mice reduces expression of sclerostin by osteocytes: a novel mechanism for hormonal control of osteoblastogenesis. **146**(11):4577-4583

Burr DB, Hirano T, Turner CH, Hotchkiss C, Brommage R, Hock JMJJoB, Research M. 2001. Intermittently administered human parathyroid hormone (1–34) treatment increases intracortical bone turnover and porosity without reducing bone strength in the humerus of ovariectomized cynomolgus monkeys. **16**(1):157-165.

Cheloha RW, Gellman SH, Vilardaga J-P, Gardella TJ. 2015. PTH receptor-1 signalling—mechanistic insights and therapeutic prospects. *Nature Reviews Endocrinology* **11**(12):712-724. DOI:https://doi.org/10.1038/nrendo.2015.139.

Cortet B. 2011. Bone repair in osteoporotic bone: postmenopausal and cortisone-induced osteoporosis. *Osteoporos Int* **22**(6):2007-2010. DOI:https://doi.org/10.1007/s00198-011-1612-3.

Dempster DW, Compston JE, Drezner MK, Glorieux FH, Kanis JA, Malluche H, Meunier PJ, Ott SM, Recker RR, Parfitt AMJJob, Bone mrtojotASf, Research M. 2013. Standardized nomenclature, symbols, and units for bone histomorphometry: a 2012 update of the report of the ASBMR Histomorphometry Nomenclature Committee. **28**(1):2.

Fox J, Miller M, Newman M, Recker RR, Turner C, Smith SJB. 2007. Effects of daily treatment with parathyroid hormone 1–84 for 16 months on density, architecture and biomechanical properties of cortical bone in adult ovariectomized rhesus monkeys. **41**(3):321-330https://www.sciencedirect.com/science/article/pii/S8756328207003997?via%3Dihub.

Fox J, Miller M, Recker RR, Turner C, Smith SJCTI. 2007. Effects of treatment of ovariectomized adult rhesus monkeys with parathyroid hormone 1-84 for 16 months on trabecular and cortical bone structure and biomechanical properties of the proximal femur. **81**:53-63https://link.springer.com/content/pdf/10.1007/s00223-007-9036-y.pdf.

Garcia P, Histing T, Holstein J, Klein M, Laschke M, Matthys R, Ignatius A, Wildemann B, Lienau J, Peters AJECM. 2013. Rodent animal models of delayed bone healing and non-union formation: a comprehensive review. **26**(1):12.

Gomes-Ferreira P, de Oliveira D, Frigério P, de Souza Batista F, Grandfield K, Okamoto RJOI. 2020. Teriparatide improves microarchitectural characteristics of peri-implant bone in orchiectomized rats. **31**:1807-1815https://link.springer.com/content/pdf/10.1007/s00198-

460   020-05431-y.pdf.

461 Hock J, Gera IJJoB, Research M. 1992. Effects of continuous and intermittent administration and
462   inhibition of resorption on the anabolic response of bone to parathyroid hormone. **7**(1):65-
463   72.

464 Huja SS, Fernandez SA, Hill KJ, Li Y. 2006. Remodeling dynamics in the alveolar process in
465   skeletally mature dogs. *Anat Rec A Discov Mol Cell Evol Biol* **288**(12):1243-1249.
466   DOI:https://doi.org/10.1002/ar.a.20396.

467 Jilka RLJB. 2007. Molecular and cellular mechanisms of the anabolic effect of intermittent PTH.
468   **40**(6):1434-
469   1446https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1995599/pdf/nihms25729.pdf.

470 Jung J, Shim G-J, Kim M, Yoon Y, Kim J-E, Jue S-S, Al-Nawas B, Kwon Y-DJJoC-MS. 2021. Effect
471   and timing of parathyroid hormone analog administration for preventing medication-related
472   osteonecrosis of the jaws in a murine model. **49**(8):719-
473   725https://www.sciencedirect.com/science/article/pii/S1010518221000809?via%3Dihub.

474 Jung J, Yoo HY, Kim GT, Lee JW, Lee YA, Kim DY, Kwon YDJJoB, Research M. 2017. Short-term
475   teriparatide and recombinant human bone morphogenetic protein-2 for regenerative
476   approach to medication-related osteonecrosis of the jaw: a preliminary study. **32**(12):2445-
477   2452.

478 Kakehashi H, Ando T, Minamizato T, Nakatani Y, Kawasaki T, Ikeda H, Kuroshima S, Kawakami A,
479   Asahina IJIjoo, surgery m. 2015. Administration of teriparatide improves the symptoms of
480   advanced bisphosphonate-related osteonecrosis of the jaw: preliminary findings.
481   **44**(12):1558-
482   1564https://www.sciencedirect.com/science/article/pii/S0901502715012011?via%3Dihub.

483 Krishnan V, Bryant HU, Macdougald OA. 2006. Regulation of bone mass by Wnt signaling. *J Clin*
484   *Invest* **116**(5):1202-1209. DOI:https://doi.org/10.1172/jci28551.

485 Lee J-H, Davaatseren M, Lee SJE, Metabolism. 2020. Rare PTH gene mutations causing parathyroid
486   disorders: a review. **35**(1):64-70.

487 Lee J-H, Lee S. 2022. Chapter Four - The parathyroid glands and parathyroid hormone: Insights from
488   PTH gene mutations. In: Litwack G, editor. *Vitamins and Hormones*. Academic Press:79-108.

489 Lee M, Partridge NCJCoin, hypertension. 2009. Parathyroid hormone signaling in bone and kidney.
490   **18**(4):298.

491 Lindsay R, Krege J, Marin F, Jin L, Stepan JJOi. 2016. Teriparatide for osteoporosis: importance of
492   the   full   course.   **27**:2395-
493   2410https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4947115/pdf/198_2016_Article_3534.p
494   df.

495 McClung MR, San Martin J, Miller PD, Civitelli R, Bandeira F, Omizo M, Donley DW, Dalsky GP,

Eriksen EFJAoim. 2005. Opposite bone remodeling effects of teriparatide and alendronate in increasing bone mass. **165**(15):1762-1768.

Mosekilde L, Danielsen CC, Søgaard CH, McOsker JE, Wronski TJ. 1995. The anabolic effects of parathyroid hormone on cortical bone mass, dimensions and strength—assessed in a sexually mature, ovariectomized rat model. *Bone* **16**(2):223-230. DOI:https://doi.org/https://doi.org/10.1016/8756-3282(94)00033-V.

Neer RM, Arnaud CD, Zanchetta JR, Prince R, Gaich GA, Reginster JY, Hodsman AB, Eriksen EF, Ish-Shalom S, Genant HK, Wang O, Mitlak BH. 2001. Effect of parathyroid hormone (1-34) on fractures and bone mineral density in postmenopausal women with osteoporosis. *N Engl J Med* **344**(19):1434-1441. DOI:https://doi.org/10.1056/nejm200105103441904.

Noh M, Che X, Jin X, Lee D-K, Kim H-J, Park DR, Lee SY, Lee H, Gardella T, Choi J-Y, Lee S. 2024. Dimeric [R25C]PTH(1–34) Activates the Parathyroid Hormone-1 Receptor in vitro and Stimulates Bone Formation in Osteoporotic Female Mice. *bioRxiv*:2024.2003.2013.584815. DOI:https://doi.org/10.1101/2024.03.13.584815.

Paggiosi MA, Yang L, Blackwell D, Walsh JS, McCloskey E, Peel N, Eastell RJOI. 2018. Teriparatide treatment exerts differential effects on the central and peripheral skeleton: results from the MOAT study. **29**:1367-1378https://link.springer.com/content/pdf/10.1007/s00198-018-4445-5.pdf.

Park D, Kitaura Y, Dean T, Ohba S, Gardella TJ, Chung U-i, Lee SY, Lee S. 2021. Therapeutic Potentials of Dimeric Cys25PTH(1–34) Peptide for Osteoporosis and Fracture Healing of the Bones- Buy One, Get One Free. *Journal of the Endocrine Society* **5**(Supplement_1):A240-A240. DOI:https://doi.org/10.1210/jendso/bvab048.488.

Robinson ST, Shyu PT, Guo XEJB. 2021. Mechanical loading and parathyroid hormone effects and synergism in bone vary by site and modeling/remodeling regime. **153**:116171https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8499476/pdf/nihms-1739600.pdf.

Ruggiero SL, Dodson TB, Aghaloo T, Carlson ER, Ward BB, Kademani D. 2022. American Association of Oral and Maxillofacial Surgeons' Position Paper on Medication-Related Osteonecrosis of the Jaws—2022 Update. *Journal of Oral and Maxillofacial Surgery* **80**(5):920-943. DOI:https://doi.org/https://doi.org/10.1016/j.joms.2022.02.008.

Sato M, Westmore M, Ma YL, Schmidt A, Zeng QQ, Glass EV, Vahle J, Brommage R, Jerome CP, Turner CHJJoB, Research M. 2004. Teriparatide [PTH (1–34)] strengthens the proximal femur of ovariectomized nonhuman primates despite increasing porosity. **19**(4):623-629https://asbmr.onlinelibrary.wiley.com/doi/pdfdirect/10.1359/JBMR.040112?download=true.

Schipani E, Langman C, Hunzelman J, Le Merrer M, Loke K, Dillon M, Silve C, Juppner HJTJoCE, Metabolism. 1999. A novel parathyroid hormone (PTH)/PTH-related peptide receptor

mutation in Jansen's metaphyseal chondrodysplasia. **84**(9):3052-3057

Setiawati R, Rahardjo P. 2019. Bone development and growth. *Osteogenesis and bone regeneration*

10.

Silva BC, Bilezikian JPJCoip. 2015. Parathyroid hormone: anabolic and catabolic actions on the

skeleton. **22**:41-

50https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5407089/pdf/nihms854430.pdf.

Tsai JN, Uihlein AV, Lee H, Kumbhani R, Siwila-Sackman E, McKay EA, Burnett-Bowie S-AM,

Neer RM, Leder BZJTL. 2013. Teriparatide and denosumab, alone or combined, in women

with postmenopausal osteoporosis: the DATA study randomised trial. **382**(9886):50-

56https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4083737/pdf/nihms-595764.pdf.

Uusi-Rasi K, Semanick LM, Zanchetta JR, Bogado CE, Eriksen EF, Sato M, Beck TJ. 2005. Effects

of teriparatide [rhPTH (1-34)] treatment on structural geometry of the proximal femur in

elderly osteoporotic women. *Bone* **36**(6):948-958.

DOI:https://doi.org/10.1016/j.bone.2005.03.003.

Yamamoto T, Hasegawa T, Sasaki M, Hongo H, Tsuboi K, Shimizu T, Ota M, Haraguchi M, Takahata

M, Oda KJE. 2016. Frequency of teriparatide administration affects the histological pattern of

bone formation in young adult male mice. **157**(7):2604-2620

Yamane H, Takakura A, Shimadzu Y, Kodama T, Lee J-W, Isogai Y, Ishizuya T, Takao-Kawabata R,

Iimura TJPO. 2017. Acute development of cortical porosity and endosteal naïve bone

formation from the daily but not weekly short-term administration of PTH in rabbit.

12(4):e0175329https://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0175329

&type=printable.

Yu W, Su JJOD. 2020. The effects of different doses of teriparatide on bisphosphonate⬚related

osteonecrosis of the jaw in mice. **26**(3):609-

620https://onlinelibrary.wiley.com/doi/pdfdirect/10.1111/odi.13275?download=true.

Zebaze R, Takao-Kawabata R, Peng Y, Zadeh AG, Hirano K, Yamane H, Takakura A, Isogai Y,

Ishizuya T, Seeman EJB. 2017. Increased cortical porosity is associated with daily, not weekly,

administration of equivalent doses of teriparatide. **99**:80-

84https://www.sciencedirect.com/science/article/pii/S8756328217301114?via%3Dihub.

A

ovariectomy | tooth extraction | implant placement | rhPTH or PTH analog injection | Sacrifice

0 wks | 12 wks | 24 wks | 26 wks | 36 wks

B subcutaneous Injection

C

3
2
1

P
R

D μCT

3 2 1

1. Implant
2. Implant+Gap formation
3. Implant+Gap formation +Bone graft

E

| implant | implant + gap | implant + gap + bone graft |

BMD

Bone Volume

Trabecular Number

Trabecular Thickness

Trabecular Separation

vehicle rhPTH ^Cys25^PTH

	vehicle	rhPTH (1-34)	dimeric ^{Cys25}PTH(1–84)
implant	A	B	C
implant+Gap	D	E	F
implant+Gap+bone graft	G	H	I

	vehicle	rhPTH (1-34)	dimeric ^{Cys25}PTH(1–84)
gap + bone graft	A	B	C
	D	E	F
gap	G	H	I
	J	K	L