Title: Whole-Genome Sequencing of Three Extremophile *Bacillus sp.* Strains Isolated from the Atacama Desert

Authors: Nicole T Cavanaugh[1], Girish Kumar[2], Matteo Couto Frignani[1], Nathan Thewedros[1], Marcello Twahirwa[1], Carlos Riquelme[3], André O Hudson[2], Yunrong Chai[1], Veronica Godoy-Carter[1*]

Affiliations:
[1]Northeastern University College of Science, Department of Biology, Boston, MA
[2]Rochester Institute of Technology College of Science, Thomas H. Gosnell School of Life Sciences, Rochester, NY
[3]Universidad de Antofagasta, Departamento de Biotecnología, Antofagasta, Chile
[*]Corresponding Author: v.godoycarter@northeastern.edu

Abstract:

The Atacama Desert is home to bacteria that use biofilms as a means of protecting themselves
against the harsh environment. We cultured and sequenced the genomes of three *Bacillus sp.*
isolates from the soil of the Atacama Desert. This information will inform research in survival
mechanisms of eubacteria in the Atacama Desert.

Announcement:

The Atacama Desert is one of the most ancient, most highly irradiated, and driest deserts in the
world (1). Despite this, it is home to a variety of bacterial life. Bacteria in this region are shown
to have developed adaptations to survive this harsh, irradiated environment such as high levels
of pigment production and biofilm formation (2). In the past, most research addressing survival
mechanisms in microbes from the Atacama have focused on archaea. We are studying how
eubacteria have developed adaptations to survive in such a desert environment.

To study how biofilms impact bacterial survival, we started by culturing and further isolating a
variety of bacterial colonies from the desert soil and screening for biofilm-forming isolates.
Bacterial colonies were purified after plating 100 µL of a soil water slurry on Reasoner's 2A
(R2A) agar medium and incubating plates at 28 ºC for 48-72 hours. To identify the isolates, we
performed 16S rRNA gene sequencing on the V3/4 variable regions using the following primers:
5'-CCTACGGGNGGCWGCAG-3' and 5'-GACTACHVGGGTATCTAATCC-3'. Using National
Society for Biotechnology Information (NCBI) BLASTN version 2.15.0, three *Bacillus sp.* were
identified (3, 4). The *Bacillus* genus contains Gram-positive bacteria that are found throughout
the environment, many of which are known to produce biofilms (5, 6, 7, 8, 9, 10). To test biofilm
formation, strains were grown at 37 ºC to OD600 = 1.0 in Luria-Bertani (LB) broth and 2 µl of
culture was spotted on LB agar plates supplemented with 1% glycerol and 0.1 mM $MnSO_4$
(LBGM, biofilm inducing medium) (11). Colony biofilms were incubated for 48 hours.

28
29
30 **Figure 1: Environmental strains isolated from the Atacama Desert form colony biofilms**.
31 *Colony biofilms of 0102A (A), 0209A (B), 0909A (C), and 3610 (*B. subtilis *lab strain) (D) grown*
32 *on LBGM medium (11) after 48 hours of incubation.*
33
These *Bacillus sp.* were selected for further genomic analysis. Genomic DNA was isolated from
a 2-mL LB culture using the Wizard Genomic DNA Purification Kit (Promega, USA) following kit
instructions for Gram-positive bacteria. Purified DNA was quantified using a NanoDrop
spectrophotometer (Thermo-Fisher), DNA libraries were prepared using the Nextera XT library
preparation kit (Illumina) and sequenced using the Illumina MiSeq System at the Genomics Lab,
Rochester Institute of Technology. Raw reads were trimmed using Trimmomatic Galaxy version
0.38.1 (12). The trimmed reads were assembled de novo using Unicycler Galaxy version
0.5.0+galaxy1 (13). Quality analysis was performed using Quast Galaxy version 5.2.0+galaxy1
(14) and FastQC Galaxy version 0.74+galaxy0 (15). Quality assessments are outlined in Table
1. All programs used were run at default parameters unless otherwise noted. Analysis was
performed using the Prokaryotic Genome Analysis Pipeline (PGAP) version 6.6 to reveal the
total number of genes, protein coding sequences, rRNAs, and tRNAs in each assembly (Table
1) (16, 17, 18). Further analysis of the *eps* operon in each genome using NCBI BLASTN Galaxy
Version 2.14.1+galaxy and taxonomy analysis by the Automated Multi Locus Species Tree
(autoMLST) showed that these three isolates were most closely related to *Bacillus vallismortis*
and Bacillus mojavensis (3-4, 19).
50

Isolate	Most Closely Related Species	Total Bases (Mbp)	Read Length (bp)	Number of Reads (FastQC)	Number of Contigs	N50 (bp)	GC Content	Total Genes	Protein Coding Sequences	rRNA Operons	tRNAs
0102A	*B. vallismortis*	813.2	4,192,438	4,406,089	27	350,970	43.87%	4296	4067	3	76
0209A	*B. vallismortis*	980.4	4,167,904	5,239,849	19	1,048,338	43.98%	4208	3989	3	73
0909A	*B. mojavensis*	857.1	3,899,559	4,586,376	19	930,776	43.66%	4061	3870	3	67

51
Table 1: Summary of quality assessments for each strain. *A summary of the identities and*
quality assessments of each strain 0102A, 0209A, and 0909A. For each strain, the total bases
and number of reads was determined using FastQC Galaxy version 0.74+galaxy0. Read length,
number of contigs, N50, and GC content were assessed using Quast Galaxy version
5.2.0+galaxy1. Numbers of total genes, protein coding sequences, rRNAs, and tRNAs were
determined using the Prokaryotic Genome Analysis Pipeline (PGAP) version 6.6.
58

Data Availability:
The WGS projects for *Bacillus subtilis* strains 0102A, 0209A, and 0909A have been deposited in
GenBank under BioProject number PRJNA1071648. Strain 0102A is deposited under
BioSample number SAMN39706723 and SRA number SRR27839954. PGAP analysis can be
found under accession number JBAIZW000000000. Strain 0209A is deposited under
BioSample number SAMN39706724 and SRA number SRR27839953. PGAP analysis can be
found under accession number JBAIZV000000000. Strain 0909A is deposited under BioSample
number SAMN39706725 and SRA number SRR27839952. PGAP analysis can be found under
accession number JBAIZU000000000.
The NCBI BLASTN webserver was used to identify the 0102A, 0209A, and 0909A 16s rRNA
gene sequence. All programs found on the Galaxy platform can be found at usegalaxy.org.
70
Acknowledgements:
We would like to thank the National Science Foundation (MCB1651732) and the NU Global
Experience Office for their support. Thank you to the RIT Genomics Lab. V.G. was supported by
the HHMI Inclusive Excellence grant from the NU Biology Department. M.C.F, N.T., and M.T.
were supported by NU Undergraduate Research and Fellowships Office PEAK Awards. N.C. is
supported by the NSF Graduate Research Fellowship Program (1938052).
77

References:

[1] Bull, A. T., Andrews, B. A., Dorador, C., & Goodfellow, M. (2018). Introducing the Atacama

Desert. *Antonie van Leeuwenhoek*, *111*(8), 1269–1272. https://doi.org/10.1007/s10482-018-

1100-2

[2] Reverdy, A., Hathaway, D., Jha, J., Michaels, G., Sullivan, J., Mac-Adoo, D. D., Riquelme, C.,

Chai, Y., & Godoy, V. G. (2020). *Insights into the diversity and survival strategies of soil*

bacterial isolates from the Atacama Desert (p. 2020.09.24.312199). bioRxiv.

https://doi.org/10.1101/2020.09.24.312199

[3] Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J.

(1997). Gapped BLAST and PSI-BLAST: A new generation of protein database search

programs. *Nucleic Acids Research*, *25*(17), 3389–3402. https://doi.org/10.1093/nar/25.17.3389

[4] Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L.

(2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1), 421.

https://doi.org/10.1186/1471-2105-10-421

[5] Arnaouteli, S., Bamford, N. C., Stanley-Wall, N. R., & Kovács, Á. T. (2021). Bacillus subtilis biofilm formation and social interactions. *Nature Reviews Microbiology*, *19*(9), Article 9. https://doi.org/10.1038/s41579-021-00540-9

[6] Beauregard, P. B., Chai, Y., Vlamakis, H., Losick, R., & Kolter, R. (2013). Bacillus subtilis biofilm induction by plant polysaccharides. *Proceedings of the National Academy of Sciences*, *110*(17), E1621–E1630. https://doi.org/10.1073/pnas.1218984110

[7] Branda, S. S., González-Pastor, J. E., Ben-Yehuda, S., Losick, R., & Kolter, R. (2001). Fruiting body formation by Bacillus subtilis. *Proceedings of the National Academy of Sciences*, *98*(20), 11621–11626. https://doi.org/10.1073/pnas.191384198

[8] Diomande, S., NGUYEN-THE, C., Guinebretière, M.-H., Broussolle, V., & Brillard, J. (2015). Role of fatty acids in Bacillus environmental adaptation. *Frontiers in Microbiology*, *6*. https://www.frontiersin.org/articles/10.3389/fmicb.2015.00813

[9] Fritze, D. (2004). Taxonomy of the Genus Bacillus and Related Genera: The Aerobic Endospore-Forming Bacteria. *Phytopathology®*, *94*(11), 1245–1248. https://doi.org/10.1094/PHYTO.2004.94.11.1245

[10] Vlamakis, H., Chai, Y., Beauregard, P., Losick, R., & Kolter, R. (2013). Sticking together: Building a biofilm the Bacillus subtilis way. *Nature Reviews Microbiology*, *11*(3), Article 3. https://doi.org/10.1038/nrmicro2960

[11] Shemesh, M., & Chai, Y. (2013). A Combination of Glycerol and Manganese Promotes Biofilm Formation in Bacillus subtilis via Histidine Kinase KinD Signaling. *Journal of Bacteriology*, *195*(12), 2747–2754. https://doi.org/10.1128/JB.00028-13

[12] Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

[13] Wick, R. R., Judd, L. M., Gorrie, C. L., & Holt, K. E. (2017). Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLOS Computational Biology*, *13*(6), e1005595. https://doi.org/10.1371/journal.pcbi.1005595

[14] Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D., & Gurevich, A. (2018). Versatile genome assembly evaluation with QUAST-LG. *Bioinformatics*, *34*(13), i142–i150. https://doi.org/10.1093/bioinformatics/bty266

[15] *Babraham Bioinformatics—FastQC A Quality Control tool for High Throughput Sequence Data*. (n.d.). Retrieved February 2, 2024, from https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

[16] Haft, D. H., DiCuccio, M., Badretdin, A., Brover, V., Chetvernin, V., O'Neill, K., Li, W., Chitsaz, F., Derbyshire, M. K., Gonzales, N. R., Gwadz, M., Lu, F., Marchler, G. H., Song, J. S., Thanki, N., Yamashita, R. A., Zheng, C., Thibaud-Nissen, F., Geer, L. Y., … Pruitt, K. D. (2018). RefSeq: An update on prokaryotic genome annotation and curation. *Nucleic Acids Research*, *46*(D1), D851–D860. https://doi.org/10.1093/nar/gkx1068

[17] Li, W., O'Neill, K. R., Haft, D. H., DiCuccio, M., Chetvernin, V., Badretdin, A., Coulouris, G., Chitsaz, F., Derbyshire, M. K., Durkin, A. S., Gonzales, N. R., Gwadz, M., Lanczycki, C. J., Song, J. S., Thanki, N., Wang, J., Yamashita, R. A., Yang, M., Zheng, C., … Thibaud-Nissen, F. (2021). RefSeq: Expanding the Prokaryotic Genome Annotation Pipeline reach with protein family model curation. *Nucleic Acids Research*, *49*(D1), D1020–D1028. https://doi.org/10.1093/nar/gkaa1105

[18] Tatusova, T., DiCuccio, M., Badretdin, A., Chetvernin, V., Nawrocki, E. P., Zaslavsky, L., Lomsadze, A., Pruitt, K. D., Borodovsky, M., & Ostell, J. (2016). NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Research*, *44*(14), 6614–6624. https://doi.org/10.1093/nar/gkw569

[19] Alanjary, M., Steinke, K., & Ziemert, N. (2019). AutoMLST: An automated web server for generating multi-locus species trees highlighting natural product potential. *Nucleic Acids Research*, *47*(W1), W276–W282. https://doi.org/10.1093/nar/gkz282