<u>Title:</u>

Targeted Viromes and Total Metagenomes Capture Distinct Components of Bee Gut Phage

Communities

<u>Authors:</u>

Dino Lorenzo Sbardellati[1], Rachel Lee Vannette[2]

<u>Author information:</u>

[1]Microbiology Graduate Group, University of California Davis, Davis, California, USA.

[2]Department of Entomology and Nematology, University of California Davis, Davis, California,

USA

<u>Corresponding author</u>

Dino L. Sbardellati

Phone: 530-752-3379

Email: <u>dlsbardellati@ucdavis.edu</u>

<u>Abstract:</u>

Despite being among the most abundant biological entities on earth, bacteriophage

(phage) remain an understudied component of host-associated systems. One limitation to

studying host-associated phage is the lack of consensus on methods for sampling phage

communities. Here, we compare paired total metagenomes and viral size fraction metagenomes

(viromes) as methods for investigating the dsDNA viral communities associated with the GI tract

of two bee species: the European honey bee *Apis mellifera* and the eastern bumble bee *Bombus*

impatiens. We find that viromes successfully enriched for phage, thereby increasing phage

recovery, but only in honey bees. In contrast, for bumble bees, total metagenomes recovered

greater phage diversity. Across both bee species, viromes better sampled low abundance and low

occupancy phage, while total metagenomes were biased towards sampling temperate phage and

the most prominent phage. Additionally, many of the phage captured by total metagenomes were

absent altogether from viromes. Comparing between bees, we show that phage communities in

commercially reared bumble bees are significantly reduced in diversity compared to honey bees,

likely reflecting differences in bacterial titer and diversity. In a broader context, these results

highlight the complementary nature of total metagenomes and targeted viromes, especially when

applied to host-associated environments. Overall, we suggest that studies interested in assessing

total communities of host-associated phage should consider using both approaches. However,

given the constraints of virome sampling, total metagenomes may serve to sample phage

communities with the understanding that they will preferentially sample dominant and temperate

phage.

<u>Keywords:</u>

bacteriophage, bee microbiome, host-associated microbiome, microbial ecology, viral ecology

<center>Introduction</center>

Bacteriophage (phage) are the viruses which target bacteria. Phage are hypothesized to be among the most abundant biological entities on earth[1–3] and to be important modulators of microbial communities [2,4]. In marine and soil environments, where these prokaryotic viruses are relatively well studied, phage shape the microbiomes they infect by lysing dominant bacteria [5–7], influencing nutrient turnover and cycling[8,9], and vectoring functional genes between bacterial hosts[10–13]. While the basis for much of our understanding of phage ecology stems from the study of marine and soil environments, there is a growing interest in understanding the role of phage in host-associated systems[14,15]. However, host-associated environments and free-living soil and marine environments represent different microbial ecosystems with different selective pressures. As such, insights gained from studying free-living phage communities may not always apply to communities of host-associated phage. Specifically, while the best methods for sampling free-living phage communities have been empirically compared[16,17], little work has evaluated the best methods for sampling host-associated phage[18]. Moreover, given the importance of microbiomes to host health, understanding phage community dynamics and how phage differ between related animal hosts is a key priority.

Like their environmental counterparts, phage infecting host-associated bacteria (hereafter 'host-associated' phage) can affect the abundance, composition, and function of host-derived microbial communities. However, unlike free-living phage, host-associated phage must also adapt to their close association with a host and its microbiome. For example, the comparatively high-density bacterial communities associated with the human gut[19,20], mucosal layer of corals[21], and other host environments is suggested to favor integrase encoding temperate phage, as opposed to obligately lytic phage[22–24]. Phage targeting bacteria adhered to gut mucosa have also

been shown to evolve the ability to adhere to and persist in the animal mucosa where their target

bacteria reside[25]. Given the fundamental differences in the ecology of free-living and host-

associated phage communities[22], it remains unclear whether sampling methods developed for

free-living communities are also appropriate for other habitat types.

Two methods are frequently used for describing dsDNA viral communities: bioinformatic

mining of total metagenomes and targeted viral enrichments (viromes). In bioinformatic mining

of total metagenomes, total genomic DNA is extracted from a sample, amplified, sequenced

using 'shotgun' or untargeted methods, assembled into metagenomes, and phage computationally

mined[26,27]. This approach is cheap, easy to perform, and offers simultaneous characterization of

all DNA in a sample (including bacteria). However, only a small minority of the total

metagenomic data generated will originate from phage, resulting in a relatively shallow sampling

of phage communities. An alternative method, targeted virome sequencing, leverages the

physical characteristics of viruses to select for the phage and virus-like particle fraction of a

sample prior to nucleic acid extraction and sequencing[28,29]. With more sequencing devoted to

viruses, targeted viromes can recover a greater diversity of phage relative to total metagenomes.

This is illustrated by previous comparisons of viromes and total metagenomes in soil and

freshwater samples[16,17]. However, whether viromes always outperform total metagenomes is not

clear. Work applying these two methods to host-associated (human gut) and low-biomass (deep-

sea marine) samples has produced conflicting results[17,18]. One explanation for these

discrepancies is the biases associated with total metagenomes and targeted viromes[30,31]. For

example, viromes remove bacterial cells prior to sequencing and, as a result, miss integrated

temperate phage[32,33]. This is especially relevant in host-associated systems where temperate

phage are abundant[32]. Total metagenomes may also be preferable to viromes logistically, given

that viromes demand a relatively large biomass and are labor-intensive to generate, which can

limit sample sizes in low-biomass systems.

Insects have become valuable models for exploring host-microbe interactions[34–36]. Social

bees in particular house a simple (5 - 9 taxa), highly conserved, socially transmitted, gut bacterial

community[37,38] which contributes to host nutrient acquisition and pathogen defense[39–41]. These

features position social bees as an excellent model for studying not only host-microbe

interactions[42,43], but tripartite host-microbe-phage interactions as well[44]. A series of recent studies

using targeted virome approaches have shown that honey bees host a diversity of novel phage

which target core bee gut bacteria[45–47]. While this work has advanced our understanding of bee

phage, no work has evaluated how sampling method influences recovered bee phage

communities. Given the large biomass required for viromes, total metagenomes may be

advantageous for surveying the phage communities associated with bees and other biomass

limited systems. Additionally, describing the phage associated with other social bees will develop

our understanding of how phage communities differ between related hosts. Lastly, broader work

comparing the role and diversity of phage among small invertebrate hosts will hinge on

appropriate sampling methods, so comparing the performance of viromes and total

metagenomics is particularly important for enabling future comparative study.

Here, we evaluate the phage communities inferred by applying total metagenomes and

targeted viromes to managed *Apis mellifera* and commercially raised *Bombus impatiens* gut

material. We hypothesize that targeted viromes will enrich for phage sequences and capture a

greater diversity of phage, relative to bioinformatic mining of total metagenomes, and that phage

will differ between host bee species. To interrogate these hypotheses, we first examine how host

bee species and sampling method impact sequencing read data and viral enrichment. We then

validate our sampling and computational methodology through rarefaction and by comparing our

phage sequences to those previously described in honey bees. Next, to test how phage

community differs across host bee and sampling methods, we compare phage community

diversity, structure, and composition. We then use gene content, occupancy-abundance plots,

qPCR, and bacterial community profiling to delve into what contributes to the apparent

differences in phage communities. Overall, we find that honey bees host more phage than do

bumble bees, but that the method which captured the most phage differed by bee species. This

suggests that viromes and total metagenomes are each biased in their own way and appear to

sample different populations of phage. As a result, we propose these techniques are

complementary in describing the full diversity of host-associated phage.

<center>Methods</center>

<u>Bee rearing, sampling, and experimental design</u>

A detailed description of our methods (including sample collection, and bench and

computational work) is provided in the supplemental materials. Briefly, bees were harvested

from three colonies of managed honey bees (*Apis mellifera*) and three colonies of commercially

reared bumble bees (*Bombus impatiens*, Koppert Biological Systems; Howell, MI, USA).

Bumble bee colonies were maintained in the laboratory and provided with honey bee collected

pollen and artificial nectar (Koppert Biological Systems) *ad libitum*, as per manufacturer's

recommendation. Honey bees were sampled from colonies maintained at the University of

California, Davis apiary and were collected from inside the colony using a handheld bee vacuum

between January 30th – February 13th 2023, 8:00am – 10:00am.

Bees were anesthetized via a 60 sec CO_2 exposure and then euthanized via decapitation.

Immediately following sacrifice, the mid-hindgut section was dissected from 100 bees (all

collected from the same colony) and pooled for same day virome extraction. Bees destined for

total metagenomic DNA extraction were collected the same day as virome samples and were

stored at -20°C pending mid-hindgut dissection on the following day. All dissections took place

in sterile PBS using ethanol and flame sterilized forceps. Each honey and bumble bee colony

generated one targeted virome and three total metagenomes (**Fig. 1**).

Sample preparation and DNA extraction

Phage enrichments for targeted virome sequencing were carried out following a protocol

adapted from previous publications[28,29,48] and is described fully in the supplement methods.

Briefly, bee gut pools were combined with 10 ml of Protein-supplemented phosphate-buffered

saline (PPBS: 2% BSA, 10% PBS, 1% K-citrate, 150 mM MgSO4), homogenized, and submitted

to a series of washes aimed at eluting phage particles. Phage were enriched via centrifugation,

0.22 um filtration, and ultracentrifugation. Lastly, phage pellets were resuspended in 200 uL

water and DNase treated.

For total metagenomes, mid-hind guts of individual bees were dissected and

homogenized and used for extraction. DNA was extracted from both virome and total

metagenome samples using a DNeasy PowerSoil Pro kit (Qiagen, Hilden, Germany) following

the manufacturer's instructions with the optional Qiagen Vortex Adapter 15 min bead beating

step.

Library construction and DNA sequencing

Extracted DNA from samples and kit blank negative controls was submitted to the

University of California Davis Genome Center for library prep and paired end 150 bp shotgun

metagenomic sequencing. Libraries were prepared using a Kapa Hyper prep kit (Kapa

Biosystems-Roche, Basel, Switzerland) with Illumina TruSeq adapters (Illumina, San Diego,

CA) and a target insert size of 350 bp. Sequencing took place on an Illumina NovaSeq. All

samples were sequenced on the same run.

<u>Read processing, viral prediction, and vOTU table construction</u>

Briefly, all demultiplexed *.fastq.gz* files from the Davis Genome Center were trimmed

and quality filtered using Trim-Galore[49] and Trimmomatic[50]. Reads aligning to negative samples

or the genomes of *A. mellifera* and *B. impatiens* were removed using Bowtie2[51]. Cleaned reads

were k-merized for complexity comparisons using Sourmash[52]. The program metaSPADES was

used to construct assemblies. Assemblies ≥5 kb were passed to Virsorter2[53] for phage prediction.

Phage with ≥95% nucleotide identity were collapsed into viral operational taxonomic units

(vOTUs) using CD-HIT[54] and were annotated with Prokka[55] using the PHROGs database[56].

vOTUs were classified as temperate when they were predicted to encode an integrase gene.

Finally, CoverM[57] was used to map cleaned reads against vOTUs. The resulting coverage table

was normalized to coverage per million reads.

<u>Bacterial community and density measurements</u>

Bacterial communities were predicted directly from cleaned total metagenomic reads

with Kraken2[58] and Bracken[59] using default parameters. Bacterial copy number was quantified

from the DNA extracts used for metagenome sequencing via a qPCR protocol adapted from

Christensen et al[60]. Each master mix solution contained: 5ul SSO Advanced Universal SYBR

Supermix 41 (Biorad, Hercules, CA, USA), 0.3ul of primers (10uM) targeting the 16S region of

the rRNA gene (799F= 5'-AACMGGATTAGATACCCKG-3';1115R= 37 5'-

AGGGTTGCGCTCGTTG-3'), 3.4ul Molecular grade water, and 1ul of template DNA (diluted

1:1000 in molecular grade water). Reactions were performed in triplicate for each sample.

<u>Statistical and ecological analyses</u>

Statistical analyses were conducted in R[61] v4.2.3. Briefly, alpha diversity, beta diversity,

and PERMANOVAs were compared between host and sampling method using vegan[62] and

phyloseq[63]. Linear mixed effect models were built using lme4[64], post-hoc tests were run using

emmeans[65], and T-tests were performed using base R. Gene-sharing networks were generated

outside of R using the program vConTACT2[66,67]. Genome alignment plots were constructed

using Clinker[68].

Phage-host predictions were performed using a custom CRISPR-spacer analysis. In short,

we built bacterial metagenome assembled genomes (MAGs) from our total metagenomic dataset

using metaSPADES and MetaBat2[69], combined these MAGs with NCBI assemblies of common

bee gut bacteria, mined all these sequences for CRISPRs using MINCED[70], and then aligned

those CRISPRs to our vOTUs using Blastn[71]. Lastly, to improve the number of vOTUs with host

assignments, we combined phage-host assignments generated by our CRISPR analysis with the

clusters predicted by our vConTACT2 network to infer the hosts of whole phage clusters,

following Rosso et al[45]. Bacterial copy number was compared between species using linear

models.

<u>Results</u>

<u>Bumble bee viromes produce fewer high-quality reads than honey bee viromes</u>

Our sequencing effort generated a total of 419,884,845 raw read pairs, 2.47%

(10,385,967) of which came from extraction negatives. While the number of raw read pairs did

not differ significantly between samples (**Fig. S1;** Bee $t_{5.40}$=0.54 p=0.61, Type $t_{8.47e23}$=1.58

p=0.11, Bee:Type $t_{1.07e23}$=-1.38 p>0.17), there were differences in the number of high-quality

read pairs remaining after quality control (Bee $t_{5.81}$=2.87 p<0.05, Type $t_{2.75e25}$=-0.33 p=0.74,

Bee:Type $t_{2.75e25}$=4.69 p<0.001) (**Fig. 2A**). After filtering, honey bee viromes averaged 15.56

million high-quality read pairs per sample, while bumble bee viromes averaged only 5.49. Total

metagenome samples generated more similar numbers of high-quality reads per sample, with an

average of 8.59 and 5.85 million read pairs generated from honey and bumble bee samples,

respectively.

<u>Bumble bee virome reads are less complex than other samples</u>

Next, we measured the complexity of read libraries to further investigate how bee species

and sampling method influenced the data we generated. This was done by comparing the number

and occurrence of 31bp sized k-mers amongst each set of high-quality read libraries (**Fig. 2B**).

The results of this analysis suggest that, across the range of k-mer occurrences, bumble bee

viromes consistently contain fewer k-mers of a given occurrence than did other sample types.

Moreover, viewing only the y-intercept of this plot (**Fig. 2C**) shows that bumble bee viromes

contain a lower number of singleton k-mers, relative to all other sample types (Bee $t_{5.34}$=2.49

p=0.052, Type t_{16}=-6.47 p<0.001, t_{16}=5.14 p<0.001,). This decreased number of singleton k-

mers, and the tendency towards lower k-mer occurrences suggest that bumble bee viromes are

less complex compared to the other sample types in this dataset.

<u>Targeted virome assemblies are enriched in phage</u>

To evaluate the size and number of assemblies produced from each read library, and to

quantify the extent to which viromes enriched for phage sequences, we visualized the

distribution of assembly sizes in each sample, as well as the percentage of assemblies in a sample

predicted to encode phage sequences (**Fig. 2D**). A total of 4,278 and 37 contigs ≥5kb were

assembled from honey and bumble bee viromes, respectively. After passing to VirSorter2, these

contigs yielded 2,382 honey bee and 24 bumble bee putative phage sequences. Meanwhile,

honey and bumble bee total metagenomes produced 16,615 and 14,335 contigs ≥5kb,

respectively, which yielded 404 and 140 putative phage sequences. This means that 53.94% of

honey bee and 60.57% of bumble bee virome contigs ≥5kb were predicted to be viral, while only

2.42% and 2.13% of honey and bumble bee total metagenome contigs ≥5kb were annotated as

viral. The enrichment of phage sequences in viromes, relative to total metagenomes, provide

validation that viromes successfully enrich for phage when applied to bee guts.

Next, we used rarefaction to assess if the different number of putative phage sequences

recovered by honey and bumble bee viromes (**Fig. 2D**) was a result of differences in sampling

depth (**Fig. 2A**). This analysis shows that even at equal sampling depths, honey bee viromes

consistently recover >40x the number of phage recovered from bumble bee viromes (**Fig. S2A**

and S2B).

<u>Phage target core bee bacteria and resemble previously described honey bee phage</u>

After measuring the ability of viromes to enrich for phage sequences, we sought to

characterize the phage communities we detected in terms of similarity to previously described

bee phage, putative bacterial hosts, and predicted taxonomy. To do this, we first constructed

vOTUs by collapsing putative phage sequences at a 95% average nucleotide identity. This

reduced the number of putative phage from 2,950 to 2,560. We then built gene-sharing networks

comparing our vOTUs to those previously described by other bee phage studies (**Fig. 3A**). In this

network, many of the vOTUs from the current study clustered with those described by other bee

phage studies, suggesting they are related at roughly the genus level[66,67]. For example, viral

cluster 29 contains sequences recovered by the current study, Rosso et al[45], and Deboutte et al[46]

(**Fig. S3**). We interpret these results as further validating our computational methodologies and

further supporting the result that bumble bees host reduced diversity phage communities.

Next, we assigned bacterial hosts to our vOTUs. To focus this and all downstream

analyses on only the most abundant phage, we filtered vOTUs to retain only those with a

coverage ≥5, resulting in 2,481 vOTUs across 24 samples. Overall, we assigned hosts to 29.91%

(742) of the 2,481 vOTUs appearing in our dataset (**Fig. 3B**). In terms of relative abundance,

phage with assigned hosts made up an average of 89.34% and 44.75% of bumble bee viromes

and total metagenomes and 50.54% and 43.89% of honey bee viromes and total metagenomes,

respectively. Notably, the high proportion of phage with assigned hosts in bumble bee viromes is

likely confounded by the fact that these samples had an extraordinarily diminished phage

diversity.

Phage predicted to target the core social bee bacterial genera *Lactobacillus*,

Bifidobacterium, and *Gilliamella* were the most abundant of those with identified hosts in our

dataset (**Fig. 3B**). However, the relative abundance of phage targeting these genera varied by bee

species and sampling method (**Table S1**). Phage targeting *Lactobacillus* were most abundant in bumble bees, especially bumble bee viromes, while *Bifidobacterium* targeting phage were most abundant in honey bee samples, particularly viromes. *Gilliamella* phage were not recovered in bumble bee viromes, but were found in most bumble bee total metagenomes, and were approximately 2x more abundant in honey bee total metagenomes than they were in honey bee viromes. Phage targeting the core bumble bee symbiont *Schmidhempelia* were exclusively found in bumble bee total metagenomes. We also found signatures of colony-level variation, for example phage targeting *Bombilactobacillus* were detected in metagenomes from individuals sampled from colony 1, but no individuals from colonies 2 or 3.

We assigned family level taxonomy to our phage if they clustered with reference phage in gene-sharing networks. In total, we only assigned taxonomy to 9.83% (244) of 2,481 vOTUs (**Fig. S4**). The large majority of classified vOTUs belonged to Siphoviridae (104), Myoviridae (94), and Podoviridae (40), with a smaller number being classified as Autographiviridae (3) and Zobellviridae (1) and multiple assignment (2).

Bee species differ in phage diversity and composition

Next we tested if vOTU diversity and community composition differed between bee species and sampling method. While the diversity and richness of communities followed similar trends (**Fig. 4A and 4B**), there were substantial quantitative differences between these two metrics. When we compared Shannon's diversity, bee species, sampling method, and their interaction all significantly affected vOTU diversity (**Fig. 4A**; Bee t_{20}=2.315, p=0.0313; Type t_{20}=-5.992, p<0.001; Bee:Type t_{20}=8.004, p<0.001). Meanwhile, only the interaction between bee species and sampling method significantly impacted community richness (**Fig. 4B**; Bee

t_{20}=0.690, p=0.498; Type t_{20}=-0.160, p=0.874; Bee:Type t_{20}=10.227, p<0.001). Overall, these

results suggest that sampling method can significantly influence the alpha diversity of inferred

phage communities, but that this can differ in magnitude and direction according to bee species.

In honey bees, 12.1% (290/2,392) of all vOTUs were shared between virome and total

metagenome samples. Similarly, in bumble bees, 12.9% (12/93) of vOTUs were detected by both

sampling methods. Meanwhile, only a few vOTUs were detected in both bee species (0.121%;

Fig. 4C). Additionally, vOTU community composition was predicted primarily by host bee

species, and secondarily based on sample method (**Figure 4D**; Bray-Curtis PERMANOVA Bee

R^2=0.143 $F_{1,20}$=4.17 p<0.001; Method R^2=0.0859 $F_{1,20}$=2.506 p<0.001; Bee:Method R^2=0.0859

$F_{1,20}$=2.505 p<0.001).

<u>Bacterial community variation predicts phage community variation</u>

We hypothesized that differences in phage richness and community composition between

bee species may be driven by differences in bacterial abundance and composition. First, we used

qPCR to compare the number of bacteria found in the guts of honey and bumble bees (**Fig. 5A**)

and found that the average mid-hindgut 16S gene copy number was significantly lower in

bumble bees than it was in honey bees ($t_{9.13}$=-3.54 p<0.01).

We then used Kraken2 and Bracken to estimate the diversity and taxonomic profile of

bacterial communities (**Fig. 5B, 5C, 5D**). At the species level, bumble bee bacterial communities

hosted lower bacterial richness and greater Shannon's diversity than honey bee bacterial

communities (Richness $t_{13.78}$=-6.27 p<0.001, Shannon $t_{11.70}$=-6.27 p<0.001), suggesting fewer

bacterial taxa were present in the bumble bees we sampled, but that bumble bee bacterial

communities were more evenly divided amongst their constituent bacteria.

Lastly, we used a Mantel test to compare the relationship between bacterial and phage

community structure (**Fig. S5**). We found a high degree of similarity between the bacterial and

phage communities (R=0.74 p<0.001), meaning that individual bees with similar bacterial

communities also have similar phage communities.

<u>Total metagenomes capture prophage but miss rare phage</u>

After identifying factors which explain some of the host-specific differences in phage

communities, we sought to evaluate why sampling method (virome vs total metagenome) also

led to significant differences in inferred phage communities. For this, we first compared the gene

content of vOTUs recovered from viromes and total metagenomes. Because total metagenomes

are primarily comprised of bacterial DNA, we hypothesized that this sampling method would

enrich for temperate phage. Supporting this hypothesis, we found that 189 of the 2075 (9.11%)

vOTUs identified in viromes were predicted to encode an integrase gene, while 89 of the 406

(21.92%) vOTUs recovered from total metagenomes encoded an integrase (**Fig. S6**). Moreover,

integrase encoding phage had a higher relative abundance in total metagenomes than in viromes

(**Fig. 6A**; Bee $t_{14.813}$=-0.625 p=0.541; Type t_{16}=2.845 p<0.05; Bee:Type Type t_{16}=0.463 p=0.650).

These results suggest that, relative to viromes, total metagenomes are biased towards sampling

prophage.

Lastly, to examine how sampling methods differ in recovery of rare vs abundant phage,

we visualized the occupancy-abundance relationships of virome-specific vOTUs vs vOTUs

recovered by both methods (**Fig. 6B and 6C**). This analysis revealed that vOTUs identified in

total metagenomes were among the most abundant and prevalent vOTUs found in viromes,

suggesting that total metagenomes recover more abundant and prevalent phage, while potentially

missing rarer and lower occupancy phage.

339   <u>Discussion</u>

340   Although phage are integral members of complex microbiomes, methods for describing

phage communities are still being developed. Here we compare how two common forms of

phage community sampling, bioinformatic mining of total metagenomes and targeted sequencing

of viromes, influence the phage communities recovered from the guts of two commercially

important bee species: *Apis mellifera* and *Bombus impatiens*. Similar to the results of previous

studies[16–18], we show that sampling method significantly affected alpha diversity, beta diversity,

composition, and structure of sampled phage communities. In particular, we show that viromes

outperform total metagenomes in terms of the number of phage recovered, but that this can

depend on the specific environment being sampled. Overall, viromes were better at sampling rare

and less prevalent phage when bacterial and phage diversity was high, while total metagenomes

captured a greater diversity of phage in low biomass samples. We also show that, regardless of

sample biomass, total metagenomes were enriched for temperate phage, compared to viromes,

and were consistently able to recover phage not found in viromes. Given these results, we

suggest that viromes and total metagenomes each have limitations, are complementary, and that

choice of one method over the other likely depends on the environment being sampled.

<u>Honey bees host a core phage community</u>

357   Viewing our results within the broader context of bee phage, our analyses show that

many features of the phage communities identified here resemble those previously described in

honey bees sampled from North America and Europe[45–47]. For example, *Siphoviridae*,

Myoviridae and *Podoviridae* were among the most common viral families that we and these

previous studies identified. Similarly, *Lactobacillus*, *Gilliamella*, *Bifidobacterium*, and other core

bee gut bacteria were among the most predicted phage hosts in our and previous studies.

Notably, while previous bee phage studies found a number of phage predicted to target

Bartonella, none of our phage were predicted to target this bacterial genus. This is likely because

no available *Bartonella apis* assemblies met the strict criteria for inclusion in our CRISPR spacer

analysis.

Delving deeper, our gene-sharing network and clinker analyses show that many of our

phage share not only large genomic regions with previously described honey bee phage, but that

gene synteny in these geographically disparate viral genomes is also conserved. This

conservation between honey bee phage communities is similar to what was observed by Busby et

al[47]. Likewise, Rosso et al[45] showed that the genomes of honey bee-associated phage sampled

from Europe are able to recruit reads from Japanese honey bee total metagenomic datasets.

Altogether, these findings support the idea that, similar to the highly conserved nature of

some bee gut bacterial communities[37,38,43,72], some features of bee gut phage communities are

also conserved between bees sampled across geographically disparate populations. Future work

should focus on identifying what features of phage communities, whether individual genes,

clusters of genes, or perhaps whole genomes, are conserved between different populations of

bees.

Host bacterial communities predict phage communities

When compared to free-foraging managed honey bees, the commercially produced bumble

bees we sampled hosted significantly lower diversity phage communities. Although differences in

species biology and colony size may play a role, we attribute this low phage diversity primarily to

the low density and diversity bacterial communities these bumble bees hosted, suggesting that

phage diversity and abundance may track bacterial diversity and abundance across hosts. However,

whether such results are generalizable to wild bumble bees is unclear. In our study, bacterial 16S

rRNA copy number in bumble bee guts ranged from 10^6 to 10^7, whereas previous work in the same

species reported 10^8 - 10^9 copies of the 16S rRNA gene[37,73–75]. Given we used standard rearing

methods and diet, we hypothesize that lower bacterial titer may be due to the age of bees we

sampled. Previous work in bumble bees shows that bacterial density increases with worker age,

saturating approximately four days after worker emergence[73]. Future work could examine how

phage abundance and composition changes through worker development and if this reflects

previous observations of bacterial succession with worker age[17]. More generally, these findings

suggest that features of host bacterial communities (i.e. density, diversity, and structure) may be

used to predict phage community features. This is further supported by our Mantel analysis

showing a correlation between bacterial and phage community dissimilarity.

<u>Viromes and total metagenomes reveal complementary phage communities.</u>

There are several possible reasons as to why the phage communities inferred by viromes

and total metagenomes differ. One explanation is the way we generated viromes and how this

influenced the specific population of phage sampled. Research by Hoyles et al.[76] has shown that

passing human fecal samples through a 0.22 um filter reduces the number of phage recovered by

nearly half. As such, size filtration may have led some phage which are present in our bee guts to

have been excluded from viromes. This size exclusion may explain why we, and previous virome

vs total metagenome studies[16–18], consistently show that some phage are only found in total

metagenomes. The particular types of phage which are being excluded likely further explain the

total metagenome and virome differences we observed. We found that putative temperate phage

were more abundant and prevalent in total metagenomes than they were in viromes. This agrees

with existing literature which suggests that size filtration can specifically remove integrated

lysogenic phage, large jumbo phage, and phage adhered to bacterial cells and particles[31,33,77,78].

While it is tempting to expand these results to different environments by stating total

metagenomes always sample temperate phage missed by viromes, work in other systems implies

a more nuanced reality. In agricultural soils, Medellin et al.[16] identified only three phages that

were present in total metagenomes and absent from viromes. This suggests that the degree to

which sample processing skews viromes differs by environment. As a result, we suggest that the

relative utility of viromes and total metagenomes likely depends on the environment being

sampled. Samples with a high abundance of temperate phage, such as host-associated systems

and low-biomass environments, may benefit from total metagenomes. This is similar to the

conclusions made by Kosmopoulos et al.[17] which suggest that the choice of viromes vs total

metagenomes should be environment specific.

While our results highlight the ability of total metagenomes to recover phage missed by

viromes, they also showcase the capacity of viromes to sample a greater diversity of phage.

Similar to previous comparisons of viromes and total metagenomes in human gut, soil, and

freshwater environments[17,18,29], our honey bee viromes recovered a substantially larger number

of phage than did total metagenomes. Further, the occupancy-abundance relationships examined

here and by Medellin et al.[16] show that total metagenomes tend to be biased towards sampling

the most abundant and prevalent phage, while viromes are more successful at sampling rare

phage. Taken together, these results support previous work documenting that viromes produce a

deeper sampling of phage communities compared to total metagenomes[31,79,80].

431   Conclusions

By comparing bioinformatic mining of total metagenomes and targeted viromes across

two bee species, we found that sampling method significantly affected inferred phage

communities, but that the directionality of these differences can depend on the host being

sampled. In general, total metagenomes tended to be biased towards sampling both putatively

temperate phage and phage which were highly abundant. In samples with a relatively high

biomass (e.g., honey bees), viromes produced a greater diversity of phage and a better sampling

of rare phage. In contrast, when applied to relatively low biomass samples (e.g., bumble bees),

total metagenomes captured a greater diversity of phage than did viromes. Regardless of sample

biomass, there were always phage unique to both viromes and total metagenomes. In conclusion,

we suggest that these methods are complementary and recommend that both be used to capture

the full diversity of phage present in a gut environment. However, given that virome sampling is

not always feasible (i.e. in the case of field collected insects) total metagenomes may serve to

sample phage communities with the caveat that they will preferentially sample dominant and

temperate phage.

447   Acknowledgements

448   We thank Dr. Elina L. Nino, Joseph Tauzer, and the UC Davis Harry H. Laidlaw Jr.

449 Honey Bee Research Facility staff for help sampling managed honey bee colonies, Samantha R.

450   Eck and members of the Vannette, Emerson, and Brown labs for comments on the manuscript,

451   Anoushka Basu for help dissecting bees and the UC Davis Genome center for sequencing.

453   Conflict of interest

454   The authors have no conflict of interest to declare.

456   Funding

457   This research was supported in part by the Henry A. Jastro Scholarship and the United

458   States Department of Agriculture: National Institute of Food and Agriculture, Agriculture and

459   Food Research Initiative (award number 2023-67011-40501) awarded to DLS, as well as Hatch

460   Multistate funding to RLV and NSF DEB #1929516 to RLV.

462   Data Availability

463   The raw sequencing data generated in this study is available on NCBI under the

464   Bioproject accession number PRJNA1072153. All scripts used to process this data is publicly

465   available at https://github.com/dsbard/Bee_phage_2023.

<u>References</u>

1. Bergh, Ø., Børsheim, K. Y., Bratbak, G. & Heldal, M. High abundance of viruses found in aquatic environments. *Nature* **340**, 467–468 (1989).

2. Chibani-Chennoufi, S., Bruttin, A., Dillmann, M. L. & Brüssow, H. Phage-host interaction: An ecological perspective. *Journal of Bacteriology* vol. 186 3677–3686 Preprint at https://doi.org/10.1128/JB.186.12.3677-3686.2004 (2004).

3. Keen, E. C. A century of phage research: Bacteriophages and the shaping of modern biology. *BioEssays* **37**, 6–9 (2015).

4. Braga, L. P. P. *et al.* Impact of phages on soil bacterial communities and nitrogen availability under different assembly scenarios. *Microbiome* **8**, (2020).

5. Trubl, G. *et al.* Soil Viruses Are Underexplored Players in Ecosystem Carbon Processing. *mSystems* **3**, 1–21 (2018).

6. Emerson, J. B. *et al.* Host-linked soil viral ecology along a permafrost thaw gradient. *Nat Microbiol* **3**, 870–880 (2018).

7. Adriaenssens, E. M. *et al.* Environmental drivers of viral community composition in Antarctic soils identified by viromics. *Microbiome* **5**, (2017).

8. Fuhrman, J. A. *Marine Viruses and Their Biogeochemical and Ecological Effects*. *NATURE* vol. 399 www.nature.com (1999).

9. Suttle, C. A. Marine viruses - Major players in the global ecosystem. *Nat Rev Microbiol* **5**, 801–812 (2007).

10. Evans, C. *et al.* Shift from Carbon Flow through the Microbial Loop to the Viral Shunt in Coastal Antarctic Waters during Austral Summer. (2021) doi:10.3390/microorganisms.

489   11.   Danovaro, R. *et al.* Major viral impact on the functioning of benthic deep-sea ecosystems.

490   *Nature* **454**, 1084–1087 (2008).

491   12.   Tsao, Y. F. *et al.* Phage morons play an important role in pseudomonas aeruginosa

492   phenotypes. *J Bacteriol* **200**, (2018).

493   13.   Heyerhoff, B., Engelen, B. & Bunse, C. Auxiliary Metabolic Gene Functions in Pelagic

494   and Benthic Viruses of the Baltic Sea. *Front Microbiol* **13**, (2022).

495   14.   Dutilh, B. E. *et al.* A highly abundant bacteriophage discovered in the unknown sequences

496   of human faecal metagenomes. *Nat Commun* **5**, (2014).

497   15.   Townsend, E. M. *et al.* The Human Gut Phageome: Origins and Roles in the Human Gut

498   Microbiome. *Frontiers in Cellular and Infection Microbiology* vol. 11 Preprint at

499   https://doi.org/10.3389/fcimb.2021.643214 (2021).

500   16.   Santos-Medellin, C. *et al.* Viromes outperform total metagenomes in revealing the

501   spatiotemporal patterns of agricultural soil viral communities. *ISME Journal* (2021)

502   doi:10.1038/s41396-021-00897-y.

503   17.   Kosmopoulos, J. C., Klier, K. M., Langwig, M. V, Tran, P. Q. & Anantharaman, K.

504   Viromes vs. mixed community metagenomes: choice of method dictates interpretation of

505   viral community ecology. *bioRxiv* (2023) doi:10.1101/2023.10.15.562385.

506   18.   Zhang, F. *et al.* Critical Assessment of Whole Genome and Viral Enrichment Shotgun

507   Metagenome on the Characterization of Stool Total Virome in Hepatocellular Carcinoma

508   Patients. *Viruses* **15**, (2022).

509   19.   Sender, R., Fuchs, S. & Milo, R. Revised Estimates for the Number of Human and

510   Bacteria Cells in the Body. *PLoS Biol* **14**, (2016).

20.   Blum, W. E. H., Zechmeister-Boltenstern, S. & Keiblinger, K. M. Does soil contribute to

the human gut microbiome? *Microorganisms* **7**, (2019).

21.   Garren, M. & Azam, F. New method for counting bacteria associated with coral mucus.

Appl Environ Microbiol **76**, 6128–6133 (2010).

22.   Silveira, C. B. & Rohwer, F. L. Piggyback-The-Winner in host-Associated microbial

Communities. *npj Biofilms and Microbiomes* vol. 2 Preprint at

https://doi.org/10.1038/npjbiofilms.2016.10 (2016).

23.   Knowles, B. *et al.* Lytic to temperate switching of viral communities. *Nature* **531**, 466–

470 (2016).

24.   Mirzaei, M. K. & Maurice, C. F. Ménage à trois in the human gut: Interactions between

host, bacteria and phages. *Nature Reviews Microbiology* vol. 15 397–408 Preprint at

https://doi.org/10.1038/nrmicro.2017.30 (2017).

25.   Chin, W. H. *et al.* Bacteriophages evolve enhanced persistence to a mucosal surface.

(2022) doi:10.1073/pnas.

26.   Nelson, A. R. *et al.* Wildfire-dependent changes in soil microbiome diversity and function.

Nat Microbiol **7**, 1419–1430 (2022).

27.   Waller, A. S. *et al.* Classification and quantification of bacteriophage taxa in human gut

metagenomes. *ISME Journal* **8**, 1391–1402 (2014).

28.   Göller, P. C., Haro-Moreno, J. M., Rodriguez-Valera, F., Loessner, M. J. & Gómez-Sanz,

E. Uncovering a hidden diversity: Optimized protocols for the extraction of dsDNA

bacteriophages from soil. *Microbiome* **8**, (2020).

29.   Santos-Medellín, C. *et al.* Spatial turnover of soil viral populations and genotypes overlain

533   by cohesive responses to moisture in grasslands. *Proceedings of the National Academy of*

534   *Sciences* **119**, e2209132119 (2022).

535   30.   Parras-Moltó, M., Rodríguez-Galet, A., Suárez-Rodríguez, P. & López-Bueno, A.

536   Evaluation of bias induced by viral enrichment and random amplification protocols in

537   metagenomic surveys of saliva DNA viruses. *Microbiome* **6**, (2018).

538   31.   Garmaeva, S. *et al.* Studying the gut virome in the metagenomic era: Challenges and

539   perspectives. *BMC Biology* vol. 17 Preprint at https://doi.org/10.1186/s12915-019-0704-y

540   (2019).

541   32.   Gregory, A. C. *et al.* The Gut Virome Database Reveals Age-Dependent Patterns of

542   Virome Diversity in the Human Gut. *Cell Host Microbe* **28**, 724-740.e8 (2020).

543   33.   Avellaneda-Franco, L., Dahlman, S. & Barr, J. J. The gut virome and the relevance of

544   temperate phages in human health. *Frontiers in Cellular and Infection Microbiology* vol.

545   13 Preprint at https://doi.org/10.3389/fcimb.2023.1241058 (2023).

546   34.   Chandler, J. A., Lang, J., Bhatnagar, S., Eisen, J. A. & Kopp, A. Bacterial communities of

547   diverse Drosophila species: Ecological context of a host-microbe model system. *PLoS*

548   *Genet* **7**, (2011).

549   35.   Douglas, A. E. The molecular basis of bacterial-insect symbiosis. *Journal of Molecular*

550   *Biology* vol. 426 3830–3837 Preprint at https://doi.org/10.1016/j.jmb.2014.04.005 (2014).

551   36.   Pernice, M., Simpson, S. J. & Ponton, F. Towards an integrated understanding of gut

552   microbiota using insects as model systems. *J Insect Physiol* **69**, 12–18 (2014).

553   37.   Hammer, T. J., Le, E., Martin, A. N. & Moran, N. A. The gut microbiota of bumblebees.

554   *Insectes Soc* **68**, 287–301 (2021).

38. Kwong, W. K. & Moran, N. A. Gut microbial communities of social bees. *Nature Reviews*

Microbiology vol. 14 374–384 Preprint at https://doi.org/10.1038/nrmicro.2016.43 (2016).

39. Raymann, K., Shaffer, Z. & Moran, N. A. Antibiotic exposure perturbs the gut microbiota

and elevates mortality in honeybees. *PLoS Biol* **15**, 1–22 (2017).

40. Marceau, T., Archer, C. R., Bulson, L. & Wilfert, L. Dose-dependent effects of antibiotic

intake on Bombus Terrestris (Linnaeus, 1758) dietary intake, survival and parasite

infection prevalence. *J Invertebr Pathol* **182**, (2021).

41. Koch, H. & Schmid-Hempel, P. Socially transmitted gut microbiota protect bumble bees

against an intestinal parasite. *Proc Natl Acad Sci U S A* **108**, 19288–19292 (2011).

42. Engel, P. *et al.* The bee microbiome: Impact on bee health and model for evolution and

ecology of host-microbe interactions. *mBio* **7**, 1–9 (2016).

43. Zheng, H., Steele, M. I., Leonard, S. P., Motta, E. V. S. & Moran, N. A. Honey bees as

models for gut microbiota research. *Lab Anim (NY)* **47**, 317–325 (2018).

44. Kwong, W. K. Bee microbiomes go viral. *Proceedings of the National Academy of*

Sciences **117**, 11197–11199 (2020).

45. Bonilla-Rosso, G., Steiner, T., Wichmann, F., Bexkens, E. & Engel, P. Honey bees harbor

a diverse gut virome engaging in nested strain-level interactions with the microbiota. *Proc*

Natl Acad Sci U S A **117**, 7355–7362 (2020).

46. Deboutte, W. *et al.* Honey-bee-associated prokaryotic viral communities reveal wide viral

diversity and a profound metabolic coding potential. doi:10.1073/pnas.1921859117/-

/DCSupplemental.

47. Busby, T. J., Miller, C. R., Moran, N. A. & Van Leuven, J. T. Global Composition of the

Bacteriophage Community in Honey Bees. *mSystems* **7**, (2022).

48.   ter Horst, A. M., Fudyma, J. D., Sones, J. L. & Emerson, J. B. Dispersal, habitat filtering,

and eco-evolutionary dynamics as drivers of local and global wetland viral biogeography.

ISME J (2023) doi:10.1038/s41396-023-01516-8.

49.   Krueger, F. TrimGalore. *Github* https://github.com/FelixKrueger/TrimGalore.

50.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

51.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

9, 357–359 (2012).

52.   Pierce, N. T., Irber, L., Reiter, T., Brooks, P. & Brown, C. T. Large-scale sequence

comparisons with sourmash. *F1000Res* **8**, (2019).

53.   Guo, J. *et al.* VirSorter2: a multi-classifier, expert-guided approach to detect diverse DNA

and RNA viruses. *Microbiome* **9**, 1–13 (2021).

54.   Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: Accelerated for clustering the next-

generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

55.   Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–

2069 (2014).

56.   Terzian, P. *et al.* PHROG: Families of prokaryotic virus proteins clustered using remote

homology. *NAR Genom Bioinform* **3**, 1–12 (2021).

57.   Woodcroft, B. J. CoverM. *Github* https://github.com/wwood/CoverM.

58.   Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

Genome Biol **20**, (2019).

59.   Lu, J., Breitwieser, F. P., Thielen, P. & Salzberg, S. L. Bracken: Estimating species

abundance in metagenomics data. *PeerJ Comput Sci* **2017**, (2017).

60.   Christensen, S. M. *et al.* Microbial Metamorphosis: Symbiotic bacteria and fungi

proliferate during diapause and may enhance overwintering survival in a solitary bee.

bioRxiv 2023.11.02.565352 (2023) doi:10.1101/2023.11.02.565352.

61.   R Core Team. R: A Language and Environment for Statistical Computing. Preprint at

https://www.r-project.org/ (2019).

62.   Oksanen, A. J. *et al.* Veganl: community ecology package. R package vegan, vers. 2.2-1.

(2015).

63.   McMurdie, P. J. & Holmes, S. Phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, (2013).

64.   Bates, D., Mächler, M., Bolker, B. M. & Walker, S. C. Fitting linear mixed-effects models

using lme4. *J Stat Softw* **67**, (2015).

65.   Russell V. Lenth. emmeans: Estimated Marginal Means, aka Least-Squares Means.

https://CRAN.R-project.org/package=emmeans.

66.   Bolduc, B. *et al.* vConTACT: An iVirus tool to classify double-stranded DNA viruses that

infect Archaea and Bacteria. *PeerJ* **2017**, (2017).

67.   Bin Jang, H. *et al.* Taxonomic assignment of uncultivated prokaryotic virus genomes is

enabled by gene-sharing networks. *Nat Biotechnol* **37**, 632–639 (2019).

68.   Gilchrist, C. L. M. & Chooi, Y. H. Clinker & clustermap.js: Automatic generation of gene

cluster comparison figures. *Bioinformatics* **37**, 2473–2475 (2021).

69.   Kang, D. D. *et al.* MetaBAT 2: An adaptive binning algorithm for robust and efficient

genome reconstruction from metagenome assemblies. *PeerJ* **2019**, (2019).

70.   Skennerton, C. T. Minced. *Github* https://github.com/ctSkennerton/minced.

71.   Camacho, C. *et al.* BLAST+: Architecture and applications. *BMC Bioinformatics* **10**,

624   (2009).

625   72.   Handy, M. Y. *et al.* Incipiently social carpenter bees (Xylocopa) host distinctive gut

626   bacterial communities and display geographical structure as revealed by full-length

627   PacBio 16S rRNA sequencing. *Mol Ecol* **32**, 1530–1543 (2023).

628   73.   Hammer, T. J., Easton-Calabria, A. & Moran, N. A. Microbiome assembly and

629   maintenance across the lifespan of bumble bee workers. *Mol Ecol* **32**, 724–740 (2023).

630   74.   Wang, L. *et al.* Dynamic Changes of Gut Microbial Communities of Bumble Bee Queens

631   through Important Life Stages. (2019) doi:10.1128/mSystems.

632   75.   Kwong, W. K. *et al.* Dynamic microbiome evolution in social bees. *Sci Adv* **3**, e1600513

633   (2017).

634   76.   Hoyles, L. *et al.* Characterization of virus-like particles associated with the human faecal

635   and caecal microbiota. *Res Microbiol* **165**, 803–812 (2014).

636   77.   López-Pérez, M., Haro-Moreno, J. M., Gonzalez-Serrano, R., Parras-Moltó, M. &

637   Rodriguez-Valera, F. Genome diversity of marine phages recovered from Mediterranean

638   metagenomes: Size matters. *PLoS Genet* **13**, (2017).

639   78.   Gregory, A. C. *et al.* The Gut Virome Database Reveals Age-Dependent Patterns of

640   Virome Diversity in the Human Gut. *Cell Host Microbe* **28**, 724-740.e8 (2020).

641   79.   Emerson, J. B. Soil Viruses: A New Hope. *mSystems* **4**, (2019).

642   80.   Trubl, G., Hyman, P., Roux, S. & Abedon, S. T. Coming-of-age characterization of soil

643   viruses: A user's guide to virus isolation, detection within metagenomes, and viromics.

644   *Soil Systems* vol. 4 1–34 Preprint at https://doi.org/10.3390/soilsystems4020023 (2020).

Main Figures

Figure 1: Graphical representation of the sampling scheme and methods used in this research. A total of three bumble bee and three honey bee colonies were sampled. From each colony of bee, we generated three total metagenomes and one targeted virome. Total metagenomes were sampled from individual bees, while targeted viromes were produced from the pooled guts of 100 bees. This sampling resulted in nine bumble bee total metagenomes (light blue), three bumble bee viromes (dark blue), nine honey bee total metagenomes (light orange), and three honey bee viromes (dark orange).

Figure 2: Figures describing sequencing and assembly quality. Color denotes sample groups. **A)** Boxplots describing sequencing depth. The x-axis shows sample groups. The y-axis displays the total number of high-quality reads produced. **B)** Line plots describing sequence complexity of the reads shown in panel **A**, as measured by the frequency distributions of 31bp sized k-mers. The x-axis presents k-mer occurrence (i.e. how abundant a particular k-mer was in a given sample's read), while the y-axis shows the number of k-mers with a certain occurrence. **C)** Boxplots describing the number of unique 31bp sized k-mers present in the read libraries of each group of samples. The x-axis shows sample groups. The y-axis shows the number of unique k-mers. **D)** Jittered dot-plots describing the length distributions of contigs assembled from each sample. Only contigs >= 1Kbp bp are shown. Individual samples are shown on the y-axis. The x-axis shows contig length. Each point represents a single contig. A dotted red line is drawn at 5Kbp. For each sample, a green square describes phage enrichment (the number of phage identified, divided by number of contigs assessed for phage, times 100).

Figure 3: Figures describing phage identified in our dataset, relative to those identified previously, and the overall phage communities found in individual samples. **A)** Weighted gene-sharing network of all non-singleton vOTUs identified in our study. Individual nodes are phage. Nodes are connected by edges when phage share genes. Nodes are colored based on the predicted bacterial host of phage. Phage with "Known" as a predicted host are previously described bee phage. **B)** Stacked bar-plot describing the community of phage found in each sample based on predicted host. Both panel **A** and **B** use the same color palette.

Figure 4: Figures describing the alpha and beta diversity of the phage communities identified in our samples. Blue represent samples taken from bumble bees. Orange represents samples taken from honey bees. Lighter colors are total metagenomes. Darker colors are viromes. **A)** Boxplots describing the **A)** Shannon's diversity and **B)** vOTU richness associated with each of our sample types. The x-axis groups samples. The y-axis shows diversity and richness scores. **C)** Euler diagram describing phage community overlaps between each of our sample types. Numbers correspond to the number of phage present in each section of the graph (i.e. 290 phage were found in both honey bee viromes and total metagenomes). Circle size is proportional to number of phage. **C)** Non-metric multidimensional scaling (NMDS) ordination describing the Bray-Curtis dissimilarity of all samples on our dataset. Individual points are samples. Color represents sample type. The closer two points are to each other, the more similar they are in phage community.

Figure 5: Honey and bumble bees differ in bacterial community density, diversity, and composition. **A)** Boxplots displaying the 16S rRNA gene copy number present in the mid-hindgut region of sampled honey and bumble bees. **B)** Boxplot presenting the number of observed bacterial species in honey and bumble bee. **C)** Boxplot showing the species level Shannon's diversity of honey and bumble bee gut bacterial communities. **D)** Stacked bar plot describing honey and bumble bee bacterial communities at the genus level.

Figure 6: vOTUS sampled by viromes and total metagenomes differ in integrase content, occupancy, and abundance. **A)** Stacked bar plot showing the relative abundance of temperate and lytic vOTUs in virome and total metagenome samples. **B)** Density distributions showing the average relative abundance of all vOTUs found in viromes. vOTUs unique to viromes are shown in blue. vOTUs found in both viromes and total metagenomes are shown in red. **C)** Dot plot showing the occupancy-abundance relationship of all virome vOTUs. The x-axis describes average relative abundance, while the y-axis shows the percent occupancy of each vOTU. As before, vOTUs unique to viromes are shown in blue and those found in both viromes and total metagenomes are shown in red. **D)** Stacked bar plot describing the percent of vOTUs found in one, two, or three samples. The x-axis represents the percent of vOTUs. The y-axis the occupancy. Similar to B and C, vOTUs unique to viromes are shown in blue and those found in both viromes and total metagenomes are shown in red