One to rule them all? Assessing the performance of sustainable forest management indicators against multitaxonomic data for biodiversity conservation

Impact statement

Biodiversity indicators used to assess the state of Europe's forests perform unequally; a combination of indicators is more informative

Keywords

Sustainable forest management; taxonomic indicators; multi-taxa diversity; forest structure

Authors

Yoan Paillet, Univ. Grenoble Alpes, INRAE, Lessem, 2 rue de la Papeterie, BP76, 38402 Saint Martin d'Hères, cedex, France, yoan.paillet@inrae.fr, 0000-0001-7232-7844

Livia Zapponi, National Research Council, Institute of BioEconomy, Via Francesco Biasi 75, 38098 S. Michele all'Adige (TN) Italy, livia.zapponi@cnr.it, 0000-0003-4033-9001

Peter Schall, University of Göttingen, Silviculture and Forest Ecology of the Temperate Zones Büsgenweg 1, 37077 Göttingen, Germany, peter.schall@forst.uni-goettingen.de, 0000-0003-4808-818X

Jean-Matthieu Monnet, Univ. Grenoble Alpes, INRAE, Lessem, 2 rue de la Papeterie, BP76, 38402 Saint Martin d'Hères, cedex, France, jean-matthieu.monnet@inrae.fr, 0000-0002-9948-9891

Christian Ammer, University of Göttingen, Silviculture and Forest Ecology of the Temperate Zones Büsgenweg 1, 37077 Göttingen, Germany, christian.ammer@forst.uni-goettingen.de, 0000-0002-4235-0135

Lorenzo Balducci, Sapienza University of Rome, Piazzale Aldo Moro, 5, 00185 Rome, Italy, lorenzo.balducci@uniroma1.it, 0000-0001-9111-5679

Steffen Boch, WSL Swiss Federal Research Institute, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland, steffen.boch@wsl.ch, 0000-0003-2814-5343

Gediminas Brazaitis, Vytautas Magnus University Agriculture Academy Studentu g. 11, 53361 Akademija, Lithuania, gediminas.brazaitis@vdu.lt, 0000-0003-0234-9292

Alessandro Campanaro, Council for Agricultural Research and Economics, Research Centre for Plant Protection and Certification Via Lanciola 12/a, Florence, Italy, alessandro.campanaro@crea.gov.it , 0000-0003-4043-6362

Francesco Chianucci, CREA - Research Centre for Forestry and Wood Viale Santa Margherita 80, Arezzo, Italy, fchianucci@gmail.com, 0000-0002-5688-2060

Inken Doerfler, Carl von Ossietzky Universität Oldenburg, Department Plant sociology and nature Conservation Ammerländer Heerstraße 114-118 26129 Oldenbur, Germany, inkendoerfler@gmail.com, 0000-0001-9624-1922

Markus Fischer, University of Bern, Institute of Plant Sciences Altenbergrain 21, 3013 Bern, Switzerland, markus.fischer@ips.unibe.ch, 0000-0002-5589-5900

Marion Gosselin, INRAE, UR EFNO Domaine des Barres, 45290 Nogent-sur-Vernisson, France, marion.gosselin@inrae.fr, 0000-0003-0161-5368

Martin M. Gossner, WSL Swiss Federal Research Institute, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland, martin.gossner@wsl.ch, 0000-0003-1516-6364

Jacob Heilmann-Clausen, Center for Macroecolgy, Evolution and Climate, Globe Institute, University of Copenhagen, Universitetsparken 15, 2100 København, Denmark, jheilmann-clausen@sund.ku.dk, 0000-0003-4713-6004

Jeňýk Hofmeister, Czech University of Life Sciences Prague, Department of Forest Ecology, Faculty of Forestry and Wood Sciences Kamýcká 129, 165 00 Praha - Suchdol, Czech Republic, jenyk.hofmeister@email.cz, 0000-0002-3915-5056

Jan Hošek, Ecological Services Tichá 784/4, 268 01 Hořovice, Czech Republic, hosek@ekologickesluzby.cz

Kirsten Jung, University of Ulm, Evolutionary Ecology and Conservation Genomics, Albert-Einstein Allee 11, 89069 Ulm, Germany, kirsten.jung@uni-ulm.de

Sebastian Kepfer-Rojas, University of Copenhagen, Department of Geosciences and Natural Resource Management Rolighedsvej 23, 1958 Frederiksberg, Denmark, skro@ign.ku.dk, 0000-0002-1681-2877

Peter   Odor, Hun-Ren Centre for Ecological Research, Institute of Ecology and Botany, Alkotmány u. 2-4., H-2163 Vácrátót; University of Sopron, Institute of Environmental Protection and Nature Conservation, Bajcsy-Zsilinszky u. 4., H-9400 Sopron, Hungary, odor.peter@ecolres.hu, 0000-0003-1729-8897

83   **Flóra   Tinya**, HUN-REN Centre for Ecological Research, Institute of Ecology and Botany
84   Alkotmány u. 2-4., H-2163 Vácrátót, Hungary, tinya.flora@ecolres.hu, 0000-0002-4271-9676
85
86   **Giovanni Trentanovi**, Research Institute on Terrestrial Ecosystems - National Research
87   Council Via Madonna del Piano n. 10 50019 Sesto Fiorentino (Florence), Italy,
88   giovanni.trentanovi@cnr.it, 0000-0001-8236-4163
89
90   **Giorgio Vacchiano**, University of Milan, Department of Agricultural and Environmental
91   Science via Celoria 2, 20133 Milan, Italy, gvacchiano@gmail.com, 0000-0001-8100-0659
92
93   **Kris   Vandekerkhove**, Research Institute for Nature and Forest (INBO), Gaverstraat 4,
94   9500 Geraardsbergen, Belgium, kris.vandekerkhove@inbo.be, 0000-0003-1954-692X
95
96   **Wolfgang W. Weisser**, Technical University of Munich, Terrestrial Ecology Research
97   Group, Department of Life Science Systems, School of Life Sciences Hans-Carl-von-
98   Carlowitz-Platz 2, 85354 Freising, Germany, wolfgang.weisser@tum.de, 0000-0002-2757-
99   8959
100
101 **Michael Wohlwend**, Albert-Ludwigs-Universität Freiburg, Chair of Wildlife Ecology and
102 Management, Stefan-Meier-Straße 76, 79106 Freiburg, Germany,
103 michael.wohlwend@wildlife.uni-freiburg.de, 0000-0001-7230-6058
104
105 **Sabina Burrascano**, Sapienza University of Rome, Piazzale Aldo Moro, 5, 00185 Rome,
106 Italy, sabina.burrascano@uniroma1.it, 0000-0002-6537-3313
107

Acknowledgements

This work was funded by the EU Framework Programme Horizon 2020 through the COST

Association (www.cost.eu): COST Action CA18207: BOTTOMS-UP – Biodiversity Of

Temperate forest Taxa Orienting Management Sustainability by Unifying Perspectives. The

authors are thankful to all those experts contributing to the data here harmonized and

resumed that were not listed as data contributors.

Abstract

Several regional initiatives and reporting efforts assess the state of forest biodiversity through broad-scale indicators based on data from national forest inventories. Although valuable, these indicators are essentially indirect and evaluate habitat quantity and quality rather than biodiversity *per se*. Besides, most of these indicators are applicable at regional or national scales, while their use at a more local level is difficult. Therefore, their link to biodiversity may be weak, which decreases their usefulness for decision-making.

For several decades, Forest Europe indicators assessed the state of European forests, in particular its biodiversity. However, no extensive study has been conducted to date to assess the performance – ie. their capacity to reflect variations in biodiversity – of these indicators against multitaxonomic data. We hypothesized that no single biodiversity indicator from Forest Europe can represent overall forest biodiversity, but that several – eventually combined – indicators would reflect habitat quality for at least some taxa in a comprehensive way. We tested the set of indicators proposed by Forest Europe against the species richness of six taxonomic and functional groups (tracheophytes, epixylic and epiphytic bryophytes, birds, saproxylic beetles, saproxylic non-lichenized fungi and epixylic and epiphytic lichenized fungi) across several hundreds of plots over Europe. We showed that, while some indicators perform relatively well across groups (e.g. deadwood volume), no single indicator represented all biodiversity at once, and that a combination of several indicators performed better. Surprisingly, some indicators showed weak links with the biodiversity of the six taxonomic and functional groups.

Forest Europe indicators were chosen for their availability and ease of understanding for most people. However, our analyses showed that there are still gaps in the monitoring framework, and that surveying certain taxa along with stand structure is necessary to support policymaking and tackle forest biodiversity loss at the large scale.

Keywords

Introduction

Monitoring biodiversity in the face of global change is a challenge in many ecosystems across the world (IPBES 2019; Lindenmayer and Likens 2010). Despite large-scale initiatives such as GEOBON (Group on Earth Observation Biodiversity Observation Network 2008) and collaborative networks (Burrascano et al. 2023), no standard biodiversity monitoring scheme has been agreed in Europe or elsewhere. Long-term biodiversity monitoring hence currently relies on a limited number of initiatives that are used to assess the impact of climate change (e.g. Jiguet et al. 2012), habitat loss (Betts et al. 2017) or pollution (Rigal et al. 2023). Initiatives combining monitoring of biodiversity, types of pressure and ecosystem-level variables remain quite rare, or may concern only single pressure vs. single taxon assessment (e.g. Proença et al. 2017; Weber et al. 2004), or may use different methods for overall assessments (European Environment 2020). However, different taxonomic and functional groups likely respond differently to a given pressure, which challenges prioritization of policy initiatives and tools (Pereira and David Cooper 2006).

Forest ecosystems are no exception to the rule, and lack broad-scale standardized biodiversity and pressure data assessment. In Europe, forestry is reported as one of the main pressures to forest habitats (European Environment 2020), and Sustainable Forest Management, i.e., managing forests to maintain and enhance their economic, social and environmental values, is one of the main assets to cope with this pressure (European Commission 2021). Expectations and targets, progress towards them, as well as different elements of sustainable forest management, e.g. novel management methods, best practices, are broadly defined and articulated in the framework of a series of agreed Criteria and Indicators that are applied from the global to the regional, national and subnational levels (Forest Europe 2015). Each criterion relates to a key element of sustainability, and may be described by one or more indicators.

Indicators are measurable parameters that contribute to monitor the status and changes of forests in quantitative, qualitative and descriptive terms in relation to each criterion (FAO 2024). In the Sustainable Forest Management framework, indicators are developed

accounting for: 1) a review of existing forest information; 2) 'Bottom-up' stakeholder engagement, i.e., priorities for forest sector performance; 3) 'Top down' alignment to international Criteria and Indicators (Linser and O'Hara 2019).

Given this basis, it comes as no surprise that National Forest Inventories were the first choice to develop nationwide indicators of Sustainable Forest management (Simons et al. 2021), since they are widely available (point 1), and were originally designed to assess wood stock and production, thus may indicate the sector performance (point 2), even if they do not provide a comprehensive indication for the international criteria linked to sustainability (point 3). At the European level, data from National Forest Inventories have been aggregated and published every five years for more than three decades (Forest Europe 2020) and provide a set of indicators to inform sustainable forest management. For European forests, the Criterion 4 "*Maintenance, conservation and appropriate enhancement of biological diversity in forest ecosystems*" is the one that should help decision makers to assess progress towards biodiversity-friendly forest management.

As for other criteria, the desired indication on biodiversity sustainability is not ensured, since most of the indicators are indirect (structural) proxies derived from National Forest Inventories or land cover maps (e.g. deadwood or forest fragmentation), while only two indicators involve other species than trees (namely 4.8 Threatened forest species, 4.10 Common forest bird species, Table 1). This approach follows the general trend of conservation programs and researches of relying heavily on habitat attributes and area-based metrics, mainly in relation to the relative ease with which these measures can be collected and analyzed, compared with complex ecological or population processes (Marshall et al. 2020). As a matter of fact, despite the large use of National Forest Inventories data to indirectly assess the state and evolution of biodiversity (Chirici et al. 2012; Heym et al. 2021; Reise et al. 2019), these were often found to lack links with forest biodiversity (Gao et al. 2015; Paillet et al. 2018; Storch et al. 2023; Zeller et al. 2022). Despite a large corpus of individual studies and few syntheses (Zeller et al. 2023), no global and systematic assessment of the correlations between multi-taxonomic biodiversity and

Forest Europe indicators has been attempted. This is particularly worrying since the
European system of Criteria and Indicators is widely used as a reference across several
regions, as well as for global initiatives (Linser and O'Hara 2019).

In this context, our goal was to provide an assessment of the performance of biodiversity
indicators. Here, we define performance as the capacity of the indicator to reflect the
variations of one or more *indicanda*. This encompasses the magnitude of the correlation
between a given indicator and an *indicanda*, but also the respective fit of each indicator for a
given indicanda as well as the relative fit of individual vs. combined indicators. We intended
to support potential improvements in the global reporting processes for biodiversity
conservation and sustainable forest management. In other words, we aimed to: i) assess the
link between indicators and several *indicanda* (i.e. taxonomic and functional groups) based
on a large dataset combining species and forest characteristic data; ii) identify the indicators
that performed best and universally - i.e. for all groups; iii) define, if possible, the most
effective combination of indicators for forest biodiversity.

To address these aims, we tested the performance of Forest Europe's biodiversity indicators
against six taxonomic and functional groups. We used a three step approach: we first tested
individual indicators against the species richness of the six groups to assess the magnitude
of the links between indicators and *indicanda*; second, we used a model selection approach
to assess whether one single indicator outperformed the others; third, we tested whether a
combination of indicators performed better than single ones by searching for the best and
most parsimonious combination of biodiversity indicators. We used a multi-taxonomic
database (Burrascano et al. 2023; Trentanovi et al. 2023) combining species inventories and
forest structure to analyze the correlations between Forest Europe's biodiversity indicators
with the biodiversity of six groups (tracheophytes, epixylic and epiphytic bryophytes, birds,
saproxylic beetles, saproxylic non-lichenized fungi and epixylic and epiphytic lichenized
fungi).

Materials and Methods

Database

We used the database gathered within the framework of the COST action "BOTTOMS-UP"

(CA18207 – Biodiversity Of Temperate forest Taxa Orienting Management Sustainability by

Unifying Perspectives). In a nutshell, this unique database spatially combines sampling units

for both forest measurements and species of at least 3 taxa out of which at least one animal

(Burrascano et al. 2023). It merges 34 different datasets from 12 European countries and

more than 3500 sampling units. It covers a wide range of forest categories and

environmental conditions but without apparent bias concerning the temperature and

precipitation (Appendix 1, see Burrascano et al. 2023 for further details).

It is particularly well suited to test correlations between Forest Europe's biodiversity

indicators and biodiversity, since the former can be extracted either from forest

measurement, species inventories themselves as well as external (cartographic) data. From

this database, we extracted the six most represented taxonomic and functional groups,

namely tracheophytes, epixylic and epiphytic bryophytes (hereafter bryophytes), birds,

saproxylic beetles, saproxylic non-lichenized fungi (hereafter fungi) and epixylic and

epiphytic lichenized fungi (hereafter lichens).

241
Figure 1: Spatial distribution of the sampling sites. Gray areas are covered by forests with a

tree cover greater than 40% according to Kempeneers et al. (2011). The size of the dot is

proportional to the number of sampling units.

While tracheophytes and birds were inventoried without any specific selection of the guilds

targeted, only epiphytic and epixylic bryophytes and lichens were included (sampled on

living trees and deadwood), and saproxylic fungi and beetles (dependent on deadwood

substrates or on other organisms inhabiting deadwood).

Since the database is the result of the merging of different research projects, sampling

designs and protocols varied across datasets (Burrascano et al. 2023; Trentanovi et al.

2023). Therefore, we standardized species richness (number of species per sampling plot)

by dividing it by the asymptotic gamma richness at the site level, with site representing a

homogeneous sampling area with a maximum size of a few thousand hectares. We derived

site gamma diversity through plot-based rarefaction curves using the R package iNEXT

(Hsieh et al. 2016). Sites with less than 6 sampling units were discarded from the final

database since the estimation of the gamma richness was judged non-reliable. Although

species richness represents only a partial measure of biodiversity (Noss 1990), it was the

simplest, widely used and most reliable metric that could be standardized for sampling effort.

Exploratory analyses revealed especially large deadwood volumes linked with small plots

(nugget effect). To avoid strong leveraging from these outliers (Zuur et al. 2010), we limited

the maximum deadwood volume per sampling plot to 500 m^3/ha, a value that corresponds to

the maximal volumes observed in primeval forest of Europe (e.g. Christensen et al. 2005).

The final data distribution per taxonomic and functional groups is shown in Table 1 and

Figure 1.

Taxa	I4.2.Regeneration			I4.3.Naturalness			Total
	Coppice	Natural	Planting	Plantation	Semi-natural	Unmanaged	
Tracheophytes	10	734	94	94	550	194	838
Bryophytes	3	353	66	66	198	158	422
Beetles	7	408	94	94	255	160	509
Birds	10	721	94	94	537	194	825
Fungi	8	502	94	94	347	163	604
Lichens	10	332	92	92	333	9	434

Table 1: Number of plots used to fit the models distributed per taxonomic and functional

groups and Regeneration origin or Naturalness (Forest Europe 2020).

Indicators' calculation

We used the State of Europe's Forests (Forest Europe 2020) definitions to calculate the

corresponding metrics for the indicators included in the Criterion 4 (Table 2). Since it was

necessary to "translate" the definition into calculable values to analyze the relationship

between indicator values and *indicanda* (i.e. standardized species richness), we proceeded

as follows:

-    indicator 4.6 *Genetic resources* was excluded since it was not possible to translate

this indicator at the plot or site levels;

-    indicators 4.1 *Diversity of tree species*, 4.4 *Introduced tree species* and 4.5

Deadwood were directly calculated from the plot measurements. Instead of

introduced tree species, we considered only invasive non-native woody species that

can have a significant impact on forest biodiversity following Campagnaro et al.

(2018), namely *Acer negundo*, *Ailanthus altissima, Prunus serotina*, *Quercus rubra*

and *Robinia pseudoacacia*. We limited to these species since there was no regional

reference list for introduced tree species and it would have been far beyond the

scope of this study to create such a list, e.g. Norway spruce (*Picea abies*) is

introduced in lowland France but native in mountains where it has also been planted

in some places. In addition, no reference list of invasive tree species in Europe was

available in the Forest Europe (2020) report;

-    indicators 4.2 *Regeneration* and 4.3 *Naturalness* were derived from declarative

metadata when merging the database (Burrascano et al. 2023). In the 4.3, forests

were considered 'undisturbed by man' when declared without intervention by the data

holder (i.e. 'abandoned', 'unmanaged', 'primeval'). Time since the last intervention

was not used here since it was not homogeneously declared. 'Semi-natural' refers to

forest plots that are under any type of forest management regime (from even-aged to

uneven-aged silvicultural systems), but with natural regeneration processes of trees.

'Plantation' forests overlap with forest plots where regeneration derives from

'Planting' activities;

-    indicators 4.8 *Threatened forest species* and 4.10 *Common forest bird species* were

derived from biodiversity sampling and reference lists. For 4.8, we used the IUCN

Red List (https://www.iucnredlist.org/) and calculated the total number of threatened

species (categories VU, EN and CR) per plot all groups together. For 4.10, we

calculated the total number per plot of the 34 forest birds classified by the Pan-

European Common Birds Monitoring Scheme (https://pecbms.info/trends-and-

indicators/indicators/);

-    indicator 4.7 *Forest fragmentation* was calculated following the protocol applied by

European Commission et al. (2019) on the Corine Land Cover 2018 layer

(https://efi.int/knowledge/maps/forest). The forest area density (FAD) at fixed

observation scale was obtained calculating the percentage of forest pixels in a 500

ha circular buffer, centered on each plot;

-    indicator 4.9 *Protected forests* was calculated based on the map of protected areas

in Europe provided by the European Environment Agency (Nationally Designated

areas - CDDA: https://www.eea.europa.eu/data-and-maps/data/nationally-

designated-areas-national-cdda-17). We extracted the proportion of Categories Ia, Ib

and II in a 500ha circular buffer around each plot. We then added these three values

so that the total proportion may be higher than 100.

Biodiversity indicators	Definition (Forest Europe 2020)	Data source	Metrics
4.1. Diversity of tree species	Area of forest and other wooded land, classified by number of tree species occurring	Plot measurements	Number of tree species per sampling unit
4.2. Regeneration	Total forest area by stand origin and area of annual forest regeneration and expansion (natural regeneration, planting, coppicing)	Metadata	3 classes: Natural regeneration Planting Coppicing
4.3. Naturalness	Area of forest and other wooded land by class of naturalness (undisturbed by man, semi-natural, plantations)	Metadata	3 classes: Unmanaged Semi-natural Plantation
4.4. Introduced tree species	Area of forest and other wooded land dominated by introduced tree species □ modified to invasive ligneous species (Campagnaro et al. 2018)	Plot measurements	Number of invasive tree species per sampling unit
4.5. Deadwood	Volume of standing deadwood and of lying deadwood on forest and other wooded land	Plot measurements	Deadwood volume per ha
4.6. Genetic resources	Area managed for conservation and utilization of forest tree genetic resources (in situ and ex situ genetic conservation) and area managed for seed production	Not assessed	
4.7. Forest fragmentation (FAD)	Area of continuous forest and of patches of forest separated by non-forest lands (European Commission et al. 2019)	European Forest Institute map of forest cover	Forest area density (FAD) within 500ha around the sampling unit
4.8. Threatened forest species	Number of threatened forest species, classified according to IUCN Red List categories, in relation to total number of forest species	IUCN Red List species list	Species richness of threatened species (categories VU, EN and CR)
4.9. Protected forests	Area of forest and other wooded land protected to conserve biodiversity, landscapes and specific natural elements, according to MCPFE categories	CDDA map of protected areas	Cumulative percentage of Ia, Ib and II categories areas
4.10. Common forest bird species	Occurrence of common breeding bird species related to forest ecosystems	PECBMS	Species richness of forest birds

Table 2: Forest Europe's biodiversity indicators definitions and data sources used to

calculate them in this paper.

Statistical analyses

We processed all analyses in R v.4.3.0 (R Core Team 2023). We used the standardized

species richness of each taxonomic and functional group as a response variable. We fitted a

generalized linear mixed model with site as a random effect on the intercept to account for

the fact that two plots from the same site were more likely to be similar than two plots from

different sites. We used the beta error distribution with logit link since standardized richness

was continuous, strictly positive with a maximum value below one. We scaled continuous

explanatory variables to improve the convergence and to make the estimates of the models

comparable. We used the glmmTMB function in the glmmTMB package (Brooks et al. 2017).

We processed the data in three steps:

- first, for each group, we fitted nine single models plus the null (intercept only) model

with each indicator as an explanatory variable (Table 2). We compared the

magnitudes of all single effects in the models and their significance and represented

this using a heatmap of the standardized effects (package ggplot2, Wickham 2016).

We tested differences between estimates of categorical variables (e.g.

4.3.naturalness) using Tukey post-hoc test. To search for potential non-linear

relationships between indicators and *indicanda*, we also fitted generalized additive

mixed models (GAMMs) with indicators as smoothers (package mgcv, Wood 2023).

We found very few differences between GLMMs and GAMMs and preferred to stick

to the former (comparisons are presented in Appendix 2).

- second, we assessed the relative performance of all single indicators by comparing

the Akaike Information Criterion corrected for small samples (AICc, Burnham and

Anderson 2002) of all single models including a null (intercept only) model.

- third, we searched for the best and most parsimonious linear combination (no

interaction) of indicators that represented biodiversity based on AICc using the

dredge function(package MuMIn, Barton 2023). In this process, we discarded the

indicator 4.2. Regeneration since it was collinear with 4.3. Naturalness and a model

containing both variables could not be fitted (see Appendix 2). When two competing

models had a difference in AICc less than 2 points, we chose the most parsimonious

one.

Results

Data and indicators' distribution

The most represented group in the database was tracheophytes (838 plots) followed by

birds (825 plots, Table 3, see Burrascano et al. 2023 for a full description of the biodiversity

dataset). All taxonomic and functional groups have been inventoried in at least 400 plots

(Table 1). The distribution in the classes Regeneration (4.3.) and Naturalness (4.4.) were,

however, strongly unbalanced (see also Table 1): the majority of sampling units were

associated with "natural regeneration", while "coppicing" - and "planting" to a lesser extent -

were underrepresented; the majority of sampling units were within semi-natural forests, but

the distribution was more balanced than for regeneration types. For quantitative indicators,

the values taken were relatively balanced between groups (Table 3) and no strong

collinearity was observed (see Appendix 3).

Taxonomic and functional groups datasets	I4.1.tree.sp (species number)	I4.4.invasive (species number)	I4.5.deadwood (m^3/ha)	I4.7.fragmentation (FAD) (%)	I4.8.threat.sp (species number)	I4.9. Protected forests (%)	I4.10.birds (species number)
Tracheophytes (n = 838)	1.96(2.06) [0-11]	0.01(0.08) [0-1]	41.83(60.72) [0-444.89]	0.85(0.16) [0.13-1]	0.25(0.56) [0-3]	14.92(33.66) [0-142.4]	3.63(1.78) [1-11]
Bryophytes (n = 422)	1.91(1.84) [0-11]	0(0.06) [0-1]	39.15(55.86) [0-461.41]	0.84(0.18) [0.06-1]	0.2(0.5) [0-3]	27.02(43.16) [0-142.4]	4.41(2.63) [1-15]
Beetles (n = 509)	1.96(2.01) [0-11]	0(0.07) [0-1]	41.1(57.2) [0-444.89]	0.83(0.17) [0.08-1]	0.26(0.59) [0-5]	16.63(35.16) [0-142.4]	4.15(2.43) [1-12]
Birds (n = 825)	2.49(2.18) [0-11]	0.01(0.08) [0-1]	42.03(64.56) [0-461.41]	0.85(0.16) [0.13-1]	0.29(0.6) [0-3]	15.92(34.06) [0-142,4]	4.14(2.38) [1-15]
Fungi (n = 604)	1.47(1.95) [0-11]	0.01(0.08) [0-1]	39.73(57.98) [0-461.41]	0.82(0,.8) [0.08-1]	0.17(0.49) [0-3]	14.55(34.22) [0-100]	4.32(2.83) [1-15]
Lichens (n = 434)	1.93(1.85) [0-11]	0(0.06) [0-1]	39.28(56.21) [0-461.41]	0.84(0.18) [0.06-1]	0.21(0.53) [0-5]	26.65(42.93) [0-142.4]	4.43(2.62) [1-15]

Table 3: Range of values for each indicator distributed over the different taxonomic and

functional datasets (mean per sampling unit (sd) [min - max]) . FAD = Forest Area Density.

Links between single indicators on indicanda

Several of the Forest Europe indicators had significant relationships with the standardized

species richness of one or more of the six taxonomic and functional groups (Figure 2, see

Appendix 3 for values of the estimates).

Figure 2: Heatmap representing the standardized estimates (slope) of standardized species

richness obtained with generalized mixed models with Beta error distribution and logit link,

using Forest Europe indicators as independent predictors. S.N = semi-natural, UNM =

unmanaged. Intercept (reference) for 4.2 Regeneration is the "Coppice" class, and

"Plantations" for 4.3 Naturalness (p<0.1, * p<0.05, ** p<0.01, *** p<0.001).

Figure 3: Relationships between standardized species richness of 6 taxonomic and functional groups and Forest Europe indicator 4.5. Total deadwood volume. Estimations are issued from generalized linear models with Beta error distribution and logit link. Plain line represents the mean estimate, ribbons the 95% confidence interval.

Deadwood (4.5, Figure 3) and Diversity of tree species (4.1) showed four positive and (marginally) significant but generally weak relationships with tracheophytes, bryophytes, fungi and lichens (Appendix 4). Other indicators showed at least three significant (or marginally significant) relationships, with varying magnitudes, i.e. Fragmentation (FAD) (4.7),

Protected areas (4.9, Figure 4) and Forest Birds (4.10). Surprisingly, the proportion of

protected areas around sampling plots had a negative effect on birds and no effect on

tracheophytes and fungi. Regarding Naturalness (4.3, Figure 5), unmanaged forest and

semi-natural forests showed higher levels of fungi, lichens and bryophytes than plantations.

The opposite was true for tracheophytes. Threatened species (4.8) showed very few

significant results (marginally positive for tracheophytes and bryophytes, positive for birds)

while Invasive species (4.4) only showed a marginally significant positive relationship with

tracheophytes. Plantation (4.2) showed a very strong negative relationship with lichens only,

and a marginally significant negative effect on tracheophytes, while there were no

differences between coppice and natural regeneration (Appendix 3).

Figure 4: Relationships between standardized species richness of 6 taxonomic and
functional groups and Forest Europe indicator 4.9 (Proportion of protected areas in a 500ha
buffer). Estimations are issued from generalized linear models with Beta error distribution.
Plain line represents the mean estimate, ribbons the 95% confidence interval.

Figure 5: Relationships between standardized species richness of 6 taxonomic and functional groups and Forest Europe indicator 4.3. Naturalness: PLA = Plantations; S.N = Semi – Natural forests; UNM = Unmanaged forests. Estimations are issued from generalized linear models with Beta error distribution and logit link. Barplots represent mean estimates, error bars are the 95% confidence intervals. Letters indicate significant differences per group based on Tukey post-hoc test.

Relative performance of the different indicators

Comparing the AICc (Table 4) for all single models revealed that the indicator based on

forest Naturalness (4.3) best explained the scaled species richness for three out of six

groups (tracheophytes, bryophytes, and lichens); Regeneration (4.2) was within 2 points of

AICc for tracheophytes and lichens. For birds and beetles, the indicator based on Forest

birds (4.10) was the best explanatory one, while for fungi, Deadwood (4.5) stood first. Also

note that the null model was never the best one and more than 2 AICc points away from the

best.

Explanatory variables	K	Tracheophytes Delta AICc	Bryophytes Delta AICc	Birds Delta AICc	Beetles Delta AICc	Fungi Delta AICc	Lichens Delta AICc
Null	3	14.5	76.9	626.2	2.6	123.1	27.4
I4.1.tree.sp	4	13.3	59.9	627.1	4.2	107.1	18.4
I4.2.regeneration	5	0.5	23.5	629.5	6.0	73.3	1.6
I4.3.naturalness	5	**0.0**	**0.0**	629.7	2.7	33.9	**0.0**
I4.4.invasive	4	13.0	77.8	627.3	4.1	125.1	29.3
I4.5.deadwood	4	14.4	52.3	624.2	4.6	**0.0**	19.3
I4.7.fragmentation (FAD)	4	4.4	70.8	610.9	1.9	121.7	29.3
I4.8.threat.sp	4	12.8	75.2	623.8	4.4	124.1	28.5
I4.9.protected forests	4	16.4	64.0	622.1	2.5	124.4	21.6
I4.10.birds	4	16.3	78.7	**0.0**	**0.0**	115.4	29.4

Table 4: Differences of Akaike Information Criteria corrected for small samples (Delta AICc)

for all generalized mixed models with Beta error distribution and logit link. K is the number of

parameters in each model. The best model (delta AICc = 0) is in bold.

Combination of indicators

We used data dredging to find the best linear (i.e. without interaction) combination of

indicators reflecting the variations in species richness of each group. Only beetles were

represented by only one indicator (Forest Birds, 4.10.), all the other ones had a combination

of 3 to 4 indicators in their best models (Table 5). In terms of indicators, Invasive species

(4.4.) was never selected. Conversely, and as observed before, Diversity of tree species

(4.1.), Naturalness (4.3.), Deadwood (4.5.) were selected four times, followed by Forest

Birds (4.10, 3 times). Finally, it is interesting to note that, except for Fragmentation (FAD,

4.7.), all indicators had positive effects, and with the remarkable exception of beetles, all

multivariate models performed better than single ones (in terms of AICc, they had at least 2

points less than single variable ones, Table 5).

Taxonomic and functional groups	Tracheophytes	Bryophytes	Birds	Beetles	Fungi	Lichens
Intercept	-0.47	-1.28	0.11	-1.03	-2.14	-1.34
I4.1.tree.sp	0.11	0.17			0.12	0.12
I4.3.naturalness	+	+			+	+
I4.4.invasive						
I4.5.deadwood	0.06	0.09			0.17	0.07
I4.7.fragmentation (FAD)	-0.072		-0.07			
I4.8.threat.sp			0.05			
I4.9.protected forests						
I4.10.birds			0.65	0.05	0.08	
Df	8	7	6	4	8	7
AICc	-1246.18	-384.61	-1678.81	-1123.87	-1743.14	-547.79
Delta AICc	0.61	1.42	0.19	0.96	1.23	1.69
AICc (best single variable models)	-1226.72	-375.82	-1648.93	-1123.87	-1671.48	-541.632

Table 5: Scaled estimates of the models selected by data dredging combining all indicators

from Forest Europe (without interaction). Estimates are issued from generalized linear mixed

models with Beta error distribution and logit link. "+" indicates the presence of the factor in

the model. The best model was the most parsimonious (lowest degrees of freedom - df) with

the lowest Akaike Information Criterion corrected for small samples (AICc). "delta" indicates

the AICc difference with the best model. logLik = logarithm of the likelihood.

Discussion

We analyzed the links between Forest Europe's biodiversity indicators and the standardized

richness of six taxonomic and functional groups on a unique multi taxonomic dataset

covering most forest types in Europe (Burrascano et al. 2023, Appendix 1). We showed that

these indicators perform unequally: while some correctly correlated with the biodiversity of

several groups, no one acted as a universal indicator, questioning their strength to predict

biodiversity in European forests. In contrast, our results reinforced the approach based on

multiple complementary indicators of the same *indicandum*. This also raises the question of

contextualization of these indicators, to better assess state and trends of biodiversity across

Europe, and opens perspectives for improvement.

Performance of current forest biodiversity indicators

While evidence on the link between some indicators and the biodiversity they are supposed

to indicate remains incomplete (Gao et al. 2015; Penone et al. 2019; Zeller et al. 2023), our

results highlighted several habitat-species relationships and habitat preferences of different

groups.

We first observed strong effects of Regeneration origin (4.2) and Naturalness (4.3) on the

biodiversity of the six groups. The biodiversity of bryophytes and fungi was higher in

unmanaged forests compared to semi-natural forests, and, including lichens, higher than in

plantation forests (Figure 5). On the contrary, plantations had a marginal positive effect on

tracheophytes compared to semi-natural and unmanaged forests, and naturalness had no

effects on birds and beetles. The negative effects of plantations, and the positive effects of

management abandonment or primeval forests have been thoroughly documented (e.g.

Chaudhary et al. 2016; Paillet et al. 2010). However, our results should be nuanced by the

fact that the distribution of the data in the different categories of regeneration and

naturalness were strongly unbalanced, with only few plots located in coppice-managed

forests and a vast majority in semi-natural forests (Table 1). That said, unmanaged forests

had a generally positive influence on biodiversity in our dataset, especially for deadwood

dependent species (epixylic bryophytes and lichens), while plantations had detrimental

effects on several other groups. Indeed, habitat provision and continuity are higher in semi-

natural forests, not to speak of unmanaged forests, which allow the persistence of dispersal

limited species (e.g. Boch et al. 2013; Boch et al. 2021). The positive response of

tracheophytes in plantations may seem surprising, but probably reflects a higher share of

disturbance tolerant herbs in more disturbed sites, as shown in several individual studies

(Boch et al. 2013). In addition, the semi-natural category encompasses a wide range of

management types (Trentanovi et al. 2023), and unmanaged forests in our dataset have

mostly been recently abandoned, which may cause a decrease in the biodiversity of

tracheophytes during the first decades after abandonment (e.g. Paillet et al. 2010).

Deadwood volume, followed by diversity of tree species, were two indicators that often

correlated with scaled species richness, both in univariate and multivariate analyses.

Deadwood and diversity of tree species have complementary roles for biodiversity (Storch et

al. 2023). Deadwood provides a substrate and a resource for numerous species that depend

on it for part of their life cycle (Lassauce et al. 2011; Müller et al. 2019; Müller et al. 2015;

Parajuli and Markwith 2023), but also for facultative species (Graf et al. 2022). Indeed, the

richness of bryophytes, fungi and lichens, and to a lesser extent birds, correlated positively

to deadwood volume in our analyses. Surprisingly however, saproxylic beetles richness did

not correlate significantly to deadwood volume despite a weak positive effect. It is likely that,

for this group in particular, deadwood does not act as a universal indicator that could be

transposed in all situations (see also Zeller et al. 2023). For example, Lassauce et al. (2011)

showed that the correlation between richness of saproxylic beetles and deadwood volume

was higher in boreal than in temperate forests, a result that was however not confirmed by

Parajuli and Markwith (2023). Müller et al. (2015) showed compensation effects between

deadwood and temperature, supporting that the influence of deadwood on saproxylic beetle

richness declines with increasing temperature, both in geographical space, but also locally in

relation to canopy openness (e.g. Koch Widerberg et al. 2012). It is also possible that the

observed deadwood volumes (mean 40 m^3/ha, Table 3) lead to relatively poor beetle

communities. However, in most cases, the mean deadwood values in our dataset were

above the average deadwood volume per European country (which was 11.5 m3/ha on

average, but varied from 2.3 to 28 m3/ha between countries, Forest Europe 2020).

Consequently, while the positive role of deadwood for biodiversity remains unchallenged, its

use as an indicator should probably be refined by a better integration of the context

(macroclimate, microclimate, position on the deadwood volume gradient) in the relation. It is

also true that deadwood quality (e.g. including size and decay stage, Vítková et al. 2018) is

as much relevant as its quantity. Some saproxylic beetle species (e.g. the red-listed species

Rosalia alpina, Campanaro et al. 2017) develop in large logs, snags and standing dead trees

within canopy gaps. In most of the sampling units we analyzed, deadwood was represented

by relatively small fragments (mostly around 15-20 cm diameter) likely deriving from

selective thinning processes occurring in closed forests (Burrascano et al. 2023). Further,

many rare taxa depend on the occurrence of tree-related microhabitats (Larrieu et al. 2018),

such as the protected umbrella species *Osmoderma eremita,* which larvae develop in large

cavities in living trees (Dubois et al. 2009). Since several tree-related microhabitats are very

scarce in managed forests (Paillet et al. 2017), and not readily quantified as part of the

deadwood pool (Müller et al. 2014), their role is not currently taken into consideration.

Diversity of tree species also showed positive correlation with the biodiversity of several

taxonomic and functional groups (bryophytes, fungi, lichens and more marginally

tracheophytes). Different tree species provide different habitat conditions for epiphytic and

saproxylic species living in forests (e.g. Boch et al. 2021; Cavard et al. 2011; Leidinger et al.

2021). These conditions encompass direct biotic interactions, different chemical properties of

the bark and wood, decomposition rates as well as differential light interception (Gosselin et

al. 2017; Zeller et al. 2023). This in turn provides niche heterogeneity for several species and

thus enhances biodiversity and confirms the role of diversity of tree species as a biodiversity

indicator.

Among the other indicators, forest fragmentation showed negative effects on tracheophytes

and birds, and a positive effect on bryophytes, and marginally on beetles and fungi. It should

be noted that since it is based on forest area density (European Commission et al. 2019), it

represents forest cover rather than its discontinuity and configuration, but has the advantage

of being calculable and homogeneous across Europe. Besides, the landscapes surrounding

our sampling units presented a high forest cover (77% on average, in a 500 ha buffer,

Table 1). Thus, the observed correlations reflect that forest area density could have been

beneficial for forest interior species that are sensitive to edge effects (bryophytes or lichens),

but detrimental to non-forest, open habitat or light demanding species, such as

tracheophytes and open-habitat birds (but see Bełcik et al. 2020, who show an increase

before a plateau in taxonomic diversity with increasing forest area in agricultural

landscapes). The inferences on the effect of fragmentation on biodiversity, based on the

Single Large vs Several Small principle (Diamond 1975), support that large continuous

habitat should have a greater effect on biodiversity conservation, compared to several small

ones. However, recent multitaxa studies proved the disproportionately high value of small

patches, harboring richer assemblages, even when focusing only on protected species (Riva

and Fahrig 2022). Besides, it has been highlighted that biodiversity conservation in human-

modified forest landscapes is better achieved by maintaining at least 40% of forest cover,

rather than focusing on fragmentation and configuration (Arroyo-Rodríguez et al. 2020).

Hence, the role of fragmentation, as it is currently estimated, should probably be

reconsidered to take into account both the amount of forest cover and the value of the

patches.

Contrary to the previous indicators, threatened species and birds are based on direct

biodiversity sampling (other than trees). These indicators directly describe the evolution of a

small part of the biodiversity, but generally correlate poorly to the richness of other taxa and

most of the results observed are linked to the fact that these indicators represent subsets of

larger groups (evidently forest birds vs. birds). However, despite several pieces of evidence

showing that congruence between taxa is generally small, especially in forests (Burrascano

et al. 2018; Westgate et al. 2017), forest birds positively correlated to beetles and fungi.

Probably these groups respond to the same favorable habitat conditions, but this was not

reflected in the multivariate analyses.

The proportion of protected forests around the plots positively influenced the richness of

bryophytes and lichens, highlighting the positive effects of protection for these groups, but

surprisingly had a negative effect on birds (see Honkanen et al. 2010). This may be related

to the consideration of only strictly protected areas in this indicator. Strictly protected areas

have no or little human interference, reducing possible heterogeneity at the landscape scale.

Finally, invasive non-native tree species did not show any correlation with biodiversity, apart

from a marginal positive effect on tracheophytes supporting high local tracheophyte diversity

to be often an indicator of disturbance. Despite the negative effects of invasive non-native

species introduction on native flora, fauna and mycota, this effect is rather limited in our

dataset since it contains very few invasive species in total (maximum one tree species per

plot) reflecting that no projects targeted the effects of plantations rich in, or dominated by,

invasive non-native tree species on biodiversity. This limited gradient probably does not

allow to show significant effects on biodiversity. In addition, an indicator on introduced

species rather than invasive ones – as it is defined in Forest Europe's indicators – may

perform better.

Limitations of Forest Europe indicators to predict biodiversity patterns and implications for

improvements

Our study is one of the first to test Forest Europe indicators against multitaxonomic data at

the European level. During the model selection process, the best single variable models

were more often involving naturalness (tracheophytes, bryophytes and lichens) followed by

forest birds (birds and beetles) and deadwood (fungi). However, the multivariate models did

perform better than single ones (except for beetles), and often the best ones combined

several indicators to reflect variations of the *indicandum*. Only beetles were best indicated by

forest birds only, but with a low magnitude, and birds by three combined indicators (including

forest birds and threatened species). The magnitude of the effects in the multivariate models

were comparable to those of single variable models (Appendix 4) which confirmed that

indicators were not collinear.

We studied only total scaled richness as a response variable, and evidently, the results may

be different for other, more specialized groups, or other metrics of biodiversity (abundance,

occurrence of individual species, functional diversity). Such an approach remains to be

tested (see e.g. Lelli et al. 2019) but was beyond the scope of the present study. In addition,

some of the references we used were probably incomplete regarding some groups: e.g.

almost no lichens are included in the list we used for red-listed species, but the proportion of

threatened species at the national levels may be high.

We limited our approach to the strict definition of the indicators as used in Forest Europe, but

higher performance could probably be reached by at least two improvements. First, adding

context to the indicators could probably reveal that they need to be adapted locally

(Chiarucci et al. 2012; Honkanen et al. 2010). Examples of context could be: elevation

(mountain vs. lowlands), biome (Mediterranean, temperate, boreal) and European forest

types. Second, most of the metrics we used are abundance metrics (apart from tree,

threatened and bird species indicators) that quantify habitat available for species. Indicators

based on diversity of resources (following the heterogeneity-diversity theory, Tews et al.

2004) could perform better, e.g. in the case of deadwood and saproxylic beetles (Bouget et

al. 2013), or tree-related microhabitats and birds and bats (Paillet et al. 2018). It would then

be interesting to assess the performance of other indicator metrics vs. the current ones in

assessing forest biodiversity.

Conclusions: towards new indicators definitions and better reporting

Even if novel agreements, such as the Kunming-Montreal Global Biodiversity Framework

(Convention on Biological Diversity 2022), continue stressing the importance of sustainable

management in forestry, there is the risk that environmental policies could be biased towards

market-related values (Pascual et al. 2023). Thus, the availability and application of reliable

biodiversity indicators is both urgent and imperative.

Many forest biodiversity indicators are proxies based on pre-existing data mostly issued from

National Forest Inventories (Tomppo et al. 2010) and more generally describe habitat

attributes or area-based metrics rather than biodiversity *per se* (Marshall et al. 2020).

Despite recent progresses based on international initiatives (namely the Essential

Biodiversity Variables, GEOBON, IPBES), monitoring the state and trends of forest

biodiversity solely based on proxies is not satisfactory: while proxies are generally easier to

measure than species themselves, they are prone to demographic effects such as extinction

debts or colonization credits. In other words, the presence of a given habitat - such as

deadwood - does not guarantee the presence of the species that depend on it (e.g. Paillet et

al. 2018). This results in a time-lag between the effects of management or policy on actual

biodiversity that may impair adaptive capacity in case of negative effects. In addition, the

response of biodiversity to a given indicator depends on the taxonomic or functional group

studied (Zeller et al. 2022), so no indicator may represent biodiversity overall. This was

confirmed by our analyses. We showed that, while some indicators performed correctly for

several taxonomic and functional groups in terms of significance and magnitude, we did not

highlight a predominant role of one indicator over the others.

Stevenson et al. (2021) claimed that indicators were often implemented without clear

considerations of their purposes and utility in terms of decision-making. We argue that, while

combinations of current Forest Europe indicators are useful to delineate general trends in

biodiversity, taking into account context and analyzing the performance of other - more

diversity-driven - metrics would help better reporting on biodiversity (e.g. Alterio et al. 2023;

Paillet et al. 2018). Such improvements would also be beneficial to the use of indicators

beyond general trends to include evaluating management and policy actions, decisions, or

set biodiversity targets (Stevenson et al. 2021), also beyond forest ecosystems. For

example, we showed that larger negative effects on biodiversity were observed in planted

forests. This poses key challenges for making the 3 billion trees planting promoted by the

European Forest Strategy for 2030 a beneficial action for biodiversity in forests (Sills et al.

2020), if high growing rate plantations such as introduced Douglas fir (*Pseudotsuga*

menziesii) or eucalyptus (*Eucalyptus* spp.) are promoted against semi-natural forests.

Conversely, promoting old-growth and unmanaged forests, as well as restoration of

monocultures and conversions towards semi-natural forests, could have a positive effect on

biodiversity. It is crucial to assess and balance these potential effects with the use of current

data available and biodiversity indicators before taking actions or to modulate actions in

favor of biodiversity against detrimental ones. Even if proxies are useful and relatively easy

to collect, they are probably not reactive enough to assess the state and dynamics of

biodiversity. We thus recommend assessing both habitat features as well as species to

clearly evaluate the effects of various pressures on biodiversity and being able to cope with

them in an adaptive way.

References

Alterio, E., Campagnaro, T., Sallustio, L., Burrascano, S., Casella, L., Sitzia, T., 2023. Forest management plans as data source for the assessment of the conservation status of European Union habitat types. Frontiers in Forests and Global Change 5.

Arroyo-Rodríguez, V., Fahrig, L., Tabarelli, M., Watling, J.I., Tischendorf, L., Benchimol, M., Cazetta, E., Faria, D., Leal, I.R., Melo, F.P.L., Morante-Filho, J.C., Santos, B.A., Arasa-Gisbert, R., Arce-Peña, N., Cervantes-López, M.J., Cudney-Valenzuela, S., Galán-Acedo, C., San-José, M., Vieira, I.C.G., Slik, J.W.F., Nowakowski, A.J., Tscharntke, T., 2020. Designing optimal human-modified landscapes for forest biodiversity conservation. Ecology Letters 23, 1404-1420.

Barton, K., 2023. MuMIn: multi-model inference. in R package version 1.47.5.

Bełcik, M., Lenda, M., Amano, T., Skórka, P., 2020. Different response of the taxonomic, phylogenetic and functional diversity of birds to forest fragmentation. Scientific Reports 10, 20320.

Betts, M.G., Wolf, C., Ripple, W.J., Phalan, B., Millers, K.A., Duarte, A., Butchart, S.H.M., Levi, T., 2017. Global forest loss disproportionately erodes biodiversity in intact landscapes. Nature 547, 441-444.

Boch, S., Prati, D., Hessenmöller, D., Schulze, E.D., Fischer, M., 2013. Richness of Lichen Species, Especially of Threatened Ones, Is Promoted by Management Methods Furthering Stand Continuity. PLoS ONE 8.

Boch, S., Saiz, H., Allan, E., Schall, P., Prati, D., Schulze, E.-D., Hessenmöller, D., Sparrius, L.B., Fischer, M., 2021. Direct and Indirect Effects of Management Intensity and Environmental Factors on the Functional Diversity of Lichens in Central European Forests, In Microorganisms.

Bouget, C., Larrieu, L., Nusillard, B., Parmain, G., 2013. In search of the best local habitat drivers for saproxylic beetle diversity in temperate deciduous forests. Biodiversity and Conservation 22, 2111-2130.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., Skaug, H.J., Maechler, M., Bolker, B.M., 2017. glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. The R Journal, 9, 378-400.

Burnham, K.P., Anderson, D.R., 2002. Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach (2nd ed).

Burrascano, S., Chianucci, F., Trentanovi, G., Kepfer-Rojas, S., Sitzia, T., Tinya, F., Doerfler, I., Paillet, Y., Nagel, T.A., Mitic, B., Morillas, L., Munzi, S., Van der Sluis, T., Alterio, E., Balducci, L., de Andrade, R.B., Bouget, C., Giordani, P., Lachat, T.,

Matosevic, D., Napoleone, F., Nascimbene, J., Paniccia, C., Roth, N., Aszalós, R., Brazaitis, G., Cutini, A., D'Andrea, E., De Smedt, P., Heilmann-Clausen, J., Janssen, P., Kozák, D., Mårell, A., Mikoláš, M., Nordén, B., Matula, R., Schall, P., Svoboda, M., Ujhazyova, M., Vandekerkhove, K., Wohlwend, M., Xystrakis, F., Aleffi, M., Ammer, C., Archaux, F., Asbeck, T., Avtzis, D., Ayasse, M., Bagella, S., Balestrieri, R., Barbati, A., Basile, M., Bergamini, A., Bertini, G., Biscaccianti, A.B., Boch, S., Bölöni, J., Bombi, P., Boscardin, Y., Brunialti, G., Bruun, H.H., Buscot, F., Byriel, D.B., Campagnaro, T., Campanaro, A., Chauvat, M., Ciach, M., Čiliak, M., Cistrone, L., Pereira, J.M.C., Daniel, R., De Cinti, B., De Filippo, G., Dekoninck, W., Di Salvatore, U., Dumas, Y., Elek, Z., Ferretti, F., Fotakis, D., Frank, T., Frey, J., Giancola, C., Gomoryová, E., Gosselin, M., Gosselin, F., Gossner, M.M., Götmark, F., Haeler, E., Hansen, A.K., Hertzog, L., Hofmeister, J., Hošek, J., Johannsen, V.K., Justensen, M.J., Korboulewsky, N., Kovács, B., Lakatos, F., Landivar, C.M., Lens, L., Lingua, E., Lombardi, F., Máliš, F., Marchino, L., Marozas, V., Matteucci, G., Mattioli, W., Møller, P.F., Müller, J., Németh, C., Ónodi, G., Parisi, F., Perot, T., Perret, S., Persiani, A.M., Portaccio, A., Posillico, M., Preikša, Ž., Rahbek, C., Rappa, N.J., Ravera, S., Romano, A., Samu, F., Scheidegger, C., Schmidt, I.K., Schwegmann, S., Sicuriello, F., Spinu, A.P., Spyroglou, G., Stillhard, J., Topalidou, E., Tøttrup, A.P., Ujházy, K., Veres, K., Verheyen, K., Weisser, W.W., Zapponi, L., Ódor, P., 2023. Where are we now with European forest multi-taxon biodiversity and where can we head to? Biological Conservation 284.

Burrascano, S., De Andrade, R.B., Paillet, Y., Ódor, P., Antonini, G., Bouget, C., Campagnaro, T., Gosselin, F., Janssen, P., Persiani, A.M., Nascimbene, J., Sabatini, F.M., Sitzia, T., Blasi, C., 2018. Congruency across taxa and spatial scales: are we asking too much of species data? Global Ecology and Biogeography 27, 980-990.

Campagnaro, T., Brundu, G., Sitzia, T., 2018. Five major invasive alien tree species in European Union forest habitat types of the Alpine and Continental biogeographical regions. Journal for Nature Conservation 43, 227-238.

Campanaro, A., Redolfi De Zan, L., Hardersen, S., Antonini, G., Chiari, S., Cini, A., Mancini, E., Mosconi, F., Rossi de Gasperis, S., Solano, E., Bologna, M.A., Sabbatini Peverieri, G., 2017. Guidelines for the monitoring of Rosalia alpina. Nature Conservation 20, 165-203.

Cavard, X., Macdonald, S.E., Bergeron, Y., Chen, H.Y.H., 2011. Importance of mixedwoods for biodiversity conservation: Evidence for understory plants, songbirds, soil fauna, and ectomycorrhizae in northern forests. Environmental Reviews 19, 142-161.

Chaudhary, A., Burivalova, Z., Koh, L.P., Hellweg, S., 2016. Impact of Forest Management on Species Richness: Global Meta-Analysis and Economic Trade-Offs. Scientific Reports 6, 23954.

Chiarucci, A., Bacaro, G., Filibeck, G., Landi, S., Maccherini, S., Scoppola, A., 2012. Scale
dependence of plant species richness in a network of protected areas. Biodiversity and
Conservation 21, 503-516.
Chirici, G., McRoberts, R.E., Winter, S., Bertini, R., Bröändli, U.B., Asensio, I.A., Bastrup-
Birk, A., Rondeux, J., Barsoum, N., Marchetti, M., 2012. National forest inventory
contributions to forest biodiversity monitoring. Forest Science 58, 257-268.
Christensen, M., Hahn, K., Mountford, E.P., Odor, P., Standovar, T., Rozenbergar, D., Diaci,
J., Wijdeven, S., Meyer, P., Winter, S., Vrska, T., 2005. Dead wood in European beech
(Fagus sylvatica) forest reserves. Forest Ecology and Management 210, 267-282.
Convention on Biological Diversity, 2022. Decision CBD/COP/DEC/15/4, ed. United Nations
Environment Programme.
Diamond, J.M., 1975. The island dilemma: Lessons of modern biogeographic studies for the
design of natural reserves. Biological Conservation 7, 129-146.
Dubois, G.F., Vignon, V., Delettre, Y.R., Rantier, Y., Vernon, P., Burel, F., 2009. Factors
affecting the occurrence of the endangered saproxylic beetle Osmoderma eremita
(Scopoli, 1763) (Coleoptera: Cetoniidae) in an agricultural landscape. Landscape and
Urban Planning 91, 152-159.
European Commission, 2021. New EU Forest Strategy for 2030, In COM(2021) 572 final.
ed. t.C. Communication from the Commission to the European Parliament, the
European Economic and Social Commitee and the Commitee of the Regions, Brussels,
Belgium.
European Commission, Joint Research Centre, Vogt, P., Riitters, K., Caudullo, G., Eckhardt,
B., Raši, R., 2019. An approach for pan-European monitoring of forest fragmentation.
Publications Office.
European Environment, A., 2020. State of nature in the EU – Results from reporting under
the nature directives 2013-2018. Publications Office.
European Environment Agency, 2007. European forest types. Categories and types for
sustainable forest management reporting and policy, In EEA Technical report. EEA,
Copenhagen, Danemark.
FAO, 2024. The State of the World's Forests 2024 – Forest-sector innovations towards a
more sustainable future. FAO, Rome, Italy.
Fick, S.E., Hijmans, R.J., 2017. WorldClim 2: new 1-km spatial resolution climate surfaces
for global land areas. International Journal of Climatology 37, 4302-4315.
Forest Europe, 2015. The updated pan-european indicators for sustainable forest
management, pp. 1-4. Ministerial Conference on the Protection of Forests in Europe,,
Zvolen, Slovak Republic.

Forest Europe, 2020. State of Europe's Forests 2020, p. 394. Ministerial Conference on the
Protection of Forests in Europe - FOREST EUROPE, Liaison Unit Bratislava.

Gao, T., Nielsen, A.B., Hedblom, M., 2015. Reviewing the strength of evidence of
biodiversity indicators for forest ecosystems in Europe. Ecological Indicators 57, 420-
434.

Gosselin, M., Fourcin, D., Dumas, Y., Gosselin, F., Korboulewsky, N., Toïgo, M., Vallet, P.,
2017. Influence of forest tree species composition on bryophytic diversity in mixed and
pure pine (Pinus sylvestris L.) and oak (Quercus petraea (Matt.) Liebl.) stands. Forest
Ecology and Management 406, 318-329.

Graf, M., Seibold, S., Gossner, M.M., Hagge, J., Weiß, I., Bässler, C., Müller, J., 2022.
Coverage based diversity estimates of facultative saproxylic species highlight the
importance of deadwood for biodiversity. Forest Ecology and Management 517, 120275.

Group on Earth Observation Biodiversity Observation Network, 2008. GEO BON Concept
Document, In GEO-V document 20. Geneva, Switzerland.

Heym, M., Uhl, E., Moshammer, R., Dieler, J., Stimm, K., Pretzsch, H., 2021. Utilising forest
inventory data for biodiversity assessment. Ecological Indicators 121, 107196.

Honkanen, M., Roberge, J.-M., Rajasärkkä, A., Mönkkönen, M., 2010. Disentangling the
effects of area, energy and habitat heterogeneity on boreal forest bird species richness
in protected areas. Global Ecology and Biogeography 19, 61-71.

Hsieh, T.C., Ma, K.H., Chao, A., 2016. iNEXT: an R package for rarefaction and
extrapolation of species diversity (Hill numbers). Methods in Ecology and Evolution 7,
1451-1456.

IPBES, 2019. Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services
(Version 1), eds E.D. Brondizio, S., J.N. Settele, H.T., p. 1144 IPBES secretariat, Bonn,
Germany.

Jiguet, F., Devictor, V., Julliard, R., Couvet, D., 2012. French citizens monitoring ordinary
birds provide tools for conservation and ecological sciences. Acta Oecologica 44, 58-66.

Kempeneers, P., Sedano, F., Seebach, L., Strobl, P., San-Miguel-Ayanz, J., 2011. Data
Fusion of Different Spatial Resolution Remote Sensing Images Applied to Forest-Type
Mapping. IEEE Transactions on Geoscience and Remote Sensing 49, 4977-4986.

Koch Widerberg, M., Ranius, T., Drobyshev, I., Nilsson, U., Lindbladh, M., 2012. Increased
openness around retained oaks increases species richness of saproxylic beetles.
Biodiversity and Conservation 21, 3035-3059.

Larrieu, L., Paillet, Y., Winter, S., Bütler, R., Kraus, D., Krumm, F., Lachat, T., Michel, A.K.,
Regnery, B., Vandekerkhove, K., 2018. Tree related microhabitats in temperate and

Mediterranean European forests: A hierarchical typology for inventory standardization. Ecological Indicators 84, 194-207.

Lassauce, A., Paillet, Y., Jactel, H., Bouget, C., 2011. Deadwood as a surrogate for forest biodiversity: Meta-analysis of correlations between deadwood volume and species richness of saproxylic organisms. Ecological Indicators 11, 1027-1039.

Leidinger, J., Blaschke, M., Ehrhardt, M., Fischer, A., Gossner, M.M., Jung, K., Kienlein, S., Kózak, J., Michler, B., Mosandl, R., Seibold, S., Wehner, K., Weisser, W.W., 2021. Shifting tree species composition affects biodiversity of multiple taxa in Central European forests. Forest Ecology and Management 498, 119552.

Lelli, C., Bruun, H.H., Chiarucci, A., Donati, D., Frascaroli, F., Fritz, Ö., Goldberg, I., Nascimbene, J., Tøttrup, A.P., Rahbek, C., Heilmann-Clausen, J., 2019. Biodiversity response to forest structure and management: Comparing species richness, conservation relevant species and functional diversity as metrics in forest conservation. Forest Ecology and Management 432, 707-717.

Lindenmayer, D.B., Likens, G.E., 2010. The science and application of ecological monitoring. Biological Conservation 143, 1317-1328.

Linser, S., O'Hara, P., 2019. Guidelines for the Development of a Criteria and Indicator Set for Sustainable Forest Management, In Timber and Forest Discussion Paper. p. 91. United Nations and the Food and Agriculture, Organization of the United Nations, Geneva, Switzerland.

Marshall, E., Wintle, B.A., Southwell, D., Kujala, H., 2020. What are we measuring? A review of metrics used to describe biodiversity in offsets exchanges. Biological Conservation 241, 108250.

Müller, J., Boch, S., Prati, D., Socher, S.A., Pommer, U., Hessenmöller, D., Schall, P., Schulze, E.D., Fischer, M., 2019. Effects of forest management on bryophyte species richness in Central European forests. Forest Ecology and Management 432, 850-859.

Müller, J., Brustel, H., Brin, A., Bussler, H., Bouget, C., Obermaier, E., Heidinger, I.M.M., Lachat, T., Förster, B., Horak, J., Procházka, J., Köhler, F., Larrieu, L., Bense, U., Isacsson, G., Zapponi, L., Gossner, M.M., 2015. Increasing temperature may compensate for lower amounts of dead wood in driving richness of saproxylic beetles. Ecography 38, 499-509.

Müller, J., Jarzabek-Müller, A., Bussler, H., Gossner, M.M., 2014. Hollow beech trees identified as keystone structures for saproxylic beetles by analyses of functional and phylogenetic diversity. Animal Conservation 17, 154-162.

Noss, R.F., 1990. Indicators for monitoring biodiversity - A hierarchical approach. Conservation Biology 4, 355-364.

Paillet, Y., Archaux, F., Boulanger, V., Debaive, N., Fuhr, M., Gilg, O., Gosselin, F., Guilbert,
E., 2017. Snags and large trees drive higher tree microhabitat densities in strict forest
reserves. Forest Ecology and Management 389, 176-186.

Paillet, Y., Archaux, F., du Puy, S., Boulanger, V., Debaive, N., Fuhr, M., Gilg, O., Gosselin,
F., Guilbert, E., 2018. The indicator side of tree microhabitats: a multi-taxon approach
based on bats, birds and saproxylic beetles. Journal of Applied Ecology 55, 2147-2156.

Paillet, Y., Bergès, L., Hjälten, J., Odor, P., Avon, C., Bernhardt-Romermann, M., Bijlsma,
R.J., De Bruyn, L., Fuhr, M., Grandin, U., Kanka, R., Lundin, L., Luque, S., Magura, T.,
Matesanz, S., Meszaros, I., Sebastia, M.T., Schmidt, W., Standovar, T., Tothmeresz, B.,
Uotila, A., Valladares, F., Vellak, K., Virtanen, R., 2010. Biodiversity Differences
between Managed and Unmanaged Forests: Meta-Analysis of Species Richness in
Europe. Conservation Biology 24, 101-112.

Parajuli, R., Markwith, S.H., 2023. Quantity is foremost but quality matters: A global meta-
analysis of correlations of dead wood volume and biodiversity in forest ecosystems.
Biological Conservation 283.

Pascual, U., Balvanera, P., Anderson, C.B., Chaplin-Kramer, R., Christie, M., González-
Jiménez, D., Martin, A., Raymond, C.M., Termansen, M., Vatn, A., Athayde, S.,
Baptiste, B., Barton, D.N., Jacobs, S., Kelemen, E., Kumar, R., Lazos, E., Mwampamba,
T.H., Nakangu, B., O'Farrell, P., Subramanian, S.M., van Noordwijk, M., Ahn, S.E.,
Amaruzaman, S., Amin, A.M., Arias-Arévalo, P., Arroyo-Robles, G., Cantú-Fernández,
M., Castro, A.J., Contreras, V., De Vos, A., Dendoncker, N., Engel, S., Eser, U., Faith,
D.P., Filyushkina, A., Ghazi, H., Gómez-Baggethun, E., Gould, R.K., Guibrunet, L.,
Gundimeda, H., Hahn, T., Harmáčková, Z.V., Hernández-Blanco, M., Horcea-Milcu, A.I.,
Huambachano, M., Wicher, N.L.H., Aydın, C.İ., Islar, M., Koessler, A.K., Kenter, J.O.,
Kosmus, M., Lee, H., Leimona, B., Lele, S., Lenzi, D., Lliso, B., Mannetti, L.M., Merçon,
J., Monroy-Sais, A.S., Mukherjee, N., Muraca, B., Muradian, R., Murali, R., Nelson,
S.H., Nemogá-Soto, G.R., Ngouhouo-Poufoun, J., Niamir, A., Nuesiri, E., Nyumba, T.O.,
Özkaynak, B., Palomo, I., Pandit, R., Pawłowska-Mainville, A., Porter-Bolland, L.,
Quaas, M., Rode, J., Rozzi, R., Sachdeva, S., Samakov, A., Schaafsma, M., Sitas, N.,
Ungar, P., Yiu, E., Yoshida, Y., Zent, E., 2023. Diverse values of nature for
sustainability. Nature 620, 813-823.

Penone, C., Allan, E., Soliveres, S., Felipe-Lucia, M.R., Gossner, M.M., Seibold, S., Simons,
N.K., Schall, P., van der Plas, F., Manning, P., Manzanedo, R.D., Boch, S., Prati, D.,
Ammer, C., Bauhus, J., Buscot, F., Ehbrecht, M., Goldmann, K., Jung, K., Müller, J.,
Müller, J.C., Pena, R., Polle, A., Renner, S.C., Ruess, L., Schönig, I., Schrumpf, M.,
Solly, E.F., Tschapka, M., Weisser, W.W., Wubet, T., Fischer, M., 2019. Specialisation

and diversity of multiple trophic groups are promoted by different forest features. Ecology Letters 22, 170-180.

Pereira, H.M., David Cooper, H., 2006. Towards the global monitoring of biodiversity change. Trends in Ecology & Evolution 21, 123-129.

Proença, V., Martin, L.J., Pereira, H.M., Fernandez, M., McRae, L., Belnap, J., Böhm, M., Brummitt, N., García-Moreno, J., Gregory, R.D., Honrado, J.P., Jürgens, N., Opige, M., Schmeller, D.S., Tiago, P., van Swaay, C.A.M., 2017. Global biodiversity monitoring: From data sources to Essential Biodiversity Variables. Biological Conservation 213, 256-263.

R Core Team, 2023. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Reise, J., Kukulka, F., Flade, M., Winter, S., 2019. Characterising the richness and diversity of forest bird species using National Forest Inventory data in Germany. Forest Ecology and Management 432, 799-811.

Rigal, S., Dakos, V., Alonso, H., Auniņš, A., Benkő, Z., Brotons, L., Chodkiewicz, T., Chylarecki, P., de Carli, E., del Moral, J.C., Domşa, C., Escandell, V., Fontaine, B., Foppen, R., Gregory, R., Harris, S., Herrando, S., Husby, M., Ieronymidou, C., Jiguet, F., Kennedy, J., Klvaňová, A., Kmecl, P., Kuczyński, L., Kurlavičius, P., Kålås, J.A., Lehikoinen, A., Lindström, Å., Lorrillière, R., Moshøj, C., Nellis, R., Noble, D., Eskildsen, D.P., Paquet, J.-Y., Pélissié, M., Pladevall, C., Portolou, D., Reif, J., Schmid, H., Seaman, B., Szabo, Z.D., Szép, T., Florenzano, G.T., Teufelbauer, N., Trautmann, S., van Turnhout, C., Vermouzek, Z., Vikstrøm, T., Voříšek, P., Weiserbs, A., Devictor, V., 2023. Farmland practices are driving bird population decline across Europe. Proceedings of the National Academy of Sciences 120, e2216573120.

Riva, F., Fahrig, L., 2022. The disproportionately high value of small patches for biodiversity conservation. Conservation Letters 15, e12881.

Sills, J., Gómez-González, S., Ochoa-Hueso, R., Pausas, J.G., 2020. Afforestation falls short as a biodiversity strategy. Science 368, 1439-1439.

Simons, N.K., Felipe-Lucia, M.R., Schall, P., Ammer, C., Bauhus, J., Blüthgen, N., Boch, S., Buscot, F., Fischer, M., Goldmann, K., Gossner, M.M., Hänsel, F., Jung, K., Manning, P., Nauss, T., Oelmann, Y., Pena, R., Polle, A., Renner, S.C., Schloter, M., Schöning, I., Schulze, E.-D., Solly, E.F., Sorkau, E., Stempfhuber, B., Wubet, T., Müller, J., Seibold, S., Weisser, W.W., 2021. National Forest Inventories capture the multifunctionality of managed forests in Germany. Forest Ecosystems 8, 5.

Stevenson, S.L., Watermeyer, K., Caggiano, G., Fulton, E.A., Ferrier, S., Nicholson, E., 2021. Matching biodiversity indicators to policy needs. Conservation Biology 35, 522-532.

Storch, F., Boch, S., Gossner, M.M., Feldhaar, H., Ammer, C., Schall, P., Polle, A., Kroiher,
F., Müller, J., Bauhus, J., 2023. Linking structure and species richness to support forest
biodiversity monitoring at large scales. Annals of Forest Science 80, 3.

Tews, J., Brose, U., Grimm, V., Tielbörger, K., Wichmann, M.C., Schwager, M., Jeltsch, F.,
2004. Animal species diversity driven by habitat heterogeneity/diversity: The importance
of keystone structures. Journal of Biogeography 31, 79-92.

Tomppo, E., Gschwantner, T., Lawrence, M., Mc Roberts, R.E., 2010. National forest
inventories. Pathways for common reporting. Springer Science, Heidelberg, Allemagne.

Trentanovi, G., Campagnaro, T., Sitzia, T., Chianucci, F., Vacchiano, G., Ammer, C., Ciach,
M., Nagel, T.A., del Río, M., Paillet, Y., Munzi, S., Vandekerkhove, K., Bravo-Oviedo, A.,
Cutini, A., D'Andrea, E., De Smedt, P., Doerfler, I., Fotakis, D., Heilmann-Clausen, J.,
Hofmeister, J., Hošek, J., Janssen, P., Kepfer-Rojas, S., Korboulewsky, N., Kovács, B.,
Kozák, D., Lachat, T., Mårell, A., Matula, R., Mikoláš, M., Nordén, B., Ódor, P., Perović,
M., Pötzelsberger, E., Schall, P., Svoboda, M., Tinya, F., Ujházyová, M., Burrascano, S.,
2023. Words apart: Standardizing forestry terms and definitions across European
biodiversity studies. Forest Ecosystems 10, 100128.

Vítková, L., Bače, R., Kjučukov, P., Svoboda, M., 2018. Deadwood management in Central
European forests: Key considerations for practical implementation. Forest Ecology and
Management 429, 394-405.

Weber, D., Hintermann, U., Zangger, A., 2004. Scale and trends in species richness:
considerations for monitoring biological diversity for political purposes. Global Ecology
and Biogeography 13, 97-104.

Westgate, M.J., Tulloch, A.I.T., Barton, P.S., Pierson, J.C., Lindenmayer, D.B., 2017.
Optimal taxonomic groups for biodiversity assessment: a meta-analytic approach.
Ecography 40, 539-548.

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis, 2 edn. Springer Cham.

Wood, S., 2023. Mixed GAM Computation Vehicle with Automatic Smoothness Estimation.
V1.9.

Zeller, L., Baumann, C., Gonin, P., Heidrich, L., Keye, C., Konrad, F., Larrieu, L., Meyer, P.,
Sennhenn-Reulen, H., Müller, J., Schall, P., Ammer, C., 2022. Index of biodiversity
potential (IBP) versus direct species monitoring in temperate forests. Ecological
Indicators 136, 108692.

Zeller, L., Förster, A., Keye, C., Meyer, P., Roschak, C., Ammer, C., 2023. What does
literature tell us about the relationship between forest structural attributes and species
richness in temperate forests? – A review. Ecological Indicators 153, 110383.

Zuur, A.F., Ieno, E.N., Elphick, C.S., 2010. A protocol for data exploration to avoid common
statistical problems. Methods in Ecology and Evolution 1, 3-14.

Appendices

Appendix 1: Distribution of plots used for the analyses according to forest type (European

Environment Agency 2007, Table 1.1), mean annual temperature (Figure 1.1) and mean

annual precipitation (Figure 1.2) derived from Worldclim 2.0 (Fick and Hijmans 2017).

Table 1.1: Distribution of the study plots in the 14 forest categories of the (European

Environment Agency 2007)

Forest category	Name	Tracheo phytes	Bryophyt es	Birds	Beetles	Fungi	Lichens
1	Boreal forest	64	0	64	0	0	0
2	Hemiboreal forest and nemoral coniferous and mixed broadleaved coniferous forest	100	18	101	36	36	35
3	Alpine coniferous forest	25	18	25	21	16	0
4	Acidophilous oak and oak-birch forest	60	56	60	57	59	0
5	Mesophytic deciduous forest	83	72	83	68	69	47
6	Beech forest	194	103	180	171	191	128
7	Mountainous beech forest	253	105	254	107	176	166
8	Thermophilous deciduous forest	9	9	9	0	7	9
9	Broadleaved evergreen forest	1	0	0	0	1	0
14	Floodplain forest	49	41	49	49	49	49

Figure 1.1: Mean annual temperatures for the different taxa datasets derived from Worldclim

2.1 (1991-2010).

Figure 1.2: Mean annual precipitation for the different taxa datasets derived from Worldclim

2.1 (1991-2010)

Appendix 2: Comparison of linear (GLMMs) vs non-linear (GAMMs) models analyses

between standardized species richness and several Forest Europe's biodiversity indicators.

964   Appendix 3: Correlation between indicators of Forest Europe from the different taxa

965   datasest. Figures on the right are Pearson correlation coefficients, biplots on the left and

966   histograms on the diagonal represent the distribution of the data. Regeneration (4.2) and

967   Naturalness (4.3) are categorical variables not represented here.

968   3.1. Tracheophytes

 3.2. Bryophytes

    3.3. Beetles

3.4. Birds

 3.5. Lichens

 Appendix 4: Standardized richness estimates table for all single generalized mixed models

982 with beta error distribution and logit link. Intercept for I4.2.Regeneration is "Coppice", and for

983 I4.3.Naturalness "Plantation" (PLA). S.N = semi-natural forest, UNM = Unmanaged forests.

984 se = standard error of the mean, pl = critical probability, (*) p<0.1, * p<0.05, ** p<0.01, ***

985 p<0.001.

Group	variable	estimate	se	p	
Tracheophytes	I4.1.tree.sp	0.0648	0.0359	0.0712	(*)
	I4.2.regeneration(natural regeneration)	0.1431	0.2606	0.5828	
	I4.2.regeneration(planting)	0.5186	0.2754	0.0597	(*)
	I4.3.naturalnessUNM	-0.3878	0.0889	<0.001	***
	I4.3.naturalnessS.N	-0.2073	0.1429	0.147	
	I4.4.invasive	0.0376	0.0193	0.0508	(*)
	I4.5.deadwood	0.0318	0.0219	0.1463	
	I4.7.fragmentation	-0.0898	0.0256	<0.001	***
	I4.8.threat.sp	0.0428	0.0221	0.0522	(*)
	I4.9.protected.forests	-0.0087	0.0416	0.8344	
	I4.10.birds	0.0122	0.0326	0.707	
Bryophytes	I4.1.tree.sp	0.2828	0.0633	<0.001	***
	I4.2.regeneration(natural regeneration)	0.4156	0.4546	0.3606	
	I4.2.regeneration(planting)	-0.5511	0.4716	0.2426	
	I4.3.naturalnessUNM	0.8606	0.128	<0.001	***
	I4.3.naturalnessS.N	1.5727	0.1745	<0.001	***
	I4.4.invasive	-0.0409	0.0388	0.2917	
	I4.5.deadwood	0.2097	0.04	<0.001	***
	I4.7.fragmentation	0.1384	0.0486	0.0044	**
	I4.8.threat.sp	0.0776	0.0402	0.0539	(*)
	I4.9.protected.forests	0.2268	0.0585	<0.001	***
	I4.10.birds	0.0235	0.0571	0.6806	
Birds	I4.1.tree.sp	0.0341	0.0323	0.2916	
	I4.2.regeneration(natural regeneration)	0.0996	0.2114	0.6375	
	I4.2.regeneration(planting)	0.0452	0.2249	0.8407	

Group	variable	estimate	se	p	
	I4.3.naturalnessUNM	0.0388	0.0785	0.6207	
	I4.3.naturalnessS.N	0.2011	0.1259	0.1104	
	I4.4.invasive	-0.0177	0.0186	0.3417	
	I4.5.deadwood	0.0402	0.0203	0.0469	*
	I4.7.fragmentation	-0.1018	0.0243	<0.001	***
	I4.8.threat.sp	0.043	0.0205	0.0365	*
	I4.9.protected.forests	-0.1031	0.041	0.012	*
	I4.10.birds	0.6521	0.0218	<0.001	***
Beetles	I4.1.tree.sp	0.0242	0.034	0.4762	
	I4.2.regeneration(natural regeneration)	0.1411	0.2247	0.53	
	I4.2.regeneration(planting)	0.1782	0.233	0.4444	
	I4.3.naturalnessUNM	-0.0338	0.0638	0.5965	
	I4.3.naturalnessS.N	-0.0749	0.109	0.4917	
	I4.4.invasive	-0.0135	0.0175	0.441	
	I4.5.deadwood	0.0045	0.0204	0.8238	
	I4.7.fragmentation	0.0435	0.0263	0.0983	(*)
	I4.8.threat.sp	0.0096	0.0196	0.6255	
	I4.9.protected.forests	-0.0526	0.0357	0.1409	
	I4.10.birds	0.05	0.023	0.0297	*
Fungi	I4.1.tree.sp	0.1476	0.0343	<0.001	***
	I4.2.regeneration(natural regeneration)	0.1607	0.2444	0.5109	
	I4.2.regeneration(planting)	-0.3632	0.2542	0.1531	
	I4.3.naturalnessUNM	0.5017	0.0691	<0.001	***
	I4.3.naturalnessS.N	0.9083	0.1434	<0.001	***
	I4.4.invasive	0.0035	0.0189	0.8526	
	I4.5.deadwood	0.2228	0.0181	<0.001	***
	I4.7.fragmentation	0.054	0.0292	0.0644	(*)
	I4.8.threat.sp	0.0226	0.022	0.3039	
	I4.9.protected.forests	0.0422	0.0466	0.366	
	I4.10.birds	0.1021	0.0324	0.0016	**
Lichens	I4.1.tree.sp	0.1714	0.0507	<0.001	***

Group	variable	estimate	se	p	
	I4.2.regeneration(natural regeneration)	0.0169	0.258	0.9479	
	I4.2.regeneration(planting)	-0.5447	0.2736	0.0465	*
	I4.3.naturalnessUNM	0.5513	0.1028	<0.001	***
	I4.3.naturalnessS.N	0.7687	0.2439	0.0016	**
	I4.4.invasive	0.0092	0.0313	0.7691	
	I4.5.deadwood	0.1139	0.0343	<0.001	***
	I4.7.fragmentation	0.0137	0.0442	0.7564	
	I4.8.threat.sp	0.0339	0.0348	0.329	
	I4.9.protected.forests	0.1778	0.061	0.0036	**
	I4.10.birds	0.0067	0.0559	0.904	