One to rule them all? Assessing the performance of Forest Europe's biodiversity indicators against multitaxonomic data

Impact statement

Biodiversity indicators used to assess the state of Europe's forests perform unequally; a combination of indicators is more informative

Keywords

Sustainable forest management; taxonomic indicators; multi-taxa diversity; forest structure

Authors

Yoan Paillet, Univ. Grenoble Alpes, INRAE, Lessem, 2 rue de la Papeterie, BP76, 38402 Saint Martin d'Hères, cedex, France, yoan.paillet@inrae.fr, 0000-0001-7232-7844

Livia Zapponi, National Research Council, Institute of BioEconomy, Via Francesco Biasi 75, 38098 S. Michele all'Adige (TN) Italy, livia.zapponi@cnr.it, 0000-0003-4033-9001

Peter Schall, University of Göttingen, Silviculture and Forest Ecology of the Temperate Zones Büsgenweg 1, 37077 Göttingen, Germany, peter.schall@forst.uni-goettingen.de, 0000-0003-4808-818X

Jean-Matthieu Monnet, Univ. Grenoble Alpes, INRAE, Lessem, 2 rue de la Papeterie, BP76, 38402 Saint Martin d'Hères, cedex, France, jean-matthieu.monnet@inrae.fr, 0000-0002-9948-9891

Christian Ammer, University of Göttingen, Silviculture and Forest Ecology of the Temperate Zones Büsgenweg 1, 37077 Göttingen, Germany, christian.ammer@forst.uni-goettingen.de, 0000-0002-4235-0135

Lorenzo Balducci, Sapienza University of Rome, Piazzale Aldo Moro, 5, 00185 Rome, Italy, lorenzo.balducci@uniroma1.it, 0000-0001-9111-5679

Steffen Boch, WSL Swiss Federal Research Institute, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland, steffen.boch@wsl.ch, 0000-0003-2814-5343

36 **Gediminas Brazaitis**, Vytautas Magnus University Agriculture Academy Studentu g. 11,
37 53361 Akademija, Lithuania, gediminas.brazaitis@vdu.lt, 0000-0003-0234-9292
38
39 **Alessandro Campanaro**, Council for Agricultural Research and Economics, Research
40 Centre for Plant Protection and Certification Via Lanciola 12/a, Florence, Italy,
41 alessandro.campanaro@crea.gov.it , 0000-0003-4043-6362
42
43 **Francesco Chianucci**, CREA - Research Centre for Forestry and Wood Viale Santa
44 Margherita 80, Arezzo, Italy, fchianucci@gmail.com, 0000-0002-5688-2060
45
46 **Inken Doerfler**, Carl von Ossietzky Universität Oldenburg, Department Plant sociology and
47 nature Conservation Ammerländer Heerstraße 114-118 26129 Oldenbur, Germany,
48 inkendoerfler@gmail.com, 0000-0001-9624-1922
49
50 **Markus Fischer**, University of Bern, Institute of Plant Sciences Altenbergrain 21, 3013 Bern,
51 Switzerland, markus.fischer@ips.unibe.ch, 0000-0002-5589-5900
52
53 **Marion Gosselin**, INRAE, UR EFNO Domaine des Barres, 45290 Nogent-sur-Vernisson,
54 France, marion.gosselin@inrae.fr, 0000-0003-0161-5368
55
56 **Martin M. Gossner**, WSL Swiss Federal Research Institute, Zürcherstrasse 111, 8903
57 Birmensdorf, Switzerland, martin.gossner@wsl.ch, 0000-0003-1516-6364
58
59 **Jacob Heilmann-Clausen**, Center for Macroecolgy, Evolution and Climate, Globe Institute,
60 University of Copenhagen, Universitetsparken 15, 2100 København, Denmark, jheilmann-
61 clausen@sund.ku.dk, 0000-0003-4713-6004
62
63 **Jeňýk Hofmeister**, Czech University of Life Sciences Prague, Department of Forest
64 Ecology, Faculty of Forestry and Wood Sciences Kamýcká 129, 165 00 Praha - Suchdol,
65 Czech Republic, jenyk.hofmeister@email.cz, 0000-0002-3915-5056
66
67 **Jan Hošek**, Ecological Services Tichá 784/4, 268 01 Hořovice, Czech Republic,
68 hosek@ekologickesluzby.cz
69
70 **Sebastian Kepfer-Rojas**, University of Copenhagen, Department of Geosciences and
71 Natural Resource Management Rolighedsvej 23, 1958 Frederiksberg, Denmark,
72 skro@ign.ku.dk, 0000-0002-1681-2877
73
74 **Peter Odor**, Hun-Ren Centre for Ecological Research, Institute of Ecology and Botany,
75 Alkotmány u. 2-4., H-2163 Vácrátót; University of Sopron, Institute of Environmental
76 Protection and Nature Conservation, Bajcsy-Zsilinszky u. 4., H-9400 Sopron, Hungary,
77 odor.peter@ecolres.hu, 0000-0003-1729-8897
78
79 **Flóra Tinya**, HUN-REN Centre for Ecoogical Research, Institute of Ecology and Botany
80 Alkotmány u. 2-4., H-2163 Vácrátót, Hungary, tinya.flora@ecolres.hu, 0000-0002-4271-9676
81

82 **Giovanni Trentanovi**, Research Institute on Terrestrial Ecosystems - National Research
83 Council Via Madonna del Piano n. 10 50019 Sesto Fiorentino (Florence), Italy,
84 giovanni.trentanovi@cnr.it, 0000-0001-8236-4163
85
86 **Giorgio Vacchiano**, University of Milan, Department of Agricultural and Environmental
87 Science via Celoria 2, 20133 Milan, Italy, gvacchiano@gmail.com, 0000-0001-8100-0659
88
89 **Kris   Vandekerkhove**, Research Institute for Nature and Forest (INBO), Gaverstraat 4,
90 9500 Geraardsbergen, Belgium, kris.vandekerkhove@inbo.be, 0000-0003-1954-692X
91
92 **Wolfgang W. Weisser**, Technical University of Munich, Terrestrial Ecology Research
93 Group, Department of Life Science Systems, School of Life Sciences Hans-Carl-von-
94 Carlowitz-Platz 2, 85354 Freising, Germany, wolfgang.weisser@tum.de, 0000-0002-2757-
95 8959
96
97 **Sabina Burrascano**, Sapienza University of Rome, Piazzale Aldo Moro, 5, 00185 Rome,
98 Italy, sabina.burrascano@uniroma1.it, 0000-0002-6537-3313
99
100

Acknowledgements

This work was funded by the EU Framework Programme Horizon 2020 through the COST

Association (www.cost.eu): COST Action CA18207: BOTTOMS-UP – Biodiversity Of

Temperate forest Taxa Orienting Management Sustainability by Unifying Perspectives. The

authors are thankful to all those experts contributing to the data here harmonized and

resumed that were not listed as data contributors.

Abstract

Most broad-scale forest biodiversity indicators are based on data from national forest

inventories and are used to assess the state of biodiversity through several regional

initiatives and reporting. Although valuable, these indicators are essentially indirect and

evaluate habitat quantity and quality rather than biodiversity *per se*. Besides, most of these

indicators are applicable at regional or national scales, while their use at a more local level is

difficult. Therefore, their link to biodiversity may be weak, which decreases their usefulness

for decision-making.

For several decades, Forest Europe indicators assessed the state of European forests, in

particular its biodiversity. However, no extensive study has been conducted to date to

assess the performance of these indicators against multitaxonomic data. We hypothesized

that – as implied by the reporting process – no single biodiversity indicator from Forest

Europe can represent overall forest biodiversity, but that several – eventually combined –

indicators would reflect habitat quality for at least some taxa in a comprehensive way. We

tested the set of indicators proposed by Forest Europe against the species richness of six

taxonomic and functional groups (tracheophytes, epixylic and epiphytic bryophytes, birds,

saproxylic beetles, saproxylic non-lichenized fungi and epixylic and epiphytic lichenized

fungi) across several hundreds of plots over Europe. We showed that, while some indicators

perform relatively well across groups (e.g. deadwood volume), no single indicator

represented all biodiversity at once, and that a combination of several indicators performed

better. Surprisingly, some indicators showed weak links with the biodiversity of the six

taxonomic and functional groups.

Forest Europe indicators were chosen for their availability and ease of understanding for

most people. However, our analyses showed that there are still gaps in the monitoring

framework, and that surveying certain taxa along with stand structure is necessary to

support policymaking and tackle forest biodiversity loss at the large scale.

Keywords

Sustainable forest management; taxonomic indicators; multi-taxa diversity; forest structure

Introduction

Monitoring biodiversity in the face of global change is a challenge in many ecosystems across the world (IPBES 2019; Lindenmayer & Likens 2010). Despite large-scale initiatives such as GEOBON (Group on Earth Observation Biodiversity Observation Network 2008) and collaborative networks (Burrascano et al. 2023), no standard biodiversity monitoring scheme has been agreed in Europe or elsewhere. Long-term biodiversity monitoring hence currently relies on a limited number of initiatives that are used to assess the impact of climate change (e.g.Jiguet et al. 2012), habitat loss (Betts et al. 2017) or pollution (Rigal et al. 2023). Initiatives combining monitoring of biodiversity, types of pressure and ecosystem-level variables remain quite rare, or may concern only single pressure vs. single taxon assessment (e.g. Proença et al. 2017; Weber et al. 2004). However, different taxonomic and functional groups likely respond differently to a given pressure, which challenges prioritization of policy initiatives and tools (Pereira & David Cooper 2006).

Forest ecosystems are no exception to the rule, despite the large use of National Forest Inventory data to indirectly assess the state and evolution of biodiversity (Chirici et al. 2012; Heym et al. 2021; Reise et al. 2019). Originally designed to assess wood stock and production, National Forest Inventories provide nationwide data on forest ecosystems that allow, notably, to assess criteria and indicators of sustainable forest management and ecosystem services (Simons et al. 2021). At the European level, data from National Forest Inventories have been aggregated and published every five years for more than three decades (Forest Europe 2020) and provide a set of indicators to inform sustainable forest management.

In this process, the Sustainable Forest Management Criterion 4 is dedicated to the "*Maintenance, conservation and appropriate enhancement of biological diversity in forest ecosystems*". Composed of ten indicators, this criterion aims at helping decision makers to assess biodiversity-friendly sustainable forest management at the national and European scales. However, most of the indicators are indirect (structural) proxies of the forest ecosystem state (e.g. deadwood or forest fragmentation), whose link with biodiversity often

lacks strong scientific evidence (Gao et al. 2015; Paillet et al. 2018; Storch et al. 2023; Zeller

et al. 2022). Indeed, only two indicators involve other species than trees (namely 4.8

Threatened forest species, 4.10 Common forest bird species, Table 1) and to date, despite a

large corpus of individual studies and few syntheses (Zeller et al. 2023), no global and

systematic assessment of the correlations between multi-taxonomic biodiversity and Forest

Europe indicators has been attempted.

In this context, our aim was to provide an *ex post* validation of these indicators and propose

improvements to the reporting process. For this, we: i) assessed the link between indicators

and several - multi-taxonomic (i.e. taxonomic and functional groups) - *indicanda*; ii) identified

the indicators that performed best and universally - i.e. for all groups; iii) defined, if possible,

the most effective combination of indicators for forest biodiversity.

To address these aims, we first tested Forest Europe's biodiversity indicators against each

taxonomic and functional group. Second, we used a model selection approach to search for

the best indicators in terms of taxonomic and functional group. Third, we tested the effect of

a combination of indicators by searching for the best and most parsimonious combination of

biodiversity indicators. We used a multi-taxonomic database (Burrascano et al. 2023;

Trentanovi et al. 2023) combining species inventories and forest structure to analyze the

correlations between Forest Europe's biodiversity indicators with the biodiversity of six

groups (tracheophytes, epixylic and epiphytic bryophytes, birds, saproxylic beetles,

saproxylic non-lichenized fungi and epixylic and epiphytic lichenized fungi).

Materials and Methods

Database

We used the database gathered within the framework of the COST action "BOTTOMS-UP"

(CA18207 – Biodiversity Of Temperate forest Taxa Orienting Management Sustainability by

Unifying Perspectives). In a nutshell, this database comprises biodiversity samplings along

with forest plot descriptions (Burrascano et al. 2023). It merges 34 different datasets from 12

European countries and more than 3500 sampling units. From this database, we extracted

the six most represented taxonomic and functional groups, namely tracheophytes, epixylic

and epiphytic bryophytes (hereafter bryophytes), birds, saproxylic beetles, saproxylic non-

lichenized fungi (hereafter fungi) and epixylic and epiphytic lichenized fungi (hereafter

lichens).

Figure 1: Spatial distribution of the sampling sites. Gray areas are covered by forests with a

tree cover greater than 40% according to Kempeneers et al. (2011). The size of the dot is

proportional to the number of sampling units.

While tracheophytes and birds were inventoried without any specific selection of the guilds

targeted, only epiphytic and epixylic bryophytes and lichens were included (sampled on

living trees and deadwood), and saproxylic fungi and beetles (dependent on deadwood

substrates or on other organisms inhabiting deadwood).

Since the database is the result of the merging of different research projects, sampling

designs and protocols varied across datasets (Burrascano et al. 2023; Trentanovi et al.

2023). Therefore, we standardized species richness (number of species per sampling plot)

by dividing it by the asymptotic gamma richness at the site level, with site representing a

homogeneous sampling area with a maximum size of a few thousand hectares. We derived

site gamma diversity through plot-based rarefaction curves using the R package iNEXT

(Hsieh et al. 2016). Sites with less than 6 sampling units were discarded from the final

database since the estimation of the gamma richness was judged non-reliable. Exploratory

analyses revealed especially large deadwood volumes linked with small plots (nugget

effect). To avoid strong leveraging from these outliers (Zuur et al. 2010), we limited the

maximum deadwood volume per sampling plot to 500 m3/ha, a value that corresponds to the

maximal volumes observed in primeval forest of Europe (e.g. Christensen et al. 2005). The

final data distribution per taxonomic and functional groups is shown in Table 1 and Figure 1.

Taxa	I4.2.Regeneration			I4.3.Naturalness			Total
	Coppice	Natural	Planting	Plantation	Semi-natural	Unmanaged	
Tracheophytes	10	734	94	94	550	194	838
Bryophytes	3	353	66	66	198	158	422
Beetles	7	408	94	94	255	160	509
Birds	10	721	94	94	537	194	825
Fungi	8	502	94	94	347	163	604
Lichens	10	332	92	92	333	9	434

Table 1: Number of plots used to fit the models distributed per taxonomic and functional

groups and Regeneration origin or Naturalness following Forest Europe (2020).

Indicators' calculation

We used the State of Europe's Forests (Forest Europe 2020) definitions to calculate the

corresponding metrics for the indicators included in the Criterion 4 (Table 2). Since it was

necessary to "translate" the definition into calculable values to analyze the relationship

between indicator values and *indicanda* (i.e. standardized species richness), we proceeded

as follows:

-    indicator 4.6 *Genetic resources* was excluded since it was not possible to translate

this indicator at the plot or site levels;

-    indicators 4.1 *Diversity of tree species*, 4.4 *Introduced tree species* and 4.5

Deadwood were directly calculated from the plot measurements. Instead of

introduced tree species, we considered only invasive woody species that can have a

significant impact on forest biodiversity following Campagnaro et al. (2018), namely

Acer negundo, *Ailanthus altissima, Prunus serotina*, *Quercus rubra* and *Robinia*

pseudoacacia (Campagnaro et al. 2018). We limited to these species since there

was no regional reference list for introduced tree species and it would have been far

beyond the scope of this study to create such a list, e.g. Norway spruce (*Abies alba*)

is introduced in lowland France but native in mountains where it has also been

planted in some places. In addition, no reference list of invasive tree species in

Europe was available in the Forest Europe (2020) report;

-    indicators 4.2 *Regeneration* and 4.3 *Naturalness* were derived from declarative

metadata when merging the database (Burrascano et al. 2023). In the 4.3, forests

were considered 'undisturbed by man' when declared without intervention by the data

holder (i.e. 'abandoned', 'unmanaged', 'primeval'). Time since the last intervention

was not used here since it was not homogeneously declared. 'Semi-natural' refers to

forest plots that are under any type of forest management regime (from even-aged to

uneven-aged silvicultural systems), but with natural regeneration processes of trees.

'Plantation' forests overlap with forest plots where regeneration derives from

'Planting' activities;

-    indicators 4.8 *Threatened forest species* and 4.10 *Common forest bird species* were

249   derived from biodiversity sampling and reference lists. For 4.8, we used the IUCN

250   Red List and calculated the total number of threatened species (categories VU, EN

251   and CR) per plot all groups together. For 4.10, we calculated the total number per

252   plot of the 34 forest birds classified by the Pan-European Common Birds Monitoring

253   Scheme (https://pecbms.info/trends-and-indicators/indicators/);

254   -   indicator 4.7 *Forest fragmentation* was calculated following the protocol applied by

255   European Commission et al. (2019) on the Corine Land Cover 2018 layer

256   (https://efi.int/knowledge/maps/forest). The forest area density (FAD) at fixed

257   observation scale was obtained calculating the percentage of forest pixels in a 500

258   ha circular buffer, centered on each plot;

259   -   indicator 4.9 *Protected forests* was calculated based on the map of protected areas

260   in Europe provided by the European Environment Agency (Nationally Designated

261   areas - CDDA: https://www.eea.europa.eu/data-and-maps/data/nationally-

262   designated-areas-national-cdda-17). We extracted the proportion of Categories Ia, Ib

263   and II according to the IUCN in a 500ha circular buffer around each plot. We then

264   added these three values so that the total proportion may be higher than 100.

Biodiversity indicators	Definition (Forest Europe 2020)	Data source	Metrics
4.1. Diversity of tree species	Area of forest and other wooded land, classified by number of tree species occurring	Plot measurements	Number of tree species per sampling unit
4.2. Regeneration	Total forest area by stand origin and area of annual forest regeneration and expansion (natural regeneration, planting, coppicing)	Metadata	3 classes: Natural regeneration Planting Coppicing
4.3. Naturalness	Area of forest and other wooded land by class of naturalness (undisturbed by man, semi-natural, plantations)	Metadata	3 classes: Unmanaged Semi-natural Plantation
4.4. Introduced tree species	Area of forest and other wooded land dominated by introduced tree species ⇨ modified to invasive ligneous species (Campagnaro et al. 2018)	Plot measurements	Number of invasive tree species per sampling unit
4.5. Deadwood	Volume of standing deadwood and of lying deadwood on forest and other wooded land	Plot measurements	Deadwood volume per ha
4.6. Genetic resources	Area managed for conservation and utilization of forest tree genetic resources (in situ and ex situ genetic conservation) and area managed for seed production	Not assessed	
4.7. Forest fragmentation (FAD)	Area of continuous forest and of patches of forest separated by non-forest lands (European Commission et al. 2019)	European Forest Institute map of forest cover	Forest area density (FAD) within 500ha around the sampling unit
4.8. Threatened forest species	Number of threatened forest species, classified according to IUCN Red List categories, in relation to total number of forest species	IUCN Red List species list	Species richness of threatened species (categories VU, EN and CR)
4.9. Protected forests	Area of forest and other wooded land protected to conserve biodiversity, landscapes and specific natural elements, according to MCPFE categories	CDDA map of protected areas	Cumulative percentage of Ia, Ib and II categories areas
4.10. Common forest bird species	Occurrence of common breeding bird species related to forest ecosystems	PECBMS	Species richness of forest birds

Table 2: Forest Europe's biodiversity indicators definitions and data sources used to

calculate them in this paper.

Statistical analyses

We processed all analyses in R v.4.3.0 (R Core Team 2023). We used the standardized

species richness of each taxonomic and functional group as a response variable. For each

group, we fitted a single generalized linear mixed model with each indicator as an

explanatory variable (9 models + the null - intercept only - model) and site as a random

effect on the intercept to account for the fact that two plots from the same site were more

likely to be similar than two plots from different sites. We used the beta error distribution with

logit link since standardized richness was continuous, strictly positive with a maximum value

below one. We scaled continuous explanatory variables to improve the convergence and to

make the estimates of the models comparable. We used the glmmTMB function in the

glmmTMB package (Brooks et al. 2017). We compared the magnitudes of all single effects

in the models and their significance and represented this using a heatmap of the

standardized effects (package ggplot2, Wickham 2016). We tested differences between

estimates of categorical variables (e.g. 4.3.naturalness) using Tukey post-hoc test. To

search for potential non-linear relationships between indicators and *indicanda*, we also fitted

generalized additive mixed models (GAMMs) with indicators as smoothers (package mgcv,

Wood 2023). We found very few differences between GLMMs and GAMMs and preferred to

stick to the first (comparisons are presented in Appendix 1).

Finally, to assess the relative performance of all indicators, we first compared the Akaike

Information Criterion corrected for small samples (AICc, Burnham & Anderson 2002) of all

single models including a null (intercept only) model. Then, we searched for the best and

most parsimonious linear combination (no interaction) of indicators that represented

biodiversity based on AICc using the dredge function (package MuMIn, Barton 2023). In this

process, we discarded the indicator 4.2. Regeneration since it was collinear with 4.3.

Naturalness and a model containing both variables could not be fitted (see Appendix 2).

When two competing models had a difference in AICc less than 2 points, we chose the most

parsimonious one.

Results

Data and indicators' distribution

The most represented group in the database was tracheophytes (838 plots) followed by

birds (825 plots, Table 3, see Burrascano et al. 2023 for a full description of the biodiversity

dataset). All taxonomic and functional groups have been inventoried in at least 400 plots

(Table 1). The distribution in the classes Regeneration (4.3.) and Naturalness (4.4.) were,

however, strongly unbalanced (see also Table 1): the majority of sampling units were

associated with "natural regeneration", while "coppicing" - and "planting" to a lesser extent -

were underrepresented; the majority of sampling units were within semi-natural forests, but

the distribution was more balanced than for regeneration types. For quantitative indicators,

the values taken were relatively balanced between groups (Table 3) and no strong

collinearity was observed (see Appendix 2 for an example on tracheophytes).

Taxonomic and functional groups	I4.1.tree.sp	I4.4.invasive	I4.5.deadwood	I4.7.fragmentation (FAD)	I4.8.threat.sp	I4.9.IUCN	I4.10.birds
Tracheophytes	1.96(2.06) [0-11]	0.01(0.08) [0-1]	41.83(60.72) [0-444.89]	0.85(0.16) [0.13-1]	0.25(0.56) [0-3]	14.92(33.66) [0-142.4]	3.63(1.78) [1-11]
Bryophytes	1.91(1.84) [0-11]	0(0.06) [0-1]	39.15(55.86) [0-461.41]	0.84(0.18) [0.06-1]	0.2(0.5) [0-3]	27.02(43.16) [0-142.4]	4.41(2.63) [1-15]
Birds	2.49(2.18) [0-11]	0.01(0.08) [0-1]	42.03(64.56) [0-461.41]	0.85(0.16) [0.13-1]	0.29(0.6) [0-3]	15.92(34.06) [0-142,4]	4.14(2.38) [1-15]
Beetles	1.96(2.01) [0-11]	0(0.07) [0-1]	41.1(57.2) [0-444.89]	0.83(0.17) [0.08-1]	0.26(0.59) [0-5]	16.63(35.16) [0-142.4]	4.15(2.43) [1-12]
Fungi	1.47(1.95) [0-11]	0.01(0.08) [0-1]	39.73(57.98) [0-461.41]	0.82(0,.8) [0.08-1]	0.17(0.49) [0-3]	14.55(34.22) [0-100]	4.32(2.83) [1-15]
Lichens	1.93(1.85) [0-11]	0(0.06) [0-1]	39.28(56.21) [0-461.41]	0.84(0.18) [0.06-1]	0.21(0.53) [0-5]	26.65(42.93) [0-142.4]	4.43(2.62) [1-15]

Table 3: Summary table for the range of values (mean per sampling unit (sd) [min - max])

taken for each indicator distributed over the different subsets. FAD = Forest Area Density.

Correlations between indicators and indicanda

Several of the Forest Europe indicators had significant relationships with the standardized

species richness of one or more of the six taxonomic and functional groups (Figure 2, see

Appendix 3 for values of the estimates).

315
316 Figure 2: Heatmap representing the standardized estimates (slope) of standardized species

317 richness obtained with generalized mixed models with Beta error distribution and logit link,

318 using Forest Europe indicators as independent predictors. S.N = semi-natural, UNM =

319 unmanaged. Intercept (reference) for 4.2 Regeneration is the "Coppice" class, and

320 "Plantations" for 4.3 Naturalness (p<0.1, * p<0.05, ** p<0.01, *** p<0.001).

Figure 3: Relationships between standardized species richness of 6 taxonomic and functional groups and Forest Europe indicator 4.5. Total deadwood volume. Estimations are issued from generalized linear models with Beta error distribution and logit link. Plain line represents the mean estimate, ribbons the 95% confidence interval.

Deadwood (4.5, Figure 3) and Diversity of tree species (4.1) showed four positive and (marginally) significant but generally weak relationships with tracheophytes, bryophytes, fungi and lichens (Appendix 3). Other indicators showed at least three significant (or marginally significant) relationships, with varying magnitudes, i.e. Fragmentation (FAD) (4.7),

Protected areas (4.9, Figure 4) and Forest Birds (4.10). Surprisingly, the proportion of

protected areas around sampling plots had a negative effect on birds and no effect on

tracheophytes and fungi. Regarding Naturalness (4.3, Figure 5), unmanaged forest and

semi-natural forests showed higher levels of fungi, lichens and bryophytes than plantations.

However, the opposite was true for tracheophytes. Threatened species (4.8) showed very

few significant results (marginally positive for tracheophytes and bryophytes, positive for

birds) while Invasive species (4.4) only showed a marginally significant positive relationship

with tracheophytes. Plantation (4.2) showed a very strong negative relationship with lichens

only, and a marginally significant negative effect on tracheophytes, while there were no

differences between coppice and natural regeneration (Appendix 3).

Figure 4: Relationships between standardized species richness of 6 taxonomic and

functional groups and Forest Europe indicator 4.9 (Proportion of protected areas in a 500ha

buffer). Estimations are issued from generalized linear models with Beta error distribution.

Plain line represents the mean estimate, ribbons the 95% confidence interval.

Figure 5: Relationships between standardized species richness of 6 taxonomic and

functional groups and Forest Europe indicator 4.3. Naturalness: PLA = Plantations; S.N =

Semi – Natural forests; UNM = Unmanaged forests. Estimations are issued from generalized

linear models with Beta error distribution and logit link. Barplots represent mean estimates,

error bars are the 95% confidence interval. Letters indicate significant differences per group

based on Tukey post-hoc test.

Relative performance of the different indicators

Comparing the AICc (Table 4) for all single models revealed that the indicator based on

forest Naturalness (4.3) best explained the scaled species richness for three out of six

groups (tracheophytes, bryophytes, and lichens); Regeneration (4.2) was within 2 points of

AICc for tracheophytes and lichens. For birds and beetles, the indicator based on Forest

birds (4.10) was the best explanatory one, while for fungi, Deadwood (4.5) stood first. Also

note that the null model was never the best one and more than 2 AICc points away from the

best.

		Tracheophytes	Bryophytes	Birds	Beetles	Fungi	Lichens
Explanatory variables	**K**	**Delta AICc**	**Delta AICc**	**Delta AICc**	**Delta AICc**	**Delta AICc**	**Delta AICc**
Null	3	14.5	76.9	626.2	2.6	123.1	27.4
I4.1.tree.sp	4	13.3	59.9	627.1	4.2	107.1	18.4
I4.2.regeneration	5	0.5	23.5	629.5	6.0	73.3	1.6
I4.3.naturalness	5	**0.0**	**0.0**	629.7	2.7	33.9	**0.0**
I4.4.invasive	4	13.0	77.8	627.3	4.1	125.1	29.3
I4.5.deadwood	4	14.4	52.3	624.2	4.6	**0.0**	19.3
I4.7.fragmentation (FAD)	4	4.4	70.8	610.9	1.9	121.7	29.3
I4.8.threat.sp	4	12.8	75.2	623.8	4.4	124.1	28.5
I4.9.IUCN	4	16.4	64.0	622.1	2.5	124.4	21.6
I4.10.birds	4	16.3	78.7	**0.0**	**0.0**	115.4	29.4

Table 4: Differences of Akaike Information Criteria corrected for small samples (Delta AICc)

for all generalized mixed models with Beta error distribution and logit link. K is the number of

parameters in each model. The best model (delta AICc = 0) is in bold.

Combination of indicators

We used data dredging to find the best linear (i.e. without interaction) combination of

indicators reflecting the variations in species richness of each group. Only beetles were

represented by only one indicator (Forest Birds, 4.10.), all the other ones had a combination

of 3 to 4 indicators in their best models (Table 5). In terms of indicators, Invasive species

(4.4.) was never selected. Conversely, and as observed before, Diversity of tree species

(4.1.), Naturalness (4.3.), Deadwood (4.5.) were selected four times, followed by Forest

Birds (4.10, 3 times). Finally, it is interesting to note that, except for Fragmentation (FAD,

4.7.), all indicators had positive effects, and with the remarkable exception of beetles, all

multivariate models performed better than single ones (in terms of AICc, they had at least 2

points less than single variable ones, Table 5).

Taxonomic and functional groups	Tracheophytes	Bryophytes	Birds	Beetles	Fungi	Lichens
Intercept	-0.47	-1.28	0.11	-1.03	-2.14	-1.34
I4.1.tree.sp	0.11	0.17			0.12	0.12
I4.3.naturalness	+	+			+	+
I4.4.invasive						
I4.5.deadwood	0.06	0.09			0.17	0.07
I4.7.fragmentation (FAD)	-0.072		-0.07			
I4.8.threat.sp			0.05			
I4.9.IUCN						
I4.10.birds			0.65	0.05	0.08	
Df	8	7	6	4	8	7
AICc	-1246.18	-384.61	-1678.81	-1123.87	-1743.14	-547.79
Delta AICc	0.61	1.42	0.19	0.96	1.23	1.69
AICc (best single variable models)	-1226.72	-375.82	-1648.93	-1123.87	-1671.48	-541.632

Table 5: Scaled estimates of the models selected by data dredging combining all indicators

from Forest Europe (without interaction). Estimates are issued from generalized linear mixed

models with Beta error distribution and logit link. "+" indicates the presence of the factor in

the model. The best model was the most parsimonious (lowest degrees of freedom - df) with

the lowest Akaike Information Criterion corrected for small samples (AICc). "delta" indicates

the AICc difference with the best model. logLik = logarithm of the likelihood.

Discussion

We analyzed the links between Forest Europe's biodiversity indicators and the biodiversity of six taxonomic and functional groups on a unique multi taxonomic dataset covering most forest types in Europe (Burrascano et al. 2023). We showed that these indicators perform unequally: while some correctly describe the biodiversity of several groups, no one acted as a universal indicator, questioning their strength to predict biodiversity in European forests. In contrast, our results reinforced the approach based on multiple complementary indicators of the same *indicandum*. This also raises the question of contextualization of these indicators, to better assess state and trends of biodiversity across Europe, and opens perspectives for improvement.

Performance of current forest biodiversity indicators

While evidence on the link between some indicators and the biodiversity they are supposed to indicate remains incomplete (Gao et al. 2015; Penone et al. 2019; Zeller et al. 2023), our results highlighted several habitat-species relationships and habitat preferences of different groups.

We first observed strong effects of Regeneration origin (4.2) and Naturalness (4.3) on the biodiversity of the six groups. The biodiversity of bryophytes and fungi was higher in unmanaged forests compared to semi natural forests, and, including lichens, higher than in plantation forests (Figure 5). On the contrary, plantations had a marginal positive effect on tracheophytes compared to semi natural and unmanaged forests, and naturalness had no effects on birds and beetles. The negative effects of plantations, and the positive effects of management abandonment or primeval forests have been thoroughly documented (e.g. Chaudhary et al. 2016; Paillet et al. 2010). However, our results should be nuanced by the fact that the distribution of the data in the different categories of regeneration and naturalness were strongly unbalanced, with only few plots located in coppice-managed forests and a vast majority in semi-natural forests (Table 1). That said, unmanaged forests had a generally positively influence on biodiversity in our dataset, especially for deadwood

dependent species (epixylic bryophytes and lichens), while plantations had detrimental

effects on several other groups. Indeed, habitat provision and continuity are higher in semi-

natural forests, not to speak of unmanaged forests, which allow the persistence of dispersal

limited species (e.g. Boch et al. 2013; Boch et al. 2021). The positive response of

tracheophytes in plantations may seem surprising, but probably reflects a higher share of

disturbance tolerant herbs in more disturbed sites, as shown in several individual studies

(Boch et al. 2013). In addition, the semi-natural category encompasses a wide range of

management types (Trentanovi et al. 2023), and unmanaged forests in our dataset have

mostly been recently abandoned, which may cause a decrease in the biodiversity of

tracheophytes during the first decades after abandonment (e.g. Paillet et al. 2010).

Deadwood volume, followed by diversity of tree species, were two indicators that often

correlated with scaled species richness, both in univariate and multivariate analyses.

Deadwood and diversity of tree species have complementary roles for biodiversity (Storch et

al. 2023). Deadwood provides a substrate and a resource for numerous species that depend

on it for part of their life cycle (Lassauce et al. 2011; Müller et al. 2019; Müller et al. 2015;

Parajuli & Markwith 2023), but also for facultative species (Graf et al. 2022). Indeed, the

richness of bryophytes, fungi and lichens, and to a lesser extent birds, correlated positively

to deadwood volume in our analyses. Surprisingly however, saproxylic beetles richness did

not correlate significantly to deadwood volume despite a weak positive effect. It is likely that,

for this group in particular, deadwood does not act as a universal indicator that could be

transposed in all situations (see also Zeller et al. 2023). For example, Lassauce et al. (2011)

showed that the correlation between richness of saproxylic beetles and deadwood volume

was higher in boreal than in temperate forests, a result that was however not confirmed by

Parajuli and Markwith (2023). Müller et al. (2015) showed compensation effects between

deadwood and temperature, supporting that the influence of deadwood on saproxylic beetle

richness declines with increasing temperature, both in geographical space, but also locally in

relation to canopy openness (e.g. Koch Widerberg et al. 2012). It is also possible that the

observed deadwood volumes (mean 40 m^3/ha, Table 3) lead to relatively poor beetle

communities. However, in most cases, the mean deadwood values in our dataset were

above the average deadwood volume per European country (which was 11.5m3/ha on

average, but varied from 2.3 to 28m3/ha between countries, Forest Europe 2020).

Consequently, while the positive role of deadwood for biodiversity remains unchallenged, its

use as an indicator should probably be refined by a better integration of the context

(macroclimate, microclimate, position on the deadwood volume gradient) in the relation. It is

also true that deadwood quality (e.g. including size and decay stage Vítková et al. 2018) is

as much relevant as its quantity. Some saproxylic beetle species (e.g. the red-listed species

Rosalia alpina, Campanaro et al. 2017) develop in large logs, snags and standing dead trees

within canopy gaps. In most of the sampling units we analyzed, deadwood was represented

by relatively small fragments (mostly around 15-20 cm diameter) likely deriving from thinning

processes occurring in closed forests (Burrascano et al. 2023). Further, many rare taxa

depend on the occurrence of tree-related microhabitats (Larrieu et al. 2018), such as the

protected umbrella species *Osmoderma eremita,* which larvae develop in large cavities in

living trees (Dubois et al. 2009). Since several tree-related microhabitats are very scarce in

managed forests (Paillet et al. 2017), and not readily quantified as part of the deadwood pool

(Müller et al. 2014), their role is not currently taken into consideration.

Diversity of tree species also showed positive correlation with the biodiversity of several

taxonomic and functional groups (bryophytes, fungi, lichens and more marginally

tracheophytes). Different tree species provide different habitat conditions for epiphytic and

saproxylic species living in forests (e.g. Boch et al. 2021; Cavard et al. 2011; Leidinger et al.

2021). These conditions encompass direct biotic interactions, different chemical properties of

the bark and wood, decomposition rates as well as differential light interception (Gosselin et

al. 2017; Zeller et al. 2023). This in turn provides niche heterogeneity for several species and

thus enhances biodiversity and confirms the role of diversity of tree species as a biodiversity

indicator.

Among the other indicators, forest fragmentation showed negative effects on tracheophytes

and birds, and a positive effect on bryophytes, and marginally on beetles and fungi. It should

be noted that since it is based on forest area density (European Commission et al. 2019), it

represents forest cover rather than its discontinuity and configuration, but has the advantage

of being calculable and homogeneous across Europe. Besides, the landscapes surrounding

our sampling units presented a high forest cover (77% on average, in a 500 ha buffer,

Table 1). Thus, the observed correlations reflect that forest area density could have been

beneficial for forest interior species that are sensitive to edge effects (bryophytes or lichens),

but detrimental to non-forest, open habitat or light demanding species, such as

tracheophytes and open-habitat birds (but see Bełcik et al. 2020, who show an increase

before a plateau in taxonomic diversity with increasing forest area in agricultural

landscapes). The inferences on the effect of fragmentation on biodiversity, based on the

Single Large vs Several Small principle (Diamond 1975), support that large continuous

habitat should have a greater effect on biodiversity conservation, compared to several small

ones. However, recent multitaxa studies proved the disproportionately high value of small

patches, harboring richer assemblages, even when focusing only on protected species (Riva

& Fahrig 2022). Besides, it has been highlighted that biodiversity conservation in human-

modified forest landscapes is better achieved maintaining at least 40% of forest cover, rather

than focusing on fragmentation and configuration (Arroyo-Rodríguez et al. 2020). Hence, the

role of fragmentation, as it is currently estimated, should be probably be reconsidered to take

into account both the amount of forest cover and the value of the patches.

Contrary to the previous indicators, threatened species and birds are based on direct

biodiversity sampling (other than trees). These indicators directly describe the evolution of a

small part of the biodiversity, but generally correlate poorly to the richness of other taxa and

most of the results observed are linked to the fact that these indicators represent subsets of

larger groups (evidently forest birds vs. birds). However, despite several pieces of evidence

showing that congruence between taxa is generally small, especially in forests (Burrascano

et al. 2018; Westgate et al. 2017), forest birds positively correlated to beetles and fungi.

Probably these groups respond to the same favorable habitat conditions, but this was not

reflected in the multivariate analyses.

The proportion of IUCN-protected areas around the plots positively influenced the richness

of bryophytes and lichens, highlighting the positive effects of protection for these groups, but

surprisingly had a negative effect on birds (see Honkanen et al. 2010).

Finally, invasive tree species did not show any correlation with biodiversity, apart from a

marginal positive effect on tracheophytes supporting high local tracheophyte diversity to be

often an indicator of disturbance. Despite the negative effects of invasive species

introduction on native flora, fauna and mycota, this effect is rather limited in our dataset

since it contains very few invasive species in total (maximum one tree species per plot)

reflecting that no projects targeted the effects of plantations rich in, or dominated by,

introduced tree species on biodiversity. This limited gradient probably does not allow to show

significant effects on biodiversity. In addition, an indicator on introduced species rather than

invasive ones – as it is defined in Forest Europe's indicators – may perform better.

Limitations of Forest Europe indicators to predict biodiversity patterns and implications for

improvements

Our study is one of the first to test Forest Europe indicators against multitaxonomic data at

the European level. During the model selection process, the best single variable models

were more often involving naturalness (tracheophytes, bryophytes and lichens) followed by

forest birds (birds and beetles) and deadwood (fungi). However, the multivariate models did

perform better than single ones (except for beetles), and often the best ones combined

several indicators to reflect variations of the *indicandum*. Only beetles were best indicated by

forest birds, but with a low magnitude, and birds by three combined indicators (including

forest birds and threatened species). The magnitude of the effects in the multivariate models

were comparable to those of single variable models (Appendix 3) which confirmed that

indicators were not collinear.

We studied only total richness as a response variable, and evidently, the results may be

different for other, more specialized groups, or other metrics of biodiversity (abundance,

occurrence of individual species, functional diversity). Such approach remains to be tested

(see e.g. Lelli et al. 2019) but was beyond the scope of the present study. In addition, some

of the references we used were probably incomplete regarding some groups: e.g. almost no

lichens are included in the list we used for red-listed species, but the proportion of

threatened species at the national levels may be high.

We limited our approach to the strict definition of the indicators as used in Forest Europe, but

higher performance could probably be reached by at least two improvements. First, adding

context to the indicators could probably reveal that they need to be adapted locally

(Chiarucci et al. 2012; Honkanen et al. 2010). Examples of context could be: elevation

(mountain vs. lowlands), biome (Mediterranean, temperate, boreal) and European forest

types. Second, most of the metrics we used are abundance metrics (apart from tree species

richness) that quantify habitat available for species. However, indicators based on diversity

of resources (following the heterogeneity-diversity theory, Tews et al. 2004) could perform

better, e.g. in the case of deadwood and saproxylic beetles (Bouget et al. 2013), or tree-

related microhabitats and birds and bats (Paillet et al. 2018). It would then be interesting to

assess the performance of other indicator metrics vs. the current ones in assessing forest

biodiversity.

Conclusions: towards new indicators definitions and better reporting

Many forest biodiversity indicators are proxies based on pre-existing data mostly issued from

National Forest Inventories (Tomppo et al. 2010). Despite recent progresses based on

international initiatives (namely the Essential Biodiversity Variables, GEOBON, IPBES),

monitoring the state and trends of forest biodiversity solely based on proxies is not

satisfactory: while proxies are generally easier to measure than species themselves, they

are prone to demographic effects such as extinction debts or colonization credits. In other

words, the presence of a given habitat - such as deadwood - does not guarantee the

presence of the species that depend on it (e.g. Paillet et al. 2018). In addition, the response

of biodiversity to a given indicator depends on the taxonomic or functional group studied

(Zeller et al. 2022), so no indicator may represent biodiversity overall. This was confirmed by

our analyses. We showed that, while some indicators performed correctly for several

taxonomic and functional groups in terms of significance and magnitude, we did not highlight

a predominant role of one indicator over the others.

Stevenson et al. (2021) claimed that indicators were often implemented without clear

considerations of their purposes and utility in terms of decision-making. We argue that, while

combinations of current Forest Europe indicators are useful to delineate general trends in

biodiversity, taking into account context and analyzing the performance of other - more

diversity-driven - metrics would help better reporting on biodiversity (e.g. Alterio et al. 2023;

Paillet et al. 2018). Such improvements would also be beneficial to the use of indicators

beyond general trends to include evaluating management and policy actions, decisions, or

set biodiversity targets (Stevenson et al. 2021). For example, we showed that larger

negative effects on biodiversity were observed in planted forests. This poses key challenges

for making the 3 billion trees planting promoted by the European Forest Strategy for 2030 a

beneficial action for biodiversity in forests (Sills et al. 2020), if high growing rate plantations

such as introduced Douglas fir (*Pseudotsuga menziesii*) or eucalyptus (*Eucalyptus* spp.) are

promoted against semi-natural forests. Conversely, promoting old-growth and unmanaged

forests, as well as restoration of monocultures and conversions towards semi-natural forests,

could have a positive effect on biodiversity. It is crucial to assess and balance these potential

effects with the use of current data available and biodiversity indicators before taking actions

or to modulate actions in favor of biodiversity against detrimental ones.

References

Alterio, E., T. Campagnaro, L. Sallustio, S. Burrascano, L. Casella, and T. Sitzia. 2023.
Forest management plans as data source for the assessment of the conservation
status of European Union habitat types. Frontiers in Forests and Global Change **5**.
Arroyo-Rodríguez, V., et al. 2020. Designing optimal human-modified landscapes for forest
biodiversity conservation. Ecology Letters **23**:1404-1420.
Barton, K. 2023. MuMIn: multi-model inference. in R package version 1.47.5.
Bełcik, M., M. Lenda, T. Amano, and P. Skórka. 2020. Different response of the taxonomic,
phylogenetic and functional diversity of birds to forest fragmentation. Scientific
Reports **10**:20320.
Betts, M. G., C. Wolf, W. J. Ripple, B. Phalan, K. A. Millers, A. Duarte, S. H. M. Butchart, and
T. Levi. 2017. Global forest loss disproportionately erodes biodiversity in intact
landscapes. Nature **547**:441-444.
Boch, S., D. Prati, D. Hessenmöller, E. D. Schulze, and M. Fischer. 2013. Richness of
Lichen Species, Especially of Threatened Ones, Is Promoted by Management
Methods Furthering Stand Continuity. PLoS ONE **8**.
Boch, S., H. Saiz, E. Allan, P. Schall, D. Prati, E.-D. Schulze, D. Hessenmöller, L. B.
Sparrius, and M. Fischer. 2021. Direct and Indirect Effects of Management Intensity
and Environmental Factors on the Functional Diversity of Lichens in Central
European Forests. Microorganisms.
Bouget, C., L. Larrieu, B. Nusillard, and G. Parmain. 2013. In search of the best local habitat
drivers for saproxylic beetle diversity in temperate deciduous forests. Biodiversity and
Conservation **22**:2111-2130.
Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H.
J. Skaug, M. Maechler, and B. M. Bolker. 2017. glmmTMB Balances Speed and
Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling.
The R Journal, **9**:378-400.
Burnham, K. P., and D. R. Anderson 2002. Model Selection and Multimodel Inference: A
Practical Information-Theoretic Approach (2nd ed).
Burrascano, S., et al. 2023. Where are we now with European forest multi-taxon biodiversity
and where can we head to? Biological Conservation **284**.
Burrascano, S., et al. 2018. Congruency across taxa and spatial scales: are we asking too
much of species data? Global Ecology and Biogeography **27**:980-990.
Campagnaro, T., G. Brundu, and T. Sitzia. 2018. Five major invasive alien tree species in
European Union forest habitat types of the Alpine and Continental biogeographical
regions. Journal for Nature Conservation **43**:227-238.
Campanaro, A., et al. 2017. Guidelines for the monitoring of Rosalia alpina. Nature
Conservation **20**:165-203.
Cavard, X., S. E. Macdonald, Y. Bergeron, and H. Y. H. Chen. 2011. Importance of
mixedwoods for biodiversity conservation: Evidence for understory plants, songbirds,
soil fauna, and ectomycorrhizae in northern forests. Environmental Reviews **19**:142-
161.
Chaudhary, A., Z. Burivalova, L. P. Koh, and S. Hellweg. 2016. Impact of Forest
Management on Species Richness: Global Meta-Analysis and Economic Trade-Offs.
Scientific Reports **6**:23954.
Chiarucci, A., G. Bacaro, G. Filibeck, S. Landi, S. Maccherini, and A. Scoppola. 2012. Scale
dependence of plant species richness in a network of protected areas. Biodiversity
and Conservation **21**:503-516.
Chirici, G., et al. 2012. National forest inventory contributions to forest biodiversity
monitoring. Forest Science **58**:257-268.
Christensen, M., et al. 2005. Dead wood in European beech (Fagus sylvatica) forest
reserves. Forest Ecology and Management **210**:267-282.
Diamond, J. M. 1975. The island dilemma: Lessons of modern biogeographic studies for the
design of natural reserves. Biological Conservation **7**:129-146.

Dubois, G. F., V. Vignon, Y. R. Delettre, Y. Rantier, P. Vernon, and F. Burel. 2009. Factors affecting the occurrence of the endangered saproxylic beetle Osmoderma eremita (Scopoli, 1763) (Coleoptera: Cetoniidae) in an agricultural landscape. Landscape and Urban Planning **91**:152-159.

European Commission, Joint Research Centre, P. Vogt, K. Riitters, G. Caudullo, B. Eckhardt, and R. Raši 2019. An approach for pan-European monitoring of forest fragmentation. Publications Office.

Forest Europe. 2020. State of Europe's Forests 2020. Page 394. Ministerial Conference on the Protection of Forests in Europe - FOREST EUROPE, Liaison Unit Bratislava.

Gao, T., A. B. Nielsen, and M. Hedblom. 2015. Reviewing the strength of evidence of biodiversity indicators for forest ecosystems in Europe. Ecological Indicators **57**:420-434.

Gosselin, M., D. Fourcin, Y. Dumas, F. Gosselin, N. Korboulewsky, M. Toïgo, and P. Vallet. 2017. Influence of forest tree species composition on bryophytic diversity in mixed and pure pine (Pinus sylvestris L.) and oak (Quercus petraea (Matt.) Liebl.) stands. Forest Ecology and Management **406**:318-329.

Graf, M., S. Seibold, M. M. Gossner, J. Hagge, I. Weiß, C. Bässler, and J. Müller. 2022. Coverage based diversity estimates of facultative saproxylic species highlight the importance of deadwood for biodiversity. Forest Ecology and Management **517**:120275.

Group on Earth Observation Biodiversity Observation Network. 2008. GEO BON Concept Document. GEO-V document 20, Geneva, Switzerland.

Heym, M., E. Uhl, R. Moshammer, J. Dieler, K. Stimm, and H. Pretzsch. 2021. Utilising forest inventory data for biodiversity assessment. Ecological Indicators **121**:107196.

Honkanen, M., J.-M. Roberge, A. Rajasärkkä, and M. Mönkkönen. 2010. Disentangling the effects of area, energy and habitat heterogeneity on boreal forest bird species richness in protected areas. Global Ecology and Biogeography **19**:61-71.

Hsieh, T. C., K. H. Ma, and A. Chao. 2016. iNEXT: an R package for rarefaction and extrapolation of species diversity (Hill numbers). Methods in Ecology and Evolution **7**:1451-1456.

IPBES. 2019. Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (Version 1). Page 1144 in E. D. Brondizio, S., and J. N. Settele, H.T., editors. IPBES secretariat, Bonn, Germany.

Jiguet, F., V. Devictor, R. Julliard, and D. Couvet. 2012. French citizens monitoring ordinary birds provide tools for conservation and ecological sciences. Acta Oecologica **44**:58-66.

Kempeneers, P., F. Sedano, L. Seebach, P. Strobl, and J. San-Miguel-Ayanz. 2011. Data Fusion of Different Spatial Resolution Remote Sensing Images Applied to Forest-Type Mapping. IEEE Transactions on Geoscience and Remote Sensing **49**:4977-4986.

Koch Widerberg, M., T. Ranius, I. Drobyshev, U. Nilsson, and M. Lindbladh. 2012. Increased openness around retained oaks increases species richness of saproxylic beetles. Biodiversity and Conservation **21**:3035-3059.

Larrieu, L., et al. 2018. Tree related microhabitats in temperate and Mediterranean European forests: A hierarchical typology for inventory standardization. Ecological Indicators **84**:194-207.

Lassauce, A., Y. Paillet, H. Jactel, and C. Bouget. 2011. Deadwood as a surrogate for forest biodiversity: Meta-analysis of correlations between deadwood volume and species richness of saproxylic organisms. Ecological Indicators **11**:1027-1039.

Leidinger, J., et al. 2021. Shifting tree species composition affects biodiversity of multiple taxa in Central European forests. Forest Ecology and Management **498**:119552.

Lelli, C., et al. 2019. Biodiversity response to forest structure and management: Comparing species richness, conservation relevant species and functional diversity as metrics in forest conservation. Forest Ecology and Management **432**:707-717.

Lindenmayer, D. B., and G. E. Likens. 2010. The science and application of ecological
monitoring. Biological Conservation **143**:1317-1328.
Müller, J., S. Boch, D. Prati, S. A. Socher, U. Pommer, D. Hessenmöller, P. Schall, E. D.
Schulze, and M. Fischer. 2019. Effects of forest management on bryophyte species
richness in Central European forests. Forest Ecology and Management **432**:850-859.
Müller, J., et al. 2015. Increasing temperature may compensate for lower amounts of dead
wood in driving richness of saproxylic beetles. Ecography **38**:499-509.
Müller, J., A. Jarzabek-Müller, H. Bussler, and M. M. Gossner. 2014. Hollow beech trees
identified as keystone structures for saproxylic beetles by analyses of functional and
phylogenetic diversity. Animal Conservation **17**:154-162.
Paillet, Y., F. Archaux, V. Boulanger, N. Debaive, M. Fuhr, O. Gilg, F. Gosselin, and E.
Guilbert. 2017. Snags and large trees drive higher tree microhabitat densities in strict
forest reserves. Forest Ecology and Management **389**:176-186.
Paillet, Y., F. Archaux, S. du Puy, V. Boulanger, N. Debaive, M. Fuhr, O. Gilg, F. Gosselin,
and E. Guilbert. 2018. The indicator side of tree microhabitats: a multi-taxon
approach based on bats, birds and saproxylic beetles. Journal of Applied Ecology
55:2147-2156.
Paillet, Y., et al. 2010. Biodiversity Differences between Managed and Unmanaged Forests:
Meta-Analysis of Species Richness in Europe. Conservation Biology **24**:101-112.
Parajuli, R., and S. H. Markwith. 2023. Quantity is foremost but quality matters: A global
meta-analysis of correlations of dead wood volume and biodiversity in forest
ecosystems. Biological Conservation **283**.
Penone, C., et al. 2019. Specialisation and diversity of multiple trophic groups are promoted
by different forest features. Ecology Letters **22**:170-180.
Pereira, H. M., and H. David Cooper. 2006. Towards the global monitoring of biodiversity
change. Trends in Ecology & Evolution **21**:123-129.
Proença, V., et al. 2017. Global biodiversity monitoring: From data sources to Essential
Biodiversity Variables. Biological Conservation **213**:256-263.
R Core Team. 2023. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria.
Reise, J., F. Kukulka, M. Flade, and S. Winter. 2019. Characterising the richness and
diversity of forest bird species using National Forest Inventory data in Germany.
Forest Ecology and Management **432**:799-811.
Rigal, S., et al. 2023. Farmland practices are driving bird population decline across Europe.
Proceedings of the National Academy of Sciences **120**:e2216573120.
Riva, F., and L. Fahrig. 2022. The disproportionately high value of small patches for
biodiversity conservation. Conservation Letters **15**:e12881.
Sills, J., S. Gómez-González, R. Ochoa-Hueso, and J. G. Pausas. 2020. Afforestation falls
short as a biodiversity strategy. Science **368**:1439-1439.
Simons, N. K., et al. 2021. National Forest Inventories capture the multifunctionality of
managed forests in Germany. Forest Ecosystems **8**:5.
Stevenson, S. L., K. Watermeyer, G. Caggiano, E. A. Fulton, S. Ferrier, and E. Nicholson.
2021. Matching biodiversity indicators to policy needs. Conservation Biology **35**:522-
532.
Storch, F., et al. 2023. Linking structure and species richness to support forest biodiversity
monitoring at large scales. Annals of Forest Science **80**:3.
Tews, J., U. Brose, V. Grimm, K. Tielbörger, M. C. Wichmann, M. Schwager, and F. Jeltsch.
2004. Animal species diversity driven by habitat heterogeneity/diversity: The
importance of keystone structures. Journal of Biogeography **31**:79-92.
Tomppo, E., T. Gschwantner, M. Lawrence, and R. E. Mc Roberts 2010. National forest
inventories. Pathways for common reporting. Springer Science, Heidelberg,
Allemagne.
Trentanovi, G., et al. 2023. Words apart: Standardizing forestry terms and definitions across
European biodiversity studies. Forest Ecosystems **10**:100128.

Vítková, L., R. Bače, P. Kjučukov, and M. Svoboda. 2018. Deadwood management in
Central European forests: Key considerations for practical implementation. Forest
Ecology and Management **429**:394-405.
Weber, D., U. Hintermann, and A. Zangger. 2004. Scale and trends in species richness:
considerations for monitoring biological diversity for political purposes. Global
Ecology and Biogeography **13**:97-104.
Westgate, M. J., A. I. T. Tulloch, P. S. Barton, J. C. Pierson, and D. B. Lindenmayer. 2017.
Optimal taxonomic groups for biodiversity assessment: a meta-analytic approach.
Ecography **40**:539-548.
Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer Cham.
Wood, S. 2023. Mixed GAM Computation Vehicle with Automatic Smoothness Estimation.
V1.9.
Zeller, L., et al. 2022. Index of biodiversity potential (IBP) versus direct species monitoring in
temperate forests. Ecological Indicators **136**:108692.
Zeller, L., A. Förster, C. Keye, P. Meyer, C. Roschak, and C. Ammer. 2023. What does
literature tell us about the relationship between forest structural attributes and
species richness in temperate forests? – A review. Ecological Indicators **153**:110383.
Zuur, A. F., E. N. Ieno, and C. S. Elphick. 2010. A protocol for data exploration to avoid
common statistical problems. Methods in Ecology and Evolution **1**:3-14.
756

757 **Appendices**

758 Appendix 1: Comparison of linear (GLMMs) vs non-linear (GAMMs) models analyses

759 between standardized species richness and several Forest Europe's biodiversity indicators.

Appendix 2: Correlation between indicators of Forest Europe from the tracheophytes

dataset. Figures on the right are Pearson correlation coefficients, biplots on the left and

histograms on the diagonal represent the distribution of the data. Regeneration (4.2) and

Naturalness (4.3) are categorical variables not represented here.

Appendix 3: Standardized richness estimates table for all single generalized mixed models

with beta error distribution and logit link. Intercept for I4.2.Regeneration is "Coppice", and for

I4.3.Naturalness "Plantation" (PLA). S.N = semi-natural forest, UNM = Unmanaged forests.

se = standard error of the mean, pl = critical probability, (*) p<0.1, * p<0.05, ** p<0.01, ***

p<0.001.

Group	variable	estimate	se	p	
Tracheophytes	I4.1.tree.sp	0.0648	0.0359	0.0712	(*)
	I4.2.regeneration(natural regeneration)	0.1431	0.2606	0.5828	
	I4.2.regeneration(planting)	0.5186	0.2754	0.0597	(*)
	I4.3.naturalnessUNM	-0.3878	0.0889	<0.001	***
	I4.3.naturalnessS.N	-0.2073	0.1429	0.147	
	I4.4.invasive	0.0376	0.0193	0.0508	(*)
	I4.5.deadwood	0.0318	0.0219	0.1463	
	I4.7.fragmentation	-0.0898	0.0256	<0.001	***
	I4.8.threat.sp	0.0428	0.0221	0.0522	(*)
	I4.9.IUCN	-0.0087	0.0416	0.8344	
	I4.10.birds	0.0122	0.0326	0.707	
Bryophytes	I4.1.tree.sp	0.2828	0.0633	<0.001	***
	I4.2.regeneration(natural regeneration)	0.4156	0.4546	0.3606	
	I4.2.regeneration(planting)	-0.5511	0.4716	0.2426	
	I4.3.naturalnessUNM	0.8606	0.128	<0.001	***
	I4.3.naturalnessS.N	1.5727	0.1745	<0.001	***
	I4.4.invasive	-0.0409	0.0388	0.2917	
	I4.5.deadwood	0.2097	0.04	<0.001	***
	I4.7.fragmentation	0.1384	0.0486	0.0044	**
	I4.8.threat.sp	0.0776	0.0402	0.0539	(*)
	I4.9.IUCN	0.2268	0.0585	<0.001	***
	I4.10.birds	0.0235	0.0571	0.6806	
Birds	I4.1.tree.sp	0.0341	0.0323	0.2916	
	I4.2.regeneration(natural regeneration)	0.0996	0.2114	0.6375	
	I4.2.regeneration(planting)	0.0452	0.2249	0.8407	

Group	variable	estimate	se	p	
	I4.3.naturalnessUNM	0.0388	0.0785	0.6207	
	I4.3.naturalnessS.N	0.2011	0.1259	0.1104	
	I4.4.invasive	-0.0177	0.0186	0.3417	
	I4.5.deadwood	0.0402	0.0203	0.0469	*
	I4.7.fragmentation	-0.1018	0.0243	<0.001	***
	I4.8.threat.sp	0.043	0.0205	0.0365	*
	I4.9.IUCN	-0.1031	0.041	0.012	*
	I4.10.birds	0.6521	0.0218	<0.001	***
Beetles	I4.1.tree.sp	0.0242	0.034	0.4762	
	I4.2.regeneration(natural regeneration)	0.1411	0.2247	0.53	
	I4.2.regeneration(planting)	0.1782	0.233	0.4444	
	I4.3.naturalnessUNM	-0.0338	0.0638	0.5965	
	I4.3.naturalnessS.N	-0.0749	0.109	0.4917	
	I4.4.invasive	-0.0135	0.0175	0.441	
	I4.5.deadwood	0.0045	0.0204	0.8238	
	I4.7.fragmentation	0.0435	0.0263	0.0983	(*)
	I4.8.threat.sp	0.0096	0.0196	0.6255	
	I4.9.IUCN	-0.0526	0.0357	0.1409	
	I4.10.birds	0.05	0.023	0.0297	*
Fungi	I4.1.tree.sp	0.1476	0.0343	<0.001	***
	I4.2.regeneration(natural regeneration)	0.1607	0.2444	0.5109	
	I4.2.regeneration(planting)	-0.3632	0.2542	0.1531	
	I4.3.naturalnessUNM	0.5017	0.0691	<0.001	***
	I4.3.naturalnessS.N	0.9083	0.1434	<0.001	***
	I4.4.invasive	0.0035	0.0189	0.8526	
	I4.5.deadwood	0.2228	0.0181	<0.001	***
	I4.7.fragmentation	0.054	0.0292	0.0644	(*)
	I4.8.threat.sp	0.0226	0.022	0.3039	
	I4.9.IUCN	0.0422	0.0466	0.366	
	I4.10.birds	0.1021	0.0324	0.0016	**
Lichens	I4.1.tree.sp	0.1714	0.0507	<0.001	***

Group	variable	estimate	se	p	
	I4.2.regeneration(natural regeneration)	0.0169	0.258	0.9479	
	I4.2.regeneration(planting)	-0.5447	0.2736	0.0465	*
	I4.3.naturalnessUNM	0.5513	0.1028	<0.001	***
	I4.3.naturalnessS.N	0.7687	0.2439	0.0016	**
	I4.4.invasive	0.0092	0.0313	0.7691	
	I4.5.deadwood	0.1139	0.0343	<0.001	***
	I4.7.fragmentation	0.0137	0.0442	0.7564	
	I4.8.threat.sp	0.0339	0.0348	0.329	
	I4.9.IUCN	0.1778	0.061	0.0036	**
	I4.10.birds	0.0067	0.0559	0.904	