The transformation of sensory to perceptual braille letter representations in the visually deprived brain

Marleen Haupt[1,¶,*], Monika Graumann[1,2,¶], Santani Teng[3], Carina Kaltenbach[1], Radoslaw M. Cichy[1,2,4]

[1] Department of Education and Psychology, Freie Universität Berlin, Berlin, Germany
[2] Berlin School of Mind and Brain, Faculty of Philosophy, Humboldt-Universität zu Berlin, Berlin, Germany
[3] Smith-Kettlewell Eye Research Institute, San Francisco, USA
[4] Bernstein Center for Computational Neuroscience Berlin, Berlin, Germany

[¶] these authors contributed equally
[*] correspondence to: marleen.haupt@gmail.com

Acknowledgements:
We thank all of our participants for taking part in our experiments. We also thank Agnessa Karapetian, Johannes Singer and Siying Xie for their valuable comments on the manuscript. We acquired EEG and fMRI data at the Center for Cognitive Neuroscience (CCNB), Freie Universität Berlin, and we thank the HPC Service of ZEDAT, Freie Universität Berlin, for computing time [1].

Funding:
The study was supported by German Research Council grants (CI241/1-1, CI241/3-1, CI241/7-1) and European Research Council grants (ERC-StG-2018-803370) to R.M.C. The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Author contributions:
M.G., S.T., and R.M.C. designed research. M.G. and C.K. collected data. M.H. and M.G. analyzed data. M.H. and R.M.C. wrote the manuscript. M.G. and S.T. provided feedback on the manuscript. R.M.C acquired funding.

Competing interests:
The authors declare no competing interests.

1. Abstract

Experience-based plasticity of the human cortex mediates the influence of individual experience on cognition and behavior. The complete loss of a sensory modality is among the most extreme such experiences. Investigating such a selective, yet extreme change in experience allows for the characterization of experience-based plasticity at its boundaries.

Here, we investigated information processing in individuals who lost vision at birth or early in life by probing the processing of braille letter information. We characterized the transformation of braille letter information from sensory representations depending on the reading hand to perceptual representations that are independent of the reading hand.

Using a multivariate analysis framework in combination with fMRI, EEG and behavioral assessment, we tracked cortical braille representations in space and time, and probed their behavioral relevance.

We located sensory representations in tactile processing areas and perceptual representations in sighted reading areas, with the lateral occipital complex as a connecting "hinge" region. This elucidates the plasticity of the visually deprived brain in terms of information processing.

Regarding information processing in time, we found that sensory representations emerge before perceptual representations. This indicates that even extreme cases of brain plasticity adhere to a common temporal scheme in the progression from sensory to perceptual transformations.

Ascertaining behavioral relevance through perceived similarity ratings, we found that perceptual representations in sighted reading areas, but not sensory representations in tactile processing areas are suitably formatted to guide behavior.

Together, our results reveal a nuanced picture of both the potentials and limits of experience-dependent plasticity in the visually deprived brain.

2. Introduction

Human brains vary due to individual experiences. This so-called experience-based plasticity of the human cortex mediates cognitive and behavioral adaptation to changes in the environment [2]. Typically, plasticity reflects learning from species-typical experiences. However, plasticity also results from species-atypical changes to experience like the loss of a sensory modality.

Sensory loss constitutes a selective, yet large-scale change in experience that offers a unique experimental opportunity to study cortical plasticity at its boundaries [3]. One deeply investigated case of sensory loss is blindness, i.e., the lack of visual input to the brain. Previous research has shown that cortical structures most strongly activated by visual input in sighted brains are activated by a plethora of other cognitive functions in visually deprived brains [4], including braille reading [5–11]. However, overlapping functional responses alone cannot inform us about the nature of the observed activations, i.e., what kind of information they represent and thus what role they play in cognitive processing.

To elucidate the nature of information processing in the visually deprived brain, we investigate the tactile braille system in individuals who lost vision at birth or early in life (hereafter blind participants). Braille readers commonly use both hands, requiring their brain to transform sensory tactile input into a hand-independent perceptual format. We made use of this practical everyday requirement to experimentally characterize the transformation of sensory to perceptual braille letter representations. We operationalize sensory braille letter representations as representations coding information specific to the hand that was reading (hand-dependent). In contrast, we operationalize perceptual braille letter representations as representations coding information independent of which hand was reading (hand-independent).

53 Combining this operationalization with fMRI and EEG in a multivariate analysis framework [12–15], we
54 determine the cortical location and temporal emergence of sensory and perceptual representations.
55 Lastly, to ascertain the functional role of the identified representations, we relate them to behavioral
56 similarity ratings [16–19].
57

3. Results

59
Figure 1. Stimuli, experimental design, pattern extraction and multivariate analysis framework.
A) We presented eight braille letters (B,C,D,L,M,N,V,Z) to participants on braille cells. Two additional letters (E
and O) served as catch trials and were excluded from all analyses.
B) Top: During the fMRI session, braille letters were presented for 500ms with an inter-stimulus interval (ISI) of
2500ms. Participants were instructed to respond to catch trials by pressing a button with their foot.
Bottom: During a separate EEG session, braille letters were presented for 500ms. The ISI following regular trials
lasted 500ms, the ISI following catch trials lasted 1,100ms to avoid movement confounds. Participants were
instructed to respond to catch trials by pressing a foot pedal.
C) In fMRI, we extracted voxel-wise activations for every region of interest (ROI). In EEG, we extracted channel-
wise activations for every time point. In both cases, this resulted in one response vector per letter and per
experimental run.
D) For both fMRI and EEG, we divided pattern vectors into training (4 pseudo-runs) and test (1 pseudo-run) sets.
For every pair of braille letters (e.g., B and V), we trained a support vector machine (SVM) to classify between
pattern vectors related to the presentation of both letters read with the same hand. We then tested the SVM on the
left-out pattern vectors related to the presentation of the same two letters read with the same hand (within-hand
classification) or with the other hand (across-hand classification). The resulting pairwise decoding accuracies
were aggregated in a decoding accuracy matrix that is symmetric along the diagonal, with the diagonal itself being
undefined. We interpret the within-hand matrix (black) as a measure of sensory and perceptual braille letter
representations. We interpret the across-hand matrix (green) as a measure of perceptual braille letter
representations. We derive the measure of sensory braille letter representations (blue) by subtracting one matrix
from the other.
81
82 We recorded fMRI (N = 15) and EEG (N = 11) data while blind participants (see Supplementary Table
83 1) read braille letters with their left or right index finger. We delivered the braille stimuli using single
84 piezo-electric refreshable cells. This allowed participants to read braille letters without moving their
85 finger, thus avoiding finger motion artifacts in the brain signal and analyses.
86
We used a common experimental paradigm for fMRI and EEG that was adapted to the specifics of each
imaging modality. The common stimulus set consisted of ten different braille letters (Fig. 1a). Eight
letters entered the main analysis. Two letters (E and O) served as vigilance targets to which participants
responded with their foot; these trials were excluded from all analyses. The stimuli were presented in
random order, with each trial consisting of a 500ms stimulus presentation to either the right or left hand.
In fMRI, all trials were followed by an inter-stimulus interval (ISI) of 2,500ms to account for the

sluggishness of the BOLD response (Fig. 1b). In EEG, standard trials had an ISI of 500ms while catch
trials had an ISI of 1,100ms in order to avoid movement contaminations.
95
96 The common experimental paradigm for fMRI and EEG allowed us to use an equivalent multivariate
97 classification scheme to track the transformation of sensory to perceptual representations. We assessed
98 fMRI voxel patterns to reveal where sensory braille letter representations are located in the cortex.
99 Likewise, we assessed EEG electrode patterns to reveal the temporal dynamics of braille letter
100 representations (Fig. 1c).
101
We operationalized sensory versus perceptual representations as hand-dependent versus hand-
independent braille letter representations, respectively. To measure perceptual representations, we
trained classifiers on brain data recorded during stimulation of one hand and tested the classifiers on
data recorded during the stimulation of the other hand (Fig. 1d). We refer to this analysis as across-hand
classification. It reveals perceptual braille letter representations that are independent of the specific hand
being stimulated.
108
109 To assess sensory representations, we used a two-step procedure. In a first step, we trained and tested
110 classifiers on brain data recorded during stimulation of the same hand (Fig. 1d). We refer to this analysis
111 as within-hand classification. It reveals both sensory and perceptual braille letter representations. Thus,
112 to further isolate sensory representations from perceptual representations, in a second step, we
113 subtracted across-hand classification from within-hand classification results.
114

 3.1 Spatial dynamics of braille letter representations

 Figure 2. Spatial dynamics of braille letter representations.
A) Rendering of regions of interests associated with tactile processing (S1: primary somatosensory cortex, S2: secondary somatosensory cortex, aIPS: anterior intraparietal sulcus, pIPS: posterior intraparietal sulcus, insula) and sighted reading (EVC: early visual cortex, V4: visual area 4, LOC: lateral occipital complex, LFA: letter form area, VWFA: visual word form area).
B) Sensory and perceptual (left), sensory (middle) and perceptual (right) braille letters representations in tactile processing and sighted reading ROIs (N = 15, two-tailed Wilcoxon signed-rank test, $P < 0.05$, FDR corrected; Stars below bars indicate significance above chance. Error bars represent 95% confidence intervals. Dots represent single subject data.
C) fMRI searchlight results for sensory (blue) and perceptual (green) braille letter representations (N = 15, height threshold $P < 0.001$, cluster-level FWE corrected $P < 0.05$, colored voxels indicate significance). Results for combined sensory and perceptual representations are in Supplementary Figure 1.

We started the analyses by determining the locations of sensory and perceptual braille letter representations in the visually deprived brain using fMRI. We focused our investigation on two sets of cortical regions based on previous literature: tactile processing areas and sighted reading areas (Fig. 2a). Given the tactile nature of braille, we expected braille letters to be represented in the tactile processing stream encompassing somatosensory cortices (S1 and S2), intra-parietal cortex (IPS), and insula [20]. Given that visual reading information is processed in a ventral processing stream [21–23], and

136 that braille reading has been observed to elicit activations along those nodes [5–9], we investigated the
137 sighted processing stream ranging from early visual cortex (EVC) [24] over V4 [25] and the lateral occipital
138 complex (LOC) [26] to the letter form area (LFA) [27] and visual word form area (VWFA) [25,26,28,29].
139
140 We hypothesized (H1) that sensory braille letter information is represented in tactile processing areas
141 (H1.1), while perceptual braille letter representations are located in sighted reading areas (H1.2). To
142 test H1, we conducted within-hand and across-hand classification of braille letters in the above-
143 mentioned areas for tactile processing and sighted reading (Fig. 2a) in a region-of-interest (ROI)
144 analysis.
145
146 We found that within-hand classification of braille letters was significantly above chance in regions
147 associated with both tactile processing (S1, S2, aIPS, pIPS, insula) and sighted reading (EVC, LOC,
148 LFA, VWFA) (Fig. 2b left; N = 15, one-tailed Wilcoxon signed-rank test, $P < 0.05$, FDR corrected).
149 As expected, this reveals both tactile and sighted reading areas as potential candidate regions housing
150 sensory and perceptual braille letter representations.
151
152 To pinpoint the loci of sensory representations we subtracted the results of the across-hand classification
153 from the within-hand classification. We found the difference to be significant in tactile processing areas
154 (S1, S2, aIPS and pIPS) and in LOC, but not elsewhere (Fig. 2b, middle). This confirms H1.1 in that
155 sensory braille letter representations are located in tactile areas.
156
To determine the location of perceptual representations, we assessed the results of across-hand
classification of braille letters. We found significant information (Fig. 2b, right) only in sighted reading
areas (EVC, LOC, VWFA), with the exception of the insula. This confirms H1.2 in that perceptual
braille letter representations emerge predominantly in sighted reading areas. The surprising finding of
the insula can possibly be explained by the insula's heterogeneous functions [30] beyond tactile
processing.
163
164 Notably, LOC is the only region that contained both sensory and perceptual braille letter
165 representations. This suggests a "hinge" function in the transformation from sensory to perceptual
166 braille letter representations.
167
168 To ascertain whether other areas beyond our hypothesized ROIs contained braille letter representations,
169 we conducted a spatially unbiased fMRI searchlight classification analysis. The results confirmed that
170 braille letter representations are located in the assessed ROIs without revealing any additional regions
171 (Fig. 2c, N = 15, height threshold $P < 0.001$, cluster-level FWE corrected $P < 0.05$).
172
173 Together, our fMRI results revealed sensory braille letter representations in tactile processing areas and
174 perceptual braille letter representations in sighted reading areas, with LOC serving as a "hinge" region
175 between them.
176

 3.2 Temporal dynamics of braille letter representations

Figure 3. Temporal dynamics of braille letter representations.
A) EEG results for sensory and perceptual (black), sensory (blue) and perceptual (green) braille letter representations in time. Shaded areas around curves indicate standard error of the mean. Significance onsets and their 95% confidence intervals are indicated by dots and horizontal lines below curves (color-coded as curves, N = 11, 1,000 bootstraps, one-tailed Wilcoxon signed-rank test, P < 0.05, FDR corrected).
B) EEG searchlight results for sensory and perceptual (top), sensory (middle) and perceptual (bottom) braille letter representations in EEG channel space (sampled down to 10ms resolution) in 100ms intervals. Significant electrodes are marked with black circles (N = 11, one-tailed Wilcoxon signed-rank test, P < 0.05, FDR corrected across electrodes and time points).

We next determined the temporal dynamics with which braille letter representations emerge using EEG.

We hypothesized (H2) that sensory braille letter representations emerge in time before perceptual braille letter representations, analogous to the sequential processing of sensory representations before perceptual representations in the visual [15,31–33] and auditory [34] domain.

To test H2, we conducted time-resolved within- and across-hand classification on EEG data. We determined the time point at which representations emerge by finding the first time point with respect to the onset of the braille stimulation where the classification effects are significant (fifty consecutive significant time point criteria, 95% confidence intervals reported in brackets).

The EEG classification analyses revealed significant and reliable results for both within- and across-hand classification of braille letters, as well as their difference (Fig. 3a; N = 11, 1,000 bootstraps, one-tailed Wilcoxon signed-rank test, P < 0.05, FDR corrected). We found that within-hand classification became significant at 62ms (29-111ms) (Fig. 3a, blue curve). To isolate sensory representations, we subtracted the results of the across-hand classification from the within-hand classification. This difference became significant at 77ms (45-138ms) (Fig. 3a, black curve). In contrast, the across-hand classification, indicating perceptual representations, became significant later at 184ms (127-230ms) (Fig. 3a, green curve).

Importantly, the temporal dynamics of sensory and perceptual representations differed significantly. Compared to sensory representations, the significance onset of perceptual representations was delayed by 107ms (21-167ms) (N = 11, 1,000 bootstraps, one-tailed bootstrap test against zero, P= 0.012).

To approximate the sources of the temporal signals, we complemented the EEG classification analysis
with a searchlight classification analysis in EEG sensor space. In the time window of highest
decodability (~200-300ms), sensory braille letter information was decodable from widespread
electrodes across the scalp. In contrast, perceptual braille letter information was decodable later and
from overall fewer electrodes which were located over right frontal, central, left parietal and left
temporal areas (Figure 3b; N = 11, one-tailed Wilcoxon signed-rank test, P < 0.05, FDR corrected across
electrodes and time points). These additional results reinforce that sensory braille letter information is
represented in more widespread brain areas than perceptual braille letter information, corroborating our
ROI classification results.

In sum, our EEG results characterized the temporal dynamics with which braille letter representations
emerge as a temporal sequence of sensory before perceptual representations.

3.3 Relating representations of braille letters to behavior

227
Figure 4. Representational similarity of braille letters in neural and behavioral measures.
A) For fMRI (top) and EEG (bottom), we formed representational dissimilarity matrices (RDMs) from pairwise
decoding accuracies. In the behavioral experiment (middle), we presented two braille letters (e.g., B and V) on
two braille cells. For all combinations, we asked participants to read both braille letters with the same hand (e.g.,
right) and verbally rate their similarity on a scale from 1-7. We formed the behavioral RDM from those perceptual
similarity judgements for every letter pair. We then correlated the behavioral RDM (averaged over participants)
with neural RDMs (subject specific) using Spearman's R.
B) Results of RSA relating fMRI and behavior in ROIs that showed significant sensory (left) and perceptual (right)
braille letter representations in fMRI (see. Fig. 2) (N = 15, two-tailed Wilcoxon signed-rank test, P < 0.05, FDR
corrected). Stars below bars indicate significance above chance. Error bars represent 95% confidence intervals.
Dots represent single subject data.
C) Results of RSA relating EEG and behavior for sensory (blue) and perceptual (green) representations. Shaded
areas around curves indicate standard error of the mean. Significance onsets and their 95% confidence intervals
are indicated by dots and lines below curves (N = 11, 1,000 bootstraps, one-tailed Wilcoxon signed-rank test, P <
0.05, FDR corrected, color-coded as result curves).

The ultimate goal of perception is to provide an organism with representations enabling adaptive
behavior. The analyses across space and time described above identified potential candidates for such
braille letter representations in the sense that the representations distinguished between braille letters.
However, not all of these representations have to be used by the brain to guide behavior; some of these
representations might be epiphenomenal and only available to the experimenter [35–37].

Therefore, we tested the hypothesis (H3) that the sensory and perceptual braille letter representations
identified in space (H1) and time (H2) are in a suitable format to be behaviorally relevant. We used
perceived similarity as a proxy for behavioral relevance. The idea is that if two stimuli are perceived to
be similar, they will also elicit similar actions [16–19]. For this, we acquired perceived similarity ratings

from blind participants (N=19) in a separate behavioral experiment (Fig. 4a middle) in which
participants verbally rated the similarity of each pair of braille letters from the stimulus set on a scale.
256
To relate perceived similarity ratings to neural representations, we used representational similarity
analysis (RSA) [38]. RSA relates different measurement spaces (such as EEG, fMRI and behavior) by
abstracting them into a common representational similarity space. We sorted behavioral similarity
ratings into representational dissimilarity matrices (RDMs) indexed in rows and columns by the 8
braille letters used in experimental conditions (Fig. 4a). For neural data, we sorted decoding accuracies
from the previous analyses as a dissimilarity measure [32,39]. Thus, by arranging the classification results
of the EEG and fMRI results we obtained EEG RDMs for every time point and fMRI RDMs for each
ROI (for the within-hand and the across-hand analyses separately). To finally relate behavioral and
neural data in the common RDM space, we correlated the behavioral RDM with each fMRI ROI RDM
and each EEG time point RDM.
267
Considering braille letter representations in space through fMRI (Fig. 4b, N = 15, one-tailed Wilcoxon
signed-rank test, $P < 0.05$, FDR corrected), we found that previously identified perceptual
representations (i.e., identified by the across-hand analysis) in EVC, LOC, VWFA, and insula showed
significant correlations with behavior. In contrast, sensory representations (i.e., identified by the
differences between within and across-hand analyses) in S1, S2, aIPS, pIPS and LOC were not
significantly correlated with behavior. This indicates that perceptual braille letter representations in
sighted reading areas are suitably formatted to guide behavior.
275
Considering braille letter representations in time through EEG (Fig. 4c; N = 11, 1,000 bootstraps, one-
tailed Wilcoxon signed-rank test, $P < 0.05$, FDR corrected), we found significant relationships with
behavior for both sensory and perceptual representations. The temporal dynamics mirrored those of the
EEG classification analysis (Fig. 3a), in that the results related to sensory representations emerged
earlier at 220ms (167- 567ms) than the results related to perceptual representations at 466ms (249-
877ms). The onset latency differences of 240ms (-81-636ms) (N = 11, 1,000 bootstrap, one-tailed
bootstrap test against zero, $P = 0.046$) was significant. This indicates that both earlier sensory
representations and later perceptual representations of braille letters are suitable formatted to guide
behavior.
285
286 In sum, our RSA results highlighted that perceptual representations in sighted reading areas, as well as
287 initial sensory and later perceptual representations in time, are suitably formatted to guide behavior.
288

4. Discussion

We assessed experience-based brain plasticity at its boundaries by investigating the nature of braille
information processing in the visually deprived brain. For this, we assessed the transformation of
sensory to perceptual braille letter representations in blind participants. Our experimental strategy
combining fMRI, EEG, and behavioral assessment yielded three key findings about spatial distribution,
temporal emergence and behavioral relevance. First, concerning the spatial distribution of braille letter
representations, we found that sensory braille letter representations are located in tactile processing
areas while perceptual braille letter representations are located in sighted reading areas. Second,
concerning the temporal emergence of braille letter representations, we found that sensory braille letter
representations emerge before perceptual braille letter representations. Third, concerning the behavioral
relevance of representations, we found that perceptual representations identified in sighted reading
areas, as well as sensory and perceptual representations identified in time, are suitably formatted to
guide behavior.
302

4.1 The topography of sensory and perceptual braille letter representations

304 Previous research has identified the regions activated during braille reading in high detail [5–11]. However,
305 activation in a brain region alone does not indicate its functional role or the kind of information it

306 represents [40]. Here, we characterize the information represented in a region by distinguishing between
307 sensory and perceptual representations of single braille letters. Our findings extend our understanding
308 of the cortical regions processing braille letters in the visually deprived brain in five ways.
309
First, we clarified the role of EVC activations in braille reading [5–11] by showing that EVC harbors
representations of single braille letters. More specifically, our finding that EVC represents perceptual
rather than sensory braille letter information indicates that EVC representations are formatted at a higher
perceptual level rather than a tactile input level. Previous studies also found that EVC of blind
participants processes other higher-level information such as natural sounds [41,42] and language [43–52]. A
parsimonious view is that EVC in the visually deprived brain engages in higher-level computations
shared across domains, rather than performing multiple distinct lower-level sensory computations.
Importantly, higher-level computations are not limited to the EVC in visually deprived brains. Natural
sound representations [41] and language activations [53] are also located in EVC of sighted participants.
This suggests that EVC, in general, has the capacity to process higher-level information [54]. Thus, EVC
in the visually deprived brain might not be undergoing fundamental changes in brain organization [53].
This promotes a view of brain plasticity in which the cortex is capable of dynamic adjustments within
pre-existing computational capacity limits [4,53–55].
323
324 Second, we found that VWFA contains perceptual braille letter representations. By clarifying the
325 representational format of language representations in VWFA, our results support previous findings of
326 the VWFA being functionally selective for letter and word stimuli in the visually deprived brain [8,56,57].
327
328 Third, LOC represented hand-dependent and -independent braille letter information, suggesting a
329 "hinge" function between sensory and perceptual braille letter representations. We stipulate that shape
330 serves as an intermediate level representational format in between lower-level properties such as
331 specific location of tactile stimulation or dot number in braille letters and higher-level perceptual letter
332 features [58].
333
334 Fourth, the finding of letter representations of tactile origin in both VWFA and LOC indicate that the
335 functional organization of both regions is multimodal, contributing to the debate on how experience
336 from vision or other sensory modalities shapes representations along the ventral stream [58,59].
337
338 Fifth, we observed that the somatosensory cortices and intra-parietal sulci represent hand-dependent
339 but not hand-independent braille letter representations. This is consistent with previous studies reporting
340 that the primary somatosensory cortex represents the location of tactile stimulation [60], but not the
341 identity of braille words [57]. Taken together, these findings suggest that these tactile processing areas
342 represent sensory rather than higher-level features of tactile inputs in visually deprived brains.
343
The involvement of the insula in processing braille letter information is more difficult to interpret. Based
on previous studies in the sighted brain, the insula plays a role in tactile memory [61–63] and multisensory
integration [64–67]. Both aspects could have contributed to our findings as braille letters are retrievable
from long-term memory but are also inherently nameable and linked to auditory experiences. A future
study could disambiguate the contributions of tactile memory and multisensory integration by
presenting meaningless dot arrays, that are either unnamable or paired with invented names. Insular
representations of trained, unnamable stimuli but not novel, unnamable stimuli would align with
memory requirements. Insular representations of trained, namable stimuli but not trained, unnamable
stimuli would favor audio-tactile integration.
353
4.2 Sensory representations emerge before perceptual representations
Using time-resolved multivariate analysis of EEG data [15,31,33], we showed that hand-dependent, sensory
braille letter representations emerge in time before hand-independent, perceptual representations. Such
sequential multi-step processing in time is a general principle of information processing in the human

brain, also known in the visual [15,31,32] and auditory [34] domain. Together, these findings suggest that the
human brain, even in extreme instances of species-atypical cortical plasticity, honors this principle.
360
While braille letter reading follows the same temporal processing sequence as its visual counterpart, it
operates on a different time scale. Our results indicate that braille letter classification peaks substantially
later in time (~200ms for hand-dependent and ~390ms for hand-independent representations) than
previously reported classification of visually presented words, letters, objects, and object categories
(e.g., ~125ms for location-dependent and ~175ms for location-independent representations) [15,31,32,68,69].
This discrepancy raises the question which factors could limit the speed of processing braille letters.
Importantly, this delay is not a consequence of slower cortical processing in the tactile domain [70]. We
find that tactile information reaches the cortex fast: we can classify which hand was stimulated as early
as 35ms after stimulation onset (Supplementary Figure 2). Thus, the delay relates directly to the
identification of braille letters.
371
A compelling explanation for the temporal processing properties of braille letter information are the
underlying reading mechanics. Braille reading is slower than print reading [71,72] even if participants are
fluent braille readers. This slowing is specific to braille reading and does not translate to other types of
information intake in the visually deprived brain, e.g., auditory information. Blind participants have
higher listening rates [73] and better auditory discrimination skills [74] than sighted participants, indicating
more efficient auditory processing [75,76]. Together, this result pattern suggests that the temporal dynamics
with which braille letter representations emerge are limited by the particular efficiency of the braille
letter system, rather than the capacity of the brain. To test this idea, future studies could compare the
temporal dynamics of braille letter and haptic object representations: a temporal processing delay
specific to braille letters would support the hypothesis.
382

4.3 Representations identified in space and time guide behavior

Our results clarified that perceptual rather than sensory braille letter representations identified in space
are suitably formatted to guide behavior. However, we acknowledge that this finding is task-dependent.
Arguably, general similarity ratings of braille letters depend more on intake-independent (e.g., dot
arrangements or linguistic similarities such as pronunciation) than intake-dependent features
(similarities in stimulation location on finger). Future behavioral assessments could ask participants to
assess similarity separately based on only stimulation location or linguistic features. We would predict
that similarity ratings based on stimulation location are related to sensory representations while
similarity ratings based on linguistic features are related to perceptual representations of braille letters.
392
Concerning temporal dynamics, our results reveal that sensory representations of braille letters are
relevant for behavior earlier than perceptual ones. Interestingly, the similarity between perceptual
braille letter representations and behavioral similarity ratings emerges in a time window in which
transcranial magnetic stimulation (TMS) over the VWFA affects braille letter reading in sighted braille
readers [77]. This implies that around 320-420ms after the onset of reading braille, visually deprived and
sighted brains utilize braille letter representations for performing tasks such as letter identification.
Applying a comparable TMS protocol not only to sighted but also non-sighted braille readers would
elucidate whether this time window of behavioral relevance can be generalized to braille reading,
independent of visual experience.
402

4.4 Conclusions

Our investigation of experience-based plasticity at its boundaries due to the loss of the visual modality
reveals a nuanced picture of its potential and limits. On the one hand, our findings emphasize how
plastic the brain is by showing that regions typically processing visual information adapt to represent
perceptual braille letter information. On the other hand, our findings illustrate inherent limits of brain
plasticity. Brain areas represent information from atypical inputs within the boundaries of their pre-

existing computational capacity and the progression from sensory to perceptual transformations adheres
to a common temporal scheme and functional role.

5. Methods

5.1 Participants

We conducted three separate experiments with partially overlapping participants: an fMRI, an EEG,
and a behavioral experiment. All experiments were approved by the ethics committee of the Department
of Education and Psychology of the Freie Universität Berlin and were conducted in accordance with
the Declaration of Helsinki. 16 participants completed the fMRI experiment. One person was excluded
due to technical problems during the recording, leaving a total of 15 participants in the fMRI experiment
(mean age 39 years, *SD*=10, 9 female). 11 participants participated in the EEG experiment (*N*=11, mean
age 44 years, *SD*=10, 8 female). The participant pools of the EEG and fMRI experiments overlapped
by five participants. Out of a total of 21 participants, 19 participants (excluding one fMRI and one EEG
participant) completed an additional behavioral task in which they rated the perceived similarity of
braille letter pairs. All participants were blind since birth or early childhood (≤ 3 years, for details see
Supplementary Table 1). All participants provided informed consent prior to the studies and received a
monetary reward for their participation.

5.2 Experimental stimuli and design

In all experiments, we presented braille letters (B, C, D, L, M, N, V, Z; Figure 2a) to the left and right
index fingers of participants using piezo-actuated refreshable braille cells (https://metec-
ag.de/index.php) with two modules of 8 pins each. We only used the top 6 pins from each module to
present letters from the braille alphabet. The modules were taped to the clothes of a participant for the
fMRI experiment and on the table for the EEG and behavioral experiment. This way, participants could
read in a comfortable position with their index fingers resting on the braille cells to avoid motion
confounds. We instructed participants to read letters regardless of whether the pins stimulated their right
or left index finger. We presented all eight letters to both hands, resulting in 16 experimental conditions
(8 letters × 2 hands). In addition, two braille letters (E, O) were included as catch stimuli and participants
were instructed to respond to them by pressing a button (fMRI) or pedal (EEG) with their foot. Catch
trials were excluded from further analysis due to confounding motor and sensory signals.

5.3 Experimental procedures

5.3.1 fMRI experiment

The fMRI experiment consisted of two sessions. 15 participants completed the first fMRI session,
during which we recorded a structural image (~4 min.), a localizer run (7 min) and 10 runs of the main
experiment (56 minutes). The total duration of the first session was 67 minutes excluding breaks. Eight
of these 15 subjects completed a second fMRI recording session, in which we recorded an additional
15 runs of the main experiment (85 minutes). We did not record any structural images or localizer runs
in the second session, resulting in a total duration of 85 minutes excluding breaks.

5.3.1.1. fMRI main experiment

During the fMRI main experiment, we presented participants with letters on braille cells and asked them
to respond to occasionally appearing catch letters. We presented letters for 500ms, with a 2500ms inter-
stimulus-interval (ISI; see Figure 2b top). Each regular trial - belonging to one of the 16 experimental
conditions - was repeated 5 times per run (run duration: 337 s) in random order. Regular trials were
interspersed every ~20 trials with a catch trial, such that a catch trial occurred about once per minute.
In addition, every 3rd to 5th trial (equally probable) was a null trial where no stimulation was given. In
total, one run consisted of 80 regular trials, 5 catch trials and 22 null trials, amounting to a total of 107
trials per run.
454

To ensure that participants were able to read letters with both hands and understood the task instructions,
participants first completed an experimental run outside the scanner.

5.3.1.2 fMRI localizer experiment

To define regions-of-interest (ROIs), we performed a separate localizer experiment prior to the main
fMRI experiment with tactile stimuli in four experimental conditions: braille letters read with the left
hand, braille letters read with the right hand, fake letters read with the left hand and fake letters read
with the right hand. The letters presented in the braille conditions were 16 letters from the alphabet
excluding the letters used in the main experiment. The stimuli in the fake letter conditions were 16
tactile stimuli that were each composed of 8 dots, deviating from the standard 6-dot configuration in
the braille alphabet.
465
The localizer experiment consisted of a single run lasting 432s, comprising 5 blocks of presentation of
braille letters left, braille letters right, fake letters left, fake letters right and blank blocks as baseline.
Each stimulation block was 14.4s long, consisting of 18 different letters presentations (500ms on,
300ms off) including two one-back repetitions that participants were instructed to respond to by
pressing a button with their foot. We presented stimulation blocks in random order and regularly
interspersed them with blank blocks.

5.3.2 EEG experiment

The EEG experiment consisted of two sessions. 11 participants completed the first session and 8 of
these participants completed a second session. The total duration of each session was 59 minutes
excluding breaks.
476
The experimental setup was similar to that for fMRI but adapted to the specifics of EEG. We presented
braille letters for 500ms with a 500ms ISI on regular trials. In catch trials, the letters were presented for
500ms with a 1100ms ISI to avoid contamination of movement on subsequent trials (see Figure 2b
bottom). Each of the 16 experimental conditions was presented 170 times per session. Regular trials
were interspersed every 5^{th} to 7^{th} trial (equally probable) with a catch trial. In total, one EEG recording
session consisted of 2720 regular trials and 541 catch trials, amounting to a total of 3261 trials. 2
participants completed additional trials due to technical problems leading to a total of 190 and 180
repetitions per stimulus, accordingly.
485
Prior to the experiment, participants completed a short screening task during which each letter of the
alphabet was presented for 500ms to each hand in random order. Participants were asked to verbally
report the letter they had perceived to assess their reading capabilities with both hands using the same
presentation time as in the experiment. The average performance for the left hand was 89% correct (SD
= 10) and for the right hand it was 88% correct (SD = 13).

5.3.3 Behavioral letter similarity ratings

In a separate behavioral experiment, participants judged the perceived similarity of the braille letters
used in the neuroimaging experiments. For this task, participants sat at a desk and were presented with
two braille cells next to each other. Each pair of letters was presented once, and participants compared
them with the same finger. The rating was without time constraints, meaning participants decided when
they rated the stimuli. Participants were asked to verbally rate the similarity of each pair of braille letters
on a scale from 1 = very similar to 7 = very different and the experimenter noted down their responses.

5.4 fMRI data acquisition, preprocessing and preparation

5.4.1 fMRI acquisition

We acquired MRI data on a 3-T Siemens Tim Trio scanner with a 12-channel head coil. We obtained
structural images using a T1-weighted sequence (magnetization-prepared rapid gradient-echo, 1 mm^3
voxel size). For the main experiment and the localizer run, we obtained functional images covering the

entire brain using a T2*-weighted gradient-echo planar sequence (TR=2s, TE=30ms, flip angle=70°, 3
mm^3 voxel size, 37 slices, FOV=192mm, matrix size=64×64, interleaved acquisition).

5.4.2 fMRI preprocessing

For fMRI preprocessing, we used tools from FMRIB's Software Library (FSL,
www.fmrib.ox.ac.uk/fsl). We excluded non-brain tissue from analysis using the Brain Extraction Tool
(BET)[78] and motion corrected the data using MCFLIRT [79]. We did not apply high or low-pass temporal
filters. We spatially smoothed fMRI localizer data with an 8mm FWHM Gaussian kernel. We registered
functional images to the high-resolution structural scans and to the MNI standard template using FLIRT
[80]. We carried out all further fMRI analyses in MATLAB R2021a (www.mathworks.com).

5.4.3 Univariate fMRI analysis

For all univariate fMRI analyses, we used SPM12 (http://www.fil.ion. ucl.ac.uk/spm/). For the main
experiment, we modelled the fMRI responses to the 16 experimental conditions for each run using a
general linear model (GLM). The onsets and durations of each image presentation entered the GLM as
regressors and were convolved with a hemodynamic response function (hrf). Six movement parameters
(pitch, yaw, roll, x-, y-, z-translation) entered the GLM as nuisance regressors. For each of the 16
conditions we converted GLM parameter estimates into *t*-values by contrasting each parameter estimate
against the implicit baseline. This resulted in 16 condition specific *t*-value maps per run and participant.
520
For the localizer experiment, we modelled the fMRI response to the 5 experimental conditions entering
block onsets and durations as regressors of interest and movement parameters as nuisance regressors
before convolving with the hrf. From the resulting three parameter estimates we generated two
contrasts. The first contrast served to localize activations in primary (S1) and secondary (S2)
somatosensory cortex and was defined as letters & fake letters > baseline. The second contrast served
to localize activations in early visual cortex (EVC), V4, lateral occipital complex (LOC), letter form
area (LFA), visual word form area (VWFA), anterior intra-parietal sulcus (aIPS), posterior intra-parietal
sulcus (pIPS) and insula and was defined as letters > fake letters. In sum, this resulted in two *t*-value
maps for the localizer run per participant.

5.4.4 Definition of regions-of-interest

To identify regions along the sighted reading and tactile processing pathway, we defined regions of
interest (ROIs) in a two-step procedure. We first constrained ROIs by anatomical masks using brain
atlases, in each case combining regions across both hemispheres. We included 5 ROIs from the sighted
reading pathway: EVC (merging the anatomical masks of V1, V2 and V3), V4, LOC, LFA and the
VWFA. We also included 5 ROIs from the tactile processing pathway: S1, S2, aIPS (merging the
anatomical masks of IPS3, IPS4, IPS5), pIPS (merging the anatomical masks of IPS0, IPS1 and IPS2)
and the insula. For EVC, V4, LOC, aIPS and pIPS, we used masks from the probabilistic Wang atlas
[81]. For LFA, we defined the mask using the MarsBaR Toolbox (https://marsbar-toolbox.github.io/) with
a 10 mm radius around the center voxel at MNI coordinates X=-40; Y=-78 and Z=-18 [27]. We also
defined the VWFA mask using the MarsBaR Toolbox with a 10 mm radius around the center voxel at
MNI coordinates X=-44; Y=-57 and Z=-13 [25] and converted from Talairach to MNI space using the
MNI<->Talairach Tool (https://bioimagesuiteweb.github.io/bisweb-manual/tools/mni2tal.html). We
created the mask for S1 by merging the sub-masks of BA1, BA2 and BA3 from the WFU PickAtlas
(https://www.nitrc.org/projects/wfu_pickatlas/) and the mask for S2 by merging the sub-masks
operculum 1-4 from the Anatomy Toolbox [82]. Lastly, we extracted the mask for the insula from the
WFU PickAtlas. The smallest mask included 321 voxels. Therefore, in a second step, we selected the
321 most activated voxels of the participant-specific localizer results within each of the masks, using
the letters & fake letters > baseline contrast for S1 and S2 and the letters > fake letters for the remaining
ROIs. This yielded participant-specific definitions for all ROIs.

5.5 EEG data acquisition and preprocessing

We recorded EEG data using an EASYCAP 64-channel system and a Brainvision actiCHamp amplifier at a sampling rate of 1,000 Hz. The electrodes were placed according to the standard 10-10 system. The data was filtered online between 0.03 and 100 Hz and re-referenced online to FCz.

We preprocessed data offline using the EEGLAB toolbox version 14 [83]. We incorporated a low-pass filter with a cut-off at 50 Hz and epoched trials between -100 ms and 999 ms with respect to stimulus onset, resulting in 1,100 1ms data points per epoch. We baseline-corrected the epochs by subtracting the mean of the 100ms prestimulus time window from the epoch. We re-referenced the data offline to the average reference. To clean the data from artifacts such as eye blinks, eye movements and muscular contractions, we used independent component analysis as implemented in the EEGLAB toolbox. We used SASICA [84] to guide the visual inspection of components for removal. We identified components related to horizontal eye movements using two lateral frontal electrodes (F7-F8). During five recordings (1 participant first session, 4 participants second session), additional external electrodes were available that allowed for the direct recording of the horizontal electro-oculogram to identify and remove components related to horizontal eye movements. For blink artifact detection based on the vertical electro-oculogram, we used two frontal electrodes (Fp1 and Fp2). As a final step, we applied multivariate noise normalization to improve the signal-to-noise ratio (SNR) and reliability of the data [39], resulting in subject-specific trial-based time courses of electrode activity.

5.6 Braille letter classification from brain measurements

To determine the amount of information about braille letter identity present in brain measurements, we used a multivariate classification scheme [12–15]. We conducted subject-specific braille letter classification in two ways. First, we classified between letter pairs presented to one reading hand, i.e., we trained and tested a classifier on brain data recorded during the presentation of braille stimuli to the same hand (either the right or the left hand). This yields a measure of hand-dependent braille letter information in neural measurements. We refer to this analysis as within-hand classification. Second, we classified between letter pairs presented to different hands in that we trained a classifier on brain data recorded during the presentation of stimuli to one hand (e.g., right), and tested it on data related to the other hand (e.g., left). This yields a measure of hand-independent braille letter information in neural measurements. We refer to this analysis as across-hand classification.

All classification analyses were carried out in MATLAB R2021a (www.mathworks.com) and relied on binary c-support vector classification (C-SVC) with a linear kernel as implemented in the libsvm toolbox [85] (https://www.csie.ntu.edu.tw/cjlin/libsvm). Furthermore, all analyses were conducted in a participant specific manner. The next section describes the multivariate fMRI and EEG analyses in more detail.

5.6.1 Spatially resolved multivariate fMRI analysis

We conducted both a ROI-based and a spatially unbiased volumetric searchlight procedure [86]. For each ROI included in the ROI-based analysis, we extracted and arranged *t*-values into pattern vectors for each of the 16 conditions and experimental runs. If participants completed only one session, the analysis was conducted on 10 runs. If participants completed both sessions, the 10 runs from session 1 and 15 runs from session 2 were pooled and the analysis was conducted across 25 runs. To increase the SNR, we randomly assigned run-wise pattern vectors into bins and averaged them into pseudo-runs. For participants with one session, the bin size was 2 runs, resulting in 5 pseudo-runs. If participants completed 2 sessions and thus had 25 runs, the bin size was 5 runs resulting in 5 pseudo-runs. Thus, in both cases, each participant ended up with five pseudo-run pattern vectors that entered the classification analysis. We then performed 5-fold leave-one-pseudo-run-out-cross validation, training on 4 and testing on one pseudo-trial per classification iteration.

We will first describe the classification procedure for braille letters within-hand and then for the classification of braille letters across-hand.

5.6.2 fMRI ROI-based classification of braille Letters within-hand

For the classification of braille letters within-hand, we assigned four pseudo-trials corresponding to the data from two braille letters of the same hand (e.g., right) to the training set. We then tested the SVM on the remaining, fifth pseudo-trial corresponding to data from the same two braille letters of the same hand (e.g., right) as in the training set but using held-out data for the testing set. This yielded percent classification accuracy (50% chance level) as output. Equivalent SVM training and testing was repeated for all combinations of letter pairs within each hand.

With 8 letters that were all classified pairwise once per hand, this resulted in 28 pairwise classification accuracies per hand. We averaged accuracies across condition pairs and hands, yielding a measure of hand-dependent braille letter information for each ROI and participant separately

5.6.3 fMRI ROI-based classification of braille Letters across-hand

The classification procedure of braille letters across reading hands was identical to the classification procedure within-hand with the important difference that the training data always came from one hand (e.g., right) and the testing data from the other hand (e.g., left).

With 8 letters that were all classified pairwise once across two hands, this resulted again in 28 pairwise classification accuracies across-hand per training-testing direction (i.e., train left, test right and vice versa). We averaged accuracies across condition pairs and training-testing directions, yielding a measure of hand-independent braille letter information for each ROI and participant separately.

5.6.4 fMRI searchlight classification of braille Letters

The searchlight procedure was conceptually equivalent to the ROI-based analysis. For each voxel v_i in the 3D t-value maps, we defined a sphere with a radius of 4 voxels centered around voxel v_i. For each condition and run, we extracted and arranged the t-values for each voxel of the sphere into pattern vectors. Classification of braille letters across-hand proceeded as described above. This resulted in one average classification accuracy for voxel v_i. Iterated across all voxels this yielded a 3D volume of classification accuracies across the brain for each participant separately.

5.6.5 Time-resolved classification of braille letters within-hand from EEG data

To determine the timing with which braille letter information emerges in the brain, we conducted time-resolved EEG classification [15,33]. This procedure was conceptually equivalent to the fMRI braille letter classification in that it classified letter pairs either within or across-hand and was conducted separately for each participant.

For each time point of the epoched EEG data, we extracted 63 EEG channel activations and arranged them into pattern vectors for each of the 16 conditions. Participants who completed one session had 170 trials per condition and participants who completed two sessions had 340 trials per condition. To increase the SNR, we randomly assigned the trials into bins and averaged them into new pseudo-trials. For participants with one session, the bin size was 34 trials, resulting in 5 pseudo-trials. If participants completed 2 sessions and thus had 340 trials, the bin size was 68 trials resulting in 5 pseudo-trials. In both cases, each participant ended up with five pseudo-run pattern vectors that entered the classification analysis. We then performed 5-fold leave-one-pseudo-run-out-cross validation, training on 4 and testing on one pseudo-trial per classification iteration. This procedure was repeated 100 times with random assignment of trials to pseudo-trials, and across all combinations of letter pairs and hands. We averaged results across condition pairs, folds, iterations and hands, yielding a decoding accuracy time course reflecting how much hand-dependent braille letter information was present at each time point in each participant.

 5.6.6 Time-resolved classification of braille letters across-hand from EEG data

The classification procedure for braille letters across-hand was identical to the classification of braille
letters within-hand with the crucial difference that training and testing data always came from separate
hands and results were averaged across condition pairs, folds, iterations and training-testing directions.
Averaging results yielded a decoding accuracy time course reflecting how much hand-independent
braille letter information was present at each time point in each participant.

5.6.7 Time-resolved EEG searchlight in sensor space

We conducted an EEG searchlight analysis resolved in time and sensor space (i.e., across 63 EEG
channels) to gain insights into which EEG channels contributed to the results of the time-resolved
analysis described above. For the EEG searchlight, we conducted the time-resolved EEG classification
as described above with the following difference: For each EEG channel c_i, we conducted the
classification procedure on the four closest channels surrounding c. The classification accuracy was
stored at the position of c. After iterating across all channels and down-sampling the time points to a
10ms resolution, this yielded a classification accuracy map across all channels and time points in 10ms
steps for each participant.

5.7 Representational similarity analysis of brain data and behavioral letter similarity ratings

To determine the subset of neural braille letter representations identified that is relevant for behavior [16–
19], we compared perceptual letter similarity ratings to braille letter representations identified from EEG
and fMRI signals using representational similarity analysis (RSA) [38]. RSA characterizes the
representational space of a measurement space (e.g., fMRI or EEG data) with a representational
dissimilarity matrix (RDM). RDMs aggregate pairwise distances between responses to all experimental
conditions, thereby abstracting from the activity patterns of measurement units (e.g., fMRI voxels or
EEG channels) to between-condition dissimilarities. The rationale of the approach is that neural
measures and behavior are linked if their RDMs are similar.
671
672 We constructed RDMs for behavior, fMRI and EEG as follows.
673
674 For behavior, we arranged the perceptual similarity judgments averaged across participants (indicated
675 by participants on a scale from 1 = very similar to 7 = very different) into an RDM format. All RDMs
676 were averaged over both hands and had the dimensions 8 letters x 8 letters.
677
678 For both fMRI and EEG, we used the classification results from the conducted within-hand and across-
679 hand classifications as a measure of (dis-)similarity relations between braille letters. Classification
680 accuracies can be interpreted as a measure of dissimilarity because two conditions have a higher
681 classification accuracy when they are more dissimilar[32,39]. Thus, we assembled participant-specific
682 RDMs for each fMRI ROI and EEG time point in the time course from decoding accuracies.
683
In a final step we correlated (Spearman's *R*) the lower triangular part of the respective RDMs (without
the diagonal [84]). For fMRI, this resulted in one correlation value per participant and ROI. For EEG, this
analysis resulted in one correlation time course per participant.

5.8 Statistical testing

5.8.1 Wilcoxon signed-rank test

We performed non-parametric one-tailed Wilcoxon signed-rank tests to test for above-chance
classification accuracy for ROIs in the fMRI classification, for time points in the EEG classification,
for time points and channels in the EEG searchlight, and for ROI and time courses in the RSA. In each
case the null hypothesis was that the observed parameter (i.e., classification accuracy, correlation) came
from a distribution with a median of chance level performance (i.e., 50% for pairwise classification; 0
correlation). The resulting *P*-values were corrected for multiple comparisons using the false discovery

rate (FDR) [87] at 5% level if more than one test was conducted. This was done a) across ROIs in the ROI
classification and fMRI-behavior RSA , b) across time points in the EEG classification and EEG-
behavior RSA, and c) across time points and channels in the EEG searchlight.

5.8.2 Bootstrap tests

We used bootstrapping to compute 95% confidence intervals for onset latencies (the first 50 consecutive
significant timepoints after trial onset) of EEG time courses as well as for determining the significance
of onset latencies. In each case we sampled the participant pool 1,000 times with replacement calculated
the statistic of interest for each sample.
703
For the EEG onset latency differences, we bootstrapped the latency difference between the onsets of
the time courses of hand-dependent or hand-independent letter representations. This yielded an
empirical distribution that could be compared to zero. To determine whether onset latencies differences
in the EEG time courses were significantly different from zero, we computed the proportion of values
that were equal to or smaller than zero and corrected them for multiple comparisons using FDR at
P=0.05.

5.8.3 Other statistical tests

For the fMRI searchlight classification results, we applied a voxel-wise height threshold of P= 0.001.
The resulting P-values were corrected for multiple comparisons using the family-wise error (FWE) at
the 5% level.

5.9 Data availability

The raw fMRI and EEG data are available on OpenNeuro via https://openneuro.org/datasets/ds004956
and https://openneuro.org/datasets/ds004951/. The preprocessed fMRI, EEG and behavioral data as
well as the results of the ROI classification, time classification and RSA can be accessed on OSF via
https://osf.io/a64hp/.

5.10 Code availability

The code used in this study is available on Github via
https://github.com/marleenhaupt/BrailleLetterRepresentations/.
722

6. References

1. Bennett, L., Melchers, B. & Proppe, B. Curta: A General-purpose High-Performance
Computer at ZEDAT, Freie Universität Berlin. (2020).

2. Pascual-Leone, A., Amedi, A., Fregni, F. & Merabet, L. B. The plastic human brain cortex.
Annu. Rev. Neurosci. **28**, 377–401 (2005).

3. Ricciardi, E., Bottari, D., Ptito, M., Röder, B. & Pietrini, P. The sensory-deprived brain as a
unique tool to understand brain development and function. *Neurosci. Biobehav. Rev.* **108**, 78–
82 (2020).

4. Bedny, M. Evidence from Blindness for a Cognitively Pluripotent Cortex. *Trends Cogn. Sci.*
21, 637–648 (2017).

5. Büchel, C., Price, C. & Friston, K. A multimodal language region in the ventral visual
pathway. *Nature* **394**, 274–277 (1998).

6. Burton, H., Sinclair, R. J. & Agato, A. Recognition memory for Braille or spoken words: An
fMRI study in early blind. *Brain Res.* **1438**, 22–34 (2012).

7. Rączy, K. *et al.* Orthographic Priming in Braille Reading as Evidence for Task-specific
Reorganization in the Ventral Visual Cortex of the Congenitally Blind. *J. Cogn. Neurosci.* **31**,
1065–1078 (2019).

8.   Reich, L., Szwed, M., Cohen, L. & Amedi, A. A ventral visual stream reading center
independent of visual experience. *Curr. Biol.* **21**, 363–368 (2011).

9.   Sadato, N. Neural networks for Braille reading by the blind. *Brain* **121**, 1213–1229 (1998).

10.   Uhl, F., Franzen, P., Lindinger, G., Lang, W. & Deecke, L. On the functionality of the visually
deprived occipital cortex in early blind persons. *Neurosci. Lett.* **124**, 256–259 (1991).

11.   Sadato, N. *et al.* Activation of V1 by Braille reading in blind subjects. *Nature* **380**, 526–528
(1996).

12.   Cichy, R. M. *et al.* Probing principles of large-scale object representation: Category preference
and location encoding. *Hum. Brain Mapp.* **34**, 1636–1651 (2013).

13.   Cichy, R. M., Chen, Y. & Haynes, J. D. Encoding the identity and location of objects in
human LOC. *Neuroimage* **54**, 2297–2307 (2011).

14.   Carlson, T. A., Hogendoorn, H., Fonteijn, H. & Verstraten, F. A. Spatial coding and invariance
in object-selective cortex. *Cortex* **47**, 14–22 (2011).

15.   Isik, L., Meyers, E. M., Leibo, J. Z. & Poggio, T. The dynamics of invariant object recognition
in the human visual system. *J. Neurophysiol.* **111**, 91–102 (2014).

16.   Cichy, R. M., Kriegeskorte, N., Jozwik, K. M., van den Bosch, J. J. F. & Charest, I. The
spatiotemporal neural dynamics underlying perceived similarity for real-world objects.
Neuroimage **194**, 12–24 (2019).

17.   Bankson, B. B., Hebart, M. N., Groen, I. I. A. & Baker, C. I. The temporal evolution of
conceptual object representations revealed through models of behavior, semantics and deep
neural networks. *Neuroimage* **178**, 172–182 (2018).

18.   Mur, M. *et al.* Human object-similarity judgments reflect and transcend the primate-IT object
representation. *Front. Psychol.* **4**, 1–22 (2013).

19.   Charest, I. *et al.* Unique semantic space in the brain of each beholder predicts perceived
similarity. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 14565–14570 (2014).

20.   Dijkerman, H. C. & de Haan, E. H. F. Somatosensory processes subserving perception and
action. *Behav. Brain Sci.* **30**, 189–201 (2007).

21.   DiCarlo, J. J. & Cox, D. D. Untangling invariant object recognition. *Trends Cogn. Sci.* **11**,
333–341 (2007).

22.   DiCarlo, J. J., Zoccolan, D. & Rust, N. C. How does the brain solve visual object recognition?
Neuron **73**, 415–434 (2012).

23.   Goodale, M. A. & Milner, A. D. Separate visual pathways for perception and action. *Trends
Neurosci.* **15**, 20–25 (1992).

24.   Dehaene, S., Cohen, L., Sigman, M. & Vinckier, F. The neural code for written words: A
proposal. *Trends Cogn. Sci.* **9**, 335–341 (2005).

25.   Cohen, L. *et al.* The visual word form area: spatial and temporal characterization of an initial
stage of reading in normal subjects and posterior split-brain patients. *Brain* **123**, 291–307
(2000).

26.   Neudorf, J., Gould, L., Mickleborough, M. J. S., Ekstrand, C. & Borowsky, R. Unique, Shared,
and Dominant Brain Activation in Visual Word Form Area and Lateral Occipital Complex
during Reading and Picture Naming. *Neuroscience* **481**, 178–196 (2022).

27.   Thesen, T. *et al.* Sequential then interactive processing of letters and words in the left fusiform
gyrus. *Nat. Commun.* **3**, 1–8 (2012).

28.   McCandliss, B. D., Cohen, L. & Dehaene, S. The visual word form area: Expertise for reading
in the fusiform gyrus. *Trends Cogn. Sci.* **7**, 293–299 (2003).

29.   Dehaene, S. & Cohen, L. The unique role of the visual word form area in reading. *Trends*
Cogn. Sci. **15**, 254–262 (2011).

30.   Uddin, L. Q., Nomi, J. S., Hébert-Seropian, B., Ghaziri, J. & Boucher, O. Structure and
Function of the Human Insula. *J. Clin. Neurophysiol.* **34**, 300–306 (2017).

31.   Carlson, T., Tovar, D. A., Alink, A. & Kriegeskorte, N. Representational dynamics of object
vision: The first 1000 ms. *J. Vis.* **13**, 1–19 (2013).

32.   Cichy, R. M., Pantazis, D. & Oliva, A. Resolving human object recognition in space and time.
Nat. Neurosci. **17**, 455–462 (2014).

33.   Carlson, T. A., Hogendoorn, H., Kanai, R., Mesik, J. & Turret, J. High temporal resolution
decoding of object position and category. *J. Vis.* **11**, 1–17 (2011).

34.   Lowe, M. X. *et al.* Cochlea to categories: The spatiotemporal dynamics of semantic auditory
representations. *Cogn. Neuropsychol.* (2021) doi:10.1080/02643294.2022.2085085.

35.   de-Wit, L., Alexander, D., Ekroll, V. & Wagemans, J. Is neuroimaging measuring information
in the brain? *Psychon. Bull. Rev.* **23**, 1415–1428 (2016).

36.   Reddy, L. & Kanwisher, N. Category Selectivity in the Ventral Visual Pathway Confers
Robustness to Clutter and Diverted Attention. *Curr. Biol.* **17**, 2067–2072 (2007).

37.   Williams, M. A., Dang, S. & Kanwisher, N. G. Only some spatial patterns of fMRI response
are read out in task performance. *Nat. Neurosci.* **10**, 685–686 (2007).

38.   Kriegeskorte, N., Mur, M. & Bandettini, P. Representational similarity analysis - connecting
the branches of systems neuroscience. *Front. Syst. Neurosci.* **2**, 1–28 (2008).

39.   Guggenmos, M., Sterzer, P. & Cichy, R. M. Multivariate pattern analysis for MEG: A
comparison of dissimilarity measures. *Neuroimage* **173**, 434–447 (2018).

40.   Haynes, J. D. & Rees, G. Decoding mental states from brain activity in humans. *Nat. Rev.*
Neurosci. **7**, 523–534 (2006).

41.   Vetter, P. *et al.* Decoding Natural Sounds in Early "Visual" Cortex of Congenitally Blind
Individuals. *Curr. Biol.* **30**, 3039-3044.e2 (2020).

42.   Mattioni, S. *et al.* Categorical representation from sound and sight in the ventral occipito-
temporal cortex of sighted and blind. *Elife* **9**, 1–33 (2020).

43.   Burton, H., Snyder, A. Z., Diamond, J. B. & Raichle, M. E. Adaptive changes in early and late
blind: A fMRI study of verb generation to heard nouns. *J. Neurophysiol.* **88**, 3359–3371
(2002).

44.   Burton, H., Diamond, J. B. & McDermott, K. B. Dissociating cortical regions activated by
semantic and phonological tasks: A fMRI study in blind and sighted people. *J. Neurophysiol.*
90, 1965–1982 (2003).

45.   Röder, B., Stock, O., Bien, S., Neville, H. & Rösler, F. Speech processing activates visual
cortex in congenitally blind humans. *Eur. J. Neurosci.* **16**, 930–936 (2002).

46.   Bedny, M., Pascual-Leone, A., Dodell-Feder, D., Fedorenko, E. & Saxe, R. Language
processing in the occipital cortex of congenitally blind adults. *Proc. Natl. Acad. Sci. U. S. A.*
108, 4429–4434 (2011).

47.   Lane, C., Kanjlia, S., Omaki, A. & Bedny, M. "Visual" cortex of congenitally blind adults
responds to syntactic movement. *J. Neurosci.* **35**, 12859–12868 (2015).

48.   Dietrich, S., Hertrich, I. & Ackermann, H. Ultra-fast speech comprehension in blind subjects
engages primary visual cortex, fusiform gyrus, and pulvinar - a functional magnetic resonance
imaging (fMRI) study. *BMC Neurosci.* **14**, (2013).

49.   Watkins, K. E., Coullon, G. S. L. & Bridge, H. Language and nonverbal auditory processing in

the occipital cortex of individuals who are congenitally blind due to anophthalmia.
Neuropsychologia **173**, 108304 (2022).

50.   Abboud, S., Engemann, D. A. & Cohen, L. Semantic coding in the occipital cortex of early
833   blind individuals. *bioRxiv* 539437 (2019) doi:10.1101/539437.

834   51.   Amedi, A., Raz, N., Pianka, P., Malach, R. & Zohary, E. Early 'visual' cortex activation
835   correlates with superior verbal memory performance in the blind. *Nat. Neurosci.* **6**, 758–766
836   (2003).

837   52.   Burton, H. Visual Cortex Activity in Early and Late Blind People. *J. Neurosci.* **23**, 4005–4011
838   (2003).

839   53.   Seydell-Greenwald, A., Wang, X., Newport, E. L., Bi, Y. & Striem-Amit, E. Spoken language
840   processing activates the primary visual cortex. *PLoS One* **18**, 1–22 (2023).

841   54.   Pascual-Leone, A. & Hamilton, R. The metamodal organization of the brain. *Prog. Brain Res.*
842   **134**, 427–445 (2001).

843   55.   Makin, T. R. & Krakauer, J. W. Against cortical reorganisation. 1–43 (2023).

844   56.   Striem-Amit, E., Dakwar, O., Reich, L. & Amedi, A. The large-scale organization of 'visual'
845   streams emerges without visual experience. *Cereb. Cortex* **22**, 1698–1709 (2012).

846   57.   Liu, Y.-F., Rapp, B. & Bedny, M. Reading Braille by Touch Recruits Posterior Parietal
847   Cortex. *J. Cogn. Neurosci.* **35**, 1593–1616 (2023).

848   58.   Amedi, A., Malach, R., Hendler, T., Peled, S. & Zohary, E. Visuo-haptic object-related
849   activation in the ventral visual pathway. *Nat. Neurosci.* **4**, 324–330 (2001).

850   59.   Bedny, M. *et al.* How does visual experience shape representations and transformations along
851   the ventral stream ? *CCN Gener. Advers. Collab.* (2021).

852   60.   Sanchez-Panchuelo, R. M. *et al.* Within-digit functional parcellation of brodmann areas of the
853   human primary somatosensory cortex using functional magnetic resonance imaging at 7 tesla.
854   *J. Neurosci.* **32**, 15815–15822 (2012).

855   61.   Bonda, E., Petrides, M. & Evans, A. Neural systems for tactual memories. *J. Neurophysiol.* **75**,
856   1730–1737 (1996).

857   62.   Burton, H. & Sinclair, R. J. Attending to and Remembering Tactile Stimuli. *J. Clin.*
858   *Neurophysiol.* **17**, 575–591 (2000).

859   63.   Reed, C. L., Shoham, S. & Halgren, E. Neural Substrates of Tactile Object Recognition: An
860   fMRI Study. *Hum. Brain Mapp.* **21**, 236–246 (2004).

861   64.   Banati, R. The functional anatomy of visual-tactile integration in man: a study using positron
862   emission tomography. *Neuropsychologia* **38**, 115–124 (2000).

863   65.   Hadjikhani, N. & Roland, P. E. Cross-modal transfer of information between the tactile and
864   the visual representations in the human brain: A positron emission tomographic study. *J.*
865   *Neurosci.* **18**, 1072–1084 (1998).

866   66.   Prather, S. C., Votaw, J. R. & Sathian, K. Task-specific recruitment of dorsal and ventral
867   visual areas during tactile perception. *Neuropsychologia* **42**, 1079–1087 (2004).

868   67.   Naghavi, H. R., Eriksson, J., Larsson, A. & Nyberg, L. The claustrum/insula region integrates
869   conceptually related sounds and pictures. *Neurosci. Lett.* **422**, 77–80 (2007).

870   68.   Ling, S., Lee, A. C. H., Armstrong, B. C. & Nestor, A. How are visual words represented?
871   Insights from EEG-based visual word decoding, feature derivation and image reconstruction.
872   *Hum. Brain Mapp.* **40**, 5056–5068 (2019).

873   69.   Teng, S., Cichy, R., Pantazis, D. & Oliva, A. The neural dynamics of letter perception in blind
874   and sighted readers. *J. Vis.* **15**, 126 (2015).

70.   Müller, F. *et al.* A thalamocortical pathway for fast rerouting of tactile information to occipital
cortex in congenital blindness. *Nat. Commun.* **10**, 1–9 (2019).

71.   Bola, Ł. *et al.* Braille in the sighted: Teaching tactile reading to sighted adults. *PLoS One* **11**,
1–13 (2016).

72.   Brysbaert, M. How many words do we read per minute? A review and meta-analysis of
reading rate. *J. Mem. Lang.* **109**, 104047 (2019).

73.   Bragg, D., Bennett, C., Reinecke, K. & Ladner, R. A large inclusive study of human listening
rates. *Conf. Hum. Factors Comput. Syst. - Proc.* **2018**-**April**, 1–12 (2018).

74.   Hötting, K. & Röder, B. Auditory and auditory-tactile processing in congenitally blind
humans. *Hear. Res.* **258**, 165–174 (2009).

75.   Röder, B., Rösler, F., Hennighausen, E. & Näcker, F. Event-related potentials during auditory
and somatosensory discrimination in sighted and blind human subjects. *Cogn. Brain Res.* **4**,
77–93 (1996).

76.   Muchnik, C., Efrati, M., Nemeth, E., Malin, M. & Hildesheimer, M. Central Auditory Skills In
Blind And Sighted Subjects. *Scand. Audiol.* **20**, 19–23 (1991).

77.   Bola, Ł. *et al.* Functional hierarchy for tactile processing in the visual cortex of sighted adults.
Neuroimage **202**, (2019).

78.   Smith, S. M. Fast robust automated brain extraction. *Hum. Brain Mapp.* **17**, 143–155 (2002).

79.   Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved optimization for the robust and
accurate linear registration and motion correction of brain images. *Neuroimage* **17**, 825–841
(2002).

80.   Jenkinson, M. & Smith, S. A global optimisation method for robust affine registration of brain
images. *Med. Image Anal.* **5**, 143–156 (2001).

81.   Wang, L., Mruczek, R. E. B., Arcaro, M. J. & Kastner, S. Probabilistic maps of visual
topography in human cortex. *Cereb. Cortex* **25**, 3911–3931 (2015).

82.   Eickhoff, S. B. *et al.* A new SPM toolbox for combining probabilistic cytoarchitectonic maps
and functional imaging data. *Neuroimage* **25**, 1325–1335 (2005).

83.   Delorme, A. & Makeig, S. EEGLAB: An open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. *J. Neurosci. Methods* **134**, 9–21 (2004).

84.   Chaumon, M., Bishop, D. V. M. & Busch, N. A. A practical guide to the selection of
independent components of the electroencephalogram for artifact correction. *J. Neurosci.*
Methods **250**, 47–63 (2015).

85.   Chang, C.-C. & Lin, C.-J. Libsvm: A library for support vector machines. *ACM Trans. Intell.*
Syst. Technol. **2**, 1–27 (2011).

86.   Kriegeskorte, N., Goebel, R. & Bandettini, P. Information-based functional brain mapping.
Proc. Natl. Acad. Sci. **103**, 3863–3868 (2006).

87.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).