Targeted genotyping-by-sequencing of potato and data analysis with R/polyBreedR

Jeffrey B. Endelman*[1], Moctar Kante[2], Hannele Lindqvist-Kreuze[2], Andrzej Kilian[3], Laura M. Shannon[4], Maria V. Caraza-Harter[1], Brieanne Vaillancourt[5], Kathrine Mailloux[5], John P. Hamilton[5], C. Robin Buell[5]

[1]Department of Plant & Agroecosystem Sciences, University of Wisconsin–Madison, Madison, WI, USA.

[2]Genomics, Genetics and Crop Improvement, International Potato Center, Lima, Peru.

[3]Diversity Arrays Technology Pty Ltd, University of Canberra, Bruce, ACT, Australia.

[4]Department of Horticultural Science, University of Minnesota, Saint Paul, MN, USA.

[5]Center for Applied Genetic Technologies, University of Georgia, Athens, GA, USA

*Corresponding author (endelman@wisc.edu)

J.B.E. ORCID 0000-0003-0957-4337

M.K. ORCID 0000-0003-4669-3132

H.L-K. ORCID 0000-0002-6523-8824

L.M.S. ORCID 0000-0003-3935-4909

M.C-H. ORCID 0000-0002-5934-1526

B.V. ORCID 0000-0002-6795-5173

J.P.H. ORCID 0000-0002-8682-5526

C.R.B. ORCID 0000-0002-6727-4677

Core Ideas

1. A mid-density, targeted genotyping-by-sequencing (GBS) assay was developed for potato.

2. The GBS assay includes markers for resistance to potato virus Y, golden cyst nematode, and potato wart.

3. The GBS assay includes multi-allelic markers for potato maturity and tuber shape.

4. The polyBreedR software has functions for manipulating and imputing polyploid marker data in Variant Call Format.

5. Linkage Analysis was more accurate than the Random Forest method when imputing from 2K to 10K markers.

ABSTRACT

Mid-density targeted genotyping-by-sequencing (GBS) combines trait-specific markers with thousands of genomic markers at an attractive price for linkage mapping and genomic selection. A 2.5K targeted GBS assay for potato was developed using the DArTag^{TM} technology and later expanded to 4K targets. Genomic markers were selected from the potato Infinium^{TM} SNP array to maximize genome coverage and polymorphism rates. The DArTag and SNP array platforms produced equivalent dendrograms in a test set of 298 tetraploid samples, and 83% of the common markers showed good quantitative agreement, with RMSE (root-mean-squared-error) less than 0.5. DArTag is suited for genomic selection candidates in the clonal evaluation trial, coupled with imputation to a higher density platform for the training population. Using the software polyBreedR, an R package for the manipulation and analysis of polyploid marker data, the RMSE for imputation by linkage analysis was 0.15 in a small half-diallel population (N=85), which was significantly lower than the RMSE of 0.42 with the Random Forest method. Regarding high-value traits, the DArTag markers for resistance to potato virus Y, golden cyst nematode, and potato wart appeared to track their targets successfully, as did multi-allelic markers for maturity and tuber shape. In summary, the potato DArTag assay is a transformative and publicly available technology for potato breeding and genetics.

1   INTRODUCTION

Targeted genotyping-by-sequencing (GBS) has become an essential technology for molecular plant breeding. As with restriction site-associated DNA (RAD) sequencing (Baird et al., 2008; Elshire et al., 2011), targeted GBS is based on sequencing a reduced representation of the genome. A key difference is that targeted GBS uses a fixed set of primer pairs or oligonucleotide baits, with the number of targets designed based on the application and price point (Campbell et al., 2015; Gasc et al., 2016; Ali et al., 2016). DArTag is a targeted GBS method based on PCR with molecular inversion probes (Hardenbol et al. 2003) and scalable to thousands of targets (Hardigan et al., 2023; Zhao et al., 2023). As part of the CGIAR Excellence in Breeding platform, DArTag panels were developed for wheat, maize, rice, cowpea, pigeon pea, common bean, groundnut (peanut), sorghum, and potato (Excellence in Breeding, 2022). This article describes the design and validation of the first potato DArTag panel, which had 2.5K targets, as well as a second design project, which extended the assay to 4K targets.

Before DArTag, there was no comparable "mid-density" genotyping service for potato. The main genotyping platform for genetic mapping and genomic selection in potato has been an Infinium^{TM} SNP array, which was originally developed with 8303 markers and then expanded to 12K (Version 2) based on the same discovery panel of 6 varieties (Hamilton et al., 2011; Felcher et al., 2012). The 22K V3 array incorporated new SNPs from a larger discovery panel of 83 tetraploid varieties (Uitdewilligen et al., 2013; Vos et al., 2015), and the 31K V4 array added markers from yet another discovery pool (Sharma and Bryan, 2017). To maintain backwards compatibility with existing marker data, the genomic markers for DArTag were selected from the potato Infinium array.

In addition to genomic markers, the DArTag design includes "essential markers" that are

prioritized during the final marker selection and primer design process. For potato, our initial

priority was identifying markers with high diagnostic value (i.e., haplotype-specificity) for key

resistance genes in a wide variety of genetic backgrounds. When the V1 assay was designed in

2020, KASP markers for the Ry_{adg} (Herrera et al., 2018) and Ry_{sto} (Nie et al., 2016) resistance

genes against potato virus Y (PVY) were being widely utilized through a "low-density"

genotyping service of the Excellence in Breeding platform. These two markers were therefore

obvious candidates to include in the V1 DArTag panel. When the V2 DArTag panel was

designed in 2023, a number of additional traits were targeted with essential markers.

R package polyBreedR (https://github.com/jendelman/polyBreedR) was initiated in 2020 to

publicly distribute software used for genomics-assisted breeding of potato (Endelman et al. 2017;

Endelman et al. 2018). Since elite potato germplasm is autotetraploid, all polyBreedR functions

work for both diploid and tetraploid marker data, and most work for higher ploidy too. The focus

of the original functions was SNP array data, which was more commonly used than GBS in

potato and other polyploids due to the high read depth needed to differentiate heterozygotes with

different allele dosage (Uitdewilligen et al., 2013). The required depth for < 5% error in a

tetraploid ranges from 30 to 60, depending on the population structure and other assumptions

(Gerard et al. 2018; Matias et al. 2019). Because the number of genome fragments is much

smaller with mid-density, targeted GBS than RAD-seq, read depth is less of a limitation. In

parallel to developing the DArTag assay for potato, new functions were added to polyBreedR to

facilitate the manipulation and imputation of marker data in Variant Call Format (VCF).

2 MATERIALS AND METHODS

2.1 Genomic markers

SNPs were selected from the 22K V3 SNP array (Felcher et al., 2012; Vos et al., 2015) for the 2.5K V1 DArTag set, and additional SNPs were selected from the V4 31K SNP array for the 4K V2 DArTag set. Physical positions were based on the DMv6.1 reference genome (Pham et al., 2020). Genetic map positions (in cM) were interpolated from the map positions reported in Endelman and Jansky (2016). The interpolated "Marey" map of cM vs. bp was constrained to be monotone nondecreasing (Figure S1) using an I-spline basis with 12 degrees of freedom, generated with R/splines2 (Wang and Yan, 2021). Non-negative basis coefficients were computed by minimizing the mean-squared error with R/CVXR (Fu et al., 2020). The script is available as function *interpolate_cM* in R/MapRtools (Endelman, 2023a). Initially, SNPs were selected based on discretizing the genome into 1 cM bins, and within each bin, SNPs were prioritized based on minor allele frequency (MAF) in a collection of US and CIP germplasm. After saturating the genome, additional SNPs were selected sequentially based on the ad-hoc score $d + 10 \times MAF$, where d is cM distance to the closest selected SNP.

Germplasm for evaluating V1 DArTag came from the International Potato Center (CIP) and University of Wisconsin breeding programs. Data for 703 tetraploid samples are provided as VCF File S1, with the year of submission for each sample recorded in File S2. The function *dart2vcf* in polyBreedR generates a VCFv4.3 compliant file from the two standard DArTag CSV files ("Allele_Dose_Report" and "Allele_match_counts_collapsed"). polyBreedR function *gbs* was used to replace the original DArTag genotype calls (FORMAT field GT) with those from function *flexdog* in R/updog, using the "norm" prior (Gerard et al., 2018). Three parameters of

the beta-binomial model (SE = sequencing error, AB = allelic bias, OD = overdispersion) were

stored for each variant. Both functions utilize R package vcfR (Knaus and Grunwald, 2017).

One submission of tetraploid (N=323) and diploid (N=52) samples was used to evaluate the

4K V2 DArTag assay. VCF File S3 contains the target SNP data, and CSV File S4 is the MADC

("missing allele discovery counts") file from DArT, which contains read counts for 81bp

haplotypes (similar to a TagsByTaxa file in TASSEL; Glaubitz et al. 2014). Genotype calls were

made separately for each ploidy group using the *gbs* function, and the resulting VCF files were

then combined with *bcftools merge* (Danecek et al. 2021); the model parameters in INFO are

from the tetraploid set. To verify ploidy, genotype calls were made for the diploid samples based

on the tetraploid model parameters, using the "model.fit=FALSE" option of function *gbs*, which

in turn uses *flexdog* with "update_bias", "update_seq", and "update_od" set to FALSE.

A comparison of DArTag vs. SNP array genotypes was conducted using 298 clones for V1

DArTag and 78 clones for V2 DArTag. XY intensity values and genotype calls are provided in

File S5 for 15,187 markers from the V4 SNP array, based on a normal mixture model estimated

with R/fitPoly (Voorrips et al., 2011; Zych et al., 2019). The parameter file for the normal

mixture model is distributed with the polyBreedR package as "potato_V4array_model.csv" and

was used to convert Genome Studio Final Reports to VCF with function *array2vcf*. This function

also requires a VCF map definition file to convert from B allele dosage to ALT dosage, which is

distributed as "potato_V4array.vcf" with polyBreedR. The common markers between DArTag

and the SNP array, including matching REF/ALT, were identified using *bcftools isec*.

2.2 Imputation

Two methods were compared for the accuracy of imputing SNP array markers from

DArTag: Random Forest (RF) and Linkage Analysis (LA). Method RF was implemented as

polyBreedR function *impute_L2H*, using the R/randomForest package (Liaw and Wiener, 2002).

The number of trees was set at 100 by monitoring the out-of-bag error, using the default number

of variables randomly sampled at each split, which is $\sqrt{m}$ for m classification variables. Each

marker was imputed separately, using the m closest markers as candidate prediction variables.

Results were generated for m=10, 25, 50, and 100, and the lowest error (Table 1) was observed at

25, but the optimal marker number will vary by dataset. Method LA was implemented as

polyBreedR function *impute_LA*, using the software PolyOrigin (Zheng et al., 2021) and default

parameters. Imputation error was measured using leave-one-family-out cross-validation in a five-

parent half-diallel population (pedigree in File S6). The parent codes in Table 1 are P1=W6609-

3, P2=W12078-76, P3=W13NYP102-7, P4=W14NYQ4-1, P5=W14NYQ9-2. The high density

(10K) phased parental genotypes are in File S7.

2.3 Trait markers

A set of six interconnected F1 populations was used to assess the accuracy of genotype calls

for the V2 DArTag trait markers Ryadg_chr11_2499502 and H1_chr05_52349069. Parental

phasing and haplotype reconstruction utilized PolyOrigin (Zheng et al., 2021), and binary trait

locus (BTL) analysis utilized R/diaQTL (Amadeu et al., 2021). The pedigree, genomic marker,

and dominant trait marker files needed for diaQTL are Files S8, S9, S10, respectively. Validation

of trait marker Sli_chr12_2372490 was based on the Sli_898 KASP marker (Clot et al., 2020;

Kaiser et al., 2021).

Trait markers CDF1.2_chr05_4488015 and CDF1.4_chr05_4488021 target two different 7

bp insertions of *CDF1* (Kloosterman et al., 2013; Gutaker et al., 2019) and contain equivalent

information in the DArT MADC file. The 81-bp haplotypes were aligned using MUSCLE v3.8

(Edgar, 2004). DArTag read counts for *CDF1* alleles 1, 2, and 4 were tabulated with polyBreedR

function *madc* and validated against genotypes determined via whole-genome sequencing with

NovaSeq 2x150 reads (Song and Endelman, 2023).

Genome assemblies of *S. tuberosum* dihaploids were used to validate markers for *OFP20*, a

major gene affecting tuber shape (Wu et al. 2018). High molecular weight DNA was extracted

from tissue culture plantlets using a CTAB isolation method and Qiagen Genomic tips (Hilden,

Germany), followed by an Amicon filter (MilliporeSigma, Burlington, MA) buffer exchange

(Vaillancourt et al., 2019) or Takara NucleoBond HMW DNA kit (Takara, Kusatsu, Shiga,

Japan). Genome assembly used hifiasm v0.16.1-r375 (Cheng et al. 2021, 2022) with PacBio HiFi

Sequel II (Menlo Park, CA) reads from the University of Minnesota Genomics Center. Contigs

less than 50kb were discarded using seqkit v2.3.0 (Shen et al. 2016), followed by Ragtag v2.1.0

(Alonge et al. 2019) to scaffold with DM 1-3 516 R44 v6.1 (Pham et al., 2020).

A multiple sequence alignment of 19 *OFP20* haplotypes (File S11) was generated using

MUSCLE v3.8. Alleles 1–7 and M6_ScOFP20 were reported by van Eck et al. (2022), and the

remaining haplotypes come from the dihaploids. The frequency of *OFP20.1* was approximated

by ALT frequency at marker OFP20_M6_CDS_994 (994 bp in M6 CDS). For allele *OFP20.8,*

which was discovered in the dihaploids (i.e., not in the FASTA file from van Eck et al. (2022)),

allele frequency was approximated by REF frequency at marker OFP20_M6_CDS_24; this only

works in populations without the M6_ScOFP20 allele. Marker OFP20_M6_CDS_171 was used

to report allele depth for allele 2 (ALT) vs. alleles 3 and 7 combined (REF); alleles 1 and 8 were

not detected by this marker. Marker OFP20_M6_CDS_75 was supposed to capture an indel at

82 bp that differentiates alleles 3 and 7, but neither haplotype was present in the MADC File S4.

3   RESULTS

3.1 Genomic markers

Version 1 (V1) of the potato DArTag GBS assay contained 2501 genomic SNPs, which

were selected from the 22K V3 potato SNP array to maximize genome coverage and

polymorphism rates (i.e., high minor allele frequency). The number of genomic markers per

chromosome ranged from 176 on chr12 to 272 on chr01. The mean distance between adjacent

markers was 0.35 cM, with the largest gap of 4.77 cM located on chr11 (Figure S2).

Analysis of 703 tetraploid samples, from three submissions across three years (2020-2022),

revealed variability in the amount of sequencing data per sample. In 2020, the total depth (DP

sum over markers) was consistent across samples, with mean 0.53M/sample and standard

deviation 0.07M (Figure 1). The distribution in 2021 was bimodal, with the two modes

corresponding to different plates. The lower mode was 0.63M, while the higher mode was

0.96M. The average total depth in 2022 was similar to 2020, at 0.53M/sample, but the standard

deviation was higher, at 0.17M.

When sample DP was summarized by marker, the data were more consistent across years

(Figure 2). The 10^{th} percentile for mean sample DP was 32, 53, and 24 in years 2020, 2021, and

2022, respectively (Fig. 2A). Despite the observed differences in total DP per sample (Figure 1),

there was a consistent relationship between the mean (μ) and standard deviation (σ) for sample

DP (Fig. 2B). The relationship between these quantities in a Poisson distribution is $\mu = \sigma^{0.5}$,

which is a straight line with slope 0.5 on a log-log plot (dashed line in Fig. 2B). The observed

data were overdispersed (i.e., more variable) compared to the Poisson, with slope 0.79 (SE 0.00),

meaning that $\mu \approx \sigma^{0.8}$.

Tetraploid genotype calls were made with R package *updog* (Gerard et al. 2018), which

provides estimates of allelic bias (AB) for each marker—a parameter that measures the relative

probability of observing the REF vs. ALT allele. When AB=1, or equivalently $\log_2(AB) = 0$,

there is no bias. When AB=2, or equivalently $\log_2(AB)=1$, the REF allele is twice as likely to be

observed in a balanced heterozygote. 10% of the markers exhibited bias $|(AB)| > 1$ (Figure S3),

but many of these still appeared to have reliable clustering (Figure 3).

V1 DArTag and SNP array genotypes were compared for 1865 common markers across 298

tetraploid clones. Both platforms identified two groups of genetically identical clones, one pair

and one threesome, originating from the same F1 populations (Figure S4). This is not uncommon

in potato breeding due to how single plant selection is conducted in the first field year. After

removing duplicates, the two marker profiles (GBS & array) for every clone were paired under

hierarchical clustering (Figure S5), indicating close agreement.

For a quantitative comparison, several measures of error were computed for each marker

(File S12). Classification error (CE), which is the proportion of samples with different genotype

calls, was calculated for both tetraploid (4x) and pseudo-diploid (2x) genotypes (where

differences in heterozygote allele dosage are ignored). There was a sharp bend in the cumulative

distribution for 2x CE at approximately 0.1 error (Figure 4), with 1647 markers below this

threshold (88% of those tested). As expected, fewer markers (1302) satisfied 4x CE < 0.1

because of the difficulty discriminating between heterozygous genotypes. For 4x genotypes, the

root-mean-squared-error (RMSE) of allele dosage is potentially more meaningful than CE, and

1547 markers had RMSE < 0.5 (Figure 4), a somewhat arbitrary threshold selected because it

represents the midpoint between integer dosages.

Version 2 (V2) of the potato DArTag GBS assay was designed in 2023 and contains 3893

genomic SNPs, of which 2144 were included in V1. The additional SNPs were selected from the

31K V4 potato SNP array using the same criteria as before. GBS and SNP array genotypes were

compared for 2608 common markers across 78 clones (40 tetraploid, 38 diploid). Given the

small number of tetraploids, only the 2x CE criterion was computed, and 2341 markers had 2x

CE < 0.1 (Figure S6; File S13).

3.2 Imputation and ploidy analysis

A key role for the DArTag genomic markers is to facilitate imputation to higher density

platforms for genomic selection. Among the 298 clones genotyped with both the SNP array and

V1 DArTag is a five-parent half-diallel population of 85 clones, with F1 family sizes between 1

and 20 (Table 1). The accuracy of two imputation methods was compared by leave-one-family-

out cross-validation: Random Forest (RF) vs. Linkage Analysis (LA). Linkage analysis uses a

genetic model of recombination and phased parental genotypes to reconstruct progeny in terms

of parental haplotypes. The RMSE for imputing 10K SNP array genotypes from DArTag was

always lower with LA compared to RF (Table 1), with overall means of 0.15 and 0.42,

respectively.

The *check_ploidy* function of polyBreedR was originally developed to use SNP array

markers to differentiate diploid from tetraploid samples based on the following principle: when a

bi-allelic genotype model developed for tetraploids is applied to a diploid sample, ideally there

would be no simplex (AAAB) or triplex (ABBB) heterozygotes, only duplex (AABB) calls

because both AB and AABB genotypes have 1:1 allele ratios. The proportion of calls that are

simplex or triplex can therefore be used to discriminate ploidies. This method has been used

extensively during recent haploid induction crosses of tetraploid potato, where the desired result

is a diploid haploid, aka dihaploid, containing only the maternal chromosomes, but sometimes

the progeny are tetraploid or aneuploid (Amundson et al. 2020; Busse et al. 2021).

The *check_ploidy* function was extended to allow VCF file input and the R/updog model for

genotype calling of GBS markers. When applied to the V2 DArTag dataset (File S3), the 323

tetraploids were clearly separated from the 52 diploids (Figure 5).

3.3 Trait markers

The V1 DArTag assay had only two trait markers, targeting two different resistance genes

(Ry_{adg}, Ry_{sto}) for potato virus Y (PVY), which is the most economically important viral pathogen

of potato. Both variants had previously been targeted with KASP markers, and for 93 samples

genotyped with both KASP and V1 DArTag, there were 2 discrepancies for presence/absence of

Ry_{adg} (Table S1). Both PVY markers were carried forward to the V2 DArTag assay, and four

clones tested positive for Ry_{sto}: three were expected based on previous testing, and the fourth was

plausible based on its pedigree (Table S2). Many samples in the V2 submission tested positive

for Ry_{adg}, which was expected given its high frequency in US chip processing germplasm, but the

allele dosages seemed too high—eight samples were even homozygous tetraploids. To

investigate further, a partial diallel population (N=123) within the dataset was analyzed (Figure

S7). Treating the Ry_{adg} marker as a dominant trait, joint linkage analysis identified which

parental haplotypes carry the R gene (Figure S8), and corrected dosages were determined by

reconstructing the progeny in terms of parental haplotypes (Figure 6). Five triplex and two

quadriplex calls for Ry_{adg} were corrected down to duplex, and the average upward bias was 0.24

dosage.

Besides the two PVY markers, the V2 DArTag assay has five additional trait markers with

reliable results (Table 2). Like Ry_{adg}, the golden cyst nematode resistance gene *H1* was common

in US chip processing germplasm, but diallel analysis indicated the *H1* marker calls were more

accurate, with an average bias of only 0.05 dosage (Figure 6).

Little is known about resistance to potato wart disease (*S. endobioticum*) in US germplasm,

but given the prevalence of the disease in other parts of the world (Obidiegwu et al., 2014), it has

become a higher priority for molecular breeding. One trait marker targets the *Sen3* resistance

gene, which was detected in four individuals with a common parent, AW07791-2rus. Based on

pedigree information, the resistance appears to have been inherited from the maternal parent,

PALB0303-1 (Elison et al., 2021).

Another trait marker targets *Sli*, a non-S locus F-box protein that disrupts the gametophytic

incompatibility system and allows for the development of diploid, inbred lines (Ma et al., 2022;

Eggers et al., 2022). The GBS marker showed perfect agreement with prior knowledge for 28

diploid samples based on KASP marker screening (Table S3).

A trait marker for the maturity gene *CDF1* targets the location of the 7 bp indel variants that

differentiate alleles 2 and 4 from wild-type alleles, collectively designated group 1. Because of

the multi-allelic nature of this variant, correct interpretation requires use of the DArT "missing

allele discovery count" (MADC) file, which contains read counts for 81 bp haplotypes

surrounding each target variant. Five CDF1 haplotypes were detected in the population (Figure

7A): three were full-length variants of CDF1.1 (Ref, Other1, Other2), one was CDF1.4 (Alt), and

one was CDF1.2 (Other3). The validity of the assay was confirmed by comparing the read counts

with samples of known CDF1 genotype (Figure 7B), with the complication that CDF1.3, which

has an 865 bp transposon insertion at the same position, is not detected. As a result, samples with

zero (or near zero, due to sequencing error) counts are interpreted as homozygous for allele 3.

And since clones selected under long-day conditions are typically not homozygous wild-type,

when CDF1.1 alleles are detected but not alleles 2 or 4, the predicted genotype is 1/3.

Several markers were included in the V2 panel to target *OFP20*, an ovate family protein

with a major effect on tuber shape (Wu et al. 2018). This is a complex locus with dozens of

predicted alleles (van Eck et al. 2022), so the following approach to interpreting the DArTag

markers may not work in all germplasm groups. Marker OFP20_M6_CDS_994 was used to

estimate the frequency of *OFP20.1*, which is the most common allele in cultivated germplasm

and promotes elongated shape (van Eck et al. 2022). *OFP20.1* was present at a higher frequency

in the russet (N=21) vs. chip (N=300) samples from the UW breeding program (Fig. 8A), which

is consistent with the long vs. round tuber phenotypes required for those market types. Marker

OFP20_M6_CDS_24 was used to estimate the frequency of *OFP20.8*, which was present in 13%

of the chip samples. Together with OFP20_M6_CDS_171, which provided information about

presence/absence of *OFP20* alleles 2, 3, and 7, the DArTag markers were able to correctly

predict five different *OFP20* genotypes (Fig. 8B).

4   DISCUSSION

The potato DArTag assay has several applications in potato breeding. For its price point, an ideal stage of deployment is the first clonal evaluation trial (CET), which typically occurs in the second field year of potato breeding and may have several thousand clones. The DArTag genomic markers provide a genetic fingerprint that can be used to correct pedigree errors (Muñoz et al., 2014; Endelman et al., 2017) and provide a reference genotype for quality control. The clonal trial entries are also candidates for genomic selection, both as potential clonal varieties and as parents to begin the next breeding cycle (Slater et al., 2016; Wu et al., 2023). Limited phenotyping for some traits occurs in the CET, and a genomic relationship matrix computed from DArTag markers could enable a multi-location trial to better estimate genetic values for the target population of environments, i.e., "sparse testing" (Endelman et al. 2014; Jarquin et al. 2020).

Based on previous studies, higher selection accuracy is expected if DArTag markers are first imputed to higher density (Cleveland and Hickey, 2013; Gorjanc et al., 2017). The exploitation of pedigree or family structure during marker imputation in diploids is well documented, with a range of methods and software available depending on the structure of the dataset (Meuwissen and Goddard, 2010; Swarts et al., 2014; Hickey et al., 2015; Whalen et al., 2018; Whalen et al., 2020). The present study has confirmed our hypothesis that linkage analysis is also beneficial for imputation in autopolyploids. DArTag panels are available for several autopolyploid crops besides potato, including alfalfa, blueberry, and sweetpotato (Breeding Insight, 2023), so the software developed for this study should benefit other breeding communities. Based on the current functionality of the PolyOrigin software (Zheng et al., 2021), only bi-allelic SNPs were used for imputation, but the DArTag MADC file offers the possibility of using multi-allelic

haplotypes as markers, which are generally more informative for linkage analysis (Luo et al.,

2001).

Besides more genomic markers, a major advantage of V2 compared to V1 DArTag is the

additional trait markers (Table 2). It is very valuable to select for resistance to three important

pests of potato—PVY, wart, and golden cyst nematode—with the same assay used for genomic

selection. Notably absent from this list is potato late blight, caused by the pathogen *P. infestans*.

Trait marker blb1_chr08_51070621 was designed to target the *RB/Rpi-blb1* gene (Song et al.,

2003; van der Vossen et al., 2003) based on a SNP in the 3´UTR that worked well as a KASP

marker (Sorensen et al., 2023). However, no haplotypes were detected in the V2 DArTag

experiment for three positive samples from the KASP study. The V2 assay also targeted two

genes affecting tuber skin color: *f3´5´h* (Jung et al., 2005) and *an2* (Jung et al., 2009). Both loci

have complex allelic series (Hoopes et al. 2022), and more information is needed about their

functional effects to guide selection. For tuber shape, a trait marker for the most common long

allele (*OFP20.1*) can have an immediate impact on parent selection in the russet market type,

where round alleles are undesirable due to their partial dominance.

Table 3 summarizes the new or updated functions in polyBreedR associated with this

research. Examples of their usage on a sample dataset can be found in Vignette 3 of the package

and the Supplemental Methods file, which provides code to generate the main figures and tables.

 FIGURE CAPTIONS

Figure 1. Total depth per sample, in million (M) read counts, for three submissions of potato V1

DArTag.

Figure 2. (A) Distribution of the mean sample depth (DP) for V1 DArTag markers. (B) Log-log

plot of the relationship between the standard deviation and mean for sample DP. Individual

marker points are shown only for 2021 to maintain legibility. Combining the data across years,

the overall regression line (not shown) has slope 0.79 (SE 0.00) and $R^2 = 0.99$.

Figure 3. Examples of DArTag markers without (A) vs. with (B) allelic bias. Dashed lines

correspond to possible tetraploid allele ratios when there is no allelic bias (1:0, 3:1, 1:1, 1:3, 0:1).

(A) solcap_snp_c2_36615 with bias = -0.2. (B) PotVar0072076 with bias = 1.8.

Figure 4. Empirical cumulative distribution for the error between the V1 DArTag and SNP array

on 1865 common markers. CE = classification error. RMSE = root-mean-squared-error. 2x =

pseudo-diploid genotypes. 4x = tetraploid genotypes.

Figure 5. Violin plot illustrating ploidy discrimination with the V2 DArTag assay, based on the

proportion of simplex or triplex markers with a tetraploid genotype calling model. The diploid

potato samples had much lower values of this parameter.

Figure 6. Original vs. corrected genotypes for the trait markers Ryadg_chr11_2499502 and

H1_chr05_52349069. The original genotypes were based on R/updog with a "norm" prior and

then corrected based on linkage analysis.

Figure 7. (A) Multiple sequence alignment of the DArTag haplotypes discovered for trait marker

CDF1.4_chr05_448021. Haplotypes Ref, Other1, Other2 are CDF1.1 alleles, while Alt is

CDF1.4 and Other3 is CDF1.2. (B) Haplotype read counts for samples with known CDF1

genotype.

Figure 8. (A) Distribution of sample allele frequencies for OFP20.1 in round chip (N=300) vs. long russet (N=21) germplasm. (B) Comparison of known OFP20 genotypes with V2 DArT markers. Allele frequency (AF) of OFP20.1 was approximated by ALT frequency at marker OFP20_M6_CDS_994. AF of OFP20.8 was approximated by REF frequency at marker OFP20_M6_CDS_24. Allele depth (AD) at OFP20_M6_CDS_171 was used to distinguish allele 2 (ALT) from alleles 3 and 7 (REF).

TABLES

Table 1. Half-diallel population with five parents. Above diagonal: F1 population sizes; Below diagonal: imputation root-mean-squared-error with linkage analysis (blue, top) vs. random forest (red, bottom).

	P1	P2	P3	P4	P5
P1		3	5	9	20
P2	0.14 0.43		8	1	9
P3	0.15 0.40	0.17 0.45		7	11
P4	0.13 0.42	0.14 0.38	0.13 0.41		12
P5	0.16 0.43	0.16 0.44	0.14 0.40	0.15 0.42	

397   **Table 2.** Validated trait markers in the V2 DArTag assay.

Marker	Target Gene (Trait)	Functional Allele	Reference
Rysto_chr12_2352742	Ry_{sto} (PVY)	REF[a]	Nie et al. (2016)
Ryadg_chr11_2499502	Ry_{adg} (PVY)	ALT	Herrera et al. (2018)
H1_chr05_52349069	*H1* (golden cyst nematode)	ALT	Meade et al. (2020)
Sen3_chr11_2563398	*Sen3* (wart)	ALT	Prodhomme et al. (2019)
Sli_chr12_2372490	*Sli* (self-compatibility)	ALT	Clot et al. (2020)
CDF1.4_chr05_4488021	*CDF1* (maturity)	N/A	Gutaker et al. (2019)
OFP20_M6_CDS_24 OFP20_M6_CDS_171 OFP20_M6_CDS_994	*OFP20* (tuber shape)	N/A	van Eck et al. (2022)

398   [a] The REF/ALT designation for Rysto_chr12_2352742 is reversed compared to the original design file
399   based on the DMv6.1 reference genome. As a result, the functional allele is REF.

403   **Table 3.** New or updated functions in R/polyBreedR developed for this research.

Function	Description
dart2vcf	converts DArTag data files to VCF
array2vcf	converts Genome Studio Final Report to VCF file
gbs	adds genotype calls to a VCF file based on the R/updog model
impute_L2H	imputes low to high density markers by Random Forest
impute_LA	imputes low to high density markers by Linkage Analysis
check_ploidy	discriminates diploid from tetraploid samples

SUPPLEMENTAL FILES

During peer review, the supplemental files are available from the Dryad Digital Repository at

https://datadryad.org/stash/share/tdvUt18gBCz6bJ568DaE7mLrWtq55kZzJa_C1uLYSfQ. The

permanent link for the supplemental files after publication is

https://doi.org/10.5061/dryad.8pk0p2nw4.

File S1. Potato DArTag V1 data for 703 samples (VCF).

File S2. Metadata with year submission for the samples in File S1 (CSV).

File S3. Potato DArTag V2 data for 375 samples (VCF).

File S4. DArT Missing Allele Discovery Counts for the samples in File S3 (CSV).

File S5. Potato V4 SNP array data for 298 samples (VCF).

File S6. Pedigree for diallel population in the V1 DArTag dataset (CSV).

File S7. Phased parental genotypes for the diallel population in File S6 (CSV).

File S8. Pedigree for diallel population in the V2 DArTag dataset (CSV).

File S9. Parental genotype probabilities for the diallel population in File S8 (CSV).

File S10. Trait marker phenotypes for the diallel population in File S8 (CSV).

File S11. Sequence alignment and percent identity matrix for *OFP20* (DOCX).

File S12. Marker concordance between V1 DArTag and the SNP array (CSV).

File S13. Marker concordance between V2 DArTag and the SNP array (CSV).

AUTHOR CONTRIBUTIONS

Jeffrey B. Endelman: Conceptualization, Resources, Investigation, Formal analysis, Software, Supervision, Writing – original draft. **Moctar Kante**: Conceptualization, Resources, Investigation, Formal analysis, Writing – original draft. **Hannele Lindqvist-Kreuze**: Conceptualization, Resources, Supervision. **Andrzej Kilian**: Methodology. **Laura M. Shannon:** Conceptualization, Supervision. **Maria V. Caraza-Harter:** Resources. **Brieanne Vaillancourt:** Formal analysis, Data curation. **Kathrine Mailloux:** Investigation, Resources. **John P. Hamilton:** Formal analysis. **C. Robin Buell:** Conceptualization, Supervision. **All authors**: Writing – review & editing.

ACKNOWLEDGMENTS

Development of the V1 DArTag panel was supported by the CGIAR Excellence in Breeding Platform and Crops to End Hunger Initiative. The USDA National Institute of Food & Agriculture (NIFA) Award 2019-51181-30021 supported development of the V2 DArTag panel, with additional support from PepsiCo for the development of genomic resources to validate markers. Genotyping of UW-Madison potato breeding lines was supported by USDA NIFA Awards 2020-51181-32156 and 2021-34141-35447. We thank D. DeKoeyer for suggesting the marker for *Sen3* resistance.

CONFLICT OF INTEREST STATEMENT

J. Endelman is a member of the editorial board for The Plant Genome. A. Kilian is an employee of Diversity Arrays Technology, the company that provides the DArTag genotyping service.

DATA AVAILABILITY STATEMENT

Supplemental Files S1 – S13, which contain the marker and pedigree data needed to reproduce

the results of this study, will be available from the Dryad Digital Repository at

https://doi.org/10.5061/dryad.8pk0p2nw4 upon publication. Upon manuscript acceptance,

PacBio HiFi sequencing data will be available via the NCBI Sequence Read Archive under

BioSamples SAMN38982152, SAMN38982165, SAMN38982166, SAMN38982167, and

SAMN38982169, and Illumina sequencing data will be available via the NCBI Sequence Read

Archive under BioSamples SAMN39419651, SAMN39670896, SAMN39670897, and

SAMN39670898.

REFERENCES

Ali, O.A., O'Rourke, S.M., Amish, S.J., Meek, M.H., Luikart, G., Jeffres, C., & Miller, M.R. (2016). Rad capture (Rapture): Flexible and efficient sequence-based genotyping. *Genetics*, *202*, 389–400. https://doi.org/10.1534/genetics.115.183665

Alonge, M., Soyk, S., Ramakrishnan, S. (2019). RaGOO: fast and accurate reference-guided scaffolding of draft genomes. *Genome Biology*, *20*, 224. https://doi.org/10.1186/s13059-019-1829-6

Amadeu, R.R., Muñoz, P.R., Zheng, C., & Endelman, J.B. (2021). QTL mapping in outbred tetraploid (and diploid) diallel populations. *Genetics*, *219*, iyab124. https://doi.org/10.1093/genetics/iyab124

Amundson, K.R., Ordoñez, B., Santayana, M., Tan, E.H., Henry, I.M., Mihovilovich, E., Bonierbale, M., & Comai, L. (2020). Genomic outcomes of haploid induction crosses in potato (Solanum tuberosum L.). *Genetics*, *214*, 369–380. https://doi.org/10.1534/genetics.119.302843

Baird, N.A., Etter, P.D., Atwood, T.S., Currey, M.C., Shiver, A.L., Lewis, Z.A., Selker, E.U., Cresko, W.A., & Johnson, E.A. (2008). Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLoS ONE*, *3*, e3376. https://doi.org/10.1371/journal.pone.0003376

Breeding Insight (2023). Open Source Genetic Marker Panels. https://breedinginsight.org/breeding-solutions/open-source-dartag-marker-panels/ (Accessed 29 December 2023).

Busse, J.S., Jansky, S.H., Agha, H.I., Schmitz Carley, C.A., Shannon, L.M., & Bethke, P.C. (2021). A high throughput method for generating dihaploids from tetraploid potato. *American Journal of Potato Research, 98,* 304–314. https://doi.org/10.1007/s12230-021-09844-1

Campbell, N.R., Harmon, S.A., & Narum, S.R. (2015). Genotyping-in-Thousands by sequencing (GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing. *Molecular Ecology Resources*, *15*, 855–867. https://doi.org/10.1111/1755-0998.12357

Cheng, H., Concepcion, G.T., Feng, X., Zhang, H., & Li H. (2021). Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nature Methods, 18*, 170-175. https://doi.org/10.1038/s41592-020-01056-5

Cheng, H., Jarvis, E.D., Fedrigo, O., Koepfli, K.P., Urban, L., Gemmell, N.J., & Li, H. (2022). Haplotype-resolved assembly of diploid genomes without parental data. *Nature Biotechnology, 40*, 1332–1335. https://doi.org/10.1038/s41587-022-01261-x

Cleveland, M.A., & Hickey, J.M. (2013). Practical implementation of cost-effective genomic selection in commercial pig breeding using imputation. *Journal of Animal Science*, *91*, 3583–3592. https://doi.org/10.2527/jas2013-6270

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane, T., McCarthy, S.A., & Davies, R.M. (2021). Twelve years of SAMtools and BCFtools. *GigaScience*, *10*. https://doi.org/10.1093/gigascience/giab008

Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, *32*, 1792–1797. https://doi.org/10.1093/nar/gkh340

Eggers, E.J., van der Burgt, A., van Heusden, S.A.W., de Vries, M.E., Visser, R.G.F., Bachem, C.W.B., & Lindhout, P. (2021). Neofunctionalisation of the Sli gene leads to self-

compatibility and facilitates precision breeding in potato. *Nature Communications*, *12*. https://doi.org/10.1038/s41467-021-24267-6

Elison, G.L., Novy, R.G., & Whitworth, J.L. (2021). Russet Potato Breeding Clones with Extreme Resistance to Potato Virus Y Conferred by Rychc as well as Resistance to Late Blight and Cold-Induced Sweetening. *American Journal of Potato Research*, *98*, 411–419. https://doi.org/10.1007/s12230-021-09852-1

Elshire, R.J., Glaubitz, J.C., Sun, Q., Poland, J. a, Kawamoto, K., Buckler, E.S., & Mitchell, S.E. (2011). A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PloS ONE*, *6*, e19379. https://doi.org/10.1371/journal.pone.0019379

Endelman, J.B. (2023a). R/MapRtools. https://github.com/jendelman/MapRtools (version 0.32)

Endelman, J.B., Atlin, G.N., Beyene, Y., Semagn, K., Zhang, X., Sorrells, M.E., & Jannink, J.L. (2014). Optimal design of preliminary yield trials with genome-wide markers. *Crop Science*, *54*, 48–59. https://doi.org/10.2135/cropsci2013.03.0154

Endelman, J.B., & Jansky, S.H. (2016). Genetic mapping with an inbred line-derived F2 population in potato. *Theoretical and Applied Genetics*, *129*, 935–943. https://doi.org/10.1007/s00122-016-2673-7

Endelman, J.B., Schmitz Carley, C.A., Bethke, P.C., Coombs, J.J., Clough, M.E., da Silva, W.L., De Jong, W.S., Douches, D.S., Frederick, C.M., Haynes, K.G., Holm, D.G., Miller, J.C., Muñoz, P.R., Navarro, F.M., Novy, R.G., Palta, J.P., Porter, G.A., Rak, K.T., Sathuvalli, V.R., Thompson, A.L., & Yencho, G.C. (2018). Genetic variance partitioning and Genome-wide prediction with allele dosage information in autotetraploid potato. *Genetics*, *209*, 77–87. https://doi.org/10.1534/genetics.118.300685

Endelman, J.B., Schmitz Carley, C.A., Douches, D.S., Coombs, J.J., Bizimungu, B., De Jong, W.S., Haynes, K.G., Holm, D.G., Miller, J.C., Novy, R.G., Palta, J.P., Parish, D.L., Porter, G.A., Sathuvalli, V.R., Thompson, A.L., & Yencho, G.C. (2017). Pedigree Reconstruction with Genome-Wide Markers in Potato. *American Journal of Potato Research*, *94*, 184–190. https://doi.org/10.1007/s12230-016-9556-y

Excellence in Breeding (2022). Mid-density genotyping service. https://excellenceinbreeding.org/toolbox/services/mid-density-genotyping-service (accessed 29 December 2023)

Felcher, K.J., Coombs, J.J., Massa, A.N., Hansey, C.N., Hamilton, J.P., Veilleux, R.E., Buell, C.R., & Douches, D.S. (2012). Integration of two diploid potato linkage maps with the potato genome sequence. *PloS ONE*, *7*, e36347. https://doi.org/10.1371/journal.pone.0036347

Fu, A., Narasimhan, B., & Boyd, S. (2020). CVXR: An R package for disciplined convex optimization. *Journal of Statistical Software*, *94*, 1–34. https://doi.org/10.18637/jss.v094.i14

Gasc, C., Peyretaillade, E., & Peyret, P. (2016). Sequence capture by hybridization to explore modern and ancient genomic diversity in model and nonmodel organisms. *Nucleic Acids Research*, *44*, 4504–4518. https://doi.org/10.1093/nar/gkw309

Gerard, D., Ferrão, L.F.V., Garcia, A.A.F., & Stephens, M. (2018). Genotyping polyploids from messy sequencing data. *Genetics*, *210*, 789–807. https://doi.org/10.1534/genetics.118.301468

Glaubitz, J.C., Casstevens, T.M., Lu, F., Harriman, J., Elshire, R.J., Sun, Q., & Buckler, E.S. (2014). TASSEL-GBS: A high capacity genotyping by sequencing analysis pipeline. *PLoS ONE, 9,* e90346. https://doi.org/10.1371/journal.pone.0090346

Gorjanc, G., Battagin, M., Dumasy, J.F., Antolin, R., Gaynor, R.C., & Hickey, J.M. (2017). Prospects for cost-effective genomic selection via accurate within-family imputation. *Crop Science*, *57*, 216–228. https://doi.org/10.2135/cropsci2016.06.0526

Gutaker, R.M., Weiß, C.L., Ellis, D., Anglin, N.L., Knapp, S., Luis Fernández-Alonso, J., Prat, S., & Burbano, H.A. (2019). The origins and adaptation of European potatoes reconstructed from historical genomes. *Nature Ecology and Evolution*, *3*, 1093–1101. https://doi.org/10.1038/s41559-019-0921-3

Hamilton, J.P., Hansey, C.N., Whitty, B.R., Stoffel, K., Massa, A.N., Deynze, A. Van, Jong, W.S. De, Douches, D.S., & Buell, C.R. (2011). Single nucleotide polymorphism discovery in elite North American potato germplasm. *BMC Genomics*, *302*. https://doi.org/10.1186/1471-2164-12-302

Hardenbol, P., Banér, J., Jain, M., Nilsson, M., Namsaraev, E.A., Karlin-Neumann, G.A., Fakhrai-Rad, H., Ronaghi, M., Willis, T.D., Landegren, U., & Davis, R.W. (2003) Multiplexed genotyping with sequence-tagged molecular inversion probes. *Nature Biotechnology, 21*, 673–678. https://doi.org/10.1038/nbt821

Hardigan, M.A., Feldmann, M.J., Carling, J., Zhu, A., Kilian, A., Famula, R.A., Cole, G.S., & Knapp, S.J. (2023). A medium-density genotyping platform for cultivated strawberry using DArTag technology. *Plant Genome*, *16*. https://doi.org/10.1002/tpg2.20399

Herrera, M. del R., Vidalon, L.J., Montenegro, J.D., Riccio, C., Guzman, F., Bartolini, I., & Ghislain, M. (2018). Molecular and genetic characterization of the Ryadg locus on chromosome XI from Andigena potatoes conferring extreme resistance to potato virus Y. *Theoretical and Applied Genetics*, *131*, 1925–1938. https://doi.org/10.1007/s00122-018-3123-5

Hickey, J.M., Gorjanc, G., Varshney, R.K., & Nettelblad, C. (2015). Imputation of single nucleotide polymorphism genotypes in biparental, backcross, and topcross populations with a hidden markov model. *Crop Science*, *55*, 1934–1946. https://doi.org/10.2135/cropsci2014.09.0648

Jarquin, D., Howard, R., Crossa, J., Beyene, Y., Gowda, M., Martini, J.W.R., Pazaran, G.C., Burgueño, J., Pacheco, A., Grondona, M., Wimmer, V., & Prasanna, B.M. (2020). Genomic prediction enhanced sparse testing for multi-environment trials. *G3: Genes, Genomes, Genetics*, *10*, 2725–2739. https://doi.org/10.1534/g3.120.401349

Jung, C.S., Griffiths, H.M., De Jong, D.M., Cheng, S., Bodis, M., & De Jong, W.S. (2005). The potato P locus codes for flavonoid 3',5'-hydroxylase. *TAG. Theoretical and Applied Genetics, 110*, 269–75. https://doi.org/10.1007/s00122-004-1829-z

Jung, C.S., Griffiths, H.M., De Jong, D.M., Cheng, S., Bodis, M., Kim, T.S., & De Jong, W.S. (2009). The potato developer (D) locus encodes an R2R3 MYB transcription factor that regulates expression of multiple anthocyanin structural genes in tuber skin. *Theoretical and Applied Genetics*, *120*, 45–57. https://doi.org/10.1007/s00122-009-1158-3

Kloosterman, B., Abelenda, J. a, Gomez, M.D.M.C., Oortwijn, M., de Boer, J.M., Kowitwanich, K., Horvath, B.M., van Eck, H.J., Smaczniak, C., Prat, S., Visser, R.G.F., & Bachem, C.W.B. (2013). Naturally occurring allele diversity allows potato cultivation in northern latitudes.. *Nature*, *495*, 246–50. https://doi.org/10.1038/nature11912

Knaus, B.J., & Grünwald, N.J. (2017). VCFR: a package to manipulate and visualize variant call format data in R. *Molecular Ecology Resources*, *17*, 44–53. https://doi.org/10.1111/1755-0998.12549

Liaw, A., & Wiener, M. (2002). Classification and regression by randomForest. *R News*, *2*, 18–
22. https://doi.org/10.1177/154405910408300516

Luo, Z.W., Hackett, C.A., Bradshaw, J.E., McNicol, J.W., & Milbourne, D. Construction of a
Genetic Linkage Map in Tetraploid Species Using Molecular Markers. *Genetics, 157,* 1369–
1385.

Ma, L., Zhang, C., Zhang, B., Tang, F., Li, F., Liao, Q., Tang, D., Peng, Z., Jia, Y., Gao, M.,
Guo, H., Zhang, J., Luo, X., Yang, H., Gao, D., Lucas, W.J., Li, C., Huang, S., & Shang, Y.
(2021). A nonS-locus F-box gene breaks self-incompatibility in diploid potatoes. *Nature*
Communications, *12*. https://doi.org/10.1038/s41467-021-24266-7

Matias, F.I., Xavier Meireles, K.G., Nagamatsu, S.T., Lima Barrios, S.C., Borges do Valle, C.,
Carazzolle, M.F., Fritsche-Neto, R., & Endelman, J.B. (2019). Expected Genotype Quality
and Diploidized Marker Data from Genotyping-by-Sequencing of *Urochloa spp.*
Tetraploids. *The Plant Genome*, *12*. https://doi.org/10.3835/plantgenome2019.01.0002

Meuwissen, T., & Goddard, M. (2010). The use of family relationships and linkage
disequilibrium to impute phase and missing genotypes in up to whole-genome sequence
density genotypic data. *Genetics*, *185*, 1441–1449.
https://doi.org/10.1534/genetics.110.113936

Muñoz, P.R., Resende, M.F.R., Huber, D.A., Quesada, T., Resende, M.D.V., Neale, D.B.,
Wegrzyn, J.L., Kirst, M., & Peter, G.F. (2014). Genomic relationship matrix for correcting
pedigree errors in breeding populations: Impact on genetic parameters and genomic
selection accuracy. *Crop Science*, *54*, 1115–1123.
https://doi.org/10.2135/cropsci2012.12.0673

Nie, X., Sutherland, D., Dickison, V., Singh, M., Murphy, A.M., & De Koeyer, D. (2016).
Development and validation of high-resolution melting markers derived from Rysto STS
markers for high-throughput marker-assisted selection of potato carrying Rysto.
Phytopathology, *106*, 1366–1375. https://doi.org/10.1094/PHYTO-05-16-0204-R

Obidiegwu, J.E., Flath, K., & Gebhardt, C. (2014). Managing potato wart: A review of present
research status and future perspective. *Theoretical and Applied Genetics*, *127*, 763–780.
https://doi.org/10.1007/s00122-014-2268-0

Pham, G.M., Hamilton, J.P., Wood, J.C., Burke, J.T., Zhao, H., Vaillancourt, B., Ou, S., Jiang,
J., & Robin Buell, C. (2020). Construction of a chromosome-scale long-read reference
genome assembly for potato. *GigaScience*, *9*. https://doi.org/10.1093/gigascience/giaa100

Prodhomme, C., Esselink, D., Borm, T., Visser, R.G.F., Van Eck, H.J., & Vossen, J.H. (2019).
Comparative Subsequence Sets Analysis (CoSSA) is a robust approach to identify haplotype
specific SNPs; Mapping and pedigree analysis of a potato wart disease resistance gene *Sen3*.
Plant Methods, *15*. https://doi.org/10.1186/s13007-019-0445-5

Shen, W., Le, S., Li, Y., & Hu, F. (2016). SeqKit: a cross-platform and ultrafast toolkit for
FASTA/Q file manipulation. *PLoS ONE, 11,* e0163962.
https://doi.org/10.1371/journal.pone.0163962

Sharma, S.K., & Bryan, G.J. (2017). Genome Sequence-Based Marker Development and
Genotyping in Potato. In S. Kumar Chakrabarti, C. Xie & J. Kumar Tiwari (Eds.), *The*
Potato Genome. Compendium of Plant Genomes (pp. 307–326). Springer.

Slater, A.T., Cogan, N.O.I., Forster, J.W., Hayes, B.J., & Daetwyler, H.D. (2016). Improving
Genetic Gain with Genomic Selection in Autotetraploid Potato. *The Plant Genome*, *9*.
https://doi.org/10.3835/plantgenome2016.02.0021

Song, J., Bradeen, J.M., Kristine Naess, S., Raasch, J.A., Wielgus, S.M., Haberlach, G.T., Liu, J.,
Kuang, H., Austin-Phillips, S., Robin Buell, C., Helgeson, J.P., & Jiang, J. (2003). Gene *RB*
cloned from *Solanum bulbocastanum* confers broad spectrum resistance to potato late blight.
Proceedings of the National Academy of Sciences, 100, 9128–9133.
Song, L., & Endelman, J.B. (2023). Using haplotype and QTL analysis to fix favorable alleles in
diploid potato breeding. *The Plant Genome, e20339.* https://doi.org/10.1002/tpg2.20339
Sorensen, P.L., Christensen, G., Karki, H.S., & Endelman, J.B. (2023). A KASP Marker for the
Potato Late Blight Resistance Gene *RB/Rpi-blb1*. *American Journal of Potato Research*,
100, 240–246. https://doi.org/10.1007/s12230-023-09914-6
Swarts, K., Li, H., Romero Navarro, J.A., An, D., Romay, M.C., Hearne, S., Acharya, C.,
Glaubitz, J.C., Mitchell, S., Elshire, R.J., Buckler, E.S., & Bradbury, P.J. (2014). Novel
Methods to Optimize Genotypic Imputation for Low-Coverage, Next-Generation Sequence
Data in Crop Plants. *The Plant Genome*, *7*.
https://doi.org/10.3835/plantgenome2014.05.0023
Uitdewilligen, J.G.A.M.L., Wolters, A.A., Bjorn, B.D., Borm, T.J.A., Visser, R.G.F., & Eck,
H.J. Van. (2013). A Next-Generation Sequencing Method for Genotyping- by-Sequencing
of Highly Heterozygous Autotetraploid Potato, *PLoS ONE, 8*, e0141940.
https://doi.org/10.1371/journal.pone.0062355
Vaillancourt, B., & Buell, C.R. (2019). High molecular weight DNA isolation method from
diverse plant species for use with Oxford Nanopore sequencing. *bioRxiv*, 783159.
https://doi.org/10.1101/783159.
Voorrips, R.E., Gort, G., & Vosman, B. (2011). Genotype calling in tetraploid species from bi-
allelic marker data using mixture models.. *BMC Bioinformatics*, *12*, 172.
https://doi.org/10.1186/1471-2105-12-172
Vos, P.G., Uitdewilligen, J.G.A.M.L., Voorrips, R.E., Visser, R.G.F., & van Eck, H.J. (2015).
Development and analysis of a 20K SNP array for potato (Solanum tuberosum): an insight
into the breeding history. *Theoretical and Applied Genetics*, *128*, 2387–2401.
https://doi.org/10.1007/s00122-015-2593-y
van der Vossen, E., Sikkema, A., Te Lintel Hekkert, B., Gros, J., Stevens, P., Muskens, M.,
Wouters, D., Pereira, A., Stiekema, W., & Allefs, S. (2003). An ancient R gene from the
wild potato species Solanum bulbocastanum confers broad-spectrum resistance to
Phytophthora infestans in cultivated potato and tomato. *Plant Journal*, *36*, 867–882.
https://doi.org/10.1046/j.1365-313X.2003.01934.x
van Eck, H.J., Oortwijn, M.E.P., Terpstra, I.R., van Lieshout, N.H.M., van der Knaap, E.,
Willemsen, J.H., & Bachem, C.W.B. (2022). Engineering of tuber shape in potato (Solanum
tuberosum) with marker assisted breeding or genetic modification using StOFP20. *Research
Square*, *PREPRINT*. https://doi.org/10.21203/rs.3.rs-1807189/v1
Wang, W., & Yan, J. (2021). Shape-Restricted Regression Splines with R Package splines2.
Journal of Data Science, 498–517. https://doi.org/10.6339/21-JDS1020
Whalen, A., Gorjanc, G., & Hickey, J.M. (2020). AlphaFamImpute: High-accuracy imputation in
full-sib families from genotype-by-sequencing data. *Bioinformatics*, *36*, 4369–4371.
https://doi.org/10.1093/bioinformatics/btaa499
Whalen, A., Ros-Freixedes, R., Wilson, D.L., Gorjanc, G., & Hickey, J.M. (2018). Hybrid
peeling for fast and accurate calling, phasing, and imputation with sequence data of any
coverage in pedigrees. *Genetics Selection Evolution*, *50*. https://doi.org/10.1186/s12711-
018-0438-2

Wu, P.Y., Stich, B., Renner, J., Muders, K., Prigge, V., & van Inghelandt, D. (2023). Optimal
implementation of genomic selection in clone breeding programs—Exemplified in potato: I.
Effect of selection strategy, implementation stage, and selection intensity on short-term
genetic gain. *Plant Genome*, *16*. https://doi.org/10.1002/tpg2.20327
Wu, S., Zhang, B., Keyhaninejad, N., Rodríguez, G.R., Kim, H.J., Chakrabarti, M., Illa-
Berenguer, E., Taitano, N.K., Gonzalo, M.J., Díaz, A., Pan, Y., Leisner, C.P., Halterman, D.,
Buell, C.R., Weng, Y., Jansky, S.H., van Eck, H., Willemsen, J., Monforte, A.J., Meulia, T.,
& van der Knaap, E. (2018). A common genetic mechanism underlies morphological
diversity in fruits and other plant organs. *Nature Communications*, *9*.
https://doi.org/10.1038/s41467-018-07216-8
Zhao, D., Mejia-Guerra, K.M., Mollinari, M., Samac, D., Irish, B., Heller-Uszynska, K., Beil,
C.T., & Sheehan, M.J. (2023). A public mid-density genotyping platform for alfalfa
(Medicago sativa L.). *Genetic Resources*, *4*, 55–63.
https://doi.org/10.46265/genresj.EMOR6509
Zheng, C., Amadeu, R.R., Munoz, P.R., & Endelman, J.B. (2021). Haplotype reconstruction in
connected tetraploid F1 populations. *Genetics*, *219*. https://doi.org/10.1093/genetics/iyab106
Zych, K., Gort, G., Maliepaard, C.A., Jansen, R.C., & Voorrips, R.E. (2019). FitTetra 2.0 -
Improved genotype calling for tetraploids with multiple population and parental data
support. *BMC Bioinformatics*, *20*. https://doi.org/10.1186/s12859-019-2703-y

Figure 1. Total depth per sample, in million (M) read counts, for three submissions of potato V1 DArTag.

Figure 2. (A) Distribution of the mean sample depth (DP) for V1 DArTag markers. (B) Log-log plot of the relationship between the standard deviation and mean for sample DP. Individual marker points are shown only for 2021 to maintain legibility. Combining the data across years, the overall regression line (not shown) has slope 0.79 (SE 0.00) and $R^2 = 0.99$.

Figure 3. Examples of DArTag markers without (A) vs. with (B) allelic bias. Dashed lines correspond to possible tetraploid allele ratios when there is no allelic bias (1:0, 3:1, 1:1, 1:3, 0:1). (A) solcap_snp_c2_36615 with bias = -0.2. (B) PotVar0072076 with bias = 1.8.

Figure 4. Empirical cumulative distribution for the error between the V1 DArTag and SNP array on 1865 common markers. CE = classification error. RMSE = root-mean-squared-error. 2x = pseudo-diploid genotypes. 4x = tetraploid genotypes.

Figure 5. Violin plot illustrating ploidy discrimination with the V2 DArTag assay, based on the proportion of simplex or triplex markers with a tetraploid genotype calling model. The diploid potato samples had much lower values of this parameter.

Figure 6. Original vs. corrected genotypes for the trait markers Ryadg_chr11_2499502 and H1_chr05_52349069. The original genotypes were based on R/updog with a "norm" prior and then corrected based on linkage analysis.

A
```
     CDF1.4_chr05_4488021|Ref   GAAGCGGCTAAAAGCTCTATATGGTCAACACTAG-------GTAT
  CDF1.4_chr05_4488021|Other1   GAAGCGGCTAAAAGCTCTATATGGTCCACACTAG-------GTAT
  CDF1.4_chr05_4488021|Other2   GATGCGGCTAAAAGCTCTATATGGTCAACACTAG-------GTAT
     CDF1.4_chr05_4488021|Alt   GAAGCGGCTAAAAGCTCTATATGGTCAACACTAGGTATCCCGTAT
  CDF1.4_chr05_4488021|Other3   GAAGCGGCTAAAAGCTCTATATGGTCAACACTAGTCACTAGGTAT
```

B

	CDF1.1 Count	CDF1.2 Count	CDF1.4 Count	CDF1 Genotype
Atlantic	23	0	34	1/3/4/4
W13069-5-DH088	87	0	64	1/4
W14NYQ29-5-DH024	77	0	0	1/3
RioColorado-DH005	122	81	0	1/2
W9968-5-DH151	0	18	0	2/3
W2x001-22-45	0	0	0	3/3

Figure 7. (A) Multiple sequence alignment of the DArTag haplotypes discovered for trait marker CDF1.4_chr05_448021. Haplotypes Ref, Other1, Other2 are CDF1.1 alleles, while Alt is CDF1.4 and Other3 is CDF1.2. (B) Haplotype read counts for samples with known CDF1 genotype.

	OFP20 Genotype	AF Allele 1	AF Allele 8	AD Alleles 2,3+7
CO99076-6R-DH002	1/1	1.00	0.00	0,0
W14NYQ29-5-DH024	3/3	0.00	0.00	0,240
W14NYQ9-2-DH137	1/3	0.49	0.00	0,115
W8890-1R-DH002	1/7	0.64	0.00	0,70
W9968-5-DH084	1/8	0.44	0.60	0,1

Figure 8. (A) Distribution of sample allele frequencies for OFP20.1 in round chip (N=300) vs. long russet (N=21) germplasm. (B) Comparison of known OFP20 genotypes with V2 DArT markers. Allele frequency (AF) of OFP20.1 was approximated by ALT frequency at marker OFP20_M6_CDS_994. AF of OFP20.8 was approximated by REF frequency at marker OFP20_M6_CDS_24. Allele depth (AD) at OFP20_M6_CDS_171 was used to distinguish allele 2 (ALT) from alleles 3 and 7 (REF).

Supplemental Figures and Tables

Endelman *et al.* Targeted genotyping-by-sequencing of potato and data analysis with R/polyBreedR

Figure S1. Marey Map of the potato genome. Horizontal axis is the DMv6.1 reference genome position (Pham et al., 2020).

Figure S2. Distribution of the 2501 genomic markers for V1 DArTag.

Figure S3. Distribution of allele bias (AB) estimates, where AB=1 indicates no bias, and values greater than 1 indicate bias toward the REF allele

SNP array

GBS

Figure S4. Hierarchical clustering based on SNP array (top) or GBS (bottom) data. Both platforms identified two groups of genetically identical clones, one pair and one threesome, originating from the same F1 populations

Figure S5. Joint clustering of SNP array (blue) and GBS (orange) samples. The two marker profiles for every clone were paired.

Figure S6. Empirical cumulative distribution for 2x classification error for 2608 common markers between the V4 SNP array and V2 DArTag.

Figure S7. Five-parent partial diallel. Graphical output from PolyOrigin shows the number of progeny per biparental F1 population.

Figure S8. Additive effect estimates for parental haplotypes in the five-parent partial diallel. Positive values indicate presence of the R gene. From left to right, the result indicates the parental dosage of Ryadg is 0, 1, 1. 2, 1, and for H1 the parental dosage is 1, 2, 0, 2, 1. Parents W14NYQ9-2 and W15NYR11-13 were included in the V2 DArTag submission, and their genotype calls agree with these predictions (FileS3).

Table S1. Comparison of KASP (rows) vs. V1 DArTag (columns) markers targeting *Ry$_{adg}$* (snpST00073).

	Absent	Present
Absent	7	1
Present	1	84

Table S2. Positive samples for marker Rysto_chr12_2352742 in V2 DArTag.

id	mother	father
W17079-16rus	Payette Russet	AW07791-2rus
W17081-2rus	Payette Russet	W9742-3rus
A12304-1sto	A96953-13sto	Clearwater Russet
W6511-1R	Kankan	W2275-9R

Table S3. Results for marker Sli_chr12_2372490 in V2 DArTag.

id	ALT dosage
W9968-5-DH027	0
CO99076-6R-DH002	0
CO99076-6R-DH033	0
W14NYQ29-5-DH024	0
W14NYQ9-2-DH119	0
W14NYQ9-2-DH132	0
W8890-1R-DH003	0
W9968-5-DH084	0
W9968-5-DH022	0
W13069-5-DH088	0
RioColorado-DH003	0
W9968-5-DH151	0
W12078-76-DH352	0
W12078-76-DH099	0
W13058-4-DH002	0
W14NYQ9-2-DH137	0
W14NYQ9-2-DH146	0
RioColorado-DH005	1
W15263-50R-DH001	1
W15263-50R-DH011	1
W8890-1R-DH002	1
W9426-3R-DH005	1
W9426-3R-DH037	1
W2x150-24	1
W2x113-3	1
W2x001-22-45	2
W2x082-(14/20)-13	2
W2x082-(14/20)-13-2	2

Supplemental Methods

Endelman et al. (2024)

This document shows how the main figures and tables were generated on a MacOS system. R packages polyBreedR and diaQTL can be installed from their GitHub sites. Other R packages are available on CRAN. The `impute_LA` function utilizes PolyOrigin, which is available from its GitHub site and requires installation of Julia. Command line calls to bcftools are used to manipulate VCF files.

```r
library(vcfR)
```

```
##
##    *****       ***   vcfR   ***       *****
##    This is vcfR 1.15.0
##      browseVignettes('vcfR') # Documentation
##      citation('vcfR') # Citation
##    *****       *****     *****       *****
```

```r
library(polyBreedR)
library(ggplot2)
library(tidyr)
library(ggpubr)
library(diaQTL)
library(data.table)
library(stringr)
```

Figure 1

```r
data1 <- read.vcfR("FileS1-DArTagV1.vcf.gz",verbose=F)
id1 <- colnames(data1@gt)[-1]
m1 <- nrow(data1@gt) #2503

DP <- extract.gt(data1,element="DP",as.numeric=T)
metadata <- read.csv("FileS2-V1metadata.csv")
metadata <- metadata[match(id1,metadata$id),]

pdat <- metadata
pdat$year <- factor(pdat$year)
pdat$DP <- apply(DP,2,sum)/1e6
ggplot(pdat,aes(x=DP,fill=year)) + geom_density(alpha=0.5) +
  theme_bw() + scale_fill_brewer(palette="Set1",name="") +
  scale_x_continuous(name="Total Depth per Sample (M)",breaks=seq(0,1.4,by=0.2),
                     label=seq(0,1.4,by=0.2))
```

Figure 2

```r
mDP <- t(apply(DP,1,function(z){tapply(z,metadata$year,mean)}))
ix <- which(apply(mDP,1,mean) > 1)
pdat <- pivot_longer(data.frame(marker=rownames(DP)[ix],mDP[ix,]),2:4)
pdat$year <- factor(gsub("X","",pdat$name),
                    levels=as.character(2020:2022),ordered=T)

p1 <- ggplot(pdat,aes(x=value,fill=year)) + geom_density(alpha=0.5) +
  theme_bw() + scale_x_log10(name="",limits=c(1,1e4)) +
  scale_fill_brewer(palette="Set1",name="")

sDP <- t(apply(DP,1,function(z){tapply(z,metadata$year,sd)}))
tmp <- pivot_longer(data.frame(marker=rownames(DP)[ix],sDP[ix,]),2:4)
pdat$sd <- tmp$value
pdat21 <- pdat[pdat$year=="2021",]

p2 <- ggplot(mapping=aes(x=value,y=sd,colour=year)) +
  geom_point(data=pdat21,shape=1,size=0.5,colour="grey50") +
  stat_smooth(data=pdat,method="lm") +
  scale_color_brewer(name="",palette="Set1",drop=FALSE) +
  geom_abline(slope=0.5,intercept=0,linetype=2) + theme_bw() +
  scale_x_log10(name="Mean Sample DP",limits=c(1,1e4)) +
  scale_y_log10(name="Std. Dev. DP")

ggarrange(p1,p2,common.legend = TRUE,labels=c("A","B"),ncol=1)

## `geom_smooth()` using formula = 'y ~ x'
```

```r
#Regression model
data2 <- data.frame(sd=log10(pdat$sd),
                    mu=log10(pdat$value))
ix <- which(data2$mu > 1 & data2$mu < 4)
summary(lm(sd~mu-1,data2[ix,]))
```

```
##
## Call:
## lm(formula = sd ~ mu - 1, data = data2[ix, ])
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.36268 -0.10602  0.01168  0.10615  0.56770
##
## Coefficients:
##     Estimate Std. Error t value Pr(>|t|)
## mu 0.7885795  0.0007635    1033   <2e-16 ***
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.1462 on 7226 degrees of freedom
## Multiple R-squared:  0.9933, Adjusted R-squared:  0.9933
## F-statistic: 1.067e+06 on 1 and 7226 DF,  p-value: < 2.2e-16
```

Figure 3

```r
AB <- extract.info(data1,element="AB",as.numeric=T)
logAB <- log(AB)/log(2)

GT <- extract.gt(data1,element="GT")
geno <- GT2DS(GT,n.core=2)

AD <- extract.gt(data1,element="AD")
REF <- t(apply(AD,1,function(z){as.numeric(sapply(strsplit(z,split=","),"[[",1))}))
ALT <- t(apply(AD,1,function(z){as.numeric(sapply(strsplit(z,split=","),"[[",2))}))
markers <- rownames(AD)

#panel A
k=1
p1 <- ggplot(data.frame(x=REF[1,],y=ALT[1,],dosage=factor(geno[1,])),aes(x=x,y=y,colour=dosage)) +
  geom_point()  + theme(text=element_text(size=14)) + xlab("REF") + ylab("ALT") +
  geom_abline(intercept=0,slope=1,linetype=2) + geom_abline(intercept=0,slope=3,linetype=2) +
  geom_abline(intercept=0,slope=1/3,linetype=2) + geom_hline(yintercept=0,linetype=2) +
  geom_vline(xintercept=0,linetype=2)

#panel B
k=8
p2 <- ggplot(data.frame(x=REF[k,],y=ALT[k,],dosage=factor(geno[k,])),aes(x=x,y=y,colour=dosage)) +
  geom_point() + theme(text=element_text(size=14)) + xlab("REF") + ylab("ALT") +
  geom_abline(intercept=0,slope=1,linetype=2) + geom_abline(intercept=0,slope=3,linetype=2) +
  geom_abline(intercept=0,slope=1/3,linetype=2) + geom_hline(yintercept=0,linetype=2) +
  geom_vline(xintercept=0,linetype=2)

ggarrange(p1,p2,ncol=1,common.legend = TRUE,labels = c("A","B"),vjust = 0.8)
```

Figure 4

```
#Array comparison
system("bcftools isec -p dir1 -Oz FileS1-DArTagV1.vcf.gz FileS5-SNParrayV4.vcf.gz")

compare1 <- read.vcfR("dir1/0002.vcf.gz",verbose=F)
compare2 <- read.vcfR("dir1/0003.vcf.gz",verbose=F)
GT1 <- extract.gt(compare1,element="GT")
GT2 <- extract.gt(compare2,element="GT")
id <- intersect(colnames(GT1),colnames(GT2))
geno.dart <- GT2DS(GT1[,id],n.core=2)
geno.dart.2x <- GT2DS(GT1[,id],n.core=2,diploidize=T)
geno.array <- GT2DS(GT2[,id],n.core=2)
geno.array.2x <- GT2DS(GT2[,id],n.core=2,diploidize=T)

m <- nrow(geno.array)
V1error <- data.frame(compare1@fix[,c("CHROM","POS","ID")],RMSE.4x=numeric(m),
```

```
                    CE.4x=numeric(m),CE.2x=numeric(m))
for (i in 1:m) {
  V1error$RMSE.4x[i] <- sqrt(mean((geno.array[i,]-geno.dart[i,])^2,na.rm=T))
  V1error$CE.4x[i] <- mean(as.integer(geno.array[i,]!=geno.dart[i,]))
  V1error$CE.2x[i] <- mean(as.integer(geno.array.2x[i,]!=geno.dart.2x[i,]))
}
f1 <- ecdf(V1error$RMSE.4x)
f2 <- ecdf(V1error$CE.4x)
f3 <- ecdf(V1error$CE.2x)

x <- seq(0,1,by=0.01)
m <- length(x)
pdat <- data.frame(Type=factor(rep(c("2x CE","4x CE","4x RMSE"),each=m)),
                   x=x,
                   y=round(c(f3(x),f2(x),f1(x))*1865))
ggplot(data=pdat,aes(x=x,y=y,colour=Type)) + geom_line() +
  theme_bw() + theme(text=element_text(size=13)) + xlab("Error") +
  ylab("# Markers") + theme(legend.title = element_blank()) +
  theme(legend.position="inside",legend.position.inside = c(0.7,0.5)) +
  scale_x_continuous(breaks=seq(0,1,by=0.1),labels=seq(0,1,by=0.1))
```

```
#Number of markers passing thresholds
sum(V1error$CE.2x <= 0.1,na.rm=T)
```

```
## [1] 1649
```

```
sum(V1error$CE.4x <= 0.1,na.rm=T)
```

```
## [1] 1302
```

```
sum(V1error$RMSE.4x <= 0.5,na.rm=T)
```

```
## [1] 1547
```

Figure 5

```
#apply tetraploid genotoype calling model to all samples
gbs(in.file="FileS3-DArTagV2.vcf.gz",out.file="tmp.vcf.gz",
    ploidy=4,model.fit=FALSE)
```

```
## Progress: 1000 markers
## Progress: 2000 markers
## Progress: 3000 markers
## Progress: 3915 markers
```

```
#get ploidy from original file
data <- read.vcfR("FileS3-DArTagV2.vcf.gz",verbose=F)
GT <- extract.gt(data)
ploidy <- sapply(gregexpr("/",GT[2,]),length)+1
table(ploidy)
```

```
## ploidy
##   2   4
##  52 323
```

```
#convert to bgzip and eliminate scaffolds not in DM reference genome
system("gunzip tmp.vcf.gz -c | bgzip -c - > tmp2.vcf.gz")
system("bcftools index tmp2.vcf.gz")
chr <- paste(paste0("chr",str_pad(1:12,width=2,pad="0")),collapse=",")
system(sub("CHR",chr,"bcftools view -r CHR tmp2.vcf.gz -Oz > check_ploidy.vcf.gz"))

ans <- check_ploidy(vcf.file = "check_ploidy.vcf.gz")
pdat <- data.frame(Ploidy=factor(ploidy), y=apply(ans$mat,1,max))
ggplot(pdat,aes(x=Ploidy,y=y)) + geom_violin() + theme_bw() +
  ylab("Proportion Simplex or Triplex Markers")
```

Table 1

```r
#subset to good markers
ix <- which(V1error$RMSE.4x <= 0.5 & V1error$CE.2x <= 0.1)
fwrite(V1error[ix,c("CHROM","POS")],sep="\t",col.names=F,file="mark.txt")

ped <- read.csv("FileS6-W19diallel_pedigree.csv")
progeny.id <- ped$id[ped$pop > 0]
fwrite(data.frame(progeny.id),sep="\t",col.names = F,file="id.txt")

#subset VCF to diallel population
system("bcftools view -S id.txt -R mark.txt FileS1-DArTagV1.vcf.gz -Oz > dia.vcf.gz")
```

```r
phased_parents <- read.csv("FileS7-W19diallel_phased_parents.csv")
marks <- phased_parents$marker
array <- read.vcfR("FileS5-SNParrayV4.vcf.gz",verbose=FALSE)
geno.array <- GT2DS(extract.gt(array)[marks,progeny.id],n.core=2)

RMSE <- data.frame(ped[match(1:10,ped$pop),2:4],RF=numeric(10),LA=numeric(10))
for (i in 1:10) {
  ix <- which(ped$pop==i)
  VP <- ped$id[ix]
  ans <- impute_L2H(high.file = "FileS5-SNParrayV4.vcf.gz",
                    low.file="W19diallel_low.vcf.gz",
                    out.file = "dia_RF.csv",
                    params=list(format="GT",n.tree=100,n.mark=25,exclude=VP),
                    n.core=2)
  tmp <- read.csv("dia_RF.csv",as.is=T,check.names=F)
  imputed <- as.matrix(tmp[,-(1:3)])
```

```
  rownames(imputed) <- tmp$marker
  RMSE$RF[i] <- sqrt(mean((geno.array[,VP]-imputed[marks,VP])^2,na.rm=T))
}
```

```
## Training Set N=77
## Training Set N=82
## Training Set N=84
## Training Set N=78
## Training Set N=76
## Training Set N=76
## Training Set N=74
## Training Set N=73
## Training Set N=65
## Training Set N=80
```

```
impute_LA(high.file="FileS7-W19diallel_phased_parents.csv",
          low.file="dia.vcf.gz",low.format="GT",
          ped.file="FileS6-W19diallel_pedigree.csv",out.file="dia_LA.csv")

imputed <- as.matrix(read.csv("W19dia_LA.csv",as.is=T,check.names=F)[,-(1:3)])
for (i in 1:10) {
  ix <- which(ped$pop==i)
  VP <- ped$id[ix]
  RMSE$LA[i] <- sqrt(mean((geno.array[,VP]-imputed[,VP])^2,na.rm=T))
}

knitr::kable(RMSE,digits=2)
```

	pop	mother	father	RF	LA
6	1	W12078-76	W13NYP102-7	0.45	0.17
14	2	W12078-76	W6609-3	0.43	0.14
17	3	W14NYQ4-1	W12078-76	0.37	0.14
18	4	W14NYQ4-1	W13NYP102-7	0.41	0.13
25	5	W14NYQ4-1	W6609-3	0.42	0.13
34	6	W14NYQ9-2	W12078-76	0.44	0.16
43	7	W14NYQ9-2	W13NYP102-7	0.40	0.14
54	8	W14NYQ9-2	W14NYQ4-1	0.42	0.15
66	9	W14NYQ9-2	W6609-3	0.43	0.16
86	10	W6609-3	W13NYP102-7	0.40	0.15

Figure 6

```
data <- read_data(genofile="FileS9-W21diallel_genofile.csv",ploidy=4L,
                  pedfile="FileS8-W21diallel_ped.csv",
                  phenofile="FileS10-W21diallel_trait.csv",dominance = 2,n.core=2)
```

```
## 123 individuals with pedigree and genotype data
## 123 individuals with pedigree, genotype and phenotype data
## Preparing genotype data...
## Preparing for polygenic effects...
```

```
v2 <- read.vcfR("FileS3-DArTagV2.vcf.gz",verbose=F)
geno <- GT2DS(extract.gt(v2,"GT")[c("Ryadg_chr11_2499502","H1_chr05_52349069"),],
```

```
            n.core=2)

H1 <- haplo_get(data,marker="PotVar0123022")
H1.1 <- apply(H1[,c(2,6,7,15,16,18)],1,sum)
table(factor(geno["H1_chr05_52349069",names(H1.1)],levels=0:4),factor(H1.1,levels=0:4))
```

```
##
##      0  1  2  3  4
##   0 23  0  0  0  0
##   1  0 61  3  0  0
##   2  0 10 21  3  0
##   3  0  0  2  0  0
##   4  0  0  0  0  0
```

```
Ry <- haplo_get(data,marker="solcap_snp_c1_4322")
Ry.1 <- round(apply(Ry[,c(7,11,14,15,20)],1,sum))
table(factor(geno["Ryadg_chr11_2499502",names(Ry.1)],levels=0:4),factor(Ry.1,levels=0:4))
```

```
##
##      0  1  2  3  4
##   0 24  0  0  0  0
##   1  0 60  2  0  0
##   2  0 23  7  0  0
##   3  0  0  5  0  0
##   4  0  0  2  0  0
```

Figure 7

```
ans <- madc("FileS4-DArTagV2_madc.csv",marker="CDF1")
id <- c("Atlantic","W13069-5-DH088","W14NYQ29-5-DH024","RioColorado-DH005",
        "W9968-5-DH151","W2x001-22-45")
knitr::kable(ans[id,])
```

	CDF1.1	CDF1.2T	CDF1.2C	CDF1.4
Atlantic	23	0	0	34
W13069-5-DH088	87	0	0	64
W14NYQ29-5-DH024	77	0	0	0
RioColorado-DH005	122	81	0	0
W9968-5-DH151	0	18	0	0
W2x001-22-45	0	0	0	0

Figure 8

```
ans <- madc("FileS4-DArTagV2_madc.csv",marker="OFP20")
id <- c("CO99076-6R-DH002","W14NYQ29-5-DH024","W14NYQ9-2-DH137","W8890-1R-DH002",
        "W9968-5-DH084")
knitr::kable(ans[id,])
```

	AF1	AF8	AD2	AD3&7
CO99076-6R-DH002	1.00	0.0	0	0
W14NYQ29-5-DH024	0.00	0.0	0	240
W14NYQ9-2-DH137	0.49	0.0	0	115

	AF1	AF8	AD2	AD3&7
W8890-1R-DH002	0.64	0.0	0	70
W9968-5-DH084	0.44	0.6	0	1