Tuft cells transdifferentiate to neural-like progenitor cells in the progression of pancreatic

cancer.

Daniel J. Salas-Escabillas[1,2]; Megan T. Hoffman, PhD[3]; Jacee S. Moore, MS[2]; Sydney M. Brender,

MS[2]; Hui-Ju Wen, PhD[2]; Simone Benitz, PhD[2]; Erick T. Davis[2]; Dan Long[2]; Allison M. Wombwell[2];

Nina G. Steele, PhD[2]; Rosalie C. Sears, PhD[4]; Ichiro Matsumoto, PhD[5]; Kathleen E. DelGiorno, PhD[6];

Howard C. Crawford, PhD* [2, 7]

[1] Cancer Biology, University of Michigan, Ann Arbor, MI

[2] Department of Surgery, Henry Ford Health, Detroit, MI

[3] Department of Immunology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA

[4] Department of Molecular and Medical Genetics, Oregon Health & Science University, Portland, OR

[5]Monell Chemical Senses Center, Philadelphia, PA

[6]Department of Cell and Developmental Biology, Vanderbilt University School of Medicine, Nashville,

TN

[7] Cancer Biology Program, Wayne State University, Detroit, MI

*Corresponding Author:

Howard C. Crawford, PhD, Department of Surgery, Henry Ford Health System, Detroit, Michigan,

USA

Email: hcrawfo1@hfhs.org

Summary

Pancreatic ductal adenocarcinoma (PDA) is partly initiated through the transdifferentiation of acinar cells to metaplastic ducts that act as precursors of neoplasia and cancer. Tuft cells are solitary chemosensory cells not found in the normal pancreas but arise in metaplasia and neoplasia, diminishing as neoplastic lesions progress to carcinoma. Metaplastic tuft cells (mTCs) function to suppress tumor progression through communication with the tumor microenvironment, but their fate during progression is unknown. To determine the fate of mTCs during PDA progression, we have created a lineage tracing model that uses a tamoxifen-inducible tuft-cell specific Pou2f3$^{CreERT/+}$ driver to induce transgene expression, including the lineage tracer tdTomato or the oncogene *Myc*. mTC lineage trace models of pancreatic neoplasia and carcinoma were used to follow mTC fate. We found that mTCs, in the carcinoma model, transdifferentiate into neural-like progenitor cells (NRPs), a cell type associated with poor survival in PDA patients. Using conditional knock-out and overexpression systems, we found that *Myc* activity in mTCs is necessary and sufficient to induce this Tuft-to-Neuroendocrine-Transition (TNT).

Introduction

Pancreatic ductal adenocarcinoma (PDA), the most common form of pancreatic cancer, is a devastating disease with a 5-year survival rate of only 13% [1]. Effective treatment options are few, and reliable early detection methods needed to enhance treatment effectiveness are non-existent. PDA is initiated by oncogenic mutations in the *KRAS* gene, causing irreversible transdifferentiation of normal acinar cells into metaplastic ducts. Metaplastic ducts then progress to pancreatic intraepithelial neoplasia (PanIN) [2] and, with the acquisition of mutations in tumor suppressor genes such as *TP53*, *CDKN2A*, or *SMAD4* [3, 4], progress to invasive carcinoma. Metaplastic ducts are a heterogeneous population of reactive epithelial cells [5] comprised of at least 6 distinct identities: gastric neck cells [6], gastric pit-like cells, gastric chief-like cells, ductal senescent cells, metaplastic neuroendocrine cells (mNECs), and specialized chemosensory cells known as metaplastic tuft cells (mTCs) [7].

Normal tuft cells are solitary chemosensory cells found throughout the body that sense and respond to environmental cues to maintain tissue homeostasis. Functionally, they use the canonical gustatory signal transduction pathway, with specific stimulants and responses depending on context and the organ in which they reside [8-10]. In pancreatic neoplasia, metaplastic tuft cells (hereafter referred to as mTCs) have been proposed to be a class of tumor progenitor cells [9, 11]. Still, functionally, they suppress progression to PDA via inhibition of anti-inflammatory immune cells through crosstalk with the tumor microenvironment [8, 10]. mTCs are identified by staining for markers such as Acetylated-*Tuba1a*[10], *Dclk1*, *Vav1*, *Cox1*[9], *Cox2*[12-14], *Trpm5* [15], and *Pou2f3*, the latter being the master transcriptional regulator of tuft cell genesis [15, 16]. mTCs are not typically found in the normal pancreas and only appear after prolonged damage, such as in chronic pancreatitis [17] or the initiation of neoplasia [9, 11], arising from the transdifferentiation of acinar cells. As neoplasia progresses to

invasive carcinoma, the mTC population gradually disappears [9, 11], challenging their putative

role as classic tumor progenitor cells.

In the current study, to address the hypothesis that mTCs act as tumor progenitor cells

in pancreatic cancer, we lineage-traced their fate during disease progression in mouse

models of pancreatic disease. To accomplish this, we used a dual recombinase model

wherein tumorigenesis is initiated by the expression of FlpO recombinase knocked into the

pancreas-specific *Ptf1a* locus, driving the expression of an Frt-STOP-Frt oncogenic *Kras*$^{G12D/+}$

allele and the deletion an Frt-flanked allele of *Trp53* [18]. This model allowed us to use an

inducible *Pou2f3*$^{CreERT/+}$ driver [19] to target the expression of a *ROSA26*$^{LSL-TdTomato}$ lineage

trace reporter specifically to mTCs [16, 20] in both neoplasia and carcinoma, indelibly labeling

mTCs and their progeny. In the *Ptf1a*$^{FlpO/+}$*; Kras*^{G12D} neoplasia model, mTCs and other

metaplastic cellular subtypes, including neuroendocrine-like enteroendocrine cells [7, 21, 22]

arise from acinar cells, independent from one another. However, in the *Ptf1a*$^{FlpO/+}$*; Kras*$^{FSF-}$

$^{G12D/+}$*; Trp53*$^{Frt-Exons\ 2\ to\ 5-Frt/+}$ carcinoma model, we found that acinar cell-derived mTCs

transdifferentiate to neural-like progenitor cells (NRPs) [21], also known as ductal

neuroendocrine cells [23]. NRPs in PDA patients correlate with worse survival, are resistant to

chemotherapy, and their genesis is associated with elevated *MYC* activity [21, 23]. Indeed,

modulating the expression of *Myc* to mTCs demonstrated that *Myc* activity was necessary

and sufficient to drive the transdifferentiation of mTCs to NRPs.

RESULTS

Pou2f3-CreERT specifically targets recombination in metaplastic tuft cells in the pancreas.

Tuft cells are not found in the normal mouse pancreas but arise from the transdifferentiation of acinar cells in conditions of tissue damage and early neoplasia [24]. As neoplasia progresses, metaplastic tuft cells (mTCs) decrease in number and are absent in carcinoma [9, 11, 24]. To determine the fate of mTCs during neoplastic progression, we developed a novel dual recombinase lineage trace system, wherein Ptf1a$^{Flpo/+}$ [25] initiates expression of oncogenic Kras from a Kras$^{Frt-Stop-Frt-G12D}$ allele (*KF*) to drive neoplasia [18] (Fig. 1A, 1D). When combined with a conditional knock-out trp53 allele with exons 2-5 flanked by Frt recombination sites (Trp53$^{Frt-Exons\ 2\ to\ 5-Frt/+}$), lesions rapidly progress from neoplasia to carcinoma (*KPF*) (Fig. 1B, 1D). Within this system, we introduced a Cre^{ERT} driver knocked into the *Pou2f3* locus [19], which encodes the transcription factor master regulator of tuft cell genesis [15, 16, 20, 26, 27]. *Pou2f3$^{CreERT/+}$* was used to drive recombination and expression of a *ROSA26$^{LSL-tdTomato}$* reporter (tdTom) [28] in a tamoxifen-inducible manner (P2f3T) [8, 20, 27], allowing us to lineage trace mTCs temporally in our neoplasia (*KF-P2f3T)* and carcinoma (*KPF-P2f3T)* models.

To determine the fidelity of our systems in labeling tuft cells, we treated *KF-P2f3T* with cerulein at 8-9 weeks of age to induce mild pancreatitis and accelerate transformation, then allowed the mice to recover for 3 weeks to optimize mTC genesis [9, 11]. The *KPF-P2f3T* carcinoma model induced sufficient transformation and mTC formation at 11-12 weeks of age with no additional treatment (Fig. 1C). Cerulein-treated *KF-P2f3T* and 11–12-week-old *KPF-P2f3T* mice were treated with tamoxifen to induce tdTom reporter activity, and the pancreata

harvested 1 day later. We co-stained tissues for tdTom and the tuft cell markers Pou2f3,

Acetylated-α-tubulin, Cox1, Dclk1, Trpm5, and Vav1 (Fig. 1E and Supp. Fig. 1A, 1B) by

immunofluorescence (IF). In both models, we found co-expression of tdTom exclusively in

cells expressing mTC markers (Supp. Fig. 1A, 1B), confirming the system's specificity and

allowing us to trace mTC fate throughout the progression of PDA.

Metaplastic tuft cells change identity in pancreatic adenocarcinoma.

To determine the fate of the mTC population over time, we collected tissue from our

KF-P2f3T and *KPF-P2f3T* mice 3 days and 7 weeks post-tamoxifen. In the *KF-P2f3T*

neoplasia model, all tdTom[+] cells co-expressed tuft cell markers (Fig. 2A). The population of

mTCs dropped off slightly over this period but largely remained stable (Fig. 2B, *Black*). In

contrast, in the *KPF-P2f3T* carcinoma model, the mTC population dropped drastically by 7

weeks, consistent with their disappearance as PanIN progresses to PDA [9, 11] (Fig. 2A and 2B,

Black). We found no evidence that mTC attrition was associated with apoptosis by co-

staining for cleaved caspase 3 (CC3) (Fig. 1F). Instead, we noted that tdTom[+] cells persisted

in this model but no longer expressed mTC markers (Fig. 2A), suggesting that mTCs take on

a new cellular identity.

Quantitation of the tdTom expression throughout multiple pancreata revealed that, in

the *KPF-P2f3T* carcinoma model, the Cox1[+] mTC population dropped from an average of

15% at 3 days post-tamoxifen to an average of 3% at 7 weeks post-tamoxifen (Fig. 2B,

Black). In contrast, the mTC progeny population, defined as tdTom[+] cells not expressing

Cox1, increased by ~5-fold from an average of 4% of pancreatic ducts at 3 days post-

tamoxifen to an average of nearly 17% at 7 weeks post-tamoxifen, reflecting the degree of

mTC loss (Fig. 2B, *Pink*). These data support mTCs adopting a new identity in carcinomas.

125 **Transdifferentiation of mTCs to metaplastic neuroendocrine cells.**

126   To investigate the identity of the mTC progeny, we co-stained for tdTom together with

127 markers of the known identities of other metaplastic ductal cells defined by *Schlesinger et al.*

128 and *Benitz et al.*, including gastric pit-like (Gkn1$^+$), gastric neck-like (Lectin GS-II$^+$) [29], and

129 metaplastic neuroendocrine cells (Syp$^+$) [7, 9, 17].

130   The lack of co-expression of tdTom with Gkn1 or Lectin GS-II suggested that *KPF-*

131 *P2f3T* tdTom$^+$ mTC progeny had not taken on gastric pit-like cell or gastric neck-like cell

132 identities (Fig. 2C and Supp. Fig. 2A). Instead, mTC progeny stained positive for Syp,

133 revealing the acquisition of a metaplastic neuroendocrine cell (mNEC) identity (Fig. 2C,

134 *Yellow arrows*). To determine if tdTom expression in cells with a neuroendocrine phenotype

135 could be due to previously unnoticed recombination in pancreatic islets, we confirmed no co-

136 localization of tdTom co-staining with Insulin (Ins) or in surrounding islet cells (Supp. Fig. 2B).

137 Quantitation of the different time points for the *KPF-P2f3T* model confirmed a decrease of

138 tdTom$^+$/Cox1$^+$ mTCs, (Fig. 2E, *Black*), with a concomitant ~4-fold increase of tdTom$^+$/Syp$^+$

139 mNECs, (Fig. 2E, *Pink*) by 7 weeks post-tamoxifen (Fig. 2D, 2E). Interestingly, while there is

140 a mNEC population in the metaplastic ducts in the *KF-P2f3T* neoplasia model, none

141 expressed tdTom, suggesting that they did not derive from mTCs (Fig. 2D, 2E, *Green*) and

142 had a distinct origin from those in the *KPF-P2f3T* mice.

143

mNECs depend on the mTC population to develop in PDA but not PanIN models.

Lineage tracing experiments show that mNECs in both neoplasia and carcinoma

models are initially derived from acinar cells [9] (Supp. Fig 2C) but differ in that carcinoma

mNECs derive from an mTC intermediate. As an independent test of these divergent origins,

we took advantage of our *Pou2f3*[CreERT/+] driver being a knock-in to the *Pou2f3* allele, thus

inactivating endogenous Pou2f3 activity [20]. By creating a model homozygous for

Pou2f3[CreERT/CreERT], we effectively ablated *Pou2f3* (Supp. Fig. 3A, 3B), preventing mTC

formation [8] and, therefore, any mTC progeny. Consistent with our previous observations,

Cox1[+] mTC and Syp[+] mNEC populations were easily identifiable in the pancreatic lesions of

our KF; Pou2f3[CreERT/+] neoplasia model. Pou2f3 ablation in these mice eliminated the mTC

population, as expected, while the mNEC population persisted (Fig. 3A), supporting their

independent origins. In contrast, in the KPF; Pou2f3[CreERT/CreERT] carcinoma mice, we found

neither mTCs nor mNECs, confirming a shared dependency on Pou2f3 function (Fig. 3A,

Green arrows).

Lineage tracing and Pou2f3 ablation data in KPF mice support a model where mTCs

give rise to mNECs. However, it was also possible that in the KPF model, both cell types

express *Pou2f3* and require its activity. mNECs do not express *Pou2f3* in *Kras*[G12D]-driven

neoplasia [7, 22, 27] but have not been examined in the KPF carcinoma model. To address this,

we performed co-immunofluorescence for Pou2f3, together with Cox1 (mTCs) and Syp

(mNECs). As expected, Pou2f3 was readily detectable in mTCs in both KF and KPF

pancreata (Fig 1E, 3B). In contrast, we found no examples of Pou2f3[+] mNECs in either

system (Fig. 3B). The lack of identifiable recombination in early mNECs suggests that the

increasing population of tdTom[+] mNEC cells is dependent on the prior genesis of the mTC

population.

mTCs transdifferentiate into mNECs in PDA.

169   The observation that mTCs act as a precursor cell to mNECs suggested two possible

170 models for this relationship: mTCs act as a stem-like cell, giving rise to mNEC daughter cells,

171 or alternatively, they directly transdifferentiate to mNECs. To distinguish these possibilities,

172 we tested if mNEC genesis had undergone DNA replication, consistent with them arising from

173 an asymmetric cell division of mTCs. mTCs rarely proliferate in neoplasia models [30], but their

174 proliferative capacity has not been examined in a carcinoma model.

175   Initially, we co-stained for Ki67 and Syp and found that, like mTCs, mNECs were not in

176 the cell cycle in *KF-P2f3T* or *KPF-P2f3T* mice (Fig. 3C, *Black arrows*). To test if a rare cell

177 division event was necessary for mNEC genesis, we used a long-term EdU incorporation

178 assay to determine if mNECs had gone through a prior cell division [9, 11]. *KPF-P2f3T* mice

179 were treated with tamoxifen once daily for 5 days at 11-12 weeks of age. We treated mice

180 with EdU every 12 hours for 3 days and collected pancreata 12 hours after the last injection.

181 Although we identified several EdU+ cells in the tissue (*Magenta arrows)*, neither Cox1$^+$

182 mTCs (*Green arrows)* nor Syp$^+$ TNTs (*White arrows*) had incorporated EdU (Fig. 3D), and,

183 given the relatively long-time frame to allow for incorporation, were unlikely to have

184 proliferated prior to their appearance.

185   To test the alternative that mTC-derived mNECs arise from the direct

186 transdifferentiation of mTCs, we hypothesized that there would be cells that exist in mid-

187 transition, co-expressing markers of both mTCs (Cox1) and mNECs (Syp). To identify

188 transitionary cells, we co-stained Cox1 and Syp in *KPF-P2f3T* mice at 7 weeks post-

189 tamoxifen treatment. Although rare, we identified cells expressing both Cox1 and Syp (Fig.

190 3E, *Yellow arrows*) exclusively in *KPF-P2f3T* mice. These transitionary cells were also EdU

negative (Supp. Fig. 3C, *Yellow arrows*). Collectively, our data support a "Tuft-to-

Neuroendocrine Transition" (TNT) transdifferentiation event, analogous to an epithelial-to-

mesenchymal transition, exclusively in our pancreatic carcinoma model.

Neoplastic mNECs have an enteroendocrine phenotype.

While we have found mNECs in the pancreata of both neoplasia and carcinoma

models, they appear to have distinct cellular origins. We confirmed that mNECs and mTCs in

both models derive from acinar cells [17, 22] using an acinar cell lineage trace of *Pt1fa*[Cre/+]*;*

ROSA26[LSL-YFP/+] in neoplasia and carcinoma models. We show co-localization with GFP (to

mark acinar cells and progeny) and either Cox1 (mTCs) or Syp (mNECs) for both models

(Supp. Fig. 2C). Although mNECs derive from acinar cells in neoplasia and carcinoma, in the

KPF-P2f3T carcinoma model, mNECs require the genesis of an mTC intermediate. This

unexpected observation led us to ask if mNECs in the *KF-P2f3T* neoplasia model and the

KPF-P2f3T carcinoma model are distinct identities.

Two types of mNECs are described in human pancreatic disease: neuroendocrine-like

cells [21], also designated enteroendocrine cells (EECs) [17, 22], marked by their expression of

islet cell hormones, and neural-like progenitor cells (NRPs) [21], also called ductal

neuroendocrine cells [7, 23], marked by the expression of genes associated with neural

development. EECs are distinct from tuft cells in their origins in pancreatic disease [17, 22] in

that they do not rely on Pou2f3 activity [17, 22, 31], and express markers of normal islet cells [21, 22].

To determine if mNECs in our models bore an EEC phenotype, we examined islet cell

hormone expression [21] by co-IF of wide-spectrum cytokeratin (PanCK) and Syp together with

either glucagon (Gcg), insulin (Ins), somatostatin (Sst), pancreatic polypeptide (Ppy), and

ghrelin (Ghrl), markers of α, β, δ, γ, and ε islet cells, respectively [32].

Since the KF mNECs do not express a lineage tracer, we confined our analysis to

PanCK+/Syp+ cells to distinguish them from singular islet cells that can become trapped

within the diseased tissue (Figs. 4A-E & Supp. Fig. 3D). In the *KF-P2f3T* neoplasia model,

we consistently found isolated mTC-independent mNECs expressing islet cell identity

markers apart from the β cell markers Ins and serotonin (5-HT) (Fig. 4B and Supp. Fig. 2B).

In contrast, we found no islet hormone expression in the mTC-derived mNECs in the *KPF-*

P2f3T mouse. These data indicated that mNECs in our KF-P2f3T neoplasia model, but not in

our *KPF-P2f3T* carcinoma model, are likely EECs, similar to those found in other models of

pancreatic disease [17, 21, 22].

Cells derived from TNT persist in pancreatic carcinoma.

Having established that mNECs in KF neoplasia are likely EECs, whereas those in the

KPF carcinoma are not, we hypothesized that the TNT-derived mNECs are neural-like

progenitor cells (NRPs), as identified in human PDA [21]. In human PDA, NRPs correlate with

poor survival and drug resistance [23]. Initially, we asked if our TNT mNECs persist in frank

carcinoma, as NRPs do in patients. To test this, we extended the trace time post-tamoxifen

treatment to 12 weeks and examined tdTom expression in areas of undifferentiated

carcinoma, confirmed by histology (data not shown). As expected, tuft cells were absent in

regions of carcinoma, as indicated by IHC for Dclk1 (Fig. 5A, *Left*). However, when we

stained adjacent sections for tdTom, we easily found mTC progeny cells scattered throughout

the tumors (Fig. 5A, *Right*). Using co-IF, we confirmed that the identity of these mTC

progeny were indeed Syp-expressing mNECs (Fig. 5B). We conclude that not only are mTCs

undergoing TNT, but the mTC-derived mNECs persist within dedifferentiated carcinoma,

consistent with NRPs in PDA patients [21].

237 **TNT gives rise to Nrxn3⁺ neural-like progenitor cells in pancreatic carcinoma.**

238   Neurexin 3 (Nrxn3) is a marker of neural-like progenitor cells (NRPs) linked with poor

239 prognosis and chemotherapy resistance in multiple cancers, including colon, lung, and

240 pancreatic cancer [21, 33, 34]. The persistence of mNECs in carcinoma led us to examine the

241 expression of Nrxn3 in the TNT-derived mNECs in the KPF carcinoma model. KF neoplasia

242 and KPF carcinoma tissues were co-stained for Nrxn3, Syp, and PanCK to distinguish

243 mNECs with an NRP identity. As expected, *KF-P2f3T* EECs did not express Nrxn3 (Fig. 6A

244 and 6B, *Top*). However, in the *KPF-P2f3T* carcinoma model, Nrxn3, Syp, and PanCK co-

245 localization confirmed that TNT-derived mNECs cells were indeed NRPs both in areas of

246 neoplasia (Fig. 6A and 6B, *Bottom*) and carcinoma (Fig. 6C).

247 **The Tuft to Neuroendocrine transition is dependent upon *Myc* activity.**

248   Pancreatic parenchyma-wide *Myc* activity induces the formation of mNECs in a KC

249 model of neoplasia [23], similar to the prostate and lung [35-38]. NRPs in patient samples are

250 marked by high levels of MYC activity in cancers of the pancreas and other tissues [21, 23, 35, 36].

251 We hypothesized that targeting *Myc* activity specifically to mTCs in the *KF-P2f3T* neoplasia

252 mouse, where TNT is not otherwise observed, would be sufficient to induce NRP genesis. To

253 accomplish this, we crossed the *Myc* Cre-dependent overexpression mouse line, ROSA26[LSL-

254 cMyc/+], into the *KF-P2f3T* model (designated KF-MOE) (Supp. Fig. 4A). We treated these mice

255 with cerulein to induce transformation and mTC genesis, then treated with tamoxifen 3 weeks

256 later to activate the tdTom reporter and induce upregulation of *Myc* specifically in the mTCs.

257 As previously observed, the control *KF-P2f3T* mice had no mTC-derived mNECs. In

258 contrast, we readily found mTC-derived mNEC genesis in KF-MOE tissue as early as 3 days

259 post-induction (Fig. 7A and Supp. Fig. 4C), *Yellow arrows*), with a 6-fold increase in the

tdTom$^+$ mNEC population 7 weeks post-tamoxifen treatment (Fig 7C, *Pink*). Finally, to

determine if the mNECs in the *KF-MOE* mice were NRPs, we co-stained tissue 7 weeks post-

tamoxifen for Nrxn3, Syp, and tdTom (Fig. 7B). Indeed, we consistently found NRP identity in

the neuroendocrine population induced by overexpressing *Myc* in mTCs. Thus, *Myc*

expression targeted to mTCs is sufficient to induce transdifferentiation to NRPs in a neoplasia

model where NRPs are not observed otherwise.

Having shown that Myc activity is sufficient to drive TNT, we tested if it was necessary

for the process. To accomplish this, we crossed a conditional *Myc* knock-out allele (*Myc$^{fl/+}$*)

into the *KPF-P2f3T* model (designated *KPF-P2M$^{fl/+}$*) (Supp. Fig. 4B). These mice were

treated with tamoxifen to simultaneously activate the tdTom reporter and reduce *Myc* activity

specifically in the mTCs. As previously, the *KPF-P2f3T* control at 7 weeks post-tamoxifen

showed a lack of mTCs and an increased population of mTC-derived mNECs (Fig. 7D). In

contrast, in the *KPF-P2M$^{fl/+}$* pancreata, we could easily identify Cox1$^+$/tdTom$^+$ mTCs at 7

weeks post-tamoxifen (Fig. 7D&E) and a lack of Syp$^+$/tdTom$^+$ mTC-derived mNECs.

Unexpectedly, these mice had a significant population of Syp$^+$/tdTom$^-$ mTC-independent

mNECs (Fig. 7D; Supp. Fig 4D). To test if the mTC-independent mNECs were EECs, we

combined antibodies for Ghrl, Gcg, Sst, Ppy, and Ins to identify any hormone-positive cells

simultaneously and co-stained for tdTom (Fig. 7E). Indeed, the tdTom$^-$ mNECs in the *KPF-*

P2M$^{fl/+}$ model expressed islet hormones, confirming their identity as EECs. We conclude that

Myc activity is necessary and sufficient to induce mTC transdifferentiation into NRPs.

Discussion

In the pancreas, prolonged injury or expression of oncogenic *Kras* leads to acinar-to-

ductal metaplasia, with metaplastic ducts comprised of at least 6 different cell types: gastric

pit-like cells, gastric chief-like cells, gastric SPEM/neck-like cells [6], senescent duct cells,

metaplastic neuroendocrine cells (mNECs), and metaplastic tuft cells (mTCs) [7, 9, 17]. The

functions of these different metaplastic cell types have only begun to be recognized. For

instance, normal tuft cells in other organs combat infection [14] and promote wound healing [11,

30, 39]. Consistent with their overall role in maintaining normal tissue homeostasis, mTCs

suppress the progression of pancreatic neoplasia to adenocarcinoma [8, 10]. In contrast,

mNECs in the pancreas are associated with tumor cell-neuron cross talk and promote tumor

cell proliferation [40].

While mTCs functionally hinder the progression of neoplasia to pancreatic cancer, they

also have been theorized to be tumor progenitor cells in the pancreas[11, 30, 41], intestine [42], and

stomach [43]. Doublecortin-like kinase (Dclk1), a common tuft cell marker, is also a marker of

progenitor cells in the pancreas and intestine [11, 30, 42]. Dclk1 is a microtubule kinase highly

expressed in neurons and is linked with neuronal survival and migration in other organs [44].

Besides Dclk1, mTCs express other stem cell markers such as CD133, CD24, and CD44 [11],

which, together with their low proliferation rate [11, 30, 42, 45], support their role as putative cancer

stem cells. Consistent with a facultative progenitor cell function, depleting mTCs delays

tissue repair after experimental pancreatitis [30, 42]. However, contrary to their proposed role as

cancer-initiating cells in PDA, mTC numbers diminish as neoplasia progresses to carcinoma

[9, 11], inconsistent with their mandatory role in the genesis of new tumors.

We theorized many possible reasons why mTCs disappear through progression. With

little evidence that they are dying or being outcompeted by more proliferative cell types, we

focused on the possibility that they were changing their cellular identity. Using a unique dual

recombinase mouse model to trace mTC fate through progression, we found that mTCs were

stable in $Kras^{G12D}$-driven neoplasia but transdifferentiated to a mNEC identity in a $Kras^{G12D}$;

Tp53[+/-] carcinoma model. Importantly, we cannot distinguish whether it is progression to a

less differentiated adenocarcinoma in general or the deletion of a *Tp53* allele specifically that

is responsible for driving this transition. However, given that we do not observe mTCs or

their progeny in carcinomas derived from *Kras*[G12D] expression only (data not shown), this

suggests an association with general tumor progression, agnostic to specific tumor

suppressor gene mutations.

Pancreatic tumor cells expressing neural/neuroendocrine markers in human PDA and

are comprised of two different subsets: neuroendocrine-like or enteroendocrine cells (EECs)

[7, 22] and neural-like progenitor cells (NRPs) [21], the latter of which is associated with poor

outcomes. EECs express hormone markers typically confined to islet cells [17, 21, 22], whereas

NRPs express markers of neural progenitor cells, including Nrxn3 [21]. We found that mNECs

in *Kras*[G12D]-driven neoplasia arise independent of mTCs and take on an islet hormone-

positive EEC identity, consistent with previous data showing EECs in metaplasia are

independent of mTCs and do not depend on *Pou2f3* to develop [22]. In contrast, mNECs in

Kras[G12D]; *Trp53*[+/-] carcinoma do derive from mTCs and have an Nrxn3[+] NRP phenotype.

EECs share a seemingly similar fate as mTCs in that they are present at early stages of ADM

but diminish as PDA progresses to dedifferentiated carcinoma [22]. Our data support an

association of EECs with early oncogenesis as they are largely confined to our neoplasia

model and absent in our carcinoma model. Unexpectedly, ablation of a single allele of *Myc* in

the carcinoma model mTCs allowed EECs to reemerge, suggesting an inverse relationship

between EECs and NRPs.

Mouse mNECs (both EECs and NRPs) derive from acinar cells [17, 22, 23], but NRPs

require an mTC intermediate. NRPs are associated with high *Myc* activity in PDA and other

carcinomas such as lung, colon, and prostate cancer [21, 23, 35, 36]. Supporting the orthologous

nature of mouse NRP cells, upregulating *Myc* expression specifically in mTCs in our

neoplasia model was indeed sufficient to drive NRP transdifferentiation whereas reducing

Myc activity in the mTCs of our carcinoma model inhibited the process. Together, these data

suggest that *Myc* is both necessary and sufficient for a "tuft to neural progenitor transition"

(TNT) in PDA. NRPs contribute to tumorigenicity [46] and advanced carcinomas frequently

express neural characteristics [47], consistent with their association with poor outcomes in PDA

patients [21, 23] and our observation that NRPs are long-lived, persisting within poorly

differentiated carcinomas. Even so, we have not yet observed conditions under which murine

NRPs proliferate so as to constitute a significant population of carcinoma cells. It is possible

that NRPs being resistant to chemotherapy may be selected for confronted with treatment [21,

23], and possibly proliferate subsequently in that environment.

The coexistence of tuft cells and neuroendocrine cellular subpopulations is a

characteristic of several carcinoma types. Small cell lung cancer (SCLC) can be classified

into subtypes that include a tuft cell-like subtype based on POU2F3 expression and two

neuroendocrine subtypes based on ASCL1 and NEUROD1 expression [26]. Tuft cell-like SCLC

transitions to neuroendocrine subtypes as SCLC tumors lose *Tp53* and have an increased

activity of *Myc* [37]. In addition to this neural transition, these cells are characterized by the

gain of EMT markers in SCLC [35, 37]. Neuroendocrine prostate cancer (NEPC) NRPs are also

driven by *Myc* expression and *Tp53* deletion [38, 48]. In the setting of prostate cancer, NEPC is

derived from epithelial cells[36] in prostate adenocarcinoma after the treatment of hormone

therapy and is associated with worse outcomes [38].

In summary, our study reveals a novel type of transdifferentiation of metaplastic tuft

cells to neural-like progenitor cells, shifting their role from suppressing tumor progression

through their chemosensory function in early-stage disease, to being associated with poor

outcomes in the later stages after TNT. We found that the TNT requires *Myc* activity [49-51] and

possibly the loss of *Tp53* function[37], associations that have been observed in other tumor

types [27, 35-37]. While this does not settle the debate as to whether mTCs can act as tumor

progenitor cells, the NRP progeny of mTCs can persist long term as part of a poorly

differentiated carcinoma, though their lack of proliferation under unperturbed conditions

makes it unlikely that they contribute significantly to overall tumor burden. The observation

that NRPs are resistant to chemotherapy and their numbers are associated with worse

outcome and tumor recurrence[21, 23] suggest that, when challenged, their role may shift to a

more traditional tumor initiating cell. Other studies suggest that mNECs instead play a

supportive role for other carcinoma cells, promoting their expansion and possibly contributing

to their survival under stress [40], though whether these are functions common to both EECs

and NRPs has not been determined. While the specific functions of NRPs in PDA remain

under investigation, understanding the mechanisms controlling their genesis suggests that

one possible way to target these aggressive carcinoma cells is to reverse the TNT process to

restore their mTC tumor suppressing role.

Methods

Mouse Models

mTC Lineage tracing

Pou2f3$^{CreERT/+}$ [19], *ROSA26$^{LSL-TdTomato/+}$* [28]. These were combined to generate

Ptf1a$^{FlpO/+}$; FSF-KRAS$^{G12D/+}$ [18, 52]; *Pou2f3$^{CreERT/+}$* [19]; *ROSA26$^{LSL-TdTomato/+}$* (*KF-P2f3T*) and

Ptf1a$^{FlpO/+}$; FSF-KRAS$^{G12D/+}$; Trp53$^{Frt-Exons\ 2\ to\ 5-Frt/+}$ [18, 52]; *Pou2f3$^{CreERT/+}$* [19]; *ROSA26$^{LSL-TdTomato/+}$*

(*KPF-P2f3T*) mice. We induced pancreatic damage in the 8- to 9-week-old *KF-P2f3T* mice

with a once-daily intraperitoneal injection of cerulein (46-1-50; American Peptide Company,

Inc, Sunnyvale, CA) at a dose of 250 μg/kg for a total of five days. To induce recombination

of ROSA26[LSL-TdTomato/+] in *KF-P2f3T* and *KPF-P2f3T* models, 11- to 12-week-old mice were

treated with 5mg of Tamoxifen (T5648; Millipore-Sigma, St Louis, MO) via oral gavage, once

a day, for 5 days. Tamoxifen was dissolved in corn oil before treating mice. Pancreata were

then harvested 1-day post-tamoxifen, three days post-tamoxifen, and seven weeks post-

tamoxifen to capture different stages of PDA in mice.

Acinar-derived mTCs and mNECs

Ptf1a[Cre+]; LSL-KRAS[G12D/+]; LSL-Trp53[R172H/+]; ROSA26[LSL-YFP/+] [9] were combined to

generate *Ptf1a[Cre+]; LSL-KRAS[G12D/+], ROSA26[LSL-YFP/+]* (KCY) and *Ptf1a[Cre+]; LSL-KRAS[G12D/+];*

LSL-Trp53[R172H/+]; ROSA26[LSL-YFP/+] (KPCY) mice to allow lineage trace of epithelial

compartment. KCY mice were treated with Cerulein at 9 weeks old and allowed to recover

for 3 weeks. KCY and KPCY mice were treated with Tamoxifen to induce YFP expression in

Ptf1a+ cells at twelve weeks of age.

mTC Ablation

A homozygous CreERT knock-in inserted into the *Pou2f3* locus was used to induce

metaplastic tuft cell ablation in KF and KPF mice. 8- to 9-week-old *Ptf1a[FlpO/+]; FSF-*

KRAS[G12D/+] [18, 52]*; Pou2f3[CreERT/CreERT]* (KF; Pou2f3[CreERT/CreERT]) mice were treated with cerulein

via intraperitoneal injection at a dose of 250 μg/kg for a total of 5 days to induce pancreatic

damage*. Ptf1a[FlpO/+]; FSF-KRAS[G12D/+]; Trp53[Frt-Exons 2 to 5-Frt/+]*; [18, 52] *Pou2f3[CreERT/CreERT]* (KPF;

Pou2f3[CreERT/CreERT]) did not require cerulein to cause damage. Pancreata were harvested at

12 weeks of age to determine ablation of mTCs in both models.

cMyc overexpression in mTCs

To Investigate the role of *cMyc* in the transdifferentiation of mTCs into mNECs, we

bred mouse lines to create the genotype: $Ptf1a^{FlpO/+}$, $KRAS^{FSF-G12D/+}$, $Pou2f3^{CreERT/+ 16, 20}$,

$ROSA26^{LSL-TdTomato/+}$, $ROSA26^{LSL-cMyc/+ 23}$. These were combined to generate KF-MOE mice

comprised of $Ptf1a^{FlpO/+}$; $FSF-KRAS^{G12D/+ 18, 52}$; $Pou2f3^{CreERT/+ 20}$; $ROSA26^{LSL-TdTomato/+}$;

$ROSA26^{LSL-cMyc/+ 23}$. KF-MOE mice overexpressed *cMyc* and tdTomato reporter specifically

in mTCs in the KF neoplasia model. Pancreatic damage was induced in 8- to 9-week-old KF-

MOE mice with a once-daily intraperitoneal injection of cerulein (46-1-50; American Peptide

Company, Inc, Sunnyvale, CA) at a dose of 250 µg/kg for a total of 5 days. To induce

recombination of $ROSA26^{LSL-TdTomato/+}$ and $ROSA26^{LSL-cMyc/+ 23}$ in the KF-MOE model, eleven-

to twelve-week-old mice were treated with 5mg of Tamoxifen (T5648; Millipore-Sigma, St

Louis, MO) via oral gavage, once a day, for 5 days. Tamoxifen was dissolved in corn oil

before treating mice. We harvested pancreata 3 days post-tamoxifen and seven weeks post-

tamoxifen to investigate the role of *cMyc* in mTC-derived mNECs.

cMyc flox in mTCs

To Investigate the role of *cMyc* in the transdifferentiation of mTCs into mNECs, we

bred mouse lines to create the genotype: $Ptf1a^{FlpO/+}$, $KRAS^{FSF-G12D/+}$, $Trp53^{Frt-Exons\ 2\ to\ 5-Frt/+\ 18,}$

^{52}, $Pou2f3^{CreERT/+ 16, 20}$, $ROSA26^{LSL-TdTomato/+}$, $Myc^{Flox/+\ 23}$. These were combined to generate

$KPF-P2M^{Fl/+}$ mice comprised of $Ptf1a^{FlpO/+}$; $FSF-KRAS^{G12D/+ 18, 52}$; $Trp53^{Frt-Exons\ 2\ to\ 5-Frt/+}$

$Pou2f3^{CreERT/+ 20}$; $ROSA26^{LSL-TdTomato/+}$; $Myc^{Flox/+\ 23}$. $KPF-P2M^{Fl/+}$ mice reduced *Myc* activity

and expressed the tdTomato reporter specifically in mTCs in the KPF PDA carcinoma model.

To induce recombination of $ROSA26^{LSL-TdTomato/+}$ and $Myc^{Flox/+\ 23}$ in the $KPF-P2M^{Fl/+}$ model,

eleven- to twelve-week-old mice were treated with 5mg of Tamoxifen (T5648; Millipore-

Sigma, St Louis, MO) via oral gavage, once a day, for 5 days. Tamoxifen was dissolved in

corn oil before treating mice. We harvested pancreata seven weeks post-tamoxifen to

investigate the role of *Myc* in mTC-derived mNECs.

Immunohistochemistry (IHC)

Mouse pancreas tissues were fixed overnight in a Z-fix solution (NC9050753; Anatech

Ltd, Battle Creek, MI). The pancreas tissues were processed using a Leica ASP300S tissue

processor (Buffalo Grove, IL). Sections (4 μm) of paraffin-embedded tissue were stained for

target proteins (Table 1) of interest using the Discovery Ultra XT auto Stainer (Ventana

Medical Systems Inc, Tucson, AZ). IHC slides were imaged and stitched together by a

3DHisTech scanner (Perkin Elmer, Seattle, WA) using a ×20 objective lens.

Immunofluorescence (IF)

Tissues were prepared as Z-fixed paraffin-embedded blocks, as stated above. In

increased dilutions, sectioned tissue was deparaffinized using xylene and a series of ethanol.

We boiled the slides in a pH 6.0 Citrate buffer solution for 5 minutes. Slides were allowed to

cool at room temperature for 20 mins, then washed in 0.1% tritonX + PBS solution for 15

mins three times for a total of 45 mins. The sections were then blocked in a donkey block

solution of 5% donkey serum/1% bovine serum albumin (BSA) in PBS for 1 hour. Following

the block, sections were incubated overnight with primary antibodies, as stated in **Supp.**

Table 1. The following day, we washed in a 0.1% TritonX + PBS solution for 15 mins thrice.

Then Alexa Fluor-conjugated secondary antibodies (1:500, Invitrogen, Carlsbad, CA) were

added and incubated at RT for 1hr in the dark. Slides were then washed once more with

0.1% TritonX + PBS + Hoechst 33342 (62249, 1:10000, Thermo Scientific, Carlsbad, CA)

solution for 15 mins thrice to remove excess secondary and label nuclei. Lastly, slides were

briefly rinsed in deionized water, mounted with Prolong Diamond antifade mountant (P36961;

Fisher), and dried overnight at room temperature before imaging.

EdU Proliferation Assay:

The "Click-iT™ Plus EdU Cell Proliferation Kit for Imaging, Alexa Fluor™ 488 dye"

(ThermoFisher, Catalog number: C10637) was used to label cells that have undergone

proliferation for a specific time period [53]. We dissolved EdU in PBS to a concentration of

1mg/mL. *KPF-P2f3T* mice were treated with tamoxifen at 11- to 12- weeks of age once daily

for 5 days. Mice were treated with 100uL of EdU injected via I.P. injection every 12 hours for

3 days post tamoxifen labeling. We harvested the pancreata from each mouse the following

morning. We fixed and stained each pancreas as mentioned in the IF protocol above just

until after secondary antibodies were added and washed in 0.1% TritonX + PBS + Hoechst

33342 (62249, 1:10000, Thermo Scientific, Carlsbad, CA) solution for 15 mins three times to

removed excess secondary and label nuclei. We detected EdU by making a cocktail using

the following components provided by the kit: 440uL of 1x Click-iT Reaction buffer, 10uL of

Copper Protectant, 1.2uL of Alexa Fluor Picolyl, and 50uL of Reaction buffer additive.

Reagents must be added in the order listed. 100uL of the cocktail was added to each slide

and incubated for 30 minutes before washing with 3% BSA in PBS. We rinsed the slides with

ddH2O, mounted them with Prolong Diamond antifade mountant (P36961; Fisher), and dried

them overnight at room temperature before imaging.

Quantitation

HALO Software (Indica Labs, Corrales, NM) was used to quantify whole tissue

scanned sections. Quantification excluded blood vessels, lymph nodes, and

adipose/connective tissue. We took multiple sections from each tissue throughout the block

and averaged them to produce a data point. Through this software, we can train the

algorithm to recognize specific areas of tissue and allow us to determine marker expression

in different regions of the pancreas using the whole section. We did image analysis as per

Indica Labs' recommendations.

Acknowledgements

The authors would like to thank Marina Pasca di Magliano for helpful discussions. This work

was supported by R01CA235141, U01CA224145 and U01CA274154 (to HCC). DJSE was

supported by the University of Michigan's Rackham Merit Fellowship, NCI F31CA271576,

and NCI T32CA009676. KED is supported by the Vanderbilt Ingram Cancer Center Support

Grant (NIH/NCI P30CA068485), the Vanderbilt-Ingram Cancer Center SPORE in

Gastrointestinal Cancer (NIH/NCI P50CA236733), the Vanderbilt Digestive Disease

Research Center (NIH/NIDDK P30DK058404), an American Gastroenterological Association

Research Scholar Award (AGA2021-13-02), NIH/NIGMS R35GM142709, The Department of

Defense (DOD W81XWH2211121), The Sky Foundation, Inc (AWD00000079), and Linda's

Hope (Nashville, TN). RCS is supported by: NCI U01CA224012 and R01s CA186241,

CA196228 and DoD PA210068. SB was supported by a postdoctoral fellowship from the

German Research Foundation (BE 7224/1-1) and a research grant provided by the Sky

Foundation. NGS is supported by NCI R00 CA263154 and the Henry Ford Health + Michigan

State University Collaboration grant.

References:

1.	Society, A.C., *Cancer Facts & Figures 2024.* American Cancer Society, 2024.
2.	Chu, L.C., M.G. Goggins, and E.K. Fishman, *Diagnosis and Detection of Pancreatic Cancer.* The Cancer Journal, 2017. **23**(6): p. 333-342.
3.	Pihlak, R., J. Weaver, J. Valle, and M. McNamara, *Advances in Molecular Profiling and Categorisation of Pancreatic Adenocarcinoma and the Implications for Therapy.* Cancers, 2018. **10**(1): p. 17.
4.	Sahin, I.H., C.A. Iacobuzio-Donahue, and E.M. O'Reilly, *Molecular signature of pancreatic adenocarcinoma: an insight from genotype to phenotype and challenges for targeted therapy.* Expert Opinion on Therapeutic Targets, 2016. **20**(3): p. 341-359.
5.	*Pancreatic Cancer: Molecular Characterization, Clonal Evolution and Cancer Stem Cells.* Biomedicines, 2017. **5**(4): p. 65.
6.	Benitz, S., *et al.*, *ROR2 regulates cellular plasticity in pancreatic neoplasia and adenocarcinoma.* 2023, Cold Spring Harbor Laboratory.
7.	Schlesinger, Y., *et al.*, *Single-cell transcriptomes of pancreatic preinvasive lesions and cancer reveal acinar metaplastic cells' heterogeneity.* Nature Communications, 2020. **11**(1).
8.	Delgiorno, K.E., *et al.*, *Tuft Cells Inhibit Pancreatic Tumorigenesis in Mice by Producing Prostaglandin D2.* Gastroenterology, 2020. **159**(5): p. 1866-1881.e8.
9.	Delgiorno, K.E., *et al.*, *Identification and Manipulation of Biliary Metaplasia in Pancreatic Tumors.* Gastroenterology, 2014. **146**(1): p. 233-244.e5.
10.	Hoffman, M.T., *et al.*, *The Gustatory Sensory G-Protein GNAT3 Suppresses Pancreatic Cancer Progression in Mice.* Cellular and Molecular Gastroenterology and Hepatology, 2021. **11**(2): p. 349-369.
11.	Bailey, J.M., *et al.*, *DCLK1 Marks a Morphologically Distinct Subpopulation of Cells With Stem Cell Properties in Preinvasive Pancreatic Cancer.* Gastroenterology, 2014. **146**(1): p. 245-256.
12.	Nevo, S., N. Kadouri, and J. Abramson, *Tuft cells: From the mucosa to the thymus.* Immunology Letters, 2019. **210**: p. 1-9.
13.	Hoover, B., V. Baena, M.M. Kaelberer, F. Getaneh, S. Chinchilla, and D.V. Bohórquez, *The intestinal tuft cell nanostructure in 3D.* Scientific Reports, 2017. **7**(1).
14.	Howitt, M.R., *et al.*, *Tuft cells, taste-chemosensory cells, orchestrate parasite type 2 immunity in the gut.* Science, 2016. **351**(6279): p. 1329-1333.
15.	Yamashita, J., M. Ohmoto, T. Yamaguchi, I. Matsumoto, and J. Hirota, *Skn-1a/Pou2f3 functions as a master regulator to generate Trpm5-expressing chemosensory cells in mice.* PLOS ONE, 2017. **12**(12): p. e0189340.
16.	Yamaguchi, T., *et al.*, *Skn-1a/Pou2f3 is required for the generation of Trpm5-expressing microvillous cells in the mouse main olfactory epithelium.* BMC Neuroscience, 2014. **15**(1): p. 13.
17.	Ma, Z., *et al.*, *Single-Cell Transcriptomics Reveals a Conserved Metaplasia Program in Pancreatic Injury.* Gastroenterology, 2022. **162**(2): p. 604-620 e20.
18.	Garcia, P.E., *et al.*, *Differential Contribution of Pancreatic Fibroblast Subsets to the Pancreatic Cancer Stroma.* Cell Mol Gastroenterol Hepatol, 2020. **10**(3): p. 581-599.
19.	McGinty, J.W., H.-A. Ting, T.E. Billipp, M.S. Nadjsombati, D.M. Khan, N.A. Barrett, H.-E. Liang, I. Matsumoto, and J. Von Moltke, *Tuft-Cell-Derived Leukotrienes Drive Rapid Anti-helminth Immunity in the Small Intestine but Are Dispensable for Anti-protist Immunity.* Immunity, 2020. **52**(3): p. 528-541.e7.
20.	Matsumoto, I., M. Ohmoto, M. Narukawa, Y. Yoshihara, and K. Abe, *Skn-1a (Pou2f3) specifies taste receptor cell lineage.* Nature Neuroscience, 2011. **14**(6): p. 685-687.
21.	Hwang, W.L., *et al.*, *Single-nucleus and spatial transcriptome profiling of pancreatic cancer identifies multicellular dynamics associated with neoadjuvant treatment.* Nature Genetics, 2022. **54**(8): p. 1178-1191.

22. Caplan, L.R., V. Vavinskaya, D.G. Gelikman, N. Jyotsana, V.Q. Trinh, K.P. Olive, M.C.B. Tan,
and K.E. Delgiorno, *Enteroendocrine Cell Formation Is an Early Event in Pancreatic*
Tumorigenesis. Frontiers in Physiology, 2022. **13**.
23. Farrell, A.S.*, et al.*, *MYC regulates ductal-neuroendocrine lineage plasticity in pancreatic*
ductal adenocarcinoma associated with poor outcome and chemoresistance. Nature
Communications, 2017. **8**(1).
24. Wu, C.-Y.C.*, et al.*, *PI3K Regulation of RAC1 Is Required for KRAS-Induced Pancreatic*
Tumorigenesis in Mice. Gastroenterology, 2014. **147**(6): p. 1405-1416.e7.
25. Wen, H.-J., S. Gao, Y. Wang, M. Ray, M.A. Magnuson, C.V.E. Wright, M.P. Di Magliano, T.L.
Frankel, and H.C. Crawford, *Myeloid Cell-Derived HB-EGF Drives Tissue Recovery After*
Pancreatitis. Cellular and Molecular Gastroenterology and Hepatology, 2019. **8**(2): p. 173-192.
26. Baine, M.K.*, et al.*, *SCLC Subtypes Defined by ASCL1, NEUROD1, POU2F3, and YAP1: A*
Comprehensive Immunohistochemical and Histopathologic Characterization. Journal of
Thoracic Oncology, 2020. **15**(12): p. 1823-1835.
27. Huang, Y.-H.*, et al.*, *POU2F3 is a master regulator of a tuft cell-like variant of small cell lung*
cancer. Genes & Development, 2018. **32**(13-14): p. 915-928.
28. Madisen, L.*, et al.*, *A robust and high-throughput Cre reporting and characterization system for*
the whole mouse brain. Nat Neurosci, 2010. **13**(1): p. 133-40.
29. Burclaff, J., S.G. Willet, J.B. Sáenz, and J.C. Mills, *Proliferation and Differentiation of Gastric*
Mucous Neck and Chief Cells During Homeostasis and Injury-induced Metaplasia.
Gastroenterology, 2020. **158**(3): p. 598-609.e5.
30. Westphalen, C.B.*, et al.*, *Dclk1 Defines Quiescent Pancreatic Progenitors that Promote Injury-*
Induced Regeneration and Tumorigenesis. Cell Stem Cell, 2016. **18**(4): p. 441-455.
31. Worthington, J.J., F. Reimann, and F.M. Gribble, *Enteroendocrine cells-sensory sentinels of*
the intestinal environment and orchestrators of mucosal immunity. Mucosal Immunology,
2018. **11**(1): p. 3-20.
32. Riedel, M.J., A. Asadi, R. Wang, Z. Ao, G.L. Warnock, and T.J. Kieffer, *Immunohistochemical*
characterisation of cells co-producing insulin and glucagon in the developing human pancreas.
Diabetologia, 2012. **55**(2): p. 372-381.
33. Hu, L.*, et al.*, *Genome-Wide Association Study of Prognosis in Advanced Non–Small Cell*
Lung Cancer Patients Receiving Platinum-Based Chemotherapy. Clinical Cancer Research,
2012. **18**(19): p. 5507-5514.
34. Lim, U.*, et al.*, *Susceptibility variants for obesity and colorectal cancer risk: The multiethnic*
cohort and PAGE studies. International Journal of Cancer, 2012. **131**(6): p. E1038-E1043.
35. Ireland, A.S.*, et al.*, *MYC Drives Temporal Evolution of Small Cell Lung Cancer Subtypes by*
Reprogramming Neuroendocrine Fate. Cancer Cell, 2020. **38**(1): p. 60-78.e12.
36. Lee, J.K.*, et al.*, *N-Myc Drives Neuroendocrine Prostate Cancer Initiated from Human Prostate*
Epithelial Cells. Cancer Cell, 2016. **29**(4): p. 536-547.
37. Mollaoglu, G.*, et al.*, *MYC Drives Progression of Small Cell Lung Cancer to a Variant*
Neuroendocrine Subtype with Vulnerability to Aurora Kinase Inhibition. Cancer Cell, 2017.
31(2): p. 270-285.
38. Yin, Y.*, et al.*, *N-Myc promotes therapeutic resistance development of neuroendocrine prostate*
cancer by differentially regulating miR-421/ATM pathway. Molecular Cancer, 2019. **18**(1).
39. Barker, N., *Adult intestinal stem cells: critical drivers of epithelial homeostasis and*
regeneration. Nature Reviews Molecular Cell Biology, 2014. **15**(1): p. 19-33.
40. Sinha, S.*, et al.*, *PanIN Neuroendocrine Cells Promote Tumorigenesis via Neuronal Cross-talk.*
Cancer Research, 2017. **77**(8): p. 1868-1879.
41. Koga, H., Y. Ikezono, and T. Torimura, *Pancreatic DCLK1 marks quiescent but oncogenic*
progenitors: a possible link to neuroendocrine tumors. Stem Cell Investigation, 2016. **3**: p. 37-
37.
42. Nakanishi, Y.*, et al.*, *Dclk1 distinguishes between tumor and normal stem cells in the intestine.*
Nature Genetics, 2013. **45**(1): p. 98-103.

43.   Saqui-Salces, M., T.M. Keeley, A.S. Grosse, X.T. Qiao, M. El-Zaatari, D.L. Gumucio, L.C.
Samuelson, and J.L. Merchant, *Gastric tuft cells express DCLK1 and are expanded in*
hyperplasia. Histochemistry and Cell Biology, 2011. **136**(2): p. 191-204.
44.   Nawabi, H.*, et al.*, *Doublecortin-Like Kinases Promote Neuronal Survival and Induce Growth*
Cone Reformation via Distinct Mechanisms. Neuron, 2015. **88**(4): p. 704-719.
45.   May, R.*, et al.*, *Identification of a novel putative pancreatic stem/progenitor cell marker*
DCAMKL-1 in normal mouse pancreas. Am J Physiol Gastrointest Liver Physiol, 2010. **299**(2):
p. G303-10.
46.   Xu, L., M. Zhang, L. Shi, X. Yang, L. Chen, N. Cao, A. Lei, and Y. Cao, *Neural stemness*
contributes to cell tumorigenicity. Cell & Bioscience, 2021. **11**(1).
47.   Zhang, Z.*, et al.*, *Similarity in gene-regulatory networks suggests that cancer cells share*
characteristics of embryonic neural cells. Journal of Biological Chemistry, 2017. **292**(31): p.
12842-12859.
48.   Beltran, H.*, et al.*, *The role of lineage plasticity in prostate cancer therapy resistance.* Clinical
Cancer Research, 2019: p. clincanres.1423.
49.   Dai, M.-S., Y. Jin, J.R. Gallegos, and H. Lu, *Balance of Yin and Yang: Ubiquitylation-Mediated*
Regulation of p53 and c-Myc. Neoplasia, 2006. **8**(8): p. 630-644.
50.   Feng, Y.C.*, et al.*, *c-Myc inactivation of p53 through the pan-cancer lncRNA MILIP drives*
cancer pathogenesis. Nature Communications, 2020. **11**(1).
51.   Ho, J.S.L., W. Ma, D.Y.L. Mao, and S. Benchimol, *p53-Dependent Transcriptional Repression*
of c-myc Is Required for G$_1$ Cell Cycle Arrest. Molecular and Cellular Biology,
2005. **25**(17): p. 7423-7431.
52.   Wu, J.*, et al.*, *Generation of a pancreatic cancer model using a Pdx1-Flp recombinase knock-*
in allele. PLOS ONE, 2017. **12**(9): p. e0184984.
53.   Zhang, H.*, et al.*, *Glucose-Mediated Repression of Menin Promotes Pancreatic β-Cell*
Proliferation. Endocrinology, 2012. **153**(2): p. 602-611.
616
617

Figure Legends:

Figure 1. Lineage trace model shows tuft cell specific expression of tdTomato.

Genetic strategy to trace the fate of metaplastic tuft cells (mTCs) in pancreatic cancer progression

using a dual recombinase neoplasia (*KF-P2f3T*) model **(A)** and carcinoma (*KPF-P2f3T*) model **(B)**. **C)**

Treatment regimen for neoplasia and carcinoma Models. The models have both pancreatic

progression and reporter specifically in tuft cells. The *KF-P2f3T* model utilizes Cerulein once daily,

followed by 2 weeks recovery, then once daily tamoxifen treatment for 5 days. The *KPF-P2f3T* model

is aggressive as it contains one null p53 allele. Cerulein is not necessary and so only Tamoxifen is

treated once daily for 5 days. Mice are allowed to age for different time points to determine lineage of

metaplastic tuft cells in progression of pancreatic cancer. **D)** Dclk1 in Brown identifies tuft cells in

neoplasia model (left) and carcinoma model (right). **E)** Specific expression of tdTomato reporter in

both models. White represents Hoechst staining to mark the nucleus, tdTomato in red to show

recombination of tdTomato (tdTom), Pou2f3, the master regulator of tuft cells in Green, and

Acetylated alpha tubulin in Magenta to mark filamentous actin in the tufts of the mTCs. **F)** Dual IHC of

tdTom (Teal) and CC3 (Brown) stained *KF-P2f3T* and *KPF-P2f3T* mice to determine cell death of

mTCs in neoplasia and carcinoma. Black arrows indicate tdTom+ cells in lesions. Scale bars = 10 μm

unless otherwise noted.

Figure 2. Lineage tracing mTCs reveals transdifferentiation to metaplastic neuroendocrine

cells (mNECs).

Dual recombinase models show fate of mTCs in progression of pancreatic cancer. **A)** Mice harvested

3 days post-tamoxifen treatment for labeling mTCs. Hoechst stains nucleus of cells (White), Cox1 is

used to identify mTCs (Green), tdTom reporter to determine fate of mTCs (Red) and acetylated-α-

tubulin (Magenta) marks filamentous actin on tuft of mTCs. **B)** Quantification of Percentage of mTC

(White) and mTC Progeny (Pink) among ductal cells at different time points post-tamoxifen treatment

in *KF-P2f3T* and *KPF-P2f3T* models. mTCs are identified by Cox1 and tdTom, mTC Progeny are only

tdTom positive. (n = 3/group) p-values were calculated using a two-way ANOVA on Prism GraphPad

7. * = $p < 0.05$, ** = $p < 0.01$. **C)** Co-staining of tdTomato (Red) expression with remaining

metaplastic ductal cell subsets. In Green, Gkn1, Lectin GS-II, and Synaptophysin (Syp) mark gastric

pit-like, gastric neck-like, and neuroendocrine cells respectively in pancreatic metaplastic ducts. **D)** 7

weeks post-tamoxifen labeling of neoplasia and carcinoma models exposes shift of populations.

Cox1 (Green) marks mTCs, tdTomato (Red) identifies reporter, Syp (White) marks neuroendocrine

cells, and Blue represents DAPI Staining to identify the nucleus of a cell. **E)** Quantification of

percentage of mTCs (Cox1+ tdTom+) in White, mTC-derived mNECs (Syp+ tdTom+) in Pink, mTC-

independent mNECs (Syp+ tdTom-) in Green. (n = 3/group). p-values were calculated using a two-

way ANOVA on Prism GraphPad 7. * = $p < 0.05$, ** = $p < 0.01$. Scale bars = 10 μm unless otherwise

noted.

Figure 3. Characterizing the tuft cell to neuroendocrine cell transdifferentiation (TNT).

IF staining of Cox1 (Red) and Syp (Green) in neoplasia and carcinoma models determines that

mNECs in the *KPF-P2f3T* model relies on the mTC population. **A)** Cox1 (Red) and Syp (Green)

markers are stained in Pou2f3 double knock-in models of pancreatic neoplasia and carcinoma: (Left)

KF; Pou2f3[CreERT/CreERT] and (Right) KPF; Pou2f3[CreERT/CreERT], correspondingly to investigate the role of

tuft cells in mNEC genesis. Green Arrows denote mNECs, Red Asterisks denote stromal Cox1[+]

staining identifying Cox1[+] Immune cells. **B)** Pou2f3, the master regulator of tuft cells (Cyan) is stained

with Cox1, marker of tuft cells (Red), and Syp, neuroendocrine marker (Green) to determine if mNECs

also express Pou2f3. **C)** Dual IHC of Syp (Teal) and Ki67 (Brown) of neoplasia and carcinoma models

of pancreas to determine proliferation of mNECs. **D)** Co-IF of mTC marker, Cox1 (Green), lineage

tracer, tdTom (Red), mNEC marker, Syp (White), incorporated EdU (Magenta) to mark proliferating

cells, and nuclear marker, Hoechst (Blue) in *KPF-P2f3T* mice treated with EdU over the course of 3

days post-tamoxifen treatment. **E)** Yellow arrows mark rare transitionary cells that are marked with

both Syp (Green) and Cox1 (Red). Green arrow identifies cells that are only expressing Syp, mNECs.

Red arrow marks cells only expressing Cox1, mTCs. Scale bars = 10 µm unless otherwise noted.

Figure 4. Identifying distinctions between mNECs in the progression of PDA.

Neoplasia and carcinoma PDA models were stained with markers of different islet compartments to

determine differences between mNECs between the *KF-P2f3T* and *KPF-P2f3T* models. **B)** Islets and

lesions from *KF-P2f3T* and *KPF-P2f3T* are shown to determine co-localization of Syp (Red) and

markers of the different compartments of the Islet (Green). PanCK (Cyan) was used to distinguish

islet cells from mNECs, and Hoechst was used to mark the nucleus of each cell. Glucagon (Gcg),

Insulin (Ins), Somatostatin (Sst), Ghrelin (Ghrl), and Pancreatic Polypeptide (Pp) were stained to

identify the α cells, β cells, δ cells, ε cells and γ cells respectively. Scale bars = 10 µm unless

otherwise noted.

Figure 5. mTC-derived mNECs are identified in PDA Carcinoma.

IHC and IF are used to confirm the identity of mTC Progeny in dedifferentiated carcinoma of *KPF-*

P2f3T model. **A)** IHC of dedifferentiated carcinoma stained with Dclk1 (*Left*) and tdTomato (*Right*) in

Brown. **B)** Dedifferentiated carcinoma of *KPF-P2f3T* model is stained with Hoechst (Cyan), Syp

(White), and tdTomato (Red) to determine identity of mTC progeny. Scale bars = 50 µm

Figure 6. Identification of mTC-derived Nrxn3+ NRPs in Carcinoma

Co-staining of Syp (Red), Nrxn3 (Green), PanCK (Magenta), and Hoechst (White) in *KF-P2f3T*

neoplasia model and *KPF-P2f3T* carcinoma model at **A)** 3 days post-tamoxifen, **B)** 7 weeks post-

tamoxifen, and **C)** *KPF-P2f3T* dedifferentiated carcinoma. Scale bars = 10 µm unless otherwise

noted.

Figure 7. cMyc Mediated development of mNECs.

Overexpression and knockdown of *Myc* in neoplasia and carcinoma models show dependence of *Myc*

expression on transition of mTCs to NRPs in PDA progression. **A)** *KF-MOE* 7 weeks post-tamoxifen

treatment are stained with tdTom to identify cells recombined by tamoxifen treatment and cells

overexpressing *Myc* (Red), Syp to mark neuroendocrine cells (Green), Acetylated-α-tubulin to mark

filamentous actin of cells (Magenta), and Hoechst to identify nucleus of all cells (White) to detect

mTC-derived mNECs. **B)** KF-MOE 7 weeks post-tamoxifen treatment tissue is stained for co-IF to

determine expression of Syp (Green), tdTom (Red), Nrxn3 (Magenta), and Hoechst (White). **C)**

Quantification of mTCs (Cox1+ tdTom+) in White, mTC derived mNECs (Syp+ tdTom+) in Pink and

mTC independent mNECs (Syp+ tdTom-) in Green for *KF-P2f3T* and *KF-MOE* models. Mice were

sac'd after 3 days and 7 weeks post tamoxifen treatment to label tuft cells with tdTomato and

overexpress cMyc specifically in tuft cells. (n = 3/group) p-values were calculated using a two-way

ANOVA on Prism GraphPad 7. * = $p < 0.05$. D) Quantification of mTCs (White), mTC Derived

mNECs (Pink), and mTC Independent, (Green) in *KPF-P2f3T* and *KPF-P2M^{Fl/+}* models. Mice were

sac'd at 7 weeks post tamoxifen treatment to label tuft cells with tdTom and flox *cMyc* expression in

tuft cells specifically. (n = 3/group) p-values were calculated using a two-way ANOVA on Prism

GraphPad 7. * = $p < 0.05$, **= $p<0.005$. **E)** Co-staining of Cox1 (Green), tdTom (Red), PanCK

(Magenta), and Hoechst (White) in *KPF-P2M^{Fl/+}* at 7 weeks post-tamoxifen treatment to identify co-

localization of mTC marker, Cox1, with tdTom (*Top*). Co-staining of combined antibodies for Ghrl,

715   Gcg, Sst, Ppy, and Ins to mark cells expressing "Hormones" to identify EECs (Green), tdTom used to

716   identify cells recombined and *Myc* knockout cells (Red), PanCK to mark pancreatic epithelium/lesions

717   (Magenta), and Hoechst to mark nucleus of cells (White) in *KPF-P2M$^{Fl/+}$* at 7 weeks post-tamoxifen

718   treatment to identify EECs and tdTom+ cells as separate cell populations when *Myc* is conditionally

719   knocked out in carcinoma model. Scale bars = 10 µm unless otherwise noted.

Figure 1

Figure 2

Figure 3

A

KF;Pou2f3^CreERT/CreERT

KPF;Pou2f3^CreERT/CreERT

Hoechst · Syp · Cox1 · Merged

B KPF-P2f3T

Pou2f3 · Syp · Cox1 · Merged

C KF-P2f3T · KPF-P2f3T

Syp · Ki67

D Cox1 · tdTom · Syp · EdU · Hoechst · Merged

E KPF-P2f3T

Hoechst · Syp · Cox1 · Merged

Figure 4

Figure 5

A **KPF-P2f3T**

B **KPF-P2f3T**

Figure 6

A 3 Days Post-Tamoxifen

Syp | Nrxn3 | PanCK | Hoechst | Merged

KF-P2f3T

KPF-P2f3T

B 7 weeks Post-Tamoxifen

Syp | Nrxn3 | PanCK | Hoechst | Merged

KF-P2f3T

KPF-P2f3T

C Dedifferentiated carcinoma

Syp | Nrxn3 | PanCK | Hoechst | Merged

KPF-P2f3T

Figure 7

A KF-MOE 7 Wks — tdTom | Syp | Acetyl-α-Tubulin | Merged

B KF-MOE 7 Wks — Syp | tdTom | Nrxn3 | Hoechst | Merged

C KF-P2f3T Vs. KF-MOE

%Ductal Cells

Time Point Post-Tamoxifen — KF-P2f3T (3d, 7wks), KFMOE (3d, 7wks)

- □ mTCs (Cox1⁺ tdTom⁺)
- ■ mTC Derived mNECs (Syp⁺ tdTom⁺)
- ■ mTC Independent mNECs (Syp⁺ tdTom⁻)

D KPF-P2f3T Vs KPF-P2M^Fl/+

%Ductal Cells

7 weeks Post-Tamoxifen — KPF-P2f3T, KPF-P2M^Fl/+

- □ mTCs (Cox1⁺ tdTom⁺)
- ■ mTC Derived mNECs (Syp⁺ tdTom⁺)
- ■ mTC Independent mNECs (Syp⁺ tdTom⁻)

E KPF-P2M^Fl/+ 7 Wks

Row 1: Cox1 | tdTom | PanCK | Hoechst | Merged

Row 2: Hormones | tdTom | PanCK | Hoechst | Merged