How muscle synergies fail to solve the muscle redundancy problem during human reaching

AS Korol and V Gritsenko

Keywords: *Muscle coactivation, EMG, gravitational torque component, dynamical torque component, active muscle torque.*

Anna S Korol
Department of Neuroscience, School of Medicine, West Virginia University
Rockefeller Neuroscience Institute, West Virginia University
Morgantown, USA
ORCID: 0000-0001-9699-721X

Valeriya Gritsenko
Department of Human Performance, Division of Physical Therapy, School of Medicine, West Virginia University
Department of Neuroscience, School of Medicine, West Virginia University
Rockefeller Neuroscience Institute, West Virginia University
Morgantown, USA
ORCID: 0000-0002-6408-9433

Abstract

The production of movement involves integrating biomechanical, neural, and environmental factors. The biomechanics is complex enough that neural sensorimotor circuits must embed its dynamics for efficient and robust control. However, a problem of redundancy exists, i.e., the problem of choosing among multiple muscles and combinations of joint angles that are possible for a given desired hand position or motion. This problem may be resolved by reducing the dimensionality of the space of motor commands by the central nervous system, i.e., through muscle synergies or motor primitives. Other studies have obtained muscle synergies using decomposition methods. However, we posit that it is not sufficient to show the existence of a low dimensional space, one needs to demonstrate the utility of the obtained synergies in controlling movement. Here we defined a muscle synergy as a single control signal producing specific force direction. We then tested a hypothesis that such muscle synergies exist using dimensionality reduction method. Our approach takes advantage of the close relationship between the temporal profiles of muscle activity observed with electromyography and the joint moments they produce during reaching derived from motion capture. We recorded electromyography of 12 muscles and the kinematics of both arms in 14 right-handed participants performing reaching movements in multiple directions from different starting positions. We used principal component analysis to evaluate the contribution of individual muscles to supporting the arm against gravity and producing propulsive forces. Results show that the joint torques in specific directions (flexion or extension) required to move toward a target were not produced by consistent muscle groups in most conditions as would be expected from the muscle synergy definition outlined above. We further show that both agonistic and antagonistic muscles coactivate in flexible muscle groups but to a different extent between the dominant and non-dominant arms. Our main findings indicate that the nervous system solves the problem of choosing which muscles to activate and when by taking into account limb dynamics rather than reducing the dimensionality through muscle synergies. Furthermore, our data supports the idea of two neural controllers that target different muscle groups in the arm and hand for gross postural and fine goal-directed control of reaching.

Keywords: electromyography, neuromechanics, limb dynamics, hand dominance, gravity, postural forces

Introduction

The production of movement involves integrating biomechanical, neural, and environmental factors. Muscle recruitment by the central nervous system (CNS) causes motion or lack thereof, such as when maintaining posture, through the nonlinear production of forces by muscles about the degrees of freedom (DOFs) of the joints. Body inertia, the number of DOFs, and external forces then shape the resulting posture and movement. This biomechanics is complex enough that neural sensorimotor circuits must embed its dynamics for efficient and robust control [1–3]. However, a problem of redundancy exists, i.e., the problem of choosing among multiple muscles and combinations of joint angles that are possible for a given desired hand position or motion. This problem may be resolved by reducing the dimensionality of the space of motor commands by the CNS, i.e., through muscle synergies or motor primitives. This concept implies that muscles (or motoneurons in the spinal cord) are recruited in groups that represent a specific action so that a combination of the smaller number of synergies can produce a larger number of different movements [4–8]. The anatomical organization of muscles that form agonistic and antagonistic groups support the idea of modular control targets [9]. The spatial organization of motoneuron pools in the spinal cord also captures the functional relationships of the muscles they innervate, further supporting modularity [10]. Indeed, the neural activation of muscles observed with electromyography (EMG) is reducible to a low dimensional space for certain types of movements, including reaching [4,8,11]. However, we posit that it is not sufficient to show the existence of a low dimensional space, one needs to demonstrate the utility of the obtained synergies in controlling movement. For example, if a muscle synergy represents a given controlled force direction, then it needs to scale with the amount of that force across movements in different directions. For example, a single control signal can activate both brachioradialis and biceps muscles to produce elbow flexion moment. A decomposition method can identify this control signal as a synergy represented by the relative contribution of this synergy to each muscle activity profile for a given movement (scores) in a lower dimensional space. Assuming that this low dimensional space is aligned with the joint moments needed to produce these movements [12], then we expect the synergy scores to scale with the amplitude of the force that that synergy controls. We will test this idea in this study using EMG recorded during human center-out reaching movements.

The temporal profile of the envelope of surface EMG is closely related to the force the corresponding muscle is producing in response to neural activation [13]. More recently we have shown that for reaches with the dominant hand towards visual targets in three-dimensional space, the EMG profiles are closely related to the active joint moments that underly the movements [12]. Specifically, the static component of EMG that underlies postural forces needed to support the arm in specific position and during transitions between positions is closely related to the components of joint moments that include gravity terms in the equations of motion. This suggests that the static component of EMG envelope from a given muscle reflects the muscle's contribution to counteracting gravity load on the joints it spans. Moreover, the residual phasic component of EMG is closely related to the residual joint moment that underlies the acceleration and deceleration forces toward the reaching goal after gravity-related

component is subtracted. This suggests that the phasic component of EMG envelope
from a given muscle reflects the muscle's contribution to propulsion in a given direction.
Here we will test the generalizability of these relationships between EMGs and joint
moments to the non-dominant limb and across different workspaces.
In this study, healthy human participants performed unconstrained pointing
movements toward visual targets located equidistantly from one of two central targets.
This resulted in reaching movements in multiple directions performed by either
dominant (right) or non-dominant (left) arm that spanned the typical reaching
workspace. We collected motion capture and EMG data and ran dynamic simulations
with individualized inertial models of the arm to compute joint moments and their
components. We tested two hypotheses. The first hypothesis was that the muscle
activity profiles reflect the forces needed to produce the movement. This hypothesis
tests the generalizability of conclusions from Olesh et al. [12]. The second hypothesis
was that muscle synergies captured with dimensionality reduction methods represent
force directions controlled by the CNS.
95

Methods

Data Collection

The experimental protocol was approved by the Institutional Review Board of
West Virginia University (Protocol #1311129283). The recruitment for this study started
in March 1st, 2014 and ended on July 25th, 2016. Participants provided a written
informed consent prior the start of experiments. A second member of the investigative
team witnessed the signing of the consent form.
Fourteen healthy participants (mean ± std age: 35.43 ± 17.62 years, 6 females
and 8 males) performed a modified center-out reaching task by pointing to visual targets
in virtual reality (software Vizard by Worldviz, Oculus Rift) [12]. All participants reported
they were right-handed and had no neurological or musculoskeletal conditions that
could alter movement. Participants repeated reaching movements between a central
target and one of 14 targets located along a sphere, with 8 targets placed equidistantly
on a horizontal circle parallel with the floor and 8 targets placed equidistantly on a
vertical circle parallel to the body sagittal plane, with 2 targets sharing locations in both
circles as in Olesh *et al.* [12]. Participants were instructed to point to targets with the
index finger, while holding their hand palm down, as quickly and accurately as they can
without moving their trunk and wrist. The distances toward the targets were normalized
for each subject based on their arm segment lengths to minimize the inter-subject
variability in angular kinematics. Each movement to and from each target was repeated
15 times in a randomized order.
The center-out and return movements were repeated with each arm starting at
one of two locations of the central target. The central target was placed at the same
distance relative to the participant's shoulder scaled to their arm length. This further
minimized the inter-subject variability in angular kinematics. In the lateral starting
position, the central target was placed at a location that positioned the shoulder at 0
angle of all degrees of freedom and elbow at 90 degrees, so that the upper arm was

parallel to the trunk and the forearm was parallel to the floor (Fig. 1, pictograms in top
corners, view from above). In the medial starting position, the central target was placed
at a location that positioned the hand at the midline of the body and at the same
distance from the trunk as in the lateral starting position (Fig. 1, pictograms in top
center, view from above). The medial starting position was the same for reaches with
both right and left arms representing common medial workspace. Thus, four conditions
were created based on the location of the central target and the arm used to make
reaches, i.e., LLat and RLat for reaches by left and right arm respectively in their
respective lateral workspaces and LMed and RMed for the reaches by left and right arm
respectively in the common medial workspace.

Movement was recorded and visualized in virtual reality using the Impulse
system (PhaseSpace Inc) at the temporal resolution of 480 Hz. We tracked the location
of 8 light emitting diodes placed on the bony landmarks of the trunk, arm, and hand with
8 cameras as described in detail in Olesh et al. [12]. Motion capture and EMG data
were imported into MATLAB (MathWorks Inc) and processed using custom scripts.

Surface electromyography (EMG) was recorded at the temporal resolution of
2,000 Hz using MA400-28 (MotionLab Systems). EMG was captured from 12 arm
muscles: the clavicular head of pectoralis (Pec), teres major (TerM), anterior deltoid
(ADel), posterior deltoid (PDel), the long and lateral heads of triceps (TriL and TriS), the
short and long heads of biceps (BiS and BiL), brachioradialis (Brd), flexor capri radialis
(FCR), flexor carpi ulnaris (FCU), and extensor capri radialis (ECR). EMG recordings
were temporally synchronized with the motion capture and virtual reality systems during
reaching as described in Talkington *et al.* [14].

Data Analysis

Motion capture data were low pass filtered at 10 Hz and interpolated with a cubic
spline. Arm kinematics was obtained from motion capture by defining local coordinate
systems representing the trunk, arm, forearm, and hand. Joint angles representing 5
DOFs of the arm were derived using linear algebra, namely shoulder flexion-extension,
shoulder abduction-adduction, shoulder internal-external rotation, elbow flexion-
extension, forearm pronation-supination, and wrist flexion-extension (Fig. 1A, example
for one DOF). Active muscle torques and their components were calculated from
kinematic data using a dynamic model of the arm with 5 DOFs (Simulink, MathWorks)
as described in [12]. The individual height, weight, and segment lengths together with
the published anthropometric proportions were used to scale the model inertia to
individual body sizes [15]. Inverse simulations were run with the angular kinematics as
input and the active torques produced by muscles, termed muscle torques (MT), as
output. Simulations were ran with and without gravity force to estimate the components
of muscle torque responsible for supporting the arm against gravity (postural component
of muscle torque or MT_P) and for intersegmental dynamics compensation (dynamic
component of muscle torque or MT_D) as described in detail in Olesh *et al.* [12]. The
simulations were run for each trial. Torque profiles were normalized in time, averaged
per movement direction per DOF and per subject, and down sampled to 100 samples
(Fig. 1B, example for one DOF). Maximum torque amplitudes were determined per DOF

per participant's arm and used to scale the amplitudes of averaged profiles across
movement directions and workspaces.
EMG recordings were high-pass filtered at 20 Hz to remove motion artifacts,
rectified, and low-pass filtered at 10 Hz [16]. Next, EMG recordings were normalized to
movement duration, down sampled to 100 samples, and averaged per movement
direction. Maximum contraction values were determined for each muscle and participant
across averaged EMG timeseries per movement direction and used to scale the
amplitudes of averaged profiles (Fig. 1C).
Principal Component Analysis (PCA) was used to reduce the dimensionality of
EMG and torque data similar to Olesh *et al.* [12]. Normalized and demeaned EMG
profiles from 12 muscles of 28 center-out reaching movements (14 center-out and 14
return movements between each pair of targets) were combined into one matrix, A_{nxm},
where n = 100 (time) and m = 336 (movement directions x muscle). PCA was applied to
the data from each of the four conditions (LLat, RLat, LMed, and RMed) separately.
Thus, four A_{nxm} matrices were obtained for each subject. Similarly, normalized and
demeaned muscle torque profiles (MT, MT_P, and MT_D) from 5 DOFs of 28 center-out
reaching movements were combined into one matrix, B_{nxm}, where n = 100 (time) and m
= 140 (movement directions x DOF). Since the analysis included three types of torques
(MT, MT_P, MT_F) in the four conditions, twelve B_{nxm} matrices were obtained for each
subject. PCA was performed using *pca* function from MATLAB Statistics and Machine
Learning Toolbox. Each matrix was centered, and a default singular value
decomposition algorithm was applied to obtain principal component eigenvectors (Fig.
2A, B), the variance accounted for (VAF) by each principal component (Fig. 2C), and
scores that are the representations of the matrix in the principal component space. It is
important to note that reaching in the same direction leftward or rightward from the
starting location rely on different muscles due to the bilateral symmetry of our body.
That is why the reaches that include leftward movements to the targets on the left from
the central target performed by the left hand were compared to the mirror rightward
movements to the right from the central target performed by the right hand and vice
versa.
197

Statistics

To test the hypotheses, we performed two types of statistical tests. The first test
compared the profiles of the 1st and 2nd principal components obtained from EMG to the
principal components obtained from muscle torques using a correlation analysis (*corr*
function in MATLAB Signal Processing Toolbox). We used repeated measures analysis
of variance (RM ANOVA, *ranova* function in MATLAB Statistics Toolbox) to test the null
hypotheses that the coefficients of determination (R^2) from all conditions and data types
come from the same distribution. Post-hoc multiple comparisons were done using
multcompare function in MATLAB Statistics Toolbox. We applied Bonferroni-Sidak
correction to correct the alpha for familywise error, so that the adjusted alpha for the
comparison between 6 combinations between 4 conditions = 0.0085, for the comparison
between 7 meaningful combinations of two principal components = 0.0073.
The second test evaluated how muscles co-activate across changing starting
postures and movement directions. The rationale is that reaching in different directions

is caused by different amplitudes of postural or dynamic muscle torques. Therefore, the
muscles that cause the actions in the appropriate directions will coactivate together in
proportion to the difference in forces between movements. This implies that there
should be a correlation between the activation amplitude of two muscles that work
together across multiple movement directions, i.e., classical muscle synergies should
emerge. To test this idea, we have utilized a linear correlation analysis (*corr* and *corrplot*
functions in MATLAB Statistics and Econometrics Toolboxes) to quantify the linear
relationships between principal component scores from pairs of muscles across
movements. First, we extracted the scores of the 1st principal component from EMG and
concatenated them into matrix C_{nxm}, where n = 28 (movement directions) and m = 12
(muscles) per condition per subject. Then we evaluated the statistical significance of the
linear relationships between scores of 66 muscle pairs (Fig. 4). We applied Bonferroni-
Sidak correction to correct the alpha for familywise error, so that the adjusted alpha =
0.0008. We repeated this analysis for the 2nd principal component from EMG.
Four older adults out of the total of 14 participants did not complete the reaching
in the medial workspace (RMed and LMed). Therefore, the data from 10 adults (9 young
adults, and one older adult; mean age = 26.3 ± 11.2 years) who completed all 4 tasks
are reported here. The same statistical analysis was performed on all participants
across the two workspace s and reported in Results. Supplementary figures show the
results of the same analysis with inclusion of all individuals across the LLat and RLat
tasks (Supplementary Figures S1 and S2).
233
Results

All participants were able to complete the center out task with either left or right
arms with consistent kinematics (Fig. 1A). We expected that movements with left and
right arms with the same starting positions and kinematics in forward, backward, up, or
down directions would rely on similar forces produced by the same muscles. These
forces are captured here with muscle torques that are further subdivided into a linear
combination of postural and dynamic components. The muscle torque profiles
accompanying the corresponding movements had very similar trajectories across
conditions (Fig. 1B). The were also static offsets in the postural components of muscle
torques underlying matching movement directions. For example, holding the initial
position at the beginning of the movement required larger muscle torques about the
right shoulder flexion/extension DOF (RMed and RLat conditions) compared to that for
the left shoulder in matching conditions (LMed and LLat conditions respectively, Fig.
1B). This is likely due to the motor redundancy, i.e., the different combinations of joint
angles and torques that can result in the same position of the hand. From our earlier
work we knew that important features of muscle activity can be captured by the postural
and dynamic components of muscle torques [12]. However, in that study we focused
our analysis on the movements performed by the dominant arm in the lateral
workspace. Here the analysis encompasses different workspaces and both arms, which
enables us to evaluate the generalizability of this conclusion. We found that intra- and
inter-subject variability of EMG profiles were very low. The average standard deviation
of EMG profiles across movement repetitions for muscles in the dominant right arm
(normalized to maximal EMG of each muscle) was 12% ± 1% across 14 subjects. For

the muscles in the non-dominant left arm the inter-trial variability was 22% ± 2% across
14 subjects. This shows that the muscle activation profiles were more consistent across
repetitions of the same movement type for the dominant arm compared to the non-
dominant arm. Across 14 subjects, the average normalized standard deviation of EMG
profiles of muscles in the dominant right arm was 11% ± 4%. For the muscles in the
non-dominant arm the inter-subject variability was 11% of maximal EMG ± 3%.
Moreover, the EMG profiles also broadly reflected the differences in the joint torques
across conditions (Fig. 1C). In the example for the reach forwards and up in Fig. 1, the
amplitude of anterior deltoid activity increased in parallel with the increases in the
postural component of the shoulder flexion muscle torque across conditions (Fig. 1, MT_P
and ADel). Furthermore, the initial propulsive flexion moment in the dynamic muscle
torque profile was preceded by a burst in the activation of the long head of biceps (Fig.
1, MT_D and BiL, end of the burst is visualized). However, EMG profiles are notoriously
noisy and difficult to interpret. Therefore, in the following analysis we will broadly test
the generalizability of conclusions from Olesh et al. [12] by focusing on the salient
features in EMG obtained with PCA as described in Methods. To test the hypotheses,
we will compare those salient features in EMG with the corresponding features in
muscle torques.
The temporal profile of the 1^{st} principal component (eigenvector E_1) obtained
from EMG matched the 1^{st} principal component (M_1) obtained from muscle torques (MT)
and the 1^{st} principal component (P_1) obtained from the postural components only of the
muscle torques (MT_P; Fig 2A). This feature captures the posture-related changes in
EMG that underly gradually increasing or decreasing EMG profiles. The temporal profile
of the 2^{nd} principal component (E_2) obtained from EMG matched the 2^{nd} principal
component (M_2) obtained from muscle torques and the 1^{st} principal component (D_1)
obtained from the dynamic components only of the muscle torques (MT_D) although with
a phase shift (Fig 2B). This feature captures the focal movement-related changes in
EMG that underly bursts in EMG profiles. The VAF by the 1^{st} principal components
obtained from muscle torques (M_1, P_1, and D_1) was very high (Fig. 2C), the mean values
across conditions and all participants were 90% for M_1, 97% for P_1, and 88% for D_1
(Table 1 contains values per condition). This shows that the profile of muscle torques
was very similar across joints and DOFs, indicating a high degree of intralimb
coordination. P_1 captured more variance across conditions and DOFs than M_1, similarly
D_1 captured more variance across conditions and DOFs than M_2 despite similar profiles.
This shows that subdividing muscle torques into postural and dynamic components
results in more reliable features compared to those obtained with PCA of muscle
torques. As expected, the EMG VAF by the 1^{st} principal component was lower than the
muscle torque VAF by the 1^{st} principal component due to higher noise in EMG data (Fig.
2C), the mean VAF across conditions and participants was 53% for E_1, but the opposite
was true for the 2^{nd} principal component: mean VAF by E_2 was 15% while the mean
VAF for M_2 was 7%. Moreover, both EMG and muscle torque VAF by the corresponding
principal components were similar across limbs and workspace s (Fig. 2C). This
indicates that highly stereotypical muscle torques during reaching movements explain a
high portion of the overall variance in EMG profiles in multiple arm muscles. The same
conclusion was reached in Olesh et al. [12] from a smaller subset of data. Moreover, our
results further support the idea that E_1 captures the muscle recruitment necessary for

generating postural forces to support the arm against gravity, while E_2 captures the
muscle recruitment necessary for generating dynamic forces for propulsion during
acceleration and deceleration phases of reach. This supports the first hypothesis that
the muscle activity profiles reflect the forces needed to produce the movement.
The degree of similarity between features obtained with PCA was compared
statistically using cross correlation analysis as described in Methods. RM ANOVA on
the coefficients of determination (R^2) between principal component profiles showed a
significant main effect F(15, 120) = 11.39, p = 0.00. Post-hoc tests have shown that
there were no differences between the R^2 values across conditions (Table 2) and that
the temporal profiles of E_1, M_1, and P_1 were more similar to each other than to E_2, M_2, or
D_1 and vice versa (Fig. 3A; Table 3). Similar results were observed when data from all
14 individuals who performed reaches from only the lateral workspace were analyzed
(Supplementary Fig. S1). The RM ANOVA main effect was significant (F(7, 84) = 12.62,
p = 0.00) and the post-hoc test did not reject the null hypothesis that the data from LLat
and RLat conditions comes from the same distribution (MSE = 0.02 ± SE 0.02; p =
0.41). The temporal profiles of E_1, M_1, and P_1 were more similar to each other than to
E_2, M_2, or D_1 and vice versa (Supplementary Fig. S1A; Table 4). This further supports
our first hypothesis and the idea that the first and second principal components of EMG
represent postural and dynamic forces. These conclusions generalize across both limbs
and workspace s. This may be useful for removing noise from muscle activity signals
and increasing the interpretability of surface EMG recordings.
Additionally, we observed a consistent phase lead of the D_1 components onto the
E_2 component (Fig. 3B, Supplementary Fig. S1B). This phase lead can potentially be
taken advantage of to predict the changes in the underlying muscle forces or
contractions from dynamical simulations prior to observable motion and improve
performance of real-time control applications.
The second hypothesis was that muscle synergies captured with dimensionality
reduction methods represent force directions controlled by the CNS. The rationale for
the test of this hypothesis is that the synergies must reflect the dynamic requirements of
each reaching movement. This means that for movements requiring a larger change in
the postural moments, a consistent group of muscles will have higher scores for the
principal component representing that synergy (E_1) in the corresponding movement and
the same group of muscles will have lower scores in other movements with smaller
changes in postural moments. The same logic applies to the synergy responsible for the
generation of dynamic forces (E_2). This means that if the primary features in EMG (E_1
and E_2) represent muscle synergies, we expect significant linear correlations between
scores of pairs of muscles across movement directions. We performed this linear
correlation analysis by comparing scores between pairs of muscle across reaching in
multiple directions (Fig. 4 shown an example for one condition and participant). The
scores represent the projection of EMG onto the principal component vectors. When the
scores are close to 0, this means that the components do not capture much variance in
the corresponding EMG profile. For all combinations of muscles, movements, and
participants, about 21% of the E_1 scores were low, i.e., values were below 5% of
maximal score (RLat: 25%, LLat: 18%, LMed: 16%, and RMed 24%). About 26% of E_2
scores were low (RLat: 27%, LLat: 26%, LMed: 22%, and RMed: 28%). The low scores
are likely noise. Therefore, no relationships between the low score values are expected,

as seen for example in the insignificant correlation between low scores of BicL and ADel
in the example shown in Fig. 4. Despite the noise, we observed multiple correlations
between E_1 scores in certain conditions. For reaching within the left lateral workspace,
68% of muscles (45 muscle pairs out of 66) in the non-dominant arm showed moderate
correlations between their E_1 scores ($R^2 \geq 0.5$; Fig. 5A). For reaching within the medial
workspace, 35% of muscles (23/66 pairs) in the non-dominant arm showed moderate
correlations between their E_1 scores (Fig. 5B). In contrast, only 12% of muscles in the
dominant arm showed moderate correlations when reaching within the right lateral
workspace (Fig. 5C). For reaching within the shared medial workspace, 32% of muscles
in the dominant arm showed moderate correlations (Fig. 5D), similar to that for the non-
dominant arm (Fig. 5B). Overall, more correlations were observed between E_1 scores in
muscles of the non-dominant arm. There were fewer muscle pairs with consistently
correlated E_2 scores (Fig. 6). The correlations between E_2 scores were much lower, so
that on average none of the muscle pairs showed moderate correlations across all 4
conditions. However, despite the higher percentage of low E_2 scores, there were
significant correlations between select muscle pairs (see below) with more correlations
occurring between E_2 scores in muscles of the dominant arm compared to the non-
dominant arm, in contrast to the correlations between E_1 scores. The same results were
obtained from the data across 14 participants who performed the RLat and LLat
conditions only (Supplementary Figures S2). The larger number of subjects included in
the analysis across lateral workspaces reduced the number of significant correlations
between E_1 scores in the muscles of the non-dominant arm only (Fig. 5A vs.
Supplementary Figure S2A). Overall, these results reject the second hypothesis that
specific force directions are controlled through muscle synergies.
Correlations between scores were overwhelmingly positive, indicating that not
only agonistic but also antagonistic muscles, such as biceps and triceps, changed their
activity together. The coactivation between muscles of the non-dominant arm
associated with counteracting gravity load on the limb (correlations between E_1 scores)
was the broadest in the LLat condition. Two salient muscle groups comprised primarily
the proximal muscles (TerM, ADel, PDel, TriL, and TriS) and distal muscles (BiL, Brd,
FCR, FCU, and ECR), both highly coupled (Fig. 5A). In contrast, the coactivation
between muscles of the dominant arm in RLat condition consisted of three much
smaller muscle groups (PDel/TriS, TriS/BicL/Brd, and FCR/FCU) with only the first two
groups coupled (Fig. 5C). These three small clusters were the only consistent groups of
coactivating muscles across both workspaces and both limbs. For forces needed to
propel the arm, the E_2 scores for three pairs of muscles of the dominant arm positively
correlated for reaching in the lateral workspace (R^2 for BicL/TriS = 0.49, Brd/TriS = 0.45,
and Brd/BicL = 0.45). There were 6 significantly correlated muscle pairs in RLat
condition that were similar across at least 5/10 subjects (Brd/ECR, FCR/FCU, Brd/BicL,
Brd/TriS, BicL/BicS, and BicL/TriS) and 4 significantly correlated muscle pairs in RMed
condition (Brd/ECR, FCR/FCU, Brd/TriS, and TriS/TriL). For the non-dominant arm,
there were 2 significantly correlated muscle pairs in LLat condition that were similar
across 6/10 subjects (Brd/TriS and BicL/TriS) and 1 significantly correlated muscle pair
in LMed condition (BicL/TriS). Note that in these reaching movements with the arm
pronated, the muscles performing the antigravity action would be ADel, BiL, Brd, and
ECR, while the rest would produce joint moments in the direction of gravity. Here we

observed consistent coactivation of not only antigravity agonists, but also antagonists,
such as biceps and triceps. Such coactivation of antagonistic muscles modulates joint
stiffness, suggesting that this may be the control strategy for supporting the body
against gravity.
399

Discussion

We observed that highly stereotypical muscle torques during reaching
movements explain a high portion of the overall variance in EMG profiles in multiple arm
muscles. We have shown that the first principal component of EMG captures the muscle
recruitment necessary for generating postural forces to support the arm against gravity,
while the second principal component of EMG captures the muscle recruitment
necessary for generating dynamic forces for propulsion during acceleration and
deceleration phases of reach (Fig. 2, 3, Supplementary Fig. S1). This supports our first
hypothesis showing that the first principal components of EMG represent postural and
dynamic forces. This shows that subdividing muscle torques into postural and dynamic
components captures muscle contributions to postural and propulsive forces. These
conclusions, originally derived from a smaller subset of movements by the dominant
arm in one area of reaching workspace [12], are generalizable to movements by the
non-dominant arm and across a larger reaching workspace that overlaps between both
arm arms. This may be useful for removing noise from muscle activity signals and
increasing the interpretability of surface EMG recordings.
Moreover, we observed biomechanically expected correlations between scores
of muscle pairs across movements in very few conditions (Fig. 5, 6, Supplementary Fig.
S2). This shows that when joint torques in specific directions (flexion or extension) were
required to reach toward a target, muscles were *not* recruited in consistent groups in
most conditions. This result rejects the second hypothesis that specific force directions
are controlled through muscle synergies. When correlations between scores were
observed when producing postural forces by the non-dominant arm, they were
overwhelmingly positive, despite the antagonistic actions between multiple muscle pairs
in our dataset. This suggests that the arm is supported against gravity not through the
production of reaction forces with antigravity muscles, but rather through the modulation
of limb stiffness.
It is possible that the reason we have not obtained consistent muscle synergies
across movements is that surface EMG was too noisy and variable across the different
movements and individuals. However, our variability data reported in Results shows that
the temporal profiles of EMG were very consistent across all these conditions,
especially for the dominant arm with the fewest observed coactivations. Another
evidence of the consistency of EMG profiles is the large percentage of variance that is
captured in EMG by the first two principal components (Fig. 2) and by the components
obtained from muscle torques (Fig. 3), which are highly stereotypical during reaching
movements. Therefore, it seems highly unlikely that the inter-movement or inter-subject
variability can explain the lack of consistent coactivation of muscle pairs for the
production of joint torque in a given direction in most conditions.
Muscle synergies are often proposed as a method of reducing the dimensionality
of neural control space and overcoming the redundancy of the musculoskeletal system

[4–8]. We have posited that it is not sufficient to show the existence of a low
dimensional space, one needs to demonstrate the utility of the obtained synergies in
performing specific movements. Here we define muscle synergies as groups of muscles
controlled with a single control signal that produces joint torque about one or mode
DOFs in a specific direction. This definition is similar to force fields first defined by Bizzi
[17]. This definition implies that when movements require a force vector in the same
direction but of different amplitude, the recruitment level of the muscle synergy needed
to produce that force vector should also vary in scale with the amplitude of the force
vector across those movements. In other words, if a simple synergy exists for a single
input recruiting both biceps and brachioradialis to produce elbow flexion torque, then
these muscles should activate together less in movements needing lower elbow flexion
torque and more in movements needing higher elbow flexion torque. This logic is what
motivated our second hypothesis, which we tested using the regression analysis shown
in Fig. 4. The level of coactivation of muscle synergies was captured by the PCA
scores. Therefore, examining changes in PCA scores across different movement
directions, i.e. across different force directions underlying these movements, reveals the
pattern of synergistic coactivation of muscles. Surprisingly, we found consistent
coactivation of very few pairs of muscles and primarily in the component of muscle
activation responsible for overcoming gravity (Fig. 5, Supplementary Fig. 2A, B). Limited
consistent coactivations were observed in the phasic component of distal muscles
responsible for acceleration/deceleration forces (Fig. 6, Supplementary Fig. 2C, D). This
means that when joint moments in the same direction but different amplitudes need to
be produced for movements toward different targets, it is accomplished by activation of
different combinations of muscles requiring separate neural recruitment signals. These
results contradict the idea of muscle synergies driving specific force directions and lead
us to reject the second hypothesis.
Our results support the idea of internal models embedding body dynamics.
Internal models are defined as neural networks in the brain that simulate the dynamics
of the body and its interaction with the external world [1,3,18–21]. Forward models
predict the consequences of a given action, for example, predicting sensory feedback
without delay from an efferent copy of a corticospinal motor command [22,23]. Inverse
models capture information in the opposite direction [3,24]. They convert a motor plan
into the motor execution output of the motor system that converges on the motoneurons
in the spinal cord. The motor plan is thought to be developed in the premotor areas,
where neural activity has many kinematic features in the extrinsic head-centered
reference frame resolving the kinematic redundancy problem [25–27]. The motor
execution output is thought to originate in the primary motor cortex, where neural
activity has many dynamic features in the intrinsic reference frame [28–30]. The
transformations from motor plan into motor execution may potentially resolve the
muscle redundancy problem by embedding biomechanical constraints. These
constraints may take the form of embedded muscle anatomy relationships, such as
known ratios between moments arms of muscles about a given DOF [31].
Reaching movements are often separated into two distinct actions 1) the
transport action that involves gross stereotypical movement of the whole arm to
transport and orient the hand and 2) the manipulation action that involves fine
movements of the hand. This separation is reflected in the anatomical organization of

muscles that form *proximal* and *distal* groups that could serve as *separate control*
targets [9]. Interestingly, we observed here distinct correlations between the principal
component scores of proximal and distal muscle pairs, particularly in the non-dominant
arm for the gross postural component and in the dominant arm for the propulsive
component. The proximal group of muscles including deltoids, biceps short head, and
both heads of triceps, co-activated in proportion to the gravity-related forces acting on
the limb (top left corners of correlation matrices in Fig. 5A, B and Supplementary Fig.
S2A). The distal group of muscles included wrist flexor muscles and brachioradialis, it
co-activated to a lesser degree in proportion to the propulsion-related forces acting on
the limb, sometimes including the long head of biceps or the short head of triceps
(bottom right corners of correlation matrices in Fig. 5, 6, and Supplementary Fig. S2).
These clusters of proximal and distal muscle coactivations suggests that there may be
separate control signals responsible for activating proximal and distal musculature,
which are coordinated together when wrist stabilization is required. Altogether, this
evidence supports the idea of two interacting neural controllers targeting proximal and
distal muscle groups in the arm and hand performing gross postural and fine goal-
directed motor control during reaching [32]. This idea is supported by the neural
stimulation experiments showing that cortical muscle representations are organized
somatotopically based on their proximal-to-distal anatomy, the famous motor
homunculus [33,34]. Non-invasive stimulation of human motor cortex can facilitate
differentially proximal and distal muscles depending upon the demands of a particular
task, i.e., reaching vs. grasping [35]. Distinct white matter tracts demonstrate some
preferential contribution for proximally-controlled gross movements versus distally-
controlled fine motor skills [36]. The proximal controller may be more energetically
favorable for controlling whole arm impedance to support the limb against gravity [37].
The distal controller may be more attuned to stabilizing the hand during transport and
compensating for forces from external objects during their manipulation. The proposed
proximal and distal controllers may also underlie the differences in controlling limb
dynamics and adaptation to novel limb dynamics between dominant from nondominant
arms [38,39]. Overall, our data supports the idea of two neural controllers that target
different muscle groups in the arm and hand for gross postural and fine goal-directed
control of reaching.
518
519

Acknowledgements

VG was supported by NIGMS grants P20GM109098 and P30GM103503. ASK was
supported by a fellowship from NIGMS T32 AG052375. This work was supported in part
by the Office of the Assistant Secretary of Defense for Health Affairs through the
Restoring Warfighters with Neuromusculoskeletal Injuries Research Program
(RESTORE) under Award No. W81XWH-21-1-0138. Opinions, interpretations,
conclusions, and recommendations are those of the author and are not necessarily
endorsed by the Department of Defense.
528

Author Contributions

ASK contributed to conceptualization, data curation, data analysis, and writing. VG contributed to conceptualization, methodology development, data curation, data analysis, and writing.

Tables

Table 1. Variance accounted for across reaching conditions.

	Left lateral	Left medial	Right medial	Right lateral	Total
E_1	53.46 ± 8.83	48.88 ±10.51	57.05 ±5.05	53.91 ±8.43	53.33 ± 8.21
M_1	89.51 ± 4.35	88.33 ± 6.55	92.35 ± 3.77	91.04 ± 5.33	90.31 ± 5.00
P_1	98.09 ± 0.73	96.46 ± 3.70	96.63 ± 5.88	98.19 ± 1.02	97.34 ± 2.83
E_2	13.90 ± 4.58	13.39 ± 4.11	15.84 ± 4.93	16.79 ± 6.70	14.98 ± 5.08
M_2	6.99 ± 2.81	8.43 ± 4.93	5.49 ± 3.04	6.94 ± 4.67	6.96 ± 3.86
D_1	78.39 ± 3.83	90.25 ± 4.08	92.99 ± 2.85	91.27 ± 7.00	88.23 ± 4.44

Values are means ± standard deviation across 10 individuals.

Table 2. Post-hoc comparisons of Pearson's correlation coefficients across conditions.

Condition 1	Condition 2	MSE ± SE	p value
Llat	Lmed	-0.02 ± 0.03	0.95
Llat	Rlat	0.02 ± 0.03	0.91
Llat	Rmed	0.01 ± 0.03	0.99
Lmed	Rlat	0.03 ± 0.04	0.82
Lmed	Rmed	0.02 ± 0.03	0.88
Rlat	Rmed	-0.01 ± 0.02	0.95

MSE – mean squared error, SE – standard error of the mean. Adjusted alpha = 0.0085.

Table 3. Post-hoc comparisons of Pearson's correlation coefficients between EMG- and torque-based components across four conditions and 10 participants.

Condition 1	Condition 2	MSE ± SE	p value
E_1/M_1	E_1/P_1	-0.01 ± 0.01	0.7739
E_1/M_1	E_2/M_2	0.19 ± 0.03	**0.0018**
E_1/M_1	E_2/D_1	0.29 ± 0.05	**0.0014**
E_1/P_1	E_2/M_2	0.20 ± 0.04	**0.0068**
E_1/P_1	E_2/D_1	0.31 ± 0.06	**0.0069**
E_2/M_2	E_1/P_1	-0.20 ± 0.04	**0.0069**
E_2/M_2	E_2/D_1	-0.10 ± 0.04	0.1772

MSE – mean squared error, SE – standard error of the mean. Bold p values show significant difference with adjusted alpha = 0.0073.

Table 4. Post-hoc comparisons of Pearson's correlation coefficients between EMG- and torque-based components across two conditions and 14 participants.

Condition 1	Condition 2	MSE ± SE	p value
E_1/M_1	E_1/P_1	0.00 ± 0.01	0.9999
E_1/M_1	E_2/M_2	0.15 ± 0.03	**0.0006**
E_1/M_1	E_2/D_1	0.23 ± 0.05	**0.0016**
E_1/P_1	E_2/M_2	0.15 ± 0.04	**0.0044**
E_1/P_1	E_2/D_1	0.23 ± 0.05	**0.0028**
E_2/M_2	E_1/P_1	-0.15 ± 0.04	**0.0044**
E_2/M_2	E_2/D_1	-0.08 ± 0.04	0.2435

MSE – mean squared error, SE – standard error of the mean. Bold p values show significant difference with adjusted alpha = 0.0073.

Figure Legends

Fig 1. **Examples of kinematics, dynamics, and muscle activity profiles during reaching.** Each plot shows profiles per condition averaged across repetitions (n = 15) of the same reaching movement in one direction by one participant. (**A**) Profiles (solid lines) and standard deviation (shaded areas) of shoulder flexion-extension angle and angular velocity. (**B**) Shoulder flexion-extension torque (t) profiles that caused the movements in (**A**). (**C**) Normalized electromyography (EMG) profiles of anterior deltoid (ADel), triceps long (TriL), and biceps long (BiL) that accompanied the movements in (**A**). Shaded areas show the standard error for the mean across repetitions (n = 10) of the same movement.

Fig 2. Principal components of EMG and muscle torques. Data from the four conditions indicated by the pictograms are shown in plots arranged in columns. The data is from 10 participants who performed all 4 conditions. (**A**) Solid lines show profiles of the 1^{st} principal component obtained from EMG (E_1), muscle torque (M_1), and the postural component of muscle torque (P_1). (**B**) Solid lines show profiles of the 2^{nd} principal component obtained from EMG (E_2) and muscle torque (M_2), and the 1^{st} principal component obtained from the dynamic component of muscle torque (D_1). (C) Variance accounted for (VAF) by principal components obtained from EMG and muscle torques.

Fig 3. Correlations between the temporal profiles of principal components derived from EMG and muscle torques. Data (N=10) from the four conditions indicated by the pictograms are shown in plots arranged in columns. (A) Dots show individual R^2 values from cross-correlations between profiles, red lines show median values, and grey boxes show interquartile ranges. Brackets with stars indicate significant relationships with correction for familywise error. (B) Dots show individual lag values from cross correlations, red lines show median values, and grey boxes show interquartile ranges. Open dots are outliers. E_1 and E_2 are the 1^{st} and 2^{nd} principal components respectively obtained from EMG; M_1 and M_2 are the 1^{st} and 2^{nd} principal

584 components respectively obtained from muscle torques; P_1 and D_1 are the 1st principal
585 components obtained from the postural and dynamic components of muscle torque
586 respectively.
587
Fig 4. Example linear regressions between scores across reaching directions.
Example regression matrix between E_1 scores of 12 muscles across 28 right-handed
reaches within the medial workspace (RMed condition) for a single participant. The
coordinates of each dot represent principal component scores for two muscles for a
single movement direction. Histograms show the distribution of the scores for a given
muscle across reaching directions. Solid lines show least-squares linear regression, red
lines indicate significant relationships with correction for familywise error, so that the
adjusted alpha = 0.0008.
596
Fig 5. Posture-related muscle coactivation across all conditions. Heatmaps show
coefficients of determination (R^2) from regressions shown in Fig. 4 averaged across 10
participants. Blue colors represent minor or no significant relationships between E_1
coefficients across movement directions, while red and yellow colors represent
moderate and strong relationships. Pictograms indicate conditions, i.e., LLat (A), LMed
(B), RLat (C), and RMed (D). Muscles are abbreviated as described in Methods.
603
Fig 6. Propulsion-related muscle coactivation across all conditions. Heatmaps
show coefficients of determination (R^2) from regressions shown in Fig. 4 averaged
across 10 participants. Blue colors represent minor or no significant relationships
between E_2 coefficients across movement directions, while red and yellow colors
represent moderate and strong relationships. Pictograms indicate conditions, i.e., LLat
(A), LMed (B), RLat (C), and RMed (D). Muscles are abbreviated as described in
Methods.
611
Supplementary Figure S1. Correlations between the temporal profiles of principal
components derived from EMG and muscle torques across both limbs. Data (N =
14) from the two conditions indicated by the pictograms are shown in plots arranged in
columns. (A) Dots show individual R^2 values from cross-correlations between profiles,
red lines show median values, and grey boxes show interquartile ranges. Brackets with
stars indicate significant relationships with correction for familywise error. (B) Dots show
individual lag values from cross correlations, red lines show median values, and grey
boxes show interquartile ranges. E_1 and E_2 are the 1st and 2nd principal components
respectively obtained from EMG; M_1 and M_2 are the 1st and 2nd principal components
respectively obtained from muscle torques; P_1 and D_1 are the 1st principal components
obtained from the postural and dynamic components of muscle torque respectively.
623
Supplementary Figure S2. Posture- and propulsion-related muscle coactivation
across the two lateral conditions. Heatmaps show coefficients of determination (R^2)
averaged across 14 participants. Blue colors represent minor or no significant
relationships between E_1 (A, B) and E_2 (C, D) coefficients across movement directions,
while red and yellow colors represent moderate and strong relationships. Pictograms

indicate conditions, i.e., LLat (A, C) and RLat (B, D). Muscles are abbreviated as
described in Methods.
631

References

1. Kawato M, Furukawa K, Suzuki R. A hierarchical neural-network model for control
and learning of voluntary movement. Biol Cybern. 1987;57: 169–185.
doi:10.1007/BF00364149

2. Wolpert DM, Ghahramani Z. Computational principles of movement neuroscience.
Nat Neurosci. 2000;3: 1212–1217. doi:10.1038/81497

3. Wolpert DM, Kawato M. Multiple paired forward and inverse models for motor
control. Neural Networks. 1998;11: 1317–1329. doi:10.1016/S0893-6080(98)00066-
5

4. d'Avella A, Lacquaniti F. Control of reaching movements by muscle synergy
combinations. Front Comput Neurosci. 2013;7. doi:10.3389/fncom.2013.00042

5. Mussa-Ivaldi FA. Modular features of motor control and learning. Current Opinion in
Neurobiology. 1999;9: 713–717. doi:10.1016/S0959-4388(99)00029-X

6. Thoroughman KA, Shadmehr R. Learning of action through adaptive combination of
motor primitives. Nature. 2000;407: 742–747. doi:10.1038/35037588

7. Bizzi E, Tresch MC, Saltiel P, d'Avella A. New perspectives on spinal motor
systems. Nat Rev Neurosci. 2000;1: 101–108. doi:10.1038/35039000

8. d'Avella A, Saltiel P, Bizzi E. Combinations of muscle synergies in the construction
of a natural motor behavior. Nat Neurosci. 2003;6: 300–308. doi:10.1038/nn1010

9. Gritsenko V, Hardesty RL, Boots MT, Yakovenko S. Biomechanical constraints
underlying motor primitives derived from the musculoskeletal anatomy of the human
arm. PLoS ONE. 2016;11: 1–18. doi:10.1371/journal.pone.0164050

10. Taitano RI, Yakovenko S, Gritsenko V. Muscle anatomy is reflected in the spatial
organization of the spinal motoneuron pools. Commun Biol. 2024;7: 97.
doi:10.1038/s42003-023-05742-w

11. Tresch MC, Jarc A. The case for and against muscle synergies. Current Opinion
in Neurobiology. 2009;19: 601–607. doi:10.1016/j.conb.2009.09.002

12. Olesh EV, Pollard BS, Gritsenko V. Gravitational and dynamic components of
muscle torque underlie tonic and phasic muscle activity during goal-directed
reaching. Front Hum Neurosci. 2017;11: 474. doi:10.3389/fnhum.2017.00474

13. Luca CJD. The Use of Surface Electromyography in Biomechanics. Journal of
Applied Biomechanics. 1997;13: 135–163. doi:10.1123/jab.13.2.135

14. Talkington WJ, Pollard BS, Olesh EV, Gritsenko V. Multifunctional setup for
studying human motor control using transcranial magnetic stimulation,
electromyography, motion capture, and virtual reality. JoVE. 2015; 52906.
doi:10.3791/52906

15.   Winter DA. Biomechanics and Motor Control of Human Movement. 4th ed. Wiley;
2009.   Available: https://www.wiley.com/en-
us/Biomechanics+and+Motor+Control+of+Human+Movement%2C+4th+Edition-p-
9780470398180

16.   Hermens HJ, Merletti R, Freriks B. SENIAM□: European recommendations for
surface electromyography. 2d edition. Enschede, The Netherlands: Roessingh
Research and Development; 1999.

17.   Hogan N, Bizzi E, Mussa-Ivaldi FA, Flash T. Controlling multijoint motor behavior.
Exerc Sport Sci Rev. 1987;15: 153–190.

18.   Cisek P. Internal Models. In: Binder MD, Hirokawa N, Windhorst U, editors.
Encyclopedia of Neuroscience. Berlin, Heidelberg: Springer; 2009. pp. 2009–2012.
doi:10.1007/978-3-540-29678-2_2528

19.   Wolpert D, Ghahramani Z, Jordan M. An internal model for sensorimotor
integration. Science. 1995;269: 1880–1882. doi:10.1126/science.7569931

20.   Hwang EJ, Shadmehr R. Internal models of limb dynamics and the encoding of
limb state. J Neural Eng. 2005;2: S266–S278. doi:10.1088/1741-2560/2/3/S09

21.   Zago M, Lacquaniti F. Visual perception and interception of falling objects: a
review of evidence for an internal model of gravity. J Neural Eng. 2005;2: S198–
S208. doi:10.1088/1741-2560/2/3/S04

22.   Haggard P, Whitford B. Supplementary motor area provides an efferent signal for
sensory   suppression.   Cognitive   Brain   Research.   2004;19:   52–58.
doi:10.1016/j.cogbrainres.2003.10.018

23.   Chapman CE. Active versus passive touch: factors influencing the transmission
of somatosensory signals to primary somatosensory cortex. Can J Physiol
Pharmacol. 1994;72: 558–570. doi:10.1139/y94-080

24.   Schweighofer N, Arbib MA, Kawato M. Role of the cerebellum in reaching
movements in humans. I. Distributed inverse dynamics control. European Journal of
Neuroscience. 1998;10: 86–94. doi:10.1046/j.1460-9568.1998.00006.x

25.   Crammond DJ, Kalaska JF. Prior Information in Motor and Premotor Cortex:
Activity During the Delay Period and Effect on Pre-Movement Activity. Journal of
Neurophysiology. 2000;84: 986–1005. doi:10.1152/jn.2000.84.2.986

26.   Cisek P, Kalaska JF. Neural correlates of mental rehearsal in dorsal premotor
cortex. Nature. 2004;431: 993–996. doi:10.1038/nature03005

27.   Scott SH, Sergio LE, Kalaska JF. Reaching Movements With Similar Hand Paths
but Different Arm Orientations. II. Activity of Individual Cells in Dorsal Premotor
Cortex and Parietal Area 5. Journal of Neurophysiology. 1997;78: 2413–2426.
doi:10.1152/jn.1997.78.5.2413

28.   Scott SH, Kalaska JF. Reaching Movements With Similar Hand Paths But
Different Arm Orientations. I. Activity of Individual Cells in Motor Cortex. Journal of
Neurophysiology. 1997;77: 826–852. doi:10.1152/jn.1997.77.2.826

29. Ashe J. Force and the motor cortex. Behav Brain Res. 1997;87: 255–269.
doi:10.1016/s0166-4328(97)00752-3

30. Kalaska JF. From Intention to Action: Motor Cortex and the Control of Reaching
Movements. In: Sternad D, editor. Progress in Motor Control: A Multidisciplinary
Perspective. Boston, MA: Springer US; 2009. pp. 139–178. doi:10.1007/978-0-387-
77064-2_8

31. Gritsenko V, Hardesty RL, Boots MT, Yakovenko S. Biomechanical Constraints
Underlying Motor Primitives Derived from the Musculoskeletal Anatomy of the
Human Arm. Jones KE, editor. PLoS ONE. 2016;11: e0164050.
doi:10.1371/journal.pone.0164050

32. Lemon R, Johansson R, Westling G. Corticospinal control during reach, grasp,
and precision lift in man. J Neurosci. 1995;15: 6145–6156.
doi:10.1523/JNEUROSCI.15-09-06145.1995

33. Penfield W, Rasmussen T. The cerebral cortex of man: A Clinical Study of
Localization of Function. 1950.

34. Wassermann EM, McShane LM, Hallett M, Cohen LG. Noninvasive mapping of
muscle representations in human motor cortex. Electroencephalography and Clinical
Neurophysiology/ Evoked Potentials. 1992. doi:10.1016/0168-5597(92)90094-R

35. Schieppati M, Trompetto C, Abbruzzese G. Selective facilitation of responses to
cortical stimulation of proximal and distal arm muscles by precision tasks in man.
Journal of Physiology. 1996;491: 551–562. doi:10.1113/jphysiol.1996.sp021239

36. Lawrence DG, Kuypers HG. The functional organization of the motor system in
the monkey. I. The effects of bilateral pyramidal lesions. Brain□: a journal of
neurology. 1968;91: 1–14.

37. Nguyen HP, Dingwell JB. Proximal Versus Distal Control of Two-Joint Planar
Reaching Movements in the Presence of Neuromuscular Noise. Journal of
Biomechanical Engineering. 2012;134: 061007. doi:10.1115/1.4006811

38. Sainburg RL. Evidence for a dynamic-dominance hypothesis of handedness. Exp
Brain Res. 2002;142: 241–258. doi:10.1007/s00221-001-0913-8

39. Bagesteiro LB, Sainburg RL. Handedness: Dominant Arm Advantages in Control
of Limb Dynamics. Journal of Neurophysiology. 2002;88: 2408–2421.
doi:10.1152/jn.00901.2001