Differential Impact of Biomechanical Constraints on Control Signal Dimensionality for Gravity Support Versus Propulsion

AS Korol and V Gritsenko

Keywords: *Muscle co-contraction, EMG, gravitational torque component, dynamical torque component, active muscle torque.*

Anna S Korol
Department of Neuroscience, School of Medicine, West Virginia University
Rockefeller Neuroscience Institute, West Virginia University
Morgantown, USA
ORCID: 0000-0001-9699-721X

Valeriya Gritsenko
Department of Human Performance, Division of Physical Therapy, School of Medicine, West Virginia University
Department of Neuroscience, School of Medicine, West Virginia University
Rockefeller Neuroscience Institute, West Virginia University
Morgantown, USA
ORCID: 0000-0002-6408-9433

Abstract

Neural control of movement has to overcome the problem of redundancy in the multidimensional musculoskeletal system. The problem can be solved by reducing the dimensionality of the control space of motor commands, i.e., through muscle synergies or motor primitives. Evidence for this solution exists, multiple studies have obtained muscle synergies using decomposition methods. These synergies vary across different workspaces and are present in both dominant and non-dominant limbs. Here we explore the dimensionality of control space by examining muscle activity patterns across reaching movements in different directions starting from different postures performed bilaterally by healthy individuals. We further explore the effect of biomechanical constraints on the dimensionality of control space. We are building on top of prior work showing that muscle activity profiles can be explained by applied moments about the limb joints that reflect the biomechanical constraints. These muscle torques derived from motion capture represent the combined actions of muscle contractions that are under the control of the nervous system. Here we test the generalizability of the relationship between muscle torques and muscle activity profiles across different starting positions and between limbs. We also test a hypothesis that the dimensionality of control space is shaped by biomechanical constraints. We used principal component analysis to evaluate the contribution of individual muscles to producing muscle torques across different workspaces and in both dominant and non-dominant limbs. Results generalize and support the hypothesis. We show that the muscle torques that support the limb against gravity are produced by more consistent combinations of muscle co-contraction than those that produce propulsion. This effect was the strongest in the non-dominant arm moving in the lateral workspace on one side of the body.

Keywords: electromyography, neuromechanics, limb dynamics, hand dominance, gravity, postural forces

Introduction

The production of movement involves integrating biomechanical, neural, and environmental factors. Muscle recruitment by the central nervous system (CNS) causes motion or lack thereof, such as when maintaining posture, through the nonlinear production of forces by muscles about the degrees of freedom (DOFs) of the joints. Body inertia, the number of DOFs, and external forces then shape the resulting posture and movement. This biomechanics is complex enough that neural sensorimotor circuits must embed its dynamics for efficient and robust control [1–3]. However, a problem of redundancy exists, i.e., the problem of choosing among multiple muscles and combinations of joint angles that are possible for a given desired hand position or motion. This problem may be resolved by reducing the dimensionality of the space of motor commands by the CNS, a concept known as muscle synergies or motor primitives. This concept implies that muscles (or motoneurons in the spinal cord) are recruited in groups that represent a specific action so that a combination of the smaller number of synergies can produce a larger number of different movements [4–8]. The anatomical organization of muscles that form agonistic and antagonistic groups that can be activated by fewer control signals constraining the dimensionality of control space [9]. The spatial organization of motoneuron pools in the spinal cord also captures the functional relationships of the muscles they innervate, further supporting the idea of reduced dimensionality of control space [10]. Indeed, the neural activation of muscles observed with surface electromyography (EMG) is reducible to a low dimensional space for certain types of movements, including reaching [4,8,11]. However, the dimensionality reduction can be done using different mathematical methods applied to data organized in different ways [12,13]. These methods all obtain control space solutions with lower dimensions than the original dataset, but their neuromuscular or biomechanical underpinnings are not always clear.

Animal studies have worked out the basic modular and hierarchal organization of the neural motor control system that comprises nested feedback loops from sensory afferents to spinal motor neurons that encompass progressively more delayed processing circuits including the spinal reflexes, reticular formation in the brainstem, and primary sensory and motor cortices in the brain to name a few [14,15] (Fig 1A). Relating the synergies obtained with decomposition methods to the dynamic neural control units is not trivial. The commonly used Principal Component Analysis (PCA) can be applied across muscles, task conditions, and time in two ways, to obtain temporally-variable synergies (temporal synergies) or temporally invariant synergies (spatial synergies) [16]. A recent study has shown that the temporal dynamics of the spinal central pattern generator that controls rhythmic movements is captured well using temporal synergies obtained with PCA of EMG during locomotion [17]. We have also shown that the temporal dynamics of muscle torques needed to support the limb against gravity and for propulsion are well described by the 1st and 2nd temporal synergies respectively obtained with PCA. These two components of muscle action may rely on postural mechanisms governed by the pontomedullary reticular formation and the phasic velocity-dependent output of the primary motor cortex respectively [18–21]. For reaching, the temporal synergies have been shown to be more accurately capturing the temporal aspects of muscle activity and potentially neural control dimensionality [16].

A

corticospinal tract

reticulospinal tract

motoneurons

B

Pec

PDel
TerM

ADel

BicL
BicS

TriL
TriS

Brd

ECR

FCR
FCU

The temporal profile of the envelope of surface EMG is closely related to the force
the corresponding muscle is producing in response to neural activation [22]. More recently
we have shown that for reaches with the dominant hand towards visual targets in three-
dimensional space, the EMG profiles are closely related to the muscle torques that cause
motion [23]. Specifically, the static component of EMG that underlies postural forces
needed to support the arm in specific position and during transitions between positions is
closely related to the components of muscle torques that include gravity terms in the
equations of motion. This suggests that the static component of EMG envelope from a
given muscle reflects the muscle's contribution to counteracting gravity load on the joints
it spans. Moreover, the residual phasic component of EMG is closely related to the
residual muscle torque that underlies the acceleration and deceleration forces toward the
reaching goal after gravity-related component is subtracted. This suggests that the phasic
component of EMG envelope from a given muscle reflects the muscle's contribution to
propulsion toward the target and stopping there. Here we will test the generalizability of
these relationships between muscle activity and torque profiles to the non-dominant limb
and across different workspaces.
Dimensionality reduction of control space can be observed through the co-
contraction of multiple muscles. For example, a single control signal can activate both the
brachioradialis and biceps muscles to produce an elbow flexion moment. This is the co-
contraction of agonistic muscles. Moreover, that same control signal activating the triceps
muscle, which opposes the action of the elbow flexors mentioned above, will both reduce
the elbow flexion torque and increase the stiffness of elbow and shoulder joints. This is
the co-contraction of antagonistic muscles. Both types of co-contraction are observed
during reaching. The agonistic co-contraction together with reciprocal activation of
antagonistic muscles contribute directly to muscle torques computed with inverse models
and are likely facilitated through spinal reflexes and CPG. For example, stimulation of the
primary afferents from the medial gastrocnemius muscle produces EPSPs not only in the
motoneurons innervating that same muscle but also in motoneurons innervating lateral
gastrocnemius and soleus, contributing to agonistic co-contraction [24]. The levels of
antagonistic co-contraction are thought to vary with the dynamical requirements of the
movement that vary with reaching in different directions from different starting postures
[21,25]. While the CNS is thought to control whole limb stiffness through co-contraction
[26,27], we do not know how this co-contraction signal is generated by the nervous
system to help reduce the dimensionality of control space.
In this study, healthy human participants performed unconstrained pointing
movements toward visual targets located equidistantly from one of two central targets.
The reaching movements in multiple directions were performed withing the typical
reaching workspace by either dominant (right) or non-dominant (left) arm. We collected
motion capture and EMG data and ran dynamic simulations with individualized inertial
models of the arm to compute muscle torques and their components. We first tested the
generalizability of the conclusions from Olesh et al. [23] by comparing principal
components obtained from muscle torques with EMG profiles across limbs and
workspaces. Here PCA derived scores that represent the projection of each EMG profile
in the principal component space. We applied PCA with the assumption of invariant
activation profiles to obtain temporal synergies as described above. The scores derived
this way will vary across movement directions and muscles but remain constant across

time. Then when muscle activity scales together across movement directions we expect
the scores of the corresponding muscles to be linearly related. When the muscles whose
scores are correlated are agonists, this will capture agonistic co-contraction. Similarly,
when the muscles whose scores are correlated are antagonists, this will capture
antagonistic co-contraction. Overall, correlations between muscle scores will reveal co-
contraction. The more muscles are co-contracting, the lower is the dimensionality of
control. The levels of co-contraction were compared between limbs and workspaces.
Using this method, we tested the hypothesis that the dimensionality of control space is
shaped by biomechanical constraints.
128

Materials and Methods

Data Collection

The experimental protocol was approved by the Institutional Review Board of West
Virginia University (Protocol #1311129283). Perspective participants were recruited
through fliers distributed around Morgantown, WV. The data collection started on March
12th, 2014, and ended on July 25th, 2016. Participants provided a written informed consent
prior the start of experiments. A second member of the investigative team witnessed the
signing of the consent form.
Nine healthy participants (mean age ± SD = 22.78 ± 0.67 years, 3 females and 6
males, sex assigned at birth) performed a modified center-out reaching task by pointing
to visual targets in virtual reality (software Vizard by Worldviz, Oculus Rift) [23]. All
participants reported they were right-handed and had no neurological or musculoskeletal
conditions that could alter movement. Participants repeated reaching movements
between a central target and one of 14 targets located along a sphere, with 8 targets
placed equidistantly on a horizontal circle parallel with the floor and 8 targets placed
equidistantly on a vertical circle parallel to the body sagittal plane, with 2 targets sharing
locations in both circles as in Olesh *et al.* [23]. Participants were instructed to point to
targets with the index finger, while holding their hand palm down, as quickly and
accurately as they can without moving their trunk and wrist. The distances toward the
targets were normalized for each subject based on their arm segment lengths to minimize
the inter-subject variability in angular kinematics. Each movement to and from each target
was repeated 15 times in a randomized order.
The center-out and return movements were repeated with each arm starting at one
of two locations of the central target. The central target was placed at the same distance
relative to the participant's shoulder scaled to their arm length. This further minimized the
inter-subject variability in angular kinematics. In the lateral starting position, the central
target was placed at a location that positioned the shoulder at 0 angle of all degrees of
freedom and elbow at 90 degrees, so that the upper arm was parallel to the trunk and the
forearm was parallel to the floor (Fig 2, pictograms in top corners, view from above). In
the medial starting position, the central target was placed at a location that positioned the
hand at the midline of the body and at the same distance from the trunk as in the lateral
starting position (Fig 2, pictograms in top center, view from above). The medial starting
position was the same for reaches with both right and left arms representing common
medial workspace. Thus, four conditions were created based on the location of the central
target and the arm used to make reaches, i.e., LLat and RLat for reaches by left and right

arm respectively in their respective lateral workspaces and LMed and RMed for the
reaches by left and right arm respectively in the common medial workspace.
Movement was recorded and visualized in virtual reality using the Impulse system
(PhaseSpace Inc) at the temporal resolution of 480 Hz. We tracked the location of 9 light
emitting diodes placed on the bony landmarks of the trunk, arm, and hand (Fig 1B) with
8 cameras as described in detail in Olesh et al. [23]. EMG was recorded at the temporal
resolution of 2,000 Hz using MA400-28 (MotionLab Systems). EMG was captured from
12 arm muscles: the clavicular head of pectoralis (Pec), teres major (TerM), anterior
deltoid (ADel), posterior deltoid (PDel), the long and lateral heads of triceps (TriL and
TriS), the short and long heads of biceps (BiS and BiL), brachioradialis (Brd), flexor capri
radialis (FCR), flexor carpi ulnaris (FCU), and extensor capri radialis (ECR). EMG
recordings were temporally synchronized with the motion capture and virtual reality
systems during reaching as described in Talkington *et al.* [28].
177

Data Analysis

All data analysis was carried out in MATLAB 2023b (MathWorks Inc). Motion
capture data were low pass filtered at 10 Hz and interpolated with a cubic spline. Arm
kinematics was obtained from motion capture by defining local coordinate systems
representing the trunk, arm, forearm, and hand. Joint angles representing 5 DOFs of the
arm were derived using linear algebra, namely shoulder flexion-extension, shoulder
abduction-adduction, shoulder internal-external rotation, elbow flexion-extension, forearm
pronation-supination, and wrist flexion-extension (Fig 2A, example for one DOF). Active
muscle torques and their components were calculated from kinematic data using a
dynamic model of the arm with 5 DOFs (Simulink, MathWorks Inc) as described in [23].
The individual height, weight, and segment lengths together with the published
anthropometric proportions were used to scale the model inertia to individual body sizes
[29]. Inverse simulations were run with the angular kinematics as input and the active
torques produced by muscles, termed muscle torques (MT), as output. Simulations were
ran with and without gravity force to estimate the components of muscle torque
responsible for supporting the arm against gravity (postural component of muscle torque
or MT_P) and for intersegmental dynamics compensation (dynamic component of muscle
torque or MT_D) as described in detail in Olesh *et al.* [23]. The simulations were run for
each trial. Torque profiles were normalized in time, averaged per movement direction per
DOF and per subject, and down sampled to 100 samples (Fig 2B, example for one DOF).
Maximum torque amplitudes were determined per DOF per participant's arm and used to
scale the amplitudes of averaged profiles across movement directions and workspaces.
EMG recordings were high-pass filtered at 20 Hz to remove motion artifacts,
rectified, and low-pass filtered at 10 Hz [30]. Next, EMG recordings were normalized to
movement duration, down sampled to 100 samples, and averaged per movement
direction. Maximum contraction values were determined for each muscle and participant
across averaged EMG timeseries per movement direction and used to scale the
amplitudes of averaged profiles (Fig 2C).
Principal Component Analysis (PCA) was used to reduce the dimensionality of
EMG and torque data similar to Olesh *et al.* [23]. Normalized and demeaned EMG profiles
from 12 muscles of 28 center-out reaching movements (14 center-out and 14 return
movements between each pair of targets) were combined into one matrix, $A_{n \times m}$, where n

= 100 (time) and m = 336 (movement directions x muscle). PCA was applied to the data
from each of the four conditions (LLat, RLat, LMed, and RMed) separately. Thus, four
A_{nxm} matrices were obtained for each subject. Similarly, normalized and demeaned
muscle torque profiles (MT, MT_P, and MT_D) from 5 DOFs of 28 center-out reaching
movements were combined into one matrix, B_{nxm}, where n = 100 (time) and m = 140
(movement directions x DOF). Since the analysis included three types of torques (MT,
MT_P, MT_D) in the four conditions, twelve B_{nxm} matrices were obtained for each subject.
PCA was performed using *pca* function from MATLAB Statistics and Machine Learning
Toolbox. Each matrix was centered, and a default singular value decomposition algorithm
was applied to obtain principal component eigenvectors (Fig 3A, B), the variance
accounted for (VAF) by each principal component (Fig 3C), and scores that are the
representations of the matrix in the principal component space. It is important to note that
reaching in the same direction leftward or rightward from the starting location rely on
different muscles due to the bilateral symmetry of our body. That is why the reaches that
include leftward movements to the targets on the left from the central target performed by
the left hand were compared to the mirror rightward movements to the right from the
central target performed by the right hand and vice versa.
227
Statistics and Reproducibility
The first test of the generalizability of conclusions compared the profiles of the 1st
and 2nd principal components obtained from EMG to the principal components obtained
from muscle torques using a correlation analysis (*corr* function in MATLAB Signal
Processing Toolbox). We used repeated measures analysis of variance (RM ANOVA,
ranova function in MATLAB Statistics Toolbox) to test the null hypotheses that the
coefficients of determination (R^2) from all conditions and data types come from the same
distribution. The between-subject factor was Sex (assigned at birth), the within-subject
factors were Conditions (4 levels at LLat, RLat, LMed, and RMed) and Components (4
levels comparing E_1 vs. M_1, E_2 vs. M_2, E_1 vs. P_1, and E_2 vs. D_1). Post-hoc multiple
comparisons were done using *multcompare* function in MATLAB Statistics Toolbox. We
applied Bonferroni-Sidak correction to correct the alpha for familywise error, so that the
adjusted alpha for the comparison between 6 combinations of 4 conditions = 0.0085 and
for the comparison between 6 meaningful combinations of two principal components =
0.0085. The ethnicity – based statistical analysis was not performed due to small sample
size.
To test the hypothesis, we evaluated how muscles co-activate across changing
starting postures and movement directions. The rationale is that reaching in different
directions is caused by different amplitudes of postural or dynamic torques. Therefore,
given that the EMG variance is captured well by these torques (1st test above), the
corresponding features in EMG are also changing with the amplitudes of the torques. For
example, if the amplitude of shoulder flexion postural torque was larger during movement
upward compared to the movement forward, then the 1st principal component score of
the anterior deltoid (AD, shoulder flexor) EMG will also be larger during movement upward
compared to the movement forward. If the scores of multiple muscles change together
across movement directions, then these muscles are co-contracting together and can be
activated by a single control signal. The advantage of this method of calculating co-
contraction is that it does not depend on the assumptions underlying the scaling of EMG

amplitude. As long as the relative changes in the EMG amplitudes across movement
directions are preserved, the method works.
This method to calculate co-contraction was implemented using a linear correlation
analysis (*corr* and *corrplot* functions in MATLAB Statistics and Econometrics Toolboxes)
aimed at quantifying the linear relationships between principal component scores from
pairs of muscles across movements. First, we extracted the scores of the 1st principal
component from EMG and concatenated them into matrix C_{nxm}, where n = 28 (movement
directions) and m = 12 (muscles) per condition per subject. Positive correlations indicate
co-contraction of muscles, which negative correlations indicate reciprocal activation of
muscles, i.e., if the EMG activity related to postural or dynamic torques increases in one
muscle it decreases in another muscle. We evaluated the statistical significance of the
linear relationships represented by the Coefficients of Determination (R^2) between scores
of 66 muscle pairs (Fig 5). We applied Bonferroni-Sidak correction to correct the alpha
for familywise error, so that the adjusted alpha = 0.0008. We repeated this analysis for
the 2nd principal component from EMG that captures the dynamic torque feature in EMG.
271

Results

Participants performed center-out and return movements with each arm starting at
one of two locations of the central target. Four conditions were created based on the
location of the central target and the arm used to make reaches, i.e., LLat and RLat for
reaches by left and right arm respectively in their respective lateral workspaces and LMed
and RMed for the reaches by left and right arm respectively in the common medial
workspace. All participants were able to complete the center out task with either left or
right arms with consistent kinematics (Fig 2A). Due to bilateral symmetry, we expect that
movements with left and right arms with the same starting positions and kinematics in
forward, backward, up, or down directions rely on similar forces produced by the same
muscles. For the same reason movements leftward or rightward from the starting
positions by one limb will be similar to the rightward and leftward movements respectively
by the other limb. The forces underlying these movements were inferred from motion
capture using dynamic modeling that calculated muscle torques about each DOF of the
shoulder, elbow, and wrist joints. These torques were further subdivided into a linear
combination of postural and dynamic torques. The muscle torque profiles accompanying
the corresponding movements had very similar trajectories across conditions (Fig 2B).
However, there were static offsets in the postural torques underlying matching movement
directions. For example, holding the initial position at the beginning of the movement
required larger muscle torques about the right shoulder flexion/extension DOF (RMed and
RLat conditions) compared to that for the left shoulder in matching conditions (LMed and
LLat conditions respectively, Fig 2B). This is likely due to the motor redundancy or the
multiple combinations of joint angles and torques that can result in the same position of
the hand.
From our earlier work we knew that important features of muscle activity can be
captured by the postural and dynamic torques [23]. However, in that study we focused
our analysis on the movements performed by the dominant arm in the lateral workspace.
Here the analysis encompasses different workspaces and both arms, which enabled us
to evaluate the generalizability of this conclusion. We found that intra- and inter-subject
variability of EMG profiles were very low. The average standard deviation of EMG profiles

across movement repetitions for muscles in the dominant right arm (normalized to
maximal EMG of each muscle) was 12% ± 1% (standard deviation across subjects, SD).
The corresponding variability of EMG of muscles in the non-dominant left arm was 22%
± 2%. This shows that the muscle activation profiles were more consistent across
repetitions of the same movement type for the dominant arm compared to the non-
dominant arm. Moreover, the EMG profiles also broadly reflected the differences in the
joint torques across conditions (Fig 2C). In the example for the reach forwards and up in
Fig 2, the amplitude of anterior deltoid activity increased in parallel with the increases in
the postural component of the shoulder flexion muscle torque across conditions (Fig 2,
MT_P and ADel). Furthermore, the initial propulsive flexion moment in the dynamic muscle
torque profile was preceded by a burst in the activation of the long head of biceps (Fig 2,
MT_D and BiL, end of the burst is visualized). However, EMG profiles are notoriously noisy
and difficult to interpret. Therefore, in the following analysis we will broadly test the
generalizability of conclusions from Olesh et al. [23] by focusing on the salient features in
EMG obtained with principal component analysis (PCA; see details in Methods).
The temporal profile of the 1^{st} principal component (eigenvector E_1) obtained from
EMG matched the 1^{st} principal component (M_1) obtained from muscle torques (MT) and
the 1^{st} principal component (P_1) obtained from the postural torques only (MT_P; Fig 3A).
The 1^{st} principal component captures the gradually increasing or decreasing EMG or
torque profiles associated with the changes in posture from the starting position to the
final position at the reach target. The high variance captured by P_1 indicates that the
profiles of muscle torques were very similar across joints and DOFs, indicating a high
degree of intralimb coordination. As expected, the variance accounted for (VAF) by the
1^{st} principal component in EMG (E_1 = 53%) was lower than that for muscle torque (M_1 =
90%; Fig 3C) due to higher variability of EMG profiles compared to torque profiles.
The temporal profile of the 2^{nd} principal component (E_2) obtained from EMG
matched the 2^{nd} principal component (M_2) obtained from muscle torques and the 1^{st}
principal component (D_1) obtained from the dynamic torques only (MT_D) although with a
phase shift (Fig 3B). This feature captures the acceleration/deceleration-related changes
in torques that underly bursts in EMG profiles. The VAF by the 1^{st} principal components
obtained from muscle torques (M_1, P_1, and D_1) was very high (Fig 3C), the mean values
across conditions and all participants were 90% for M_1, 97% for P_1, and 88% for D_1 (Table
1 contains values per condition).
Interestingly, P_1 captured more variance across conditions and DOFs than M_1,
similarly D_1 captured more variance across conditions and DOFs than M_2 despite similar
profiles. This shows that subdividing muscle torques into postural and dynamic
components results in more reliable features compared to those obtained with PCA of
muscle torques. More VAF in EMG was captured by the 2^{nd} principal component (E_2 =
16%) than that in muscle torque (M_2 = 7%; Fig 3C). This indicates that highly stereotypical
muscle torques during reaching movements explain a high portion of the overall variance
in EMG profiles. The same conclusion was reached in Olesh et al. [23] from a smaller
subset of data. Moreover, our results further support the idea that E_1 captures the muscle
recruitment necessary for generating forces to counteract gravity, while E_2 captures the
muscle recruitment necessary for generating forces for accelerating and decelerating the
arm toward the target. This generalizes our earlier conclusions that the static and phasic

components of EMG envelope from a given muscle reflects the muscle's contribution to
gravity compensation and propulsion toward the target respectively.
The degree of similarity between features obtained with PCA across torques and
EMG was analyzed using cross correlation analysis as described in Methods. One-tailed
repeated measures analysis of variance (RM ANOVA) on the coefficients of determination
(R^2) between principal component profiles showed a significant main effect ($F(15, 105)$ =
9.36, p = 0.00) but accounting for sex reduced differences between conditions and
components ($F(15, 105)$ = 1.21, p = 0.27). Two-tailed post-hoc tests have shown no
differences between males and females and between conditions (Table 2). There were
significant differences between signal types from which components were derived
indicating that the temporal profiles of E_1, M_1, and P_1 were more similar to each other than
to E_2, M_2, or D_1 and vice versa (Table 3; Fig 4A). This further supports the idea that the
1^{st} and 2^{nd} principal components of EMG represent the contribution of the muscles to
postural and dynamic torques. These conclusions generalize across both limbs and
across medial and lateral workspaces. The profiles of postural and dynamic torques
captured most of variability in muscle activity profiles. Thus, this method of EMG
decomposition can be used to remove noise from EMG signals and increase the
interpretability of surface EMG recordings.
We have observed a consistent phase lead of the D_1 components onto the E_2
component (Fig 4B). This phase lead can potentially be taken advantage of to predict the
changes in the underlying muscle forces or contractions from dynamical simulations prior
to observable motion and improve performance of real-time control applications.
To test the hypothesis that the dimensionality of control space is shaped by
biomechanical constraints we quantified co-contraction using correlations between PCA
scores from EMG. The method relies on the established relationships between torque
and EMG components. These relationships imply that for movements requiring a larger
change in the postural torques, higher scores for the principal component representing
that feature (E_1) will be observed in one or more muscles in the same movement. If the
scores change together across movement directions in several muscles, these muscles
are co-contracting and can be activated by the same neural control signal. This means
that if the primary features in EMG (E_1 and E_2) represent low dimensional control signals,
we expect significant linear correlations between corresponding scores of multiple
muscles. We performed this linear correlation analysis per arm and workspace (Fig 5
shows an example for one condition and participant).
The scores represent the projection of EMG onto the principal component vectors.
When the scores are close to 0, this means that the components do not capture much
variance in the corresponding EMG profile. For all combinations of muscles, movements,
and participants, about 21% of the E_1 scores were low, i.e., values were below 5% of
maximal score (RLat: 24%, LLat: 19%, LMed: 17%, and RMed 22%). About 26% of E_2
scores were low (RLat: 27%, LLat: 26%, LMed: 22%, and RMed: 28%). The low score
means that the EMG profile for a given muscle in a given movement is not captured well
by PCA. Therefore, no relationships between the low score values are expected, as seen
for example in the insignificant correlation between low scores of BicL and ADel in the
example shown in Fig 5. However, we did observe multiple significant correlations
between E_1 scores in certain conditions. For reaching within the left lateral workspace,
61% of muscles (40 muscle pairs out of 66) in the non-dominant arm showed moderate

correlations between their E_1 scores ($R^2 \geq 0.5$; Fig 6A). For reaching within the medial
workspace, 41% of muscles (27/66 pairs) in the non-dominant arm showed moderate
correlations between their E_1 scores (Fig 6B). In contrast, only 11% of muscles in the
dominant arm showed moderate correlations when reaching within the right lateral
workspace (Fig 6C). For reaching within the shared medial workspace, 29% of muscles
in the dominant arm showed moderate correlations (Fig 6D), closer but lower than that
for the non-dominant arm (Fig 6B). Overall, more correlations were observed between E_1
scores in muscles of the non-dominant arm in 7 out of 9 individuals (Figs S1-S9 top row).
This shows that the forces needed to support the arm against gravity can be generated
by few control signals evidenced by the large number of co-contracting muscles,
particularly in the non-dominant arm. In contrast, there were much fewer muscle pairs
with consistently correlated E_2 scores (Fig 7). This was true for all individuals (Figs S1-
S9 bottom row). This suggests that the generation of the propulsive forces requires higher
dimensionality of control than that of the postural forces evidenced by the lack of
consistent co-contraction patterns across muscles. Overall, these results support the
hypothesis that the biomechanics shapes the dimensionality of control space.
Correlations between scores were overwhelmingly positive, indicating that not only
agonistic but also antagonistic muscles, such as biceps and triceps, changed their activity
together. Note that in these reaching movements with the arm pronated, the muscles
performing the antigravity action would be ADel, BiL, Brd, and ECR, while the rest would
produce joint moments in the direction of gravity (Fig. 1B). There were the most
correlations between E_1 scores in the non-dominant arm in the LLat condition. Two salient
muscle groups comprised proximal muscles (TerM, ADel, PDel, TriL, and TriS) and distal
muscles (BiL, Brd, FCR, FCU, and ECR), both highly coupled (Fig 6A). In contrast, the
correlations between E_1 scores in the dominant arm in the corresponding RLat condition
consisted of three much smaller muscle groups (PDel/TriS//BicL/Brd, Brd/ECR, and
FCR/FCU) with only the first two groups coupled through Brd (Fig 6C). The three small
clusters were the only consistent groups of coactivating muscles across both workspaces
and both limbs. This shows that both agonistic and antagonistic muscles are recruited to
produce forces that support the limb against gravity, indicating that limb stiffness can help
reduce the dimensionality of neural control space.
The co-contraction patterns for generating dynamic torques were much less
consistent across conditions and individuals. The correlations between E_2 scores in the
dominant arm in the RLat condition, only 4 pairs of muscles were consistently positively
correlated across individuals with <0.5 R^2 (R^2 for BicL/TriS = 0.46, Brd/TriS = 0.44,
Brd/BicL = 0.45, and ECR/Brd = 0.46). In individuals the patterns of co-contraction were
even less consistent (Figs S1-S9, bottom row). There were 6 significantly correlated
muscle pairs in RLat condition that were similar across at least 5/9 subjects (Brd/ECR,
FCR/FCU, Brd/BicL, Brd/TriS, BicL/BicS, and BicL/TriS) and 3 significantly correlated
muscle pairs in RMed condition (Brd/ECR, Brd/TriS, and BicL/TriS). For the non-dominant
arm, there were 2 significantly correlated muscle pairs in LLat condition that were similar
across 5/9 subjects (Brd/TriS and BicL/TriS) and 1 significantly correlated muscle pair in
LMed condition (BicL/TriS). The co-activation of most of these muscle groups, with the
exception of Brd/TriS and BicL/TriS, was agonistic. Moreover, some instances of negative
correlations between antagonistic muscle pairs were observed in individuals (ADel/TerM;
PDel/Pec, BicL/PDel, and ECR/FCU; Figs S1-S9 bottom row). This shows that more

complex control signals coordinating both co-contraction and reciprocal activation of muscles are needed to produce forces that propel the limb in different directions from different starting locations.

Discussion

We observed that highly stereotypical muscle torques during reaching movements explain a high portion of the overall variance in EMG profiles in multiple arm muscles. We have shown that the first principal component of EMG captures the muscle recruitment necessary for generating forces to counteract the force of gravity, while the second principal component of EMG captures the muscle recruitment necessary for generating dynamic forces for accelerating and decelerating the arm toward the target (Figs 3 and 4). These conclusions, originally derived from a smaller subset of movements by the dominant arm in one area of reaching workspace [23], are generalizable to movements by the non-dominant arm and across a larger reaching workspace that overlaps between both arm arms. This is useful for understanding the dimensionality of neural control space.

We have also observed correlations between scores of muscle pairs across movements indicating co-contraction but which and how many muscles co-contracted varied under deferent conditions (Figs 6, 7, and S1-S9). We have shown that the forces needed to support the arm against gravity can be generated by few control signals evidenced by the substantial number of co-contracting muscle pairs, particularly in the non-dominant arm. This is due to the force of gravity always pointing in the same direction, so that the counteracting forces can be produced by consistent groups of muscles. We have also shown that the generation of the propulsive forces require higher dimensionality of control than that of the postural forces evidenced by the lack of consistent co-contraction patterns across muscles. This is likely due to the direction-specific forces needed to make the movements to the different targets, which require different combinations of muscle forces. These results support the second hypothesis that the biomechanics shapes the dimensionality of control space.

It is possible that the reason we have not obtained consistent co-contraction patterns across movements is that the surface EMG was too noisy and variable across the different movements and individuals. However, our variability data reported in Results shows that the temporal profiles of EMG were very consistent across all these conditions, especially for the dominant arm with the fewest observed co-contractions. Another evidence of the consistency of EMG profiles is the large percentage of variance that is captured in EMG by the first two principal components (Fig 3) and by the components obtained from muscle torques (Fig 4), which are highly stereotypical during reaching movements. Therefore, it seems highly unlikely that the inter-movement or inter-subject variability can explain the lack of consistent co-contraction across different conditions.

The neural control strategies underlying muscle recruitment during different phases of movement can be quantitatively described through not only muscle torques but also mechanical impedance. Muscles are organized in a complex pattern to generate opposing moments around each joint DoF as determined by intrinsic muscle properties [31] and muscle moment arms. Therefore, neural muscle recruitment causes not only muscle torques but also the counterbalanced forces that define the stiffness and viscosity components of impedance, which in turn determine the reaction of the musculoskeletal system to perturbations [32]. The impedance parameters that are under the control of the

nervous system can be measured by assessing changes in the co-contraction of
antagonistic muscles [26]. Both stiffness and viscosity components of mechanical
impedance define how far the joints move in response to an applied external force [32].
Muscle recruitment defines the functional aspects of joint impedance that can and do
change with task demands [33,34]. Here too we observed that the component of EMG
that is responsible for supporting the limb against gravity is present in not only antigravity
muscles but also in their antagonists and that the amplitude of this component changes
together across movement directions indicating co-contraction (Fig 6). Other studies have
shown that the amount of muscle co-activation increases in movements with unstable
loads compared to those with stable loads [35] and the impedance of the whole limb
increases to stabilize movement when learning novel dynamical tasks [33]. The joint
stiffness component of impedance has been shown to be higher during unconstrained
movements compared to constrained ones indicating, unsurprisingly, that unconstrained
movements are inherently less stable [36]. Limb impedance increases when learning to
control an unstable object and when movement stability is challenged by a robot [37,38].
Wrist stiffness increases during unstable dynamical loads compared to stable ones [39]
and when the accuracy demands of the task are high [40]. Moreover, we observed more
co-contraction in the non-dominant limb compared that in the dominant limb, supporting
earlier observations that that there is hemispheric specialization in the compensation for
limb dynamics. The non-dominant limb motor control has been shown to be ore
specialized for impedance control [41–43] Overall, this suggests that the compensation
for gravity load on the arm during reaching is accomplished through a low-dimensional
control space by controlling limb stiffness. Low-dimensional control of posture through
stereotypical postural adjustments and other reflexes has been shown to involve the
reticulospinal tract [18–20] (Fig. 1, green).
Our results also support the idea of internal models embedding body dynamics.
Internal models are defined as neural networks in the brain that simulate the dynamics of
the body and its interaction with the external world [1,3,44–47]. Forward models predict
the consequences of a given action, for example, predicting sensory feedback without
delay from an efferent copy of a corticospinal motor command [48,49]. Inverse models
capture information in the opposite direction [3,50]. They convert a motor plan into the
motor execution output of the motor system that converges on the motoneurons in the
spinal cord. The motor execution output is thought to be forces in the intrinsic reference
frame that are closely related to individual muscle forces or moments [51,52]. The motor
plan is thought to be developed in the premotor areas, where neural activity has many
kinematic features in the extrinsic head-centered reference frame resolving the kinematic
redundancy problem [53–55]. The motor execution output is thought to originate in the
primary motor cortex, where neural activity has many dynamic features in the intrinsic
reference frame [51,56–58]. The transformations from motor plan into motor execution
may potentially resolve the muscle redundancy problem by embedding biomechanical
constraints. These constraints may take the form of embedded muscle anatomy
relationships, such as known ratios between moments arms of muscles about a given
DOF [59]. Here too we have observed that a large amount of variance in the muscle
activity profiles are captured by muscle torques. Interestingly, the component of muscle
activity that reflects propulsion was not associated with consistent patterns of antagonistic
co-contraction across multiple movement directions (Fig 7). Instead, smaller subsets of

mainly agonistic co-contraction and reciprocal activation of antagonists was observed(Figs 7, S1-S9). This provides evidence that control solutions with higher dimensionality are needed for generating these forces. Internal models could support high-dimensional control, because the control solutions are thought to be calculated on the fly by the neural networks that embed the equations of motion and biomechanical constraints. These calculations likely involve the higher-order neural structures whose output converges on the corticospinal tract [21,60–62] (Fig. 1, purple).

Conclusions

Results show that across multiple movement directions and starting positions and in both arms muscle activity broadly reflects the temporal evolution of muscle torques. We suggest that the muscles generate forces to counteract gravity through a simplified set of control signals, leading to coordinated contraction among various muscles, which adjusts the stiffness of the limb. Conversely, the forces that drive the limb towards a target demand more intricate control signals that vary with the direction of movement.

Acknowledgements

We would like to acknowledge the contributions of Dr. E.V. Olesh and Dr. A.B. Thomas to the collection and preliminary analysis of the reported data. V.G. was supported by NIGMS grants P20GM109098 and P30GM103503. ASK was supported by a fellowship from NIGMS T32 AG052375. This work was supported in part by the Office of the Assistant Secretary of Defense for Health Affairs through the Restoring Warfighters with Neuromusculoskeletal Injuries Research Program (RESTORE) under Award No. W81XWH-21-1-0138. Opinions, interpretations, conclusions, and recommendations are those of the author and are not necessarily endorsed by the Department of Defense.

Author Contributions

A.S.K. contributed to conceptualization, data curation, data analysis, and writing. V.G. contributed to conceptualization, methodology development, data curation, data analysis, and writing. All co-authors have read and edited the manuscript and agree with its content.

Competing Interests

The authors declare no competing interests.

Data Availability Statement

The data used for statistical analysis, such as principal components and scores for electromyography and muscle torques, are shared online [63].

Tables

Table 1. Variance accounted for across reaching conditions.

	Left lateral	Left medial	Right medial	Right lateral	Total
E_1	53.72 ± 9.33	49.74 ± 10.78	56.65 ± 5.18	51.96 ± 6.09	53.02 ± 7.85
M_1	89.19 ± 4.50	87.66 ± 6.57	92.03 ± 3.85	91.04 ± 5.65	89.98 ± 5.14
P_1	98.16 ± 0.75	96.46 ± 3.93	96.51 ± 6.23	98.32 ± 1.00	97.36 ± 2.98

E_2	14.42 ± 4.55	13.73 ± 4.21	16.32 ± 5.00	17.62 ± 6.55	15.52 ± 5.08
M_2	7.10 ± 2.96	8.89 ± 5.00	5.66 ± 3.18	6.86 ± 4.94	7.13 ± 4.02
D_1	78.52 ± 4.04	90.19 ± 4.33	92.94 ± 3.02	91.36 ± 7.42	88.25 ± 4.70

Values are means ± standard deviation across 9 individuals.

Table 2. Post-hoc differences between the coefficients of determination across males and females and across limbs and workspaces across 9 participants.

Sex		MSE ± SE	p value
F	M	-0.10 ± 0.04	0.06
Conditions		MSE ± SE	p value
LLat	LMed	-0.02 ± 0.03	0.95
LLat	RLat	0.01 ± 0.03	0.96
LLat	RMed	0.01 ± 0.04	1.00
LMed	RLat	0.03 ± 0.04	0.88
LMed	RMed	0.02 ± 0.04	0.91
RLat	RMed	-0.01 ± 0.02	0.98

MSE – mean squared error, SE – standard error of the mean. Adjusted alpha = 0.0085.

Table 3. Post-hoc differences between the coefficients of determination based on EMG- and torque-based components across 9 participants.

Components		MSE ± SE	p value
E_1/M_1	E_1/P_1	-0.01 ± 0.01	0.9036
E_1/M_1	E_2/M_2	0.19 ± 0.04	**0.0048**
E_1/M_1	E_2/D_1	0.29 ± 0.05	**0.0040**
E_1/P_1	E_2/M_2	0.20 ± 0.05	0.0155
E_1/P_1	E_2/D_1	0.30 ± 0.06	**0.0066**
E_2/M_2	E_2/D_1	-0.10 ± 0.05	0.2367

MSE – mean squared error, SE – standard error of the mean. Bold p values show significant difference with adjusted alpha = 0.0085.

Figure Legends

Figure 1. Schematic of neuromuscular system. A. Basic components of the neural circuit involved in muscle activation. Drawing is not to scale. **B.** Innervation of muscles recorded during the experiment and their arrangement on the arm. Abbreviations are as follows: the clavicular head of pectoralis (Pec), teres major (TerM), anterior deltoid (ADel), posterior deltoid (PDel), the long and lateral heads of triceps (TriL and TriS), the short and long heads of biceps (BiS and BiL), brachioradialis (Brd), flexor capri radialis (FCR), flexor carpi ulnaris (FCU), and extensor capri radialis (ECR). Blue dots indicate locations of LED markers for motion capture.

Figure 2. **Examples of kinematics, dynamics, and muscle activity profiles during reaching.** Each plot shows profiles per condition averaged across repetitions (n = 15) of the same reaching movement in one direction by one participant. (**A**) Profiles (solid lines) and standard deviation (shaded areas) of shoulder flexion-extension angle and angular velocity. (**B**) Shoulder flexion-extension torque (t) profiles that caused the movements in

(**A**). (**C**) Normalized electromyography (EMG) profiles of anterior deltoid (ADel), triceps long (TriL), and biceps long (BiL) that accompanied the movements in (**A**). Shaded areas show the standard error for the mean across repetitions (n = 15) of the same movement.

Figure 3. Principal components of EMG and muscle torques. Data from the four conditions indicated by the pictograms are shown in plots arranged in columns. The data is from 9 participants who performed all 4 conditions. (**A**) Solid lines show profiles of the 1^{st} principal component obtained from EMG (E_1), muscle torque (M_1), and the postural component of muscle torque (P_1). (**B**) Solid lines show profiles of the 2^{nd} principal component obtained from EMG (E_2) and muscle torque (M_2), and the 1^{st} principal component obtained from the dynamic component of muscle torque (D_1). (C) Variance accounted for (VAF) by principal components obtained from EMG and muscle torques.

Figure 4. Correlations between the temporal profiles of principal components derived from EMG and muscle torques. Data (N = 9) from the four conditions indicated by the pictograms are shown in plots arranged in columns. (A) Dots show individual R^2 values from cross-correlations between profiles, red lines show median values, and grey boxes show interquartile ranges. Brackets with stars indicate significant relationships with correction for familywise error. (B) Dots show individual lag values from cross correlations, red lines show median values, and grey boxes show interquartile ranges. Open dots are outliers. E_1 and E_2 are the 1^{st} and 2^{nd} principal components respectively obtained from EMG; M_1 and M_2 are the 1^{st} and 2^{nd} principal components respectively obtained from muscle torques; P_1 and D_1 are the 1^{st} principal components obtained from the postural and dynamic components of muscle torque respectively.

Figure 5. Example linear regressions between scores across reaching directions. Example regression matrix between E_1 scores of 12 muscles across 28 right-handed reaches within the medial workspace (RMed condition) for a single participant. The coordinates of each dot represent principal component scores for two muscles for a single movement direction. Histograms along the diagonal show the distribution of the scores for a given muscle across reaching directions. Solid lines show least-squares linear regression, red lines indicate significant relationships with correction for familywise error, so that the adjusted alpha = 0.0008.

Figure 6. Gravity-related muscle co-contraction across all conditions. Heatmaps show coefficients of determination (R^2) from regressions shown in Fig 4 averaged across 9 participants. Red and orange colors represent strong and moderate relationships respectively between E_1 coefficients across movement directions. Pictograms indicate conditions, i.e., LLat (A), LMed (B), RLat (C), and RMed (D). Muscles are abbreviated as described in Methods.

Figure 7. Dynamic torque-related muscle co-contraction across all conditions. Heatmaps show coefficients of determination (R^2) from regressions shown in Fig 4 averaged across 9 participants. Red, orange and darker blue colors represent moderate and strong relationships between E_2 coefficients across movement directions. Pictograms

642   indicate conditions, i.e., LLat (A), LMed (B), RLat (C), and RMed (D). Muscles are
643   abbreviated as described in Methods.
644
Supporting information
646
**Supplementary Figures S1-S9. Gravity and dynamic torque-related muscle co-
contraction across all conditions for each participant.** Heatmaps show coefficients of
determination (R^2). Red, orange and darker blue colors represent moderate and strong
relationships between E_1 (A, B) and E_2 (C, D) coefficients across movement directions;.
Pictograms indicate conditions, i.e., LLat (A, C) and RLat (B, D). Muscles are abbreviated
as follows: the clavicular head of pectoralis (Pec), teres major (TerM), anterior deltoid
(ADel), posterior deltoid (PDel), the long and lateral heads of triceps (TriL and TriS), the
short and long heads of biceps (BiS and BiL), brachioradialis (Brd), flexor capri radialis
(FCR), flexor carpi ulnaris (FCU), and extensor capri radialis (ECR).
656

References

1. Kawato M, Furukawa K, Suzuki R. A hierarchical neural-network model for control and
learning of voluntary movement. Biol Cybern. 1987;57: 169–185.
doi:10.1007/BF00364149

2. Wolpert DM, Ghahramani Z. Computational principles of movement neuroscience.
Nat Neurosci. 2000;3: 1212–1217. doi:10.1038/81497

3. Wolpert DM, Kawato M. Multiple paired forward and inverse models for motor control.
Neural Networks. 1998;11: 1317–1329. doi:10.1016/S0893-6080(98)00066-5

4. d'Avella A, Lacquaniti F. Control of reaching movements by muscle synergy
combinations. Front Comput Neurosci. 2013;7. doi:10.3389/fncom.2013.00042

5. Mussa-Ivaldi FA. Modular features of motor control and learning. Current Opinion in
Neurobiology. 1999;9: 713–717. doi:10.1016/S0959-4388(99)00029-X

6. Thoroughman KA, Shadmehr R. Learning of action through adaptive combination of
motor primitives. Nature. 2000;407: 742–747. doi:10.1038/35037588

7. Bizzi E, Tresch MC, Saltiel P, d'Avella A. New perspectives on spinal motor systems.
Nat Rev Neurosci. 2000;1: 101–108. doi:10.1038/35039000

8. d'Avella A, Saltiel P, Bizzi E. Combinations of muscle synergies in the construction of
a natural motor behavior. Nat Neurosci. 2003;6: 300–308. doi:10.1038/nn1010

9. Gritsenko V, Hardesty RL, Boots MT, Yakovenko S. Biomechanical constraints
underlying motor primitives derived from the musculoskeletal anatomy of the human
arm. PLoS ONE. 2016;11: 1–18. doi:10.1371/journal.pone.0164050

10. Taitano RI, Yakovenko S, Gritsenko V. Muscle anatomy is reflected in the spatial
organization of the spinal motoneuron pools. Commun Biol. 2024;7: 97.
doi:10.1038/s42003-023-05742-w

11. Tresch MC, Jarc A. The case for and against muscle synergies. Current Opinion in
Neurobiology. 2009;19: 601–607. doi:10.1016/j.conb.2009.09.002

12. Russo M, D'Andola M, Portone A, Lacquaniti F, d'Avella A. Dimensionality of joint
torques and muscle patterns for reaching. Front Comput Neurosci. 2014;8.
doi:10.3389/fncom.2014.00024

13. Tresch MC, Cheung VCK, d'Avella A. Matrix Factorization Algorithms for the
Identification of Muscle Synergies: Evaluation on Simulated and Experimental Data
Sets. Journal of Neurophysiology. 2006;95: 2199–2212. doi:10.1152/jn.00222.2005

14. Purves D, editor. Neuroscience. 3rd ed. Sunderland, Mass: Sinauer Associates,
Publishers; 2004.

15. Kandel ER, Koester J, Mack S. Principles of Neural Science. Sixth edition.
Siegelbaum S, editor. New York: McGraw Hill; 2021.

16. Brambilla C, Atzori M, Müller H, d'Avella A, Scano A. Spatial and Temporal Muscle
Synergies Provide a Dual Characterization of Low-dimensional and Intermittent
Control of Upper-limb Movements. Neuroscience. 2023;514: 100–122.
doi:10.1016/j.neuroscience.2023.01.017

17. Hanna K, Yakovenko S. The same muscle synergies are used to control symmetric
and asymmetric locomotion. 2021 10th International IEEE/EMBS Conference on
Neural Engineering (NER). Italy: IEEE; 2021. pp. 840–843.
doi:10.1109/NER49283.2021.9441083

18. Brownstone RM, Chopek JW. Reticulospinal Systems for Tuning Motor Commands.
Front Neural Circuits. 2018;12: 30. doi:10.3389/fncir.2018.00030

19. Hardesty RL, Ellaway PH, Gritsenko V. The Human Motor Cortex Contributes to
Gravity Compensation to Maintain Posture and During Reaching. Journal of
Neurophysiology. 2023;129: 83–101. doi:10.1152/jn.00367.2021

20. Glover IS, Baker SN. Cortical, Corticospinal, and Reticulospinal Contributions to
Strength Training. J Neurosci. 2020;40: 5820–5832.
doi:10.1523/JNEUROSCI.1923-19.2020

21. Gritsenko V, Kalaska JF, Cisek P. Descending Corticospinal Control of
Intersegmental Dynamics. Journal of Neuroscience. 2011;31: 11968–11979.
doi:10.1523/JNEUROSCI.0132-11.2011

22. Luca CJD. The Use of Surface Electromyography in Biomechanics. Journal of
Applied Biomechanics. 1997;13: 135–163. doi:10.1123/jab.13.2.135

23.   Olesh EV, Pollard BS, Gritsenko V. Gravitational and dynamic components of
715   muscle torque underlie tonic and phasic muscle activity during goal-directed
716   reaching. Front Hum Neurosci. 2017;11: 474. doi:10.3389/fnhum.2017.00474

717   24.   Burke RE, Rymer WZ. Relative strength of synaptic input from short-latency
718   pathways to motor units of defined type in cat medial gastrocnemius. Journal of
719   Neurophysiology. 1976;39: 447–458. doi:10.1152/jn.1976.39.3.447

720   25.   Hardesty RL, Boots MT, Yakovenko S, Gritsenko V. Computational evidence for
721   nonlinear feedforward modulation of fusimotor drive to antagonistic co-contracting
722   muscles. Sci Rep. 2020;10: 10625. doi:10.1038/s41598-020-67403-w

723   26.   Hogan N. Adaptive control of mechanical impedance by coactivation of antagonist
724   muscles. IEEE Transactions on Automatic Control. 1984;29: 681–690.
725   doi:10.1109/TAC.1984.1103644

726   27.   Hogan N. The mechanics of multi-joint posture and movement control | SpringerLink.
727   Biological Cybernetics. 1985;52: 315–335.

728   28.   Talkington WJ, Pollard BS, Olesh EV, Gritsenko V. Multifunctional setup for studying
729   human motor control using transcranial magnetic stimulation, electromyography,
730   motion capture, and virtual reality. JoVE. 2015; 52906. doi:10.3791/52906

731   29.   Winter DA. Biomechanics and Motor Control of Human Movement. 4th ed. Wiley;
732   2009. Available: https://www.wiley.com/en-
733   us/Biomechanics+and+Motor+Control+of+Human+Movement%2C+4th+Edition-p-
734   9780470398180

735   30.   Hermens HJ, Merletti R, Freriks B. SENIAM: European recommendations for
736   surface electromyography. 2d edition. Enschede, The Netherlands: Roessingh
737   Research and Development; 1999.

738   31.   Zajac FE. Muscle and tendon: properties, models, scaling, and application to
739   biomechanics and motor control. Critical Reviews in Biomedical Engineering.
740   1989;17: 359–411.

741   32.   Mizrahi J. Mechanical Impedance and Its Relations to Motor Control, Limb
742   Dynamics, and Motion Biomechanics. J Med Biol Eng. 2015;35: 1–20.
743   doi:10.1007/s40846-015-0016-9

744   33.   Milner TE, Franklin DW. Impedance control and internal model use during the initial
745   stage of adaptation to novel dynamics in humans: Initial stage of motor adaptation.
746   The Journal of Physiology. 2005;567: 651–664. doi:10.1113/jphysiol.2005.090449

747   34.   Franklin DW, Liaw G, Milner TE, Osu R, Burdet E, Kawato M. Endpoint Stiffness of
748   the Arm Is Directionally Tuned to Instability in the Environment. Journal of
749   Neuroscience. 2007;27: 7705–7716. doi:10.1523/JNEUROSCI.0968-07.2007

35. Milner TE. Adaptation to destabilizing dynamics by means of muscle cocontraction.
Exp Brain Res. 2002;143: 406–416. doi:10.1007/s00221-002-1001-4

36. Damm L, McIntyre J. Physiological Basis of Limb-Impedance Modulation During
Free and Constrained Movements. Journal of Neurophysiology. 2008;100: 2577–
2588. doi:10.1152/jn.90471.2008

37. Rancourt D, Hogan N. Stability in force-production tasks. Journal of Motor Behavior.
2001. doi:10.1080/00222890109603150

38. Burdet E, Osu R, Franklin DW, Milner TE, Kawato M. The central nervous system
stabilizes unstable dynamics by learning optimal impedance. Nature. 2001;414:
446–449.

39. De Serres SJ, Milner TE. Wrist muscle activation patterns and stiffness associated
with stable and unstable mechanical loads. Exp Brain Res. 1991;86: 451–458.
doi:10.1007/BF00228972

40. Wong J, Wilson ET, Malfait N, Gribble PL. Limb Stiffness Is Modulated With Spatial
Accuracy Requirements During Movement in the Absence of Destabilizing Forces.
Journal of Neurophysiology. 2009;101: 1542–1549. doi:10.1152/jn.91188.2008

41. Schaffer JE, Sainburg RL. Interlimb differences in coordination of unsupported
reaching   movements.   Neuroscience.   2017;350:   54–64.
doi:10.1016/j.neuroscience.2017.03.025

42. Sainburg RL. Evidence for a dynamic-dominance hypothesis of handedness. Exp
Brain Res. 2002;142: 241–258. doi:10.1007/s00221-001-0913-8

43. Bagesteiro LB, Sainburg RL. Handedness: Dominant Arm Advantages in Control of
Limb   Dynamics.   Journal   of   Neurophysiology.   2002;88:   2408–2421.
doi:10.1152/jn.00901.2001

44. Cisek P. Internal Models. In: Binder MD, Hirokawa N, Windhorst U, editors.
Encyclopedia of Neuroscience. Berlin, Heidelberg: Springer; 2009. pp. 2009–2012.
doi:10.1007/978-3-540-29678-2_2528

45. Wolpert D, Ghahramani Z, Jordan M. An internal model for sensorimotor integration.
Science. 1995;269: 1880–1882. doi:10.1126/science.7569931

46. Hwang EJ, Shadmehr R. Internal models of limb dynamics and the encoding of limb
state. J Neural Eng. 2005;2: S266–S278. doi:10.1088/1741-2560/2/3/S09

47. Zago M, Lacquaniti F. Visual perception and interception of falling objects: a review
of evidence for an internal model of gravity. J Neural Eng. 2005;2: S198–S208.
doi:10.1088/1741-2560/2/3/S04

48. Haggard P, Whitford B. Supplementary motor area provides an efferent signal for
sensory   suppression.   Cognitive   Brain   Research.   2004;19:   52–58.
doi:10.1016/j.cogbrainres.2003.10.018

49. Chapman CE. Active versus passive touch: factors influencing the transmission of
somatosensory signals to primary somatosensory cortex. Can J Physiol Pharmacol.
1994;72: 558–570. doi:10.1139/y94-080

50. Schweighofer N, Arbib MA, Kawato M. Role of the cerebellum in reaching
movements in humans. I. Distributed inverse dynamics control. European Journal of
Neuroscience. 1998;10: 86–94. doi:10.1046/j.1460-9568.1998.00006.x

51. Scott SH, Kalaska JF. Reaching Movements With Similar Hand Paths But Different
Arm Orientations. I. Activity of Individual Cells in Motor Cortex. Journal of
Neurophysiology. 1997;77: 826–852. doi:10.1152/jn.1997.77.2.826

52. Scott SH, Gribble PL, Graham KM, Cabel DW. Dissociation between hand motion
and population vectors from neural activity in motor cortex. Nature. 2001;413: 161–
165. doi:10.1038/35093102

53. Crammond DJ, Kalaska JF. Prior Information in Motor and Premotor Cortex: Activity
During the Delay Period and Effect on Pre-Movement Activity. Journal of
Neurophysiology. 2000;84: 986–1005. doi:10.1152/jn.2000.84.2.986

54. Cisek P, Kalaska JF. Neural correlates of mental rehearsal in dorsal premotor cortex.
Nature. 2004;431: 993–996. doi:10.1038/nature03005

55. Scott SH, Sergio LE, Kalaska JF. Reaching Movements With Similar Hand Paths but
Different Arm Orientations. II. Activity of Individual Cells in Dorsal Premotor Cortex
and Parietal Area 5. Journal of Neurophysiology. 1997;78: 2413–2426.
doi:10.1152/jn.1997.78.5.2413

56. Ashe J. Force and the motor cortex. Behav Brain Res. 1997;87: 255–269.
doi:10.1016/s0166-4328(97)00752-3

57. Kalaska JF. From Intention to Action: Motor Cortex and the Control of Reaching
Movements. In: Sternad D, editor. Progress in Motor Control: A Multidisciplinary
Perspective. Boston, MA: Springer US; 2009. pp. 139–178. doi:10.1007/978-0-387-
77064-2_8

58. Evarts EV. Relation of pyramidal tract activity to force exerted during voluntary
movement.   Journal   of   Neurophysiology.   1968;31:   14–27.
doi:10.1152/jn.1968.31.1.14

59. Gritsenko V, Hardesty RL, Boots MT, Yakovenko S. Biomechanical Constraints
Underlying Motor Primitives Derived from the Musculoskeletal Anatomy of the
Human   Arm.   Jones   KE,   editor.   PLoS   ONE.   2016;11:   e0164050.
doi:10.1371/journal.pone.0164050

821 60. Kawato M. Internal models for motor control and trajectory planning. Current Opinion
822   in Neurobiology. 1999;9: 718–727. doi:10.1016/S0959-4388(99)00028-8

823 61. Glover IS, Baker SN. Both Corticospinal and Reticulospinal Tracts Control Force of
824   Contraction. J Neurosci. 2022;42: 3150–3164. doi:10.1523/JNEUROSCI.0627-
825   21.2022

826 62. Lemon R, Johansson R, Westling G. Corticospinal control during reach, grasp, and
827   precision lift in man. J Neurosci. 1995;15: 6145–6156. doi:10.1523/JNEUROSCI.15-
828   09-06145.1995

829 63. Korol A, Gritsenko V. Dataset for How muscle synergies fail to solve the muscle
830   redundancy problem during human reaching. [object Object]; 2024. p. 1577914
831   Bytes. doi:10.6084/M9.FIGSHARE.25360231.V2

832