Natural Variation in Age-Related Dopamine Neuron Degeneration is Glutathione-Dependent and Linked to Life Span

By

Colin R. Coleman[1], Judit Pallos[1], Alicia Arreola-Bustos[1], Lu Wang[2], Daniel Raftery[3], Daniel E.L. Promislow[4,5] and Ian Martin[1*]

[1] Jungers Center for Neurosciences, Department of Neurology, Oregon Health and Science University, Portland, Oregon, USA

[2] Department of Environmental and Occupational Health Sciences, University of Washington, Seattle, Washington, USA

[3] Northwest Metabolomics Research Center, Department of Anesthesiology and Pain Medicine, University of Washington, Seattle, Washington, USA

[4] Department of Laboratory Medicine and Pathology, University of Washington School of Medicine, Seattle, Washington, USA

[5] Department of Biology, University of Washington School of Medicine, Seattle, Washington, USA

* Corresponding author
Ian Martin, PhD
Jungers Center for Neurosciences Research
Department of Neurology
Oregon Health and Science University
3181 SW Sam Jackson Park Road
Portland, OR 97239
Phone: 503-494-9140
Fax: 503-494-7358
Email: martiia@ohsu.edu

SUMMARY

Aging is the biggest risk factor for Parkinson's disease (PD), suggesting that age-related changes in the brain promote dopamine neuron vulnerability. It is unclear, however, whether aging alone is sufficient to cause significant dopamine neuron loss and if so, how this intersects with PD-related neurodegeneration. Here, through examining a large collection of naturally varying *Drosophila* strains, we find a strong relationship between life span and age-related dopamine neuron loss. Naturally short-lived strains exhibit a loss of dopamine neurons but not generalized neurodegeneration, while long-lived strains retain dopamine neurons across age. Metabolomic profiling reveals lower glutathione levels in short-lived strains which is associated with elevated levels of reactive oxygen species (ROS), sensitivity to oxidative stress and vulnerability to silencing the familial PD gene *parkin*. Strikingly, boosting neuronal glutathione levels via glutamate-cysteine ligase (GCL) overexpression is sufficient to normalize ROS levels, extend life span and block dopamine neurons loss in short-lived backgrounds, demonstrating that glutathione deficiencies are central to neurodegenerative phenotypes associated with short longevity. These findings may be relevant to human PD pathogenesis, where glutathione depletion is frequently reported in idiopathic PD patient brain. Building on this evidence, we detect reduced levels of GCL catalytic and modulatory subunits in brain from PD patients harboring the LRRK2 G2019S mutation, implicating possible glutathione deficits in familial LRRK2-linked PD. Our study across *Drosophila* and human PD systems suggests that glutathione plays an important role in the influence of aging on PD neurodegeneration.

INTRODUCTION

The movement disorder in Parkinson's disease (PD) results from a progressive degeneration and loss of dopamine neurons in the substantia nigra. The single biggest risk factor for developing PD is age, suggesting that aging promotes a loss of dopamine neuron health and viability (1, 2). Aging leads to a decline in molecular, cellular and physiological function that impacts brain health and homeostasis (3). Cell stressors such as oxidative stress, metabolic dysfunction and genomic instability all increase with age in many species including humans (4), and dopamine neurons are considered particularly vulnerable to oxidative stress because of their high intrinsic production of reactive oxygen species (ROS) through dopamine metabolism and autonomous pacemaking activity (5). Although aging is the biggest risk factor for PD, the extent to which dopamine neuron senescence contributes to neurodegeneration and PD onset is still unknown. For example, it is unclear whether aging alone is sufficient to cause subclinical dopaminergic neurodegeneration in the absence of PD-associated genetic and/or environmental factors. Further, many PD-linked genetic mutations or neurotoxins cause PD in only a subset of people (6-8), suggesting that there may be influential aging factors governing whether disease manifests, yet the identity of these are unknown. Humans of advanced age without PD are usually reported to experience minor or no loss of substantia nigra dopamine neurons (9-11) with some exceptions (12), despite a robust age-related increase in nigral alpha-synuclein protein (13). Few studies in animal models have focused on the impact of aging alone on dopamine neuron health and viability, although these data can frequently be gleaned from control groups comprising wild-type animals examined at various ages in parallel with animals

harboring, for example, mutations in familial PD genes. These control group data typically suggest minor or no loss of dopamine neurons upon aging, whether in the mouse SN (14-16) or fly central brain (17-20), and this also appears to be the case in non-human primates (21-23). When C57BL/6 mice were aged over a longer period to 120 weeks, they exhibited reduced striatal DA levels, locomotor impairment and altered SN mitochondrial morphology compared to 30-week-old mice, but no overt loss of SN dopamine neurons (24). Hence, limited data from humans and animal models is largely consistent with only minor effects of aging on dopamine neuron viability.

One important limitation of prior animal studies is that they have mostly been conducted using animals of limited genetic diversity, for example rodents belonging to an inbred standard laboratory background strain such as C57BL/6 mice. Genetic and environmental factors are strong drivers of aging heterogeneity and hence individuals of the same chronological age in an outbred population may vary substantially in the extent to which they manifest individual features of biological aging (4). As a result, certain aging-related traits may be masked in animal studies restricted to a single inbred genetic background. Conversely, studies of genetically diverse populations may enable clearer observation of these traits and thus provide an important foundation for gaining mechanistic insight into their origin. We asked whether natural genetic variation can be harnessed to better understand dopamine neuron vulnerability to aging. We used the *Drosophila* Genetic Reference Panel (DGRP), a large set of genetically and phenotypically diverse fly strains that have each been inbred to homozygosity and fully sequenced, harboring millions of polymorphisms that can affect complex traits (25). Multiple studies have demonstrated extensive variation in life span among the DGRP

strains (26-28), consistent with aging heterogeneity in this collection of fly strains that can be leveraged to study age-dependent traits. Indeed, DGRP strains have been shown to vary widely in several parameters associated with aging, including visual senescence (29), decline in fecundity (27, 30) and climbing ability (30, 31) as well as susceptibility to neurodegeneration triggered by expression of pathogenic proteins such as $A\beta_{42}$ and tau (32) or mutant LRRK2 (33).

Here, we took two cohorts of flies representing the top 15 and bottom 15 DGRP inbred strains ranked by longevity to assess whether the influence of natural genetic variation on survival also affects dopamine neuron health in aged animals. We observe a strong inverse correlation between strain life span and age-related loss of tyrosine hydroxylase (TH)-expressing neurons. This loss is associated with elevated ROS levels as well as vulnerability to silenced expression of *parkin*, a gene whose loss-of-function mutations are linked to familial PD. Unbiased metabolic profiling uncovers glutathione deficits in short-lived strains and boosting glutathione levels is sufficient to normalize ROS levels, block the age-related loss of TH-expressing neurons and extend life span in these strains, suggesting a link between brain glutathione levels, dopamine neuron health and organismal longevity. We further extend our findings to human PD, observing reduced expression of the key glutathione biosynthetic enzyme glutamate-cysteine ligase (GCL) in human PD brain tissue from patients harboring the LRRK2 G2019S mutation.

RESULTS

Age-related dopamine neuron health and viability is associated with life span

The approximately 200 inbred fly strains in the DGRP collection exhibit major variability in life span (26-28). This suggests that the natural genetic variance existing between these strains is a strong driver of aging heterogeneity. We took advantage of this natural variation to determine whether dopamine neuron health and viability are also divergent in this collection of strains. We chose a subset of DGRP strains consisting of the shortest-lived 15 lines and the longest- lived 15 lines previously reported (26) to assess the number of TH-expressing dopamine neurons in aged flies. Since life span data were only available for female life span (26, 27), we focused our analysis on female flies, choosing lines based on life span data reported in Ivanov et al. (26). We first confirmed a large variation in mean female life span across the 27 strains we could grow, ranging from 21 days (RAL-757) to 86 days (RAL-821) (Fig. 1A and 1B and Table S1). Female mean life spans of the strains used in our study correlate moderately well with their previously reported life spans (r = 0.57 p = 0.001, Fig. S1A), supporting the heritability of this trait. Flies from the same DGRP strains grown and aged in parallel exhibit a strong correlation between mean life span and number of TH-expressing dopamine neurons in the protocerebral posterior lateral 1 (PPL1) cluster (Fig. 1C and D). These 30-day-old flies were assessed at the age where all but two strains retained at least 50% survival (Table S1). Interestingly, dopamine neurons within the PPL1 cluster are lost in numerous genetic (20, 34-36) and pesticide (37, 38) models of PD, typically after a period of aging, consistent with an interaction between genetic/environmental factors and an inherent vulnerability to aging of this cell population suggested by our data.

Figure 1. Age-related dopamine neuron degeneration correlates with life span and locomotor dysfunction. (A), Kaplan-Meier survival curves for 27 DGRP strains (see Table S1 for details and strain IDs). Each line represents a single genotype. (B), mean life span $\pm$ standard error for the 27 strains in (A). SEM life span $\leq$ 2.0 days; log-rank across 27 strains (χ^2 = 2866, p <1e^{-15}). Strain IDs in Table S1. (C), TH-positive dopamine neuron number in the PPL1 cluster (n=7-12 brains/genotype), plotted against strain mean life span with six DGRP strains annotated, Spearman r and p values annotated. See Table S1 for strain IDs. (D), Confocal projection images of PPL1 dopamine neurons for 30-day-old flies belonging to the six DGRP strains annotated in (C). Scale bar, 10µM. (E), PPL1 cell counts in 1-day-old and 30-day-old flies for all DGRP strains with less than the average cell counts across all 27 strains at 30 days (dotted line). Two-way ANOVA for effect of age (p<0.0001) and DGRP strain (ns), Šídák's multiple comparison test, ***p<0.001 (n=5-12 brains/genotype/age). (F) Long-lived strain PPL1 counts at 30 days. (G) locomotor function in negative geotaxis assays correlates with PPL1 dopamine neuron counts in aged flies (n= 4 groups of 25 flies/genotype for geotaxis and 7-12 brains/genotype, Spearman r and p values annotated). Strain IDs in Table S1. Data are mean $\pm$ SEM.

In contrast, no other dopamine neuron cluster counts correlate with mean life span (Fig.

S1B-F). Informatively, dopamine neuron deficits in the PPL1 cluster are age-dependent

as most strains with deficits at 30 days of adulthood exhibit near-maximal PPL1 counts

at 1 day of adulthood (Fig. 1E). Long-lived strains generally retain near-maximal PPL1

counts at 30 days of age (Fig. 1F), hence, we did not assess them at a younger age.

As in mammals, locomotor function in *Drosophila* is thought to be influenced by

dopamine signaling (39) and previous studies indicate a role for PPL1 cluster neurons

(40). In support of this, we observe that silencing the *ple* gene, which encodes TH,

using several PPL1 cluster-specific split-GAL4 drivers impairs locomotor function in

negative geotaxis assays (Fig. S2A). Each individual split-GAL4 driver is reported to

express in only a small subset of PPL1 neurons (41), hence the magnitude of climbing

defect caused by silencing *ple* across all PPL1 dopamine neurons is unknown and likely

underestimated by the deficits seen here. Since PPL1 dopamine neuron viability is

important in *Drosophila* locomotor function, we queried whether strain locomotor

performance is impaired upon age-related dopamine neuron loss. Indeed, negative

geotaxis performance of 4-week-old flies correlates moderately with PPL1 neuron

viability across the same 27 DGRP strains (Fig. 1G), but not with other dopamine

neuron clusters (Fig. S2B-F), thus supporting a role for PPL1 neurons in age-related

locomotor performance. As can be inferred, negative geotaxis scores additionally

correlate well with strain mean life span (Fig. S2G), consistent with prior reports linking

activity levels with mean life span (30).

Although the association between life span and dopaminergic neurodegeneration

appears largely restricted to a single dopamine neuron cluster (PPL1), we next probed

whether shorter-lived strains exhibiting PPL1 dopamine neuron loss experience broader

neurodegeneration upon aging. We first assessed brain vacuole formation as a marker

of general brain neurodegeneration (42) in the three shortest-lived strains exhibiting the

most pronounced age-related PPL1 dopamine neuron loss (RAL-727, RAL-911 and

RAL-356) and three long-lived strains with the highest PPL1 dopamine neuron counts

(RAL-335, RAL-379 and RAL-821). As a positive control for vacuolization, we included a

swiss cheese (*sws*) mutant known to display broad age-related neurodegeneration (42).

Figure 2. No generalized neurodegeneration in short-lived DGRP strains. (A) Brain
vacuoles (see Methods) assessed in short-lived and long-lived DGRP strains aged for 5
weeks (inset magnified images show vacuoles adjacent to asterisks). Occasional
vacuoles in DGRP brains were minor in number (B) and area (C) compared to *sws*[4]
heterozygous mutant females aged 2 weeks. Individual ANOVAs for effect of genotype:
Vacuole number (p<0.01 without *sws*; p<0.0001 with *sws*, n=6-13 brains/genotype);
vacuole area (p<0.01 without *sws*; p<0.0001 with *sws*), Dunnett's post-hoc analysis (****
p<0.0001, n=6-13 brains/genotype).

Both vacuole area and number were modest in all DGRP strains when compared to *sws*

(Fig. 2A-C), and did not significantly correlate with PPL1 dopamine neuron loss across

the six DGRP strains (Fig. S3A). Further, 5-HT-positive serotonin neuron counts do not

differ significantly between 4-week-old short- and long-lived strains (Fig. S3B),

suggesting no change in viability of this independent monoaminergic cell population.

Taken together, our data indicate that short life span arising from natural genetic

variation in *Drosophila* correlates robustly with loss of PPL1 dopamine neuron viability

across a subset of DGRP strains, and with a motor performance task dependent on PPL1 neuron function, but does not appear to be associated with general neurodegeneration.

Metabolomics reveals diverging glutathione levels in vulnerable strains

Metabolism is thought to be a central driver of animal and human aging, whereby the metabolome provides insight into aging-related gene expression changes and the influence of environment on the senescence of cells, tissues and biological systems (4, 43). Consistent with this, life span has been extensively linked to dietary intake in model organism species including flies (43-46) and recent GWAS on the DGRP have additionally highlighted a role for metabolic function in life span determination (26-28). We reasoned that metabolic factors may promote age-related loss of dopamine neurons found in short-lived strains and may be uncovered by metabolic profiling. To assess whether strains that exhibit age-related loss of dopamine neuron viability diverge metabolically from strains that retain dopamine neurons across age, we profiled targeted metabolite levels of the same three short-lived and long-lived strains described above, at 3 weeks of age. We were able to confidently assess 147 aqueous metabolites across all six genotypes (Data S1). Principal component analysis (PCA) shows clear clustering of biological replicates within each genotype, although the short-lived strains and long-lived strains are not readily separable (Fig. 3A). Nonetheless, we find that 64 out of 147 metabolites measured are significantly different (false discovery rate (FDR)<0.1) between grouped short-lived and long-lived DGRP strains (Data S1).

Figure 3. Metabolomic profiles of short- and long-lived strains. (A) Principal components 1 and 2 of head tissue samples, colored by DGRP strain (strain numbers annotated). Ellipses are 95% confidence interval regions. Sample replicates cluster well within a genotype and are not readily separable by life span group. (B) Pathway analysis of significantly differing metabolites, integrating enrichment analysis (y-axis) and pathway topology analyses (x-axis). (C) Volcano plot of 147 metabolites assessed, showing the most downregulated and upregulated metabolites based on $\log_2$(fold-change) at a FDR<0.1 threshold (dotted line). (D-E) head total glutathione content in DGRP strain females measured at the indicated ages (n=2-4 replicates of 25 heads/replicate) and correlated to PPL1 dopamine neuron counts or mean life span. Strain IDs for (D-E) are provided in Methods.

For these differing metabolites, we carried out a metabolite enrichment analysis

and identified the most significantly impacted metabolic pathways, which are cysteine

and methionine metabolism and aminoacyl-tRNA biosynthesis. Factoring in pathway

topology and relative importance of metabolites within a pathway to our analysis, the

biggest pathway impacts are arginine biosynthesis and glycine, serine and threonine metabolism (Fig. 3B). Metabolite-level analysis of significantly differing metabolites reveals that reduced and oxidized glutathione are the most downregulated metabolites in short-lived fly strains when rank-ordered by fold-change, while N-6-trimethyllysine was the most upregulated metabolite (Fig. 3C and Data S1). Lower reduced and oxidized glutathione content in short-lived strains suggests that total glutathione levels may be divergent across DGRP strains in a manner that associates with life span and age-related dopamine neuron loss. The importance of glutathione as an intracellular antioxidant and the predominant brain antioxidant in many species including flies prompted us to measure total glutathione levels in an expanded set of DGRP strain fly heads (see methods for strain IDs). In support of our metabolomics data, total glutathione levels in 1-week-old females correlate robustly with PPL1 dopamine neuron viability of 30d-old flies and moderately with life span (Fig. 3D), while no correlation exists when flies reach 4 weeks of age (Fig. 3E). Most strains manifest an age-related reduction in total glutathione levels (mean across 13 tested strains = $20.5 \pm 3.4\%$). An age-related decrease in glutathione seen here in flies mirrors observations from rat brain (47) and mouse brain (48), where in mice an age-related decrease in total glutathione levels was recorded for male and female mice across most brain regions, including midbrain.

Boosting glutathione biosynthesis rescues age-related dopamine neuron loss and enhances life span in short-lived strains

Prior studies have shown a major influence of glutathione levels on *Drosophila* survival. Increasing total glutathione levels by overexpressing the catalytic subunit of glutamate-cysteine ligase (Gclc) or feeding flies with the cysteine donor N-acetyl-cysteine can extend life span up to 50%, while RNAi-mediated Gclc silencing truncates life span by a similar magnitude (49-51). Given the robust link between Gclc expression, Gcl activity and glutathione levels, we decided to examine whether neuronal Gclc overexpression could protect dopamine neurons from age-related attrition in our short-lived DGRP strains. We first outcrossed our core six DGRP strains to a standard laboratory control strain, w^{1118}, in order to determine whether DGRP strains retain life span and dopamine neuron characteristics after single generation outcrossing. We observe good correlation between outcrossed and parental DGRP strains for PPL1 dopamine neuron viability (Fig. S4A), supporting the validity of an outcrossing approach to assess Gclc overexpression effects on dopamine neuron viability. We next outcrossed our core three short-lived DGRP strains plus one long-lived DGRP strain (RAL-821) to flies overexpressing Gclc pan-neuronally. This results in a modest increase in total head glutathione content relative to outcross controls (Fig. S4B), but a strong rescue of TH-positive cell counts for the PPL1 dopamine neuron cluster across all three short-lived DGRP backgrounds (Fig. 4A and B), indicating that age-related attrition of dopamine neurons across these naturally varying strains is dependent on neuronal glutathione levels. Consistent with a prior study (50), we also observe a life span extension via pan-neuronal Gclc overexpression, and this extension appears to be restricted to short-lived genetic backgrounds (Fig. 4C and Table S2), suggesting that further elevating glutathione levels in long-lived strains has no impact on longevity.

Figure 4. Glutathione dependency of age-related dopamine neuron loss (A) PPL1 dopamine neuron rescue in short-lived DGRP backgrounds following pan-neuronal Gclc overexpression, quantified in (B), two-way ANOVA for effect of Gclc overexpression (p <0.0001) and DGRP strain (p <0.0001) with Bonferroni's post-test (** p <0.01, **** p <0.0001, n =10-12 brains/genotype). Genotypes are *elav-GAL4/+* (ctrl) and *elav-GAL4/UAS-gclc* (Gclc) in the DGRP background indicated. (C) Survival is enhanced in three short-lived strains following pan-neuronal GCLc overexpression, but not in a long-lived strain (individual log-rank tests).

Divergent glutathione levels are associated with glutathione biosynthetic enzyme expression

De novo biosynthesis of glutathione is principally regulated by the rate-limiting enzyme Gcl and by the availability of its substrates, in particular, cysteine (52). The product of Gcl activity, γ-glutamylcysteine, is subsequently conjugated to glycine by glutathione synthase (Gss) to form glutathione. While our metabolomics pathway analysis indicates that *cysteine and methionine metabolism* is divergent between short- and long-lived strains, we did not uncover any variance in levels of cysteine itself that could readily explain differences in glutathione content (Data S1). Glycine levels appear

slightly elevated in short-lived strains (Data S1), while glutamate levels were not

detected in these samples. To determine if the differing total glutathione levels in short-

and long-lived strains results from altered glutathione biosynthesis, we probed

expression of Gcl catalytic and modulatory subunits (*gclc* and *gclm*) as well as

glutathione synthase (*Gss*) in the same 13 DGRP strains (Fig. S5A). This uncovered a

positive relationship between glutathione levels and gclc expression (approaching

statistical significance at r = 0.54, p = 0.06), while neither *gclm* nor *Gss* levels correlate

with glutathione levels (Fig. S5A and B). We also assessed expression of gamma

glutamyl transpeptidase (*Ggt-1*), an extracellular enzyme that breaks down glutathione

exported from astrocytes to cysteinylglycine, which then supplies neurons with cysteine

and glycine substrates for glutathione synthesis (53), although there is no apparent

relationship between *Ggt-1* expression and glutathione (Fig. S5A). Hence, divergent

glutathione levels may be at least partly explained by expression of the catalytic subunit

of Gcl, the rate-limiting enzyme in glutathione biosynthesis.

ROS levels are augmented in age-related dopamine neuron vulnerability

Considering that glutathione's antioxidant activity is crucial for neutralizing

damaging intracellular ROS and lipid peroxides, our findings linking lower glutathione

levels to shorter longevity and dopaminergic neurodegeneration implicate oxidative

stress as a possible underlying mechanism. In support of this, we find that total H_2O_2

levels increase with age selectively in the heads of short-lived strains, while H_2O_2 levels

remain stable in long-lived strains over the same aging period (Fig. 5A). Importantly,

Figure 5. ROS levels and oxidative stress sensitivity in short-lived backgrounds
(A) Head H_2O_2 levels correlate inversely with strain mean life span at 10 days of age, but not at 2 days, for three short-lived (RAL-727, RAL-911, RAL-356) and three long-lived (RAL-335, RAL-379, RAL-821) DGRP backgrounds (n=3 groups of 10 heads/genotype). (B) Head H_2O_2 levels are reduced following Gclc overexpression (two-way ANOVA for effect of Gclc (p <0.01) and DGRP background (p <0.0001), Šídák's multiple comparison test, ** p <0.01, *** p <0.001, n =3 groups of 10 heads/genotype). Genotypes are *elav-GAL4/+* (ctrl) and *elav-GAL4/UAS-GCLc* (GCLc) in the DGRP background indicated. (C) Correlation of mean life span and survival on H_2O_2-containing food for the same six DGRP backgrounds as in (A). (D) PPM neurons in three short-lived and three long-lived DGRP backgrounds outcrossed to *TH-GAL4* (ctrl) or to *TH-GAL4/UAS-parkin-IR* (dopamine neuron-specific *parkin* RNAi line 37509). All three short-lived backgrounds and one long-lived background (RAL-335) exhibit dopamine neuron loss upon *parkin* knock-down, quantified in (E), two-way ANOVA for effect of parkin knock-down, p <0.0001, and DGRP background, p <0.01, interaction p <0.0001, Šídák's multiple comparison test, * p <0.05, ** p <0.01, *** p <0.001, n=10-13 brains/genotype.

pan-neuronal *gclc* overexpression is sufficient to attenuate the elevated H_2O_2 levels

seen in the heads of aged short-lived backgrounds (Fig. 5B), thus linking a rescue of

dopamine neuron viability and life span in short-lived lines with ROS amelioration.

Elevated ROS levels may render short-lived strains prone to oxidative stress.

Consistent with an underlying vulnerability, short-lived strains exhibit impaired survival

upon exposure to H_2O_2-containing food is relative to long-lived strains (Fig. 5C).

Additionally, silencing expression of *parkin* in all dopamine neurons induces age-related

degeneration of PPM dopaminergic neurons in all three short-lived DGRP strains, while

two out of three long-lived strains appear resilient to the loss of *parkin* and retain

dopamine neurons (Fig. 5D and E and Fig. S6A). Abundant evidence indicates that

Parkin loss-of-function, which is linked to familial PD, results in oxidative stress (54-57),

hence these results further corroborate a vulnerability of short-lived strains to oxidative

stress. Silencing *parkin* does not significantly exacerbate the loss of PPL1 neurons in

any DGRP lines tested (Fig. S6B), suggesting that PPL1 degeneration is not impacted

by *parkin* loss-of-function to the same extent as PPM neurons.

Glutathione biosynthesis enzyme expression is reduced in human LRRK2

G2019S PD brain

Deficiencies in reduced and/or total glutathione are an early biochemical feature

detected in PD patient brain (58-61). In a recent high-powered study applying in-depth

quantitative proteomics to substantia nigra tissue from 15 idiopathic PD patients and 15

healthy controls, GCLc downregulation was observed in SN from PD patient brain

relative to healthy controls via the authors' bootstrap ROC-based analysis (62). No

changes in GCLm, GSS and GGT were detected in this study (62). Most familial PD-

linked genetic mutations, including the common pathogenic LRRK2 G2019S mutation,

have been associated with enhanced oxidative stress sensitivity (6, 63-65). We queried

whether mutant LRRK2 G2019S PD brain shows evidence of altered glutathione

biosynthetic capacity, hypothesizing that LRRK2 G2019S might interact with an

underlying glutathione deficiency to promote neurodegeneration. Interestingly, we find a

significant reduction in both catalytic and modulatory GCL subunits in cortical extracts

from LRRK2 G2019S PD brain relative to tissue from healthy age-matched controls

(Figure 6A and B), raising the possibility that glutathione biosynthesis may be reduced

in PD associated with mutant LRRK2-mediated neurodegeneration.

Figure 6. GCL levels reduced in LRRK2 G2019S PD brain. (A) Levels of GCLc and
GCLm in human brain cortical extracts from PD patients harboring LRRK2 G2019S are
different from age-matched healthy controls, quantified in (B) relative to β-actin and
normalized, analyzed by multiple unpaired Student's t-tests with Benjamini, Krieger and
Yekutieli correction for multiple comparisons, * q <0.05, n=5 per group.

DISCUSSION

We leveraged natural genetic variation across a multitude of inbred *Drosophila* strains to obtain mechanistic insight into aging-dependent dopaminergic neurodegeneration. Our major findings are that age-related degeneration of a dopamine neuron subset correlates strongly with life span across naturally varying strains (Fig. 1), and that dopamine neuron degeneration may result at least in part when animals are predisposed to oxidative stress due to relatively low glutathione antioxidant capacity and elevated ROS levels. When compared to naturally long-lived DGRP strains, the short-lived strains characterized here generally harbor lower total glutathione levels in head tissue (Fig. 3), higher ROS levels, and are more sensitive to exogenous oxidants and to loss of *parkin* expression (Figs. 4 and 5). Boosting glutathione levels through neuronal overexpression of the Gcl catalytic subunit (Gclc) is sufficient to attenuate ROS levels, prevent age-related dopamine neuron loss and enhance life span in short-lived strains (Figs. 4 and 5). Flies overexpressing Gclc via a neuronal Gal4 driver were previously shown to manifest life span extension over controls harboring *GAL4* or *UAS* transgenes alone (50), thus attributing this phenotype to Gclc overexpression and not genetic background. Our studies build on these findings, by demonstrating that short-lived strains expressing Gclc pan-neuronally have enhanced survival and are additionally protected from ROS accumulation and age-related dopamine neuron loss. Collectively, these findings are consistent with short-lived strains having a baseline vulnerability to oxidative stress that impacts aging, both at the level of dopamine neuron viability and life span. One significant aspect of these findings is that they underscore a potential mechanistic link between dopamine neuron degeneration and reduced glutathione

levels, which is an early biochemical feature seen in human PD brain tissue (58-61).

There are multiple lines of evidence implicating glutathione deficiencies in PD

pathogenesis. Prior studies on human PD brain show a depletion of total and/or

reduced glutathione in substantia nigra relative to healthy controls (58-61), and a

greater glutathione depletion is often seen in substantia nigra relative to other brain

regions (58-61). Outside of brain, total glutathione levels in blood are found to correlate

inversely with PD severity and may also have utility as a disease biomarker in

prodromal PD (66, 67). Further, PD has been associated with polymorphisms in a

number of glutathione-S-transferase genes important for glutathione function (68-70).

While vulnerability to oxidative stress is an established consequence of pathogenic

mutations in familial PD-linked genes such as *LRRK2*, *Parkin*, *PINK1*, and *DJ-1* (64,

71), it is not yet known whether glutathione depletion is a risk factor for disease onset in

familial PD mutation carriers. We demonstrate here that PD patients harboring the

LRRK2 G2019S mutation exhibit reduced levels of GCL catalytic and modulatory

subunit expression in postmortem brain tissue. This warrants further investigation into

plasma and CNS glutathione levels in LRRK2 mutation carriers, both in patients with PD

and in those without PD in order to assess whether glutathione levels diverge between

these groups in a manner that could be relevant to disease onset.

     We opted for a metabolic analysis to delineate differences between short- and

long-lived strains influencing dopamine neuron aging rather than attempting to identify

gene expression differences between strains. This is because aging is heavily impacted

by metabolism (72, 73) which is a dynamic product of both genetic and environmental

factors. Hence, one benefit of metabolomics is that it provides insight on the biological

influence of environment as well as gene expression on age-related dopamine neuron degeneration. This approach yielded important differences in glutathione levels that, through subsequent genetic studies, we show may relate to the expression of Gclc, a rate limiting enzyme in glutathione biosynthesis.

ROS play a physiological role in redox-dependent signaling pathways, but when ROS levels outweigh cellular antioxidant capacity, oxidative stress ensues (74). An age-related increase in macromolecular oxidative damage is observed in humans (75) and model organisms such as *Drosophila* (76-78) and may occur in part due to higher ROS production from dysfunctional mitochondria (79-81). While antioxidant overexpression studies have produced mixed results in animal models (79, 82-84), we previously showed that genetically blocking the function of key cellular antioxidants such as the superoxide dismutases SOD1 or SOD2 often truncates life span and healthspan in flies (85, 86). The same can be seen in mice (87), supporting the role of ROS in aging across multiple species.

A wealth of evidence indicates that oxidative stress in the brain plays a central role in neurodegenerative disease (88), and certain neuronal populations may be more susceptible to oxidative stress due to their intrinsic properties (5). Dopamine neurons are thought to experience high levels of oxidative stress (89, 90) because they extend highly branched axonal arbors requiring abundant mitochondria, a primary ROS source (91), for metabolic support and because metabolism or auto-oxidation of DA directly produces harmful ROS and DA quinones (92). Indeed, depletion of cytosolic dopamine protects against dopamine neuron death in genetic (17, 93) and rotenone (94, 95) *Drosophila* PD models. Animal studies have established a role for oxidative stress

sensitivity in Parkinson's disease (PD)-related neurodegeneration caused by

environmental toxins (96) and familial PD mutations (97). In particular, *Parkin* and

PINK1 are thought to be important in protecting cells against oxidative stress and

mitochondrial dysfunction and evidence in flies suggests that oxidative stress is

important in *parkin* mutant degenerative phenotypes (20, 57, 71). We therefore

postulated that loss of *parkin* may be more likely to cause overt dopamine neuron

degeneration in 'at-risk' genetic backgrounds exhibiting heightened baseline ROS levels

and lower glutathione antioxidant capacity, such as the short-lived DGRP strains

characterized here. Indeed, all short-lived strains assessed display a loss of PPM

dopamine neurons upon *parkin* silencing, while two out of three long-lived strains show

no loss of these neurons (Fig. 5). These findings may help clarify why there are

numerous reports in flies both supporting (17, 20, 98) and refuting (56, 99, 100)

dopamine neuron loss following loss of *parkin* expression, as genetic background may

be pivotal in determining whether this phenotype manifests. PPM neuron loss in one of

three long-lived backgrounds assessed (RAL-379, Fig. 5) complicates the interpretation

of our findings somewhat and suggests that, while the underlying mechanisms are not

yet clear, not all long-lived fly backgrounds are resistant to the effects of *parkin* loss on

dopamine neurons. As (i) resistance to oxidative stress and ROS levels for RAL-379

align with the other long-lived backgrounds tested (ii) these phenotypes are divergent

from the elevated ROS and peroxide sensitivity of short-lived strains and (iii) *parkin* is

important for mitochondrial function beyond oxidative stress, one speculative

explanation is that this strain may harbor sensitivity to deficits in mitochondrial quality

control independent of oxidative stress, thus rendering it more susceptible to loss of

parkin. Although *parkin* silencing impacts PPM neuron counts, it does not appear to worsen PPL1 neuron loss in our short-lived strains, suggesting that either PPL1 cells are refractory to further increases in oxidative stress resulting from loss of *parkin*, or that different neuron clusters may respond to individual oxidative stress stimuli in a non-uniform manner. Our assessment of dopamine neuron degeneration consisted of identifying a loss of neurons with detectable tyrosine hydroxylase (TH) signal, and we note the possibility that surviving neurons with no TH signal may exist, yet the loss of dopamine synthesis capacity in these neurons would render them non-functional and degenerative.

While a major contribution of oxidative stress to PD has been established, identifying therapeutic strategies to combat oxidative stress that are clinically beneficial in PD remains a paramount goal. Our data from flies and from human PD postmortem brain in the context of existing literature reinforce the role of glutathione in maintaining brain health. Glutathione is a major brain antioxidant thought to be a first line of defense in the protection of dopamine neurons from oxidative stress (53, 101). Glutathione promotes redox homeostasis by neutralizing a range of reactive oxygen and nitrogen species, maintaining the reduced state of protein sulfhydryls, and by protecting cells from xenobiotic toxins (53, 101). While it remains unclear what causes glutathione depletion in PD, this is likely to render the nervous system susceptible to oxidative stress capable of promoting neurodegeneration (53, 102). In animal models, blocking glutathione production by inducible GCLc silencing can cause dopaminergic neurodegeneration in mice (103, 104) and rats (105), consistent with the GCLc downregulation we observe in human PD brain contributing to neurodegeneration

(Figure 6). Downregulation of GCLc has also been reported in a high-powered proteomics study on idiopathic PD brain nigral tissue (62). Yet, changes in glutathione biosynthesizing enzymes including GCLc are not seen in human PD brain with the same consistency as glutathione depletion (58), hence, alternative mechanisms are likely to be involved.

Glutathione augmentation has received considerable attention as a potential PD therapeutic strategy based on a wealth of supportive evidence across a range of PD animal models (106-109). Supplementation with glutathione or precursor molecules such as N-acetylcysteine (NAC) have been the focus of several completed and ongoing clinical trials for PD treatment. Intranasal administration of glutathione can increase total brain glutathione levels (110), although Phase II clinical trials testing the efficacy of intranasal glutathione on PD symptoms were inconclusive due to placebo effects (111). Intravenous NAC administration boosts brain total glutathione levels in PD patients (112) and a recent clinical trial showed that combined intravenous and oral administration of NAC may have a beneficial effect in PD, as it results in significantly increased dopamine transporter binding and improved PD symptoms (113). While encouraging, this study had several limitations and larger scale efficacy studies are needed to provide clarity on the potential benefits of NAC in PD.

The cause of glutathione depletion in human PD brain is not well understood, with existing theories of reduced biosynthesis or increased efflux of glutathione conjugates lacking consistent supportive evidence (53). The apparent deficit of total glutathione seen here in short-lived strains may be at least partly related to lower expression of Gclc (Fig. S5) which could negatively impact the biosynthetic activity of

Gcl. We do not observe any alterations in glutathione precursor substrates (Data S1) or other biosynthetic enzymes (Fig. S5) that can explain this deficit. It is also possible that an increase in glutathione conjugates resulting from oxidative stress could contribute to lower free glutathione levels in short-lived strains. Glutathione conjugates to toxic metabolites such as 4-hydroxynonenal and oxidatively modified proteins via protein S-glutathionylation in response to oxidative stress (114, 115). These conjugates would not be detected in the pool of free total glutathione, and the elevated ROS measured in short-lived strains may promote their formation, potentially accounting for the differences in observed glutathione levels. Interestingly, the increase in expression of the extracellular enzyme Ggt-1 in short-lived strains is consistent with an attempt to utilize astrocyte-released glutathione as a substrate for neuronal glutathione synthesis via extracellular breakdown and parallels an elevation of GGT1 observed in human PD postmortem brain (58), suggesting a possible compensatory mechanism.

In summary, our study harnesses natural genetic variation to examine the relationship between life span and dopamine neuron degeneration at the molecular level. Our findings support a pivotal role for glutathione deficits in age-related dopamine neuron loss occurring across a collection of naturally varying strains. Lower glutathione as a driver for age-related dopamine neurodegeneration may also be relevant to human PD, where brain glutathione depletion is detected early on in disease (53, 101) and suggests that glutathione could be an important component driving the influence of aging on PD neurodegeneration.

ACKNOWLEDGMENTS

We thank the Northwest Metabolomics Research Center for performing *Drosophila* sample extraction and metabolite analysis, the OHSU Advanced Light Microscopy Core for subsidized confocal microscope use, OHSU Neuropathology Division and Randy Woltjer for providing post-mortem brain tissue, Ben Harrison (University of Washington) for assistance with metabolomic sample preparation, Bill Orr (Southern Methodist University) for providing the *UAS-GCLc* fly line and Doris Kretzschmar (OHSU) for providing the *sws^4* line.

This work was supported by National Institute on Aging Grant T32AG055378 to C.R.C and to J.P., National Institute on Aging Grant P30AG013280, R01AG063371 and R56AG049494 to D.E.L.P., NIH instrumentation grant S10 OD021562 to D.R., National Institutes of Health Grant P30NS061800 to the OHSU Advanced Light Microscopy Core, and OHSU Neurology Foundation Funds to I.M.

AUTHOR CONTRIBUTIONS

C.R.C - Conceptualization, Investigation, Formal analysis, Writing manuscript

J.P.- Conceptualization, Investigation, Formal analysis, Writing manuscript

A.A-B.- Investigation, Formal analysis, Writing manuscript

L.W.- Formal analysis

D.R.- Supervision, Funding acquisition

D.E.L.P.- Conceptualization, Supervision, Funding acquisition

I.M. - Conceptualization, Investigation, Formal analysis, Writing manuscript, Supervision, Funding acquisition.

DECLARATION OF INTERESTS

The authors declare no competing interests.

METHODS

Drosophila stocks and culture

The following DGRP strains were used in this study and obtained from the Bloomington *Drosophila* Stock Center: RAL-409, RAL-911, RAL-177, RAL-765, RAL-727, RAL-356, RAL-83, RAL-239, RAL-703, RAL-596, RAL-461, RAL-310, RAL-373, RAL-639, RAL-426, RAL-371, RAL-502, RAL-820, RAL-313, RAL-360, RAL-379, RAL-382, RAL-714, RAL-819, RAL-136, RAL-335, RAL-821. *sws⁴* mutant flies have been characterized elsewhere (42) and were a gift from Doris Kretzschmar. *UAS-GCLc* flies were a gift from Bill Orr and have been previously described (line GCLc5 from (50)). PPL1-specific split-GAL4 drivers *MB058B-GAL4* (68278), *MB060B-GAL4* (68279), *MB304B-GAL4* (68367), *MB065B-GAL4* (68281), *MB438B-GAL4* (68326), *MB296B-GAL4* (68308), along with *UAS-ple-IR-1* (25796), *UAS-ple-IR-2* (76089), *UAS-parkin-IR* (31259 and 37509), *elav^{C155}-GAL4* (458) and *TH-GAL4* (8848) were obtained from the Bloomington *Drosophila* Stock Center. All flies were reared and aged at 25°C and 60% relative humidity under a 12 h light-dark cycle on standard food medium.

Targeted metabolomic assay

Three biological replicates of three DGRP strains designated 'short-lived' (RAL-727, RAL-911 and RAL-356) and three strains designated 'long-lived' (RAL-335, RAL-379, RAL-821) were aged to three weeks then flash frozen in liquid nitrogen in rapid succession and pulse vortexed to detach heads which were subsequently collected on dry ice (25 female fly heads/genotype/replicate).

Sample Preparation:

Aqueous metabolites for targeted LC-MS profiling of *Drosophila* head samples were extracted using a protein precipitation method similar to the one described elsewhere (116, 117). Samples were first homogenized in 200 µL purified deionized water at 4 °C, and then 800 µL of cold methanol containing 124 µM 6C13-glucose and 25.9 µM 2C13-glutamate was added (reference internal standards were added to the samples in order to monitor sample prep). Afterwards the samples were vortexed, stored for 30 minutes at -20 °C, sonicated in an ice bath for 10 minutes, centrifuged (14,000 rpm, 15 min and 4 °C), and then 600 µL of supernatant was collected from each sample. Lastly, recovered supernatants were dried in a SpeedVac and reconstituted in 0.5 mL of LC-matching solvent containing 17.8 µM 2C13-tyrosine and 39.2 3C13-lactate (reference internal standards were added to the reconstituting solvent in order to monitor LC-MS performance). Samples were transferred into LC vials and placed into a temperature controlled autosampler for LC-MS analysis.

LC-MS Assay

Targeted LC-MS metabolite analysis was performed on a duplex-LC-MS system composed of two Shimadzu UPLC pumps, CTC Analytics PAL HTC-xt temperature-controlled auto-sampler and AB Sciex 6500+ Triple Quadrupole MS equipped with ESI ionization source (117). UPLC pumps were connected to the auto-sampler in parallel and were able to perform two chromatography separations independently from each other. Each sample was injected twice on two identical analytical columns (Waters XBridge BEH Amide XP) performing separations in hydrophilic interaction liquid

chromatography (HILIC) mode. While one column was performing separation and MS data acquisition in ESI+ ionization mode, the other column was getting equilibrated for sample injection, chromatography separation and MS data acquisition in ESI- mode. Each chromatography separation was 18 minutes (total analysis time per sample was 36 minutes). MS data acquisition was performed in multiple-reaction-monitoring (MRM) mode. LC-MS system was controlled using AB Sciex Analyst 1.6.3 software. Measured MS peaks were integrated using AB Sciex MultiQuant 3.0.3 software. The LC-MS assay was targeting 363 metabolites (plus 4 spiked reference internal standards). Up to 157 metabolites (plus 4 spiked standards) were measured across the study set, and 148 were measured with <20% missingness across all the samples. In addition to the study samples, two sets of quality control (QC) samples were used to monitor the assay performance as well as data reproducibility. One QC [QC(I)] was a pooled human serum sample used to monitor system performance and the other QC [QC(S)] was from pooled study samples and this QC was used to monitor data reproducibility. Each QC sample was injected per every 10 study samples. The data were well reproducible with a median CV of 4.25 % for QC(I) and 5.20 % for QC(S), respectively.

Analysis

Raw data were $\log_2$-transformed and median normalized. After excluding metabolites with $\geq$ 20% missingness and $\geq$ 30% CV in quality control samples, 147 metabolites remained. We used a quantile regression approach for the imputation of left-censored missing data (QRILC), which has been suggested as the favored imputation method for left-censored Missing Not at Random (MNAR) data (Runmin Wei and Ni 2018). Briefly,

QRILC performs the feature-level imputation by replacing missing values with random draws from a truncated distribution with parameters estimated using quantile regression. This is implemented in the R imputeLCMD package.

We used a simple linear model to test for significant effects of life span group (short-lived versus long-lived) on metabolomic data. This was applied to the normalized and imputed metabolomic data using the Bioconductor limma package (118). The limma package uses empirical Bayes moderated statistics, which improves power by 'borrowing strength' between metabolites in order to moderate the residual variance (119).

PCA analysis and metabolite pathway analysis were carried out in MetaboAnalyst. The targeted pathway analysis function was applied to significantly different metabolite KEGG IDs (FDR<0.1), including Fisher's Exact Test for Over Representation Analysis and Relative-betweenness Centrality for Pathway Topology Analysis, and uploading a reference metabolome with the KEGG IDs of all metabolites measured in the assay.

Glutathione content

Flies (25/genotype/replicate) were flash frozen in the morning (9-11 am) at the indicated ages in order to minimize any potential effect of circadian rhythm on glutathione levels (120). Frozen flies were pulse-vortexed to separate heads and the heads were collected into a microcentrifuge tube containing 100 mM phosphate buffer. The heads were homogenized and centrifuged (17,000 g, 10 min, 4°C) and processed for measuring

total glutathione using the Invitrogen Glutathione Colorimetric Detection Kit and a SpectraMax i3 Microplate Reader for measuring OD (405nm). Wells containing fly head homogenate without assay reagents were used to control for any background absorbance due to eye pigment, and this background was subtracted from OD (405nm) readings. Male and female flies of the following DGRP strains were assessed: RAL-136 RAL-177, RAL-335, RAL-356, RAL-360, RAL-373, RAL-379, RAL-409, RAL-727, RAL-765, RAL-819, RAL-821 and RAL-911.

H_2O_2 content

Levels of H_2O_2 in DGRP fly heads were determined by Amplex Red assay, using the Amplex Red Hydrogen Peroxide Kit (Invitrogen). Flies (10/genotype/replicate) were flash-frozen in liquid nitrogen and pulse-vortexed to separate heads which were collected on dry ice. Heads were homogenized on ice in sodium phosphate buffer and centrifuged (17,000 g, 10 min, 4°C). The Amplex Red working solution was then applied to 50 μL aliquots of homogenate supernatant or to H_2O_2 standards and OD (560nm) was recorded after 60 min incubation at room temperature in a SpectraMax i3 Microplate Reader. Controls consisting of 50 μL head homogenate supernatant and phosphate buffer in place of Amplex Red working solution were also measured at 60 min in order to subtract any background OD (560nm) signal arising from eye pigment. H_2O_2 concentrations were interpolated from the standard curve and normalized to protein concentration following BCA assay.

Brain vacuole assessment

Vacuolization was assessed in brains following a previously described protocol (121) with slight modification as follows: Heads of 5-week-old flies were pre-fixed in 4% PFA (pH 7.4) for 15 min and the dissected brains were post-fixed in 4% PFA (pH 7.4) for 20 min, both times at RT on a nutator. Phalloidin (DyLight 650 Phalloidin, Cell Signaling Technology) and DAPI staining were performed sequentially, with phalloidin incubation as described (121) followed by 15 min incubation of brains with DAPI (1:1000). PBS-T was used at 0.3% Triton-X-100 throughout the protocol. Confocal z-stacks were obtained on a Zeiss LSM 900 microscope and vacuole analysis was carried out in Fiji as described (121). As described in the published protocol, only spheroidal gaps in tissue were included in the analysis, with a minimum vacuole area threshold of 8 μm^2. Tissue gaps that were tubular in nature when examining across serial confocal z-stack planes were attributed to trachea and not included.

H_2O_2 survival assay

Mated 1-day-old females were collected under brief anesthesia and kept for one days on standard food medium, then transferred to SY food containing 5% H_2O_2 (1.5% agar/5% sucrose/10% yeast (W/V) and 5% H_2O_2 (V/V)). Dead flies were counted every 12 h until all flies were dead.

Quantitative real-time PCR

Total RNA was isolated from fly heads (n = 3-6 biological replicates of 8-17 heads) using TRIzol reagent and quantified on NanoDrop 2000c. Four hundred nanograms of total

RNA was reverse transcribed using Superscript IV first-strand synthesis system, using random hexamers. Quantitative real-time PCR assays were performed in PowerUp SYBR Green Master Mix (Applied Biosystems) using a QuantStudio 3 Real Time PCR system under the following sequence of conditions: 50 °C for 2 min, 95 °C for 10 min then 40 cycles (c) of 15 s at 95 °C and 1 min at 55 °C. Actin 5C and Rpl32 were used as housekeeping genes for normalization.

Fly brain immunostaining for dopamine and serotonin neurons

Brains (12–16/genotype) were harvested, fixed, and permeabilized with 4% PFA in 0.3% PBS-T, pH 7.4, then blocked in 5% normal donkey serum for 1 h at room temperature. Brains were incubated in anti-tyrosine hydroxylase (1:1000, Immunostar) or anti-serotonin (1:1000, Immunostar) for three nights on a nutator at 4°C. After extensive washing, brains were then incubated with Alexa Fluor 488 secondary antibody (1:2000) or Alexa Fluor 568 secondary antibody (1:2000) for 3 nights at 4°C. Brains were washed extensively and whole-mounted in SlowFade Gold Antifade mounting medium. Confocal z-stack images of the stained brains were acquired on Zeiss LSM 900 at a 1 μm slice interval and dopamine neurons with any detectable tyrosine hydroxylase or serotonin neurons with any detectable serotonin were counted. While Dopamine neurons belonging to the protocerebral posterior lateral 1 (PPL1), protocerebral posterior lateral 2 (PPL2), protocerebral posterior medial 1/2 (PPM1/2), protocerebral posterior medial 3 (PPM3) and protocerebral anterior lateral (PAL) clusters were counted. Serotonin neurons belonging to the SP1, SP2, IP and LP1 clusters were counted.

Human LRRK2 G2019S cortical tissue Western blotting

De-identified human cortical tissue from five patients (3 female, 2 male) with clinician-diagnosed PD harboring LRRK2 G2019S and five age-matched control tissue (3 female, 2 male), all with a post-mortem interval range 2-21 h, was provided by OHSU Neuropathology and Sun Banner Health. Samples were extracted in RIPA buffer, electrophoresed on 4-20% Tris-Glycine gradient gels and transferred to nitrocellulose membrane for immunoblotting using the following antibodies: GCLc polyclonal (1:500, Proteintech), GCLm polyclonal (1:500, Proteintech), β-Actin-HRP (1:5000, Sigma Millipore).

Adult survival

Adult flies (0–3 days of age, 100–490 flies per genotype, selecting against newly-eclosed flies) were collected under brief anesthesia and transferred to fresh food vials at 25 flies per vial. Flies were then transferred to fresh food vials every 3–4 days, and the number of dead or censored (escaped or stuck in food) flies was recorded at each transfer, until all flies were dead. Mean life span was derived from Kaplan-Meier analyses carried out in SPSS.

Negative geotaxis behavior

Cohorts of 100 female flies (0–3 days old, selecting against flies with visible signs of recent eclosion) were collected under brief anesthesia and transferred to fresh food vials (25 flies/vial). Flies were aged for four weeks, transferring to fresh food every 3-4 days. For testing, flies were transferred to empty vials, given 1 min to rest and then

tapped to the bottom of the vial three times within a 1 sec interval to initiate climbing. The position of each fly was captured in a digital image 4 sec after climbing initiation, and the measurement was repeated three times. Automated image analysis was performed using the particle analysis tool on Image J to derive x–y coordinates for each fly, thus deriving height climbed as previously described (122). The performance of flies in a single vial was calculated from the average height climbed by all flies in that vial to generate a single datum (N = 1). Performance of each line was then derived from the average scores of 4 vials tested for the line (N = 4).

Statistical analysis

Quantified data are mean $\pm$ SEM, and individual data points are plotted for data with n <10. Sample sizes were determined based on evidence from pilot experiments. Statistical analysis details for each individual experiment are described in figure legends, including the number of flies or number of groups of flies used (n), statistical tests used and associated p values. All statistical analyses were performed using GraphPad Prism except log-rank (Mantel-Cox) tests for life span comparisons performed in SPSS, metabolomic analysis performed in Bioconductor limma and metabolite pathway analysis performed in MetaboAnalyst.

REFERENCES

1. T. J. Collier, N. M. Kanaan, J. H. Kordower, Ageing as a primary risk factor for Parkinson's disease: evidence from studies of non-human primates. *Nat Rev Neurosci* **12**, 359-366 (2011).

2. C. Coleman, I. Martin, Unraveling Parkinson's Disease Neurodegeneration: Does Aging Hold the Clues? *J Parkinsons Dis* **12**, 2321-2338 (2022).

3. M. P. Mattson, T. V. Arumugam, Hallmarks of Brain Aging: Adaptive and Pathological Modification by Metabolic States. *Cell Metab* **27**, 1176-1199 (2018).

4. C. Lopez-Otin, M. A. Blasco, L. Partridge, M. Serrano, G. Kroemer, Hallmarks of aging: An expanding universe. *Cell* **186**, 243-278 (2023).

5. P. Gonzalez-Rodriguez, E. Zampese, D. J. Surmeier, Selective neuronal vulnerability in Parkinson's disease. *Progress in brain research* **252**, 61-89 (2020).

6. D. G. Healy *et al.,* Phenotype, genotype, and worldwide genetic penetrance of LRRK2-associated Parkinson's disease: a case-control study. *The Lancet. Neurology* **7**, 583-590 (2008).

7. S. Costello, M. Cockburn, J. Bronstein, X. Zhang, B. Ritz, Parkinson's disease and residential exposure to maneb and paraquat from agricultural applications in the central valley of California. *Am J Epidemiol* **169**, 919-926 (2009).

8. X. Reed, S. Bandres-Ciga, C. Blauwendraat, M. R. Cookson, The role of monogenic genes in idiopathic Parkinson's disease. *Neurobiology of disease* **124**, 230-239 (2019).

9. P. A. Alladi *et al.*, Absence of age-related changes in nigral dopaminergic neurons of Asian Indians: relevance to lower incidence of Parkinson's disease. *Neuroscience* **159**, 236-245 (2009).

10. Y. Chu, K. Kompoliti, E. J. Cochran, E. J. Mufson, J. H. Kordower, Age-related decreases in Nurr1 immunoreactivity in the human substantia nigra. *J Comp Neurol* **450**, 203-214 (2002).

11. N. Kubis *et al.*, Preservation of midbrain catecholaminergic neurons in very old human subjects. *Brain* **123 (Pt 2)**, 366-373 (2000).

12. G. Rudow *et al.*, Morphometry of the human substantia nigra in ageing and Parkinson's disease. *Acta Neuropathol* **115**, 461-470 (2008).

13. Y. Chu, J. H. Kordower, Age-associated increases of alpha-synuclein in monkeys and humans are associated with nigrostriatal dopamine depletion: Is this the target for Parkinson's disease? *Neurobiology of disease* **25**, 134-149 (2007).

14. X. Chen *et al.*, Parkinson's disease-linked D620N VPS35 knockin mice manifest tau neuropathology and dopaminergic neurodegeneration. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 5765-5774 (2019).

15. T. Kitada, Y. Tong, C. A. Gautier, J. Shen, Absence of nigral degeneration in aged parkin/DJ-1/PINK1 triple knockout mice. *J Neurochem* **111**, 696-702 (2009).

16. Y. Xiong *et al.*, Robust kinase- and age-dependent dopaminergic and norepinephrine neurodegeneration in LRRK2 G2019S transgenic mice. *Proceedings of the National Academy of Sciences of the United States of America* **115**, 1635-1640 (2018).

17. T. K. Sang *et al.*, A Drosophila model of mutant human parkin-induced toxicity demonstrates selective loss of dopaminergic neurons and dependence on cellular dopamine. *J Neurosci* **27**, 981-992 (2007).

18. Z. Liu *et al.*, A Drosophila model for LRRK2-linked parkinsonism. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 2693-2698 (2008).

19. D. G. Ordonez, M. K. Lee, M. B. Feany, alpha-synuclein Induces Mitochondrial Dysfunction through Spectrin and the Actin Cytoskeleton. *Neuron* **97**, 108-124 e106 (2018).

20. A. J. Whitworth *et al.*, Increased glutathione S-transferase activity rescues dopaminergic neuron loss in a Drosophila model of Parkinson's disease. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 8024-8029 (2005).

21. N. M. Kanaan, J. H. Kordower, T. J. Collier, Age-related accumulation of Marinesco bodies and lipofuscin in rhesus monkey midbrain dopamine neurons:

relevance to selective neuronal vulnerability. *J Comp Neurol* **502**, 683-700 (2007).

22.   T. J. Collier *et al.*, Aging-related changes in the nigrostriatal dopamine system and the response to MPTP in nonhuman primates: diminished compensatory mechanisms as a prelude to parkinsonism. *Neurobiology of disease* **26**, 56-65 (2007).

23.   A. L. McCormack *et al.*, Aging of the nigrostriatal system in the squirrel monkey. *J Comp Neurol* **471**, 387-395 (2004).

24.   S. Noda, S. Sato, T. Fukuda, N. Tada, N. Hattori, Aging-related motor function and dopaminergic neuronal loss in C57BL/6 mice. *Mol Brain* **13**, 46 (2020).

25.   T. F. Mackay *et al.*, The Drosophila melanogaster Genetic Reference Panel. *Nature* **482**, 173-178 (2012).

26.   D. K. Ivanov *et al.*, Longevity GWAS Using the Drosophila Genetic Reference Panel. *J Gerontol A Biol Sci Med Sci* **70**, 1470-1478 (2015).

27.   M. F. Durham, M. M. Magwire, E. A. Stone, J. Leips, Genome-wide analysis in Drosophila reveals age-specific effects of SNPs on fitness traits. *Nature communications* **5**, 4338 (2014).

28.   W. Huang *et al.*, Context-dependent genetic architecture of Drosophila life span. *PLoS Biol* **18**, e3000645 (2020).

29.   M. A. Carbone *et al.*, Genetic architecture of natural variation in visual senescence in Drosophila. *Proceedings of the National Academy of Sciences of the United States of America* **113**, E6620-E6629 (2016).

30.   X. Zhao *et al.*, The metabolome as a biomarker of aging in Drosophila melanogaster. *Aging Cell* **21**, e13548 (2022).

31.   K. A. Wilson *et al.*, GWAS for Lifespan and Decline in Climbing Ability in Flies upon Dietary Restriction Reveal decima as a Mediator of Insulin-like Peptide Production. *Curr Biol* **30**, 2749-2760 e2743 (2020).

32.   M. Yang *et al.*, Using Drosophila to identify naturally occurring genetic modifiers of amyloid beta 42- and tau-induced toxicity. *G3 (Bethesda)* **13** (2023).

33.   S. Lavoy, V. G. Chittoor-Vinod, C. Y. Chow, I. Martin, Genetic Modifiers of Neurodegeneration in a Drosophila Model of Parkinson's Disease. *Genetics* **209**, 1345-1356 (2018).

34.   H. S. Wang *et al.*, In vivo evidence of pathogenicity of VPS35 mutations in the Drosophila. *Mol Brain* **7**, 73 (2014).

35.   I. Martin *et al.*, Ribosomal protein s15 phosphorylation mediates LRRK2 neurodegeneration in Parkinson's disease. *Cell* **157**, 472-485 (2014).

36.   J. Park *et al.*, Mitochondrial dysfunction in Drosophila PINK1 mutants is complemented by parkin. *Nature* **441**, 1157-1161 (2006).

37.   A. Chaudhuri *et al.*, Interaction of genetic and environmental factors in a Drosophila parkinsonism model. *J Neurosci* **27**, 2457-2467 (2007).

38.   H. Coulom, S. Birman, Chronic exposure to rotenone models sporadic Parkinson's disease in Drosophila melanogaster. *J Neurosci* **24**, 10993-10998 (2004).

39.   R. G. Pendleton, A. Rasheed, T. Sardina, T. Tully, R. Hillman, Effects of tyrosine hydroxylase mutants on locomotor activity in Drosophila: a study in functional genomics. *Behav Genet* **32**, 89-94 (2002).

40.   A. Vaccaro, A. R. Issa, L. Seugnet, S. Birman, A. Klarsfeld, Drosophila Clock Is Required in Brain Pacemaker Neurons to Prevent Premature Locomotor Aging Independently of Its Circadian Function. *PLoS Genet* **13**, e1006507 (2017).

41.   Y. Aso *et al.*, The neuronal architecture of the mushroom body provides a logic for associative learning. *Elife* **3**, e04577 (2014).

42.   D. Kretzschmar, G. Hasan, S. Sharma, M. Heisenberg, S. Benzer, The swiss cheese mutant causes glial hyperwrapping and brain degeneration in Drosophila. *J Neurosci* **17**, 7425-7432 (1997).

43.   G. A. Soultoukis, L. Partridge, Dietary Protein, Metabolism, and Aging. *Annu Rev Biochem* **85**, 5-34 (2016).

44.   R. C. Grandison, M. D. Piper, L. Partridge, Amino-acid imbalance explains extension of lifespan by dietary restriction in Drosophila. *Nature* **462**, 1061-1064 (2009).

45.   P. Kapahi *et al.*, Regulation of lifespan in Drosophila by modulation of genes in the TOR signaling pathway. *Curr Biol* **14**, 885-890 (2004).

46.   S. M. Solon-Biet *et al.*, The ratio of macronutrients, not caloric intake, dictates cardiometabolic health, aging, and longevity in ad libitum-fed mice. *Cell Metab* **19**, 418-430 (2014).

47.   S. K. Sandhu, G. Kaur, Alterations in oxidative stress scavenger system in aging rat brain and lymphocytes. *Biogerontology* **3**, 161-173 (2002).

48.   J. Ding *et al.*, A metabolome atlas of the aging mouse brain. *Nature communications* **12**, 6021 (2021).

49.   J. M. Luchak, L. Prabhudesai, R. S. Sohal, S. N. Radyuk, W. C. Orr, Modulating longevity in Drosophila by over- and underexpression of glutamate-cysteine ligase. *Ann N Y Acad Sci* **1119**, 260-273 (2007).

50.   W. C. Orr *et al.*, Overexpression of glutamate-cysteine ligase extends life span in Drosophila melanogaster. *The Journal of biological chemistry* **280**, 37331-37338 (2005).

51.   C. Brack, E. Bechter-Thuring, M. Labuhn, N-acetylcysteine slows down ageing and increases the life span of Drosophila melanogaster. *Cell Mol Life Sci* **53**, 960-966 (1997).

52.   P. Maher, The effects of stress and aging on glutathione metabolism. *Ageing Res Rev* **4**, 288-314 (2005).

53.   H. L. Martin, P. Teismann, Glutathione--a review on its role and significance in Parkinson's disease. *FASEB J* **23**, 3263-3272 (2009).

54.   D. H. Hyun *et al.*, Effect of wild-type or mutant Parkin on oxidative damage, nitric oxide, antioxidant defenses, and the proteasome. *The Journal of biological chemistry* **277**, 28572-28577 (2002).

55.   J. J. Palacino *et al.*, Mitochondrial dysfunction and oxidative damage in parkin-deficient mice. *The Journal of biological chemistry* **279**, 18614-18622 (2004).

56.   Y. Pesah *et al.*, Drosophila parkin mutants have decreased mass and cell size and increased sensitivity to oxygen radical stress. *Development* **131**, 2183-2194 (2004).

57. J. C. Greene, A. J. Whitworth, L. A. Andrews, T. J. Parker, L. J. Pallanck, Genetic and genomic studies of Drosophila parkin mutants implicate oxidative stress and innate immune responses in pathogenesis. *Human molecular genetics* **14**, 799-811 (2005).

58. J. Sian *et al.*, Glutathione-related enzymes in brain in Parkinson's disease. *Ann Neurol* **36**, 356-361 (1994).

59. J. Sian *et al.*, Alterations in glutathione levels in Parkinson's disease and other neurodegenerative disorders affecting basal ganglia. *Ann Neurol* **36**, 348-355 (1994).

60. R. K. Pearce, A. Owen, S. Daniel, P. Jenner, C. D. Marsden, Alterations in the distribution of glutathione in the substantia nigra in Parkinson's disease. *J Neural Transm (Vienna)* **104**, 661-677 (1997).

61. E. Sofic, K. W. Lange, K. Jellinger, P. Riederer, Reduced and oxidized glutathione in the substantia nigra of patients with Parkinson's disease. *Neurosci Lett* **142**, 128-130 (1992).

62. Y. Jang *et al.*, Mass Spectrometry-Based Proteomics Analysis of Human Substantia Nigra From Parkinson's Disease Patients Identifies Multiple Pathways Potentially Involved in the Disease. *Mol Cell Proteomics* **22**, 100452 (2023).

63. D. A. Loeffler, A. C. Klaver, M. P. Coffey, J. O. Aasly, P. A. LeWitt, Increased Oxidative Stress Markers in Cerebrospinal Fluid from Healthy Subjects with Parkinson's Disease-Associated LRRK2 Gene Mutations. *Front Aging Neurosci* **9**, 89 (2017).

64. H. N. Nguyen *et al.*, LRRK2 mutant iPSC-derived DA neurons demonstrate increased susceptibility to oxidative stress. *Cell stem cell* **8**, 267-280 (2011).

65. P. Reinhardt *et al.*, Genetic correction of a LRRK2 mutation in human iPSCs links parkinsonian neurodegeneration to ERK-dependent changes in gene expression. *Cell stem cell* **12**, 354-367 (2013).

66. L. K. Mischley *et al.*, Glutathione as a Biomarker in Parkinson's Disease: Associations with Aging and Disease Severity. *Oxid Med Cell Longev* **2016**, 9409363 (2016).

67. S. Charisis *et al.*, Plasma Glutathione and Prodromal Parkinson's Disease Probability. *Mov Disord* **37**, 200-205 (2022).

68. A. Menegon, P. G. Board, A. C. Blackburn, G. D. Mellick, D. G. Le Couteur, Parkinson's disease, pesticides, and glutathione transferase polymorphisms. *Lancet* **352**, 1344-1346 (1998).

69. Y. J. Li *et al.*, Revealing the role of glutathione S-transferase omega in age-at-onset of Alzheimer and Parkinson diseases. *Neurobiol Aging* **27**, 1087-1093 (2006).

70. C. Perez-Pastene *et al.*, Association of GST M1 null polymorphism with Parkinson's disease in a Chilean population with a strong Amerindian genetic component. *Neurosci Lett* **418**, 181-185 (2007).

71. M. R. Cookson, Parkinsonism due to mutations in PINK1, parkin, and DJ-1 and oxidative stress and mitochondrial pathways. *Cold Spring Harb Perspect Med* **2**, a009415 (2012).

72. J. M. Hoffman, Y. Lyu, S. D. Pletcher, D. E. L. Promislow, Proteomics and metabolomics in ageing research: from biomarkers to systems biology. *Essays Biochem* **61**, 379-388 (2017).

73. C. D. Wiley, J. Campisi, The metabolic roots of senescence: mechanisms and opportunities for intervention. *Nat Metab* **3**, 1290-1301 (2021).

74. T. Finkel, N. J. Holbrook, Oxidants, oxidative stress and the biology of ageing. *Nature* **408**, 239-247 (2000).

75. S. H. Benavides, A. J. Monserrat, S. Farina, E. A. Porta, Sequential histochemical studies of neuronal lipofuscin in human cerebral cortex from the first to the ninth decade of life. *Arch Gerontol Geriatr* **34**, 219-231 (2002).

76. R. S. Sohal, A. Agarwal, S. Agarwal, W. C. Orr, Simultaneous overexpression of copper- and zinc-containing superoxide dismutase and catalase retards age-related oxidative damage and increases metabolic potential in Drosophila melanogaster. *The Journal of biological chemistry* **270**, 15671-15674 (1995).

77. D. Toroser, W. C. Orr, R. S. Sohal, Carbonylation of mitochondrial proteins in Drosophila melanogaster during aging. *Biochemical and biophysical research communications* **363**, 418-424 (2007).

78. J. Jacobson *et al.*, Biomarkers of aging in Drosophila. *Aging Cell* **9**, 466-477 (2010).

79. R. S. Sohal, W. C. Orr, The redox stress hypothesis of aging. *Free Radic Biol Med* **52**, 539-555 (2012).

80. N. Sun, R. J. Youle, T. Finkel, The Mitochondrial Basis of Aging. *Mol Cell* **61**, 654-666 (2016).

81. M. J. Jackson, A. McArdle, Age-related changes in skeletal muscle reactive oxygen species generation and adaptive responses to reactive oxygen species. *J Physiol* **589**, 2139-2145 (2011).

82. V. I. Perez *et al.*, The overexpression of major antioxidant enzymes does not extend the lifespan of mice. *Aging Cell* **8**, 73-75 (2009).

83. W. C. Orr, R. S. Sohal, Extension of life-span by overexpression of superoxide dismutase and catalase in Drosophila melanogaster. *Science* **263**, 1128-1130 (1994).

84. S. E. Schriner *et al.*, Extension of murine life span by overexpression of catalase targeted to mitochondria. *Science* **308**, 1909-1911 (2005).

85. I. Martin, M. A. Jones, M. Grotewiel, Manipulation of Sod1 expression ubiquitously, but not in the nervous system or muscle, impacts age-related parameters in Drosophila. *FEBS Lett* **583**, 2308-2314 (2009).

86. I. Martin *et al.*, Sod2 knockdown in the musculature has whole-organism consequences in Drosophila. *Free Radic Biol Med* **47**, 803-813 (2009).

87. S. Elchuri *et al.*, CuZnSOD deficiency leads to persistent and widespread oxidative damage and hepatocarcinogenesis later in life. *Oncogene* **24**, 367-380 (2005).

88. M. T. Lin, M. F. Beal, Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* **443**, 787-795 (2006).

89. L. F. Burbulla *et al.*, Dopamine oxidation mediates mitochondrial and lysosomal dysfunction in Parkinson's disease. *Science* **357**, 1255-1261 (2017).

90. J. N. Guzman *et al.*, Oxidant stress evoked by pacemaking in dopaminergic neurons is attenuated by DJ-1. *Nature* **468**, 696-700 (2010).

91.   R. S. Balaban, S. Nemoto, T. Finkel, Mitochondria, oxidants, and aging. *Cell* **120**, 483-495 (2005).

92.   T. G. Hastings, The role of dopamine oxidation in mitochondrial dysfunction: implications for Parkinson's disease. *J Bioenerg Biomembr* **41**, 469-472 (2009).

93.   S. S. Park, E. M. Schulz, D. Lee, Disruption of dopamine homeostasis underlies selective neurodegeneration mediated by alpha-synuclein. *Eur J Neurosci* **26**, 3104-3112 (2007).

94.   F. Bayersdorfer, A. Voigt, S. Schneuwly, J. A. Botella, Dopamine-dependent neurodegeneration in Drosophila models of familial and sporadic Parkinson's disease. *Neurobiology of disease* **40**, 113-119 (2010).

95.   H. O. Lawal *et al.*, The Drosophila vesicular monoamine transporter reduces pesticide-induced loss of dopaminergic neurons. *Neurobiology of disease* **40**, 102-112 (2010).

96.   S. M. Goldman, Environmental toxins and Parkinson's disease. *Annu Rev Pharmacol Toxicol* **54**, 141-164 (2014).

97.   I. Martin, V. L. Dawson, T. M. Dawson, Recent advances in the genetics of Parkinson's disease. *Annu Rev Genomics Hum Genet* **12**, 301-325 (2011).

98.   G. H. Cha *et al.*, Parkin negatively regulates JNK pathway in the dopaminergic neurons of Drosophila. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 10345-10350 (2005).

99.   E. Dumitrescu, J. M. Copeland, B. J. Venton, Parkin Knockdown Modulates Dopamine Release in the Central Complex, but Not the Mushroom Body Heel, of Aging Drosophila. *ACS Chem Neurosci* **14**, 198-208 (2023).

100.   Y. Pesah *et al.*, Whole-mount analysis reveals normal numbers of dopaminergic neurons following misexpression of alpha-Synuclein in Drosophila. *Genesis* **41**, 154-159 (2005).

101.   M. Smeyne, R. J. Smeyne, Glutathione metabolism and Parkinson's disease. *Free Radic Biol Med* **62**, 13-25 (2013).

102.   R. Dringen, Metabolism and functions of glutathione in brain. *Prog Neurobiol* **62**, 649-671 (2000).

103. W. Feng *et al.*, Gclc deficiency in mouse CNS causes mitochondrial damage and neurodegeneration. *Human molecular genetics* **26**, 1376-1390 (2017).

104. S. J. Chinta *et al.*, Inducible alterations of glutathione levels in adult dopaminergic midbrain neurons result in nigrostriatal degeneration. *J Neurosci* **27**, 13997-14006 (2007).

105. M. Garrido *et al.*, Glutathione depletion and overproduction both initiate degeneration of nigral dopaminergic neurons. *Acta Neuropathol* **121**, 475-485 (2011).

106. E. Luna *et al.*, Differential alpha-synuclein expression contributes to selective vulnerability of hippocampal neuron subpopulations to fibril-induced toxicity. *Acta Neuropathol* **135**, 855-875 (2018).

107. A. Virel, I. Dudka, R. Laterveer, S. Af Bjerken, (1)H NMR profiling of the 6-OHDA parkinsonian rat brain reveals metabolic alterations and signs of recovery after N-acetylcysteine treatment. *Mol Cell Neurosci* **98**, 131-139 (2019).

108. A. Rahimmi, F. Khosrobakhsh, E. Izadpanah, M. R. Moloudi, K. Hassanzadeh, N-acetylcysteine prevents rotenone-induced Parkinson's disease in rat: An investigation into the interaction of parkin and Drp1 proteins. *Brain Res Bull* **113**, 34-40 (2015).

109. S. J. Chinta, J. M. Kumar, H. Zhang, H. J. Forman, J. K. Andersen, Up-regulation of gamma-glutamyl transpeptidase activity following glutathione depletion has a compensatory rather than an inhibitory effect on mitochondrial complex I activity: implications for Parkinson's disease. *Free Radic Biol Med* **40**, 1557-1563 (2006).

110. L. K. Mischley *et al.*, Central nervous system uptake of intranasal glutathione in Parkinson's disease. *NPJ Parkinsons Dis* **2**, 16002 (2016).

111. L. K. Mischley, R. C. Lau, E. G. Shankland, T. K. Wilbur, J. M. Padowski, Phase IIb Study of Intranasal Glutathione in Parkinson's Disease. *J Parkinsons Dis* **7**, 289-299 (2017).

112. M. J. Holmay *et al.*, N-Acetylcysteine boosts brain and blood glutathione in Gaucher and Parkinson diseases. *Clin Neuropharmacol* **36**, 103-106 (2013).

113. D. A. Monti *et al.*, N-Acetyl Cysteine Is Associated With Dopaminergic Improvement in Parkinson's Disease. *Clin Pharmacol Ther* **106**, 884-890 (2019).

114. W. Volkel *et al.*, Glutathione conjugates of 4-hydroxy-2(E)-nonenal as biomarkers of hepatic oxidative stress-induced lipid peroxidation in rats. *Free Radic Biol Med* **38**, 1526-1536 (2005).

115. D. Oppong, W. Schiff, M. C. Shivamadhu, Y. H. Ahn, Chemistry and biology of enzymes in protein glutathionylation. *Curr Opin Chem Biol* **75**, 102326 (2023).

116. K. Kurup *et al.*, Calorie restriction prevents age-related changes in the intestinal microbiota. *Aging (Albany NY)* **13**, 6298-6329 (2021).

117. J. P. Meador, L. F. Bettcher, M. C. Ellenberger, T. D. Senn, Metabolomic profiling for juvenile Chinook salmon exposed to contaminants of emerging concern. *Sci Total Environ* **747**, 141097 (2020).

118. M. E. Ritchie *et al.*, limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

119. G. K. Smyth, Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* **3**, Article3 (2004).

120. L. M. Beaver *et al.*, Circadian regulation of glutathione levels and biosynthesis in Drosophila melanogaster. *PLoS One* **7**, e50454 (2012).

121. J. A. Behnke, C. Ye, K. H. Moberg, J. Q. Zheng, A protocol to detect neurodegeneration in Drosophila melanogaster whole-brain mounts using advanced microscopy. *STAR Protoc* **2**, 100689 (2021).

122. M. A. Jones *et al.*, A forward genetic screen in Drosophila implicates insulin signaling in age-related locomotor impairment. *Exp Gerontol* **44**, 532-540 (2009).

SUPPLEMENTARY INFORMATION

Supplementary Figs S1-S6

Supplementary Tables S1 and S2

Data File S1

Figure S1. (A) Correlation of female mean life span for 27 DGRP strains (see Table S1 for strain IDs) used in this study (100 flies per genotype) with female life span of the same strains in (26). Pearson *r* and *p* annotated. (B-F), Correlation of mean life span (this study) in the 27 DGRP strains with TH-immunopositive neuron counts from the PAL (B), PPM1/2 (C), PPM3 (D), PPL2 (E) individual clusters or with counts from all clusters including PPL1 combined (F). Spearman *r* and *p* annotated. Dopamine neuron counts are from 7-12 brains/genotype.

Figure S2. (A) Deficits in climbing performance upon PPL1 dopamine neuron-specific silencing of the *ple* gene encoding TH. Two independent *ple* RNAi lines were crossed to each PPL1 neuron-specific GAL4 driver or to w^{1118} as control. Two-way ANOVA for effect of GAL4 driver (p<0.0001) and *ple* RNAi line (p<0.01), Dunnett's post-hoc test for *ple-IR-1*(#) or *ple-IR-2* (*) vs. ctrl (# p<0.05, ** or ## p<0.01, *** or ### p<0.001, n = 4 groups of 25 flies/genotype). (B-F) Correlation plots for negative geotaxis behavior and dopamine neuron counts within four separate clusters (B-E) or for all clusters (PAL, PPM1/2, PPM3, PPL1, PPL2) combined. DGRP strain IDs are provided in Table S1. (G) Correlation of negative geotaxis performance (n=4 groups of 25 flies/genotype) with mean life span across DGRP backgrounds. Spearman *r* and *p* annotated.

Figure S3. (A) Correlation matrix heat map for DGRP strain (RAL-727, RAL-911, RAL-356, RAL-335, RAL-379, RAL-821) PPL1 dopamine neuron counts, mean life span, brain vacuole area and vacuole number. Pearson r with *p* value in parentheses. (B) 5-HT neuron counts (SP1, SP2, IP and LP1 clusters) in 4-week-old female flies, ANOVA, ns (n=4-14 brains/genotype).

Figure S4. (A) PPL1 dopamine neuron counts of DGRP strains (RAL-727, RAL-911, RAL-356, RAL-335, RAL-379, RAL-821) outcrossed to w^{1118} relative to the parental inbred strain. Dopamine neurons for outcrossed strains were assessed at 6 weeks of age (B) Total glutathione measurements in female heads of DGRP strains indicated outcrossed to *elav-GAL4* (ctrl) or to *elav-GAL4; UAS-gclc*. There were significant effects of *gclc* overexpression and DGRP strain on glutathione levels (two-way ANOVA for effect of GCLc (p <0.0001) and DGRP strain (p <0.0001), Šídák's multiple comparison test, *p <0.05, n =4 groups of 25 heads/genotype).

A

 B

Figure S5. (A) correlation matrix heat map for the relationship between glutathione and mRNA levels of *gclc*, *gclm*, *Gss* and *Ggt-1* in 1-week-old female flies. Spearman *r* with *p* value in parentheses. (B) Correlation of glutathione levels with *gclc* mRNA levels in 1-week-old flies. mRNA levels are normalized to *Actin 5C*.

Figure S6. (A) *Parkin* RNAi verification by ubiquitous *A5c-GAL4*-driven expression of two independent *UAS-parkin-IR* lines (line ID in parenthesis), ANOVA for effect of RNAi, p <0.01, Dunnett's post-test, ** p <0.01, n = 3 groups of 15 heads/genotype. (B) No significant effect of dopamine neuron-specific (*TH-GAL4*-driven) *parkin* RNAi (line 37509) on PPL1 dopamine neuron counts (two-way ANOVA for effect of *parkin* RNAi (ns) and DGRP background (p <0.0001), n =10-13 brains/genotype.

Table S1. 27 DGRP strains (relates to data in Figures 1, S1 and S2)

Line ID	Mean LS	Std. error	Median LS	Std. error	Maximum
RAL-757	21	0.7	23	1.1	33
RAL-911	29	1.0	30	1.4	44
RAL-409	29	1.0	26	0.8	61
RAL-177	35	0.7	54	1.0	44
RAL-727	38	1.1	40	1.7	58
RAL-765	40	1.0	40	1.4	52
RAL-356	41	1.2	44	1.6	65
RAL-83	45	1.1	44	1.1	65
RAL-239	46	1.1	44	1.2	80
RAL-703	47	0.9	47	1.0	68
RAL-461	47	1.1	52	0.5	65
RAL-596	47	1.2	52	0.8	65
RAL-310	51	1.4	52	1.3	72
RAL-639	51	1.2	54	1.0	65
RAL-426	53	1.1	54	1.1	72
RAL-373	55	1.1	54	0.9	80
RAL-502	57	2.0	58	3.1	97
RAL-801	58	0.9	61	1.6	68
RAL-820	58	1.1	61	0.9	83
RAL-371	60	1.0	58	1.5	90
RAL-313	64	1.2	65	0.6	83
RAL-379	69	1.9	72	1.5	101
RAL-714	69	1.9	72	1.8	94
RAL-382	69	1.2	72	0.7	83
RAL-136	71	1.8	75	3.9	101
RAL-360	71	0.9	72	0.8	87
RAL-819	72	1.5	75	1.5	97
RAL-335	76	1.1	75	2.0	90
RAL-821	86	1.8	87	1.9	115

Table S2. Life span analyses for *gclc* overexpression

DGRP background	Genotype	Mean LS	Std. error	Log-rank *p* value
RAL-727	*elav-GAL4/+*	68	1.9	
RAL-727	*elav-GAL4/UAS-gclc*	77	1.3	<0.001
RAL-911	*elav-GAL4/+*	49	1.8	
RAL-911	*elav-GAL4/UAS-gclc*	61	1.4	<0.001
RAL-356	*elav-GAL4/+*	58	1.2	
RAL-356	*elav-GAL4/UAS-gclc*	64	1.4	<0.001
RAL-821	*elav-GAL4/+*	76	1.6	
RAL-821	*elav-GAL4/UAS-gclc*	75	1.6	0.257