Pathogen and pest risks to vegetatively propagated crops in humanitarian contexts: geographic priorities for Cameroon and Ethiopia

Romaric A. Mouafo-Tchinda[1,2,3]*, Berea A. Etherton[1,2,3]*, Aaron I. Plex Sulá[1,2,3], Jorge Andrade-Piedra[4], Kwame Ogero[5], Bonaventure A. Omondi[6], Margaret A. McEwan[7], Paul M. Tene Tayo[8], Dieudonné Harahagazwe[9], Mihiretu Cherinet[10], Setegn Gebeyehu[11], Louise Sperling[12], Karen A. Garrett[1,2,3]*

[1] Plant Pathology Department, University of Florida, Gainesville, FL, USA
[2] Global Food Systems Institute, University of Florida, Gainesville, FL, USA
[3] Emerging Pathogens Institute, University of Florida, Gainesville, FL, USA
[4] International Potato Center (CIP), Lima, Peru
[5] International Potato Center (CIP), Mwanza, Tanzania
[6] Alliance of Bioversity International and CIAT, Cotonou, Benin
[7] International Potato Center (CIP) Africa Regional Office, Nairobi, Kenya and Wageningen University and Research, Netherlands
[8] Biotechnology Center, University of Yaoundé, Yaoundé, Cameroon
[9] International Potato Center (CIP), Yaoundé, Cameroon
[10] International Potato Center (CIP), Hawassa, Ethiopia
[11] International Potato Center (CIP), Addis Ababa, Ethiopia
[12] SeedSystem, Sherman, CT, USA
* corresponding author(s)

R. A. Mouafo-Tchinda and B. A. Etherton made equivalent contributions

Emails, respectively: rmouafotchinda@ufl.edu*, betherton@ufl.edu*, plexaaron@ufl.edu, j.andrade@cgiar.org, k.ogero@cgiar.org, b.omondi@cgiar.org, margaret.mcewan08@gmail.com, martialtene@gmail.com, d.harahagazwe@cgiar.org, m.cherinet@cgiar.org, setegn.gebeyehu@cgiar.org, sperling@seedsystem.org, karengarrett@ufl.edu*

ORCID, respectively: 0000-0003-1343-4820, 0000-0002-4127-4484, 0000-0001-7317-3090, 0000-0001-7617-0506, 0000-0002-5141-6781, 0000-0001-9711-5645, 0000-0001-8510-0526, 0000-0002-8895-5615, 0000-0002-1337-1041, 0000-0002-5947-7175, 0000-0003-4610-1934, 0000-0002-2134-2638, 0000-0002-6578-1616

35 **Keywords:** humanitarian crises, expert knowledge elicitation, banana, cassava, potato
36 sweetpotato, network analysis

38 **Funding:** CGIAR Seed Equal Research Initiative, USAID Bureau for Humanitarian Assistance

40 Acknowledgements

We appreciate support from the USAID Bureau for Humanitarian Assistance and the CGIAR Seed

Equal Research Initiative; we thank all donors and organizations which globally support its work

through their contributions to the CGIAR Trust Fund. The opinions expressed in this article are

those of the authors and do not necessarily reflect the view of the USAID Bureau for

Humanitarian Assistance. We also appreciate the contributions of experts on systems in

Cameroon and Ethiopia who participated in the expert knowledge elicitation.

Abstract

Cameroon and Ethiopia face complex humanitarian crises that disrupt the production of key food security crops like banana and plantain, cassava, potato, and sweetpotato. These vegetatively propagated crops are prone to the buildup of pests and diseases causing seed degeneration. In this study, we address seed degeneration in the context of natural and human-driven crises, to provide insights that can guide the decisions of humanitarian groups to reduce seed degeneration. We identified locations in Cameroon and Ethiopia that are likely to be particularly important for pathogen and pest spread in cropland network analyses. We analyzed cropland networks in each country to develop a map of locations that are candidate priorities for disease and pest management and surveillance. For example, locations in central and western Oromia, Ethiopia, are likely important for potato pathogen and pest spread given the locations' roles in the cropland network. Next, we used expert knowledge elicitation to characterize crop health risk in these systems. For example, experts reported with high consensus that there are formal and informal imports of cassava from Gabon to Cameroon, which pose a risk to cassava production in Cameroon through seedborne pathogens. These results can inform regional and national surveillance and management strategies and provide a baseline for more in-depth studies in Cameroon and Ethiopia. The results can also inform decision-making in the humanitarian sector, for designing on-the-ground actions to avoid unintentional spread of pathogens and pests during agricultural recovery interventions.

1.0 Introduction

1.1 The importance of root, tuber, and banana crops for low-income countries and challenges for seed health

In the face of climate change and increasing natural and human-driven disasters, staple food crops like roots, tubers, and bananas (RTBs) are essential for food security (Scott, 2021; Thiele et al., 2017). RTBs are widely consumed, with a market value of $339 billion annually, providing nutrients for food-insecure populations in low-income countries (Petsakos et al., 2019; Scott, 2021). RTBs are vegetatively propagated, with planting materials that include banana (*Musa* spp.) suckers, cassava (*Manihot esculenta*) stakes, potato (*Solanum tuberosum*) tubers, and sweetpotato (*Ipomoea batatas*) vines. Vegetative propagation increases the risk of seed degeneration, or the build-up of pests and pathogens in seed, leading to subsequent infestation and infection (Gibson & Kreuze, 2015; Jacobsen et al., 2019; Thomas-Sharma et al., 2017; Thomas-Sharma et al., 2016). Seed degeneration can be exacerbated by informal trade if growers unknowingly spread pests and pathogens to other growers through casual or unregulated trade (Scott, 2021; Sperling & Almekinders, 2023; Sperling et al., 2020). Informal trade of RTB crops is the primary method for growers in low-income countries to acquire seed, providing little or no phytosanitary control (Sperling & McGuire, 2010). Formal seed trade can provide certified disease-free seed, but often at a high price, which can deter growers even when formal seed systems are available. As groups such as FAO and USAID emphasize improved seed aid, there is a knowledge gap for effective improvement for RTB crops. In this study, we focus on seed systems in Cameroon and Ethiopia, two countries which rely heavily on informal seed trade and on RTB crops for both income generation and food security (Hirpa et al., 2010; OCHA, 2023a). Prioritizing the resilience of RTB seed systems in these two countries can help to meet the challenges of natural and human-driven

disasters. Disasters can challenge RTB crop health if sufficient attention is not given to managing pest and

disease risk in humanitarian responses.

1.2 Ongoing disasters and humanitarian needs in Cameroon and Ethiopia

Disaster plant pathology (Etherton *et al.,* in review) addresses the key drivers of plant disease

associated with disasters, and the solutions available during these crises. Cameroon and Ethiopia face a

wide range of humanitarian challenges, prompting the analysis of these country's seed systems in the

context of disaster plant pathology. In Cameroon, humanitarian crises include natural disasters that affect

nearly every part of the country, such as flooding, deforestation, desertification/droughts, and landslides

(Bang, 2021; Bang & Balgah, 2022) **(Figure 1, I)**. Additionally, there are human-induced challenges

including armed conflicts like the Boko Haram insurgency in the Far North region, the separatist conflict in

the Northwest and Southwest regions, societal challenges related to the influx of refugees from the

Central African Republic in the East, Adamawa and North regions, and political instability across the nation

(Bang & Balgah, 2022; Mbah et al., 2023). The agricultural communities in these areas are affected by

these compounding crises, diminishing already limited resources. Growers are abandoning their fields due

to these natural and human-induced crises, with the risk of food shortages affecting their families (FAO,

2021a, 2021b; OCHA, 2023a). Cameroon hosts more than 450,000 refugees and asylum seekers every

year, and the demand for humanitarian aid continues to grow (Macrotrends, 2023; OCHA, 2022).

Humanitarian access, though, remains a major challenge due to administrative obstacles, insecurity

instilled by armed groups, and infrastructure damage (ECPHAO, 2023).

Humanitarian crises in Ethiopia are often associated with drought, armed conflict, pest outbreaks,

and political instability, of which drought is a main factor reducing agricultural production and increasing

malnutrition among the affected communities (Alasow et al., 2023; CDP, 2023; Wubaye et al., 2023)

(Figure 1, II). The frequency of droughts affecting cropland in Ethiopia has nearly doubled in the last 50

years (Alasow et al., 2023) and the severe drought that began in 2020 has continued into 2023, with

Ethiopia enduring five failed rainy seasons (Alasow et al., 2023; CDP, 2023; Wubaye et al., 2023). In

addition to drought, Ethiopia has experienced several armed conflicts, including the Tigray civil conflict,

and disputes between the Ethiopian military and the militias in Amhara, creating a dire situation for more

than five million people across these regions (CDP, 2023). The compounded effects of climate shocks,

armed conflicts, and related displacement, continue to create unprecedented humanitarian needs, with

more than 20 million people requiring food assistance in the year 2023 (OCHA, 2023b). In addition to a

loss of assets and planting material, a significant proportion of growers affected due to multiple disasters

cannot save seed, causing a staggering demand for seed supply as part of the emergency response. The

Figure 1. Maps of Cameroon (I) and Ethiopia (II) with representative types of disaster and humanitarian issues in the regions of the country. Each icon represents a type of disaster or major conflict affecting agricultural production, where red 'X's represent refugees issues and/or internally displaced people (IDP). These icons are placed approximately and do not reflect exact location or every reported disaster. Data adapted from Bang and Balgah (2022), IOM (2023), and Raleigh et al. (2010)

affected population may also need various types of emergency recovery support including fertilizer, farm

tools, and cash to resume agricultural activities.

1.3 The need for analysis of crop health risk to inform humanitarian aid interventions

The humanitarian imperative goes beyond saving lives and addressing immediate needs for food,

water, health care, and shelter. It aims to establish responses that foster livelihood recovery and enhance

resilience over both the short and medium term (Hunt et al., 2018; USAID, 2022). After natural disasters

occur, the relevance of agricultural interventions depends on several factors: the affected population,

displacement, seasonal timing, available resources, and security considerations (e.g., whether growers can

safely engage in cultivation amidst ongoing stressors). A key priority is ensuring the availability of disease-

free seed, through market interventions or direct distribution, and this should be, at a minimum, quality

seed of known low pathogen and pest levels (Nduwimana et al., 2022; Sperling et al., 2022).

There is often limited Information about the range of pathogens and pests that are a concern for plant

health, and their potential spread through planting materials and other means, even when studies have

addressed specific species in specific locations. For example, in Cameroon there have been targeted

studies of diseases and pests of RTB crops, such as weevils, nematodes, sigatoka, and banana bunchy top

disease on banana (Fogain et al., 1998; Ngatat et al., 2022; Okolle et al., 2009), cassava viruses, cassava

whitefly, cassava root rot, cassava green mite, cassava root mealybug, cassava anthracnose, and cassava

bacterial blight on cassava (Doungous et al., 2022; Poubom et al., 2005; Tize et al., 2021), and potato

tuber moth, aphids, potato viruses, leaf miners, bacterial wilt, and late blight on potato (Deloko et al.,

2021; Konje et al., 2019). In Ethiopia, studies have also been conducted on the incidence and spread of

some important diseases and pests of potato and sweetpotato, such as potato bacterial wilt (Tessema &

Seid, 2023), viral diseases of sweetpotato (Buko et al., 2020; Tesfaye et al., 2011) and sweetpotato weevil

(Beyene, 2015). However, we are not aware of publications evaluating the general risk of geographic

spread of these pathogens and pests in Cameroon and Ethiopia. Knowledge of key pathogens and pests is

important for ensuring both seed quality and availability of crop varieties that have needed resistance

genes, when resistance is available. When there is the risk of introducing a new pathogen or pest to a

country or region, zero tolerance for the presence of this new threat is important. However, often the

source and health status of potential seed are unknown and seed regulatory systems may not function to

provide assurance of seed quality. This highlights the importance of rapid risk assessment tools to identify

regions likely to be important in pathogen and pest spread, target economically consequential pests and

pathogens, and assess the role of informal trade in the spread of pathogens during human population

displacements (Carvajal-Yepes et al., 2019; Thomas-Sharma et al., 2017; Xing et al., 2020). This

information is important both for identifying risks for emergency seed aid and for longer-term

development strategies.

In this study, we use two complementary approaches to rapid risk assessment and mitigation for

decision-support of pathogen and pest management. These tools, when used before, during, or after

disasters, can help humanitarian groups prioritize seed aid allocation and mitigate the disease and pest

threats to agricultural communities. Cropland connectivity analysis provides a map of the likely role of

geographic locations in the spread of pathogens and pests based on the landscape of crop host availability

for crop-specific pathogens and pests (Margosian et al., 2009; Xing et al., 2020). Locations likely to act as

hubs or bridges for pathogen or pest movement are highlighted in the analysis, and these are candidate

priorities for management interventions and seed testing (Andersen et al., 2019; Andersen Onofre et al.,

2021; Buddenhagen et al., 2017; Buddenhagen et al., 2022). The second approach is expert knowledge

elicitation, which can provide a summary of crop experts' knowledge about the current RTB seed systems

and identify gaps in domain knowledge (Andersen Onofre et al., 2021; Thomas-Sharma et al., 2017).

Cropland connectivity analysis and expert knowledge elicitation can be used for rapid risk assessments

and to provide a baseline for more detailed follow-up studies as resources allow.

The objectives of this study are to 1) identify locations likely to be important for pathogen and pest

spread in potato and sweetpotato, given the locations' roles in each country's network of crop host

availability, 2) evaluate disease and pest risk in Cameroon and Ethiopia using expert knowledge elicitation,

and 3) identify potential risks to the spread of pathogens and pests through informal and formal seed

trade. We expand these analyses for Cameroon to include two more major food security crops: banana

and plantain, and cassava. We brought together a group of experts in both Cameroon and Ethiopia to

collect data regarding the RTB seed systems currently being used. We contextualize these analyses in the

humanitarian risk scenarios in Cameroon and Ethiopia that result from armed conflicts and extreme

weather events, to translate the results as input for decision-making by humanitarian organizations,

donors, and national programs to potentially shape regulatory policy as well as to guide specific practical

action on the ground.

2.0 Methods

2.1 Cropland connectivity for pathogens and pests

We used a cropland connectivity analysis to map locations likely to be important in outbreaks of

pathogens and pests of banana and plantain (or simply banana, henceforth), cassava, potato and

sweetpotato. Crop-specific harvested area across Sub-Saharan Africa (International Food Policy Research,

2020) for banana, cassava, potato and sweetpotato was used to evaluate a cropland connectivity risk

index (CCRI) as described by Xing et al. (2020), with code available in the geohabnet package in R (Plex

Sulá *et al.,* in review). The CCRI was evaluated separately for each crop, based on networks of potential

spread among hosts for crop-specific pathogens or pests. The CCRI was defined as the weighted sum of

four network metrics, to capture a range of potential roles a location may have during an epidemic or pest

invasion: 1/2 the node betweenness centrality to account for a location's potential role as a bridge for

spread between regions, 1/6 node strength to represent a location's potential importance as a hub for

spread, 1/6 the sum of a node's nearest neighbors' node degrees to account for the role of neighbors'

neighbors, and 1/6 eigenvector centrality to account for both quantity and quality of links in an invasion

network.

Cameroon and Ethiopia were the focus countries in these analyses. The cropland within each

country and in the immediate surrounding countries (±0.5° latitude and longitude) were included.

Locations with harvested areas greater than zero were retained as nodes, and a gravity model was used

to evaluate the relative likelihood of dispersal between each pair of nodes (Jongejans et al., 2015). The

dispersal kernel was based on an inverse power law model, $d_{ij}^{-\beta}$ in some components of a sensitivity

analysis, and in other components of the sensitivity analysis based on a negative exponential model,

$e^{-\gamma d_{ij}}$, where d_{ij} represents the distance between two nodes i and j, and the β and γ parameters

indicate the likelihood of pathogen movement across that distance (Andersen Onofre et al., 2021; Xing et

al., 2020). For example, the relative likelihood of pathogen movement between two nodes i and j for $\beta =$

1 is higher than for $\beta = 2$; similarly, movement for $\gamma = 1$ is more likely than for $\gamma = 2$. A range of β and

γ values were evaluated in the sensitivity analysis, as described in Xing et al. (2020), and the mean results

across these analyses are reported to provide a general connectivity measure for each location. We

generated maps of the CCRI using R (version 4.3.2) for each crop for both Cameroon and Ethiopia using

the raster package (version 3.5-15) (Hijmans, 2023). Higher CCRI scores indicate locations that are more

likely to play an important role in an epidemic or invasion based on host availability and pathogen or pest

dispersal, according to their geographic proximity and likely role in an epidemic or invasion network.

2.2 Expert knowledge elicitation

We developed a survey for use in expert knowledge elicitation for Cameroon and Ethiopia to capture the knowledge of RTB experts currently working in the field. Considering the limited database on seed systems in humanitarian situations, the goal was to build an interactive interface specifically designed for expert knowledge elicitation, in the broader context of the Rapid Risk assessment for Mitigation (R2M) toolbox (garrettlab.com/r2m/). Expert knowledge elicitation can be used to assess the risk of pest and pathogen spread through seed in humanitarian areas, and to help countries formulate rational management strategies against invasive crop pests and pathogens.

The first stage of survey development was to review the content of the country-specific versions of the survey with the organizing team to make sure it was appropriately matched to the situation in each country. The surveys focused on RTB pathogens and pests causing high yield loss, with potentially important impacts on vulnerable human populations in humanitarian situations. Expert knowledge elicitation was developed to capture expert knowledge in Cameroon, in four surveys covering banana, cassava, potato and sweetpotato. Similarly, two surveys were designed for experts in Ethiopia, focusing on potato and sweetpotato. Surveys were deployed using Google Forms (Vasantha Raju & Harinarayana, 2016). Experts from Cameroon and Ethiopia participated in a one-day workshop in November 2022 in Yaoundé, Cameroon, and in March 2023 in Addis Ababa, Ethiopia. In Cameroon, we recruited 18 experts for cassava, 13 for potato, 10 for banana and 7 for sweetpotato. In Ethiopia, we recruited 20 experts for potato, and 9 for sweetpotato. Experts known to have experience with these crops were selected from governmental organizations, the private sector, NGOs and CGIAR centers (IITA, CIP, and Alliance Biodiversity-CIAT). Experts were invited to complete the surveys individually, according to their crop expertise. In each survey, experts were asked to identify the major pests and diseases, characterize the crop-specific seed system, and identify the trade routes for seeds and harvested products within the country and with neighbouring countries. After experts completed responses individually, experts were

then asked to discuss their choices in small groups and come to a consensus, similarly to the Sheffield

protocol of expert knowledge elicitation (Gosling, 2018; O'Hagan, 2019).

Answers from experts were compiled and organized by administrative zone boundaries in each

country. For Cameroon, the regions are Far North, North, Adamawa, East, Centre, South, Littoral, West,

Northwest, and Southwest. For Ethiopia, the regional states are Tigray, Afar, Amhara, Benishangul-

Gumuz, Somali, Oromia, Gambela, Southwestern, Southern Nations, Addis Ababa, Dire Dawa and Harari.

An initial list of key pathogens and pests was drawn from previous studies carried out in both countries,

and based on preliminary discussions with the organizing team. The pathogens and pests identified by

experts to be present by administrative zones were compiled, and represented as "major pest and

pathogen communities" using the R packages ggplot2 (version 3.3.6) (Wickham, 2016) and leaflet (version

2.1.1) (Graul & Graul, 2016). Pathogen and pest communities reported by experts as present in a region

were visualized in mapped pie charts. For both countries, pest and pathogen community maps were

created for potato and sweetpotato, and for Cameroon also for cassava and banana. Note that a caveat

for comparison of locations is that locations with less crop production tend to have less attention by

experts, so lower expert experience in a region in a sense represents 'lower sampling effort'; that is,

regions and regional states where experts have less experience will tend to have lower numbers of

pathogen and pest species reported.

2.3 Within-country informal seed trade

Experts were asked about the informal seed trade of potato, sweetpotato, cassava, and banana

between regions of a country. For each potential link between a pair of regions, the link weight was the

proportion of experts who agreed that there was informal trade between that pair. This proportion was

used as a proxy for the level of informal seed trade between a pair of regions. The node strength, or the

sum of link weights, of informal seed trade was then calculated for each region using the igraph R package

(version 1.5.0) (Csárdi et al., 2023). These estimates of informal trade were compiled for each crop in both

countries and represented as networks using the ggraph R package (version 2.1.0) (Pedersen, 2022).

These informal seed trade networks between regions were then plotted on a map of each country with

political administrative boundaries, using the rgeoboundaries R package (version 1.2.9000) (Dicko, 2023).

2.4 Formal and informal trade in neighboring countries

Experts were asked about the informal trade of potato, sweetpotato, cassava and banana seed

with neighboring countries. Experts individually indicated which countries they thought imported seed

from and exported seeds to Cameroon or Ethiopia for each crop. The proportion of experts who thought

trade occurred between two countries was used as an "international trade rating". For example, if 9 of the

12 experts reported that informal exports of potato seed occur from Cameroon to Gabon, then this

directed link would have a weight of 0.75, indicating higher relative confidence in the occurrence of trade.

These ratings of informal imports and exports were collected and averaged for each crop-country

combination and represented as networks using the ggraph R package (version 2.1.0) (Pedersen, 2022).

Similarly, FAO import and export data were analyzed as formal trade networks between Cameroon,

Ethiopia, and their respective neighboring countries (FAOSTAT, 2021). Products manufactured by roasting

or sterilization were excluded in this trade data. FAO trade data were assembled for each crop, where the

trade volume (in metric tons) between each pair of countries was averaged across 2006 to 2020 (FAOSTAT,

2021). For simpler visual comparison with the levels of informal trade, the averaged formal trade volume

for each crop was rescaled between 0 and 1 to provide a "formal trade ratio," the proportion of formal

trade occurring relative to the reported informal trade occurring between Ethiopia, Cameroon, and their

respective neighboring countries. For example, if the maximum formal trade volume of potato for Ethiopia

was with Kenya, and was approximately 12 tons, then the trade volume for each Ethiopian potato trading

partner country was divided by 12 tons to rescale the volume.

3. Results

3.1 Cropland connectivity

Crop landscape connectivity was used to identify locations likely to play important roles in the spread of pests and pathogens based on host availability and general dispersal models. We evaluated cropland connectivity for potato and sweetpotato in Cameroon and Ethiopia, and also for banana and cassava in Cameroon.

In Cameroon, our analysis of crop landscape connectivity indicated that, for banana pathogens and pests, locations in the Southwest, West, Northwest, Centre, South, East, and Adamawa regions are likely most important to the risk of establishment and spread, based on host availability **(Figure 2, I)**. For cassava, many locations in the regions of Cameroon, except in the Southwest region, are at risk of establishment and spread of pathogens and pests given cassava cropland connectivity within country and their proximity to highly connected large cassava croplands in neighbouring countries like Nigeria and Chad **(Figure 2, I)**. For potato, locations in West, Nothwest, and Adamawa regions are the most exposed to the risk of establishment and spread of pathogens and pests **(Figure 2, I)**. These high-risk potato cropping areas are fairly small, reflecting the limited extent of potato production across Cameroon and their geographic isolation with more extensive potato production in neighbouring countries. For sweetpotato, locations in West, Nothwest, Adamawa, North and Far-Noth regions are likely the most important to the risk of establishment and spread of pathogens and pests **(Figure 2, I)**, based on host availability, with locations in the Northwest and West having the highest cropland connectivity across these regions.

For Ethiopian potato production, locations in a large expanse across central Ethiopia have higher cropland connectivity, spanning across Amhara, Oromia, Sidama and into the Southern Nations, Nationalities and Peoples (SNNP), with the highest CCRI scores occurring in the SNNP and central Amhara

Figure 2. Maps of cropland connectivity risk indices for I) banana, cassava, potato, and sweetpotato in Cameroon and II) potato and sweetpotato in Ethiopia. The cropland connectivity risk index is based on networks of potential spread across a landscape of crop hosts; locations with higher cropland connectivity are likely to be more important for pathogen and pest establishment and spread, based on host availability.

regions (**Figure 2, II**). For Ethiopian sweetpotato, key locations include western Oromia and Benishangul-

Gumuz, the border between SNNP and Sidama, and eastern Oromia including Harari and Dire Dawa

(**Figure 2, II**). The highest CCRI scores for Ethiopian sweetpotato were on the border between Sidama and

the SNNP. Neighbouring countries were not strongly connected to Ethiopia in potato and sweetpotato

cropland networks, as the reported harvested seed and production in neighbouring countries were

relatively low (**Figure 2, II**).

3.2 Expert knowledge elicitation: Pathogen and pest communities

For banana in Cameroon, the southern expanse was reported to have the highest major pathogen and pest communities, with the South zone having a community of at least eight major pests and pathogens **(Figure 3, I).** Anthracnose (caused by *Colletotrichum musae*) was reported present in every region across Cameroon except the North and Far North; burrowing nematodes and yellow sigatoka were reported present in every region except the North, Far North and Adamawa. Banana seed degeneration was reported to be associated with the burrowing nematode, BBTV, banana weevil, anthracnose (caused by *Colletotrichum musae*), and rhizome soft rot (caused by *Pectobacterium*).

For cassava, the East and Centre regions were reported to have the highest major pathogen and pest communities, with the East, Centre, and South regions reported to have the presence of all 13 pathogens which were included in the expert knowledge elicitation survey **(Figure 3, I).** Cassava green mite, cassava root rot, cassava mosaic disease, cassava mealybugs and cassava whitefly were reported as present in every region across Cameroon. Cassava seed degeneration was reported to be associated with CMV, cassava root rot (caused by *Fusarium* spp.), cassava anthracnose disease, ACMV, *Colletotrichum* spp., and cassava whitefly.

For potato, the western border was reported by experts to have the highest major pathogen and pest communities, with the Northwest and Southwest zones having pathogen and pest communities of at least 8 pathogens **(Figure 3, I).** Early blight (caused by *Alternaria solani*), PVY, potato aphids and Fusarium dry rot were reported to be present in seven of the ten regions across Cameroon, having the highest reported presence by experts. The smallest major pathogen and pest communities were reported for potato producers in Cameroon. Potato seed degeneration was reported to be associated with Fusarium dry rot, late blight (caused by *Phytophthora infestans*), PVY, PLRV, and potato cyst nematodes.

For sweetpotato, the West region was reported to have the largest major pathogen and pest

communities, with the South zone having the highest with six reported pathogens **(Figure 3, I).** The

Northwest, West, Centre, and Littoral zones each had reported pathogen and pest communities of at least

five pathogens. Early blight (caused by *A. solani*) and bacterial wilt (caused by *Ralstonia solanacearum*)

were both reported as present in every region across Cameroon except the North and the Southwest.

Sweetpotato seed degeneration was reported to be associated with sweetpotato viruses, Alternaria blight,

Fusarium root rot, and bacterial wilt.

For Ethiopian potato, central zones were reported in expert knowledge elicitation to have the

highest major pest and pathogen communities, with Oromia having the highest community of at least

twelve reported pathogens and pests **(Figure 3, II).** The SNNP, Southwestern, Amhara and Tigray regions

each had reported pathogen communities of at least ten pathogens. Experts reported that PLRV was

present in every region across Ethiopia. PVY and potato aphids were reported as present in every region

except Afar and Gambela, respectively. Experts selected the key causes of seed degeneration in Ethiopia

to be *Phytophthora infestans*, PVY, PLRV, potato aphids, early blight, and bacterial wilt in potato.

For Ethiopian sweetpotato, southern zones were reported by experts to have the highest pest and

pathogen communities, with Sidama and the SNNP both reporting pathogen communities of at least 8

pathogens **(Figure 3, II)**. Alternaria blight and the sweetpotato weevil were reported in every region across

Ethiopia, except for Somali and Addis Ababa, respectively. Experts also reported that some of these

pathogens and pests caused seed degeneration in both crops in Ethiopia. Experts selected the key causes

of seed degeneration as sweetpotato viruses, sweetpotato weevil, *Alternaria bataticola*, and sweetpotato

aphids.

Figure 3. Maps of the reported presence of major diseases and pests in (I) Cameroon and (II) Ethiopia, from expert knowledge elicitation. Pie charts indicate whether a disease or pest was reported to be present in each region. Note that the absence of a report in the expert knowledge elicitation may be influenced by lower research efforts in regions with less crop production.

     3.3 Expert knowledge elicitation: Within-country informal seed trade

We analyzed the network of seed exchange between regions in Cameroon and in Ethiopia, as

reported by experts for potato and sweetpotato, to assess the risk of pathogen and pest spread through

seed between regions. Experts reported high levels of exchange, suggesting a high risk for seedborne

pathogen and pest spread between regions in both countries. For informal potato and sweetpotato seed

trade in Cameroon, all the regions were reported to be interconnected. There was high seed trade

strength reported in the West region for potato **(Figure 4, I)** and in the Centre and Littoral regions for

sweetpotato **(Figure 4, I).** In Ethiopia, all the regions were also reported to be interconnected except the

Figure 4. Informal seed trade networks between regions in I) Cameroon, and II) Ethiopia as reported in expert knowledge elicitation. Node size is proportional to node strength, or the sum of link weights. Link weights are the proportion of experts that agreed on the occurrence of informal trade between the linked regions.

Somali region. Seed trade strength was reported to be high in the Oromia and Amhara regions for potato

(Figure 4, II) and in the Southwestern, SNNP, and Oromia regions for sweetpotato (Figure 4, II).

For banana and cassava in Cameroon, regions were reported to be well linked, with high regional

informal seed trade strength for banana in the Southwest, Littoral, East, West and South regions (Figure 4,

I) and for cassava in the South, Centre, Littoral, West and East regions (Figure 4, I). The experts also

reported that, due to the current crisis in the Northwest and Southwest regions of Cameroon and the

conflict with Boko Haram in the Far-North region, growers have moved to other regions and neighboring

countries with their goods, including seed and food. They also highlighted the vulnerable situation of

many of these growers working in neighboring regions and countries.

3.4 Formal and informal trade with neighboring countries

We used expert knowledge elicitation and FAO trade data to characterize the informal and formal

trade networks, respectively, linking Cameroon and Ethiopia with their respective neighboring countries.

These networks were generated for potato and sweetpotato for both Cameroon and Ethiopia, and also for

banana and cassava for Cameroon.

For informal potato trade in Cameroon, 38% of experts reported informal import of potato seed

from Nigeria to Cameroon, and 30% reported informal export of potato seed from Cameroon to Gabon

(Figure 5, I). Informal export of potato seed to all neighboring countries from Cameroon was reported by

at least one expert, and no experts reported that the Congo Republic was importing potato seed into

Cameroon. For formal trade, exports and imports were reported between the Congo Republic, the Central

African Republic and Gabon, with the highest trade volume average from Gabon imports.

For informal sweetpotato imports, 28% of experts reported that Gabon, Nigeria, and the Central

African Republic imported sweetpotato seed to Cameroon, and 28% of experts reported that Cameroon

exported sweetpotato seed to Gabon and the Central African Republic (Figure 5, I). No informal imports to

Cameroon were reported to occur from Chad, Equatorial Guinea, the Congo Republic and Chad, and no

informal exports were reported to occur to Equatorial Guinea. For the formal trade, the highest trade

volume occurred from Gabon imports to Cameroon. Formal bidirectional trade occurred between

Cameroon and Gabon, the Central African Republic, and the Congo Republic, and formal exports occurred

to Equatorial Guinea. No informal or formal trade was reported to occur with Chad **(Figure 5, I)**.

For informal trade of banana seed in Cameroon, 40% of experts reported exports to Gabon, and

30% reported imports from Gabon and exports to Nigeria **(Figure 5, I)**. No informal imports were reported

from Equatorial Guinea or Chad. The largest formal trade was exports to Chad, and secondly to Eritrea. No

formal imports were reported from the Congo Republic, Chad, or Equatorial Guinea, and no formal

exports occurred to the Central African Republic **(Figure 5, I)**.

For informal cassava trade in Cameroon, 72% of experts reported exports to Gabon, the highest

reports across all surveys, and 50% reported imports from Nigeria **(Figure 5, II)**. Informal imports and

exports were reported by experts for all neighboring countries except for informal imports from Chad

(Figure 5, I). For formal trade of cassava, FAO reported the highest trade volume occurring through Gabon

imports. Formal imports were from Congo and Gabon, and formal exports were to the Central African

Republic, Gabon, and the Congo Republic **(Figure 5, I)**.

For the informal Ethiopian potato seed trade, 30% of experts reported potato seed export to

Djibouti, and 25% reported exports to Somalia and Sudan **(Figure 5, II)**. No experts reported informal

imports from Somalia, Eritrea, and South Sudan. For formal trade, the highest trade volume was Ethiopian

exports to Somalia. No formal imports or exports occurred between South Sudan and Eritrea, and no

formal imports occurred from Sudan, Somalia, or Kenya.

Figure 5. Formal and informal trade networks with neighboring countries for I) Cameroon and II) Ethiopia. Informal networks were reported in expert knowledge elicitation, and formal networks were based on FAO trade databases. Link width represents the relative confidence in the occurrence of trade (based on the proportion of experts reporting it), and color represents the trade type reported (formal or informal).

406   Informal trade of sweetpotato seed was reportedly rare from Ethiopia, where no exports of

407   sweetpotato seed were reported **(Figure 5, II)**. Unlike exports, informal imports were reported from all

408   neighboring countries except Eritrea, with only 11% of experts reporting these imports. Formal seed

409   imports and exports were only reported to occur between Ethiopia and Djibouti.

410   4.0 Discussion

These analyses show the potential for crop pathogens and pests to spread quickly in Cameroon

and Ethiopia, especially across the highly-connected regions identified here, potentially moving

undetected through seed trade. Many actors may contribute to this unwanted movement: growers

sharing seed, traders selling seed, and humanitarian groups and governments providing emergency or

developmental assistance. All of these actors need to be aware of the potential for pathogen and pest

spread, and to be equipped with tools to prevent unintentional spread, especially spread of new species

or problematic populations, such as pathogen populations with pesticide tolerance or the ability to

overcome crop resistance genes. Information about which geographic locations are likely to be particularly

important in pathogen and pest spread is an important first step in developing priorities and strategies for

investing limited resources to protect crop health. Seed testing and charting specific sources of clean

planting materials are the next line of defense and are essential for humanitarian interventions.

The spread of crop pathogens and pests is driven by a complex interplay of factors including

environmental conduciveness, the formal and informal movement of agricultural commodities, as well as

the density and connectivity of crops (Andersen Onofre et al., 2021; Buddenhagen et al., 2022; Garrett et

al., 2022; Margosian et al., 2009; Nduwimana et al., 2022; Xing et al., 2020). Complete information about

these factors is rarely available when decisions must be made about the emergency movement of food

and planting material in humanitarian crises, but the analyses we illustrate here can be performed

relatively quickly to provide baseline information about pathogen and pest risk. Geographic pathogen and

pest risk assessments (Andersen Onofre et al., 2021; Thomas-Sharma et al., 2017) are a key component of

agricultural resilience and food security in countries affected by natural or human-driven disasters

(Etherton et al. in review). These analyses help to identify the most vulnerable cropping regions and

regions likely important for disease establishment or spread. Analyses such as the ones reported here, and

the next generations of such analyses as tools are continually improved, should be routinely conducted in

crisis regions to help humanitarian organizations, donors, and governments assess the risks associated

with diseases and pests. This information is an important component of input to establish effective and

appropriate humanitarian responses that minimize risks, and ensure that affected populations have access

to sufficient nutritious food during and following crises. In this study, we assessed the key species and key

locations important for pathogen and pest spread in banana, cassava, potato, and sweetpotato for

Cameroon and potato and sweetpotato for Ethiopia. This information is intended as input to support

decision-making to form aid policies in response to crises and to prioritize considerations in humanitarian

implementation on the ground, and for those along the informal trade chains, such as growers, traders,

and producers.

The cropland connectivity analyses identified regions likely to be important for the establishment

and spread of crop-specific pathogens and pests, based on crop host availability. We found that locations

in the West, Northwest, and Adamawa regions of Cameroon presented the highest risk of pathogen and

pest spread for all banana, cassava, potato and sweetpotato crops, as did locations in the Oromia, Amhara

and SNNP regions of Ethiopia for potato and sweetpotato crops. For Cameroon, informal movement of

cassava and sweetpotato, should also be monitored, particularly to and from Nigeria, as Nigeria has high

CCRI scores for these crops as well and may be a refuge for immigrants during humanitarian crises. For

Ethiopia, neighboring countries had low CCRI scores, suggesting migrants from Ethiopia bringing their

seed pose a lower threat to the immediate surrounding cropping regions. Other examples of cropland

connectivity analyses are the identification of locations likely to be important for the spread of potato

pathogens and pests in Kenya and Ecuador (Buddenhagen et al., 2022) and the Republic of Georgia

(Andersen Onofre et al., 2021). Cropland connectivity analysis could be used before, during, and after

crises to identify candidate priority locations where investment efforts are likely to have a greater impact

on slowing pathogen or pest spread. Maps of connectivity (CCRI) show surrounding regions that may also

play an important role in epidemics, which may be priorities for providing clean seed to fleeing

populations who are planting in new locations, or for seed testing to mitigate disease establishment. For

example, in 2021 inter-community conflicts, linked to the scarcity of water and other resources, broke out

in Cameroon between populations in the Far North, resulting in the displacement of over 70,000 people,

35,000 of whom sought refuge in Chad. Growers fleeing with their goods (including cassava and

sweetpotato seed), based on cropland connectivity analyses, have the potential to introduce seedborne

diseases in cropland in Chad (ECPHAO, 2023; FAO, 2021b). Conversely, refugees fleeing into Cameroon

and Ethiopia may bring with them infected seed; for example, the Boko Haram conflict in northeast

Nigeria caused 138,000 refugees to flee to Far North Cameroon (ECPHAO, 2023).

The expert knowledge elicitation identified the major pathogens and pests reported by experts in

each region for both Cameroon and Ethiopia. We noted differences in the major pathogens and pests

between regions for the same crop, for example, experts reported few pests and pathogens in East

Cameroon for banana, potato, and sweetpotato, but every pest and pathogen evaluated for cassava.

Some pests and pathogens, for example, white grub in Cameroon, were only reported in Centre and South

Cameroon, and nowhere else in the country. The results of expert knowledge elicitation provide a basis

for rapid assessment in field studies. Expert knowledge elicitation has been used in other scenarios to

evaluate phytosanitary risk, for example by Andersen Onofre et al. (2021) to identify phytosanitary

strategies for protecting potato seed systems in the Republic of Georgia. Savary et al. (2019) used an

expert survey to estimate the yield losses due to pathogens and pests for five major food security crops.

This new study illustrates how expert knowledge elicitation can be used in crisis situations to identify the

main pathogens and pests that could have a major impact on food security and the incomes of vulnerable

populations. It can also be used to visualize the formal or informal trade network for seeds, harvested

products, and agricultural equipment in a country, as well as the displacement of populations during

crises, and analyze the potential risk of pathogen and pests spread between regions.

Several of the pests and pathogens identified in the expert knowledge elicitation contribute to

seed degeneration, which is a major concern in the cultivation of RTB crops in both countries. Pathogens

and pests accumulate over cycles of seed saving by growers, leading to a progressive reduction in crop

yield and quality (Navarrete et al., 2022; Thomas-Sharma et al., 2017). This problem is exacerbated in

Cameroon and Ethiopia since significant numbers of refugees and displaced people have been integrated

into local communities, adding a further challenge to accessing clean seed as well as food, itself. There is

often a lack of networks for accessing formal services and resources (ECPHAO, 2023; TGHAT, 2022). In

Cameroon, for example, 90% of the agricultural market is informal (OCHA, 2023a). Analysis of the informal

trade network reported in expert knowledge elicitation indicates that regions in both countries are well

connected, with a corresponding risk of pathogen and pest spread. Informal seed trade between regions

often leads to the introduction of new pathogens and pests to a region, which can reduce crop production

and the food security of vulnerable populations (Andersen Onofre et al., 2019; Delaquis et al., 2018;

Nduwimana et al., 2022). We identified key hubs for the potential spread of pathogens and pests in

Cameroon in the Southwest and Littoral (banana), the South, Centre, and Littoral (cassava), the Northwest

and West (potato), and the Centre and Littoral (sweetpotato). In Ethiopia, locations in the Oromia and

Amhara regions were potential key locations for potato, while locations in the Southwestern, Oromia and

SNNP regions were the potential key locations for pest and pathogen spread in sweetpotato.

The ongoing insecurity in the Northwest, Southwest, East, and Far North regions of Cameroon is

increasing displacement of people to other regions with their seed, harvested crops, and agricultural

equipment, often without knowledge of the basic phytosanitary rules that could be used during this

migration. These crises are likely to continue to increase the risk of pathogen and pest spread across

regions, leading to greater food insecurity and vulnerability for the country's population. In Ethiopia, IOM

(2023) reported that ethnic and/or territorial conflicts and drought had a huge impact on agricultural

production, causing the internal displacement of over 4.8 million people in 2022 and 2023. An assessment

of agricultural performance conducted during the main season of 2022 reported that about 1.2 million

households in four regional states of Ethiopia needed emergency seed and other input support to resume

agricultural activity in the 2023 cropping seasons (RBA, 2022). Foyou et al. (2018) also reported that in

Cameroon, the Boko Haram-induced crisis in the Far North region increased the vulnerability of the

agricultural and economic system of the region and caused the internal and external displacement of

millions of people.

Experts in this study reported that growers in both countries participate in informal trade in seed

and harvested products with growers in neighboring countries. People living in border villages often have

family members in neighboring countries and share the same local markets. Both informal and formal

exchange of agricultural commodities increases the risk of seedborne pathogen and pest spread between

Cameroon/Ethiopia and neighboring countries. For example, Etherton et al. (in review) concluded that the

risk of reintroducing the *Ralstonia solanacearum* species complex (RSSC) to Ethiopia from Kenya through

formal trade was an important consideration, as was the risk of spreading the RSSC from Ethiopia to other

neighboring countries such as South Sudan, Sudan, Djibouti, and Somalia. Experts also pointed out that,

with multiple conflicts and crises (since 2013) in Cameroon and neighboring countries (Nigeria and the

Central African Republic), populations tend to move to neighboring countries during periods of intense

conflict, return to their villages during periods of calm or harvests, and flee again as soon as conflict

intensity increases. ECPHAO (2023) reports that the ongoing separatist conflict in the Northwest and

Southwest regions has displaced 80,000 Cameroonians to neighboring villages in Nigeria, while violence in

the Central African Republic and conflict with Boko Haram in northern Nigeria led to a massive influx of

over 491,000 refugees into neighboring Cameroonian villages (ECPHAO, 2023).

The analyses here represent baselines for better understanding pathogen and pest risks in RTB

systems in Cameroon and Ethiopia. When resources are available, there are many possibilities for

improving on these analyses. For example, the cropland connectivity analyses represent a simplified view

of the risk of spread by averaging across potential dispersal parameters, to represent many potential

pathogens and pests. When dispersal is well-characterized for a particular pathogen or pest species, the

cropland connectivity analysis can be implemented specifically for that species, and potentially expanded

to a fuller habitat connectivity analysis, including other location traits such as environmental

conduciveness to the species. The cropland connectivity analysis may also be improved as crop density

maps improve in quality. Cropland connectivity analyses can also be a first step for scenario analyses for

regional pest and disease management, by providing an estimate of the potential epidemic or invasion

network for simulation experiments (Etherton et al., 2023; Garrett, 2021; Wiebe et al., 2018).

The results of expert knowledge elicitation can also be improved over time. Experts are limited by

potential biases and lack of knowledge. For example, experts may tend to have less knowledge about

locations where a particular crop is less commonly produced, so that there is less 'sampling effort' in

those locations in terms of expert knowledge. When resources are available for pathogen and pest

monitoring, new information can supplement summaries of expert knowledge. Expert knowledge

elicitation and habitat connectivity analysis are part of the developing R2M toolbox for rapid risk

assessment to support mitigation of plant pathogens and pests (garrettlab.com/r2m), designed for

continual improvement based on lessons learned in applications across countries and crop systems.

Related tools focusing on seed systems include the seed emergency response tool (Sperling et al., 2022),

the multi-stakeholder framework for RTB seed systems (Bentley et al., 2018), and other components of

the CGIAR tools4seedsystems toolbox (Andrade-Piedra et al., 2023). These tools can be used in

conjunction and incorporated into comprehensive risk management frameworks.

During times of crisis, first-order humanitarian aid should clearly be allocated to providing affected

people with basic needs such as food and water. A priority shortly after is to spur agricultural production

for growers' own food production and sale. For RTB crops, attention should focus on seed verification and

providing clean seed to growers, including growers migrating to neighboring regions or countries. Risk

analysis can be conducted prior to conflict, to create priority maps for aid. For example, providing clean

cassava seed to growers in Centre, East, and South Cameroon is a candidate priority, as these regions were

identified by experts as having many major cassava pathogens and pests. Based on cropland connectivity,

candidate priorities for clean potato seed provision in Ethiopia are locations in Amhara and central

Oromia. Following a crisis, understanding the informal and formal trade of seed can help to identify

management intervention points, for example, markets in Southwestern Ethiopia are a candidate priority

for surveying for sweetpotato diseases to stop the movement of seedborne pathogens, and similarly for

potato markets in West Cameroon. Aid, whether governmental or humanitarian, is a crucial part of

agricultural sustainability and should focus on the development of sustainable long-term strategies,

involving governmental bodies to develop more formal seed testing programs nationally. It is through the

collaboration of humanitarian groups, governmental programs, growers, and traders, themselves, that

sustainable agriculture can be achieved in the face of humanitarian disasters.

565 Statements and Declarations

566 The authors declare that they have no conflict of interest.

567 Sources

568 Alasow, A. A., Hamed, M. M., & Shahid, S. (2023). Spatiotemporal variability of drought and affected
569  croplands in the horn of Africa. *Stochastic Environmental Research and Risk Assessment*, 1-16.
570  https://doi.org/10.1007/s00477-023-02575-1
571 Andersen, K. F., Buddenhagen, C. E., Rachkara, P., Gibson, R., Kalule, S., Phillips, D., & Garrett, K. A.
572  (2019). Modeling epidemics in seed systems and landscapes to guide management strategies:
573  The case of sweet potato in northern Uganda. *Phytopathology*, *109*(9), 1519-1532.
574  https://doi.org/10.1094/PHYTO-03-18-0072-R
575 Andersen Onofre, K. F., Forbes, G. A., Andrade-Piedra, J. L., Buddenhagen, C. E., Fulton, J. C., Gatto, M.,
576  Khidesheli, Z., Mdivani, R., Xing, Y., & Garrett, K. A. (2021). An integrated seed health strategy
577  and phytosanitary risk assessment: Potato in the Republic of Georgia. *Agricultural Systems*, *191*,
578  103144. https://doi.org/10.1016/j.agsy.2021.103144
579 Andrade-Piedra, J., Dontsop, P., Dunia, D., Kulakow, P., Harahagazwe, D., Mouafo-Tchinda, R., Omondi, A.,
580  Ogero, K., Rajendran, S., & McEwan, M. (2023). Tools4SeedSystems: Working towards resilience
581  through root, tuber and banana crops in humanitarian settings. Capacity needs assessment for
582  root, tuber and banana seed interventions in humanitarian settings: Cameroon and DRC.
583  Technical Report. https://doi.org/10.4160/9789290606635
584 Bang, H. N. (2021). A gap analysis of the legislative, policy, institutional and crises management
585  frameworks for disaster risk management in Cameroon. *Progress in Disaster Science*, *11*, 100190.
586  https://doi.org/10.1016/j.pdisas.2021.100190
587 Bang, H. N., & Balgah, R. A. (2022). The ramification of Cameroon's Anglophone crisis: conceptual
588  analysis of a looming "Complex Disaster Emergency". *Journal of International Humanitarian*
589  *Action*, *7*(1), 6.
590 Bentley, J. W., Andrade-Piedra, J., Demo, P., Dzomeku, B., Jacobsen, K., Kikulwe, E., Kromann, P., Kumar, P.
591  L., McEwan, M., & Mudege, N. (2018). Understanding root, tuber, and banana seed systems and
592  coordination breakdown: a multi-stakeholder framework. *Journal of Crop Improvement*, *32*(5),
593  599-621. https://doi.org/10.1080/15427528.2018.1476998

Beyene, K. (2015). Destitution, biology, yield loss and management of sweet potato weevils (Cylas formicaries (Fabrcius) Insecta: Coleoptera) in Ethiopia. *Journal of Biology, Agriculture and Healthcare*, *5*(22), 65-72.

Buddenhagen, C. E., Hernandez Nopsa, J. F., Andersen, K. F., Andrade-Piedra, J., Forbes, G. A., Kromann, P., Thomas-Sharma, S., Useche, P., & Garrett, K. A. (2017). Epidemic network analysis for mitigation of invasive pathogens in seed systems: Potato in Ecuador. *Phytopathology*, *107*(10), 1209-1218. https://doi.org/10.1094/PHYTO-03-17-0108-FI

Buddenhagen, C. E., Xing, Y., Andrade-Piedra, J. L., Forbes, G. A., Kromann, P., Navarrete, I., Thomas-Sharma, S., Choudhury, R. A., Andersen Onofre, K. F., Schulte-Geldermann, E., Etherton, B. A., Plex Sula, A. I., & Garrett, K. A. (2022). Where to invest project efforts for greater benefit: A framework for management performance mapping with examples for potato seed health. *Phytopathology*, *112*(7), 1431-1443. https://doi.org/10.1094/PHYTO-05-20-0202-R

Buko, D. H., Gedebo, A., Spetz, C., & Hvoslef-Eide, A. K. (2020). An update of sweet potato viral disease incidence and spread in Ethiopia. *African Journal of Agricultural Research*, *16(8)*, 1116-1126. https://doi.org/10.5897/AJAR2020.14918

Carvajal-Yepes, M., Cardwell, K., Nelson, A., Garrett, K. A., Giovani, B., Saunders, D., Kamoun, S., Legg, J., Verdier, V., & Lessel, J. (2019). A global surveillance system for crop diseases. *Science*, *364*(6447), 1237-1239.

CDP. (2023). *Ethiopia humanitarian crisis*. https://disasterphilanthropy.org/disasters/ethiopia-tigray-crisis/

Csárdi, G., Nepusz, T., Traag, V., Horvát, S., Zanini, F., Noom, D., & Müller, K. (2023). igraph: Network Analysis and Visualization in R. *R package version 1.5.1*. https://doi.org/10.5281/zenodo.7682609

Delaquis, E., Andersen, K. F., Minato, N., Cu, T. T. L., Karssenberg, M. E., Sok, S., Wyckhuys, K. A., Newby, J. C., Burra, D. D., & Srean, P. (2018). Raising the stakes: cassava seed networks at multiple scales in Cambodia and Vietnam. *Frontiers in Sustainable Food Systems*, *2*, 73. https://doi.org/10.3389/fsufs.2018.00073

Deloko, D. C. T., Achiangia, N. P., Chofong, N. G., Mbulli, A. I., Anoumaa, M., Sama, L. F., & Fonkou, T. (2021). Prevalence of potato viruses on potato (Solanum tuberosum L.) grown in the Western Highlands of Cameroon. *Journal of Agriculture and Food Research*, *5*, 100192. https://doi.org/10.1016/j.jafr.2021.100192

Dicko, A. (2023). rgeoboundaries: A Client to geoBoundaries, A Political Administrative Boundaries Dataset. https://github.com/wmgeolab/rgeoboundaries

Doungous, O., Masky, B., Levai, D. L., Bahoya, J. A., Minyaka, E., Mavoungou, J. F., Mutuku, J. M., & Pita, J. S. (2022). Cassava mosaic disease and its whitefly vector in Cameroon: Incidence, severity and whitefly numbers from field surveys. *Crop protection*, *158*, 106017. https://doi.org/10.1016/j.cropro.2022.106017

ECPHAO. (2023). *Cameroon factsheet: European Civil Protection and Humanitarian Aid Operations*. https://civil-protection-humanitarian-aid.ec.europa.eu/where/africa/cameroon_en

Etherton, B. A., Choudhury, R., Alcalá-Briseño, R., Xing, Y., Sulá, A. P., Carrillo, D., Wasielewski, J., Stelinski, L., Grogan, K., & Ballen, F. (2023). Are avocados toast? A framework to analyze decision-making for emerging epidemics, applied to laurel wilt. *Agricultural Systems*, *206*, 103615. https://doi.org/10.1016/j.agsy.2023.103615

FAO. (2021a). *Emergency assistance to improve food and nutrition security of IDPS in South West region of Cameroon*. https://www.fao.org/documents/card/fr/c/CB3577EN/

FAO. (2021b). *The impact of disasters and crises on agriculture and food security*. https://www.fao.org/documents/card/en/c/cb3673en

FAOSTAT. (2021). *Trade: crops and livestock products*. https://www.fao.org/faostat/en/#home

Fogain, R., Mouliom Péfoura, A., Abadie, C., Escalant, J.-V., & Tomekpé, K. (1998). Banana and plantain
IPM in Cameroon: progress and problems. Banana IPM workshop November 1998, Nelspruit,
Afrique du Sud.

Foyou, V. E., Ngwafu, P., Santoyo, M., & Ortiz, A. (2018). The Boko Haram insurgency and its impact on
border security, trade and economic collaboration between Nigeria and Cameroon: An
exploratory study. *African Social Science Review*, *9*(1), 7.
https://digitalscholarship.tsu.edu/assr/vol9/iss1/7

Garrett, K., Bebber, D., Etherton, B., Gold, K., Sulá, A. P., & Selvaraj, M. G. (2022). Climate change effects
on pathogen emergence: artificial intelligence to translate big data for mitigation. *Annual review*
of Phytopathology, *60,* 357-378. https://doi.org/10.1146/annurev-phyto-021021-042636

Garrett, K. A. (2021). Impact network analysis and the ina R package: Decision support for regional
management interventions. *Methods in Ecology and Evolution*, *12*(9), 1634-1647.
https://doi.org/https://doi.org/10.1111/2041-210X.13655

Gibson, R., & Kreuze, J. F. (2015). Degeneration in sweetpotato due to viruses, virus-cleaned planting
material and reversion: a review. *Plant Pathology*, *64*(1), 1-15.
https://doi.org/10.1111/ppa.12273

Gosling, J. P. (2018). SHELF: the Sheffield elicitation framework. *Elicitation: The science and art of*
structuring judgement, 61-93. https://doi.org/https://doi.org/10.1007/978-3-319-65052-4_4

Graul, C., & Graul, M. C. (2016). Package 'leafletR'. *Interactive Web-Maps Based on the Leaflet JavaScript*
Library. Available online: https://github. com/chgrl/leafletR (accessed on 1 April 2016).

Hijmans, R. J. (2023). raster: Geographic data analysis and modeling. R package version 3.6-21.
https://rspatial.org/raster

Hirpa, A., Meuwissen, M. P., Tesfaye, A., Lommen, W. J., Lansink, A. O., Tsegaye, A., & Struik, P. C. (2010).
Analysis of seed potato systems in Ethiopia. *American journal of potato research*, *87*(6), 537-552.
https://doi.org/10.1007/s12230-010-9164-1

Hunt, M., Chénier, A., Bezanson, K., Nouvet, E., Bernard, C., de Laat, S., Krishnaraj, G., & Schwartz, L.
(2018). Moral experiences of humanitarian health professionals caring for patients who are
dying or likely to die in a humanitarian crisis. *Journal of International Humanitarian Action*, *3*(1),
1-13. https://doi.org/10.1186/s41018-018-0040-9

International Food Policy Research, I. (2020). *Spatially-disaggregated crop production statistics data in*
Africa south of the Sahara for 2017,   https://doi.org/10.7910/DVN/FSSKBW [Spatially-
disaggregated crop production statistics]. Harvard Dataverse, V2.
https://doi.org/doi:10.7910/DVN/FSSKBW

IOM. (2023). *More than 4.38 million people displaced in Ethiopia, more than half due to conflict: New*
IOM Report. https://ethiopia.iom.int/news/more-438-million-people-displaced-ethiopia-more-
half-due-conflict-new-iom-report

Jacobsen, K., Omondi, B., Almekinders, C., Alvarez, E., Blomme, G., Dita, M., Iskra-Caruana, M. L.,
Ocimati, W., Tinzaara, W., & Kumar, P. (2019). Seed degeneration of banana planting materials:
strategies for improved farmer access to healthy seed. *Plant Pathology*, *68*(2), 207-228.
https://doi.org/10.1111/ppa.12958

Jongejans, E., Skarpaas, O., Ferrari, M. J., Long, E. S., Dauer, J. T., Schwarz, C. M., Rauschert, E. S., Jabbour,
R., Mortensen, D. A., & Isard, S. A. (2015). A unifying gravity framework for dispersal. *Theoretical*
Ecology, *8*(2), 207-223. https://doi.org/10.1007/s12080-014-0245-5

Konje, C., Abdulai, A., Tange, A. D., Tarla, D., & Tita, M. A. (2019). Identification and management of pests
and diseases of garden crops in Santa, Cameroon. *Journal of Agriculture and Ecology Research*
International, *18*(2), 1-9. https://doi.org/10.9734/JAERI/2019/v18i230055

Macrotrends. (2023). *Cameroon Refugee Statistics 1970-2023*.
689   https://www.macrotrends.net/countries/CMR/cameroon/refugee-statistics
690   Margosian, M. L., Garrett, K. A., Hutchinson, J. S., & With, K. A. (2009). Connectivity of the American
691   agricultural landscape: assessing the national risk of crop pest and disease spread. *BioScience*,
692   *59*(2), 141-151. https://doi.org/10.1525/bio.2009.59.2.7
693   Mbah, L. T., Molua, E. L., Bomdzele, E., & Egwu, B. M. (2023). Farmers' response to maize production
694   risks in Cameroon: An application of the criticality risk matrix model. *Heliyon*, *9*(4), E15124.
695   https://doi.org/10.1016/j.heliyon.2023.e15124
696   Navarrete, I., López, V., Andrade-Piedra, J. L., Almekinders, C. J., Kromann, P., & Struik, P. C. (2022).
697   Agroecological settings and seed recycling account only partially for potato seed degeneration in
698   Ecuador. *Agronomy for Sustainable Development*, *42*(6), 109. https://doi.org/10.1007/s13593-
699   022-00840-1
700   Nduwimana, I., Sylla, S., Xing, Y., Simbare, A., Niyongere, C., Garrett, K. A., & Bonaventure Omondi, A.
701   (2022). Banana seed exchange networks in Burundi–Linking formal and informal systems.
702   *Outlook on Agriculture*, *51*(3), 334-348. https://doi.org/10.1177/00307270221103288
703   Ngatat, S., Hanna, R., Lienou, J., Ghogomu, R. T., Nguidang, S. P. K., Enoh, A. C., Ndemba, B., Korie, S.,
704   Fotso Kuate, A., & Nanga Nanga, S. (2022). Musa germplasm A and B genomic composition
705   differentially affects their susceptibility to banana bunchy top virus and its aphid vector,
706   Pentalonia nigronervosa. *Plants*, *11*(9), 1206. https://doi.org/10.3390/plants11091206
707   O'Hagan, A. (2019). Expert knowledge elicitation: subjective but scientific. *The American Statistician*,
708   *73*(sup1), 69-81. https://doi.org/10.1080/00031305.2018.1518265
709   OCHA. (2022). *Cameroon multi-stakeholder dialogue on refugees*.
710   https://reliefweb.int/report/cameroon/cameroon-multi-stakeholder-dialogue-refugees-
711   november-2022
712   OCHA. (2023a). *Cameroon humanitarian needs overview 2023*.
713   https://www.unocha.org/publications/report/cameroon/cameroon-humanitarian-needs-
714   overview-2023-march-2023
715   OCHA. (2023b). *Ethiopia: humanitarian needs overview 2023*.
716   https://reliefweb.int/report/ethiopia/ethiopia-humanitarian-needs-overview-2023
717   Okolle, J. N., Fansi, G. H., Lombi, F. M., Sama Lang, P., & Loubana, P. M. (2009). Banana entomological
718   research in Cameroon: how far and what next. *The African Journal of Plant Science and
719   Biotechnology*, *3*(1), 1-19.
720   Pedersen, T. L. (2022). ggraph: An Implementation of Grammar of Graphics for Graphs and Networks.
721   *Retrieved January*, *1*, 2018. https://CRAN.R-project.org/package=ggraph
722   Petsakos, A., Prager, S. D., Gonzalez, C. E., Gama, A. C., Sulser, T. B., Gbegbelegbe, S., Kikulwe, E. M., &
723   Hareau, G. (2019). Understanding the consequences of changes in the production frontiers for
724   roots, tubers and bananas. *Global Food Security*, *20*, 180-188.
725   https://doi.org/10.1016/j.gfs.2018.12.005
726   Poubom, C., Awah, E., Tchuanyo, M., & Tengoua, F. (2005). Farmers' perceptions of cassava pests and
727   indigenous control methods in Cameroon. *International Journal of Pest Management*, *51*(2),
728   157-164. https://doi.org/10.1080/09670870500131863
729   Raleigh, C., Linke, R., Hegre, H., & Karlsen, J. (2010). Introducing ACLED: An armed conflict location and
730   event dataset. *Journal of Peace Research*, *47*(5), 651-660.
731   https://doi.org/10.1177/002234331037891
732   RBA. (2022). *Agriculture performance assessment of 2022 meher season. Regional bureaus of agriculture
733   (Amhara, Oromia, SNNPR and Sidama).* https://panorama.solutions/en/organisation/regional-
734   bureaus-agriculture-amhara-oromia-tigray-and-snnpr

Savary, S., Willocquet, L., Pethybridge, S. J., Esker, P., McRoberts, N., & Nelson, A. (2019). The global burden of pathogens and pests on major food crops. *Nature Ecology & Evolution*, *3*(3), 430-439. https://doi.org/10.1038/s41559-018-0793-y

Scott, G. J. (2021). A review of root, tuber and banana crops in developing countries: past, present and future. *International Journal of Food Science & Technology*, *56*(3), 1093-1114. https://doi.org/10.1111/ijfs.14778

Sperling, L., & Almekinders, C. J. (2023). Informal Commercial Seed Systems: Leave, Suppress or Support Them? *Sustainability*, *15*(18), 14008. https://doi.org/10.3390/su151814008

Sperling, L., Gallagher, P., McGuire, S., March, J., & Templer, N. (2020). Informal seed traders: the backbone of seed business and African smallholder seed supply. *Sustainability*, *12*(17), 7074.

Sperling, L., & McGuire, S. (2010). Understanding and strengthening informal seed markets. *Experimental Agriculture*, *46*(2), 119-136. https://doi.org/10.1017/S0014479709991074

Sperling, L., Mottram, A., Ouko, W., & Love, A. (2022). Seed emergency response tool: Guidance for practitioners. *Produced by Mercy Corps and SeedSystem.org as a Part of the ISSD Africa Activity*.

Tesfaye, T., Feyissa, T., & Abraham, A. (2011). Survey and serological detection of sweet potato (Ipomoea batatas (L.) Lam) viruses in Ethiopia. *Journal of Applied Biosciences*, *41*, 2746-2756.

Tessema, G. L., & Seid, H. E. (2023). Potato bacterial wilt in Ethiopia: history, current status, and future perspectives. *PeerJ*, *11*, e14661. https://doi.org/10.7717/peerj.14661

TGHAT. (2022). *Tigray bureau of agriculture and natural resources, emergency and recovery support plan for the 2022/23 budget year, October 2022*. https://www.tghat.com/2023/01/12/government-of-tigray-bureau-of-agriculture-natural-resources-emergency-and-recovery-plan/

Thiele, G., Khan, A., Heider, B., Kroschel, J., Harahagazwe, D., Andrade, M., Bonierbale, M., Friedmann, M., Gemenet, D., & Cherinet, M. (2017). Roots, tubers and bananas: planning and research for climate resilience. *Open Agriculture*, *2*(1), 350-361. https://doi.org/10.1515/opag-2017-0039

Thomas-Sharma, S., Andrade-Piedra, J., Carvajal Yepes, M., Hernandez Nopsa, J., Jeger, M., Jones, R., Kromann, P., Legg, J., Yuen, J., & Forbes, G. (2017). A risk assessment framework for seed degeneration: Informing an integrated seed health strategy for vegetatively propagated crops. *Phytopathology*, *107*(10), 1123-1135. https://doi.org/10.1094/PHYTO-09-16-0340-R

Thomas-Sharma, S., Abdurahman, A., Ali, S., Andrade-Piedra, J., Bao, S., Charkowski, A., Crook, D., Kadian, M., Kromann, P., & Struik, P. C. (2016). Seed degeneration in potato: the need for an integrated seed health strategy to mitigate the problem in developing countries. *Plant Pathology*, *65*(1), 3-16. https://doi.org/10.1111/ppa.12439

Tize, I., Fotso, A. K., Nukenine, E. N., Masso, C., Ngome, F. A., Suh, C., Lendzemo, V. W., Nchoutnji, I., Manga, G., & Parkes, E. (2021). New cassava germplasm for food and nutritional security in Central Africa. *Scientific Reports*, *11*(1), 7394. https://doi.org/10.1038/s41598-021-86958-w

USAID. (2022). Strategic framework for early recovery, risk reduction, and resilience (ER4). https://www.usaid.gov/sites/default/files/2023-01/ER4_Framework-10.13.2022.pdf

Vasantha Raju, N., & Harinarayana, N. (2016). Online survey tools: A case study of Google Forms. National conference on scientific, computational & information research trends in engineering, GSSS-IETW, Mysore.

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis. *Springer-Verlag New York*, *3*(2), 180-185. https://ggplot2.tidyverse.org

Wiebe, K., Zurek, M., Lord, S., Brzezina, N., Gabrielyan, G., Libertini, J., Loch, A., Thapa-Parajuli, R., Vervoort, J., & Westhoek, H. (2018). Scenario development and foresight analysis: exploring options to inform choices. *Annual Review of Environment and Resources*, *43*, 545-570. https://doi.org/10.1146/annurev-environ-102017-030109

Wubaye, G. B., Gashaw, T., Worqlul, A. W., Dile, Y. T., Taye, M. T., Haileslassie, A., Zaitchik, B., Birhan, D. A., Adgo, E., & Mohammed, J. A. (2023). Trends in rainfall and temperature extremes in Ethiopia:

Station and agro-ecological zone levels of analysis. *Atmosphere*, *14*(3), 483.
https://doi.org/10.3390/atmos14030483

Xing, Y., Hernandez Nopsa, J. F., Andersen, K. F., Andrade-Piedra, J. L., Beed, F. D., Blomme, G., Carvajal-
Yepes, M., Coyne, D. L., Cuellar, W. J., Forbes, G. A., Kreuze, J. F., Kroschel, J., Kumar, P. L., Legg, J.
P., Parker, M., Schulte-Geldermann, E., Sharma, K., & Garrett, K. A. (2020). Global Cropland
Connectivity: A Risk Factor for Invasion and Saturation by Emerging Pathogens and Pests.
BioScience, *70*(9), 744-758. https://doi.org/10.1093/biosci/biaa067