# Early mechanisms of aortic failure in a zebrafish model for thoracic aortic dissection and rupture

Michiel Vanhooydonck MSc.[1], Maxim Verlee MD.[1], Marta Santana Silva MSc.[1], Lore

Pottie PhD.[1], Annekatrien Boel PhD.[2], Matthias Van Impe MSc.[3], Hanna De Saffel

BSc.[1], Lisa Caboor BSc.[1], Piyanoot Tapaneeyaphan MSc.[1], Anne Bonnin PhD.[4], Patrick

Segers PhD.[3], Adelbert De Clercq PhD.[5], Andy Willaert PhD.[1], Delfien Syx PhD.[1],

Patrick Sips PhD.[1a], Bert Callewaert MD. PhD.[1a*]

[1] Center for Medical Genetics Ghent (CMGG), Department of Biomolecular Medicine,

Ghent University, 9000 Ghent, Belgium

[2] Ghent-Fertility and Stem Cell Team (G-FaST), Department for Reproductive Medicine,

Ghent University, 9000 Ghent, Belgium

[3] Biophysical Models for Medical Applications (bioMMeda), Institute of Biomedical

Engineering and Technology (IBiTech), Ghent University, 9000 Ghent, Belgium.

[4] Paul Scherrer Institut, Swiss Light Source, 5232 Villigen PSI, Switzerland.

[5] Evolutionary Developmental Biology, Biology Department, Ghent University, 9000

Ghent, Belgium.

[a] Bert Callewaert and Patrick Sips contributed equally as senior authors.

*Corresponding Author: Bert Callewaert, PhD, MD

Bert.Callewaert@Ugent.be; ORCID: 0000-0002-9743-4205

Word count: 8888

## ABSTRACT

Thoracic aortic aneurysm and dissection (TAAD) associates with a high mortality rate. Despite the existence of different mouse models for TAAD, the underlying disease mechanisms remain elusive. Treatment options are limited and mainly consist of surgical repair at critical aortic diameters as current pharmacological interventions are unable to stop disease progression.

In humans, loss of function (LOF) of *SMAD3* and *SMAD6* impairs vascular homeostasis, increasing the risk for TAAD. We developed a zebrafish model for thoracic aortic dissection/rupture by targeting both ohnologs of *smad3* and *smad6*. At 10 days post fertilization, we found an increased diameter of the ventral aorta in *smad3a*$^{-/-}$;*smad3b*$^{-/-}$ double knockout zebrafish, while *smad6a*$^{-/-}$;*smad6b*$^{-/-}$ double knockout zebrafish have a reduced aortic diameter associated with early mortality. We discovered that a *smad3a*$^{-/-}$;*smad3b*$^{-/-}$;*smad6a*$^{-/-}$;*smad6b*$^{-/-}$ quadruple knockout (qKO) zebrafish model is viable and survives to adulthood, although exposure to stress leads to sudden death. Histological analysis of the adult ventral aorta shows medial elastolysis, aortic dissections and ruptures at sites exposed to high biomechanical stress. RNA-sequencing of 5 days post fertilization qKO zebrafish indicates a profile of reduced negative regulation of proteolysis and upregulation of melanogenesis, a previously unaddressed pathway in this pathology. We confirm that pharmacological modulation of tyrosinase, the enzyme responsible for the production of melanin, influences aortic morphology.

46    Overall, the qKO mutant, thus far the only known zebrafish model of thoracic aortic

47    dissection and rupture, reveals novel SMAD3/6-dependent pathways that impact

48    thoracic aortic homeostasis, in this way opening avenues for the development of novel

49    treatments in TAAD.

50 **NON-STANDARD ABBREVIATIONS AND ACRONYMS**

| | |
|---|---|
| AA3 | aortic arch 3 |
| AA4 | aortic arch 4 |
| BMPs | bone morphogenetic proteins |
| Co-SMAD | common mediator SMAD |
| CPM | counts per million |
| DEG | differentially expressed genes |
| DKO | double knockout |
| dpf | days post fertilization |
| ECM | extracellular matrix |
| FDR | false discovery rate |
| GDFs | differentiation factors |
| GO | gene ontology |
| GRCz11 | Genome Reference Consortium Zebrafish Build 11 |
| GSEA | gene set enrichment analysis |
| ID | intellectual disability |
| I-SMAD | inhibitory SMAD |
| KEGG | Kyoto Encyclopedia of Genes and Genomes |
| LDS3 | Loeys-Diets syndrome type 3 |
| LOF | loss of function |
| mpf | months post fertilization |
| MR | magnetic resonance |
| PSI | Paul Scherrer Institut |
| PTU | 1-phenyl-2-thiourea treatment |
| qKO | smad3a$^{-/-}$;smad3b$^{-/-}$;smad6a$^{-/-}$;smad6b$^{-/-}$ quadruple knockout |
| RO | reverse osmosis |
| R-SMAD | receptor-regulated SMAD |

| | |
|---|---|
| SKO | single knockout |
| *smad3a/b* DKO | *smad3a$^{-/-}$;smad3b$^{-/-}$* double knockout |
| *smad6a/b* DKO | *smad6a$^{-/-}$;smad6b$^{-/-}$* double knockout |
| TAA | thoracic aortic aneurysm |
| TAAD | thoracic aortic aneurysm and dissection |
| TAD | thoracic aortic dissection |
| TEM | transmission electron microscopy |
| TGF-β | transforming growth factor β |
| ZIRC | Zebrafish International Research Center |

## INTRODUCTION

Thoracic aortic dissection (TAD) affects 3-4 individuals per 100,000 people per year[1]. This life-threatening event is often, but not exclusively, preceded by thoracic aortic aneurysm (TAA). The disease mechanisms are incompletely understood, but evidence from genetic disorders implicates altered cell-matrix interactions (by impaired hemodynamic sensing or impaired matrix assembly), as well as aberrant transforming growth factor β (TGF-β) signaling[2–5]. Despite several mouse models for TAD[6], the initial molecular mechanisms remain obscure.

Members of the TGF-β superfamily include TGF-β proteins, bone morphogenetic proteins (BMPs), activins, and growth and differentiation factors (GDFs). Binding of TGF-β to the tetrameric TGFβ-receptor complex results in phosphorylation of the receptor-regulated (R-) SMAD2 and -3 proteins. Binding of BMP and GDF to the tetrameric BMP-receptor complex results in phosphorylation of R-SMAD1, -5, and -8. Phosphorylated R-SMAD proteins recruit the common mediator (Co-) SMAD4 to form a heterotrimeric complex and translocate to the nucleus to activate the transcription of the TGF-β or BMP target genes[7]. The TGF-β and BMP signaling is negatively regulated by the inhibitory (I-) SMAD6 and -7 proteins. I-SMAD6 primarily suppresses the pathway activated by the BMP-receptor, whereas I-SMAD7 inhibits both the TGF-β and BMP induced pathways[8].

Pathogenic variants in *SMAD3* cause Loeys-Dietz syndrome type 3 (LDS3). LDS3 typically presents with early-onset osteo-arthritis, and arterial aneurysm and dissection, mainly affecting the ascending aorta. Other features include arterial tortuosity, mitral valve prolapse and craniofacial characteristics[2,9]. Pathogenic variants in *SMAD6* are

associated with craniosynostosis[10], radio-ulnar synostosis[11] or congenital heart disease

including bicuspid aortic valve[12] with or without TAAD[13] and Shone complex with a

hypoplastic ascending aorta and arch[14]. Biallelic variants in *SMAD6* underly more

complex cardiovascular phenotypes[15]. Interestingly, identical pathogenic variants in

*SMAD6* may result in either cardiovascular anomalies, craniosynostosis or radioulnar

synostosis[13]. Because *SMAD3* and *SMAD6* have been in linkage disequilibrium

throughout evolution from teleost fish to human, it has been postulated that *SMAD3*

could have a modifier role on *SMAD6* deficiency[16]. In line with this observation, we

identified a 1.5 Mb 15q22.31-15q23 microdeletion encompassing *SMAD3* and *SMAD6*

in an 11-year-old-boy and his 33-year-old mother, both presenting with bicuspid aortic

valve and arterial tortuosity, but no aneurysms, along with skeletal and cutaneous

anomalies of LDS3, hinting that deletion of *SMAD6* may influence the phenotype

caused by *SMAD3* LOF (**Supplementary Case Presentation**[17]). Due to the teleost

specific genome duplication, zebrafish have two ohnolog copies for both *SMAD3* and

*SMAD6*[18]. We established zebrafish models deficient for each of these ohnologs

identified as *smad3a* and *smad3b,* and *smad6a* and *smad6b*, respectively. The human

SMAD3 (ENSP00000332973) protein has a sequence similarity of 97.17% with

zebrafish Smad3a (ENSDARP00000045373) and 93.4% with zebrafish Smad3b

(ENSDARP00000043454). SMAD6 (ENSP00000288840) is less conserved with a

similarity of 53.3% and 62.38% with zebrafish Smad6a (ENSDARP00000112408) and

Smad6b (ENSDARP00000091342) respectively. Smad6a and Smad6b show 58.33%

protein sequence similarity, which is considered large enough to assume similar

function[19]. The sequence similarity and the same spatial expression patterns of *Smad3a*

and *Smad3b* as well as *Smad6a* and *smad6b* suggest[20–22].

In this study, we investigated modifying effects of *smad3a/smad3b* on *smad6a/smad6b*

and vice versa which enabled us to model TAAD in zebrafish. Additionally, we identified

early molecular mechanisms in aortic failure to identify druggable pathways.

## METHODS

### Ethics statement and housing

This study was approved by the Animal Ethics Committee of the Ghent University Faculty of Medicine and Health Sciences (Permit number: ECD 14/70) and housed adhering to the general guidelines, in agreement with EU Directive 2010/63/EU for laboratory animals [23,24].

### Zebrafish lines: design and generation

Smad3a[sa2363/+] was acquired from the Zebrafish International Research Center (ZIRC). All other models were generated using CRISPR/Cas9 gene editing technology utilizing CRISPRdirect software[25–27]. Specific gBlocks for all target sites are listed in **Supplementary Table 1**. Genotyping primers are given in **Supplementary Table 2**. All variants are described using Genome Reference Consortium Zebrafish Build 11 (GRCz11) and described in **Supplementary Table 3**.

### Vasculogenesis in embryos

Zebrafish were outcrossed to the transgenic Tg(fli1:EGFP) line in order to visualize endothelial cells as previously described[28]. Head length was used as normalization factor. An overview of the acquired measurements and specific breeding schemes are given in **Supplementary Fig. 1a-c**.

### 3D reconstruction of the ventral aorta in adult zebrafish

Serial 5 μm paraffin cross sections of the entire ventral aorta of zebrafish between 6 and 13 months post fertilization (mpf) were stained with Weigert's iron hematoxylin, rinsed in water and stained with Resorcin-Fuchsin for one hour followed by two wash steps in

124 95% ethanol and one wash with reverse osmosis (RO) water. Slides were dehydrated

125 and mounted with Entellan mounting medium. Pictures of serial sections were aligned

126 via the TrakEM2 1.0[29–31] and 3D reconstructed with Mimics 24.0 software.

### *Ultrasound analysis*

128 Ultrasound analysis was performed on zebrafish of 6-7 mpf as previously described[32].

129 All measurements were performed in Vevo Lab 5.5.0.

### *Acute net handling stress induction*

131 Induction of handling stress was adapted from previously described procedures[33]. Six

132 mpf zebrafish were netted and air suspended for 90 seconds, returned to the housing

133 tank for 90 seconds, netted and air suspended for 90 seconds and returned to the

134 housing tank.

### *Alizarin Red staining for mineralized bone*

Alizarin red staining for mineralized bone was performed as previously described[34], at

13 mpf. The qKO were stained after sudden death between 6 and 13 mpf.

Ectopic bone on the sinistral skeletal elements of the vertebral bodies was scored on

abdominal and caudal vertebrae of adult zebrafish in lateral position[35].

Skull morphometrics were performed on images of the zebrafish heads in lateral

position. Acquired lengths were measured in ImageJ and are given in **Supplementary**

**Fig. 1b.** Diameter of the eye was used for normalization since eye diameter correlates

with standard length[36]. Cranial surface area was measured with standard length as

normalization factor.

### *Transcriptome analysis*

RNA sequencing was performed on five pools, each containing five qKO or WT control cousins. Following RNA extraction (RNeasy® Mini kit), libraries were prepared (TruSeq Stranded Total RNA kit from Illumina) and sequenced on a NovaSeq6000 instrument (Illumina).

Following quality control [37,38], raw reads were mapped (STAR[39]) and through R packages, additional normalization and filtering was performed[40–42]. The top list of DEG was obtain via differential expression analysis (edgeR), with the necessary adjustments to account for multiple testing[43,44]. An adjusted p-value of 0.05 and a fold change of 2 were used as thresholds to determine genes of relevance. Finally gene enrichment analysis was carried out[45]. Further details can be found in the **Supplementary Methods**.

### *Synchrotron phase contrast micro Computed tomography imaging*

Propagation-based phase-contrast synchrotron X-ray imaging was performed at the TOMCAT (X02DA) beamline of the Swiss Light Source (Paul Scherrer Institute in Villigen, Switzerland) as previously described[46]. Tomographic reconstruction was performed using the Gridrec algorithm after applying the Paganin phase retrieval method[47,48].

### *3D Biomechanical modeling to assess principal stress at systolic peak in the aortic wall*

In a previous study[46], zebrafish-specific 3D biomechanical fluid-structure interaction models of five different 13-month-old zebrafish were developed. All parameter settings

as described in[46] were maintained except for flow through right aortic arch 2, 3 and 4,

which was set to nearly zero in order to mimic the vascular organization found in qKO

mutants.

***Transmission electron microscopy***

In short, WT and qKO zebrafish of 16 mpf were euthanized, fixed in Karnovsky fixative,

followed by decalcification for three weeks. Post-fixation in 1% osmium tetroxide, 0.1M

cacodylate buffer with 8% saccharose and 0.004% $CaCl_2$ was followed by *en bloc*

staining in 1% aqueous uranyl acetate. After dehydration, the samples were embedded

in Spurr's resin. 80 nm sections were stained with uranyl acetate and lead citrate and

examined by transmission electron microscopy (TEM) (JEM 1010, JEOL) equipped with

a CCD side-mounted Veleta camera (EMSIS).

***Swimming behavior studies***

Swimming behavior of 9 mpf WT, *smad3a$^{+/-}$;smad3b$^{-/-}$;smad6a$^{+/-}$;smad6b$^{-/-}$* and qKO

zebrafish was analyzed in a custom-made dark observation chamber (Noldus) equipped

with the Basler GenICam. Data was analyzed using the EthoVision XT 17 software

(Noldus). Zebrafish were placed individually in a tank and could acclimatize for 10

minutes before a 10-minute test period, was performed in the dark. Movement is

depicted as total distance travelled during the test period, normalized for standard

length. Movement frequency was also recorded.

### *1-phenyl-2-thiourea treatment*

187 At 1 day post fertilization (dpf), 0.003% 1-phenyl-2-thiourea treatment (PTU) in E3

188 medium was administered to WT zebrafish. The medium was changed daily until 10 dpf.

189 Starting from 5 dpf, zebrafish were given dry food for 2 hours daily, after which the

190 medium was refreshed again.

### *Statistical analyses*

All statistical analyses were performed with GraphPad Prism version 10.1.1 for

Windows (GraphPad Software, Boston, Massachusetts USA, www.graphpad.com).

Kruskal-Wallis test with Dunn's multiple comparisons test against WT controls was used

to determine if there was an increase in ectopic bone on the skeletal elements of the

vertebral bodies. For PTU treatment, a two-tailed t-test was used, when standard

deviation differed significantly between the groups, Welch's correction was applied. For

all other comparisons, One-way ANOVA with Dunnett's multiple comparison test against

WT controls was used. Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple

comparison test against WT controls was used when the genotypes showed a

significantly different standard deviation, determined with the Brown-Forsythe test.

## RESULTS

### *Disruption of one of the smad3 or smad6 ohnologs does not show a cardiovascular or skeletal phenotype in zebrafish*

Using CRISPR/Cas9 gene editing technology, we generated frameshift indels in the *smad3b*, *smad6a*, and *smad6b* genes, resulting in the generation of a premature termination codon in each mutant: s*mad3b*$^{c.455\_459delinsATG/c.455\_459delinsATG}$, *smad6a*$^{c.905\_906delTG/c.905\_906delTG}$ and *smad6b*$^{c.283\_287delACGGT/c.283\_287delACGGT}$ (**Supplementary Table 3**). To study *smad3a* function we made use of the available *Smad3a*$^{sa2363/+}$ line, in which a premature termination codon is introduced (c.682G>T). For the reader's benefit, we will use the term single knockout (SKO) for *smad3a, smad3b, smad6a* and *smad6b* single homozygous knockout zebrafish. At 5 dpf, measurement of the ventral aorta in the transgenic *Tg(fli1:EGFP)* reporter background is similar for heterozygous and SKO zebrafish for each gene as compared to sibling controls (**Error! Reference source not found.**).

However, incross of *smad3a* SKO results in100% embryonic lethality at 2 dpf with severe body axis deformities and pericardial edema formation, while incross of *smad3b* SKO and *smad3a*$^{+/-}$*;smad3b*$^{-/-}$ shows offspring with normal survival, suggesting that maternal *smad3a* expression is necessary for early embryonic survival, which is supported by previously published results[20]. Incross of *smad6a* or *smad6b* SKO showed normal survival.

In 6-7 mpf adult zebrafish, ultrasound measurements show that all measured parameters of cardiac function are similar in all SKO compared heterozygous mutants and sibling controls, with an exception of *smad6a*$^{-/-}$ which shows a small increase in

normalized projected surface area of the ventricle in diastole (**Error! Reference source**
**not found.**).

In addition, skeletal evaluation and alizarin red staining for mineralized bone at 13 mpf
do not show any vertebral column deformities or an increase in ectopic bone formation
(**Figure** , **Error! Reference source not found.-6**). Skull morphometrics are normal,
except for the *smad3a* SKO, in which a decrease of the snout to frontal bone distance is
observed (**Error! Reference source not found.-8**).

### *Smad3a/b DKO and smad6a/b DKO show altered vasculogenesis*

To bypass embryonic lethality due to loss of maternal *smad3a*, we performed a cross
between a *smad3a$^{+/-}$;smad3b$^{-/-}$* female and *smad3a/b* DKO male zebrafish to obtain
50% *smad3a$^{+/-}$;smad3b$^{-/-}$* and 50% *smad3a/b* DKO zebrafish, which survive normally. At
10 dpf, the *smad3a/b* DKO show a significant increase in normalized surface area of the
aortic segment between aortic arch 3 and 4 (p = 0.0449), a reduced normalized length
of the segment (p = 0.0366) and an increased normalized mean aortic diameter (p =
0.0335). The diameter adjacent to the bulbus arteriosus remains unchanged (**Error!**
**Reference source not found.**).

The *smad6a/b* DKO show normal early survival rates, as indicated by normal Mendelian
distribution of all genotypes in 10 dpf-old offspring obtained from *smad6a$^{+/-}$;smad6b$^{+/-}$*
incrosses. Nevertheless, *smad6a/b* DKO zebrafish are unable to survive to adulthood.

*Smad6a/b* DKO zebrafish show a significant reduction of the ventral aortic diameter at
10 dpf (p<0.0001). The diameter adjacent to the bulbus arteriosus as well as the
normalized surface area of the aorta segment is decreased (p<0.0001, p<0.0001). To a

lesser extent, the diameter of the aorta segment of *smad6a*$^{+/-}$*;smad6b*$^{-/-}$ is also reduced

(p = 0.0016), while the length of the aorta segment remains unchanged in all *smad6*

knockouts (**Error! Reference source not found.**).

Ultrasound analysis at 6-7 mpf of all viable *smad3* (*smad3a*$^{+/-}$*;smad3b*$^{-/-}$, *smad3a*$^{-/-}$

*;smad3b*$^{+/-}$ and *smad3a/b* DKO) or *smad6* (*smad6a*$^{+/-}$*;smad6b*$^{-/-}$ and *smad6a*$^{-/-}$*;smad6b*$^{+/-}$

) KO combinations do not show any significant differences in cardiac parameters (**Error!**

**Reference source not found.-4**).

***Loss of smad3 ohnologs in zebrafish induces ectopic bone formation and***

***vertebral column deformations***

Alizarin red staining for mineralized bone on 13 mpf *smad3a/b* DKO zebrafish shows

ectopic bone formation on the ribs, neural arches, neural spines, haemal arches, and

haemal spines (p = 0.0012) (**Figure** ) as well as hyperlordosis, notochord mineralization

and notochord sheet mineralization (**Supplementary Fig 5)**. Overall, the thickness of

the haemal arches is increased (**Fig. 4**) and the normalized surface area of the cranial

roof is reduced compared to WT controls (p = 0.0101) (**Error! Reference source not**

**found.**). The presence of a single WT copy of either *smad3a* or *smad3b* prevents any of

these skeletal malformations to develop (**Fig. 4, Supplementary Fig. 8**). Skull

morphometrics show a mild decrease of the distance between the tip of the snout and

the most posterior part of the supraoccipital bone in s*mad3a*$^{-/-}$ zebrafish (p = 0.0486),

but not in the *smad3a/b* DKO (**Error! Reference source not found.**).

Likewise, a significant increase in ectopic bone can be seen in 13 mpf *smad6a*$^{-/-}$

*;smad6b*$^{+/-}$ (p = 0.0022) as well as in *smad6a*$^{+/-}$*;smad6*$^{-/-}$ (p = 0.0033) zebrafish,

although to a lesser extent than observed in the *smad3a/b* DKO (**Figure** ). Since the

*smad6a/b* DKO does not reach adulthood, these features cannot be analyzed in this

genotype. No premature fusion of the cranial roof bones is observed, and the

normalized surface areas of the cranial roof are similar to controls (**Error! Reference**

**source not found.**). Skull morphometrics do show a decrease in length between the tip

of the snout and tip of the frontal bone in *smad6a$^{+/-}$;smad6b$^{-/-}$* mutants (p = 0.0056)

(**Error! Reference source not found.**).

## *Loss of smad3 and smad6 ohnologs in zebrafish induces ventral aortic dissection and rupture*

To investigate potential modifying effects of *smad3* on *smad6* and vice versa, we

generated a *smad3a$^{-/-}$;smad3b$^{-/-}$;smad6a$^{-/-}$;smad6b$^{-/-}$* qKO zebrafish line

(**Supplementary Table 3**).

Unlike the *smad6a/b* DKO, qKO zebrafish reach adulthood and generate viable

offspring. Microscopic analysis of vasculogenesis at 10 dpf shows a severe decrease of

the normalized surface area of the aortic projection (p<0.0001), the normalized mean

diameter of the aortic segment between aortic arch 3 and 4 (p = 0.0241), and the

normalized diameter adjacent to the bulbus arteriosus (p = 0.0005) compared to WT

controls, similar to *smad3a$^{+/-}$;smad3b$^{-/-}$;smad6a$^{+/-}$;smad6b$^{-/-}$* zebrafish (p<0.0001,

p<0.0001, p = 0.0032, respectively). The normalized length of the aortic segment is only

decreased in the qKO (p = 0.006) (**Fig. 1**, **Supplementary Fig. 9**). In addition, the qKO

shows tortuosity of the ventral aorta, aortic arches and hypobranchial artery (**Error!**

**Reference source not found.**, **Error! Reference source not found.10**).

In adult qKO zebrafish, we observed sudden deaths, mostly after breeding, netting, or

changing the tank positions. To investigate stress as a trigger, we performed a netting

test that is harmless in WT controls, but results in a mortality rate of 60% in qKO.

Sudden death associated with a red discoloration anterior in the abdominal region.

Sectioning of the entire ventral aorta and staining with Resorcin-Fuchsin to visualize the

elastic fibers shows the presence of aortic intramural hematomas and false lumens in

multiple qKO. 3D-reconstructions of the aorta based on histological images and from

synchrotron micro-CT images of intact zebrafish, confirm the presence of regions of

ventral aortic damage (**Fig. 2** and **Supplementary movie 1-8**).

Regions of ventral aortic damage start close to a branching point (6/6 qKO, 0/4 WT) and

dissection or rupture is present at the most affected sites, (5/6 qKO, 0/4 WT). The

diameter of the aortic branches is asymmetrically distributed in all qKO zebrafish (6/6

qKO, 0/4 WT) with one side consistently showing narrower branches than the other. In

more severe cases, one or more aortic arches are missing (2/5 qKO, 0/4 WT) or

attached at the wrong side of the ventral aorta (1/6 qKO, 0/4 WT) (**Supplementary**

**Movie 9-13**). No morphological differences can be detected in WT controls

(**Supplementary Movie 14-16**) or s*mad3a/b* DKO zebrafish (**Supplementary Movie**

**17-19**).

TEM shows a severely decreased elastin deposition in the internal elastic lamina of 15

mpf qKO zebrafish. Additionally, abnormal collagen deposition is detected near the

intima of the ventral aorta (**Fig. 2**).

Cardiac ultrasound analysis in 6-7 mpf qKO zebrafish shows regurgitation on ventricular

in- ($p = 0.0009$) and outflow ($p<0.0001$), suggestive of valve dysfunction

(**Supplementary Fig. 11**). Elastin staining of paraffin sections of the valves shows

hypertrophy of the valve interstitial cells, resulting in a more rounded appearance of the

qKO bulboventricular valve in contrast to the normal heart-shaped structure in WT

control zebrafish. Similarly, the atrioventricular valve shows hypertrophy and altered

morphology of the valve interstitial cells (

**Figure**, **Supplementary Fig. 12**).

### *Aortic damage in the qKO zebrafish correlates with increased stress at branching regions*

In qKO zebrafish, asymmetric branching and reduction of the diameter of the aortic arches is apparent. To determine if the morphological changes of the aorta in adult qKO zebrafish increase the principal stress within the vessel wall and to identify which locations endure the largest differences, aortic morphology was compared using *in silico* simulation models representing 13 mpf WT and qKO zebrafish aortae. To model the severe reduction of the diameter of the qKO aortic arches, the blood flow through the right aortic arches (aortic arch 2, 3 and 4) was reduced to nearly zero in the model. The principal stress within the aortic wall was increased in all 5 qKO simulations, with the most affected region close to branching points. This corresponds to the *ex vivo* observations of the location of regions of damage in the ventral aorta of qKO zebrafish (**Error! Reference source not found.**).

### *qKO zebrafish show skeletal aberrations, but largely normal swimming behavior*

The qKO show an increased presence of ectopic bone formation (p = 0.0014). Scoliosis, hyperlordosis and hyperkyphosis, bending of the arches and ribs, as well as intervertebral ligament mineralization and notochord mineralization are apparent in the qKO zebrafish only (**Figure** , **Error! Reference source not found.**).

In qKO zebrafish, the nasal region of the skull is underdeveloped with a significantly reduced distance between the tip of the snout and tip of the frontal bone (p<0.0001).

(**Figure** , **Error! Reference source not found.**). Interestingly, qKO show normal values

for the normalized surface area of the cranial roof in contrast to the *smad3a/b* DKO,

which was significantly decreased (p = 0.0101) (**Supplementary Fig. 8).** Since qKO

embryos show a tortuous hypobranchial artery, which provides blood flow to the teeth,

tooth morphology was evaluated, but appears normal (**Error! Reference source not**

**found.**).

At 9 mpf, qKO zebrafish show a decrease in movement frequency (p = 0.0197), but not

in normalized total distance traveled compared to WT zebrafish (**Error! Reference**

**source not found.**).

***Bulk RNA sequencing of 5 dpf qKO* shows a reduction in the negative**

**regulation of proteolysis and upregulation of melanogenesis**

Bulk RNA sequencing in 5 dpf qKO and 2[nd] degree related WT control zebrafish

identified 1131 differentially expressed genes (DEG) (adjusted p-value <0.05)

(**Supplementary Table 4**). The top 10 enriched GO-terms in the upregulated gene set

mostly involve pigmentation (pigment metabolic process, pigment biosynthetic process,

melanosome organization, pigment granule organization, melanocyte differentiation,

pigmentation, developmental pigmentation). One of the central DEG driving this

enrichment is the *mitfa* transcription factor, which is known to signal downstream of

BMP and governs melanocyte differentiation and development via tyrosinase (log FC *tyr*

2.01), tyrosinase related protein 1 (log FC *tyrp1a* 1.82 and log FC *tyrp1b* 2.08), the

premelanosome protein (log FC *pmela* 1.44 and log FC *pmelb* 3.66), dopachrome

tautomerase (log FC *dct* 2.6), and the melanosomal transporter (log FC *slc24a5*

2.35)[49,50]. The top 10 enriched GO-terms in the downregulated gene set are linked to

negative regulation of proteolysis (negative regulation peptidase activity, negative

regulation proteolysis, regulation of peptidase activity, negative regulation of

endopeptidase activity, regulation of proteolysis) and immune response (regulation of

immune response, positive regulation of immune system process) (**Supplementary**

**Table 5-6**). A heatmap with stringent thresholds (p.adjust ≥ 0.0001 and |log2FC| ≥ 1)

and a volcano plot are shown in **Figure** . Annotated dysregulated genes with a known

role in aortic wall homeostasis, include *loxl3a*, *emilin3a* and *fbn2b* (elastic fiber

assembly), *vcana* and *lamc2* (vascular cell adhesion glycoproteins), *mhc1uma* (a

zebrafish specific MHC class I gene), and *uts2a* (a vasoconstrictor linked to

hypertension and heart failure in humans). Dysregulation of *omd* (a transcription factor,

activated via BMP signaling, that links to cell adhesion and bone mineralization) might

relate to the observed bone phenotypes.

***Tyrosinase inhibition increases diameter of the ventral aorta in 10 dpf***

***zebrafish***

To investigate the involvement of melanogenesis-related pathways in aortic

development, we administered 0.003% PTU, a known inhibitor of tyrosinase in the

medium of WT zebrafish embryos starting at 1 dpf. At 10 dpf, treated zebrafish show a

significant increase of normalized ventral aortic surface area (p = 0.0066) and mean

normalized aorta diameter between aortic arch 3 and 4 (p = 0.0065) compared to

vehicle control. Length of the aortic segment and diameter adjacent to the bulbus

arteriosus remained unchanged compared to non-treated sibling controls (**Figure** ).

## DISCUSSION

TGFβ and BMP signaling show significant crosstalk during vascular and skeletal development and homeostasis. Our results support the existence of modifier effects particularly on aortic phenotypes associated with *SMAD6*-deficiency, which range from bicuspid aortic valve with or without ascending aortic dilatation to Shone complex with a hypoplastic ascending aorta and aortic arch.

In the genome of mammals and teleost fish, *SMAD3* and *SMAD6* locate closely together, but in different topologically associated domains[16]. Since zebrafish express *smad3a*, *smad3b, smad6a* and *smad6b*[51–54], this model provides flexibility to investigate the interactions between these genes upon gene dosage. Although maternal *smad3a* is required during early embryogenesis, which is supported by previous findings[48], SKO show no obvious vascular or skeletal phenotype, except for mild craniofacial alterations in *smad3a*[-/-] zebrafish. However, *smad3a/b* or *smad6a/b* DKO do manifest cardiovascular and mild skeletal phenotypes, supporting compensation between the ohnologs of either smad3 or smad6, in line with their strongly conserved protein structure [19].

*Smad6a/b* and *smad3a/b* DKO show opposite vascular phenotypes. *Smad3a/b* DKO zebrafish show an increase in aortic diameter in early larval stages. This aligns with previous findings in which loss of *alk5*[-/-], the type 1 TGF-β receptor which activates SMAD3, leads to an increased diameter of the cardiac outflow tract in zebrafish embryos, followed by a failed development of the ventral aorta causing death at 7 dpf[55]. It is plausible that in *smad3a/b* DKO limited amounts of Smad2/Smad4 complexes partially rescue the phenotype preventing early death and an adult cardiovascular

phenotype, in contrast to Alk5 loss of function, in which both Smad2 and Smad3 will not

be activated[56].

S*mad6a/b* DKO show a reduced aortic diameter, and early death. The qKO also shows

a reduction of the aortic diameter and arterial tortuosity, but survives to adulthood. The

observation that the *smad3a$^{+/-}$;smad3b$^{-/-}$;smad6a$^{+/-}$;smad6b$^{-/-}$* shows a more severe

reduction in aortic diameter than the qKO might be explained by the exclusion of the

most affected qKO embryos, since clear edge detection of the ventral aorta in these

embryos is hindered due to excessive tortuosity. Hence, it seems reasonable to

generalize that smad3 loss of function increases, while lack of smad6 decreases the

diameter of the aorta. The overall net result in the qKO shows a decreased diameter of

the ventral aorta with asymmetrical branching and a high susceptibility for dissection

and rupture in adult zebrafish. It has been previously reported that zebrafish *smad6b* is

involved together with BMP in modulation of lateral sprouting of the dorsal aorta[54].

Abnormal aortic branching results in altered hemodynamics and elevated systolic wall

stress, as confirmed by our fluid-structure interaction models presented here. Areas

exposed to perturbed local flow patterns prove prone to aortic wall damage and

dissection, as demonstrated by histological and synchrotron imaging of the aorta. TEM

analysis reveals a severe reduction of elastic fiber deposition in the inter-elastic

laminae, which can be linked to a lessened resilience to hemodynamic stress.

Additionally, abnormal collagen deposition near the intima of qKO zebrafish recapitulate

increased collagen deposition in human TAAD[57], indicating that similar

pathophysiological processes are activated in our zebrafish model. The presence of an

atrioventricular and bulboventricular valve defect in the qKO is in line with the role of

SMAD6 in human bicuspid aortic valve disease[58]. It is conceivable that unopposed loss

of both *smad6* ohnologs in zebrafish leads to a failure to maintain aortic homeostasis or

to severe valve dysfunction, resulting in the premature death observed in the *smad6a/b*

DKO late juvenile zebrafish.

The spectrum of bone phenotypes observed in the different genotypes confirms the

modulatory interactions between the different *smad* genes. Indeed, *smad3a/b* DKO, but

not *smad3a$^{+/-}$;smad3b$^{-/-}$* or *smad3a$^{-/-}$;smad3b$^{+/-}$* zebrafish, show increased ectopic bone

formation, a decreased normalized cranial roof surface area, and vertebral column axis

deviations. In contrast, the *smad6a$^{+/-}$;smad6b$^{-/-}$* and the *smad6a$^{-/-}$;smad6b$^{+/-}$* zebrafish

do show an increase in ectopic bone formation, although it needs to be highlighted that

adult *smad6a/b* DKO cannot be studied since they do not survive until adulthood. The

qKO zebrafish show more ectopic bone formation, and more severe column axis

deviations. These observations suggest a larger impact of BMP signaling on skeletal

development, that is further impacted by loss of TGFβ signaling.

Enrichment analysis of bulk RNA sequencing data of qKO embryos at 5 dpf revealed

upregulation of multiple pathways related to pigmentation, including the central

transcription factor *mitfa* as well as downstream effectors in melanin production such as

*tyrp1a*, *tyrp1b*, *tyr* and *dct*. MITF is a known downstream target of BMP signaling that is

inhibited by SMAD6[59–61]. MITF also acts upstream of lysosome biogenesis and it was

previously shown that lysosomal dysfunction in zebrafish due to loss of function of

*atp6v1e1b* associates with dilatation of the ventral aorta and narrow aortic branches[28].

Finally, MITF is known to be important for cell metabolism and cell cycle regulation,

promoting mitochondrial biogenesis and having complex effects on proliferation

depending on the level of MITF activity[62]. It is conceivable that one or more of these

effects of MITF are linked to the development of the phenotype observed in our qKO

model.

The transcriptional changes of pigmentation-related pathways in the qKO model

prompted us to test the consequences of pharmacological inhibition of tyrosinase, a key

enzyme responsible for eumelanin production, at a dose commonly used in zebrafish

research to block pigment formation. Inhibition of tyrosinase via PTU treatment resulted

in an increase of the aorta diameter while a decreased aortic diameter in qKO at 10 dpf

is associated with upregulated tyrosinase expression. This strongly supports a role for

the pigment biosynthesis pathway in aortic homeostasis. Therefore, the use of PTU for

imaging purposes should be approached with caution due to the potential for

unintended effects on other organ systems, especially when evaluating cardiovascular

phenotypes. Undesired effects of PTU have been reported previously including

autophagy activation[63], activation of cytochrome P4501A1 (*cyp1a1*)[64], suppression of

retinol-binding protein 4 (*rbp4*)[65] and reduction of the diameter of the eye[66].

Despite upregulation of melanin-producing pathways, no increase of skin pigmentation

could be observed in the embryo or adult qKO zebrafish. Previous research showed

that tyrosinase follows a specific spatiotemporal expression pattern during

embryogenesis, which is not confined to skin cells[67]. At 7 dpf, small amounts of melanin

can already be detected around the dorsal aorta, while at 1 month, melanin deposition

is apparent[68]. It is therefore tempting to speculate that pigmentation-related pathways

affect vascular development, particularly since it was recently shown that tyrosinase

reduces expression of vascular endothelial growth factors[69].

The relevance of the qKO model to study the disease mechanisms of dissection is

further supported by the transcriptomic changes concordant with established models

and known pathogenetic mechanisms of TAD. Altered elastic fiber homeostasis, which

was confirmed in the qKO aortic wall using TEM, is reflected in upregulation of *Fbn2b*[70]

(logFC 1.04), a fibrillin known to be involved in endocardial morphogenesis, which might

be upregulated as a response to the coarctation in the qKO, and downregulation of

*mfap4* (logFC -3.26), an important component of the extracellular matrix (ECM) involved

in elastic fiber assembly. In zebrafish, MFAP4 also regulates the balance between

myeloid and lymphoid development[71]. Another immune cell-related gene expressed

differently in qKO zebrafish is *itgam* (LogFC -5.33), coding for a component of the

heterodimeric $\alpha_M\beta_2$ integrin expressed on leukocytes which is also known as

macrophage-1 antigen (Mac-1) or CD11b. This protein plays a crucial role in binding to

ECM components and intracellular adhesion molecules, involved in adhesion and

transmigration of leukocytes across blood vessels. The qKO transcriptomic profile also

indicates that matrix proteolysis is increased, due to downregulation of the

metalloproteinase inhibitor *timp4* (LogFC -3.28), in combination with upregulation of

*mmp11b* (logFC 1.34) and *mmp13a* (logFC 1.35). These targets are all widely

established in the pathogenesis of vascular and bone homeostasis [72–74]. *Loxl3a*, an

orthologue of the human *LOXL3*[75] important for crosslinking of elastin and collagen, is

significantly downregulated in qKO zebrafish. In mice, deletion of *Loxl3* results in

abnormal skeletal development and is expressed in the precursors of the occipital and

interparietal bones and nasal area, affected in the qKO model[76].

In conclusion, Smad3-deficiency modifies Smad6-deficient phenotypes whereby

*smad3a$^{-/-}$;smad3b$^{-/-}$;smad6a$^{-/-}$;smad6b$^{-/-}$* qKO zebrafish model aortic dissection and

rupture, resulting from defective vasculogenesis and resultant local wall stress. In

addition to known signatures of TAAD in human and mouse models, this model

implicates the pigmentation pathway in the development of the aortic phenotypes.

Follow-up studies on the precise contribution of these pathways are warranted and are

likely to lead to novel targets for therapeutic intervention in TAAD.

## ACKNOWLEDGMENTS

The authors thank the family for their participation in this research. We also would like to thank the Zebrafish Facility Ghent Core at Ghent University, and particularly Karen Vermeulen for the diligent care for the zebrafish. We acknowledge the Paul Scherrer Institut (PSI), Villigen, Switzerland for provision of synchrotron radiation beamtime at the TOMCAT beamline X02DA of the Swiss Light Source, and would like to thank Karo De Rycke, Violette Deleeuw, Marina Horvat, Yousof Mohammad Asaad Abdel-Raouf (Ghent University) and Isaac Rodriguez Rovira (University of Barcelona) for their assistance in acquiring the synchrotron imaging data. Finally, we would like to thank the Nematology Research Unit, member of the UGent TEM Core Facility and the sequencing core of UZ Ghent.

## SOURCES OF FUNDING

This work was supported by a Concerted Research Action grant from the Ghent University Special Research Fund (grant number BOF GOA019-21) to BC and PS, A research project of the Research Foundation – Flanders (grant number G0B4920N) and the Marfan Foundation Innovators Grant Program (grant number INT.DIV.2023.0008.01) to BC. Bert Callewaert is a senior clinical investigator of the Research Foundation-Flanders.

## DISCLOSURES

None

## DATA AVAILABILITY

Raw RNAseq data is available at Gene Expression Omnibus with accession number

GSE249792**.** To review GEO accession GSE249792. For reviewers, following link is

available to access the data.

https://eur03.safelinks.protection.outlook.com/?url=https%3A%2F%2Fwww.ncbi.nlm.nih.

gov%2Fgeo%2Fquery%2Facc.cgi%3Facc%3DGSE249792&data=05%7C02%7Cmichie

l.vanhooydonck%40ugent.be%7Cbeba6c62b40641ced96408dbfa282a45%7Cd7811cde

ecef496c8f91a1786241b99c%7C1%7C0%7C638378822642521272%7CUnknown%7C

TWFpbGZsb3d8eyJWIjoiMC4wLjAwMDAiLCJQIjoiV2luMzIiLCJBTiI6Ik1haWwiLCJXVCI

6Mn0%3D%7C3000%7C%7C%7C&sdata=7g1JeMJOvj4lZVmQ6FVdkN%2FA2zrRUcu

Vlw6Hcear5vs%3D&reserved=0

Enter token obmhwiqkbbsprkt into the box.

All other data will be made available upon request.

## AUTHOR CONTRIBUTIONS

Conceptualization: MVH., PSI., BC. Methodology: MVH., MVE., MSS., LP., AKB., DS.,

PSI., BC. Software: MSS., MVI. Formal analysis: MVH., MSV., MVI., DS. Investigation:

MVH., MVE., MVI., MSS., LP., AKB., HDS., LC., PT., DS. Resources: PSI., BC. Data

Curation: MVH., MSS. Writing (original draft): MVH., MVE. Writing (review & editing):

MVH., MVE., MSS., LP., AKB, MVI., HDS., LC., PT., AB., P.SE., ADC., AW., DS., PSI.,

BC. Visualization: MVH. MVE. Supervision: PSI., BC. Funding acquisition: PSI., BC.

## SUPPLEMENTARY MATERIAL

Supplementary Case Presentation

Supplementary Methods

Supplementary References

Supplementary Figures 1 - 17

Supplementary Tables 1 - 7

Supplementary Movies 1 - 19

## REFERENCES

1.      Lemaire SA, Russell L. Epidemiology of thoracic aortic dissection. *Nature reviews. Cardiology*. 2011;8(2):103–113.

2.      Van De Laar IMBH, Oldenburg RA, Pals G, Roos-Hesselink JW, De Graaf BM, Verhagen JMA, Hoedemaekers YM, Willemsen R, Severijnen LA, Venselaar H, Vriend G, Pattynama PM, Collée M, Majoor-Krakauer D, Poldermans D, et al. Mutations in SMAD3 cause a syndromic form of aortic aneurysms and dissections with early-onset osteoarthritis. *Nature Genetics*. 2011;43(2):121–126.

3.      Deleeuw V, Carlson E, Renard M, Zientek KD, Wilmarth PA, Reddy AP, Manalo EC, Tufa SF, Keene DR, Olbinado M, Stampanoni M, Kanki S, Yanagisawa H, Mosquera LM, Sips P, et al. Unraveling the role of TGFβ signaling in thoracic aortic aneurysm and dissection using Fbn1 mutant mouse models. *Matrix biology : journal of the International Society for Matrix Biology*. 2023;123:17–33.

4.      Doyle JJ, Gerber EE, Dietz HC. Matrix-dependent perturbation of TGFβ signaling and disease. *FEBS Letters*. 2012;586(14):2003–2015.

5.      Li W, Li Q, Jiao Y, Qin L, Ali R, Zhou J, Ferruzzi J, Kim RW, Geirsson A, Dietz HC, Offermanns S, Humphrey JD, Tellides G. Tgfbr2 disruption in postnatal smooth muscle impairs aortic wall homeostasis. *The Journal of Clinical Investigation*. 2014;124(2):755–767.

6.      Bellini C, Bersi MR, Caulk AW, Ferruzzi J, Milewicz DM, Ramirez F, Rifkin DB, Tellides G, Yanagisawa H, Humphrey JD. Comparison of 10 murine models reveals a distinct biomechanical phenotype in thoracic aortic aneurysms. *Journal of the Royal Society Interface*. 2017;14(130):20161036.

7.      Schmierer B, Hill CS. TGFbeta-SMAD signal transduction: molecular specificity and functional flexibility. *Nature reviews. Molecular cell biology*. 2007;8(12):970–982.

8.      Miyazawa K, Miyazono K. Regulation of TGF-β Family Signaling by Inhibitory Smads. *Cold Spring Harbor perspectives in biology*. 2017;9(3)::a022095.

9.      Liu Y, Festing M, Thompson JC, Hester M, Rankin S, El-Hodiri HM, Zorn AM, Weinstein M. Smad2 and Smad3 coordinately regulate craniofacial and endodermal development. *Developmental Biology*. 2004;270(2):411–426.

10.     Timberlake AT, Choi J, Zaidi S, Lu Q, Nelson-Williams C, Brooks ED, Bilguvar K, Tikhonova I, Mane S, Yang JF, Sawh-Martinez R, Persing S, Zellner EG, Loring E, Chuang C, et al. Two locus inheritance of non-syndromic midline

craniosynostosis via rare SMAD6 and common BMP2 alleles. *eLife*.
2016;5(September2016):e20125.

11.    Yang Y, Zheng Y, Li W, Li L, Tu M, Zhao L, Mei H, Zhu G, Zhu Y. SMAD6
is frequently mutated in nonsyndromic radioulnar synostosis. *Genetics in*
*medicine : official journal of the American College of Medical Genetics*.
2019;21(11):2577–2585.

12.    Tan HL, Glen E, Töpf A, Hall D, O'Sullivan JJ, Sneddon L, Wren C, Avery
P, Lewis RJ, ten Dijke P, Arthur HM, Goodship JA, Keavney BD. Nonsynonymous
variants in the SMAD6 gene predispose to congenital cardiovascular
malformation. *Human Mutation*. 2012;33(4):720–727.

13.    Luyckx I, MacCarrick G, Kempers M, Meester J, Geryl C, Rombouts O,
Peeters N, Claes C, Boeckx N, Sakalihasan N, Jacquinet A, Hoischen A,
Vandeweyer G, Van Lent S, Saenen J, et al. Confirmation of the role of
pathogenic SMAD6 variants in bicuspid aortic valve-related aortopathy. *European*
*Journal of Human Genetics*. 2019;27(7):1044–1053.

14.    Luyckx I, Verstraeten A, Goumans MJ, Loeys B. SMAD6-deficiency in
human genetic disorders. *NPJ Genomic Medicine*. 2022;7(1):68.

15.    Kloth K, Bierhals T, Johannsen J, Harms FL, Juusola J, Johnson MC,
Grange DK, Kutsche K. Biallelic variants in SMAD6 are associated with a complex
cardiovascular phenotype. *Human genetics*. 2019;138(6):625–634.

16.    Rajderkar S, Barozzi I, Zhu Y, Hu R, Zhang Y, Li B, Alcaina Caro A,
Fukuda-Yuzawa Y, Kelman G, Akeza A, Blow MJ, Pham Q, Harrington AN,
Godoy J, Meky EM, et al. Topologically associating domain boundaries are
required for normal genome function. *Communications Biology*. 2023;6:435.

17.    Giehl KA, Eckstein GN, Pasternack SM, Praetzel-Wunder S, Ruzicka T,
Lichtner P, Seidl K, Rogers M, Graf E, Langbein L, Braun-Falco M, Betz RC,
Strom TM. Nonsense mutations in AAGAB cause punctate palmoplantar
keratoderma type Buschke-Fischer-Brauer. *American journal of human genetics*.
2012;91(4):754–759.

18.    Glasauer SMK, Neuhauss SCF. Whole-genome duplication in teleost
fishes and its evolutionary consequences. *Molecular genetics and genomics :*
*MGG*. 2014;289(6):1045–1060.

19.    Pearson WR. An Introduction to Sequence Similarity ("Homology")
Searching. *Current protocols in bioinformatics*. 2013;chapter 3:3.1.1-3.1.8.

20.    Jia S, Ren Z, Li X, Zheng Y, Meng A. smad2 and smad3 Are Required for
Mesendoderm Induction by Transforming Growth Factor-/Nodal Signals in
Zebrafish. *Journal of Biological Chemistry*. 2008;283(4):2418–2426.

21.    Casari A, Schiavone M, Facchinello N, Vettori A, Meyer D, Tiso N, Moro
E, Argenton F. A Smad3 transgenic reporter reveals TGF-beta control of zebrafish
spinal cord development. *Developmental biology*. 2014;396(1):81–93.

22.    Pogoda HM, Meyer D. Zebrafish smad7 is regulated by Smad3 and BMP
signals. *Developmental Dynamics*. 2002;224(3):334–349.

23.    Lawrence C, Sanders GE, Varga ZM, Baumann DP, Freeman A, Baur B,
Francis M. Regulatory compliance and the Zebrafish. *Zebrafish*. 2009;6(4):453–
456.

24.    Westerfield M, Doerry E, Kirkpatrick AE, Douglas SA. Chapter 19
Zebrafish Informatics and the ZFIN Database. *Methods in Cell Biology*.
1998;60(C):339–355.

25.    Naito Y, Hino K, Bono H, Ui-Tei K. CRISPRdirect: software for designing
CRISPR/Cas guide RNA with reduced off-target sites. *Bioinformatics*.
2015;31(7):1120.

26.    Boel A, De Saffel H, Steyaert W, Callewaert B, De Paepe A, Coucke PJ,
Willaert A. CRISPR/Cas9-mediated homology-directed repair by ssODNs in
zebrafish induces complex mutational patterns resulting from genomic integration
of repair-template fragments. *DMM Disease Models and Mechanisms*.
2018;11(10):dmm035352.

27.    Boel A, Steyaert W, De Rocker N, Menten B, Callewaert B, De Paepe A,
Coucke P, Willaert A. BATCH-GE: Batch analysis of Next-Generation Sequencing
data for genome editing assessment. *Scientific Reports*. 2016;6(July):30330.

28.    Pottie L, van Gool W, Vanhooydonck M, Hanisch FG, Goeminne, Rajkovic
A, Coucke P, Sips P, Callewaert B. Loss of zebrafish atp6v1e1b, encoding a
subunit of vacuolar ATPase, recapitulates human ARCL type 2C syndrome and
identifies multiple pathobiological signatures. *PLoS genetics*.
2021;17(6):e1009603.

29.    Saalfeld S, Cardona A, Hartenstein V, Tomančák P. As-rigid-as-possible
mosaicking and serial section registration of large ssTEM datasets.
*Bioinformatics*. 2010;26(12):i57–i63.

30.    Saalfeld S, Fetter R, Cardona A, Tomancak P. Elastic volume
reconstruction from series of ultra-thin microscopy sections. *Nature methods*.
2012;9(7):717–720.

31.    Cardona A, Saalfeld S, Schindelin J, Arganda-Carreras I, Preibisch S,
Longair M, Tomancak P, Hartenstein V, Douglas RJ. TrakEM2 software for neural
circuit reconstruction. *PloS one*. 2012;7(6):e38011.

32.    Van Impe M, Caboor L, Deleeuw V, Rycke K De, Vanhooydonck M,
Backer J De, Segers P, Sips P. Application of an automated analysis framework
for pulsed-wave Doppler cardiac ultrasound measurements to generate reference
data in adult zebrafish. *American Journal of Physiology*. 2023;325(6):R782–R796.

33.    Ramsay JM, Feist GW, Varga ZM, Westerfield M, Kent ML, Schreck CB.
Whole-body cortisol response of zebrafish to acute net handling stress.
*Aquaculture*. 2009;297(1–4):157–162.

34.    Jarayseh T, Guillemyn B, De Saffel H, Bek JW, Syx D, Symoens S,
Gansemans Y, Van Nieuwerburgh F, Jagadeesh S, Raja J, Malfait F, Coucke PJ,
De Clercq A, Willaert A. A tapt1 knock-out zebrafish line with aberrant lens
development and impaired vision models human early-onset cataract. *Human
Genetics*. 2023;142(3):457–476.

35.    Bird NC, Mabee PM. Developmental morphology of the axial skeleton of
the zebrafish, Danio rerio (Ostariophysi: Cyprinidae). *Developmental Dynamics*.
2003;228(3):337–357.

36.    Parichy DM, Elizondo MR, Mills MG, Gordon TN, Engeszer RE. Normal
table of postembryonic zebrafish development: Staging by externally visible
anatomy of the living fish. *Developmental Dynamics*. 2009;238(12):2975–3015.

37.    Andrews S. Babraham Bioinformatics - FastQC A Quality Control tool for
High Throughput Sequence Data. *FastQC A Quality Control tool for High
Throughput Sequence Data*. 2010.

38.    Krueger F, James F, Ewels P, Afyounian E, Schuster-Boeckler B.
FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo. 2021.

39.    Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P,
Chaisson M, Gingeras TR. STAR: ultrafast universal RNA-seq aligner.
*Bioinformatics*. 2013;29(1):15–21.

40.    Sayols S, Scherzinger D, Klein H. dupRadar: a Bioconductor package for
the assessment of PCR artifacts in RNA-Seq data. *BMC bioinformatics*.
2016;17(1):428.

41.    Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics*.
2016;32(19):3047–3048.

42.    Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics*.
2014;30(7):923–930.

43.    Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics (Oxford, England)*. 2010;26(1):139–140.

44.    Chen Y, Lun ATL, Smyth GK. From reads to genes to pathways: Differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. *F1000Research*. 2016;5.

45.    Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics : a journal of integrative biology*. 2012;16(5):284–287.

46.    Van Impe M, Caboor L, Deleeuw V, Olbinado M, De Backer J, Sips P, Segers P. Fluid-Structure Interaction Modeling of the Aortic Hemodynamics in Adult Zebrafish: a Pilot Study based on Synchrotron X-Ray Tomography. *IEEE Transactions on Biomedical Engineering*. 2023;70(7):2101–2110.

47.    Paganin D, Mayo SC, Gureyev TE, Miller PR, Wilkins SW. Simultaneous phase and amplitude extraction from a single defocused image of a homogeneous object. *Journal of Microscopy*. 2002;206(1):33–40.

48.    Marone F, Stampanoni M. Regridding reconstruction algorithm for real-time tomographic imaging. *Journal of synchrotron radiation*. 2012;19(Pt 6):1029–1037.

49.    Hsiao JJ, Fisher DE. The roles of microphthalmia-associated transcription factor and pigmentation in melanoma. *Archives of Biochemistry and Biophysics*. 2014;563:28–34.

50.    Bian C, Li R, Wen Z, Ge W, Shi Q. Phylogenetic Analysis of Core Melanin Synthesis Genes Provides Novel Insights Into the Molecular Basis of Albinism in Fish. *Frontiers in Genetics*. 2021;12:707228.

51.    Casari A, Schiavone M, Facchinello N, Vettori A, Meyer D, Tiso N, Moro E, Argenton F. A Smad3 transgenic reporter reveals TGF-beta control of zebrafish spinal cord development. *Developmental Biology*. 2014;396(1):81–93.

52.    Pogoda HM, Meyer D. Zebrafish smad7 is regulated by Smad3 and BMP signals. *Developmental Dynamics*. 2002;224(3):334–349.

53.    De Pater E, Ciampricotti M, Priller F, Veerkamp J, Strate I, Smith K, Lagendijk AK, Schilling TF, Herzog W, Abdelilah-Seyfried S, Hammerschmidt M, Bakkers J. Bmp signaling exerts opposite effects on cardiac differentiation. *Circulation Research*. 2012;110(4):578–587.

54.    Mouillesseaux KP, Wiley DS, Saunders LM, Wylie LA, Kushner EJ, Chong DC, Citrin KM, Barber AT, Park Y, Kim JD, Samsa LA, Kim J, Liu J, Jin SW,

Bautch VL. Notch regulates BMP responsiveness and lateral branching in vessel networks via SMAD6. *Nature Communications*. 2016;7:13247.

55.     Boezio GLM, Bensimon-Brito A, Piesker J, Guenther S, Helker CSM, Stainier DYR. Endothelial TGF-β signaling instructs smooth muscle cell development in the cardiac outflow tract. *eLife*. 2020;9:1–32.

56.     Lucarelli P, Schilling M, Kreutz C, Vlasov A, Boehm ME, Iwamoto N, Steiert B, Lattermann S, Wäsch M, Stepath M, Matter MS, Heikenwälder M, Hoffmann K, Deharde D, Damm G, et al. Resolving the Combinatorial Complexity of Smad Protein Complex Formation and Its Link to Gene Expression. *Cell Systems*. 2018;6(1):75-89.e11.

57.     Wang X, LeMaire SA, Chen L, Shen YH, Gan Y, Bartsch H, Carter SA, Utama B, Ou H, Coselli JS, Wang XL. Increased Collagen Deposition and Elevated Expression of Connective Tissue Growth Factor in Human Thoracic Aortic Dissection. *Circulation*. 2006;114(SUPPL. 1):I200–I205.

58.     Bechtel JFM, Misfeld M, Schmidtke C, Sievers HH. The bicuspid aortic valve. In: *Aortic Root Surgery*.; 2010:89–101.

59.     Yang J, Wang J, Pan L, Li H, Rao C, Zhang X, Niu G, Qu J, Hou L. BMP4 is required for the initial expression of MITF in melanocyte precursor differentiation from embryonic stem cells. *Experimental cell research*. 2014;320(1):54–61.

60.     Massaous J, Hata A. TGF-B signalling through the Smad pathway. *Cell Biology*. 1997;7:187–192.

61.     Steinfeld J, Steinfeld I, Coronato N, Hampel ML, Layer PG, Araki M, Vogel-Höpker A. RPE specification in the chick is mediated by surface ectoderm-derived BMP and Wnt signalling. *Development*. 2013;140(24):4959–4969.

62.     Goding CR, Arnheiter H. Mitf—the first 25 years. *Genes and Development*. 2019;33(15–16):983–1007.

63.     Chen XK, Kwan JSK, Chang RCC, Ma ACH. 1-phenyl 2-thiourea (PTU) activates autophagy in zebrafish embryos. *Autophagy*. 2021;17(5):1222–1231.

64.     Wang W Der, Wang Y, Wen HJ, Buhler DR, Hu CH. Phenylthiourea as a weak activator of aryl hydrocarbon receptor inhibiting 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced CYP1A1 transcription in zebrafish embryo. *Biochemical Pharmacology*. 2004;68(1):63–71.

65.     Tingaud-Sequeira A, Forgue J, André M, Babin PJ. Epidermal transient down-regulation of retinol-binding protein 4 and mirror expression of apolipoprotein Eb and estrogen receptor 2a during zebrafish fin and scale development. *Developmental Dynamics*. 2006;235(11):3071–3079.

66.     Li Z, Ptak D, Zhang L, Walls EK, Zhong W, Leung YF. Phenylthiourea
Specifically Reduces Zebrafish Eye Size. *PLOS ONE*. 2012;7(6):e40132.

67.     Camp E, Lardelli M. Tyrosinase gene expression in zebrafish embryos.
*Development genes and evolution*. 2001;211(3):150–153.

68.     Miano JM, Georger MA, Rich A, De Mesy Bentley KL. Ultrastructure of
zebrafish dorsal aortic cells. *Zebrafish*. 2006;3(4):455–463.

69.     Clahsen T, Hatami N, Büttner C, Reis A, Cursiefen C. Tyrosinase reduces
expression of vascular endothelial growth factors and improves corneal graft
survival. *Investigative Ophthalmology & Visual Science*. 2022;63(7):909-A0273.

70.     Mellman K, Huisken J, Dinsmore C, Hoppe C, Stainier DY. Fibrillin-2b
regulates endocardial morphogenesis in zebrafish. *Developmental biology*.
2012;372(1):111–119.

71.     Ong SLM, De Vos IJHM, Meroshini M, Poobalan Y, Dunn & NR.
Microfibril-associated glycoprotein 4 (Mfap4) regulates haematopoiesis in
zebrafish. 2020;10:11801.

72.     Koskivirta I, Rahkonen O, Mäyränpää M, Pakkanen S, Husheem M, Sainio
A, Hakovirta H, Laine J, Jokinen E, Vuorio E, Kovanen P, Järveläinen H. Tissue
inhibitor of metalloproteinases 4 (TIMP4) is involved in inflammatory processes of
human cardiovascular pathology. *Histochemistry and cell biology*.
2006;126(3):335–342.

73.     Johansson N, Ahonen M, Kähäri VM. Matrix metalloproteinases in tumor
invasion. *Cellular and Molecular Life Sciences*. 2000;10(6):415–433.

74.     Jones JA, Ikonomidis JS. The Pathogenesis of Aortopathy in Marfan
Syndrome and Related Diseases. *Current cardiology reports*. 2010;12(2):99–107.

75.     Van Boxtel AL, Gansner JM, Hakvoort HWJ, Snell H, Legler J, Gitlin JD.
Lysyl oxidase-like 3b is Critical for Cartilage Maturation During Zebrafish
Craniofacial Development. *Matrix biology : journal of the International Society for*
*Matrix Biology*. 2011;30(2001):178–187.

76.     Santamaría PG, Dubus P, Bustos-tauler J, Floristán A, Vázquez-naharro
A, Morales S, Cano A, Portillo F. Loxl2 and Loxl3 Paralogues Play Redundant
Roles during Mouse Development. *International Journal of Molecular Sciences*.
2022;23(10):5730.

## FIGURES WITH FIGURE LEGENDS

808 Links with access to high quality figures and the supplementary movies. Lower quality

809 figures can be found below:

| | |
|---|---|
| Figure 1 | https://figshare.com/s/58540e31ae5dcfed3750 |
| Figure 2 | https://figshare.com/s/b08f40d0b6c136b31ee4 |
| Figure 3 | https://figshare.com/s/9fbc19810756ed1b34b4 |
| Figure 4 | https://figshare.com/s/07b6703f58ea0653336e |
| Figure 5 | https://figshare.com/s/2b913da4b07559cc0607 |
| Figure 6 | https://figshare.com/s/42bdead6d4b916ec7327 |
| Supplementary Figure 1 | https://figshare.com/s/1d96715da2007c11ae96 |
| Supplementary Figure 2 | https://figshare.com/s/7117068f173e59e3b7e7 |
| Supplementary Figure 3 | https://figshare.com/s/8a7ce09585ca754dedd1 |
| Supplementary Figure 4 | https://figshare.com/s/e754b6c96ea2c0588a15 |
| Supplementary Figure 5 | https://figshare.com/s/c902712aae5ffea8e862 |
| Supplementary Figure 6 | https://figshare.com/s/9bcafde922edd89090ad |
| Supplementary Figure 7 | https://figshare.com/s/00039d2f1472d980fb15 |
| Supplementary Figure 8 | https://figshare.com/s/19aa5c0c33dfbf52f76b |
| Supplementary Figure 9 | https://figshare.com/s/3b70b2174c355aafa4c8 |
| Supplementary Figure 10 | https://figshare.com/s/2ed90596b1477259fc3f |
| Supplementary Figure 11 | https://figshare.com/s/5c8b0c3c23600bd54b15 |
| Supplementary Figure 12 | https://figshare.com/s/2c1733efb6b877eed39a |
| Supplementary Figure 13 | https://figshare.com/s/2a70a063f0dbda9c8bac |
| Supplementary Figure 14 | https://figshare.com/s/58a4c79c1b31b4474c1c |
| Supplementary Figure 15 | https://figshare.com/s/5ccc12c8bf07a477ab19 |
| Supplementary Figure 16 | https://figshare.com/s/64b5c5461c87858830e5 |
| Supplementary Figure 17 | https://figshare.com/s/a950dec1eb4b32f8559a |
| Supplementary Movie 1 | https://figshare.com/s/5f3e5147898632f68274 |
| Supplementary Movie 2 | https://figshare.com/s/cdf6d880d1a241c750f4 |
| Supplementary Movie 3 | https://figshare.com/s/5b6d3fb558aeb40771d2 |
| Supplementary Movie 4 | https://figshare.com/s/f2c26de3a7403d10148c |
| Supplementary Movie 5 | https://figshare.com/s/03f83afad3ad6383aa20 |
| Supplementary Movie 6 | https://figshare.com/s/51aa1e42b4cb45ecdc60 |
| Supplementary Movie 7 | https://figshare.com/s/415c7431de05ecfc5449 |
| Supplementary Movie 8 | https://figshare.com/s/8a2169e10ff223bc971d |
| Supplementary Movie 9 | https://figshare.com/s/a3aaebcb00ccb50f9b4d |
| Supplementary Movie 10 | https://figshare.com/s/84ac6e35c61e47142339 |
| Supplementary Movie 11 | https://figshare.com/s/b87dfe1d640a284edf86 |
| Supplementary Movie 12 | https://figshare.com/s/f74f27f17b2cb7343559 |
| Supplementary Movie 13 | https://figshare.com/s/b9f1f922f31ada373b87 |
| Supplementary Movie 14 | https://figshare.com/s/9a1291cd30650944a05f |
| Supplementary Movie 15 | https://figshare.com/s/3a08c958c980bb16c948 |

| Supplementary Movie 16 | https://figshare.com/s/5cd90d5c165d52dfa319 |
| Supplementary Movie 17 | https://figshare.com/s/3bac7c34e759d5849442 |
| Supplementary Movie 18 | https://figshare.com/s/db1b70cf18c9fe058a57 |
| Supplementary Movie 19 | https://figshare.com/s/769438c9b5a3cfade65d |

**Figure 1: Altered vasculogenesis at 10 dpf in double and qKO mutants. (a-d)** Ventral view of the cardiovascular structures at 10 dpf of *smad* knockout lines crossed with a *Tg(fli1:EGFP)* reporter line. Legend: "VA" ventral aorta, "BA" bulbus arteriosus, "AA3" aortic arch 3, "AA4" aortic arch 4, "HL" head length. The projected surface area of the ventral aorta between aortic arch 3 and aortic arch 4 has been outlined with a white dashed line. **(e)** *smad3a⁺ᐟ⁻;smad3b⁻ᐟ⁻ and smad3a⁻ᐟ⁻;smad3b⁻ᐟ⁻* zebrafish of 10 dpf show an increase in normalized aorta diameter (n=29-11-7). **(f)** *smad6a⁺ᐟ⁻;smad6b⁻ᐟ⁻* as well as *smad6a⁻ᐟ⁻;smad6b⁻ᐟ⁻* show a decrease in diameter of the ventral aorta (n=15-18-14-10-23). **(g)** *smad3a⁺ᐟ⁻;smad3b⁻ᐟ⁻;smad6a⁺ᐟ⁻;smad6b⁻ᐟ⁻* and *smad3a⁻ᐟ⁻;smad3b⁻ᐟ⁻;smad6a⁻ᐟ⁻;smad6b⁻ᐟ⁻* both show a decrease in diameter of the ventral aorta (n=15-18-26). **(h,i)**

Overview of the hypobranchial artery in a WT control and qKO zebrafish line. Tortuosity

and irregular branching of the hypobranchial artery (HA) and aortic arch 1 (AA1) can be

observed **(i)** magnification of boxed area in d. **(a-d,h,i)** stack focused Z-stack pictures

obtained with ZEISS Axio Observer.Z1 microscope **(f)** One-way ANOVA with Dunnett's

multiple comparison test against WT controls. **(e,g)** Brown-Forsythe and Welch ANOVA

with Dunnett's T3 multiple comparison test against WT controls. Asterisks indicate

significant differences. $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$. Data represented

as average ± standard deviation.

**Figure 2: qKO zebrafish exhibit thoracic aortic dissection and rupture. (a, b)** qKO

zebrafish that died suddenly show similar external features compared to qKO zebrafish

that die due to stress induction. Red discoloration around the heart and gill area

(anterior abdominal area) is indicated with a white arrowhead. **(c)** Resorcin-Fuchsin

staining of WT control. **(d)** Resorcin-Fuchsin staining of the elastic fiber on a 5 µm

cross-section of the ventral aorta shows rupture of the aortic wall (red arrow). **(e,f)**

Resorcin-Fuchsin staining of the elastic fiber on 5 µm thick cross-sections of the ventral

aorta. False lumen is indicated with a black arrow. **(g,h)** 3D reconstruction using 3D-

modelling software Mimics 24.0 of the ventral aorta shows a decrease in diameter of the

aortic arches. Symmetrical branching of the aortic arches is lost in the qKO model.

Region with aortic damage has been colored black **(i,j).** The area of damage identified

on a histological resorcin-fuchsin staining of the ventral aorta can similarly be observed

on a synchrotron micro-CT scan of the same region. Damage is indicated with arrows.

**(k,l)** TEM of aortic wall in WT and qKO, respectively. Severe decrease of elastic fiber in

the inner elastic lamina is apparent. **(m)** TEM of qKO shows collagen deposition in

intima. **(k,l,m)** Legend: "VSMC" vascular smooth muscle cell, "EC" endothelial cell,

"RBC" red blood cell, "C" collagen, * indicates inner elastic lamina.

**Figure 3: Increased regurgitation and abnormal valve morphology in qKO zebrafish. (a, c)** Pulsed wave doppler of inflow **(a)** and outflow **(c)** of blood in and out of the ventricle of WT control zebrafish. **(b, d)** Pulsed wave doppler of inflow **(b)** and outflow **(d)** of blood in and out of the ventricle of qKO zebrafish. **(e)** Regurgitation on inflow is significantly increased in the qKO zebrafish model (n=17-8-7-8-8-10). **(f, g)**

Resorcin Fuchsin staining shows hypertrophy of the valve interstitial cells and abnormal

leaflet morphology in the atrioventricular valve of qKO zebrafish. **(h)** Increased

regurgitation on outflow in qKO (n=17-7-7-7-7-10). **(i,j)** Resorcin Fuchsin staining of the

bulboventricular valve in WT and qKO zebrafish shows hypertrophy of the valve and

altered orientation in the qKO. **(a-j)** Legend: "RI" regurgitation inflow, "RO" regurgitation

outflow, "A" atrium, "V" ventricle, "B" bulbus arteriosus. Black arrows indicate valves.

**(e,h)** One-way ANOVA with Dunnett's multiple comparison test against WT controls.

Asterisks indicate significant differences. ***$p<0.001$, ****$p<0.0001$. Data represented as

average ± standard deviation.

**Figure 4: qKO zebrafish show scoliosis, lordosis and ectopic bone formation in the vertebral column. (a-l)** Overview pictures of alizarin red staining for mineralized bone of abdominal and caudal vertebrae, and the skull (lateral view) of WT (**a-d**), *smad3a⁻/⁻;smad3b⁻/⁻* DKO zebrafish (**e-h**), and *smad3a⁻/⁻;smad3b⁻/⁻;smad6a⁻/⁻;smad6b⁻/⁻* qKO zebrafish (**i-l**). Ectopic bone has been indicated with white arrowheads. Scoliosis and lordosis can be observed in (**i**). **(k)** Notochord sheet mineralization can also be observed between the vertebrae. Small craniofacial aberrations are detected. **(l)** The nasal bone structures appear to be missing, resulting in a clockwise rotation of the snout. **(m-o)** Quantification of spots of ectopic bone in *smad3* (n=10-10-8-10-10-4),

877     *smad6* (n=10-4-3-8-5), *smad3/smad6* (n=10-6-13-14-15-8) genotypes. **(a-l)** Whole-

878     mount alizarin red staining for mineralized bone pictures taken with Leica M165 FC

879     Fluorescent Stereo Microscope. **(m-o)** Kruskal-Wallis test with Dunn's multiple

880     comparisons test against WT controls. $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$.

881     Data represented as average ± standard deviation.

**Figure 5: RNA sequencing data shows downregulation of negative regulators of proteolysis and upregulation of melanogenesis. (a)** Volcano Plot showing upregulated genes (red) and downregulated genes (blue) with genes of importance annotated. Thresholds: p.adjust ≥ 0.05 and |log2FC| ≥ 1; **(b)** Heatmap with top DEG results between WT and qKO. Thresholds: p.adjust ≥ 0.0001 and |log2FC| ≥ 1 ; **(c)** Top 10 hits of GO enrichment analysis for both sets of upregulated (red) and downregulated (blue) genes. Numbers inside the bars represent the corresponding adjusted p-value value.

**Figure 6: Tyrosinase inhibition increases the aorta diameter of 10 dpf zebrafish.**

**(a)** Ventral view of the vasculature of 10 dpf WT control zebrafish. **(b)** Ventral view of

the vasculature of 10 dpf WT zebrafish treated with 0.003% PTU treatment starting at

24 hpf. **(c, d)** PTU-treated zebrafish show a significant increase in normalized surface

area of the aortic segment and aortic diameter (n=15-16). **(e,f)** Normalized diameter

adjacent to the bulbus arteriosus and length of the aorta segment is similar between

treated and non-treated zebrafish (n=15-16). **(a,b)** Dashed white line indicates aorta

segment between aortic arch 3 and 4. Stack focused Z-stack pictures obtained with

ZEISS Axio Observer.Z1 microscope**. (c,d)** Unpaired two-tailed t-test with Welch's

correction. **(e,f)** Unpaired two-tailed t-test. Asterisks indicate significant differences.

**p<0.01. Data represented as average ± standard deviation.