Early mechanisms of aortic failure in a zebrafish model for thoracic aortic dissection and rupture

Michiel Vanhooydonck MSc.[1], Maxim Verlee MD.[1], Marta Santana Silva MSc.[1], Lore

Pottie PhD.[1], Annekatrien Boel PhD.[2], Matthias Van Impe MSc.[3], Hanna De Saffel

BSc.[1], Lisa Caboor BSc.[1], Piyanoot Tapaneeyaphan MSc.[1], Anne Bonnin PhD.[4], Patrick

Segers PhD.[3], Adelbert De Clercq PhD.[5], Andy Willaert PhD.[1], Delfien Syx PhD.[1],

Patrick Sips PhD.[1a], Bert Callewaert MD. PhD.[1a*]

[1] Center for Medical Genetics Ghent (CMGG), Department of Biomolecular Medicine,

Ghent University, 9000 Ghent, Belgium

[2] Ghent-Fertility and Stem Cell Team (G-FaST), Department for Reproductive Medicine,

Ghent University, 9000 Ghent, Belgium

[3] Biophysical Models for Medical Applications (bioMMeda), Institute of Biomedical

Engineering and Technology (IBiTech), Ghent University, 9000 Ghent, Belgium.

[4] Paul Scherrer Institut, Swiss Light Source, 5232 Villigen PSI, Switzerland.

[5] Evolutionary Developmental Biology, Biology Department, Ghent University, 9000

Ghent, Belgium.

[a] Bert Callewaert and Patrick Sips contributed equally as senior authors.

*Corresponding Author: Bert Callewaert, PhD, MD

Bert.Callewaert@Ugent.be; ORCID: 0000-0002-9743-4205

Word count: 8888

ABSTRACT

Thoracic aortic aneurysm and dissection (TAAD) associates with a high mortality rate. Despite the existence of different mouse models for TAAD, the underlying disease mechanisms remain elusive. Treatment options are limited and mainly consist of surgical repair at critical aortic diameters as current pharmacological interventions are unable to stop disease progression.

In humans, loss of function (LOF) of *SMAD3* and *SMAD6* impairs vascular homeostasis, increasing the risk for TAAD. We developed a zebrafish model for thoracic aortic dissection/rupture by targeting both ohnologs of *smad3* and *smad6*. At 10 days post fertilization, we found an increased diameter of the ventral aorta in *smad3a*$^{-/-}$;*smad3b*$^{-/-}$ double knockout zebrafish, while *smad6a*$^{-/-}$;*smad6b*$^{-/-}$ double knockout zebrafish have a reduced aortic diameter associated with early mortality. We discovered that a *smad3a*$^{-/-}$;*smad3b*$^{-/-}$;*smad6a*$^{-/-}$;*smad6b*$^{-/-}$ quadruple knockout (qKO) zebrafish model is viable and survives to adulthood, although exposure to stress leads to sudden death. Histological analysis of the adult ventral aorta shows medial elastolysis, aortic dissections and ruptures at sites exposed to high biomechanical stress. RNA-sequencing of 5 days post fertilization qKO zebrafish indicates a profile of reduced negative regulation of proteolysis and upregulation of melanogenesis, a previously unaddressed pathway in this pathology. We confirm that pharmacological modulation of tyrosinase, the enzyme responsible for the production of melanin, influences aortic morphology.

46   Overall, the qKO mutant, thus far the only known zebrafish model of thoracic aortic

47   dissection and rupture, reveals novel SMAD3/6-dependent pathways that impact thoracic

48   aortic homeostasis, in this way opening avenues for the development of novel treatments

49   in TAAD.

50 **NON-STANDARD ABBREVIATIONS AND ACRONYMS**

AA3	aortic arch 3
AA4	aortic arch 4
BMPs	bone morphogenetic proteins
Co-SMAD	common mediator SMAD
CPM	counts per million
DEG	differentially expressed genes
DKO	double knockout
dpf	days post fertilization
ECM	extracellular matrix
FDR	false discovery rate
GDFs	differentiation factors
GO	gene ontology
GRCz11	Genome Reference Consortium Zebrafish Build 11
GSEA	gene set enrichment analysis
ID	intellectual disability
I-SMAD	inhibitory SMAD
KEGG	Kyoto Encyclopedia of Genes and Genomes
LDS3	Loeys-Diets syndrome type 3
LOF	loss of function
mpf	months post fertilization
MR	magnetic resonance
PSI	Paul Scherrer Institut
PTU	1-phenyl-2-thiourea treatment
qKO	smad3a$^{-/-}$;smad3b$^{-/-}$;smad6a$^{-/-}$;smad6b$^{-/-}$ quadruple knockout
RO	reverse osmosis
R-SMAD	receptor-regulated SMAD

SKO	single knockout
smad3a/b DKO	*smad3a$^{-/-}$;smad3b$^{-/-}$* double knockout
smad6a/b DKO	*smad6a$^{-/-}$;smad6b$^{-/-}$* double knockout
TAA	thoracic aortic aneurysm
TAAD	thoracic aortic aneurysm and dissection
TAD	thoracic aortic dissection
TEM	transmission electron microscopy
TGF-β	transforming growth factor β
ZIRC	Zebrafish International Research Center

INTRODUCTION

Thoracic aortic dissection (TAD) affects 3-4 individuals per 100,000 people per year[1]. This life-threatening event is often, but not exclusively, preceded by thoracic aortic aneurysm (TAA). The disease mechanisms are incompletely understood, but evidence from genetic disorders implicates altered cell-matrix interactions (by impaired hemodynamic sensing or impaired matrix assembly), as well as aberrant transforming growth factor β (TGF-β) signaling[2–5]. Despite several mouse models for TAD[6], the initial molecular mechanisms remain obscure.

Members of the TGF-β superfamily include TGF-β proteins, bone morphogenetic proteins (BMPs), activins, and growth and differentiation factors (GDFs). Binding of TGF-β to the tetrameric TGFβ-receptor complex results in phosphorylation of the receptor-regulated (R-) SMAD2 and -3 proteins. Binding of BMP and GDF to the tetrameric BMP-receptor complex results in phosphorylation of R-SMAD1, -5, and -8. Phosphorylated R-SMAD proteins recruit the common mediator (Co-) SMAD4 to form a heterotrimeric complex and translocate to the nucleus to activate the transcription of the TGF-β or BMP target genes[7]. The TGF-β and BMP signaling is negatively regulated by the inhibitory (I-) SMAD6 and -7 proteins. I-SMAD6 primarily suppresses the pathway activated by the BMP-receptor, whereas I-SMAD7 inhibits both the TGF-β and BMP induced pathways[8].

Pathogenic variants in *SMAD3* cause Loeys-Dietz syndrome type 3 (LDS3). LDS3 typically presents with early-onset osteo-arthritis, and arterial aneurysm and dissection, mainly affecting the ascending aorta. Other features include arterial tortuosity, mitral valve prolapse and craniofacial characteristics[2,9]. Pathogenic variants in *SMAD6* are associated with craniosynostosis[10], radio-ulnar synostosis[11] or congenital heart disease including

bicuspid aortic valve[12] with or without TAAD[13] and Shone complex with a hypoplastic

ascending aorta and arch[14]. Biallelic variants in *SMAD6* underly more complex

cardiovascular phenotypes[15]. Interestingly, identical pathogenic variants in *SMAD6* may

result in either cardiovascular anomalies, craniosynostosis or radioulnar synostosis[13].

Because *SMAD3* and *SMAD6* have been in linkage disequilibrium throughout evolution

from teleost fish to human, it has been postulated that *SMAD3* could have a modifier role

on *SMAD6* deficiency[16]. In line with this observation, we identified a 1.5 Mb 15q22.31-

15q23 microdeletion encompassing *SMAD3* and *SMAD6* in an 11-year-old-boy and his

33-year-old mother, both presenting with bicuspid aortic valve and arterial tortuosity, but

no aneurysms, along with skeletal and cutaneous anomalies of LDS3, hinting that deletion

of *SMAD6* may influence the phenotype caused by *SMAD3* LOF (**Supplementary Case**

Presentation[17]). Due to the teleost specific genome duplication, zebrafish have two

ohnolog copies for both *SMAD3* and *SMAD6*[18]. We established zebrafish models deficient

for each of these ohnologs identified as *smad3a* and *smad3b,* and *smad6a* and *smad6b*,

respectively. The human SMAD3 (ENSP00000332973) protein has a sequence similarity

of 97.17% with zebrafish Smad3a (ENSDARP00000045373) and 93.4% with zebrafish

Smad3b (ENSDARP00000043454). SMAD6 (ENSP00000288840) is less conserved with

a similarity of 53.3% and 62.38% with zebrafish Smad6a (ENSDARP00000112408) and

Smad6b (ENSDARP00000091342) respectively. Smad6a and Smad6b show 58.33%

protein sequence similarity, which is considered large enough to assume similar

function[19]. The sequence similarity and the same spatial expression patterns of *Smad3a*

and *Smad3b* as well as *Smad6a* and *smad6b* suggest[20–22].

In this study, we investigated modifying effects of *smad3a/smad3b* on *smad6a/smad6b*

and vice versa which enabled us to model TAAD in zebrafish. Additionally, we identified

early molecular mechanisms in aortic failure to identify druggable pathways.

METHODS

Ethics statement and housing

This study was approved by the Animal Ethics Committee of the Ghent University Faculty of Medicine and Health Sciences (Permit number: ECD 14/70) and housed adhering to the general guidelines, in agreement with EU Directive 2010/63/EU for laboratory animals [23,24].

Zebrafish lines: design and generation

Smad3a[sa2363/+] was acquired from the Zebrafish International Research Center (ZIRC). All other models were generated using CRISPR/Cas9 gene editing technology utilizing CRISPRdirect software[25–27]. Specific gBlocks for all target sites are listed in **Supplementary Table 1**. Genotyping primers are given in **Supplementary Table 2**. All variants are described using Genome Reference Consortium Zebrafish Build 11 (GRCz11) and described in **Supplementary Table 3**.

Vasculogenesis in embryos

Zebrafish were outcrossed to the transgenic Tg(fli1:EGFP) line in order to visualize endothelial cells as previously described[28]. Head length was used as normalization factor. An overview of the acquired measurements and specific breeding schemes are given in **Supplementary Fig. 1a-c**.

3D reconstruction of the ventral aorta in adult zebrafish

Serial 5 µm paraffin cross sections of the entire ventral aorta of zebrafish between 6 and 13 months post fertilization (mpf) were stained with Weigert's iron hematoxylin, rinsed in water and stained with Resorcin-Fuchsin for one hour followed by two wash steps in 95%

122 ethanol and one wash with reverse osmosis (RO) water. Slides were dehydrated and

123 mounted with Entellan mounting medium. Pictures of serial sections were aligned via the

124 TrakEM2 1.0[29–31] and 3D reconstructed with Mimics 24.0 software.

Ultrasound analysis

126 Ultrasound analysis was performed on zebrafish of 6-7 mpf as previously described[32]. All

127 measurements were performed in Vevo Lab 5.5.0.

Acute net handling stress induction

129 Induction of handling stress was adapted from previously described procedures[33]. Six mpf

130 zebrafish were netted and air suspended for 90 seconds, returned to the housing tank for

131 90 seconds, netted and air suspended for 90 seconds and returned to the housing tank.

Alizarin Red staining for mineralized bone

Alizarin red staining for mineralized bone was performed as previously described[34], at 13

mpf. The qKO were stained after sudden death between 6 and 13 mpf.

Ectopic bone on the sinistral skeletal elements of the vertebral bodies was scored on

abdominal and caudal vertebrae of adult zebrafish in lateral position[35].

Skull morphometrics were performed on images of the zebrafish heads in lateral position.

Acquired lengths were measured in ImageJ and are given in **Supplementary Fig. 1b.**

Diameter of the eye was used for normalization since eye diameter correlates with

standard length[36]. Cranial surface area was measured with standard length as

normalization factor.

Transcriptome analysis

RNA sequencing was performed on five pools, each containing five qKO or WT control cousins. Following RNA extraction (RNeasy® Mini kit), libraries were prepared (TruSeq Stranded Total RNA kit from Illumina) and sequenced on a NovaSeq6000 instrument (Illumina).

Following quality control [37,38], raw reads were mapped (STAR[39]) and through R packages, additional normalization and filtering was performed[40–42]. The top list of DEG was obtain via differential expression analysis (edgeR), with the necessary adjustments to account for multiple testing[43,44]. An adjusted p-value of 0.05 and a fold change of 2 were used as thresholds to determine genes of relevance. Finally gene enrichment analysis was carried out[45]. Further details can be found in the **Supplementary Methods**.

Synchrotron phase contrast micro Computed tomography imaging

Propagation-based phase-contrast synchrotron X-ray imaging was performed at the TOMCAT (X02DA) beamline of the Swiss Light Source (Paul Scherrer Institute in Villigen, Switzerland) as previously described[46]. Tomographic reconstruction was performed using the Gridrec algorithm after applying the Paganin phase retrieval method[47,48].

3D Biomechanical modeling to assess principal stress at systolic peak in the aortic wall

In a previous study[46], zebrafish-specific 3D biomechanical fluid-structure interaction models of five different 13-month-old zebrafish were developed. All parameter settings as described in[46] were maintained except for flow through right aortic arch 2, 3 and 4, which was set to nearly zero in order to mimic the vascular organization found in qKO mutants.

Transmission electron microscopy

In short, WT and qKO zebrafish of 16 mpf were euthanized, fixed in Karnovsky fixative, followed by decalcification for three weeks. Post-fixation in 1% osmium tetroxide, 0.1M cacodylate buffer with 8% saccharose and 0.004% $CaCl_2$ was followed by *en bloc* staining in 1% aqueous uranyl acetate. After dehydration, the samples were embedded in Spurr's resin. 80 nm sections were stained with uranyl acetate and lead citrate and examined by transmission electron microscopy (TEM) (JEM 1010, JEOL) equipped with a CCD side-mounted Veleta camera (EMSIS).

Swimming behavior studies

Swimming behavior of 9 mpf WT, $smad3a^{+/-};smad3b^{-/-};smad6a^{+/-};smad6b^{-/-}$ and qKO zebrafish was analyzed in a custom-made dark observation chamber (Noldus) equipped with the Basler GenICam. Data was analyzed using the EthoVision XT 17 software (Noldus). Zebrafish were placed individually in a tank and could acclimatize for 10 minutes before a 10-minute test period, was performed in the dark. Movement is depicted as total distance travelled during the test period, normalized for standard length. Movement frequency was also recorded.

1-phenyl-2-thiourea treatment

At 1 day post fertilization (dpf), 0.003% 1-phenyl-2-thiourea treatment (PTU) in E3 medium was administered to WT zebrafish. The medium was changed daily until 10 dpf. Starting from 5 dpf, zebrafish were given dry food for 2 hours daily, after which the medium was refreshed again.

Statistical analyses

All statistical analyses were performed with GraphPad Prism version 10.1.1 for Windows

(GraphPad Software, Boston, Massachusetts USA, www.graphpad.com). Kruskal-Wallis

test with Dunn's multiple comparisons test against WT controls was used to determine if

there was an increase in ectopic bone on the skeletal elements of the vertebral bodies.

For PTU treatment, a two-tailed t-test was used, when standard deviation differed

significantly between the groups, Welch's correction was applied. For all other

comparisons, One-way ANOVA with Dunnett's multiple comparison test against WT

controls was used. Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple

comparison test against WT controls was used when the genotypes showed a

significantly different standard deviation, determined with the Brown-Forsythe test.

RESULTS

Disruption of one of the smad3 or smad6 ohnologs does not show a cardiovascular or skeletal phenotype in zebrafish

Using CRISPR/Cas9 gene editing technology, we generated frameshift indels in the *smad3b*, *smad6a*, and *smad6b* genes, resulting in the generation of a premature termination codon in each mutant: s*mad3b*[c.455_459delinsATG/c.455_459delinsATG], *smad6a*[c.905_906delTG/c.905_906delTG] and *smad6b*[c.283_287delACGGT/c.283_287delACGGT] (**Supplementary Table 3**). To study *smad3a* function we made use of the available *Smad3a*[sa2363/+] line, in which a premature termination codon is introduced (c.682G>T). For the reader's benefit, we will use the term single knockout (SKO) for *smad3a, smad3b, smad6a* and *smad6b* single homozygous knockout zebrafish. At 5 dpf, measurement of the ventral aorta in the transgenic *Tg(fli1:EGFP)* reporter background is similar for heterozygous and SKO zebrafish for each gene as compared to sibling controls (**Supplementary Fig. 2**).

However, incross of *smad3a* SKO results in100% embryonic lethality at 2 dpf with severe body axis deformities and pericardial edema formation, while incross of *smad3b* SKO and *smad3a*[+/-]*;smad3b*[-/-] shows offspring with normal survival, suggesting that maternal *smad3a* expression is necessary for early embryonic survival, which is supported by previously published results[20]. Incross of *smad6a* or *smad6b* SKO showed normal survival.

In 6-7 mpf adult zebrafish, ultrasound measurements show that all measured parameters of cardiac function are similar in all SKO compared heterozygous mutants and sibling

controls, with an exception of *smad6a*$^{-/-}$ which shows a small increase in normalized

projected surface area of the ventricle in diastole (**Supplementary Fig. 3-4**).

In addition, skeletal evaluation and alizarin red staining for mineralized bone at 13 mpf do

not show any vertebral column deformities or an increase in ectopic bone formation (**Fig.**

4, **Supplementary Fig. 5-6**). Skull morphometrics are normal, except for the *smad3a*

SKO, in which a decrease of the snout to frontal bone distance is observed

(**Supplementary Fig. 7-8**).

Smad3a/b DKO and smad6a/b DKO show altered vasculogenesis

To bypass embryonic lethality due to loss of maternal *smad3a*, we performed a cross

between a *smad3a*$^{+/-}$*;smad3b*$^{-/-}$ female and *smad3a/b* DKO male zebrafish to obtain 50%

smad3a$^{+/-}$*;smad3b*$^{-/-}$ and 50% *smad3a/b* DKO zebrafish, which survive normally. At 10

dpf, the *smad3a/b* DKO show a significant increase in normalized surface area of the

aortic segment between aortic arch 3 and 4 (p = 0.0449), a reduced normalized length of

the segment (p = 0.0366) and an increased normalized mean aortic diameter (p = 0.0335).

The diameter adjacent to the bulbus arteriosus remains unchanged (**Supplementary Fig.**

9).

The *smad6a/b* DKO show normal early survival rates, as indicated by normal Mendelian

distribution of all genotypes in 10 dpf-old offspring obtained from *smad6a*$^{+/-}$*;smad6b*$^{+/-}$

incrosses. Nevertheless, *smad6a/b* DKO zebrafish are unable to survive to adulthood.

Smad6a/b DKO zebrafish show a significant reduction of the ventral aortic diameter at 10

dpf (p<0.0001). The diameter adjacent to the bulbus arteriosus as well as the normalized

surface area of the aorta segment is decreased (p<0.0001, p<0.0001). To a lesser extent,

the diameter of the aorta segment of *smad6a$^{+/-}$;smad6b$^{-/-}$* is also reduced (p = 0.0016),

while the length of the aorta segment remains unchanged in all *smad6* knockouts

(**Supplementary Fig. 9**).

Ultrasound analysis at 6-7 mpf of all viable *smad3* (*smad3a$^{+/-}$;smad3b$^{-/-}$*, *smad3a$^{-/-}$*

;smad3b$^{+/-}$ and *smad3a/b* DKO) or *smad6* (*smad6a$^{+/-}$;smad6b$^{-/-}$* and *smad6a$^{-/-}$;smad6b$^{+/-}$*

) KO combinations do not show any significant differences in cardiac parameters

(**Supplementary Fig. 3-4**).

Loss of smad3 ohnologs in zebrafish induces ectopic bone formation and vertebral column deformations

Alizarin red staining for mineralized bone on 13 mpf *smad3a/b* DKO zebrafish shows

ectopic bone formation on the ribs, neural arches, neural spines, haemal arches, and

haemal spines (p = 0.0012) (**Fig. 4**) as well as hyperlordosis, notochord mineralization

and notochord sheet mineralization (**Supplementary Fig. 5)**. Overall, the thickness of the

haemal arches is increased (**Fig. 4**) and the normalized surface area of the cranial roof

is reduced compared to WT controls (p = 0.0101) (**Supplementary Fig. 8**). The presence

of a single WT copy of either *smad3a* or *smad3b* prevents any of these skeletal

malformations to develop (**Fig. 4, Supplementary Fig. 8**). Skull morphometrics show a

mild decrease of the distance between the tip of the snout and the most posterior part of

the supraoccipital bone in s*mad3a$^{-/-}$* zebrafish (p = 0.0486), but not in the *smad3a/b* DKO

(**Supplementary Fig. 7**).

Likewise, a significant increase in ectopic bone can be seen in 13 mpf *smad6a$^{-/-}$*

;smad6b$^{+/-}$ (p = 0.0022) as well as in *smad6a$^{+/-}$;smad6$^{-/-}$* (p = 0.0033) zebrafish, although

to a lesser extent than observed in the *smad3a/b* DKO (**Fig. 4**). Since the *smad6a/b* DKO

does not reach adulthood, these features cannot be analyzed in this genotype. No

premature fusion of the cranial roof bones is observed, and the normalized surface areas

of the cranial roof are similar to controls (**Supplementary Fig. 8**). Skull morphometrics

do show a decrease in length between the tip of the snout and tip of the frontal bone in

smad6a$^{+/-}$;smad6b$^{-/-}$ mutants (p = 0.0056) (**Supplementary Fig. 7**).

Loss of smad3 and smad6 ohnologs in zebrafish induces ventral aortic

dissection and rupture

To investigate potential modifying effects of *smad3* on *smad6* and vice versa*,* we

generated a *smad3a$^{-/-}$;smad3b$^{-/-}$;smad6a$^{-/-}$;smad6b$^{-/-}$* qKO zebrafish line

(**Supplementary Table 3**).

Unlike the *smad6a/b* DKO, qKO zebrafish reach adulthood and generate viable offspring.

Microscopic analysis of vasculogenesis at 10 dpf shows a severe decrease of the

normalized surface area of the aortic projection (p<0.0001), the normalized mean

diameter of the aortic segment between aortic arch 3 and 4 (p = 0.0241), and the

normalized diameter adjacent to the bulbus arteriosus (p = 0.0005) compared to WT

controls, similar to *smad3a$^{+/-}$;smad3b$^{-/-}$;smad6a$^{+/-}$;smad6b$^{-/-}$* zebrafish (p<0.0001,

p<0.0001, p = 0.0032, respectively). The normalized length of the aortic segment is only

decreased in the qKO (p = 0.006) (**Fig. 1**, **Supplementary Fig. 9**). In addition, the qKO

shows tortuosity of the ventral aorta, aortic arches and hypobranchial artery (**Fig. 1**,

Supplementary Fig. 10).

In adult qKO zebrafish, we observed sudden deaths, mostly after breeding, netting, or

changing the tank positions. To investigate stress as a trigger, we performed a netting

test that is harmless in WT controls, but results in a mortality rate of 60% in qKO. Sudden

death associated with a red discoloration anterior in the abdominal region. Sectioning of

the entire ventral aorta and staining with Resorcin-Fuchsin to visualize the elastic fibers

shows the presence of aortic intramural hematomas and false lumens in multiple qKO.

3D-reconstructions of the aorta based on histological images and from synchrotron micro-

CT images of intact zebrafish, confirm the presence of regions of ventral aortic damage

(**Fig. 2** and **Supplementary movie 1-8**).

Regions of ventral aortic damage start close to a branching point (6/6 qKO, 0/4 WT) and

dissection or rupture is present at the most affected sites, (5/6 qKO, 0/4 WT). The

diameter of the aortic branches is asymmetrically distributed in all qKO zebrafish (6/6

qKO, 0/4 WT) with one side consistently showing narrower branches than the other. In

more severe cases, one or more aortic arches are missing (2/5 qKO, 0/4 WT) or attached

at the wrong side of the ventral aorta (1/6 qKO, 0/4 WT) (**Supplementary Movie 9-13**).

No morphological differences can be detected in WT controls (**Supplementary Movie

14-16**) or s*mad3a/b* DKO zebrafish (**Supplementary Movie 17-19**).

TEM shows a severely decreased elastin deposition in the internal elastic lamina of 15

mpf qKO zebrafish. Additionally, abnormal collagen deposition is detected near the intima

of the ventral aorta (**Fig. 2**).

Cardiac ultrasound analysis in 6-7 mpf qKO zebrafish shows regurgitation on ventricular

in- ($p = 0.0009$) and outflow ($p<0.0001$), suggestive of valve dysfunction (**Supplementary

Fig. 11**). Elastin staining of paraffin sections of the valves shows hypertrophy of the valve

interstitial cells, resulting in a more rounded appearance of the qKO bulboventricular valve

in contrast to the normal heart-shaped structure in WT control zebrafish. Similarly, the

atrioventricular valve shows hypertrophy and altered morphology of the valve interstitial

cells (**Fig. 3**, **Supplementary Fig. 12**).

Aortic damage in the qKO zebrafish correlates with increased stress at

branching regions

In qKO zebrafish, asymmetric branching and reduction of the diameter of the aortic arches

is apparent. To determine if the morphological changes of the aorta in adult qKO zebrafish

increase the principal stress within the vessel wall and to identify which locations endure

the largest differences, aortic morphology was compared using *in silico* simulation models

representing 13 mpf WT and qKO zebrafish aortae. To model the severe reduction of the

diameter of the qKO aortic arches, the blood flow through the right aortic arches (aortic

arch 2, 3 and 4) was reduced to nearly zero in the model. The principal stress within the

aortic wall was increased in all 5 qKO simulations, with the most affected region close to

branching points. This corresponds to the *ex vivo* observations of the location of regions

of damage in the ventral aorta of qKO zebrafish (**Supplementary Fig. 13**).

qKO zebrafish show skeletal aberrations, but largely normal swimming

behavior

The qKO show an increased presence of ectopic bone formation (p = 0.0014). Scoliosis,

hyperlordosis and hyperkyphosis, bending of the arches and ribs, as well as intervertebral

ligament mineralization and notochord mineralization are apparent in the qKO zebrafish

only (**Fig. 4**, **Supplementary Fig. 14**).

In qKO zebrafish, the nasal region of the skull is underdeveloped with a significantly

reduced distance between the tip of the snout and tip of the frontal bone (p<0.0001). (**Fig.**

4, Supplementary Fig. 7). Interestingly, qKO show normal values for the normalized

surface area of the cranial roof in contrast to the *smad3a/b* DKO, which was significantly

decreased (p = 0.0101) (**Supplementary Fig. 8).** Since qKO embryos show a tortuous

hypobranchial artery, which provides blood flow to the teeth, tooth morphology was

evaluated, but appears normal (**Supplementary Fig. 15**).

At 9 mpf, qKO zebrafish show a decrease in movement frequency (p = 0.0197), but not

in normalized total distance traveled compared to WT zebrafish (**Supplementary Fig.**

16).

***Bulk RNA sequencing of 5 dpf qKO* shows a reduction in the negative**

regulation of proteolysis and upregulation of melanogenesis

Bulk RNA sequencing in 5 dpf qKO and 2nd degree related WT control zebrafish identified

1131 differentially expressed genes (DEG) (adjusted p-value <0.05) (**Supplementary**

Table 4). The top 10 enriched GO-terms in the upregulated gene set mostly involve

pigmentation (pigment metabolic process, pigment biosynthetic process, melanosome

organization, pigment granule organization, melanocyte differentiation, pigmentation,

developmental pigmentation). One of the central DEG driving this enrichment is the *mitfa*

transcription factor, which is known to signal downstream of BMP and governs

melanocyte differentiation and development via tyrosinase (log FC *tyr* 2.01), tyrosinase

related protein 1 (log FC *tyrp1a* 1.82 and log FC *tyrp1b* 2.08), the premelanosome protein

(log FC *pmela* 1.44 and log FC *pmelb* 3.66), dopachrome tautomerase (log FC *dct* 2.6),

and the melanosomal transporter (log FC *slc24a5* 2.35)[49,50]. The top 10 enriched GO-

terms in the downregulated gene set are linked to negative regulation of proteolysis

(negative regulation peptidase activity, negative regulation proteolysis, regulation of

peptidase activity, negative regulation of endopeptidase activity, regulation of proteolysis)

and immune response (regulation of immune response, positive regulation of immune

system process) (**Supplementary Table 5-6**). A heatmap with stringent thresholds

(p.adjust ≥ 0.0001 and |log2FC| ≥ 1) and a volcano plot are shown in **Fig. 5**. Annotated

dysregulated genes with a known role in aortic wall homeostasis, include *loxl3a*, *emilin3a*

and *fbn2b* (elastic fiber assembly), *vcana* and *lamc2* (vascular cell adhesion

glycoproteins), *mhc1uma* (a zebrafish specific MHC class I gene), and *uts2a* (a

vasoconstrictor linked to hypertension and heart failure in humans). Dysregulation of *omd*

(a transcription factor, activated via BMP signaling, that links to cell adhesion and bone

mineralization) might relate to the observed bone phenotypes.

Tyrosinase inhibition increases diameter of the ventral aorta in 10 dpf

zebrafish

To investigate the involvement of melanogenesis-related pathways in aortic development,

we administered 0.003% PTU, a known inhibitor of tyrosinase in the medium of WT

zebrafish embryos starting at 1 dpf. At 10 dpf, treated zebrafish show a significant

increase of normalized ventral aortic surface area (p = 0.0066) and mean normalized

aorta diameter between aortic arch 3 and 4 (p = 0.0065) compared to vehicle control.

Length of the aortic segment and diameter adjacent to the bulbus arteriosus remained

unchanged compared to non-treated sibling controls (**Fig. 6**).

DISCUSSION

TGFβ and BMP signaling show significant crosstalk during vascular and skeletal development and homeostasis. Our results support the existence of modifier effects particularly on aortic phenotypes associated with *SMAD6*-deficiency, which range from bicuspid aortic valve with or without ascending aortic dilatation to Shone complex with a hypoplastic ascending aorta and aortic arch.

In the genome of mammals and teleost fish, *SMAD3* and *SMAD6* locate closely together, but in different topologically associated domains[16]. Since zebrafish express *smad3a*, *smad3b, smad6a* and *smad6b*[51–54], this model provides flexibility to investigate the interactions between these genes upon gene dosage. Although maternal *smad3a* is required during early embryogenesis, which is supported by previous findings[48], SKO show no obvious vascular or skeletal phenotype, except for mild craniofacial alterations in *smad3a*[-/-] zebrafish. However, *smad3a/b* or *smad6a/b* DKO do manifest cardiovascular and mild skeletal phenotypes, supporting compensation between the ohnologs of either smad3 or smad6, in line with their strongly conserved protein structure [19].

Smad6a/b and *smad3a/b* DKO show opposite vascular phenotypes. *Smad3a/b* DKO zebrafish show an increase in aortic diameter in early larval stages. This aligns with previous findings in which loss of *alk5*[-/-], the type 1 TGF-β receptor which activates SMAD3, leads to an increased diameter of the cardiac outflow tract in zebrafish embryos, followed by a failed development of the ventral aorta causing death at 7 dpf[55]. It is plausible that in *smad3a/b* DKO limited amounts of Smad2/Smad4 complexes partially rescue the phenotype preventing early death and an adult cardiovascular phenotype, in contrast to Alk5 loss of function, in which both Smad2 and Smad3 will not be activated[56].

S*mad6a/b* DKO show a reduced aortic diameter, and early death. The qKO also shows

a reduction of the aortic diameter and arterial tortuosity, but survives to adulthood. The

observation that the *smad3a*$^{+/-}$*;smad3b*$^{-/-}$*;smad6a*$^{+/-}$*;smad6b*$^{-/-}$ shows a more severe

reduction in aortic diameter than the qKO might be explained by the exclusion of the most

affected qKO embryos, since clear edge detection of the ventral aorta in these embryos

is hindered due to excessive tortuosity. Hence, it seems reasonable to generalize that

smad3 loss of function increases, while lack of smad6 decreases the diameter of the

aorta. The overall net result in the qKO shows a decreased diameter of the ventral aorta

with asymmetrical branching and a high susceptibility for dissection and rupture in adult

zebrafish. It has been previously reported that zebrafish *smad6b* is involved together with

BMP in modulation of lateral sprouting of the dorsal aorta[54].

Abnormal aortic branching results in altered hemodynamics and elevated systolic wall

stress, as confirmed by our fluid-structure interaction models presented here. Areas

exposed to perturbed local flow patterns prove prone to aortic wall damage and

dissection, as demonstrated by histological and synchrotron imaging of the aorta. TEM

analysis reveals a severe reduction of elastic fiber deposition in the inter-elastic laminae,

which can be linked to a lessened resilience to hemodynamic stress. Additionally,

abnormal collagen deposition near the intima of qKO zebrafish recapitulate increased

collagen deposition in human TAAD[57], indicating that similar pathophysiological

processes are activated in our zebrafish model. The presence of an atrioventricular and

bulboventricular valve defect in the qKO is in line with the role of SMAD6 in human

bicuspid aortic valve disease[58]. It is conceivable that unopposed loss of both *smad6*

ohnologs in zebrafish leads to a failure to maintain aortic homeostasis or to severe valve

dysfunction, resulting in the premature death observed in the *smad6a/b* DKO late juvenile
zebrafish.

The spectrum of bone phenotypes observed in the different genotypes confirms the
modulatory interactions between the different *smad* genes. Indeed, *smad3a/b* DKO, but
not *smad3a$^{+/-}$;smad3b$^{-/-}$* or *smad3a$^{-/-}$;smad3b$^{+/-}$* zebrafish, show increased ectopic bone
formation, a decreased normalized cranial roof surface area, and vertebral column axis
deviations. In contrast, the *smad6a$^{+/-}$;smad6b$^{-/-}$* and the *smad6a$^{-/-}$;smad6b$^{+/-}$* zebrafish do
show an increase in ectopic bone formation, although it needs to be highlighted that adult
smad6a/b DKO cannot be studied since they do not survive until adulthood. The qKO
zebrafish show more ectopic bone formation, and more severe column axis deviations.
These observations suggest a larger impact of BMP signaling on skeletal development,
that is further impacted by loss of TGFβ signaling.

Enrichment analysis of bulk RNA sequencing data of qKO embryos at 5 dpf revealed
upregulation of multiple pathways related to pigmentation, including the central
transcription factor *mitfa* as well as downstream effectors in melanin production such as
tyrp1a, *tyrp1b*, *tyr* and *dct*. MITF is a known downstream target of BMP signaling that is
inhibited by SMAD6[59–61]. MITF also acts upstream of lysosome biogenesis and it was
previously shown that lysosomal dysfunction in zebrafish due to loss of function of
atp6v1e1b associates with dilatation of the ventral aorta and narrow aortic branches[28].
Finally, MITF is known to be important for cell metabolism and cell cycle regulation,
promoting mitochondrial biogenesis and having complex effects on proliferation
depending on the level of MITF activity[62]. It is conceivable that one or more of these

effects of MITF are linked to the development of the phenotype observed in our qKO

model.

The transcriptional changes of pigmentation-related pathways in the qKO model

prompted us to test the consequences of pharmacological inhibition of tyrosinase, a key

enzyme responsible for eumelanin production, at a dose commonly used in zebrafish

research to block pigment formation. Inhibition of tyrosinase via PTU treatment resulted

in an increase of the aorta diameter while a decreased aortic diameter in qKO at 10 dpf

is associated with upregulated tyrosinase expression. This strongly supports a role for the

pigment biosynthesis pathway in aortic homeostasis. Therefore, the use of PTU for

imaging purposes should be approached with caution due to the potential for unintended

effects on other organ systems, especially when evaluating cardiovascular phenotypes.

Undesired effects of PTU have been reported previously including autophagy activation[63],

activation of cytochrome P4501A1 (*cyp1a1*)[64], suppression of retinol-binding protein 4

(*rbp4*)[65] and reduction of the diameter of the eye[66].

Despite upregulation of melanin-producing pathways, no increase of skin pigmentation

could be observed in the embryo or adult qKO zebrafish. Previous research showed that

tyrosinase follows a specific spatiotemporal expression pattern during embryogenesis,

which is not confined to skin cells[67]. At 7 dpf, small amounts of melanin can already be

detected around the dorsal aorta, while at 1 month, melanin deposition is apparent[68]. It is

therefore tempting to speculate that pigmentation-related pathways affect vascular

development, particularly since it was recently shown that tyrosinase reduces expression

of vascular endothelial growth factors[69].

The relevance of the qKO model to study the disease mechanisms of dissection is further

supported by the transcriptomic changes concordant with established models and known

pathogenetic mechanisms of TAD. Altered elastic fiber homeostasis, which was

confirmed in the qKO aortic wall using TEM, is reflected in upregulation of *Fbn2b*[70] (logFC

1.04), a fibrillin known to be involved in endocardial morphogenesis, which might be

upregulated as a response to the coarctation in the qKO, and downregulation of *mfap4*

(logFC -3.26), an important component of the extracellular matrix (ECM) involved in

elastic fiber assembly. In zebrafish, MFAP4 also regulates the balance between myeloid

and lymphoid development[71]. Another immune cell-related gene expressed differently in

qKO zebrafish is *itgam* (LogFC -5.33), coding for a component of the heterodimeric $\alpha_M\beta_2$

integrin expressed on leukocytes which is also known as macrophage-1 antigen (Mac-1)

or CD11b. This protein plays a crucial role in binding to ECM components and intracellular

adhesion molecules, involved in adhesion and transmigration of leukocytes across blood

vessels. The qKO transcriptomic profile also indicates that matrix proteolysis is increased,

due to downregulation of the metalloproteinase inhibitor *timp4* (LogFC -3.28), in

combination with upregulation of *mmp11b* (logFC 1.34) and *mmp13a* (logFC 1.35). These

targets are all widely established in the pathogenesis of vascular and bone homeostasis

[72–74]. *Loxl3a*, an orthologue of the human *LOXL3*[75] important for crosslinking of elastin and

collagen, is significantly downregulated in qKO zebrafish. In mice, deletion of *Loxl3* results

in abnormal skeletal development and is expressed in the precursors of the occipital and

interparietal bones and nasal area, affected in the qKO model[76].

In conclusion, Smad3-deficiency modifies Smad6-deficient phenotypes whereby *smad3a^-*

^/-;smad3b^-/-;smad6a^-/-;smad6b^-/- qKO zebrafish model aortic dissection and rupture,

resulting from defective vasculogenesis and resultant local wall stress. In addition to

known signatures of TAAD in human and mouse models, this model implicates the

pigmentation pathway in the development of the aortic phenotypes. Follow-up studies on

the precise contribution of these pathways are warranted and are likely to lead to novel

targets for therapeutic intervention in TAAD.

ACKNOWLEDGMENTS

The authors thank the family for their participation in this research. We also would like to thank the Zebrafish Facility Ghent Core at Ghent University, and particularly Karen Vermeulen for the diligent care for the zebrafish. We acknowledge the Paul Scherrer Institut (PSI), Villigen, Switzerland for provision of synchrotron radiation beamtime at the TOMCAT beamline X02DA of the Swiss Light Source, and would like to thank Karo De Rycke, Violette Deleeuw, Marina Horvat, Yousof Mohammad Asaad Abdel-Raouf (Ghent University) and Isaac Rodriguez Rovira (University of Barcelona) for their assistance in acquiring the synchrotron imaging data. Finally, we would like to thank the Nematology Research Unit, member of the UGent TEM Core Facility and the sequencing core of UZ Ghent.

SOURCES OF FUNDING

This work was supported by a Concerted Research Action grant from the Ghent University Special Research Fund (grant number BOF GOA019-21) to BC and PS, A research project of the Research Foundation – Flanders (grant number G0B4920N) and the Marfan Foundation Innovators Grant Program (grant number INT.DIV.2023.0008.01) to BC. Bert Callewaert is a senior clinical investigator of the Research Foundation-Flanders.

DISCLOSURES

None

DATA AVAILABILITY

Raw RNAseq data is available at Gene Expression Omnibus with accession number GSE249792. To review GEO accession GSE249792. For reviewers, following link is available to access the data.

https://eur03.safelinks.protection.outlook.com/?url=https%3A%2F%2Fwww.ncbi.nlm.nih.gov%2Fgeo%2Fquery%2Facc.cgi%3Facc%3DGSE249792&data=05%7C02%7Cmichiel.vanhooydonck%40ugent.be%7Cbeba6c62b40641ced96408dbfa282a45%7Cd7811cdeecef496c8f91a1786241b99c%7C1%7C0%7C638378822642521272%7CUnknown%7CTWFpbGZsb3d8eyJWIjoiMC4wLjAwMDAiLCJQIjoiV2luMzIiLCJBTiI6Ik1haWwiLCJXVCI6Mn0%3D%7C3000%7C%7C%7C&sdata=7g1JeMJOvj4lZVmQ6FVdkN%2FA2zrRUcuVlw6Hcear5vs%3D&reserved=0

Enter token obmhwiqkbbsprkt into the box.

All other data will be made available upon request.

AUTHOR CONTRIBUTIONS

Conceptualization: MVH., PSI., BC. Methodology: MVH., MVE., MSS., LP., AKB., DS., PSI., BC. Software: MSS., MVI. Formal analysis: MVH., MSV., MVI., DS. Investigation: MVH., MVE., MVI., MSS., LP., AKB., HDS., LC., PT., DS. Resources: PSI., BC. Data Curation: MVH., MSS. Writing (original draft): MVH., MVE. Writing (review & editing): MVH., MVE., MSS., LP., AKB, MVI., HDS., LC., PT., AB., P.SE., ADC., AW., DS., PSI., BC. Visualization: MVH. MVE. Supervision: PSI., BC. Funding acquisition: PSI., BC.

SUPPLEMENTARY MATERIAL

Supplementary Case Presentation

Supplementary Methods

Supplementary References

Supplementary Figures 1 - 17

Supplementary Tables 1 - 7

Supplementary Movies 1 - 19

REFERENCES

1.   Lemaire SA, Russell L. Epidemiology of thoracic aortic dissection. *Nature reviews. Cardiology.* 2011;8(2):103–113.

2.   Van De Laar IMBH, Oldenburg RA, Pals G, Roos-Hesselink JW, De Graaf BM, Verhagen JMA, Hoedemaekers YM, Willemsen R, Severijnen LA, Venselaar H, Vriend G, Pattynama PM, Collée M, Majoor-Krakauer D, Poldermans D, et al. Mutations in SMAD3 cause a syndromic form of aortic aneurysms and dissections with early-onset osteoarthritis. *Nature Genetics.* 2011;43(2):121–126.

3.   Deleeuw V, Carlson E, Renard M, Zientek KD, Wilmarth PA, Reddy AP, Manalo EC, Tufa SF, Keene DR, Olbinado M, Stampanoni M, Kanki S, Yanagisawa H, Mosquera LM, Sips P, et al. Unraveling the role of TGFβ signaling in thoracic aortic aneurysm and dissection using Fbn1 mutant mouse models. *Matrix biology : journal of the International Society for Matrix Biology.* 2023;123:17–33.

4.   Doyle JJ, Gerber EE, Dietz HC. Matrix-dependent perturbation of TGFβ signaling and disease. *FEBS Letters.* 2012;586(14):2003–2015.

5.   Li W, Li Q, Jiao Y, Qin L, Ali R, Zhou J, Ferruzzi J, Kim RW, Geirsson A, Dietz HC, Offermanns S, Humphrey JD, Tellides G. Tgfbr2 disruption in postnatal smooth muscle impairs aortic wall homeostasis. *The Journal of Clinical Investigation.* 2014;124(2):755–767.

6.   Bellini C, Bersi MR, Caulk AW, Ferruzzi J, Milewicz DM, Ramirez F, Rifkin DB, Tellides G, Yanagisawa H, Humphrey JD. Comparison of 10 murine models reveals a distinct biomechanical phenotype in thoracic aortic aneurysms. *Journal of the Royal Society Interface.* 2017;14(130):20161036.

7.   Schmierer B, Hill CS. TGFbeta-SMAD signal transduction: molecular specificity and functional flexibility. *Nature reviews. Molecular cell biology.* 2007;8(12):970–982.

8.   Miyazawa K, Miyazono K. Regulation of TGF-β Family Signaling by Inhibitory Smads. *Cold Spring Harbor perspectives in biology.* 2017;9(3)::a022095.

9.   Liu Y, Festing M, Thompson JC, Hester M, Rankin S, El-Hodiri HM, Zorn AM, Weinstein M. Smad2 and Smad3 coordinately regulate craniofacial and endodermal development. *Developmental Biology.* 2004;270(2):411–426.

10.   Timberlake AT, Choi J, Zaidi S, Lu Q, Nelson-Williams C, Brooks ED, Bilguvar K, Tikhonova I, Mane S, Yang JF, Sawh-Martinez R, Persing S, Zellner EG, Loring E, Chuang C, et al. Two locus inheritance of non-syndromic midline

craniosynostosis via rare SMAD6 and common BMP2 alleles. *eLife.*
2016;5(September2016):e20125.

11. Yang Y, Zheng Y, Li W, Li L, Tu M, Zhao L, Mei H, Zhu G, Zhu Y. SMAD6
is frequently mutated in nonsyndromic radioulnar synostosis. *Genetics in*
medicine : official journal of the American College of Medical Genetics.
2019;21(11):2577–2585.

12. Tan HL, Glen E, Töpf A, Hall D, O'Sullivan JJ, Sneddon L, Wren C, Avery
P, Lewis RJ, ten Dijke P, Arthur HM, Goodship JA, Keavney BD. Nonsynonymous
variants in the SMAD6 gene predispose to congenital cardiovascular
malformation. *Human Mutation.* 2012;33(4):720–727.

13. Luyckx I, MacCarrick G, Kempers M, Meester J, Geryl C, Rombouts O,
Peeters N, Claes C, Boeckx N, Sakalihasan N, Jacquinet A, Hoischen A,
Vandeweyer G, Van Lent S, Saenen J, et al. Confirmation of the role of
pathogenic SMAD6 variants in bicuspid aortic valve-related aortopathy. *European*
Journal of Human Genetics. 2019;27(7):1044–1053.

14. Luyckx I, Verstraeten A, Goumans MJ, Loeys B. SMAD6-deficiency in
human genetic disorders. *NPJ Genomic Medicine.* 2022;7(1):68.

15. Kloth K, Bierhals T, Johannsen J, Harms FL, Juusola J, Johnson MC,
Grange DK, Kutsche K. Biallelic variants in SMAD6 are associated with a complex
cardiovascular phenotype. *Human genetics.* 2019;138(6):625–634.

16. Rajderkar S, Barozzi I, Zhu Y, Hu R, Zhang Y, Li B, Alcaina Caro A,
Fukuda-Yuzawa Y, Kelman G, Akeza A, Blow MJ, Pham Q, Harrington AN,
Godoy J, Meky EM, et al. Topologically associating domain boundaries are
required for normal genome function. *Communications Biology.* 2023;6:435.

17. Giehl KA, Eckstein GN, Pasternack SM, Praetzel-Wunder S, Ruzicka T,
Lichtner P, Seidl K, Rogers M, Graf E, Langbein L, Braun-Falco M, Betz RC,
Strom TM. Nonsense mutations in AAGAB cause punctate palmoplantar
keratoderma type Buschke-Fischer-Brauer. *American journal of human genetics.*
2012;91(4):754–759.

18. Glasauer SMK, Neuhauss SCF. Whole-genome duplication in teleost
fishes and its evolutionary consequences. *Molecular genetics and genomics :*
MGG. 2014;289(6):1045–1060.

19. Pearson WR. An Introduction to Sequence Similarity ("Homology")
Searching. *Current protocols in bioinformatics.* 2013;chapter 3:3.1.1-3.1.8.

20. Jia S, Ren Z, Li X, Zheng Y, Meng A. smad2 and smad3 Are Required for
Mesendoderm Induction by Transforming Growth Factor-/Nodal Signals in
Zebrafish. *Journal of Biological Chemistry.* 2008;283(4):2418–2426.

21. Casari A, Schiavone M, Facchinello N, Vettori A, Meyer D, Tiso N, Moro E, Argenton F. A Smad3 transgenic reporter reveals TGF-beta control of zebrafish spinal cord development. *Developmental biology*. 2014;396(1):81–93.

22. Pogoda HM, Meyer D. Zebrafish smad7 is regulated by Smad3 and BMP signals. *Developmental Dynamics*. 2002;224(3):334–349.

23. Lawrence C, Sanders GE, Varga ZM, Baumann DP, Freeman A, Baur B, Francis M. Regulatory compliance and the Zebrafish. *Zebrafish*. 2009;6(4):453–456.

24. Westerfield M, Doerry E, Kirkpatrick AE, Douglas SA. Chapter 19 Zebrafish Informatics and the ZFIN Database. *Methods in Cell Biology*. 1998;60(C):339–355.

25. Naito Y, Hino K, Bono H, Ui-Tei K. CRISPRdirect: software for designing CRISPR/Cas guide RNA with reduced off-target sites. *Bioinformatics*. 2015;31(7):1120.

26. Boel A, De Saffel H, Steyaert W, Callewaert B, De Paepe A, Coucke PJ, Willaert A. CRISPR/Cas9-mediated homology-directed repair by ssODNs in zebrafish induces complex mutational patterns resulting from genomic integration of repair-template fragments. *DMM Disease Models and Mechanisms*. 2018;11(10):dmm035352.

27. Boel A, Steyaert W, De Rocker N, Menten B, Callewaert B, De Paepe A, Coucke P, Willaert A. BATCH-GE: Batch analysis of Next-Generation Sequencing data for genome editing assessment. *Scientific Reports*. 2016;6(July):30330.

28. Pottie L, van Gool W, Vanhooydonck M, Hanisch FG, Goeminne, Rajkovic A, Coucke P, Sips P, Callewaert B. Loss of zebrafish atp6v1e1b, encoding a subunit of vacuolar ATPase, recapitulates human ARCL type 2C syndrome and identifies multiple pathobiological signatures. *PLoS genetics*. 2021;17(6):e1009603.

29. Saalfeld S, Cardona A, Hartenstein V, Tomančák P. As-rigid-as-possible mosaicking and serial section registration of large ssTEM datasets. *Bioinformatics*. 2010;26(12):i57–i63.

30. Saalfeld S, Fetter R, Cardona A, Tomancak P. Elastic volume reconstruction from series of ultra-thin microscopy sections. *Nature methods*. 2012;9(7):717–720.

31. Cardona A, Saalfeld S, Schindelin J, Arganda-Carreras I, Preibisch S, Longair M, Tomancak P, Hartenstein V, Douglas RJ. TrakEM2 software for neural circuit reconstruction. *PloS one*. 2012;7(6):e38011.

32.   Van Impe M, Caboor L, Deleeuw V, Rycke K De, Vanhooydonck M, Backer J De, Segers P, Sips P. Application of an automated analysis framework for pulsed-wave Doppler cardiac ultrasound measurements to generate reference data in adult zebrafish. *American Journal of Physiology*. 2023;325(6):R782–R796.

33.   Ramsay JM, Feist GW, Varga ZM, Westerfield M, Kent ML, Schreck CB. Whole-body cortisol response of zebrafish to acute net handling stress. *Aquaculture*. 2009;297(1–4):157–162.

34.   Jarayseh T, Guillemyn B, De Saffel H, Bek JW, Syx D, Symoens S, Gansemans Y, Van Nieuwerburgh F, Jagadeesh S, Raja J, Malfait F, Coucke PJ, De Clercq A, Willaert A. A tapt1 knock-out zebrafish line with aberrant lens development and impaired vision models human early-onset cataract. *Human Genetics*. 2023;142(3):457–476.

35.   Bird NC, Mabee PM. Developmental morphology of the axial skeleton of the zebrafish, Danio rerio (Ostariophysi: Cyprinidae). *Developmental Dynamics*. 2003;228(3):337–357.

36.   Parichy DM, Elizondo MR, Mills MG, Gordon TN, Engeszer RE. Normal table of postembryonic zebrafish development: Staging by externally visible anatomy of the living fish. *Developmental Dynamics*. 2009;238(12):2975–3015.

37.   Andrews S. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data. *FastQC A Quality Control tool for High Throughput Sequence Data*. 2010.

38.   Krueger F, James F, Ewels P, Afyounian E, Schuster-Boeckler B. FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo. 2021.

39.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 2013;29(1):15–21.

40.   Sayols S, Scherzinger D, Klein H. dupRadar: a Bioconductor package for the assessment of PCR artifacts in RNA-Seq data. *BMC bioinformatics*. 2016;17(1):428.

41.   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*. 2016;32(19):3047–3048.

42.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*. 2014;30(7):923–930.

43.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics*
(Oxford, England). 2010;26(1):139–140.

44.   Chen Y, Lun ATL, Smyth GK. From reads to genes to pathways:
Differential expression analysis of RNA-Seq experiments using Rsubread and the
edgeR quasi-likelihood pipeline. *F1000Research*. 2016;5.

45.   Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for
comparing biological themes among gene clusters. *Omics : a journal of integrative*
biology. 2012;16(5):284–287.

46.   Van Impe M, Caboor L, Deleeuw V, Olbinado M, De Backer J, Sips P,
Segers P. Fluid-Structure Interaction Modeling of the Aortic Hemodynamics in
Adult Zebrafish: a Pilot Study based on Synchrotron X-Ray Tomography. *IEEE*
Transactions on Biomedical Engineering. 2023;70(7):2101–2110.

47.   Paganin D, Mayo SC, Gureyev TE, Miller PR, Wilkins SW. Simultaneous
phase and amplitude extraction from a single defocused image of a
homogeneous object. *Journal of Microscopy*. 2002;206(1):33–40.

48.   Marone F, Stampanoni M. Regridding reconstruction algorithm for real-
time tomographic imaging. *Journal of synchrotron radiation*. 2012;19(Pt 6):1029–
1037.

49.   Hsiao JJ, Fisher DE. The roles of microphthalmia-associated transcription
factor and pigmentation in melanoma. *Archives of Biochemistry and Biophysics*.
2014;563:28–34.

50.   Bian C, Li R, Wen Z, Ge W, Shi Q. Phylogenetic Analysis of Core Melanin
Synthesis Genes Provides Novel Insights Into the Molecular Basis of Albinism in
Fish. *Frontiers in Genetics*. 2021;12:707228.

51.   Casari A, Schiavone M, Facchinello N, Vettori A, Meyer D, Tiso N, Moro
E, Argenton F. A Smad3 transgenic reporter reveals TGF-beta control of zebrafish
spinal cord development. *Developmental Biology*. 2014;396(1):81–93.

52.   Pogoda HM, Meyer D. Zebrafish smad7 is regulated by Smad3 and BMP
signals. *Developmental Dynamics*. 2002;224(3):334–349.

53.   De Pater E, Ciampricotti M, Priller F, Veerkamp J, Strate I, Smith K,
Lagendijk AK, Schilling TF, Herzog W, Abdelilah-Seyfried S, Hammerschmidt M,
Bakkers J. Bmp signaling exerts opposite effects on cardiac differentiation.
Circulation Research. 2012;110(4):578–587.

54.   Mouillesseaux KP, Wiley DS, Saunders LM, Wylie LA, Kushner EJ, Chong
DC, Citrin KM, Barber AT, Park Y, Kim JD, Samsa LA, Kim J, Liu J, Jin SW,

Bautch VL. Notch regulates BMP responsiveness and lateral branching in vessel networks via SMAD6. *Nature Communications*. 2016;7:13247.

55.	Boezio GLM, Bensimon-Brito A, Piesker J, Guenther S, Helker CSM, Stainier DYR. Endothelial TGF-β signaling instructs smooth muscle cell development in the cardiac outflow tract. *eLife*. 2020;9:1–32.

56.	Lucarelli P, Schilling M, Kreutz C, Vlasov A, Boehm ME, Iwamoto N, Steiert B, Lattermann S, Wäsch M, Stepath M, Matter MS, Heikenwälder M, Hoffmann K, Deharde D, Damm G, et al. Resolving the Combinatorial Complexity of Smad Protein Complex Formation and Its Link to Gene Expression. *Cell Systems*. 2018;6(1):75-89.e11.

57.	Wang X, LeMaire SA, Chen L, Shen YH, Gan Y, Bartsch H, Carter SA, Utama B, Ou H, Coselli JS, Wang XL. Increased Collagen Deposition and Elevated Expression of Connective Tissue Growth Factor in Human Thoracic Aortic Dissection. *Circulation*. 2006;114(SUPPL. 1):I200–I205.

58.	Bechtel JFM, Misfeld M, Schmidtke C, Sievers HH. The bicuspid aortic valve. In: *Aortic Root Surgery*.; 2010:89–101.

59.	Yang J, Wang J, Pan L, Li H, Rao C, Zhang X, Niu G, Qu J, Hou L. BMP4 is required for the initial expression of MITF in melanocyte precursor differentiation from embryonic stem cells. *Experimental cell research*. 2014;320(1):54–61.

60.	Massaous J, Hata A. TGF-B signalling through the Smad pathway. *Cell Biology*. 1997;7:187–192.

61.	Steinfeld J, Steinfeld I, Coronato N, Hampel ML, Layer PG, Araki M, Vogel-Höpker A. RPE specification in the chick is mediated by surface ectoderm-derived BMP and Wnt signalling. *Development*. 2013;140(24):4959–4969.

62.	Goding CR, Arnheiter H. Mitf—the first 25 years. *Genes and Development*. 2019;33(15–16):983–1007.

63.	Chen XK, Kwan JSK, Chang RCC, Ma ACH. 1-phenyl 2-thiourea (PTU) activates autophagy in zebrafish embryos. *Autophagy*. 2021;17(5):1222–1231.

64.	Wang W Der, Wang Y, Wen HJ, Buhler DR, Hu CH. Phenylthiourea as a weak activator of aryl hydrocarbon receptor inhibiting 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced CYP1A1 transcription in zebrafish embryo. *Biochemical Pharmacology*. 2004;68(1):63–71.

65.	Tingaud-Sequeira A, Forgue J, André M, Babin PJ. Epidermal transient down-regulation of retinol-binding protein 4 and mirror expression of apolipoprotein Eb and estrogen receptor 2a during zebrafish fin and scale development. *Developmental Dynamics*. 2006;235(11):3071–3079.

66.   Li Z, Ptak D, Zhang L, Walls EK, Zhong W, Leung YF. Phenylthiourea
Specifically Reduces Zebrafish Eye Size. *PLOS ONE*. 2012;7(6):e40132.

67.   Camp E, Lardelli M. Tyrosinase gene expression in zebrafish embryos.
Development genes and evolution. 2001;211(3):150–153.

68.   Miano JM, Georger MA, Rich A, De Mesy Bentley KL. Ultrastructure of
zebrafish dorsal aortic cells. *Zebrafish*. 2006;3(4):455–463.

69.   Clahsen T, Hatami N, Büttner C, Reis A, Cursiefen C. Tyrosinase reduces
expression of vascular endothelial growth factors and improves corneal graft
survival. *Investigative Ophthalmology & Visual Science*. 2022;63(7):909-A0273.

70.   Mellman K, Huisken J, Dinsmore C, Hoppe C, Stainier DY. Fibrillin-2b
regulates endocardial morphogenesis in zebrafish. *Developmental biology*.
2012;372(1):111–119.

71.   Ong SLM, De Vos IJHM, Meroshini M, Poobalan Y, Dunn & NR.
Microfibril-associated glycoprotein 4 (Mfap4) regulates haematopoiesis in
zebrafish. 2020;10:11801.

72.   Koskivirta I, Rahkonen O, Mäyränpää M, Pakkanen S, Husheem M, Sainio
A, Hakovirta H, Laine J, Jokinen E, Vuorio E, Kovanen P, Järveläinen H. Tissue
inhibitor of metalloproteinases 4 (TIMP4) is involved in inflammatory processes of
human cardiovascular pathology. *Histochemistry and cell biology*.
2006;126(3):335–342.

73.   Johansson N, Ahonen M, Kähäri VM. Matrix metalloproteinases in tumor
invasion. *Cellular and Molecular Life Sciences*. 2000;10(6):415–433.

74.   Jones JA, Ikonomidis JS. The Pathogenesis of Aortopathy in Marfan
Syndrome and Related Diseases. *Current cardiology reports*. 2010;12(2):99–107.

75.   Van Boxtel AL, Gansner JM, Hakvoort HWJ, Snell H, Legler J, Gitlin JD.
Lysyl oxidase-like 3b is Critical for Cartilage Maturation During Zebrafish
Craniofacial Development. *Matrix biology : journal of the International Society for
Matrix Biology*. 2011;30(2001):178–187.

76.   Santamaría PG, Dubus P, Bustos-tauler J, Floristán A, Vázquez-naharro
A, Morales S, Cano A, Portillo F. Loxl2 and Loxl3 Paralogues Play Redundant
Roles during Mouse Development. *International Journal of Molecular Sciences*.
2022;23(10):5730.

792 **FIGURES WITH FIGURE LEGENDS**

793   Links with access to high quality figures and the supplementary movies. Lower quality

794   figures can be found below:

Figure 1	https://figshare.com/s/58540e31ae5dcfed3750
Figure 2	https://figshare.com/s/b08f40d0b6c136b31ee4
Figure 3	https://figshare.com/s/9fbc19810756ed1b34b4
Figure 4	https://figshare.com/s/07b6703f58ea0653336e
Figure 5	https://figshare.com/s/2b913da4b07559cc0607
Figure 6	https://figshare.com/s/42bdead6d4b916ec7327
Supplementary Figure 1	https://figshare.com/s/1d96715da2007c11ae96
Supplementary Figure 2	https://figshare.com/s/7117068f173e59e3b7e7
Supplementary Figure 3	https://figshare.com/s/8a7ce09585ca754dedd1
Supplementary Figure 4	https://figshare.com/s/e754b6c96ea2c0588a15
Supplementary Figure 5	https://figshare.com/s/c902712aae5ffea8e862
Supplementary Figure 6	https://figshare.com/s/9bcafde922edd89090ad
Supplementary Figure 7	https://figshare.com/s/00039d2f1472d980fb15
Supplementary Figure 8	https://figshare.com/s/19aa5c0c33dfbf52f76b
Supplementary Figure 9	https://figshare.com/s/3b70b2174c355aafa4c8
Supplementary Figure 10	https://figshare.com/s/2ed90596b1477259fc3f
Supplementary Figure 11	https://figshare.com/s/5c8b0c3c23600bd54b15
Supplementary Figure 12	https://figshare.com/s/2c1733efb6b877eed39a
Supplementary Figure 13	https://figshare.com/s/2a70a063f0dbda9c8bac
Supplementary Figure 14	https://figshare.com/s/58a4c79c1b31b4474c1c
Supplementary Figure 15	https://figshare.com/s/5ccc12c8bf07a477ab19
Supplementary Figure 16	https://figshare.com/s/64b5c5461c87858830e5
Supplementary Figure 17	https://figshare.com/s/a950dec1eb4b32f8559a
Supplementary Movie 1	https://figshare.com/s/5f3e5147898632f68274
Supplementary Movie 2	https://figshare.com/s/cdf6d880d1a241c750f4
Supplementary Movie 3	https://figshare.com/s/5b6d3fb558aeb40771d2
Supplementary Movie 4	https://figshare.com/s/f2c26de3a7403d10148c
Supplementary Movie 5	https://figshare.com/s/03f83afad3ad6383aa20
Supplementary Movie 6	https://figshare.com/s/51aa1e42b4cb45ecdc60
Supplementary Movie 7	https://figshare.com/s/415c7431de05ecfc5449
Supplementary Movie 8	https://figshare.com/s/8a2169e10ff223bc971d
Supplementary Movie 9	https://figshare.com/s/a3aaebcb00ccb50f9b4d
Supplementary Movie 10	https://figshare.com/s/84ac6e35c61e47142339
Supplementary Movie 11	https://figshare.com/s/b87dfe1d640a284edf86
Supplementary Movie 12	https://figshare.com/s/f74f27f17b2cb7343559
Supplementary Movie 13	https://figshare.com/s/b9f1f922f31ada373b87
Supplementary Movie 14	https://figshare.com/s/9a1291cd30650944a05f
Supplementary Movie 15	https://figshare.com/s/3a08c958c980bb16c948

Supplementary Movie 16	https://figshare.com/s/5cd90d5c165d52dfa319
Supplementary Movie 17	https://figshare.com/s/3bac7c34e759d5849442
Supplementary Movie 18	https://figshare.com/s/db1b70cf18c9fe058a57
Supplementary Movie 19	https://figshare.com/s/769438c9b5a3cfade65d

Figure 1: Altered vasculogenesis at 10 dpf in double and qKO mutants. (a-d) Ventral view of the cardiovascular structures at 10 dpf of *smad* knockout lines crossed with a *Tg(fli1:EGFP)* reporter line. Legend: "VA" ventral aorta, "BA" bulbus arteriosus, "AA3" aortic arch 3, "AA4" aortic arch 4, "HL" head length. The projected surface area of the ventral aorta between aortic arch 3 and aortic arch 4 has been outlined with a white dashed line. **(e)** *smad3a+/-;smad3b-/- and smad3a-/-;smad3b-/-* zebrafish of 10 dpf show an increase in normalized aorta diameter (n=29-11-7). **(f)** *smad6a+/-;smad6b-/-* as well as *smad6a-/-;smad6b-/-* show a decrease in diameter of the ventral aorta (n=15-18-14-10-23). **(g)** *smad3a+/-;smad3b-/-;smad6a+/-;smad6b-/-* and *smad3a-/-;smad3b-/-;smad6a-/-;smad6b-/-* both show a decrease in diameter of the ventral aorta (n=15-18-26). **(h,i)**

Overview of the hypobranchial artery in a WT control and qKO zebrafish line. Tortuosity

and irregular branching of the hypobranchial artery (HA) and aortic arch 1 (AA1) can be

observed **(i)** magnification of boxed area in d. **(a-d,h,i)** stack focused Z-stack pictures

obtained with ZEISS Axio Observer.Z1 microscope **(f)** One-way ANOVA with Dunnett's

multiple comparison test against WT controls. **(e,g)** Brown-Forsythe and Welch ANOVA

with Dunnett's T3 multiple comparison test against WT controls. Asterisks indicate

significant differences. $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$. Data represented as

average ± standard deviation.

Figure 2: qKO zebrafish exhibit thoracic aortic dissection and rupture. (a, b) qKO

zebrafish that died suddenly show similar external features compared to qKO zebrafish

that die due to stress induction. Red discoloration around the heart and gill area (anterior

abdominal area) is indicated with a white arrowhead. **(c)** Resorcin-Fuchsin staining of WT

control. **(d)** Resorcin-Fuchsin staining of the elastic fiber on a 5 µm cross-section of the

ventral aorta shows rupture of the aortic wall (red arrow). **(e,f)** Resorcin-Fuchsin staining

of the elastic fiber on 5 μm thick cross-sections of the ventral aorta. False lumen is

indicated with a black arrow. **(g,h)** 3D reconstruction using 3D-modelling software Mimics

24.0 of the ventral aorta shows a decrease in diameter of the aortic arches. Symmetrical

branching of the aortic arches is lost in the qKO model. Region with aortic damage has

been colored black **(i,j).** The area of damage identified on a histological resorcin-fuchsin

staining of the ventral aorta can similarly be observed on a synchrotron micro-CT scan of

the same region. Damage is indicated with arrows. **(k,l)** TEM of aortic wall in WT and

qKO, respectively. Severe decrease of elastic fiber in the inner elastic lamina is apparent.

(m) TEM of qKO shows collagen deposition in intima. **(k,l,m)** Legend: "VSMC" vascular

smooth muscle cell, "EC" endothelial cell, "RBC" red blood cell, "C" collagen, * indicates

inner elastic lamina.

Figure 3: Increased regurgitation and abnormal valve morphology in qKO zebrafish. (a, c) Pulsed wave doppler of inflow **(a)** and outflow **(c)** of blood in and out of the ventricle of WT control zebrafish. **(b, d)** Pulsed wave doppler of inflow **(b)** and outflow **(d)** of blood in and out of the ventricle of qKO zebrafish. **(e)** Regurgitation on inflow is significantly increased in the qKO zebrafish model (n=17-8-7-8-8-10). **(f, g)** Resorcin

Fuchsin staining shows hypertrophy of the valve interstitial cells and abnormal leaflet

morphology in the atrioventricular valve of qKO zebrafish. **(h)** Increased regurgitation on

outflow in qKO (n=17-7-7-7-7-10). **(i,j)** Resorcin Fuchsin staining of the bulboventricular

valve in WT and qKO zebrafish shows hypertrophy of the valve and altered orientation in

the qKO. **(a-j)** Legend: "RI" regurgitation inflow, "RO" regurgitation outflow, "A" atrium, "V"

ventricle, "B" bulbus arteriosus. Black arrows indicate valves. **(e,h)** One-way ANOVA with

Dunnett's multiple comparison test against WT controls. Asterisks indicate significant

differences. ***$p < 0.001$, ****$p < 0.0001$. Data represented as average ± standard

deviation.

Figure 4: qKO zebrafish show scoliosis, lordosis and ectopic bone formation in the vertebral column. (a-l) Overview pictures of alizarin red staining for mineralized bone of abdominal and caudal vertebrae, and the skull (lateral view) of WT (**a-d**), *smad3a^-/-;smad3b^-/-* DKO zebrafish (**e-h**), and *smad3a^-/-;smad3b^-/-;smad6a^-/-;smad6b^-/-* qKO zebrafish (**i-l**). Ectopic bone has been indicated with white arrowheads. Scoliosis and lordosis can be observed in (**i**). **(k)** Notochord sheet mineralization can also be observed between the vertebrae. Small craniofacial aberrations are detected. **(l)** The nasal bone structures appear to be missing, resulting in a clockwise rotation of the snout. **(m-o)** Quantification of spots of ectopic bone in *smad3* (n=10-10-8-10-10-4), *smad6* (n=10-4-3-

862   8-5), *smad3/smad6* (n=10-6-13-14-15-8) genotypes. **(a-l)** Whole-mount alizarin red

863   staining for mineralized bone pictures taken with Leica M165 FC Fluorescent Stereo

864   Microscope. **(m-o)** Kruskal-Wallis test with Dunn's multiple comparisons test against WT

865   controls. $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$. Data represented as average ±

866   standard deviation.

Figure 5: RNA sequencing data shows downregulation of negative regulators of proteolysis and upregulation of melanogenesis. (a) Volcano Plot showing upregulated genes (red) and downregulated genes (blue) with genes of importance annotated. Thresholds: p.adjust ≥ 0.05 and |log2FC| ≥ 1; **(b)** Heatmap with top DEG results between WT and qKO. Thresholds: p.adjust ≥ 0.0001 and |log2FC| ≥ 1 ; **(c)** Top 10 hits of GO enrichment analysis for both sets of upregulated (red) and downregulated (blue) genes. Numbers inside the bars represent the corresponding adjusted p-value value.

control

PTU treated

a

b

100 µm

100 µm

c   normalized surface
    area aorta segment

**

d   normalized diameter
    aorta segment

**

e   normalized diameter adjacent
    to the bulbus arteriosus

f   normalized length
    aorta segment

Figure 6: Tyrosinase inhibition increases the aorta diameter of 10 dpf zebrafish. (a)

Ventral view of the vasculature of 10 dpf WT control zebrafish. **(b)** Ventral view of the

vasculature of 10 dpf WT zebrafish treated with 0.003% PTU treatment starting at 24 hpf.

(c, d) PTU-treated zebrafish show a significant increase in normalized surface area of the

aortic segment and aortic diameter (n=15-16). **(e,f)** Normalized diameter adjacent to the

bulbus arteriosus and length of the aorta segment is similar between treated and non-

treated zebrafish (n=15-16). **(a,b)** Dashed white line indicates aorta segment between

aortic arch 3 and 4. Stack focused Z-stack pictures obtained with ZEISS Axio Observer.Z1

microscope**. (c,d)** Unpaired two-tailed t-test with Welch's correction. **(e,f)** Unpaired two-

tailed t-test. Asterisks indicate significant differences. **p<0.01. Data represented as

average ± standard deviation.