1 **Oncogenic KRAS cells use Wnt signalling and cell dormancy to override homeostatic**
2 **cell elimination mechanisms in adult pancreas.**

3 Beatriz Salvador-Barbero[1], Markella Alatsatianos[1], Jennifer P. Morton[2, 3], Owen J. Sansom[2,
4 3], Catherine Hogan[1]*.

   1. European Cancer Stem Cell Research Institute, School of Biosciences, Cardiff
      University, Cardiff CF24 4HQ, UK.
   2. CRUK Scotland Institute, Glasgow, G61 1BD, UK.
   3. School of Cancer Sciences, University of Glasgow, Glasgow, G61 1QH, UK.

9 *Correspondence to hoganc@cardiff.ac.uk

10
11

Abstract

Epithelial tissues use homeostatic defence mechanisms to actively expel aberrant or
genetically mutant cells and prevent disease. When present in healthy tissues in low
numbers, we show that cells expressing cancer-causing mutations (KrasG12D, p53R172H)
compete with normal cells for survival and are often eliminated. Thus, tumour initiation must
require mechanisms whereby mutant cells override tissue defence mechanisms to remain in
a tissue; however, the biology of these initial events is poorly understood. Here, we use an *in
vivo* model of sporadic tumorigenesis in the adult pancreas to show that a population of
KrasG12D- or p53R172H-expressing cells are never eliminated from the epithelium. Using
RNA sequencing of non-eliminated populations and quantitative fluorescence imaging, we
show that β-catenin-independent Wnt5a signalling, and cell dormancy are key features of
surviving KrasG12D cells *in vivo*. We demonstrate that Wnt5a (and not Wnt3a) inhibits apical
extrusion of RasV12 cells *in vitro* by promoting stable E-cadherin-based cell-cell adhesions
at RasV12-normal cell-cell boundaries. Inhibition of Wnt5a signalling restores E-cadherin
dynamics at normal-mutant boundaries and apical extrusion *in vitro*. RasV12 cells arrested
in the cell cycle are not extruded and this is rescued when Wnt signalling is inhibited. In the
pancreas, Wnt signalling, E-cadherin and β-catenin are increased at cell-cell contacts
between non-eliminated KrasG12D cells and normal neighbours. Importantly, we
demonstrate that active Wnt signalling is a general mechanism required to promote
KrasG12D and p53R172H cell survival *in vivo*. Treatment with porcupine inhibitor rescues
pancreas tissue defence by switching mutant cell retention to cell expulsion. Our results
suggest that RAS mutant cells activate Wnt and a dormant cell state to avoid cell expulsion
and to survive in the adult pancreas.

Keywords: Epithelial homeostasis, cell extrusion, cell competition, oncogenic *RAS*, mutant
Tp53, pancreas, Wnt5a, cell dormancy, pancreatic cancer.

Introduction

Epithelial tissues are continuously exposed to mutational insults, and as a result, genetically mutant cells often arise in a tissue. Indeed, most human cancers start sporadically from epithelial cells carrying genetic mutations that activate oncogenes or inactivate tumour suppressor function. Remarkably, epithelial tissues protect against tumorigenesis by actively removing aberrant (*1, 2*) and genetically mutant cells (*3-9*). In general, genetically different cells compete for space and survival in tissues, resulting in the elimination of 'less fit' mutant cells via apoptosis, extrusion and/or cell differentiation (*10*). Cell competition outcomes are dependent on the genetic status of a cell relative to its local neighbourhood (*10, 11*) and extrinsic environmental factors (*12-14*) and therefore under certain conditions, competition is tipped in favour of mutant cells. This is better understood in rapidly proliferating tissues that are actively replenished via defined stem cell compartments; tumour risk increases when stem cells acquire genetic mutations that confer a competitive advantage over wild type counterparts, allowing the mutant stem cells to completely occupy the niche (*15-18*). In contrast, our understanding of how cell competition outcomes shape tumorigenesis in slow proliferating adult tissues that lack a *bona fide* stem cell compartment remains poorly detailed.

Using mouse pancreas as a model system of a slow proliferating adult tissue, we recently demonstrated that cells expressing oncogenic *Kras* (*Kras*[G12D]) are outcompeted by healthy neighbours and are eliminated from exocrine and endocrine epithelial compartments (*4*). We have previously described how interactions with normal cells trigger robust evolutionary conserved cell biology phenotypes in *Ras* transformed cells, resulting in the elimination of the mutant cells via cell segregation and cell extrusion (*4, 5, 9, 19*). Removal of KrasG12D cells from adult pancreas tissues requires local remodelling of E-cadherin-based cell-cell adhesions at normal-mutant boundaries and dynamic changes in mutant and normal cell volume (*4*). Importantly, we demonstrated that abrogation of KrasG12D cell elimination in the pancreas significantly increased the appearance of preneoplastic lesions (*4*), suggesting cell competition and the subsequent expulsion of mutant cells are disease preventative.

Activating mutations in oncogenic *RAS* are the principal driver gene event in human pancreatic ductal adenocarcinoma (PDAC; (*20*)), the most common type of human pancreatic cancer. *KRAS* mutations are detected in >90% of human tumours and *in vivo* mouse studies show that KRAS signalling is essential for disease to progress through all stages (*21*). Missense mutations in *TP53* are the second most common mutation detected in ~70% of human PDAC (*22*) and required for metastasis (*23*). How genetically mutant cells survive and grow in the competitive environment of the adult pancreas remains unclear. Here, we sought to understand the mechanisms underpinning how mutant cells avoid cell

expulsion and survive in the adult pancreas. Using *in vivo* models of sporadic tumorigenesis,

we confirm that adult pancreas tissues actively eliminate cells expressing either oncogenic

KrasG12D or p53R172H mutations; however, some mutant cells are never eliminated,

suggesting these cells have a survival advantage. We report that cell competition is

completely abrogated when tissues contain 'double mutant' cells, i.e., cells expressing both

KrasG12D and p53R172H mutations. Using bulk RNA sequencing, we identify gene

signatures of 'never-eliminated' mutant cells. We demonstrate a functional role of Wnt

signalling as an important general regulator of mutant cell fate and cell dormancy as a

second mechanism, whereby KrasG12D cells avoid cell elimination.

Results

Cells expressing p53R172H mutations are outcompeted and eliminated from adult

pancreas compartments. Cells co-expressing p53172H and KrasG12D mutations are

never eliminated.

To model sporadic pancreatic cancer, we used the pancreas-specific genetically engineered

mouse models: KC: *Pdx1-Cre^{ERT}; LSL-Kras$^{G12D/+}$; Rosa26$^{LSL-tdRFP}$*; PC: *Pdx1-Cre^{ERT}; LSL-*

Trp53$^{R172H/+}$; Rosa26$^{LSL-tdRFP}$; and KPC: *Pdx1-Cre^{ERT}; LSL-Kras$^{G12D/+}$; Trp53$^{R172H/+}$; Rosa26^{LSL-}*

^{tdRFP}. Our experimental control was *Pdx1-Cre^{ERT}; Rosa26$^{LSL-tdRFP}$* mice (Fig. 1A left). Adult

mice were treated with low-dose tamoxifen to induce transgene expression (together with

RFP reporter) in a mosaic manner in all pancreatic epithelial compartments (*4*). To monitor

genetically mutant cell fate, we measured RFP levels in tissues harvested at 7-, 35- and 70-

days post induction (p.i.) of Cre recombinase (Fig. 1A-1C). Consistent with our previously

published data (*4*), we found that low dose tamoxifen induced stochastic RFP labelling in

~20-25% of the tissue (Fig. 1C); levels of RFP fluorescence were comparable between all

genotypes at 7 days (Fig. 1B, top panels; 1C; (*4*)) and significantly decreased over time in

KrasG12D tissues (KC; Fig. 1B, lower panel; 1C, $p<0.0001$ for both 35 and 70 days p.i.).

RFP fluorescence also significantly decreased over time in tissues expressing p53R172H

mutations (PC; Fig. 1B, lower panel; 1C, $p<0.0001$ and $p=0.0173$ at 35 and 70 days p.i.

respectively). In contrast, we found no significant difference in the level of RFP in double

mutant (KPC) tissues over time (Fig. 1B, lower panel; 1C, p=0.2062 and p=0.1504 at 35 and

70 days p.i. respectively). RFP+ ducts were significantly less frequent in KrasG12D (KC) and

p53R172H (PC) tissues at the 7-day p.i. time point, compared to controls (Fig. 1D, $p=0.0038$

and $p=0.0029$ respectively), while the number of RFP+ ducts in double mutant (KPC) tissues

was comparable to controls (Fig. 1D, $p=0.2879$). We also measured a significant decrease in

the percentage of RFP+ cells per islet in p53R172H (PC) and KrasG12D (KC) tissues (Fig.

1E, $p=0.0214$ and $p=0.0134$ respectively), whereas the frequency of RFP+ cells per islet in

double mutant (KPC) tissues was also comparable to wild type controls (Fig. 1E, $p=0.5394$).

Thus, like KrasG12D-expressing cells (*4*), p53R172H single mutant cells are eliminated from

all epithelial compartments. Cells expressing both KrasG12D and p53R172H mutations are

never eliminated from endocrine/exocrine epithelial tissues.

We have previously shown that KrasG12D cells are competitively eliminated at cell

boundaries with surrounding normal cells, resulting in the elimination of small clusters (*4,*

24). Similar to our previously published results (*4*), the percentage of small clusters (<50

micron^2) significantly decreased in KrasG12D (KC; Fig. 1F, $p=0.0004$) and in p53R172H

(PC; Fig. 1F, $p=0.0316$) tissues over time. We did not see significant differences in cluster

distribution based on size in double mutant (KPC) tissues, compared to controls (Fig. 1F;

$p=0.7740$, small clusters), suggesting double mutant (KPC) cells do not respond to cell-cell

interactions with normal cells. Thus, expression of p53R172H mutations in low numbers of

cells in adult pancreas tissues triggers cell competition *in vivo*, resulting in elimination of

p53R172H cells. Co-expression of mutant p53R172H confers a survival advantage to

KrasG12D cells, abrogating competition in the tissue and allowing KrasG12D cells to avoid

cell elimination.

Transcriptional profiles of non-eliminated mutant cells indicate activation of pro-

survival signals.

We consistently found that ~10% of tissue area remains RFP-labelled in KrasG12D (KC)

and in p53R172H (PC) tissues post the 35-day time-point (Fig. 1B, 1C; (*4*)), indicating that

some mutant cells are never eliminated from the pancreas *in vivo*. Consistent with our

previously published report (*4*), we found rare Alcian blue positive lesions (SFig. 1A, left

panels) in 6/9 KrasG12D (KC) mice at 168 days p.i., suggesting some non-eliminated

KrasG12D cells progress to pancreatic intraepithelial neoplasia (PanINs) (*25*). We did not

detect PanIN lesions in p53R172H (PC) tissues. As expected, PanIN lesions were detected

more frequently in 5/5 KPC tissues analysed (SFig. 1A, right panel) and 4/5 mice developed

PDAC tumours by 168 days p.i.

We hypothesised that non-eliminated mutant cells override competitive cell

elimination signals via common mechanisms. A deeper understanding of how KrasG12D

cells avoid cell elimination may inform on the biology of early tumorigenesis. To gain an

unbiased insight into how non-eliminated mutant cells survive in tissues we performed bulk

RNA sequencing (Fig. 2A). Whole pancreata were harvested from experimental cohorts

(KC/PC/KPC/Control) at 35 days p.i. and digested to cell suspensions. We used anti-lectin

antibodies to enrich for acinar/ductal epithelial cells, as it is generally accepted that PDAC

originates from genetically mutant acinar and/or ductal epithelial cells (*26*), and FACS sorted

leptin positive RFP positive cells. To ensure we reached a minimum of 5,000 viable lectin

positive RFP positive cells per sample, we pooled cell suspensions generated from three

pancreas tissues per genotype. This was repeated to generate three samples per genotype

for sequencing. We compared differential gene expression in non-eliminated mutant cells to

wild type controls and found that KrasG12D cells (KC) predominantly upregulated genes,

while p53R172H (PC) and double mutant (KrasG12D p53R172H, KPC) cells downregulated

genes (Suppl. Table S1, S2). We found few differentially expressed genes in common

between non-eliminated mutant cells when compared to wild type control (see Suppl. Table

S1-S3).

To elucidate the biological pathways underpinning mutant cell retention in pancreas

tissues, we applied gene set enrichment analysis (GSEA_4.2.2; entire Hallmarks and KEGG

MEDICUS databases) and compared non-eliminated KrasG12D (KC) or p53R172H (PC)

transcriptomes to wild type controls. We found that KrasG12D (KC) signatures positively

correlated with MAPK (Fig. 2B, SFig. 2A) and p53 (Fig. 2B, SFig. 2B) signalling pathways,

compared to controls, whereas p53R172H (PC) signatures negatively correlated with p53

(Fig. 2C, SFig. 2C) and with Kras (Fig. 2C) and MAPK (SFig. 2D) signalling. Non-eliminated

KPC signatures also downregulated KRAS signalling (SFig. 2E), suggesting deregulation of

Kras signalling and/or p53 pathway is a general requirement for mutant cells to remain in

tissues. We found that KrasG12D (KC) signatures positively enriched for pathways

associated with stress responses (Fig. 2B, e.g., Oxidative phosphorylation, Hypoxia,

Unfolded protein response, UV response), deregulation of cell cycle (e.g., E2F targets and

G2/M checkpoint) and immune response (e.g., Il2-Stat5 signalling, TNFa via NFkB

signalling). In contrast, pathways associated with p53, cell cycle and inflammatory responses

were negatively enriched in p53R172H (PC) signatures (Fig. 2C). Notably, KrasG12D (KC)

signatures positively enriched for pro-survival and pro-tumorigenic pathways (Fig. 2B, e.g.,

Epithelial to Mesenchymal Transition (EMT), TNF-α signalling, Hypoxia, Angiogenesis,

Notch signalling, Wnt signalling). Gene signatures from both p53R172H (PC; Fig. 2C) and

KrasG12D p53R172H (KPC; SFig. 2E) negatively enriched for apoptosis, suggesting cell

survival is also a requisite for mutant cell retention in tissues.

Gene signatures of non-eliminated KrasG12D cells correlate with PDAC initiation and

with increased stemness and cellular reprogramming.

Mouse models of pancreatic cancer show that expression of oncogenic *Kras* triggers injury

and stress responses, which often translate as cellular reprogramming and a change in cell

fate (*27*). We found that KrasG12D (KC) gene signatures positively correlated with

WP_Pancreatic_Adenocarcinoma_Pathway in GSEA analysis (SFig. 2F). Consistent with

GSEA analyses, we found that gene signatures of non-eliminated PC (p53R172H) and KPC

(KrasG12D p53R172H) cells negatively correlated with pro-tumorigenic pathways (Fig. 2C,

SFig. 2E, 2G, 2H). Using published datasets (*27*), we screened transcriptomes of non-

eliminated cells and found consistent upregulation of pancreas stem and progenitor genes

(e.g., *Nkx6-1*, *Prom-1*, *Pdx-1, Hnf1a*) in non-eliminated KrasG12D (KC) populations (Fig.

3A). We also found canonical markers of spasmolytic polypeptide-expressing metaplasia

(SPEM (*28*); e.g., *Aqp5, Tff2, Dmbt1, Gkn3, Wfdc2*) were upregulated in non-eliminated

KrasG12D (KC) cells (SFig. 2I), as well as acinar genes (e.g., *Ptf1a, Rbpjl*), ductal genes

(e.g., *Sox9, Onecut1*) and mucin genes (e.g., *Muc5a*, *Spdef, Creb3l4, Creb3l1*) (SFig. 2J).

Finally,   we   found   stemness   gene   signatures   enriched   in   KrasG12D   cells

(Ramalho_stemness_up (M9473) and Malta_curated_stemness_markers (M30411) GSEA

gene sets; KC) compared with control (SFig. 2K, 2L). We did not see enrichment of these

gene signatures in p53R172H-expressing populations (SFig. 2I, 2J, 2M, 2N). We

consistently showed that ductal epithelial compartments efficiently eliminate mutant cells and

very few RFP+ cells remain in ducts at 35 days p.i. cells (Fig. 1D; (*4*)). Thus, we assume that

most RFP+ cells isolated for RNA sequencing are acinar in origin. Gene signature data imply

that non-eliminated KrasG12D populations represent differentiated acinar cells and early

embryonic pancreatic progenitors, cells undergoing acinar-ductal metaplasia, gastric pyloric

and intestinal metaplasia. The data also demonstrate that canonical KrasG12D-induced

disease-associated metaplasia is evident in tissues at very early time points (35 days post

activation of the transgene).

Non-eliminated KrasG12D mutant cells express features of cell dormancy.

Several of our data indicated that non-eliminated cells are not actively dividing in tissues

over time, e.g., we found no change in the amount of RFP fluorescence in tissues over time

(Fig. 1C), suggesting RFP+ cells are not expanding. Consistently (*4*), immunostaining for Ki-

67 showed an absence of proliferation in all tissues (Fig. 3B). GSEA analyses identified

enrichment of proliferation-related pathways (e.g., E2F targets, GF-Rtk-Ras-Erk signalling) in

non-eliminated KrasG12D cells compared to controls (KC; Fig. 2B); however, analysis of

specific gene signature data showed a general downregulation of cell cycle genes (KC; Fig.

3C). GSEA analysis also revealed enrichment of pathways indicative of dysregulation of the

cell cycle towards genomic instability (*29, 30*) (Fig. 2B, e.g., G_2M checkpoint, Htlv Tax to

spindle assembly checkpoint, Hbv-Hbx to Ras/Erk signalling, DNA-replication termination).

Cell cycle gene expression profiles of non-eliminated p53R17H cells compared to wild type

controls (PC; Fig. 3C) was like that of KrasG12D signatures and GSEA analyses indicated a

general downregulation of proliferation-related pathways (e.g., E2F targets, Myc targets,

Kras signalling up) in mutant p53-expressing cells (PC; Fig. 2C; KPC; SFig. 2E). Using β-

galactosidase as a readout, we found no evidence of cell senescence in 35-day KrasG12D

(KC) pancreas tissues (SFig. 3A). We also counted rare caspase-3 positive events in all

pancreas tissues analysed (Fig. 3D), suggesting a general absence of apoptosis. We

hypothesised that non-eliminated cells deregulate and/or arrest in the cell cycle. Using p27

(also known as Kip1) immunostaining as a readout of cell cycle arrest (*31*), we scored a

significant increase in p27 positive RFP positive nuclei in KrasG12D (KC) tissues compared

to controls (Fig. 3E; *p*=0.0237), suggesting non-eliminated KrasG12D cells are arrested at

G_1 stage of the cell cycle. Interestingly, non-eliminated KrasG12D (KC) gene signatures

were also enriched for pathways associated with cancer cell dormancy, (e.g., Oxidative

phosphorylation, Unfolded protein response, Hypoxia, Fig. 2B) (*32*). In general, cell

dormancy describes a reversible cell cycle arrested state, often triggered by cell stress (*33-

35*). Using published gene signatures of cell dormancy as reference (*36*), we found cell

dormancy (Fig. 3F) and diapause (SFig. 3B) gene signatures are enriched in non-eliminated

KrasG12D (KC) cells compared to control cells. Diapause is a temporary halt in
embryogenesis when conditions are detrimental to development (*35*). We also found NRF2
pathway (SFig. 3C) and NRF2 target genes (SFig. 3D) were positively enriched in non-
eliminated KrasG12D (KC) cells. NRF2 is a transcription factor, which is activated during cell
dormancy (*37*). Cell dormancy/diapause gene signatures or related pathways were not
significantly enriched in non-eliminated p53R172H (PC) cells (Fig 3F, SFig. 3B). These data
infer that non-eliminated mutant cells deregulate the cell cycle towards an arrested state;
however, two copies of functional p53 are required for cells to adopt a dormant or diapaused
state (*38-40*).
244
Cell cycle arrest prevents mutant cell extrusion *in vitro*.
We hypothesised that mutant cells override cell elimination *in vivo* by arresting in the cell
cycle. Indeed, extrusion of v-Src-expressing cells from normal epithelial sheets is dependent
on the cell cycle (*41*). To functionally test the requirement for cell cycle arrest in RasV12 cell
extrusion, we used previously established cell competition assays and Madin-Darby Canine
Kidney (MDCK) cells expressing GFP-tagged constitutively active oncogenic RasV12 (GFP-
RasV12) (*5, 9*). We have previously shown that when surrounded by normal MDCK cells,
GFP-RasV12-expressing MDCK cells are outcompeted and eliminated from an epithelial
sheet via cell extrusion (*5, 9*). Once extruded, GFP-RasV12 cells rapidly proliferate to form
multicellular aggregates (*5*). In contrast, MDCK cells expressing mutant p53 (p53R175H,
p53R273H) are outcompeted by normal cells via necroptosis (*42*). We induced cell cycle
arrest in GFP-RasV12 cells via *Myc* knockdown and transient transfection of two
independent c-Myc siRNAs (transfected alone or in combination; siMyc-GFP-RasV12). We
chose *c-Myc* knockdown as an approach because c-Myc is a key regulator of the cell cycle
(*43*) and depletion of Myc triggers cell dormancy (*44*). Expression of c-Myc siRNA yielded a
decrease in MYC protein levels, 48h post-transfection (SFig. 3E, 3F), an increase in p21
protein expression and decreased phospho-p38 protein levels (SFig. 3E, 3F) and a marked
reduction in GFP-RasV12 cell confluency (SFig. 3G) compared to GFP-RasV12 cells
expressing scrambled siRNA controls (siScr-GFP-RasV12). Thus, transient depletion of Myc
in GFP-RasV12 cells promotes cell cycle arrest. To assess the impact of cell cycle arrest on
normal-mutant cell-cell interactions, we used our established cell extrusion assay (*5, 9*),
whereby GFP-RasV12 cells were first transfected with siScr or siMyc oligos and then mixed
with normal MDCK cells at 1:50 ratios. At 48h post mixing, cells were fixed, stained with
phalloidin and Hoechst to visualise F-actin and nuclei respectively. We scored apical
extrusion events and found that the proportion of non-extruded GFP-RasV12 cells
significantly increased when GFP-RasV12 cells were depleted for MYC (siMyc-GFP-
RasV12; Fig. 3G, 3H; *p*=0.0003 siMyc1+2-GFP-RasV12 compared to siScr-GFP-RasV12

controls). Immunofluorescence staining using anti-Ki-67 antibodies confirmed that non-
extruded siMyc-GFP-RasV12 cells are not proliferating, whereas the minority of Ki-67
positive cells were extruded (Fig. 3I, 3J). We also used cell confrontation assays (*9*) and live-
cell imaging of normal-mutant interactions across an entire epithelial cell sheet (SFig. 3H).
We previously reported that upon collision with normal MDCK cells, GFP-RasV12 cells are
triggered to retract and segregate away from normal cells, separating via the formation of
smooth boundaries (*9*). We found that GFP-RasV12 cells depleted for MYC (siMyc1+2)
retracted less efficiently than GFP-RasV12 siScr controls (SFig. 3I, SFig.3J; *p*=0.0086).
Thus, RasV12 cells depleted for Myc and exhibiting cell cycle arrest are not extruded or
triggered to segregate by normal cells, suggesting cell cycle arrest protects RasV12 mutant
cells from cell elimination.
283

Non-eliminated KrasG12D cells activate β-catenin independent Wnt signalling *in vivo*.
The Wnt pathway positively regulates cancer cell dormancy (*45, 46*), diapause (*32*) and
stemness (*47, 48*) in different tissue contexts. Our GSEA analyses identified an enrichment
for Wnt pathway activation in non-eliminated KrasG12D (KC) transcriptomes compared to
controls (Fig. 2B, Wnt signalling modulation Lgr/Rspo, Wnt5a-Ror signalling pathway) and
increased gene expression of both Wnt target genes and components of the Wnt pathway
such as ligands and receptors (Fig. 4A). Genes commonly associated with the Wnt/β-catenin
pathway were generally downregulated in KrasG12D (KC) cells, including *Lgr5*, *Myc*, *Sox2*,
Nanog, *Esrrb* and *Wnt3a* (Fig. 4B). Similarly, GSEA analyses indicated low levels of Wnt/β-
catenin signalling in non-eliminated p53R172H (PC) cells (Fig. 2C), which is supported by a
general downregulation of Wnt pathway genes (SFig. 4A). Interestingly, both KrasG12D (KC;
SFig. 4B) and p53R172H (PC; SFig. 4C) gene signatures positively correlated with Wnt5a-
Ror signalling pathway. Wnt ligands and receptors associated with Wnt5a-Ror signalling
were increased in KrasG12D (KC) cells, including *Wnt5a*, *Ror1*, *Ror2*, *Dvl2* and *Jun* (Fig. 4C,
4D and Supplementary table S11).
To investigate active Wnt signalling *in vivo*, we first performed immunofluorescence
for CD44 which is a target of β-catenin dependant and independent signalling (*49-51*).
Consistent with previous reports (46), we observed CD44 labelling at cell membranes in
PanINs (SFig. 4D). We found that CD44 was significantly increased at RFP+: RFP- cell-cell
boundaries in KrasG12D (KC) tissues compared to RFP+: RFP- boundaries in wild type
controls (SFig. 4E, 4F; KC versus control, *p*=0.0009). In contrast, CD44 was significantly
reduced within a cluster of KrasG12D cells (KC; SFig. 4E, 4G; *p*=0.0455). We asked
whether increased CD44 at mutant-normal cell-cell boundaries was dependent on
competitive cell-cell interactions. We quantified CD44 fluorescence at cell-cell contacts
between RFP+/RFP- cells in KC tissues fixed following high dose tamoxifen induction (KC

Megadose). We previously reported that KrasG12D cells are not eliminated from megadose
tissues, where mutant cells occupy ~80% of the tissue (*4*). We found that CD44 levels were
significantly lower at mutant-normal cell boundaries in megadose tissues compared to
KrasG12D low dose tissues (KC versus KC Megadose; SFig 4E, 4F; *p*=0.0047), while CD44
levels were significantly higher between KrasG12D cells in megadose tissues (SFig. 4E, 4G;
p=0.0341). Similarly, we found that CD44 was significantly increased at RFP+: RFP- cell-cell
boundaries in p53R172H tissues (PC) compared to RFP+: RFP- boundaries in wild type
controls (SFig. 4H; PC versus control, *p*=0.047). CD44 was significantly reduced within a
cluster of p53R172H cells (SFig. 4I, PC versus control, *p*=0.0397). We next asked whether
readouts of Wnt5a signalling were evident in non-eliminated KrasG12D cells in fixed
pancreas tissues. Using quantitative immunofluorescence protocols and commercial
antibodies, we observed significant increase in WNT5A (Fig. 4E; *p*=0.0491) and DVL2 (Fig.
4F; *p*=0.0110) protein in KrasG12D (KC) cells relative to controls, suggesting Wnt5a
signalling is active in non-eliminated KrasG12D cells *in vivo*. In addition, we performed
immunofluorescence for β-catenin in tissues fixed at 35 days p.i., using nuclear localisation
as a surrogate marker of activity of the Wnt/ β-catenin signalling pathway. Nuclear β-catenin
was undetectable in nuclei of both control and KrasG12D (KC) cells (Fig. 4G, 4H; *p*=0.3035),
indicating that Wnt/β-catenin signalling is not active in non-eliminated KC cells, consistent
with our transcriptional analysis. Instead, β-catenin was significantly increased at RFP+:
RFP- cell-cell contacts in KrasG12D (KC) tissues compared to controls (Fig. 4G, 4I;
p=0.0347). Moreover, levels of E-cadherin were also significantly elevated at RFP+: RFP-
boundaries in KrasG12D (KC) tissues compared to controls (Fig. 4J, 4K; *p*=0.0348).
Interestingly, we also detected increased mRNA expression of atypical *cadherin-6* in non-
eliminated KrasG12D cell signatures (KC; SFig. 5A) and increased levels of CADHERIN-6
(K-cadherin) in RFP+ clusters in KrasG12D (KC) tissues only (SFig. 5B, 5C; *p*=0.0474).
Together, our data suggest that β-catenin-independent Wnt signalling is active, specifically
in non-eliminated KrasG12D cells and at KrasG12D-normal cell-cell contacts *in vivo*. Our
data also infer that active Wnt signalling increases cohesiveness and/or stability of E-
cadherin-based cell-cell adhesions *in vivo*, specifically at cell-cell boundaries between non-
eliminated mutant cells and normal neighbours.
339

Wnt5a inhibits RasV12 cell extrusion *in vitro* by stabilising E-cadherin-based cell-cell
adhesions.
To establish whether Wnt5a directly inhibits RAS mutant cell elimination, we returned to
MDCK cell extrusion assays (*5, 9*). We first assessed the effect of Wnt5a treatment on
normal-RasV12 cell-cell interactions. We found that the proportion of non-extruded GFP-
RasV12 cells significantly increased in the presence of Wnt5a (Fig. 5A, right panel, yellow

arrowheads; 5B, yellow bar; *p*=0.0065) compared to PBS treated controls (Fig. 5A, left

panel, white arrowheads; 5B, black bar). Remarkably, addition of Wnt3a had no effect on

RasV12 cell extrusion rates compared to PBS controls (Fig. 5A, middle panel, white

arrowheads; 5B, pink bar; *p*=0.4488). In general, Wnt3a and Wnt5a activate the β-catenin-

dependent and β-catenin-independent pathway respectively (*52, 53*), suggesting β-catenin-

independent Wnt signalling inhibits RasV12 cell extrusion.

Cell extrusion requires dynamic remodelling of E-cadherin-based cell-cell adhesions

at normal-mutant cell-cell boundaries (*5, 54, 55*). Considering both E-cadherin and β-catenin

protein were significantly increased at normal-mutant cell-cell boundaries *in vivo* (Fig. 4G, 4I-

4K), we asked whether treatment with exogenous Wnt ligands modulates adherens junctions

at normal-mutant cell-cell boundaries in MDCK assays. We analysed E-cadherin and β-

catenin by immunofluorescence in Wnt-treated co-cultures and focused on cell-cell

interactions between GFP-RasV12 that had not yet been extruded and direct normal

neighbours. In the presence of Wnt3a, β-catenin and E-cadherin staining appeared

diffuse/weak and sometimes absent at normal-mutant cell-cell contacts (Fig. 5C, middle

panels), similar to PBS-treated controls (Fig. 5C, top panels). In contrast, E-cadherin and β-

catenin staining was more defined at normal-mutant cell-cell contacts in Wnt5a treated cells

(Fig. 5C, lower panels). Quantitative analysis showed that in Wnt5a treated cells, E-cadherin

(Fig. 5D; *p*=0.0116) and β-catenin (Fig. 5E; *p*=0.0026) significantly increased at normal-

RasV12 cell-cell contacts. In contrast, Wnt3a treatment had no significant effect on E-

cadherin (Fig. 5D; *p*=0.2137) or on β-catenin (Fig. 5E; *p*=0.8111) at RasV12-normal cell-cell

contacts when compared to PBS-treated controls. Treatment with Wnt5a significantly

increased the levels of E-cadherin (SFig. 5D; *p*=0.0003) but had no effect on β-catenin levels

(SFig. 5E; *p*=0.6049) detected at cell-cell contacts between RasV12 cells, suggesting Wnt5a

signalling specifically modulates E-cadherin at cell-cell contacts. Consistently Wnt3a had no

significant effect on E-cadherin or β-catenin at cell-cell contacts between RasV12 cells

(SFig. 5D; *p*=0.0584; SFig. 5E; *p*=0.2828). Moreover, in cell confrontations assays (*9*),

Wnt5a treatment significantly reduced RasV12 cell speed (SFig. 5F; *p*=0.0268) compared to

PBS treated controls, whereas normal MDCK cell speed was unaffected (SFig. 5F;

p=0.8202), implying Wnt5a signalling specifically affects RasV12 cell cohesion.

E-cadherin endocytosis is required for apical extrusion of RasV12 cells from normal

MDCK cell sheets (*55*); therefore, we asked whether Wnt5a abrogates this process to

prevent cell extrusion. We imaged RasV12-normal cell-cell contacts at 16h (prior to cell

extrusion events (*55*)) using super resolution microscopy. At this time point, E-cadherin

staining was weakly visible/diffuse at RasV12-normal cell-cell contacts in PBS-treated

controls and was often detected in distinct intracellular puncta in both RasV12 and normal

cells (Fig. 5F, top panels). In contrast, Wnt5a treated cells showed strong, defined E-

cadherin at cell-cell contacts and E-cadherin positive puncta were less visible in RasV12

cells and neighbouring cells (Fig. 5F, lower panels). Quantification of fluorescence intensity
of intracellular E-cadherin showed a significant reduction in RasV12-cells treated with Wnt5a
compared to PBS-treated controls (Fig. 5G; p=0.0051), suggesting Wnt5a treatment blocks
E-cadherin recycling/endocytosis at RasV12-normal boundaries. Finally, we immunostained
for Caveolin-1 (CAV1) a key regulator of endocytosis (*56*) and cell extrusion (*57, 58*). CAV1
was detected in normal MDCK and RasV12 cells at cell-cell junctions (SFig. 5G). At the
timepoint when endocytosis is evident in cells (Fig. 5F), CAV1 fluorescence was significantly
reduced at RasV12-normal cell-cell contacts in Wnt5a treated cells compared to PBS
controls (SFig. 5H; p=0.0117). Taken together our data show that Wnt5a stabilises E-
cadherin-based cell-cell adhesion at normal-mutant boundaries potentially by preventing E-
cadherin internalisation. Increased E-cadherin-based cell-cell adhesion correlates with a
significant reduction in RasV12 cell extrusion and increase in RasV12 cell cohesion.

Wnt signalling is required to promote retention of mutant cells *in vitro* and *in vivo*.
Next, we set out to test whether Wnt signalling is required to prevent mutant cell elimination.
We first repeated cell extrusion experiments using siScr-GFP-RasV12 or siMyc 1+2-GFP-
RasV12 MDCK cells mixed with normal MDCK cells at 1:50 ratios, allowed cells to form cell-
cell interactions and then treated with the porcupine inhibitor WNT-974 (or DMSO).
Porcupine is an acyltransferase enzyme required for the lipidation and trafficking of Wnt
proteins, which is a necessary step for ligand secretion (*59*). By inhibiting porcupine, Wnt
ligands cannot be secreted and therefore cannot signal to cells. In MDCK cells,
overexpression of RasV12 induces expression and secretion of Wnt5a in a porcupine-
dependent manner (*60*), suggesting porcupine inhibition in our experiments blocks secretion
of endogenous Wnt, including Wnt5a. The proportion of non-extruded siMyc 1+2-GFP-
RasV12 cells significantly reduced in the presence of WNT-974 (Fig. 6A, 6B; p<0.0005),
compared to siScr-GFP-RasV12 treated cultures (Fig. 6A; p=0.6912). In cell confrontation
assays, we found that WNT-974-treated siMyc 1+2-GFP-RasV12 cells efficiently retracted
(SFig. 6A, 6B) and segregated from normal cells, compared to DMSO-treated siMyc 1+2-
GFP-RasV12 cells. Quantification of the linearity of the boundary between colliding sheets of
cells (*9*) confirmed this observation (SFig. 6C; WNT-974-treated cells: p=0.0020).
Consistently, the effects of Wnt5a on RasV12 cell extrusion rates were reversed in
the presence of porcupine inhibitor (WNT-974; Fig. 6C, 6D, hatched bars; p=0.0114) and in
the presence of the Frizzled (Fzd) receptor antagonist OMP-18R5 (*61*) (Fig. 6C, 6E, striped
bars; p=0.0004) compared to DMSO-treated controls, resulting in a significant decrease in
non-extruded GFP-RasV12 cells. Treatment with WNT-974 inhibitor also reversed Wnt5a-
induced effects on E-cadherin localisation (Fig. 6F). As expected, Wnt5a induced a
significant decrease in the level of intracellular E-cadherin fluorescence in RasV12 cells (Fig.

6G, circles; *p*=0.0048). In the presence of WNT-974 inhibitor, intracellular E-cadherin
fluorescence significantly increased (Fig. 6G, squares; *p*=0.0044 compared to Wnt5a-treated
cells) and returned to levels comparable to PBS treated controls, implying that Wnt signalling
is required to prevent E-cadherin internalisation and endocytosis. In cell confrontation
assays, Wnt5a treatment significantly decreased RasV12 cell speed (SFig. 6D, open bar;
p=0.0156); RasV12 cell speed significantly increased in the presence of WNT-974 inhibitor
(SFig. 6D, hatched bar; *p*=0.0058).
Finally, we set out to functionally test whether active Wnt signalling is necessary for
mutant cell retention *in vivo*. Using 6-8-week-old adult KrasG12D (KC) mice, we first induced
low level recombination in tissues, waited for 35 days p.i. to allow for cell competition to
proceed and for most mutant cells to be eliminated, and then administered the porcupine
inhibitor WNT-974 (or DMSO in corn oil vehicle) for a period of 4 weeks (Fig. 6H). To
validate Wnt inhibition in tissues *in vivo*, we quantified CD44 fluorescence at mutant-normal
cell boundaries and within RFP+ clusters in KrasG12D (KC) pancreas tissues treated with
WNT-974 or vehicle. We found that levels of CD44 fluorescence significantly decreased in
WNT-974 treated tissues compared to vehicle (SFig. 6E, *p*=0.0467), suggesting Wnt activity
is reduced following treatment with porcupine inhibitor. Using global RFP fluorescence per
tissue area as a readout of mutant cell fate, we measured a significant decrease in RFP
fluorescence in the WNT-974-treated KrasG12D (KC) tissues compared to vehicle-treated
animals (Fig. 6J, 6I; *p*=0.0003). We also found that porcupine inhibition induced a significant
decrease in RFP fluorescence in p53R172H (PC) tissues compared to vehicle treated
controls (SFig. 6F, 6G; *p*=<0.0001). Thus, inhibition of Wnt pathway induces the expulsion of
KrasG12D or p53R172H cells from pancreas tissues *in vivo*; however, Wnt inhibition is not
sufficient to remove all mutant cells, suggesting Wnt-independent mechanisms are also
required. Nonetheless, our data support a model whereby activation of Wnt signalling allows
mutant cells to avoid cell expulsion and to survive in the pancreas.
447
448

Discussion

Tissue homeostasis is fundamental to healthy aging of an organism and cell competition is an important regulator of tissue health. Here, we extend our previously published results (*4*) to show that like KrasG12D-expressing cells, genetically mutant p53R172H-expressing cells compete with normal cells for survival in the adult pancreas and are often expelled from the tissue. We report that cell expulsion is inefficient, and a proportion of KrasG12D- or p53R172H-expressing cells are never eliminated from the tissue, suggesting these cells have a survival advantage. Transcriptomic analyses of non-eliminated mutant cells revealed Wnt signalling as a key regulator of mutant cell fate *in vivo*. Gene expression data indicate Wnt/β-catenin signalling is inactive and instead, non-canonical Wnt5a signalling is active in non-eliminated mutant cells. Indeed, readouts of non-canonical Wnt signalling are increased in non-eliminated KrasG12D cells and at KrasG12D-normal cell-cell boundaries *in vivo*. Wnt5a signalling (and not β-catenin-dependent Wnt3a signalling) inhibited apical cell extrusion of RasV12 cells from epithelial cell sheets *in vitro*, and this effect was reversed upon inhibition of porcupine/Fzd activity. In pancreas tissues, inhibition of Wnt signalling promoted expulsion of non-eliminated mutant cells, suggesting Wnt signalling is a general mechanism activated by mutant cells to override cell elimination mechanisms in the pancreas *in vivo*.

Apical cell extrusion is an active process that requires direct interaction with normal cells to expel genetically mutant cells from an epithelium (*62*). Here, we show that Wnt5a blocks apical extrusion of RasV12 cells, allowing mutant cells to remain in the epithelium. Mechanistically, we propose that Wnt5a stabilises E-cadherin-based cell-cell adhesions at RasV12-normal cell-cell contacts, and this prevents RasV12 cell extrusion. E-cadherin is a key modulator of apical cell extrusion: RasV12 cells are not extruded when E-cadherin is depleted in normal neighbours (*5*), suggesting E-cadherin-based cell-cell adhesion is required for RasV12 cell extrusion. On the other hand, our data support previous reports (*55*) that E-cadherin is internalised as discrete puncta at RasV12-normal cell-cell boundaries, suggesting E-cadherin must be dynamically remodelled for cell extrusion to occur. Indeed, inhibition of Rab5-mediated endocytosis prevents apical extrusion of RasV12 cells from MDCK cell sheets (*55*). Here, we show that in the presence of Wnt5a, E-cadherin puncta are not detected in mutant cells; instead, the levels of E-cadherin and β-catenin are increased at RasV12-normal cell-cell contacts. Moreover, inhibition of Wnt signalling (via porcupine inhibitor or Fzd antagonism) restores appearance of E-cadherin puncta at RasV12-normal cell-cell contacts and apical extrusion of RasV12 cells. In pancreas tissues, we previously showed that E-cadherin is decreased at KrasG12D-normal cell-cell adhesions, specifically at timepoints prior to KrasG12D cell elimination (*4*). Here, we extend these findings to show

that E-cadherin and β-catenin are increased at cell-cell contacts between non-eliminated
KrasG12D cells and normal neighbours. Indeed, we have shown that under conditions
where RAS mutant cell elimination is prevented, E-cadherin increases at normal-mutant
boundaries (*4*). Together our data support a model whereby stable cell-cell adhesion
induced by Wnt5a signalling, prevents cell extrusion. Interestingly, we also found that
caveolin-1 (CAV1), an important driver of RasV12 cell extrusion (*57, 58*) is also decreased at
RasV12-normal cell-cell contacts following Wnt5a treatment. In development, Wnt5a
signalling regulates cell cohesion and tissue fluidity via caveolin-dependent and clathrin-
dependent endocytosis (*63*). Whether Wnt5a directly modulates endocytosis of E-cadherin
at RasV12-normal cell-cell boundaries requires further investigation. Similar to our results, in
human epithelial cells Wnt5a induces β-catenin membrane localisation and association with
E-cadherin increasing intercellular adhesion (*64*).

We also show that RasV12 cells in a cell cycle arrested state were not extruded from
MDCK monolayers. In pancreas, non-eliminated KrasG12D cells stained positive for p27,
and upregulated cell dormancy signatures and pathways known to induce cell dormancy
(e.g., Unfolded protein response, Wnt5a-ROR signalling, Hypoxia) (*65*). Thus, exiting the cell
cycle protects KrasG12D cells from cell elimination *in vivo* and *in vitro*. Cell stress responses
are major triggers of cell dormancy (*66*). It is possible that non-eliminated KrasG12D cells
enter a dormant state in response to competition-induced stress, overriding cell elimination
and surviving in tissues. Studies in *Drosophila melanogaster* models demonstrate that cell
competition outcomes are a consequence of cellular stress responses activated in 'loser'
cells (*67-72*). Taking these studies into consideration, our transcriptomic data infer that non-
eliminated KrasG12D cells upregulate gene signatures reminiscent of 'loser' phenotypes
(e.g., Glycolysis (*7*), Unfolded protein response (*73*), p53 (*12, 42, 74-77*), oxidative stress
(*78*), NRF2 pathway (*68, 78, 79*)). Thus, cell dormancy may act to restrict expansion of 'unfit'
KrasG12D cells in the absence of an efficient tissue clearance mechanism and without
drastically reducing cell numbers. This would be particularly relevant to adult pancreas,
which has limited proliferative capacity (*80*). Indeed, we never observe compensatory
proliferation in neighbours following mutant cell clearance (*4*) (Fig. 3B). Future studies are
required to determine whether normal-mutant interactions in the adult pancreas induce
cellular stress responses in genetically mutant cells and whether this in turn activates cell
dormancy. Interestingly, Wnt5a has been shown to induce and maintain cancer cell
dormancy in metastatic niches by negatively regulating canonical Wnt/β-catenin signalling
(*45, 46*). An exciting future direction of this work will be to elucidate whether the cell
dormancy phenotype we describe here is regulated via Wnt-dependent mechanisms
described for dormant metastatic cancer cells. Transcriptional profiling of non-eliminated

KrasG12D cells support a mechanism whereby Wnt signalling at KrasG12D-normal cell-cell
boundaries is not driving cell proliferation. We also show that inhibition of Wnt restores cell
extrusion and cell segregation of cell cycle arrested RasV12 cells from normal cells *in vitro*,
suggesting a mechanistic link between active Wnt and cell dormancy *in vivo* is plausible.

TP53 is an important regulator of cell competition. In MDCK cells (*76, 81*) and in
developing embryos (*74, 77*), mutant-normal cell competition triggers an elevation in p53
levels and p53 activity in prospective 'losers' and this is required for mutant cell elimination
(*77*). Studies in the adult hematopoietic system (*82, 83*) and oesophagus (*13, 84*) illustrate
that competition outcomes are dependent on relative rather than absolute levels of p53 and
are apparent only when cells are under mild stress. Whether cell competition outcomes we
describe here are triggered by differentials in p53 activity between mutant and wild type
neighbours requires further investigation. Interestingly, we find that co-expression of
p53R172H and KrasG12D in the same cell (KPC), provides all mutant cells with a survival
advantage and 'double mutant' KPC cells are never eliminated. In MDCK cells, ectopic
expression of p53R175H/p53R273H mutations in few cells surrounded by normal cells
triggers necroptosis and elimination of the p53 mutant cells (*42*); however, expression of
p53R273H mutations in RasV12 monolayers prevents p53 mutant cell death (*42*). Recent
studies in zebrafish larval skin (*85*) show that while RasG12V-expressing cells are actively
eliminated, cells expressing both RasG12V and Tp53R175H mutations are not eliminated
and are viable *in vivo*. Both studies support our observations and suggest 'double mutant'
KPC cells may override cell death signals *in vivo*. Indeed, our RNA sequencing data suggest
cells expressing p53R175H mutations downregulate the apoptosis pathway (Fig. 2B, SFig.
2E).

KRAS and *TP53* mutations are key drivers of pancreatic ductal adenocarcinoma.
Single cell profiling of pancreatic tumorigenesis in mouse models demonstrates that *Kras*
mutant cells are very heterogeneous and show signatures of tumorigenesis even before pre-
malignant lesions are established (*86*). Our RNA sequencing data confirm this early
transformation and cellular heterogeneity profile of KrasG12D cells. Pancreatic cancer is a
devastating disease, which is generally diagnosed at late and incurable stages. An improved
understanding of the biology of how pancreatic cancer starts and grows in adult tissues will
inform the development of new early detection methods. Our results challenge the current
understanding of how cancers start in the adult pancreas and suggest that genetically
mutant cells must override tissue homoeostasis mechanisms to survive in tissues, prior to
transformation/expansion.

555
556
557

Author Contributions

Conceptualization, CH; Data curation, BS-B; Experimental design, BS-B, CH; Funding

acquisition, CH; Investigation & Analysis: BS-B, CH, JPM, OJS; Methodology, BS-B, MA;

Validation, BS-B; Writing original draft, BS-B, CH; Writing review & editing, BS-B, CH;

Resources, JPM, OJS; Supervision, CH.

Acknowledgements

We thank Angela Marchbank and colleagues at School of Biosciences Genome Hub (Cardiff

University) for technical support with RNA sequencing experiments. We thank Toby Phesse

and Valerie Meniel (European Cancer Stem Cell Research Institute, School of Biosciences,

Cardiff University) for Wnt antagonists, PCR primers and helpful discussions on the Wnt

pathway. We thank F. Afonso (European Cancer Stem Cell Research Institute, School of

Biosciences, Cardiff University) for critical reading of the manuscript and helpful discussions.

B.S. and M.A. were supported by Cancer Research UK (CRUK) Early Detection project

grant (A27838) to C.H. O.J.S. and J.P.M. were supported by Cancer Research UK core

funding to the CRUK Scotland Institute (A17196 and A31287) to O.J.S. laboratory (A21139)

and to J.P.M. laboratory (A29996). B.S-B. is currently funded by a Pancreatic Cancer UK

Career Foundation Fellowship (176937364). This study was supported by Cancer Research

UK (CRUK) Early Detection project grant (A27838) to C.H. The manuscript was critically

reviewed by Catherine Winchester (CRUK Scotland Institute).

Methods

<u>Mouse lines, Induction of Cre recombinase *in vivo*, Inhibition of Wnt *in vivo*.</u>

Pdx1-Cre^{ERT}(*87*)*, LSL-Kras$^{G12D/+}$*(*88*)*,* LSL-*Trp53$^{R172H/+}$* (*89*)*, Rosa26$^{LSL-tdRFP}$* (*90*) mouse lines

have all been previously described. Animals were housed in conventional pathogen-free

animal facilities and experiments were conducted in accordance with UK Home Office

regulations (ASPA 1986 & EU Directive 2010) under the guidelines of Cardiff University

Animal Welfare and Ethics Committee. Mice were genotyped by PCR analysis following

standard methods and using primer sequences in (*4*). Both male and female 6-8-week-old

experimental and control mice were injected with intraperitoneal injection of tamoxifen in

corn oil as previously described (*4*). Low recombination levels were obtained with a

tamoxifen dose of 1µg/40g bodyweight once, while high recombination (megadose) was

induced by three injections of 9mg/40g over 5 days (*4*). Pancreata were harvested at 7-, 35-,

70- or 140-days post tamoxifen induction. To test the functional role of Wnt signalling *in*

vivo, 6-8-week-old experimental and control mice were induced with low dose tamoxifen,

aged to 35-days post induction of Cre recombinase (p.i.) and treated with WNT-974

1.5mg/kg or vehicle (DMSO) in corn oil by oral gavage, five days per week for four weeks. At

the end of the treatment pancreatic tissue was harvested and treated as described below.

No statistical method was used to pre-determine sample size. For most animal studies,

experiments were not randomized, and investigators were not blinded to allocation during

experiments. To test the functional role of Wnt signalling mice were randomly allocated into

vehicle or WNT-974 treated groups. All experiments were reproduced using at least three

animals of each genotype.

<u>Tissue staining.</u>

For histological observation, pancreas was harvested at specified timepoints and fixed in

10% neutral buffered formalin overnight at 4ºC. Then pancreas was dissected into two

pieces (head-body/body-tail) and embedded either in paraffin or OCT matrix. FFPE

embedded pancreas was sectioned (7µm thickness) and stained with specific antibodies

(table 1). Anti-RFP, anti-cleaved caspase-3, and anti-Ki-67 staining were performed via

immunohistochemistry (IHC) in FFPE tissue sections. Tissues sections were dewaxed and

rehydrated. For antigen retrieval, tissue sections were incubated for 15 minutes at 37ºC in

20ug/ml Proteinase K diluted in TBS-T or boiled in citrate buffer pH 6 for 15 minutes

(specified in table 1). Tissues were then blocked with 3% H_2O_2 for 20 or 10 minutes at room

temperature (depending on the antigen retrieval method) respectively and 5% NGS TBS-T

for 30-60 minutes at room temperature. Sections were then incubated overnight at 4ºC with

the primary antibody diluted in 5% NGS TBS-T (see dilutions in table 1). Tissues were

stained with the secondary antibody ImmPRESS goat anti-Rabbit (MP-7451, Vector Labs)

for 60 minutes at room temperature, followed by DAB chromogen (Peroxydase substrate kit,

Vector Labs). Sections were dehydrated and mounted with DPX (Sigma-Aldrich) and

scanned using the Axio Scan Z1 slide scanner (Zeiss) 20X magnification.

Immunofluorescence (IF) co-staining using either anti-CD44 and anti-RFP (Rockland) was

performed in FFPE tissues slices using the same antigen retrieval and primary antibody

incubations as described for IHC, followed by Alexa fluor secondary antibody incubation for

60 minutes at room temperature and Hoechst (ThermoFisher) (table 1) and mounted using

Mowiol (Sigma-Aldrich).

For β-catenin E-cadherin, anti-K-cadherin, and RFP (Creative Diagnostics) IF staining,

pancreas was embedded in butyl-methyl methacrylate plastic (BMMA) under UV following

dehydration and resin infiltration (*4*). Tissue was sectioned in 2µm thick slices and

rehydrated. Tissue staining followed the same protocol as IF in FFPE sections. Antigen

retrieval and antibody concentrations are listed in table 1.

Pancreas embedded in OCT matrix was sectioned (10µm thickness), permeabilized with

0.05% SDS (Sigma-Aldrich) and 0.05% Triton X100 (Sigma-Aldrich) solution and stained

with anti-p27, anti-Wnt5a, anti-Dvl2 primary antibodies (table 1) for 2 hours at room

temperature, Alexa fluor secondary antibody incubation for 60 minutes at room temperature

and Hoechst (ThermoFisher) (table 1), and mounted using Mowiol (Sigma-Aldrich).

Global levels of endogenous RFP were measured in 10µm formalin fixed OCT frozen

sections, as previously described (*4*). RFP positive area was averaged from 3 tissue slices

per mouse separated by at least 20µm between each section. Fluorescence imaging was

carried out on a Zeiss LSM710 confocal microscope.

To assess the presence of PanIN lesions, FFPE sections were stained with Alcian blue as

previously described (*4*). β-galactosidase activity was determined using the Cell Signaling

Senescence β-galactosidase Staining kit (9860) using the manufacturers' instructions.

<u>Image analysis.</u>

The percentage of RFP+ tissue area was determined using endogenous RFP levels

quantified as described in (*4*). Using Fiji (RRID:SCR_002285) software,(*91*) positive areas

were thresholded using RFP fluorescence. Tissue autofluorescence was used to determine

total tissue area. The percentage of RFP+ ducts were analysed using E-

cadherin/RFP/Hoechst-stained sections to identify ducts. At least 20 ducts were quantified

per mouse and ducts were considered positive when they contained at least one RFP+ cell.

The percentage of RFP+ cells in pancreatic islets were scored in FFPE IHC sections stained
with anti-RFP antibody and using 'positive cell count' tool in QuPath-0.4.1 software. RFP
positive cluster size, perimeter and area were determined in FFPE IHC sections stained with
anti-RFP antibody and using QuPath-0.4.1 software by analysing shape features of RFP
positive (DAB positive) areas. Ki-67 positive and Cleaved Caspase-3 positive cells were
analysed using QuPath-0.4.1 positive cell count feature.

Wnt5a fluorescence intensity in the tissue was quantified by masking the tissue area
(AF568) and measuring mean fluorescence intensity for Wnt5a (AF488). CD44, β-catenin
and E-cadherin fluorescence intensity at the border between RFP positive and negative cells
were quantified by measuring mean fluorescence intensity of 2.43x2.43-pixel circles (ROIs)
along the membrane (RFP+/RFP-) separated by approximately 5 pixels. CD44, Dvl2 and K-
cadherin fluorescence in clusters was quantified by masking RFP+ (AF568) areas and
quantifying mean fluorescence intensity for CD44/Dvl2/K-cadherin (AF488). β-catenin
fluorescence in nuclei was quantified by masking nuclei (Hoechst) and quantifying mean
fluorescence intensity for β-catenin (AF488) in RFP+ cells. To avoid noise associated with
tissue autofluorescence, fluorescence intensity measurements were made relative to image
global mean fluorescence intensity (AF488).

Pancreas digestion and RNA Sequencing
Pancreas tissues were harvested 35 days post low-dose tamoxifen induction. To reach the
minimum of 5000 cells and a minimum of 100ng RNA/sample for bulk RNA sequencing, we
combined three pancreas tissues of the same genotype to produce one sample per
genotype. Each sample was replicated three times to generate three biological repeats per
genotype for sequencing. Pancreas tissues were digested as previously described (*92*).
Using an EGTA-based buffer the pancreas from each mouse was inflated and slightly
digested with collagenase. The tissue was chopped and further digested using a calcium-
based buffer with collagenase. Single cell digested tissue was stained with lectin PNA-A488
(ThermoFisher) and DAPI (ThermoFisher Scientific). DAPI negative, lectin positive, RFP
positive cells were sorted using the FACS Aria Fusion sorter. A minimum of 5,000 cells were
sorted for each sample. RNA was extracted using the RNeasy Micro kit (Qiagen) following
the manufacturer's recommendations. Paired-end sequencing was performed using Sanger
sequencing Illumina 1.9. Sequenced reads Fastq files were quality-checked using FastQC.
Reads were aligned to the mouse genome (Ensembl-GRCm38 *Mus musculus*) using the
STAR package, and reads were counted using the FeatureCounts package, after removing
duplicates using MarkDuplicates from GATK. Differential expression was normalised and
calculated using Deseq2, comparing the different genotypes against the Control. Finally,

GSEA 4.2.2 software was used for enrichment analysis of different pathways and GraphPad

Prism 10.0.2 for Heatmap graphs and normalised enrichment score (NES) graphs. The

RNAseq data generated during this study are available at GEO ID: XX.

<u>MDCK cell lines and *In vitro* experiments.</u>

To assess the interactions between RasV12 and MDCK cells, MDCK-pTR GFP–RasV12

cells were combined with parental MDCK cells at a ratio of 1:50. Treatment with tetracycline

(2µg/mL) induces GFP-RasV12 expression in MDCK-pTR GFP–RasV12 cells (*5, 9*). Cells

were plated at a density of 2.5 x 10^5 cells per well in MW24 plates (Corning) carrying 18mm

diameter glass cover slips (VWR). Mixed cells were incubated for 30h or 48h at 37ºC before

fixed with 4% Formaldehyde (ThermoFisher Scientific). Cells were then permeabilized and

stained (table 1). GFP-RasV12 cells were considered non-extruded when more than 30% of

their cytoplasm was basally protruded. Three technical replicates per experiment in three

experiments (i.e., using three independent frozen stocks of parental and GFP-RasV12

MDCK cells) were performed and at least 150 GFP-RasV12 cells were quantified per

replicate.

For c-Myc silencing, tetracycline induced GFP-RasV12 cells were transfected with 100ng

siRNA oligos targeting *Myc* using Lipofectamine 3000 (Invitrogen) (siRNA siMyc1:

AAGACGUUGUGUGUUCGCCUC – as GAGGCGAACACACAACGUCUU, siRNA siMyc2:

AAUUUCAACUGUUCUCGCCGC – as GCGGCGAGAACAGUUGAAAUU and siScr)[24]. For

cell extrusion and proliferation experiments, GFP-RasV12 cells were trypsinized and mixed

with parental MDCK cells 24h after siMyc1+2 transfection and were then fixed following a

further 48h.

For experiments comparing PBS, Wnt3a and Wnt5a treatments effect, RasV12 cells were

mixed 1:50 with parental MDCK cells and once cells were set, induced with tetracycline. Two

hours post-tetracycline induction cells were treated with 1µg/mL of human recombinant

Wnt3a (Peprotech), 100ng/mL of human recombinant human/mouse Wnt5a (R&D Systems)

or PBS (control-Sigma-Aldrich). Cells were fixed 30h after PBS, Wnt3a or Wnt5a treatment

for cell extrusion analysis and for analysis of E-cadherin/β-catenin at the RasV12-MDCK

contact or within GFP-RasV12 cell clusters. E-cadherin endocytosis and Caveolin

fluorescence were assessed by fixing cells 16h after PBS/Wnt5a treatment.

For Wnt pathway inhibition experiments, tetracycline induced GFP-RasV12 cells were

transfected with siMyc1+2/siScr or treated with recombinant Wnt5a/PBS for 24h and then

mixed with parental MDCK cells. Wnt5a treatment was maintained during the whole

experiment. WNT-974 (Stratech) porcupine inhibitor (1μM) or *OMP*-18R5 (10mg/mL) was
added to the medium 8h after GFP-RasV12 and MDCK cells were mixed and plated. DMSO
treated GFP-RasV12 cells were used as control for both inhibitors. To assess cell extrusion,
cells were fixed 48h after plating. E-cadherin endocytosis was assessed by mixing RasV12
cells with MDCK parental cells, then RasV12 expression was induced by tetracycline and
treated with PBS or Wnt5a 2h after. Four hours after PBS/Wnt5a treatment, cells were
treated with DMSO or WNT-974. Cells were fixed 16h after PBS/Wnt5a treatment.

Immunofluorescent staining was performed by permeabilization of cells with 0.05% SDS
(Sigma-Aldrich) and 0.05% Triton X100 (Sigma-Aldrich) solution for 15 minutes. Cells were
stained with anti-Ki-67, anti-E-cadherin, anti-β-catenin or Caveolin primary antibodies for 2h
at room temperature, Alexa fluor secondary antibody incubation for 60 minutes at room
temperature and Hoechst (ThermoFisher) (table 1) and mounted using Mowiol (Sigma-
Aldrich). For proliferation analysis at least 30 extruded and 30 non-extruded cells were
quantified in siMyc 1+2 treated cells per experimental replicate. For E-cadherin, β-catenin or
Caveolin at least 10 GFP-RasV12 cell-clusters per experimental replicate were quantified.
Fluorescence intensity at the border between RasV12 and parental MDCK cells were
quantified by measuring mean fluorescence intensity of 2.27x2.27-pixel circles (ROIs) along
the membrane separated by approximately 5 pixels. E-cadherin endocytosis was measured
quantifying the mean fluorescence intensity of the cytoplasm of each RasV12 cell. At least
10 cells were quantified in three technical replicates per experiment in three experiments.

Confrontation assays and migration speed analysis were carried out as described (*9*). 1x10[4]
cells were plated in inserts (Ibidi) in MW24 plates (Corning), allowed to form monolayers (8h)
before removing the insert and moving the plate to IncuCyte S3 (Sartorius) for live cell
imaging. Images were captured every 15min for 48h. siMyc/siScr-GFP-RasV12 cells were
plated 24h after transfection and inserts were removed 8h post transfection. For Wnt5a
experiments cells were treated with PBS/Wnt5a for 8h before inserts were removed.
PBS/Wnt5a treatment was maintained during the whole experiment. WNT-974 porcupine
inhibitor (Stratech)/DMSO (Sigma-Aldrich) was added to the medium when inserts were
removed. Retraction was measured as the distance GFP-RasV12 cells migrated during 24h,
following initial cell-cell collision with MDCK cells. Coefficient of boundary smoothness was
measured using Fiji in cell confrontation assays at the end of the 24 h experiment and as
previously described (*9*). We performed three technical replicates per experiment in three
experiments.

Western blotting was performed using GFP-RasV12 cell lysates transfected with siMyc/siScr

siRNAs at 48h post transfection (SFig. 3E), treated for 48h with WNT-974/DMSO 24h post-

transfection (SFig. 5D). Cells were lysed using laemmli buffer (0.0625 M Tris base, 2% SDS,

10% glycerol). Proteins were separated on 12% polyacrylamide gels under reducing PAGE

conditions. Proteins were then transferred to PVDF membranes (Immobilon-P, Merck) using

wet transfer (1h 100V). Membranes were blocked in 5% milk TBS-T for 1h and antibodies

were added in the same buffer at the concentrations listed in table 1 for overnight at 4ºC or

2h at room temperature. Secondary antibodies (table 1) were added in 5% milk TBS-T and

membranes were developed using ECL luminol kit (Merck) and chemiluminescence films

(Amersham Hyperfilm ECL, GE Healthcare). Protein levels were quantified using Fiji

(RRID:SCR_002285) software 'Gels' tool.

Real time cell confluency was measured using Incucyte S3 live cell imaging instrument and

software (v2020). 1×10^3 cells siMyc/siScr-GFP-RasV12 cells were plated 24h after

transfection with siMyc/siScr siRNAs in MW96 (Corning). Images were captured every 15min

for 48h. We performed three technical replicates per experiment in three experiments.

<u>Statistical tests</u>

Statistical analyses were performed using GraphPad Prism 10 software. Normally distributed

data, as determined by the Shapiro-Wilke test or D'Agostino and Pearson test were

analysed using unpaired Student's t tests. A *p* value of <0.05 was taken as significant and a

rejection of the null hypothesis. No statistical method was used to pre-determine sample

size. Definition of *n* is in the figure legends. Gene set enrichment analysis (GSEA) was

performed using GSEA 4.2.2 software. GSEA was used to identify gene sets that were

differentially expressed between two groups of samples. Gene sets were considered

enriched if they had a false discovery rate (FDR) of less than 0.25. Heatmaps were created

by computing normalised gene counts for each individual sample into row z-scores.

Table 1

Antibody and fluorescence stains	Application	Antigen Retrieval	Dilution	Source	Identifier
Ki-67	IHC, IF	ProK 15'	1:500	Abcam	ab16667
CC-3	IHC	Citrate pH6 15'	1:300	Cell Signaling	9661s
RFP	IHC, IF	ProK 15'	1:500	Rockland	600-401-379

Lectin PNA-A488	FACS			Invitrogen	L21409
c-Myc	WB		1:500	Santa Cruz	sc788
Phospho-p38	WB		1:1000	Cell Signaling	9211s
p38a	WB		1:1000	Santa Cruz	sc136210
Vinculin	WB		1:1000	Sigma Aldrich	V4505
Phosphor-ERK	WB		1:1000		
ERK	WB		1:1000	Cell Signaling	
p21WAP/Cip	WB		1:1000	Sigma Aldrich	P1484
CD44	IF	ProK 15'	1:25	Cell Signaling	3570S
K-cadherin	IF	PoK 15'	1:50	ThermoFisher	MA1-06305
β-catenin	IF	Citrate pH6 15'	1:50	BD Biosciences	610154
RFP	IF	Citrate pH6 15'	1:500	Creative Diagnosis	DPATB-H83194
E-cadherin	IF	Citrate pH6 15'	1:1000	BD Biosciences	610182
p27	IF		1:25	Santa Cruz	SC-528
Wnt5a	IF		1:50	Santa Cruz	SC-30224
Dvl2	IF			Santa Cruz	SC-13974
goat anti-rabbit AF568	IF		1:1000	ThermoFisher	A-11011
donkey anti-mouse AF488			1:1000	ThermoFisher	A-21202
goat anti-rabbit AF488	IF		1:1000	ThermoFisher	A-11008
goat anti-rabbit HRP	WB		1:5000	Vector Laboratories	MP-7451-15
goat anti-mouse HRP	WB		1:5000	Vector Laboratories	MP-7802-15
Phalloidin	IF		1:1000	Sigma Aldrich	65906
E-cadherin	F		1:100	ThermoFisher	13-1900
Caveolin	IF		1:50	Abcam	Ad2910
Hoechst			1:1000	ThermoFisher	H3570

References

1. C. Bielmeier *et al.*, Interface Contractility between Differently Fated Cells Drives Cell
Elimination and Cyst Formation. *Curr Biol* **26**, 563-574 (2016).

2. O. Klipa, F. Hamaratoglu, Cell elimination strategies upon identity switch via
modulation of apterous in Drosophila wing disc. *PLoS Genet* **15**, e1008573 (2019).

3. S. Brown *et al.*, Correction of aberrant growth preserves tissue homeostasis. *Nature*
548, 334-337 (2017).

4. W. Hill *et al.*, EPHA2-dependent outcompetition of KRASG12D mutant cells by wild-
type neighbors in the adult pancreas. *Curr Biol* **31**, 2550-2560 e2555 (2021).

5. C. Hogan *et al.*, Characterization of the interface between normal and transformed
epithelial cells. *Nat Cell Biol* **11**, 460-467 (2009).

6. M. Kajita *et al.*, Interaction with surrounding normal epithelial cells influences
signalling pathways and behaviour of Src-transformed cells. *J Cell Sci* **123**, 171-180
(2010).

7. S. Kon *et al.*, Cell competition with normal epithelial cells promotes apical extrusion
of transformed cells through metabolic changes. *Nat Cell Biol* **19**, 530-541 (2017).

8. M. Norman *et al.*, Loss of Scribble causes cell competition in mammalian cells. *J Cell*
Sci **125**, 59-66 (2012).

9. S. Porazinski *et al.*, EphA2 Drives the Segregation of Ras-Transformed Epithelial
Cells from Normal Neighbors. *Curr Biol* **26**, 3220-3229 (2016).

10. S. M. van Neerven, L. Vermeulen, Cell competition in development, homeostasis and
cancer. *Nat Rev Mol Cell Biol* **24**, 221-236 (2023).

11. N. E. Baker, Emerging mechanisms of cell competition. *Nat Rev Genet* **21**, 683-697
(2020).

12. D. Fernandez-Antoran *et al.*, Outcompeting p53-Mutant Cells in the Normal
Esophagus by Redox Manipulation. *Cell Stem Cell* **25**, 329-341 e326 (2019).

13. K. Murai *et al.*, Epidermal Tissue Adapts to Restrain Progenitors Carrying Clonal p53
Mutations. *Cell Stem Cell* **23**, 687-699 e688 (2018).

14. A. Sasaki *et al.*, Obesity Suppresses Cell-Competition-Mediated Apical Elimination of
RasV12-Transformed Cells from Epithelial Tissues. *Cell Rep* **23**, 974-982 (2018).

15. M. P. Alcolea *et al.*, Differentiation imbalance in single oesophageal progenitor cells
causes clonal immortalization and field change. *Nat Cell Biol* **16**, 615-622 (2014).

16. D. J. Flanagan *et al.*, NOTUM from Apc-mutant cells biases clonal competition to
initiate cancer. *Nature* **594**, 430-435 (2021).

17. S. M. van Neerven *et al.*, Apc-mutant cells act as supercompetitors in intestinal
tumour initiation. *Nature* **594**, 436-441 (2021).

18. M. K. Yum *et al.*, Tracing oncogene-driven remodelling of the intestinal stem cell
niche. *Nature* **594**, 442-447 (2021).

19.   W. Hill, C. Hogan, Normal epithelial cells trigger EphA2-dependent RasV12 cell
repulsion at the single cell level. *Small GTPases* **10**, 305-310 (2019).

20.   S. Lanfredini, A. Thapa, E. O'Neill, RAS in pancreatic cancer. *Biochem Soc Trans* **47**,
961-972 (2019).

21.   M. A. Collins *et al.*, Oncogenic Kras is required for both the initiation and
maintenance of pancreatic cancer in mice. *J Clin Invest* **122**, 639-653 (2012).

22.   P. Bailey *et al.*, Genomic analyses identify molecular subtypes of pancreatic cancer.
Nature **531**, 47-52 (2016).

23.   J. P. Morton *et al.*, Mutant p53 drives metastasis and overcomes growth
arrest/senescence in pancreatic cancer. *Proc Natl Acad Sci U S A* **107**, 246-251
(2010).

24.   T. E. Woolley, W. Hill, C. Hogan, Accounting for dimensional differences in stochastic
domain invasion with applications to precancerous cell removal. *J Theor Biol* **541**,
111024 (2022).

25.   S. R. Hingorani *et al.*, Preinvasive and invasive ductal pancreatic cancer and its early
detection in the mouse. *Cancer Cell* **4**, 437-450 (2003).

26.   B. M. Flowers *et al.*, Cell of Origin Influences Pancreatic Cancer Subtype. *Cancer*
Discov **11**, 660-677 (2021).

27.   Z. Ma *et al.*, Single-Cell Transcriptomics Reveals a Conserved Metaplasia Program
in Pancreatic Injury. *Gastroenterology* **162**, 604-620 e620 (2022).

28.   K. Nozaki *et al.*, A molecular signature of gastric metaplasia arising in response to
acute parietal cell loss. *Gastroenterology* **134**, 511-522 (2008).

29.   J. Benn, R. J. Schneider, Hepatitis B virus HBx protein deregulates cell cycle
checkpoint controls. *Proc Natl Acad Sci U S A* **92**, 11215-11219 (1995).

30.   T. Kasai, Y. Iwanaga, H. Iha, K. T. Jeang, Prevalent loss of mitotic spindle checkpoint
in adult T-cell leukemia confers resistance to microtubule inhibitors. *J Biol Chem* **277**,
5187-5193 (2002).

31.   H. Toyoshima, T. Hunter, p27, a novel inhibitor of G1 cyclin-Cdk protein kinase
activity, is related to p21. *Cell* **78**, 67-74 (1994).

32.   V. A. van der Weijden, A. Bulut-Karslioglu, Molecular Regulation of Paused
Pluripotency in Early Mammalian Embryos and Stem Cells. *Front Cell Dev Biol* **9**,
708318 (2021).

33.   J. E. Davis, Jr., J. Kirk, Y. Ji, D. G. Tang, Tumor Dormancy and Slow-Cycling Cancer
Cells. *Adv Exp Med Biol* **1164**, 199-206 (2019).

34.   E. Dhimolea *et al.*, An Embryonic Diapause-like Adaptation with Suppressed Myc
Activity Enables Tumor Treatment Persistence. *Cancer Cell* **39**, 240-256 e211
(2021).

35.   A. M. Hussein, N. Balachandar, J. Mathieu, H. Ruohola-Baker, Molecular Regulators
of Embryonic Diapause and Cancer Diapause-like State. *Cells* **11**, (2022).

36.   R. S. Kim *et al.*, Dormancy signatures and metastasis in estrogen receptor positive
and negative breast cancer. *PLoS One* **7**, e35569 (2012).

37.   D. B. Fox *et al.*, NRF2 activation promotes the recurrence of dormant tumour cells
through regulation of redox and nucleotide metabolism. *Nat Metab* **2**, 318-334
(2020).

38.   S. Easwaran, M. Van Ligten, M. Kui, D. J. Montell, Enhanced germline stem cell
longevity in Drosophila diapause. *Nat Commun* **13**, 711 (2022).

39.   K. Itahana *et al.*, A role for p53 in maintaining and establishing the quiescence
growth arrest in human cells. *J Biol Chem* **277**, 18206-18214 (2002).

40.   A. J. Wiecek *et al.*, Genomic hallmarks and therapeutic implications of G0 cell cycle
arrest in cancer. *Genome Biol* **24**, 128 (2023).

41.   K. A. Anton, M. Kajita, R. Narumi, Y. Fujita, M. Tada, Src-transformed cells hijack
mitosis to extrude from the epithelium. *Nat Commun* **9**, 4695 (2018).

42.   H. Watanabe *et al.*, Mutant p53-Expressing Cells Undergo Necroptosis via Cell
Competition with the Neighboring Normal Epithelial Cells. *Cell Rep* **23**, 3721-3729
(2018).

43.   D. Dominguez-Sola, J. Gautier, MYC and the control of DNA replication. *Cold Spring
Harb Perspect Med* **4**, (2014).

44.   R. Scognamiglio *et al.*, Myc Depletion Induces a Pluripotent Dormant State Mimicking
Diapause. *Cell* **164**, 668-680 (2016).

45.   M. E. Fane *et al.*, Stromal changes in the aged lung induce an emergence from
melanoma dormancy. *Nature* **606**, 396-405 (2022).

46.   D. Ren *et al.*, Wnt5a induces and maintains prostate cancer cells dormancy in bone.
J Exp Med **216**, 428-449 (2019).

47.   J. B. Cordero, O. J. Sansom, Wnt signalling and its role in stem cell-driven intestinal
regeneration and hyperplasia. *Acta Physiol (Oxf)* **204**, 137-143 (2012).

48.   D. J. Flanagan *et al.*, Frizzled7 functions as a Wnt receptor in intestinal epithelial
Lgr5(+) stem cells. *Stem Cell Reports* **4**, 759-767 (2015).

49.   A. Abedini, C. Sayed, L. E. Carter, D. Boerboom, B. C. Vanderhyden, Non-canonical
WNT5a regulates Epithelial-to-Mesenchymal Transition in the mouse ovarian surface
epithelium. *Sci Rep* **10**, 9695 (2020).

50.   S. K. Dissanayake *et al.*, The Wnt5A/protein kinase C pathway mediates motility in
melanoma cells via the inhibition of metastasis suppressors and initiation of an
epithelial to mesenchymal transition. *J Biol Chem* **282**, 17259-17271 (2007).

51.   V. J. Wielenga *et al.*, Expression of CD44 in Apc and Tcf mutant mice implies
regulation by the WNT pathway. *Am J Pathol* **154**, 515-523 (1999).

52.   A. Kikuchi, H. Yamamoto, A. Sato, S. Matsumoto, Wnt5a: its signalling, functions and
implication in diseases. *Acta Physiol (Oxf)* **204**, 17-33 (2012).

53. B. T. MacDonald, K. Tamai, X. He, Wnt/beta-catenin signaling: components,
mechanisms, and diseases. *Dev Cell* **17**, 9-26 (2009).

54. A. G. Grieve, C. Rabouille, Extracellular cleavage of E-cadherin promotes epithelial
cell extrusion. *J Cell Sci* **127**, 3331-3346 (2014).

55. S. Saitoh *et al.*, Rab5-regulated endocytosis plays a crucial role in apical extrusion of
transformed cells. *Proc Natl Acad Sci U S A* **114**, E2327-E2336 (2017).

56. R. G. Parton, K. Simons, The multiple faces of caveolae. *Nat Rev Mol Cell Biol* **8**,
185-194 (2007).

57. A. Ohoka *et al.*, EPLIN is a crucial regulator for extrusion of RasV12-transformed
cells. *J Cell Sci* **128**, 781-789 (2015).

58. J. L. Teo *et al.*, Caveolae Control Contractile Tension for Epithelia to Eliminate Tumor
Cells. *Dev Cell* **54**, 75-91 e77 (2020).

59. D. J. Flanagan, S. A. Woodcock, C. Phillips, C. Eagle, O. J. Sansom, Targeting
ligand-dependent wnt pathway dysregulation in gastrointestinal cancers through
porcupine inhibition. *Pharmacol Ther* **238**, 108179 (2022).

60. H. K. Lin *et al.*, Caveolin-1 down-regulation is required for Wnt5a-Frizzled 2 signalling
in Ha-Ras(V12) -induced cell transformation. *J Cell Mol Med* **22**, 2631-2643 (2018).

61. D. J. Flanagan *et al.*, Frizzled-7 Is Required for Wnt Signaling in Gastric Tumors with
and Without Apc Mutations. *Cancer Res* **79**, 970-981 (2019).

62. S. Kon, Y. Fujita, Cell competition-induced apical elimination of transformed cells,
EDAC, orchestrates the cellular homeostasis. *Dev Biol* **476**, 112-116 (2021).

63. K. Puzik *et al.*, Lef1 regulates caveolin expression and caveolin dependent
endocytosis, a process necessary for Wnt5a/Ror2 signaling during Xenopus
gastrulation. *Sci Rep* **9**, 15645 (2019).

64. C. Medrek, G. Landberg, T. Andersson, K. Leandersson, Wnt-5a-CKIalpha signaling
promotes beta-catenin/E-cadherin complex formation and intercellular adhesion in
human breast epithelial cells. *J Biol Chem* **284**, 10968-10979 (2009).

65. E. Risson, A. R. Nobre, V. Maguer-Satta, J. A. Aguirre-Ghiso, The current paradigm
and challenges ahead for the dormancy of disseminated tumor cells. *Nat Cancer* **1**,
672-680 (2020).

66. M. S. Sosa, P. Bragado, J. Debnath, J. A. Aguirre-Ghiso, Regulation of tumor cell
dormancy by tissue microenvironments and autophagy. *Adv Exp Med Biol* **734**, 73-89
(2013).

67. L. Baillon, F. Germani, C. Rockel, J. Hilchenbach, K. Basler, Xrp1 is a transcription
factor required for cell competition-driven elimination of loser cells. *Sci Rep* **8**, 17712
(2018).

68. P. F. Langton, M. E. Baumgartner, R. Logeay, E. Piddini, Xrp1 and Irbp18 trigger a
feed-forward loop of proteotoxic stress to induce the loser status. *PLoS Genet* **17**,
e1009946 (2021).

69.   B. Brown, S. Mitra, F. D. Roach, D. Vasudevan, H. D. Ryoo, The transcription factor
Xrp1 is required for PERK-mediated antioxidant gene induction in Drosophila. *Elife*
10, (2021).

70.   N. Ochi *et al.*, Cell competition is driven by Xrp1-mediated phosphorylation of
eukaryotic initiation factor 2alpha. *PLoS Genet* **17**, e1009958 (2021).

71.   N. E. Baker, C. Montagna, Reducing the aneuploid cell burden - cell competition and
the ribosome connection. *Dis Model Mech* **15**, (2022).

72.   M. Kiparaki *et al.*, The transcription factor Xrp1 orchestrates both reduced translation
and cell competition upon defective ribosome assembly or function. *Elife* **11**, (2022).

73.   J. Zheng *et al.*, Differential Ire1 determines loser cell fate in tumor-suppressive cell
competition. *Cell Rep* **42**, 113303 (2023).

74.   S. Bowling *et al.*, Author Correction: P53 and mTOR signalling determine fitness
selection through cell competition during early mouse embryonic development. *Nat
Commun* **9**, 3123 (2018).

75.   A. M. Klein, D. E. Brash, P. H. Jones, B. D. Simons, Stochastic fate of p53-mutant
epidermal progenitor cells is tilted toward proliferation by UV B during preneoplasia.
Proc Natl Acad Sci U S A **107**, 270-275 (2010).

76.   L. Wagstaff *et al.*, Mechanical cell competition kills cells via induction of lethal p53
levels. *Nat Commun* **7**, 11373 (2016).

77.   G. Zhang *et al.*, p53 pathway is involved in cell competition during mouse
embryogenesis. *Proc Natl Acad Sci U S A* **114**, 498-503 (2017).

78.   I. Kucinski, M. Dinan, G. Kolahgar, E. Piddini, Chronic activation of JNK JAK/STAT
and oxidative stress signalling causes the loser cell status. *Nat Commun* **8**, 136
(2017).

79.   W. Hirose *et al.*, Selective Elimination of NRF2-Activated Cells by Competition With
Neighboring Cells in the Esophageal Epithelium. *Cell Mol Gastroenterol Hepatol* **15**,
153-178 (2023).

80.   J. Burclaff, J. C. Mills, Plasticity of differentiated cells in wound repair and
tumorigenesis, part I: stomach and pancreas. *Dis Model Mech* **11**, (2018).

81.   K. Kozyrska *et al.*, p53 directs leader cell behavior, migration, and clearance during
epithelial repair. *Science* **375**, eabl8876 (2022).

82.   T. Bondar, R. Medzhitov, p53-mediated hematopoietic stem and progenitor cell
competition. *Cell Stem Cell* **6**, 309-322 (2010).

83.   A. Marusyk, C. C. Porter, V. Zaberezhnyy, J. DeGregori, Irradiation selects for p53-
deficient hematopoietic progenitors. *PLoS Biol* **8**, e1000324 (2010).

84.   K. Murai *et al.*, p53 mutation in normal esophagus promotes multiple stages of
carcinogenesis but is constrained by clonal competition. *Nat Commun* **13**, 6206
(2022).

85.   Y. Haraoka, Y. Akieda, Y. Nagai, C. Mogi, T. Ishitani, Zebrafish imaging reveals
988   TP53 mutation switching oncogene-induced senescence from suppressor to driver in
989   primary tumorigenesis. *Nat Commun* **13**, 1417 (2022).

990   86.   C. Burdziak *et al.*, Epigenetic plasticity cooperates with cell-cell interactions to direct
991   pancreatic tumorigenesis. *Science* **380**, eadd5327 (2023).

992   87.   G. Gu, J. Dubauskaite, D. A. Melton, Direct evidence for the pancreatic lineage:
993   NGN3+ cells are islet progenitors and are distinct from duct progenitors.
994   *Development* **129**, 2447-2457 (2002).

995   88.   E. L. Jackson *et al.*, Analysis of lung tumor initiation and progression using
996   conditional expression of oncogenic K-ras. *Genes Dev* **15**, 3243-3248 (2001).

997   89.   K. P. Olive *et al.*, Mutant p53 gain of function in two mouse models of Li-Fraumeni
998   syndrome. *Cell* **119**, 847-860 (2004).

90.   H. Luche, O. Weber, T. Nageswara Rao, C. Blum, H. J. Fehling, Faithful activation of
an extra-bright red fluorescent protein in "knock-in" Cre-reporter mice ideally suited
for lineage tracing studies. *Eur J Immunol* **37**, 43-53 (2007).

91.   J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-682 (2012).

92.   M. Assi, N. Dauguet, P. Jacquemin, DIE-RNA: A Reproducible Strategy for the
Digestion of Normal and Injured Pancreas, Isolation of Pancreatic Cells from
Genetically Engineered Mouse Models and Extraction of High Quality RNA. *Front*
Physiol **9**, 129 (2018).
1008

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

KC PC KPC

Supplementary figure 1

Supplementary figure 2

Supplementary figure 3

Supplementary figure 4

Supplementary figure 5

Supplementary figure 6

Figure 1: Cells expressing p53R172H are eliminated from adult pancreas *in vivo*. Co-expression of p53R172H mutations prevents KrasG12D cell elimination *in vivo*.

A Schematic of mouse models (left), tamoxifen treatment and timepoints (right). 6–8-week-old *Pdx1-Cre^ERT; Rosa26^LSL-tdRFP* (Control), *Pdx1-Cre^ERT; Rosa26^LSL-tdRFP; LSL-Kras^G12D/+* (KC), *Pdx1-Cre^ERT; Rosa26^LSL-tdRFP; LSL-TP53^R172H/+* (PC), and *Pdx1-Cre^ERT; Rosa26^LSL-tdRFP; LSL-Kras^G12D/+; LSL-TP53^R172H/+* (KPC) mice were treated with one single low dose of tamoxifen. Pancreata were harvested at 7-, 35- or 70-days post tamoxifen induction. Illustration created with BioRender.com. **B** Representative images of endogenous RFP fluorescence showing the area occupied by RFP in Control, KrasG12D (KC), p53R172H (PC) or double mutant (KPC) pancreas tissue sections harvested at 7- days (top) and 35-days p.i. Scale bar, 500µm. **C** Percentage of RFP fluorescence per total tissue area from Control, KrasG12D (KC), p53R172H (PC) or double mutant (KPC) tissues at 7- (squares), 35- (circles) and 70-days (triangles) p.i. Each data point represents average RFP fluorescence per mouse; control n=6 (7-days p.i.), n=5 (35-days p.i.), n=4 (70-days p.i.); KC n=6 (7- and 70-days p.i.), n=4 (35-days p.i.); PC n=6 (7-days p.i.), n=5 (35- and 70-days p.i.); KPC n=6 (7-days p.i.), n=4 (35- and 70-days p.i.). The data represent mean and +/- s.d. Student's t test was used to analyse the data, ***$p<0.0001$, *$p=0.0173$, ns>0.1504. **D** Percentage of RFP+ ducts at 7-days p.i. Ducts that carried at least one RFP+ cell were quantified as positive. Data points represent mean RFP+ percentage per mouse (n=3 per group), with at least 20 ducts counted per mouse. The data represent mean and +/- s.d. Student's t test was used to analyse the data, **$p<0.004$, ns=0.2879. **E** Percentage of RFP+ cells per total number of cells in islets at 35 days p.i. SuperPlot shows all the quantified islets (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data, *$p<0.03$, ns=0.5394. **F** Percentage of RFP positive clusters grouped according to cluster area (μm^2). RFP positive cluster area was measured, and clusters were grouped according to their size in 4 groups at 7 and 35 days p.i.: 20-50 μm^2, 51-100 μm^2, 101-1000 μm^2 and more than 1000 μm^2. The graph shows the percentage of clusters in each group relative to the total number of clusters. At 35 days p.i. the percentage of 20-50 μm^2 clusters significantly decreases in KC ($p=0.0004$) and PC ($p=0.0316$), no significant differences were observed in Control or KPC tissues (p=0.7740). Student's t test was used to analyse the data. The data represent mean +/- s.d. for at least 700 clusters were quantified per mouse and 3 mice per genotype/timepoint.

Figure 2: Bulk RNA sequencing results indicate cell fate changes in p53R172H and KrasG12D mutant retained cells.

A Experimental design for tissue collection for RNA sequencing. Control, KC, PC and KPC mice were treated with low dose tamoxifen induction. RFP+ cells were isolated from pancreas

tissues harvested at 35 days p.i. for bulk RNA sequencing. Three samples were analysed per genotype, each sample was obtained by pooling the pancreas from three mice. Illustration created with BioRender.com. **B-C** Normalised enrichment scores (NES) of gene set enrichment analysis (GSEA) on the Hallmark and KEGG MEDICUS gene sets for the RNA-seq analysis of **B** KC retained cells or **C** PC retained cells. Only gene sets with an FDR of less than 0.25 were included in the graph. The complete list that contains the results of GSEA analysis is provided in supplementary tables S4-S7.

Figure 3: Cell dormancy/diapause gene signatures and cell cycle arrest are features of non-eliminated Ras mutant cells.

Heatmaps showing row z-scores representing expression of **A** pancreatic progenitor associated, **C** cell cycle associated and **F** cell dormancy (rows) in non-eliminated KrasG12D (KC), non-eliminated p53R172H (PC) or Control cells (columns). Only genes with z-scores of a foldchange of 0.5 for KC average values are shown in Cell cycle panel. The full list of genes can be found in supplementary table S10. **B** Percentage of Ki-67 positive cells/total cells in Control, KrasG12D (KC), p53R172H (PC) or double mutant (KPC) pancreas tissues harvested at 7 and 35 days p.i. The data represent mean and +/- s.d. per mouse; control n=6 (7-days p.i.), n=5 (35-days p.i.); KC n=6 (7- and 35-days p.i.); PC n=3 (7- and 35-days p.i.); KPC n=3 (7-days p.i.), n=4 (35- days p.i.) mice. **D** Percentage of cleaved caspase-3 positive cells/total cells in Control, KrasG12D (KC), p53R172H (PC) or double mutant (KPC) pancreas tissues harvested at 7 and 35 days p.i. mean +/- s.d. per mouse; control n=5 (7-days p.i.), n=4 (35-days p.i.); KC n=5 (7- and 35-days p.i.); PC n=3 (7-days p.i.), n=6 (35-days p.i.); KPC n=3 (7-days p.i.), n=4 (35- days p.i.) mice. **E** Percentage of p27 positive nuclei per total number of nuclei in each RFP+ cluster in Control or KrasG12D (KC) pancreas tissues harvested at 35 days p.i. SuperPlot shows all the quantified RFP+ clusters (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data, *p=0.0237. **G** Confocal images of GFP-RasV12 cells-expressing scrambled siRNA (siScr) or two siRNA oligos targeting endogenous *Myc* (siMyc1+2) and then mixed with non-labelled MDCK cells (1:50). Cells were stained with phalloidin to visualise F-actin (red) and Hoechst (blue). Merged images are shown. Scale bar, 100µm. **H** The proportion of non-extruded GFP-RasV12 cells transiently expressing different siRNA *Myc* oligos (siMyc1, siMyc2, siMyc1+2) relative to non-extruded GFP-RasV12 cells expressing siRNA scrambled (siScr). Data represent mean +/- s.d. from three independent experiments. At least 300 cells per point (100 cells per experiment). Student's t test was used to analyse the data, ***p<0.0005, *p<0.05. **I** Confocal images of GFP-RasV12-expressing siRNA oligos targeting endogenous Myc (siMyc1+2), mixed with non-labelled MDCK cells (1:50). Cells were stained with anti-Ki-67 antibodies (red)

and Hoechst (blue). Scale bar, 50 μm. **J** Percentage of Ki-67 positive cells extruded GFP-RasV12 cells as a fraction of total extruded cells. Non-extruded cells were not Ki-67 positive. Data represent mean +/- s.d. for 60 cells from three independent experiments. Student's t test was used to analyse the data, **p<0.005.

Figure 4: Wnt pathway is activated at normal-mutant cell-cell boundaries *in vivo*.
Heatmaps of **A** Wnt signalling pathway, **B** Wnt/β-catenin pathway or **C** Wnt5a-Ror pathway related genes (rows) in Control and KrasG12D (KC) cell transcriptomes (columns). The heatmaps show row z-scores for each gene obtained from the RNA sequencing experiment. Data from three samples (three mice pooled per sample) per genotype are shown. Only genes with z-scores of a foldchange of 2 are shown in the Wnt pathway panel. The full list of genes can be found in supplementary table S11. **D** *Wnt5a* mRNA reads relative to control reads in the three samples in Control and three samples in KrasG12D (KC) obtained from the RNA sequencing experiment. Graph represents mean +/- s.d. reads. **E** Mean fluorescence intensity of WNT5A in Control or KrasG12D (KC) pancreas tissues harvested at 35 days p.i. SuperPlot shows all the quantified tissue areas (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data, *p=0.0491. **F** Mean fluorescence intensity of DVL2 in Control or KrasG12D (KC) pancreas tissues harvested at 35 days p.i. Mean fluorescence intensity is relative to the background. SuperPlot shows all the quantified RFP+ clusters (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data, *p=0.0110. **G** Confocal images of pancreas tissues harvested at 35-days p.i. from Control, KrasG12D (KC) mice. Tissues were stained with anti-RFP (red) and anti-β-catenin (cyan) antibodies and Hoechst (blue). Dashed boxes represent digital zoom images of representative cell-cell contacts. Scale bar, 50μm. **H** Mean fluorescence intensity of β-catenin in the nuclei of RFP positive cells in Control and KrasG12D (KC) tissues harvested at 35 days p.i. Mean fluorescence intensity is relative to the background. SuperPlot shows all the quantified nuclei (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data. ns p=0.3035. **I** Mean fluorescence intensity of β-catenin at the boundary between RFP positive and negative cells in Control and KrasG12D (KC) tissues harvested at 35 days p.i. Mean fluorescence intensity is relative to the background. Fluorescence was measured in ROIs, separated by a constant distance, along the cell-cell boundary. SuperPlot shows all the quantified ROIs (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse

the data. *p=0.0347. **J** Confocal images of pancreas tissues harvested at 35-days p.i. from Control and KrasG12D (KC) mice and stained with anti-RFP (red) and anti-E-cadherin (cyan) antibodies and Hoechst (blue). Scale bar, 20μm. **K** Mean fluorescence intensity of E-cadherin at the boundary between RFP positive and RFP negative cells in Control and KrasG12D (KC) tissues harvested at 35 days p.i. E-cadherin fluorescence was quantified and reported as in I. SuperPlot shows all the quantified ROIs (smaller circles) and the mean for each mouse (n=5 samples; larger circles – lighter blue, blue, darker blue, green, and yellow). The graph shows mean and +/- s.d. for the five mice. Student's t test was used to analyse the data. *p=0.0348.

Figure 5: Wnt5a induces stable E-cadherin-based cell-cell adhesion in RasV12 cells.
A Confocal images of extruded and non-extruded cells treated with PBS, Wnt3a or Wnt5a. Fixed cells were stained with Hoechst (grey). Arrowheads indicate extruding (white) or non-extruding (yellow) GFP-RasV12 cells. Scale bar, 100 μm. **B** The proportion of non-extruded GFP-RasV12 cells treated with Wnt3a (pink) or Wnt5a (yellow) recombinant proteins relative to non-extruded GFP-RasV12 cells treated with PBS (black). The data represent mean +/- s.d. for at least 300 cells from three independent repeats. Student's t test was used to analyse the data. **p=0.0065; ns p=0.4488. **C** Confocal images of GFP-RasV12 cells mixed with MDCK cells treated with PBS, Wnt3a or Wnt5a for 30h. Cells were stained with anti-β-catenin (cyan) and anti-E-cadherin (magenta) antibodies and Hoechst (blue). Scale bar, 20μm. **D, E** Mean fluorescence intensity of E-cadherin (**D**) or β-catenin (**E**) at the boundary between GFP-RasV12 and MDCK cells treated with PBS (black), Wnt3a (pink) or Wnt5a (yellow) for 30h. Fluorescence was measured in ROIs, separated by a constant distance, along the cell-cell boundary. SuperPlot shows all the quantified ROIs (smaller circles) and the mean for each experiment (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three experiments. Student's t test was used to analyse the data. **p=0.0026; *p=0.0116; ns p>0.2. **F** Confocal images of GFP-RasV12 cells mixed with MDCK cells treated with PBS or Wnt5a for 16h. Cells were stained with anti-E-cadherin (magenta) antibody and Hoechst (blue). Scale bar, 20μm. **G** Mean fluorescence intensity of intracellular E-cadherin in GFP-RasV12 (mixed with MDCK cells) treated with PBS (black) or Wnt5a (yellow) for 16h. SuperPlot shows all the quantified GFP-RasV12 cells (smaller circles) and the mean for each experiment (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three experiments. Student's t test was used to analyse the data. *p=0.0051.

Figure 6: Wnt signalling prevents apical extrusion of RasV12 cells from normal epithelial monolayer *in vitro*. Inhibition of Wnt promotes elimination of non-eliminated cells *in vivo*.

A Confocal images of extruded and non-extruded siScr-GFP-RasV12 and siMyc 1+2-GFP-RasV12 cells for 48h with DMSO or WNT-974. Fixed cells were stained with Hoechst (blue). Scale bar, 100 μm. **B** The proportion of non-extruded GFP-RasV12 cells transiently expressing siRNA *Myc* oligos (siMyc1+2) relative to non-extruded GFP-RasV12 cells expressing siRNA scrambled (siScr). Cells were treated with DMSO (white bars) or WNT-974 porcupine inhibitor (dashed bars). The data represent mean +/- s.d. for at least 300 cells from three independent repeats. Student's t test was used to analyse the data. ***$p<0.0005$. **C** Confocal images of extruded and non-extruded GFP-RasV12 cells pre-treated with PBS/Wnt5a and treated for 48h with DMSO/WNT-974/OMP-18R5. OMP-18R5 is a Frizzled receptor antagonist. Fixed cells were stained with Hoechst (blue). Scale bar, 100 μm. **D, E.** GFP-RasV12 non-extruded cells pre-treated with PBS or Wnt5a recombinant protein 24h before mixing with MDCK cells (1:50). Six hours after mixing and plating, cells were treated with **D** DMSO or WNT-974 porcupine inhibitor or **E** OMP-18R5. The data represent mean +/- s.d. for at least 300 cells from three independent repeats. Student's t test was used to analyse the data, ***$p<0.0005$, **$p<0.005$, *$p<0.05$. **F** Confocal images of GFP-RasV12 cells mixed with MDCK cells treated with PBS+DMSO, Wnt5a+DMSO or Wnt5a+WNT-974 for 16h. Cells were stained with anti-E-cadherin (magenta) antibody. Scale bar, 20μm. **G** Mean fluorescence intensity of intracellular E-cadherin in GFP-RasV12 (mixed with MDCK cells) treated with PBS (black) or Wnt5a (yellow) in combination with DMSO (circle) or WNT-974 (square) for 16h. SuperPlot shows all the quantified GFP-RasV12 cells (smaller circles) and the mean for each experiment (n=3 samples; larger circles or squares - blue, green, and yellow). The graph shows mean and +/- s.d. for the three experiments. Student's t test was used to analyse the data. **$p<0.005$. **H** Experimental design for Wnt inhibition experiments *in vivo*. 6–8-week-old KC (*Kras*$^{G12D/+}$) mice were induced with low-dose tamoxifen. Thirty-five days later, mice were treated with the porcupine inhibitor WNT-974 or vehicle for 28 days. Illustration created with BioRender.com. **I** Representative images of endogenous RFP fluorescence in KC pancreas tissue sections harvested 28-days post treatment with vehicle or WNT-974. Scale bar, 500μm. **J** RFP fluorescence per tissue area in KC mice treated with WNT-974 or vehicle and relative to vehicle treated tissues. Data represent mean per mouse (n=4-6 samples) and +/- s.d. Student's t test was used to analyse the data, ***$p=0.0003$.

Supplementary figure 1. Non-eliminated mutant cells can progress to mucin positive lesions.

A Pancreas tissues harvested from KrasG12D (KC), p53R172H (PC) or double mutant (KPC) expressing mice at 168 days p.i. and stained for mucin (using Alcian blue). Scale bar, 500 µm. Boxed area shown magnified (lower panel).

Supplementary figure 2. Transcriptomic analysis indicates Kras and p53 pathways deregulation in retained cells. Non-eliminated KrasG12D (KC) are stem-like, whereas p53R172H-expressing populations are not.

GSEA enrichment plots showing **A** KEGG_MAPK_signaling_pathway (M10792) and **B** KEGG_p53_signaling_pathway (M6370) positive correlation in KrasG12D (KC) cells compared to Control (WT) and **C** KEGG_p53_signaling_pathway (M6370) and **D** KEGG_MAPK_signaling_pathway (M10792) negative correlation in p53R172H (PC) cells compared to Control (WT). **E** Normalised enrichment scores (NES) of gene set enrichment analysis (GSEA) on the Hallmark and KEGG Medicus gene sets for the RNA sequencing experiment analysis of KPC retained cells. Only gene sets with an FDR of less than 0.25 were included in the graph. The complete list that contains the results of GSEA analysis is provided in Supplementary Tables S8-S9. GSEA enrichment plots of WP_pancreatic_adenocarcinoma_pathway (M39732) positively correlated in **F** KrasG12D (KC) cells, and negatively correlated in **G** p53R172H (PC) cells and **H** KrasG12D p53R172H (KPC) cells compared to Control (WT). **I-J** Heatmaps showing gene expression in non-eliminated KrasG12D (KC) cells and control of **I** spasmolytic polypeptide-expressing metaplasia and **J** pancreatic lineage. Heatmaps show row z-scores for the expression of each gene for three samples (three pooled mice per sample) per genotype obtained from the RNA sequencing experiment. Genes are listed in rows, genotypes in columns. GSEA enrichment plots indicate positive correlation in KrasG12D (KC) retained cells compared to Control (WT) for **K** Enrichment of Ramalho_stemness_up (M9473) and **L** Malta_curated_stemness_markers (M30411) (embryonic, neural, and hematopoietic stem cells). GSEA enrichment plots indicate negative correlation for Ramalho_Stemness_up (M30411) in **M** p53R172H (PC) retained cells and **N** KrasG12D p53R172H (KPC) retained cells compared to Control (WT).

Supplementary figure 3. Non-eliminated KrasG12D cells upregulate cell dormancy-associated pathways. Cell cycle arrest abrogates RasV12 cell extrusion *in vitro*.

A Representative images of β-galactosidase staining of pancreas tissue harvested from a KrasG12D (KC) mouse at 35 days p.i (scale bar, 1000µm) and in PanIN lesions as positive control (scale bar, 250 µm). **B** Diapause gene expression (rows) in Control, KrasG12D (KC)

and p53R172H (PC) cell transcriptomes (columns). The heatmap shows row z-scores for each gene obtained from three samples per genotype (three pooled mice per samples) from the RNA sequencing experiment. **C** GSEA enrichment plots showing positive correlation in KrasG12D (KC) retained cells compared with Control (WT) for WP_NRF2_pathway (M39454). **D** Heatmap of expression levels of NRF2 target genes (rows) in three KrasG12D (KC) and three Control samples (three pooled mice per sample). The heatmap shows row z-scores for each gene obtained from the RNA sequencing experiment. Only genes with z-scores of a foldchange of 2.25 are shown. The full list of genes can be found in supplementary table S12. **E** Immunodetection of indicated antigens in siScr-GFP-RasV12 (siScr), siMyc1-GFP-RasV12 (siMyc1), siMyc2-GFP-RasV12 (siMyc2) and siMyc1+2-GFP-RasV12 (siMyc1+2) cells protein lysates. GFP-RasV12 cells were first transfected with scrambled siRNA (siScr) or two siRNA oligos targeting endogenous *Myc* (siMyc1, siMyc2 or combined siMyc1+2). Lysates were collected 48h after transfection. Two blots were run, and vinculin was used as loading control for each blot (vinculin band shown at the bottom of each blot). p-p38=phospho-p38; p-ERK=phospho-ERK. **F** Quantification of protein levels in the blot in E. Values are relative to siScr protein levels. **G** Real time cell confluence of GFP-RasV12 cells transfected with different siRNAs as quantified via Incucyte S3 imaging. Cell confluence was determined in cells 12h post siRNA transfection over 48h. Data indicate mean and +/- s.d. for three experiments per condition. **H** Schematic representation of cell confrontation assay experiments. Illustration created with BioRender.com. **I** Representative time-lapse images of cell confrontation assays. GFP-RasV12 cells expressing either scramble siRNA (siScr-GFP-RasV12) or combined Myc1+2 siRNA (siMyc-GFP-RasV12) confront non-labelled parental MDCK cells. Dashed lines highlight the border between GFP-RasV12 cells and MDCK cells. Solid lines indicate MDCK cells' migration front at the beginning of the experiment. Scale bar, 100µm. **J** Relative retraction distance of GFP-RasV12 cells transfected with either scrambled siRNA (siScr-GFP-RasV12; black bar) or combined Myc1+2 siRNA (siMyc-GFP-RasV12; green bar) following collision with parental MDCK cells until 24h later. Results are presented as retraction distance in siMyc-GFP-RasV12 relative to siScr-GFP-RasV12. Data represent mean and +/- s.d. measurements of three experiments. Student's t test was used to analyse the data, **p<0.005.

Supplementary figure 4. Wnt pathway is active in non-eliminated KrasG12D or p53R172H cells.

A Heatmap of Wnt signalling pathway related genes (rows) in Control and p53R712H (PC) cell transcriptomes (columns). The heatmap shows row z-scores for each gene obtained from the RNA sequencing experiment. Data from three samples (three mice pooled per sample) per genotype are shown. Only genes with z-scores of a foldchange of 2 are shown in the Wnt pathway panel. The full list of genes can be found in supplementary table S11. GSEA

enrichment analysis plots showing positive correlation in **B** KrasG12D (KC) and **C** p53R172H (PC) retained cells compared with Control (WT) for the KEGG_MEDICUS_reference_WNT5A_ROR_signaling_pathway (M47822). **D** Representative images of PanINs in pancreas tissue sections harvested from KrasG12D-expressing (KC) mice at 140 days p.i. Left: Mucin positive PanINs (blue). Scale bar, 100 μm. Right: confocal image of anti-CD44 antibody (cyan) labelling and Hoechst (blue) in PanINs in a consecutive tissue slice. Scale bar, 50 μm. **E** Confocal images of pancreas tissues harvested at 35-days p.i. from Control, KrasG12D (KC) and KrasG12D high dose tamoxifen (KC megadose) mice and stained with anti-RFP (red) and anti-CD44 (cyan) antibodies and Hoechst (blue). Scale bar, 50μm. **F** Mean fluorescence intensity of CD44 at the boundary between RFP positive and RFP negative cells in Control, KrasG12D low dose tamoxifen (KC) and KrasG12D high dose tamoxifen (KC Megadose) tissues harvested at 35 days p.i. CD44 fluorescence was quantified and reported as in figure 4I. Fluorescence was measured in ROIs, separated by a constant distance, along the cell-cell boundary. SuperPlot shows all the quantified ROIs (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data. **p=0.0047; ***p=0.0009; ns=0.2136. **G** Mean fluorescence intensity of CD44 in entire RFP positive clusters. Mean fluorescence intensity is relative to the background. Fluorescence was measured in ROIs, separated by a constant distance, along the cell-cell boundary. SuperPlot shows all the quantified RFP+ clusters (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data. *p<0.05; ns=0.3916. **H** Mean fluorescence intensity of CD44 at the boundary between RFP positive and RFP negative cells in Control and p53R172H (PC) tissues harvested at 35 days p.i. CD44 fluorescence was quantified and reported as in figure 4I. Control quantification is the same used in figure 4I. Fluorescence was measured in ROIs, separated by a constant distance, along the cell-cell boundary. SuperPlot shows all the quantified ROIs (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data. *p=0.0470. **I** Mean fluorescence intensity of CD44 in entire RFP positive clusters CD44 fluorescence was quantified and reported as in G. SuperPlot shows all the quantified clusters (circles) and the mean for each mouse (n=3 samples; triangle, square and diamond). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data. *p=0.0397.

Supplementary figure 5. Wnt5a increases RasV12 cell cohesion.

A *Cdh6* mRNA reads relative to control reads in the three samples in Control and three samples in KrasG12D (KC) obtained from the RNA sequencing experiment. Graph represents mean +/- s.d. reads. **B** Confocal images of pancreas tissues harvested at 35-days p.i. from Control and KrasG12D (KC) mice. Tissues were stained with anti-RFP (red) and anti-K-cadherin (cyan) antibodies and Hoechst (blue). Scale bar, 50µm. **C** Mean fluorescence intensity of K-cadherin clusters in Control or KrasG12D (KC) pancreas tissues harvested at 35 days p.i. SuperPlot shows all the quantified RFP+ clusters (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice Student's t test was used to analyse the data, *p=0.0474. **D, E** Mean fluorescence intensity of E-cadherin (**D**) or β-catenin (**E**) in GFP-RasV12 cell clusters (mixed with MDCK cells) treated with PBS (black), Wnt3a (pink) or Wnt5a (yellow) for 30h. SuperPlot shows all the quantified GFP-RasV12 clusters (smaller circles) and the mean for each experiment (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three experiments. Student's t test was used to analyse the data. ***p=0.0003; ns p>0.05. **F** Cell speed (µm per hour) at which RasV12 (circles) or MDCK (squares) cells treated with PBS (black) or Wnt5a (yellow) migrate. The data represent mean +/- s.d. from three independent experiment. Student's t test was used to analyse the data. *p=0.0268; ns p=0.8202. **G** Confocal images of GFP-RasV12 cells mixed with MDCK cells treated with PBS or Wnt5a for 16h. Cells were stained with anti-CAV1 (magenta). Scale bar, 20µm. **H** Mean fluorescence intensity of CAV1 at the boundary between GFP-RasV12 and MDCK cells treated with PBS (black) or Wnt5a (yellow) for 16h. Fluorescence was measured in ROIs, separated by a constant distance, along the cell-cell boundary. SuperPlot shows all the quantified ROIs (smaller circles) and the mean for each experiment (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three experiments. Student's t test was used to analyse the data. *p=0.0117.

Supplementary figure 6. Cell cycle arrest and Wnt signalling prevent GFP-RasV12 segregation from normal MDCK cells *in vitro* and pancreas cell elimination *in vivo*.

A Representative time-lapse images of cell confrontation assays. GFP-RasV12 cells expressing combined Myc1+2 siRNA (siMyc 1+2-GFP-RasV12) confront non-labelled parental MDCK cells. Assays were conducted in the presence of DMSO or WNT-974. Dashed lines highlight the border between GFP-RasV12 cells and MDCK cells. Solid lines indicate migration front of MDCK cells at the beginning of the experiment. Scale bar, 100µm. **B** Relative retraction distance of GFP-RasV12 cells transfected with scrambled siRNA (siScr; black bars) or two siRNA oligos targeting endogenous *Myc* (combined siMyc1+2: siMyc-GFP-RasV12; green bars) following collision with parental MDCK cells until 24h later. Assays were conducted in the presence of DMSO (solid bars) or WNT-974 (hatched bars). Results are

presented as retraction distance in siMyc-GFP-RasV12 relative to siScr-GFP-RasV12. Data represent mean and +/- s.d. of three experiments. Student's t test was used to analyse the data, *p<0.05. **C** Co-efficient of boundary smoothness in cell confrontation assay. Co-efficient was calculated by measuring the length of the cell-cell boundary between GFP-RasV12 and MDCK cells at the end of the experiment (dashed line in A) divided by the length of a straight line from the top to the bottom of the collision (solid line in A). Data represent mean measurements of three experiments. Student's t test was used to analyse the data, **p<0.005. **D** Cell speed (μm per hour) at which RasV12 cells treated with PBS (black) or Wnt5a (yellow) and DMSO or WNT-974 migrate. The data represent mean +/- s.d. from three independent experiment. Student's t test was used to analyse the data. *p=0.0156; **p=0.0058. **E** Mean fluorescence intensity of CD44 at the boundary between RFP positive and RFP negative cells in vehicle or WNT-974 treated KrasG12D (KC) tissues harvested at the end of the treatment (see figure 5F). CD44 fluorescence was quantified and reported as in figure 4I. Control quantification is the same used in supplementary figure 4F. SuperPlot shows all the quantified ROIs (smaller circles) and the mean for each mouse (n=3 samples; larger circles - blue, green, and yellow). The graph shows mean and +/- s.d. for the three mice. Student's t test was used to analyse the data. *p=0.0467. **F** Representative images of endogenous RFP fluorescence in PC pancreas tissue sections harvested 28-days post treatment with vehicle or WNT-974. Scale bar, 500μm. **G** RFP fluorescence per tissue area in PC mice treated with WNT-974 or vehicle and relative to vehicle treated tissues. Data represent mean and +/- s.d. per mouse. Student's t test was used to analyse the data, ***p<0.0005. **H** Schematic summarising the main findings of the study. Illustration created with BioRender.com.