# Spatial top-down proteomics for the functional characterization of human kidney

**Kevin J. Zemaitis**[1, †]
(0000-0002-3524-9776)
**James M. Fulcher**[1, †]
(0000-0001-9033-3623)
**Rashmi Kumar**[1, †]
(0000-0001-8994-5091)
**David J. Degnan**[2]
(0000-0001-5737-7173)
**Logan A. Lewis**[2]
(0000-0003-2301-300X)
**Yen-Chen Liao**[1]
(0000-0002-8124-0457)
**Marija Veličković**[1]
(0000-0003-3664-5719)
**Sarah M. Williams**[1]
(0000-0002-0175-466X)
**Ronald J. Moore**[2]
(0000-0003-2806-2855)
**Lisa M. Bramer**[2]
(0000-0002-8384-1926)
**Dušan Veličković**[1]
(0000-0001-7945-9620)
**Ying Zhu**[3]
(0000-0002-5416-0566)
**Mowei Zhou**[1, §]
(0000-0003-3575-3224)
**Ljiljana Paša-Tolić**[1, *]
(0000-0001-9853-5457)

[1] Environmental and Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA 99354, United States
[2] Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA 99354, United States
[3] Department of Microchemistry, Proteomics, Lipidomics and Next Generation Sequencing, Genentech, 1 DNA Way, San Francisco, CA 94080, United States
[§] Present address: Department of Chemistry, Zhejiang University, Hangzhou, Zhejiang, China, 310058

[†] Signifies an equal contribution
[*] Signifies corresponding author

# Abstract

*Background:*

The Human Proteome Project has credibly detected nearly 93% of the roughly 20,000 proteins which are predicted by the human genome. However, the proteome is enigmatic, where alterations in amino acid sequences from polymorphisms and alternative splicing, errors in translation, and post-translational modifications result in a proteome depth estimated at several million unique proteoforms. Recently mass spectrometry has been demonstrated in several landmark efforts mapping the human proteoform landscape in bulk analyses. Herein, we developed an integrated workflow for characterizing proteoforms from human tissue in a spatially resolved manner by coupling laser capture microdissection, nanoliter-scale sample preparation, and mass spectrometry imaging.

*Results:*

Using healthy human kidney sections as the case study, we focused our analyses on the major functional tissue units including glomeruli, tubules, and medullary rays. After laser capture microdissection, these isolated functional tissue units were processed with microPOTS (microdroplet processing in one-pot for trace samples) for sensitive top-down proteomics measurement. This provided a quantitative database of 616 proteoforms that was further leveraged as a library for mass spectrometry imaging with near-cellular spatial resolution over the entire section. Notably, several mitochondrial proteoforms were found to be differentially abundant between glomeruli and convoluted tubules, and further spatial contextualization was provided by mass spectrometry imaging confirming unique differences identified by microPOTS, and further

expanding the field-of-view for unique distributions such as enhanced abundance of a truncated

form (1-74) of ubiquitin within cortical regions.

*Conclusions:*

We developed an integrated workflow to directly identify proteoforms and reveal their

spatial distributions. Where of the 20 differentially abundant proteoforms identified as

discriminate between tubules and glomeruli by microPOTS, the vast majority of tubular

proteoforms were of mitochondrial origin (8 of 10) where discriminate proteoforms in glomeruli

were primarily hemoglobin subunits (9 of 10). These trends were also identified within ion images

demonstrating spatially resolved characterization of proteoforms that has the potential to reshape

discovery-based proteomics because the proteoforms are the ultimate effector of cellular functions.

Applications of this technology have the potential to unravel etiology and pathophysiology of

disease states, informing on biologically active proteoforms, which remodel the proteomic

landscape in chronic and acute disorders.

## Keywords

78  *Spatial proteomics, intact proteins, top-down proteomics, laser capture microdissection,*

79  *microPOTS, MALDI, glomeruli, tubules, medullary rays*

## Background

Investigation of proteoforms,[1] which includes post-translational modifications (PTMs),

splice-isoforms, and amino acid variants on a canonical protein sequence has gained increased

attention in recent years. Proteoforms holds great potential to become more precise molecular

biomarkers than their respective proteins, transcripts, and genes.[2, 3] Mass spectrometry (MS) is a

powerful tool to directly characterize proteins; however, routine bottom-up proteomics (BUP)

cannot fully capture the molecular identity of proteoforms.[4] Due to proteolytic digestion within

BUP, proteoform level information is commonly lost or is not trivial to reconstruct.[5] Fortunately,

top-down proteomics (TDP) overcomes this by avoiding enzymatic digestion and allowing for the

analysis of intact proteoforms. An exemplary use of TDP was the recent completion of the Blood

Proteoform Atlas, where roughly 30,000 proteoforms were detected across 1690 human genes.

Many of these proteoforms demonstrated high cell-type specificity.[6] Furthermore, PTMs are

known to regulate cellular functions. For example, phosphorylation plays important roles in

apoptosis and cell growth,[7] and conformation and activity of proteins are widely controlled by

glycosylation.[8] The identity and roles of all proteoforms have yet to be fully characterized,

therefore there is a present need to further contextualize which proteoforms effect function.[3] As

such, advanced TDP methods must be applied for confident and comprehensive proteoform

characterization.

Historically, the sensitivity of TDP has been problematic,[9] especially for higher molecular

mass proteins.[10] Confident identification of proteoforms therefore has required large sample inputs

for high resolution accurate mass measurements and efficient gas phase fragmentation. This has

traditionally limited TDP to bulk-scale tissue analyses and large quantities of cultured cells.

However, recent advances in MS instrumentation and bioinformatic tools have significantly

improved TDP sensitivity and coverage.[11] This progress has enabled the field of TDP to move

towards the detection and discrimination of proteoforms directly from tissues in a spatially

resolved manner. Several direct analysis techniques have been deployed at near cellular resolution

for the analysis of proteins from tissue. Some notable advancements use mass spectrometry

imaging (MSI) such as matrix assisted laser desorption ionization (MALDI),[12, 13] nanospray

desorption electrospray ionization (nanoDESI),[14-16] and liquid extraction surface analysis

(LESA).[17, 18] While many of these techniques are routinely used for probing metabolites, lipids,

and peptides within functional tissue units (FTUs),[19] and in single cell analyses,[20, 21] the spatial

analysis of intact proteoforms remains challenging in both proteoform coverage and spatial

resolution.

Notably, translation of these direct analysis techniques into preclinical studies is

particularly challenging,[22] where the lack of routine chromatographic or gas-phase separation with

various matrix effects lower the sensitivity and dynamic range in comparison to traditional liquid

chromatography tandem mass spectrometry (LC-MS/MS). In the case of MALDI-MSI analyses,

low charge states of the generated ions lead to inefficient fragmentation which poses difficulty in

proteoform identification and coverage.[23] To address the challenge in detecting and quantifying

proteoforms, the field has made considerable progress over the past decade.[24] Spatial proteomics

has also been broadly expanded using methods such as microLESA,[25] liquid microjunction

microextraction (LMJ),[26] parafilm-assisted microdissection (PAM),[26] and laser capture

microdissection (LCM)[27] to circumvent some challenges of MSI. Nanodroplet processing in one-

pot for trace samples (nanoPOTS), and its companion microPOTS, have enabled routine single

cell proteomics and BUP processing of dissected tissues.[28-30]

In a previous study, we demonstrated the coupling of nanoPOTS sample preparation with

TDP enabling the identification of over 150 proteoforms along with various PTMs from roughly

70 cultured HeLa cells.[31] This technology was further extended to spatial TDP by integrating

nanoPOTS with LCM to identify an average of 509 proteoforms per 100,000 $\mu m^2$ dissection in

brain tissue.[10] Here we further demonstrate LCM-TDP of representative regions-of-interest, which

benefits from the chromatographic separation and enables deeper coverage of proteoforms. In

tandem with MALDI-MSI we utilize the accurate mass database of proteoforms from LCM-TDP

to map proteoforms with high spatial resolution and throughput (or field-of-view). Herein, we

highlight an advanced workflow for comprehensive spatial proteomics, where LCM-microPOTS-

TDP and MALDI-MSI are integrated for high-confidence identification, near-cellular localization,

and label-free quantitation of proteoforms within human kidney tissue.

## Methodology

*Experimental Design and Motivations*

**Figure 1: Overall workflow for LCM-microPOTS-TDP and MALDI-MSI. (a)** LCM process where regions of interest are dissected from the tissue mounted on polyethylene naphthalate (PEN) membrane slides and are catapulted into microdroplets on microPOTS chips for TDP; **(b)** The isolated samples in microdroplets undergo protein extraction followed by LC-MS/MS to **(c)** generate quantitative proteoform databases. **(d)** Serial sections on ITO slides are prepared for MALDI-MSI with **(e)** UHMR HF Orbitrap detection enabling high mass resolving power and high mass accuracy which **(f)** is needed to identify intact proteoforms based on isotopic distributions using top-down databases generated from LCM-microPOTS-TDP in **(a-c)**. **(g)** Bioinformatic tools are developed to confidently annotate MALDI-MSI spectra and statistics, clustering, and segmentation are then applied.

The generalized workflow for processing fresh-frozen tissues for tandem LCM-

microPOTS-TDP and MALDI-MSI is illustrated within **Figure 1**. A critical consideration for

confident proteoform identification from MALDI-MSI is the use of serial or near serial sections

from the same tissue block for TDP, where bulk analyses from different donors can result in less

accurate databases for MALDI-MSI annotations. In depth protocols for each step reported below

have been deposited in the protocols.io repository.[32, 33] The workflow generally benefits from

efficient capture of dissected regions of interest within microdroplets on manufactured microPOTS

chips (**Figure 1a**). This enables minimal protein losses from iterative transfers and sample

processing prior to LC-MS/MS analyses and allows quantitative TDP analyses of singular FTU.

However, the throughput for LC-MS based TDP is low, where roughly a dozen dissected regions

can be analyzed within a day. Here MALDI-MSI presents utility, with routine processing of serial

sections only requiring a few hours. Importantly, a custom ultrahigh mass range (UHMR) Orbitrap

**(Figure 1e)** enables the detection of proteoforms at 1 to 2 Hz, allowing for broad profiling of entire

tissue sections within 6 to 24 hours (field-of-view dependent) with a near cellular or cellular scale

probe ($\leq 175$ um$^2$). This permits the profiling of several dozen FTUs. Proteoforms within these

FTUs can be relatively quantified leveraging a strong overlap between the several hundred

proteoforms confidently identified by LC-MS/MS and the several dozen proteoforms detected by

MALDI-MSI. Herein, healthy human tissues sections were used to exemplify the potential of

preclinical applications, which often requires an in-depth analysis of precious samples.

*Experimental Samples*

Tissue sections were obtained from Vanderbilt University Biomolecular Multimodal

Imaging Center (BIOMIC) which is a Tissue Mapping Center affiliated with the Human

BioMolecular Atlas Program (HuBMAP).[34, 35] Briefly, the tissues were cryo-sectioned at 10 µm

thickness on indium-tin-oxide (ITO) and polyethylene naphthalate (PEN) membrane slides, and

tissues were vacuum sealed shipped with desiccant,on dry ice, and stored at -80 $^\circ$C until analyses.

Further metadata regarding the donor can be found on the HuBMAP data portal as described in

the data availability section.

## *Laser Capture Microdissection and Sample Preparation*

A full step-by-step protocol for the processing of tissues via microPOTS is deposited in the

protocols.io repository.[33] Briefly, an injection molding company, ProtoLabs produced the

polypropylene (PP) microPOTS chips used for these experiments. The tissue samples were fixed

in 70 % ethanol for 1 min, followed by sequential dehydration in 95 % ethanol and 100 % ethanol

for 1 min each. Regions of interest were isolated from the kidney sections mounted on PEN

membrane slides using a PALM MicroBeam laser microdissection system (Carl Zeiss

MicroImaging, Munich, Germany). Tissue voxels with an area of 100,000 $\mu m^2$ were excised from

10 μm thick tissue sections for six replicates of three different FTUs including glomeruli, tubules,

and medullary rays. Each microPOTS well had a diameter of 2.2 mm and was preloaded with 2

μL of DMSO as the capture liquid. DMSO was evaporated by heating the chip at 70 $^{\circ}$C and protein

extraction was performed by adding 2 μL lysis buffer in each well, containing 2.5 units/μL

benzonase nuclease, 2 mM $MgCl_2$, 10 mM TCEP, 0.2 % DDM and 4 M urea in 50 mM ABC. The

mixture was incubated for 1 hr at 37 $^{\circ}$C. Following incubation, the sample was acidified by adding

500 nL of 5 % formic acid into each well and dried thoroughly in a vacuum chamber where the

chips were stored at -20 $^{\circ}$C until LC-MS/MS analysis.

## *Liquid Chromatography Mass Spectrometry Analysis*

A step-by-step protocol for TDP LC-MS/MS is deposited in the protocols.io repository.[33]

Briefly, custom trap columns (150 μm i.d., 4 cm long) and analytical columns (100 μm, i.d., 50

cm long) were packed with C2 particles (SMTC2MEB2-3-300, Separation Methods Technologies,

Newark, DE) in-house. LC separations were performed using a dual pump NanoAcquity system

(186016032, Waters, Millford, MA) equipped with mobile phase A (MPA; 0.2 % FA in water)

and mobile phase B (MPB; 0.2 % FA in ACN). For desalting, the samples were loaded over a

period of 10 min, and were washed with 95 % MPA at a flow rate of 5 µL/min. The analytical

gradient was started at 90% MPA and linearly ramped to 40% MPA over 90 min, where the

gradient was then ramped to 90 % MPA over 10 min at a flowrate of 3 µL/min.

An Orbitrap Fusion Lumos Tribid mass spectrometer (Thermo Scientific, San Jose, CA)

was operated in intact protein and full profile mode with data-dependent acquisition. For MS1

acquisitions, the instrument was set with the following parameters: a default charge state of 10,

resolution of 120K at $m/z$ 200, maximum injection time of 500 ms, 4 microscans, mass range from

$m/z$ 500 to 2000, a 250% normalized AGC target of 1E6, and 15V of in-source fragmentation. For

MS2 acquisitions, the instrument was set to dynamically exclude precursor ions 1 time within 30

s period, precursors were isolated with a 2 Da window for CID with fixed collision energy of 35

%, 10 ms activation time, and 0.25 activation Q. A resolution of 120K at $m/z$ 200, maximum

injection time of 500 ms, mass range from $m/z$ 400 to 2000, 2 microscans, and 4 data dependent

scans was used. All data were uploaded to the HuBMAP Data Portal for visualization and

inspection.

*Liquid Chromatography Mass Spectrometry Data Analysis*

A step-by-step protocol for processing TDP LC-MS/MS datasets is deposited in the

protocols.io repository.[33] Briefly, instrument files converted to .mzML were deconvoluted using

TopFD.[37] These exported files were analyzed within the TopPIC Suite (v.1.4.13.1)[38] and TDPortal

(v.4.0.0)[39]. TopPIC spectra were searched against UniProtKB with both SwissProt and TrEMBL

sequences (downloaded on June 29, 2019, containing 20,352 reviewed sequences). Default settings

were used for TDPortal analysis of the same datasets. TopPIC proteoform spectrum matches

(PrSMs) were then imported into the TopPICR companion package for improved identification

and quantitationqua,[40] where outputs from both TDPortal and TopPIC were combined within the

R environment. A quality control filtering step was included to remove samples with very low

identification rates. The set threshold was a minimum of 100 proteoform spectrum matches per

sample, which led to the exclusion of only two samples (one from medullary dissection and one

tubular dissection). All further data analysis steps were performed within the R environment,

including median normalization, KNN imputation, and visualization of results.[10, 41]

## *Sample Preparation for Mass Spectrometry Imaging*

A step-by-step protocol for sample preparation for intact protein MALDI-MSI is deposited

in the protocols.io repository..[32] Briefly, tissue sections were defrosted and desiccated under

vacuum for 30 mins followed immediately by serial washes in fresh solutions of 70% ethanol for

30 s, 100% ethanol for 30 s, Carnoy's solution (6:3:1 v/v ethanol: chloroform: glacial acetic acid)

for 2 mins, 100% ethanol for 30 s, water with 0.2% TFA for 15 s and 100 % ethanol for 30 s in

glass Coplin jars, each wash was completed with a volume of 70 mL. Following the washes, the

tissue sections were thoroughly dried by a stream of nitrogen gas. An M5 sprayer from HTX

Technologies (Chapel Hill, NC) was then used for acidification of the tissue section where a

solution of 5% acetic acid (*v/v*) in 50% ethanol was sprayed at a flow rate of 150 µL/min, nozzle

temperature of 30 °C, and spray velocity of 1250 mm/min with 10 PSI of nitrogen gas. A "CC"

pattern with 3 mm track spacing was used for spraying and a 5 s drying period was applied between

each of the five passes. Immediately after tissue acidification and purging of the loop, 15 mg/mL

2,5-dihyroxyacetophenone (DHA) in 90% acetonitrile with 0.2% TFA was deposited using the

same M5 sprayer described above. The sprayer was operated at a flow rate of 150 µL/min, nozzle

temperature of 30 °C, spraying velocity of 1300 mm/min with 10 PSI of nitrogen gas. The matrix

was applied in a "CC" pattern with a 2 mm track spacing. Coverage of DHA was calculated to be

~277 ug/cm$^2$. After deposition a recrystallization step was performed with a 5% acetic acid

solution in water at 38.5 °C, where the sample was equilibrated for 2 min prior to exposure to

vapor for 2 min with a 1 min drying period afterwards.

### Instrumentation for Mass Spectrometry Imaging

The operation of the instrumentation for MALDI-MSI is described in the protocols.io

repository..[32] Briefly, a MALDI source (Spectroglyph, LLC, Kennewick, WA) equipped with an

349 nm Explorer One Nd:YAG laser (SpectraPhysics, Stahnsdorf, Germany)[42] was mounted on a

custom Q Exactive HF Orbitrap MS. The Q Exactive Orbitrap MS was upgraded with Exactive

UHMR boards and was operated under custom privileges licenses.[13] Calibration of the instrument

was performed through electrospray of a solution of 2 mg/mL cesium iodide in 50% IPA. MALDI

imaging was carried out with a *m/z* range from 3,500 to 20,000 and a lowered noise thresholding

setting of 1.5 for acquisitions, resolution of 240k at *m/z* 200, ion accumulation time of 500 ms with

a laser frequency of 500 Hz pumped at 1.49 A. This resulted in 250 laser shots per pixel and

analysis by light microscopy showed that the laser spot size was roughly 12 µm by 15 µm. The

source pressure was regulated to 7.0 Torr with in-source trapping at -300 V.

### Data Processing for Mass Spectrometry Imaging

The data processing and bioinformatics workflow for MALDI-MSI annotations were

deposited in the protocols.io repository..[32] Briefly, FreeStyle (v.1.4, Thermo Scientific, Bremen,

DE) was used to view the reduced profile .RAW files. The positional data files (.xml) were

exported from the Spectroglyph, LLC source to SCiLS Lab software (v.2021c, Bruker Daltonics,

Bremen, DE) for generation of ion images with automatic settings. Processing and root mean

square transformation was completed for generation of the ion images. To guide the segmentation

of the regions of interest in the SCiLS Lab the tissue sections were stained with periodic acid-

Schiff (PAS) after MALDI-MSI experiments with matrix washed off prior. High-resolution bright-

filed images were overlayed with the MALDI images for manual segmentation. Mass errors for

each proteoform were also reported from a single isotope in the average spectrum, and all SMART

annotations are found within captions of all ion images.[43] Peak-by-Peak (Base Edition,

v.2021.8.1.b4, SpectroSwiss, Lausanne, CH) was used to average all pixels and convert spectra to

.mzML for further annotation within IsoMatchMS[44] within a search range of $m/z$ 4800 to 16000, a

noise threshold of 10 %, a Pearson correlation cutoff of 0.8, and all other parameters left at the

default values for intact proteins. No additional adducts (*i.e.*, sodium or potassium) were searched.

*Materials and Chemicals*

Optima LC-MS grade glacial acetic acid (99.99%), trifluoroacetic acid (TFA, purity 99.5

%), tris(2-carboxyethyl) phosphine (TCEP), n-dodecyl-beta-maltoside (DDM) detergent,

protease/phosphatase inhibitor cocktails (catalog 78,430), acetonitrile (ACN), isopropanol (IPA),

and  water were HPLC grade and were purchased from Fisher Scientific (Fair Lawn, NJ). Ethanol

(200 proof) was acquired from Decon Laboratories (King of Prussia, PA), while chloroform, 2',5'

dihydroxyacetophenone (2,5-DHA), cesium iodide (CsI), magnesium chloride ($MgCl_2$), phosphate

buffered saline (PBS) and ammonium bicarbonate (ABC) were purchased from Sigma Aldrich (St.

Louis, MO). Benzonase nuclease was purchased from EMD Millipore (Darmstadt, Germany).

**Results**

### *LCM-microPOTS-TDP Profiling of Human Kidney Proteoforms*

The routine dissection of FTUs was visualized through a combination of bright-field and fluorescence microscopy as highlighted in **Figure 2**. Here we applied LCM to dissect six replicates for each of three FTUs: glomeruli, distal/proximal tubules, and medullary rays. A pooled sample representing all anatomical regions was used to create a proteoform database which is referred to as "library" below to aid in annotation of the smaller dissected regions and MALDI-MSI features. In total 616 proteoforms were identified across all samples, with an average of 123 proteoforms annotated per sample. Of these proteoforms 136 had less than 50% missing values and therefore could be imputed and utilized in downstream multivariate analyses.

**Figure 2: Cortical and medullary dissections from human kidney for LCM-microPOTS-TDP. (a)** Shows a high-resolution image of the histological image taken post LCM processing for library production and dissection of glomeruli, tubules, and medullary rays. The inset region **(i)** represents the region of interest highlighted for **(b)**, **(c)**, and **(d)** where several glomeruli and tubules were dissected and the region is rotated within the zoomed visualization, where * denotes dissected glomeruli and ** denotes dissected tubules. The inset region **(ii)** represents the region of

interest highlighted for **(e)**, **(f)**, and **(g)** where several areas of the medullary rays were dissected, where # denotes a
medullary section which was dissected but not catapulted. **(b)** and **(e)** represent the autofluorescence image before
LCM, **(c)** and **(f)** are the grayscale bright-field images before LCM, and **(d)** and **(g)** are the bright-field images after
LCM. The scale bar for **(b)**, **(c)**, **(d)**, **(e)**, **(f)**, and **(g)** is 250 µm.

Notably, the pooled library sample, representing the aggregate of all FTUs, appeared to be

centered within the first two principal components and equally distant between all different FTUs

analyzed **(Figure 3a)**. Furthermore, the variance explained by the first principal component

appeared to be related to biologically relevant differences particularly for glomeruli and tubules

where separation was greatest, and the first principal component captured 44% of the total variance

**(Figure 3a** and **3b)**. For this reason, we focused our analysis on tubules and glomeruli **(Figure 3b-**

**d)**. Although only 115 proteoforms could be quantified between the dissected tubules and

glomeruli, these measurements were sufficient to cluster the samples into two clusters that aligned

with their origin **(Figure 3c)**. Notably, 20 of these proteoforms were differentially abundant with

an FDR less than 0.01 **(Figure 3d)**.

**Figure 3: LCM-microPOTS-TDP analysis of human kidney FTUs. (a)** Principal component analysis (PCA) of all replicates passing quality control derived from the three FTUs, as well as the aggregated library sample. (b) PCA of only tubule and glomeruli samples. **(c)** Heatmap of 115 proteoforms (y-axis) quantified between tubule and kidney samples (x-axis). Hierarchical clustering was applied to both proteoforms and samples. **(d)** Volcano plot of proteoforms quantified between tubules and glomeruli. Dotted lines indicate thresholds for adjusted p-values (< 0.01)

Of the proteoforms enriched in tubules, the majority (8 out of 10) were derived from mitochondrial proteins. This includes HSPE1, ATP5F1E, UQCRFS1, ATP5ME, and ATP5IF1. Many of these proteoforms are the full-length protein product, and in a few cases different proteoforms from the same protein showed similar trends in differential abundance between the

two FTUs, such as with full-length HSPE1 and a M(Ox) HSPE1 proteoform, which both had $\log_2$

fold-changes greater than 3.8. On the other hand, proteoforms enriched in glomeruli were almost

entirely derived from hemoglobin (9 out of 10).  Interestingly, proximal and distal tubules rely on

active transport mechanisms to reabsorb ions during filtration. Because tubules have higher energy

demands, they contain more mitochondria than any other structure in the kidney, which aligns with

our observations.[45] Glomeruli contain bundles of capillaries and therefore would be expected to

have more hemoglobin due to the increased concentration of red blood cells, again supporting our

observations.

Taken together, these proteoform-level differences highlight known functional

characteristics of these FTUs and provide validation of our quantitative approach with TopPIC

processing. To aid in the production of these databases the library and dissected regions were

searched with tandem use of the the TopPIC and TDPortal suite.[10] This enabled roughly 33%

increase in the number of annotations as demonstrated in a comparison of library overlap from an

exemplary bulk top-down proteomics dataset presented in the Venn diagram in **Figure 4**. Indeed,

the performance of a variety of TDP tools can be variable for the identification of proteoforms,[46]

and our approach for the automatic annotation and integration of MALDI-MSI relies heavily upon

congruency between LCM-microPOTS-TDP and MALDI-MSI.

### *Spatial Proteomics by MALDI-MSI within Human Kidney*

Validation of any annotation for MALDI-MSI was often a tedious event and required

manual matching and validation of isotopic profiles from proteins within the UniProt database.

While many tools can seamlessly annotate TDP datasets,[46] including those demonstrated above

for processing of LCM-microPOTS-TDP data, these tools cannot accommodate simplistic MS1

level spectra generated by MALDI-MSI. To accomplish this, we utilized *IsoMatchMS*, a purpose-

built program for MALDI-MSI annotation.[44] As shown within **Figure 4** isotopic envelopes are

automatically generated and visualized within trelliscope displays, which can then be sorted via

Pearson correlation of the theoretical and observed isotopic profiles. While it is still highly

recommended to visually inspect raw data, whether it be LC-MS/MS or MALDI-MSI, this level

of automation facilitates higher throughput within our methodology critical for broad adoption and

larger cohort studies.

**Figure 4: Bioinformatics workflow for annotation of MALDI-MSI datasets with LCM-microPOTS-TDP**
**annotations. (a)** Simplified schema of LC-MS/MS annotations pipeline where combined annotation using TDPortal
and TopPIC enabled roughly 33 % more annotations as exemplified within the venn diagram from a bulk TDP dataset,
ProForma strings from these aligned ProMex proteoform IDs are ingested into IsoMatchMS for isotopic profile
matching and display within Trelliscope. Isotopic matching of **(b)** truncated ubiquitin B (1-74), **(c)** intact ubiquitin B
(1-76), **(d)** truncated cytochrome c oxidase subunit 7C (17-63), **(e)** histone H4 (2-103) with a mass shift corresponding
to an acetylation and dimethylation (+70.0418 Da), and **(f)** hemoglobin subunit A (2-142) are overlaid onto the
averaged MALDI-MSI spectrum with annotation of matched and unmatched isotopes. For each respective window
the UniProt accession ID, theoretical *m/z*, observed *m/z*, and calculated error is presented.

361    Overall, from our LCM-microPOTS-TDP proteoform library containing 616 proteoforms,

362  several dozen MALDI-MSI features could be annotated at the protein level. This is exemplified

363  within **Figure 5**, where proteoforms are able to be annotated by high resolution accurate mass with

364  less than 3 ppm error,[47] with subsequent confirmation of experimental isotopic profiles shown

365  from the average MALDI-MSI spectrum within **Figure 4**. In the absence of an experimental

366  proteoform database, ion images produced by MALDI-MSI are often matched to *in-silico*

367  predicted intact masses of full-length proteins. Where inclusion or exclusion of PTMs or terminal

368  truncations can easily lead to false identifications. Here, we demonstrate how our LCM-

369  microPOTS-TDP database enabled facile identification of isomeric proteoforms, uncommon

370  proteoforms, or truncated proteoforms. Notable annotations within MALDI-MSI include a

371  truncated cytochrome c oxidase subunit 7C (17-63) **(Figure 5a)** and ubiquitin B (1-74) **(Figure**

372  **5b)**, and heavily modified positional isomers of histone H4 **(Figure 5f-h)**. This combination of a

373  cellular-resolution imaging with a broad field of view enables the facile profiling of FTUs such as

374  glomeruli within human tissues.

**Figure 5: Ion images from baseline resolved isotopes of intact proteoforms with IsoMatchMS annotations.**
The UniProt accession ID, amino acid sequence, PTMs, theoretical and observed *m/z*, and mass measurement error are reported for **(a)** cytochrome c oxidase subunit 7C, mitochondrial, **(b)** large ribosomal subunit protein eL39, **(c)** small ribosomal subunit protein eS30, **(d)** truncated ubiquitin B, **(e)** intact ubiquitin B, **(f)** histone H4 acetylated and dimethylated (N-Ac/K20me2), **(g)** histone H4 diacetylated, **(h)** histone H4 diacetylated and dimethylated, **(i)** histone H2A type E, and **(j)** hemoglobin subunit A. The scale bar for **(a)** through **(j)** is 1 mm. **(k)** Shows a high-resolution image of the histological image taken post MALDI-MSI acquisition. Several hotspots can be noted off the tissue and were identified as embedding artifacts which did not influence analyses. **SMART** annotation:[43] **S** (step size, spot size, total scans) = 40 µm, 12 µm x 15 µm, 86,189 pixels; **M** (molecular confidence) = MS1, <5 ppm; **A** (annotations) = MS1 matching from LCM-microPOTS-TDP; **R** (resolving power) = 35k at *m/z* 11306; **T** (time of acquisition) = 736 min.

This high spatial resolution broad field-of-view analysis is well in line as an ideal case study, where preclinical and diseased samples can possess distinct spatial heterogeneity as previously demonstrated on sections containing tubular atrophy or a renal cell carcinoma.[13] MALDI-MSI profiled several dozen glomeruli in this singular kidney tissue section. Although

distinct proteoform markers were not found in the detected mass range localized solely to

glomeruli, it was previously demonstrated larger proteoforms may be more region specific in

kidney than the histone and other small proteoforms detected here.[14, 48] Conversely smaller

peptides, glycans, and metabolites also uniquely localize to these FTUs more readily.[19]

This absence of succinct proteoform markers from FTUs in our detected mass range for

MALDI-MSI was not surprising given the inherent homogeneity of the proximal FTUs barring

any acute or chronic conditions and the detection of primarily epigenetic markers in these analyses.

Similarly, nearly ten-fold more proteoforms were annotated by LCM-microPOTS-TDP with only

20 proteoforms being discriminate between regions. Interestingly, we identified several ions with

unique localizations between distinct regions. For example, interfacial and medullary regions were

densely abundant with hemoglobin subunits **(Figure 5j)**. Furthermore, we identified a

differentially abundant truncation of ubiquitin as highlighted within **Figure 6**. This annotation

derived from the LCM-microPOTS-TDP database highlights the truncated form of ubiquitin (1-

74) missing the C-terminal diglycine residues **(Figure 6a),** which is detected alongside the intact

form (1-76) of ubiquitin **(Figure 6b)**.

**Figure 6: Detection of canonical and truncated forms of ubiquitin.** The UniProt accession ID, amino acid sequence, PTMs, theoretical and observed *m/z*, and mass measurement error are reported for: **(a)** truncated ubiquitin B (1-74), which is presented with a box-and-whisker plot of normalized abundance within both the cortex and medulla, where the AUC within ROC analyses is 0.7984 showing increased abundance within the cortex; and **(b)** intact ubiquitin B (1-76), which is presented with a box-and-whisker plot of normalized abundance within both the cortex and medulla, where the AUC within discriminative analyses is 0.5694 showing no increased abundance within the cortex or medulla. Scale bars for **(a)** and **(b)** are 3 mm. Several hotspots can be noted off the tissue and were identified as embedding artifacts which did not influence analyses. **SMART** annotation:[43] **S** (step size, spot size, total scans) = 40 µm, 12 µm x 15 µm, 86,189 pixels; **M** (molecular confidence) = MS1, <5 ppm; **A** (annotations) = MS1 matching from LCM-microPOTS-TDP; **R** (resolving power) = 35k at *m/z* 11306; **T** (time of acquisition) = 736 min.

Interestingly, this truncated form of ubiquitin (1-74) appeared to have distinct localization in the cortex of the kidney. Indeed, receiver operating characteristic (ROC) analysis of this proteoform across these two areas provided an area under the curve (AUC) of 0.7984. Furthermore, ubiquitin (1-74) has been previously noted as a CXC chemokine receptor 4 (CXCR4) agonist,[51] where CXCR4 has broad roles within cell proliferation, adherence, migration, and homing.[52] While the significance and function of this unique localization of ubiquitin (1-74) within the cortex is unknown, production of ubiquitin (1-74) can be mediated through a rapid diglycine cleavage by

424    insulin-degrading enzyme (IDE).[53] At the mRNA level IDE is significantly enriched within the

425    kidney and liver, and metabolism and degradation of insulin is a well-established function of FTUs

426    in the renal cortex.[54] The integrated approach using both LCM-microPOTS-TDP and MALDI-

427    MSI technology readily captured such protein processing events highlighting the potential for

428    discovering proteoform biomarkers.

## Discussion

We present an infrastructure for spatial TDP via the integration of LCM-microPOTS-TDP and

MALDI-MSI. As the technologies are rapidly evolving, and new instrumentation for intact protein

analyses becomes available, we envision that each step within our workflow (e.g., sample

preparation, data acquisition and analyses) will be further improved to enable deeper proteoform

coverage than what is currently attainable. The LCM-TDP presented herein, and in other reports,

can routinely detect hundreds of intact proteoforms,[10, 27] but application of BUP especially with

the nanoPOTS or microPOTS format routinely achieves greater proteomic depth with thousands

of proteins routinely characterized from a minimal 10,000 $\mu m^2$ dissected area.[55] Furthermore, the

detection of proteoforms by MALDI-MSI has been practically limited to several dozen unique

proteoforms with a high molecular mass limit of roughly 24 kDa.[12] Other MSI capable techniques

including nanospray desorption electrospray (nanoDESI) have recently expanded the upper

molecular mass ceiling of proteoforms by MSI on healthy human kidney to beyond 70 kDa using

highly sensitive charge detection MS.[14] However, there is a consideration that many studies using

nanoDESI-MSI detect highly abundant surface and cytoplasmic proteoforms,[15, 56] where MALDI-

MSI excels in detecting smaller regulatory and nuclear proteins, such as histones, at cellular

resolution with throughput that enables several entire sections to be routinely imaged weekly. This

leads to high complementarity, with specific use cases exemplified for each technological platform

and vast potential for integration.

Our integrated LCM-microPOTS-TDP and MALDI-MSI approach on kidney also expands on

previous efforts published from the Kidney Precision Medicine Project (KPMP), where the

understanding of disease etiology and pathophysiology within human kidneys is a paramount

goal.[19] Previously, LCM-microPOTS-BUP enabled the identification of 92 unique proteins which

were specific to either dissected proximal tubules or glomeruli with a total of 2,500 detected

proteins from roughly 10 to 40 cells, which provided a roadmap for identifying the roles of these

proteins in sub-structures of the renal corpuscle and nephron and validated affinity reagents in a

non-targeted manner.[57] While sensitivity and proteome depth is decreased with our TDP approach,

detection of intact proteins does enable the discovery of new biology from "old" proteins, with

even well characterized proteins, such as ubiquitin, have been identified in new forms and

potentially new functions.[58]

Regardless of caveats to various techniques, these approaches have found practical use within

laboratories, and large consortia spanning from HuBMAP, where the focus has been on multiple

organ systems,[34] as well as more directed efforts focused upon specific organs such as the Lung

Molecular Atlas Program (LungMAP)[59], and the Cellular Senescence Network (SenNet)[60] focused

upon crucial transitions of cells throughout life. Altogether, efforts such as these are helping to

build the Human Reference Atlas (HRA)[61] and Human Protein Atlas (HPA)[62], where proteoform

informed analyses will be fundamental to deciphering complex biological mechanisms down to

single cellular resolution and for advanced precision medicine;[3] exemplifying that a fusion of

proteomic practices is indeed the path forward for future functional proteomic studies. And further

on the horizon, integration of these advanced proteomic technologies with other -omics assays will

enable us to start deciphering fundamental principles of biology.

When considering broad applications of these techniques, there are clearly upfront investments

which are paramount to success. A persistent burden of method optimization for all spatial

proteomics demands a rigorous optimization of sample washes for each new tissue type analyzed

limited only by the creativity of the user,[63] and assay transferability and detection of targeted

proteoforms between organs or organisms cannot be guaranteed. Additionally, advanced LCM

approaches are now breaking into the subcellular regime,[64] and traditional methods employing

high-end MS platforms for TDP do not provide the sensitivity to bridge to this resolution. The next

decade will provide exciting developments such as the translation of charge detection to

chromatographic time scales,[65] and new instrumentation to further characterize proteoforms.[66] Yet,

our current approach can still be broadly applied to profile many signal peptides, hormones, and

other small proteoforms with high biological significance in many human tissues.

**Conclusions**

The data generated through the integration of LCM-microPOTS-TDP and MALDI-MSI

enabled the bridging of several hundred high confidence proteoform annotations to several dozen

direct near-cellular resolution proteoform maps. LCM-microPOTS-TDP provided a narrow field-

of-view focusing on several FTUs and identifying 20 discriminate proteoforms between glomeruli

and tubules relevant to the known physiological processes within the kidney. This analysis enabled

the production of a library of 616 proteoforms, showing high diversity of proteoforms within the

renal corpuscle and nephron as the FTUs cross distinct anatomical regions. Tandem application of

MALDI-MSI demonstrated the ability to broadly profile an entire tissue section within a practical

period at near cellular resolution, focusing upon the relative quantitation of distinct cortical and

medullary markers. Together these datasets offer unique insights into the physiological processes

of the kidney, such as the distinct localization of proteoform truncation events.

## 493 Declarations

### 494 *Ethics approval and consent to participate*

495 We have complied with all ethical regulations related to this study. All experiments on human

496 samples followed all relevant guidelines and regulations at host institutions and informed consent

497 for remnant tissue collection was acquired in accordance to Institutional Review Board policies at

498 BIOMIC at Vanderbilt University, a Tissue Mapping Center associated with the HuBMAP.

### 499 *Consent for publication*

500 Not applicable.

### 501 *Availability of data and materials*

502 The datasets supporting the conclusions of this article are available both on the HuBMAP data

503 portal, [https://doi.org/10.35079/HBM334.DQWS.354

504 https://doi.org/10.35079/HBM666.BBVZ.767], as well as on MassIVE [MSV000093722,

505 https://massive.ucsd.edu/ProteoSAFe/dataset.jsp?task=f7afef33eea446d99f72f3220ef53a64].

### 506 *Competing interests*

507 YZ is an employee of Genentech Inc. and shareholder of Roche. All other authors declare that they

508 have no competing interests.

## *Funding*

This research was funded by the National Institutes of Health (NIH) Common Fund, Human Biomolecular Atlas Program (HuBMAP) grant UG3CA256959 (LPT) and was performed at the Environmental Molecular Science Laboratory (EMSL) (doi.org/10.46936/staf.proj.2020.51770/60000309), a Department of Energy (DOE) Office of Science User Facility sponsored by the Office of Biological and Environmental Research program under Contract No. DE-AC05-76RL01830.

## *Authors' contributions*

KJZ, JMF, and RK wrote the original draft, where KJZ and JMF finalized formal analyses and discussion. KJZ completed all mass spectrometry imaging analyses, where KJZ and RK analyzed resulting datasets. DJD and LAL processed top-down annotations for mass spectrometry imaging, and DJD performed statistical analyses on these datasets. KJZ completed all imaging visualization. MV and DV processed all laser capture microdissection and microscopy analyses, where YCL, MZ, YZ, JMF, and LPT developed top-down proteomics methodology. JMF, MZ, and SMW completed all liquid chromatography analyses, and JMF processed all top-down datasets and completed all top-down visualization. MZ administered the project, where LPT, YZ, and MZ conceptualized the project, and LPT acquired funding and provided supervision. All contributed to final editing of the manuscript and all authors read and approved the final manuscript.

## *Acknowledgements*

The authors would like to acknowledge Jamie Allen and Dr. Jeffrey Spraggins at Vanderbilt University for providing the kidney tissue sections through the HuBMAP Consortium. Dr. Mikhail Belov at Spectroglyph, LLC, as well as Dr. Gordon Anderson and Chris Anderson at

GAA Custom Electronics, LLC, for technical support for the MALDI source. The authors would

also like to acknowledge Drs. Tobias Wörner, Kyle Fort, Maria Reinhardt-Szyba, and Alexander

Makarov of Thermo Fisher Scientific for technical guidance and licensing of the instrument.

Finally, we thank Dr. Neil Kelleher, Ryan Fellers, and Joe Greer from Northwestern University

for access and assistance with TDPortal and Matthew Monroe from Pacific Northwest National

Laboratory for assist with MassIVE upload.

# References

1.    Smith, L. M.; Kelleher, N. L.; Linial, M.; Goodlett, D.; Langridge-Smith, P.; Ah Goo, Y.; Safford,
G.; Bonilla*, L.; Kruppa, G.; Zubarev, R.; Rontree, J.; Chamot-Rooke, J.; Garavelli, J.; Heck, A.; Loo, J.;
Penque, D.; Hornshaw, M.; Hendrickson, C.; Pasa-Tolic, L.; Borchers, C.; Chan, D.; Young*, N.; Agar, J.;
Masselon, C.; Gross*, M.; McLafferty, F.; Tsybin, Y.; Ge, Y.; Sanders*, I.; Langridge, J.; Whitelegge*, J.;
Marshall, A.; The Consortium for Top Down, P., Proteoform: a single term describing protein complexity.
*Nature Methods* **2013,** *10* (3), 186-187.
2.    Aebersold, R.; Agar, J. N.; Amster, I. J.; Baker, M. S.; Bertozzi, C. R.; Boja, E. S.; Costello, C. E.;
Cravatt, B. F.; Fenselau, C.; Garcia, B. A.; Ge, Y.; Gunawardena, J.; Hendrickson, R. C.; Hergenrother, P.
J.; Huber, C. G.; Ivanov, A. R.; Jensen, O. N.; Jewett, M. C.; Kelleher, N. L.; Kiessling, L. L.; Krogan, N. J.;
Larsen, M. R.; Loo, J. A.; Ogorzalek Loo, R. R.; Lundberg, E.; MacCoss, M. J.; Mallick, P.; Mootha, V. K.;
Mrksich, M.; Muir, T. W.; Patrie, S. M.; Pesavento, J. J.; Pitteri, S. J.; Rodriguez, H.; Saghatelian, A.;
Sandoval, W.; Schlüter, H.; Sechi, S.; Slavoff, S. A.; Smith, L. M.; Snyder, M. P.; Thomas, P. M.; Uhlén,
M.; Van Eyk, J. E.; Vidal, M.; Walt, D. R.; White, F. M.; Williams, E. R.; Wohlschlager, T.; Wysocki, V. H.;
Yates, N. A.; Young, N. L.; Zhang, B., How many human proteoforms are there? *Nat Chem Biol* **2018,** *14*
(3), 206-214.
3.    Smith, L. M.; Agar, J. N.; Chamot-Rooke, J.; Danis, P. O.; Ge, Y.; Loo, J. A.; Paša-Tolić, L.; Tsybin,
Y. O.; Kelleher, N. L., The Human Proteoform Project: Defining the human proteome. *Sci Adv* **2021,** *7* (46),
eabk0734.
4.    Picotti, P.; Clément-Ziza, M.; Lam, H.; Campbell, D. S.; Schmidt, A.; Deutsch, E. W.; Röst, H.;
Sun, Z.; Rinner, O.; Reiter, L.; Shen, Q.; Michaelson, J. J.; Frei, A.; Alberti, S.; Kusebauch, U.; Wollscheid,
B.; Moritz, R. L.; Beyer, A.; Aebersold, R., A complete mass-spectrometric map of the yeast proteome
applied to quantitative trait analysis. *Nature* **2013,** *494* (7436), 266-270.
5.    Bludau, I.; Frank, M.; Dörig, C.; Cai, Y.; Heusel, M.; Rosenberger, G.; Picotti, P.; Collins, B. C.;
Röst, H.; Aebersold, R., Systematic detection of functional proteoform groups from bottom-up proteomic
datasets. *Nature Communications* **2021,** *12* (1), 3810.
6.    Melani, R. D.; Gerbasi, V. R.; Anderson, L. C.; Sikora, J. W.; Toby, T. K.; Hutton, J. E.; Butcher, D.
S.; Negrão, F.; Seckler, H. S.; Srzentić, K.; Fornelli, L.; Camarillo, J. M.; LeDuc, R. D.; Cesnik, A. J.;
Lundberg, E.; Greer, J. B.; Fellers, R. T.; Robey, M. T.; DeHart, C. J.; Forte, E.; Hendrickson, C. L.;
Abbatiello, S. E.; Thomas, P. M.; Kokaji, A. I.; Levitsky, J.; Kelleher, N. L., The Blood Proteoform Atlas: A
reference map of proteoforms in human hematopoietic cells. *Science* **2022,** *375* (6579), 411-418.

7.    Ardito, F.; Giuliani, M.; Perrone, D.; Troiano, G.; Lo Muzio, L., The crucial role of protein
phosphorylation in cell signaling and its use as targeted therapy (Review). *Int J Mol Med* **2017,** *40* (2), 271-
280.
8.    Kaszuba, K.; Grzybek, M.; Orłowski, A.; Danne, R.; Róg, T.; Simons, K.; Coskun, Ü.; Vattulainen,
I., N-Glycosylation as determinant of epidermal growth factor receptor conformation in membranes. *Proc
Natl Acad Sci U S A* **2015,** *112* (14), 4334-9.
9.    Chen, B.; Brown, K. A.; Lin, Z.; Ge, Y., Top-Down Proteomics: Ready for Prime Time? *Analytical
Chemistry* **2018,** *90* (1), 110-127.
10.   Liao, Y.-C.; Fulcher, J. M.; Degnan, D. J.; Williams, S. M.; Bramer, L. M.; Veličković, D.; Zemaitis,
K. J.; Veličković, M.; Sontag, R. L.; Moore, R. J.; Paša-Tolić, L.; Zhu, Y.; Zhou, M., Spatially Resolved Top-
Down Proteomics of Tissue Sections Based on a Microfluidic Nanodroplet Sample Preparation Platform.
*Molecular & Cellular Proteomics* **2023,** *22* (2).
11.   Toby, T. K.; Fornelli, L.; Kelleher, N. L., Progress in Top-Down Proteomics and the Analysis of
Proteoforms. *Annual Review of Analytical Chemistry* **2016,** *9* (1), 499-519.
12.   Prentice, B. M.; Ryan, D. J.; Van de Plas, R.; Caprioli, R. M.; Spraggins, J. M., Enhanced Ion
Transmission Efficiency up to m/ z 24 000 for MALDI Protein Imaging Mass Spectrometry. *Analytical
chemistry* **2018,** *90* (8), 5090-5099.
13.   Zemaitis, K. J.; Veličković, D.; Kew, W.; Fort, K. L.; Reinhardt-Szyba, M.; Pamreddy, A.; Ding, Y.;
Kaushik, D.; Sharma, K.; Makarov, A. A.; Zhou, M.; Paša-Tolić, L., Enhanced Spatial Mapping of Histone
Proteoforms in Human Kidney Through MALDI-MSI by High-Field UHMR-Orbitrap Detection. *Analytical
Chemistry* **2022,** *94* (37), 12604-12613.
14.   Su, P.; McGee, J. P.; Durbin, K. R.; Hollas, M. A. R.; Yang, M.; Neumann, E. K.; Allen, J. L.; Drown,
B. S.; Butun, F. A.; Greer, J. B.; Early, B. P.; Fellers, R. T.; Spraggins, J. M.; Laskin, J.; Camarillo, J. M.;
Kafader, J. O.; Kelleher, N. L., Highly multiplexed, label-free proteoform imaging of tissues by individual
ion mass spectrometry. *Science Advances* **2022,** *8* (32), eabp9929.
15.   Yang, M.; Unsihuay, D.; Hu, H.; Nguele Meke, F.; Qu, Z.; Zhang, Z. Y.; Laskin, J., Nano-DESI Mass
Spectrometry Imaging of Proteoforms in Biological Tissues with High Spatial Resolution. *Anal Chem* **2023,**
*95* (12), 5214-5222.
16.   Hale, O. J.; Cooper, H. J., Native Mass Spectrometry Imaging of Proteins and Protein Complexes
by Nano-DESI. *Anal Chem* **2021,** *93* (10), 4619-4627.
17.   Griffiths, R. L.; Konijnenberg, A.; Viner, R.; Cooper, H. J., Direct Mass Spectrometry Analysis of
Protein Complexes and Intact Proteins up to >70 kDa from Tissue. *Anal Chem* **2019,** *91* (11), 6962-6966.
18.   Hale, O. J.; Sisley, E. K.; Griffiths, R. L.; Styles, I. B.; Cooper, H. J., Native LESA TWIMS-MSI: Spatial,
Conformational, and Mass Analysis of Proteins and Protein Complexes. *Journal of the American Society
for Mass Spectrometry* **2020,** *31* (4), 873-879.
19.   Hansen, J.; Sealfon, R.; Menon, R.; Eadon, M. T.; Lake, B. B.; Steck, B.; Anjani, K.; Parikh, S.;
Sigdel, T. K.; Zhang, G.; Velickovic, D.; Barwinska, D.; Alexandrov, T.; Dobi, D.; Rashmi, P.; Otto, E. A.;
Rivera, M.; Rose, M. P.; Anderton, C. R.; Shapiro, J. P.; Pamreddy, A.; Winfree, S.; Xiong, Y.; He, Y.; de
Boer, I. H.; Hodgin, J. B.; Barisoni, L.; Naik, A. S.; Sharma, K.; Sarwal, M. M.; Zhang, K.; Himmelfarb, J.;
Rovin, B.; El-Achkar, T. M.; Laszik, Z.; He, J. C.; Dagher, P. C.; Valerius, M. T.; Jain, S.; Satlin, L. M.;
Troyanskaya, O. G.; Kretzler, M.; Iyengar, R.; Azeloglu, E. U., A reference tissue atlas for the human kidney.
*Sci Adv* **2022,** *8* (23), eabn4965.
20.   Rappez, L.; Stadler, M.; Triana, S.; Gathungu, R. M.; Ovchinnikova, K.; Phapale, P.;
Heikenwalder, M.; Alexandrov, T., SpaceM reveals metabolic states of single cells. *Nature Methods* **2021,**
*18* (7), 799-805.
21.   Pei, S.; Michael, A. R. H.; Stanislav, R.; Fatma Ayaloglu, B.; Joseph, B. G.; Bryan, P. E.; Ryan, T.
F.; Michael, A. C.; Jonathan, V. S.; Jared, O. K.; Neil, L. K., Top-down Proteomics of 10,000 Single Brain
Cells. *bioRxiv* **2023**, 2023.05.31.543176.

22.    Pu, F.; Chiang, S.; Zhang, W.; Ouyang, Z., Direct sampling mass spectrometry for clinical analysis.
*Analyst* **2019,** *144* (4), 1034-1051.
23.    Ryan, D. J.; Spraggins, J. M.; Caprioli, R. M., Protein identification strategies in MALDI imaging
mass spectrometry: a brief review. *Current Opinion in Chemical Biology* **2019,** *48*, 64-72.
24.    Han, J.; Permentier, H.; Bischoff, R.; Groothuis, G.; Casini, A.; Horvatovich, P., Imaging of protein
distribution in tissues using mass spectrometry: An interdisciplinary challenge. *TrAC Trends in Analytical*
*Chemistry* **2019,** *112*, 13-28.
25.    Ryan, D. J.; Patterson, N. H.; Putnam, N. E.; Wilde, A. D.; Weiss, A.; Perry, W. J.; Cassat, J. E.;
Skaar, E. P.; Caprioli, R. M.; Spraggins, J. M., MicroLESA: Integrating Autofluorescence Microscopy, In Situ
Micro-Digestions, and Liquid Extraction Surface Analysis for High Spatial Resolution Targeted Proteomic
Studies. *Analytical Chemistry* **2019,** *91* (12), 7578-7585.
26.    Delcourt, V.; Franck, J.; Quanico, J.; Gimeno, J.-P.; Wisztorski, M.; Raffo-Romero, A.; Kobeissy,
F.; Roucou, X.; Salzet, M.; Fournier, I., Spatially-Resolved Top-down Proteomics Bridged to MALDI MS
Imaging Reveals the Molecular Physiome of Brain Regions. *Mol Cell Proteomics* **2018,** *17* (2), 357-372.
27.    Lubeckyj, R. A.; Sun, L., Laser capture microdissection-capillary zone electrophoresis-tandem mass
spectrometry (LCM-CZE-MS/MS) for spatially resolved top-down proteomics: a pilot study of zebrafish
brain. *Molecular Omics* **2022,** *18* (2), 112-122.
28.    Zhu, Y.; Piehowski, P. D.; Zhao, R.; Chen, J.; Shen, Y.; Moore, R. J.; Shukla, A. K.; Petyuk, V. A.;
Campbell-Thompson, M.; Mathews, C. E.; Smith, R. D.; Qian, W.-J.; Kelly, R. T., Nanodroplet processing
platform for deep and quantitative proteome profiling of 10–100 mammalian cells. *Nature*
*Communications* **2018,** *9* (1), 882.
29.    Swensen, A. C.; Veličković, D.; Williams, S. M.; Moore, R. J.; Day, L. Z.; Niessen, S.; Hennessy,
S.; Posso, C.; Monetti, M.; Qian, W. J.; Jacobs, J.; Whiteley, L.; Zhu, Y.; Piehowski, P. D., Proteomic
Profiling of Intra-Islet Features Reveals Substructure-Specific Protein Signatures. *Mol Cell Proteomics*
**2022,** *21* (12), 100426.
30.    Gosline, S. J. C.; Veličković, M.; Pino, J. C.; Day, L. Z.; Attah, I. K.; Swensen, A. C.; Danna, V.;
Posso, C.; Rodland, K. D.; Chen, J.; Matthews, C. E.; Campbell-Thompson, M.; Laskin, J.; Burnum-
Johnson, K.; Zhu, Y.; Piehowski, P. D., Proteome Mapping of the Human Pancreatic Islet
Microenvironment Reveals Endocrine–Exocrine Signaling Sphere of Influence. *Molecular & Cellular*
*Proteomics* **2023,** *22* (8), 100592.
31.    Zhou, M.; Uwugiaren, N.; Williams, S. M.; Moore, R. J.; Zhao, R.; Goodlett, D.; Dapic, I.; Paša-
Tolić, L.; Zhu, Y., Sensitive Top-Down Proteomics Analysis of a Low Number of Mammalian Cells Using a
Nanodroplet Sample Processing Platform. *Anal Chem* **2020,** *92* (10), 7087-7095.
32.    Zemaitis, K. J.; Veličković, D.; Degnan, D.; Zhou, M.; Paša-Tolić, L., Overall protocol for 2D intact
proteoform    mapping    by    MALDI    imaging.    *protocols.io*    **2022**.
https://dx.doi.org/10.17504/protocols.io.q26g74qn8gwz/v1
33.    Zhou, M.; Fulcher, J.; Liao, Y.-C.; L, P.-T., Overall protocol for MicroPOTS LCMS top down
proteomics    of    kidney    tissue    sections.    *protocols.io*    **2022**.
dx.doi.org/10.17504/protocols.io.bp2l61oy5vqe/v1
34.    Jain, S.; Pei, L.; Spraggins, J. M.; Angelo, M.; Carson, J. P.; Gehlenborg, N.; Ginty, F.; Gonçalves,
J. P.; Hagood, J. S.; Hickey, J. W.; Kelleher, N. L.; Laurent, L. C.; Lin, S.; Lin, Y.; Liu, H.; Naba, A.; Nakayasu,
E. S.; Qian, W.-J.; Radtke, A.; Robson, P.; Stockwell, B. R.; Van de Plas, R.; Vlachos, I. S.; Zhou, M.; Ahn,
K. J.; Allen, J.; Anderson, D. M.; Anderton, C. R.; Curcio, C.; Angelin, A.; Arvanitis, C.; Atta, L.; Awosika-
Olumo, D.; Bahmani, A.; Bai, H.; Balderrama, K.; Balzano, L.; Bandyopadhyay, G.; Bandyopadhyay, S.;
Bar-Joseph, Z.; Barnhart, K.; Barwinska, D.; Becich, M.; Becker, L.; Becker, W.; Bedi, K.; Bendall, S.;
Benninger, K.; Betancur, D.; Bettinger, K.; Billings, S.; Blood, P.; Bolin, D.; Border, S.; Bosse, M.; Bramer,
L.; Brewer, M.; Brusko, M.; Bueckle, A.; Burke, K.; Burnum-Johnson, K.; Butcher, E.; Butterworth, E.;
Cai, L.; Calandrelli, R.; Caldwell, M.; Campbell-Thompson, M.; Cao, D.; Cao-Berg, I.; Caprioli, R.; Caraccio,

C.; Caron, A.; Carroll, M.; Chadwick, C.; Chen, A.; Chen, D.; Chen, F.; Chen, H.; Chen, J.; Chen, L.; Chen, L.; Chiacchia, K.; Cho, S.; Chou, P.; Choy, L.; Cisar, C.; Clair, G.; Clarke, L.; Clouthier, K. A.; Colley, M. E.; Conlon, K.; Conroy, J.; Contrepois, K.; Corbett, A.; Corwin, A.; Cotter, D.; Courtois, E.; Cruz, A.; Csonka, C.; Czupil, K.; Daiya, V.; Dale, K.; Davanagere, S. A.; Dayao, M.; de Caestecker, M. P.; Decker, A.; Deems, S.; Degnan, D.; Desai, T.; Deshpande, V.; Deutsch, G.; Devlin, M.; Diep, D.; Dodd, C.; Donahue, S.; Dong, W.; dos Santos Peixoto, R.; Duffy, M.; Dufresne, M.; Duong, T. E.; Dutra, J.; Eadon, M. T.; El-Achkar, T. M.; Enninful, A.; Eraslan, G.; Eshelman, D.; Espin-Perez, A.; Esplin, E. D.; Esselman, A.; Falo, L. D.; Falo, L.; Fan, J.; Fan, R.; Farrow, M. A.; Farzad, N.; Favaro, P.; Fermin, J.; Filiz, F.; Filus, S.; Fisch, K.; Fisher, E.; Fisher, S.; Flowers, K.; Flynn, W. F.; Fogo, A. B.; Fu, D.; Fulcher, J.; Fung, A.; Furst, D.; Gallant, M.; Gao, F.; Gao, Y.; Gaulton, K.; Gaut, J. P.; Gee, J.; Ghag, R. R.; Ghazanfar, S.; Ghose, S.; Gisch, D.; Gold, I.; Gondalia, A.; Gorman, B.; Greenleaf, W.; Greenwald, N.; Gregory, B.; Guo, R.; Gupta, R.; Hakimian, H.; Haltom, J.; Halushka, M.; Han, K. S.; Hanson, C.; Harbury, P.; Hardi, J.; Harlan, L.; Harris, R. C.; Hartman, A.; Heidari, E.; Helfer, J.; Helminiak, D.; Hemberg, M.; Henning, N.; Herr, B. W.; Ho, J.; Holden-Wiltse, J.; Hong, S.-H.; Hong, Y.-K.; Honick, B.; Hood, G.; Hu, P.; Hu, Q.; Huang, M.; Huyck, H.; Imtiaz, T.; Isberg, O. G.; Itkin, M.; Jackson, D.; Jacobs, M.; Jain, Y.; Jewell, D.; Jiang, L.; Jiang, Z. G.; Johnston, S.; Joshi, P.; Ju, Y.; Judd, A.; Kagel, A.; Kahn, A.; Kalavros, N.; Kalhor, K.; Karagkouni, D.; Karathanos, T.; Karunamurthy, A.; Katari, S.; Kates, H.; Kaushal, M.; Keener, N.; Keller, M.; Kenney, M.; Kern, C.; Kharchenko, P.; Kim, J.; Kingsford, C.; Kirwan, J.; Kiselev, V.; Kishi, J.; Kitata, R. B.; Knoten, A.; Kollar, C.; Krishnamoorthy, P.; Kruse, A. R. S.; Da, K.; Kundaje, A.; Kutschera, E.; Kwon, Y.; Lake, B. B.; Lancaster, S.; Langlieb, J.; Lardenoije, R.; Laronda, M.; Laskin, J.; Lau, K.; Lee, H.; Lee, M.; Lee, M.; Strekalova, Y. L.; Li, D.; Li, J.; Li, J.; Li, X.; Li, Z.; Liao, Y.-C.; Liaw, T.; Lin, P.; Lin, Y.; Lindsay, S.; Liu, C.; Liu, Y.; Liu, Y.; Lott, M.; Lotz, M.; Lowery, L.; Lu, P.; Lu, X.; Lucarelli, N.; Lun, X.; Luo, Z.; Ma, J.; Macosko, E.; Mahajan, M.; Maier, L.; Makowski, D.; Malek, M.; Manthey, D.; Manz, T.; Margulies, K.; Marioni, J.; Martindale, M.; Mason, C.; Mathews, C.; Maye, P.; McCallum, C.; McDonough, E.; McDonough, L.; McDowell, H.; Meads, M.; Medina-Serpas, M.; Ferreira, R. M.; Messinger, J.; Metis, K.; Migas, L. G.; Miller, B.; Mimar, S.; Minor, B.; Misra, R.; Missarova, A.; Mistretta, C.; Moens, R.; Moerth, E.; Hu, B. C., Advances and prospects for the Human BioMolecular Atlas Program (HuBMAP). *Nature Cell Biology* **2023**.

35.     Snyder, M. P.; Lin, S.; Posgai, A.; Atkinson, M.; Regev, A.; Rood, J.; Rozenblatt-Rosen, O.; Gaffney, L.; Hupalowska, A.; Satija, R.; Gehlenborg, N.; Shendure, J.; Laskin, J.; Harbury, P.; Nystrom, N. A.; Silverstein, J. C.; Bar-Joseph, Z.; Zhang, K.; Börner, K.; Lin, Y.; Conroy, R.; Procaccini, D.; Roy, A. L.; Pillai, A.; Brown, M.; Galis, Z. S.; Cai, L.; Shendure, J.; Trapnell, C.; Lin, S.; Jackson, D.; Snyder, M. P.; Nolan, G.; Greenleaf, W. J.; Lin, Y.; Plevritis, S.; Ahadi, S.; Nevins, S. A.; Lee, H.; Schuerch, C. M.; Black, S.; Venkataraaman, V. G.; Esplin, E.; Horning, A.; Bahmani, A.; Zhang, K.; Sun, X.; Jain, S.; Hagood, J.; Pryhuber, G.; Kharchenko, P.; Atkinson, M.; Bodenmiller, B.; Brusko, T.; Clare-Salzler, M.; Nick, H.; Otto, K.; Posgai, A.; Wasserfall, C.; Jorgensen, M.; Brusko, M.; Maffioletti, S.; Caprioli, R. M.; Spraggins, J. M.; Gutierrez, D.; Patterson, N. H.; Neumann, E. K.; Harris, R.; deCaestecker, M.; Fogo, A. B.; van de Plas, R.; Lau, K.; Cai, L.; Yuan, G.-C.; Zhu, Q.; Dries, R.; Yin, P.; Saka, S. K.; Kishi, J. Y.; Wang, Y.; Goldaracena, I.; Laskin, J.; Ye, D.; Burnum-Johnson, K. E.; Piehowski, P. D.; Ansong, C.; Zhu, Y.; Harbury, P.; Desai, T.; Mulye, J.; Chou, P.; Nagendran, M.; Bar-Joseph, Z.; Teichmann, S. A.; Paten, B.; Murphy, R. F.; Ma, J.; Kiselev, V. Y.; Kingsford, C.; Ricarte, A.; Keays, M.; Akoju, S. A.; Ruffalo, M.; Gehlenborg, N.; Kharchenko, P.; Vella, M.; McCallum, C.; Börner, K.; Cross, L. E.; Friedman, S. H.; Heiland, R.; Herr, B.; Macklin, P.; Quardokus, E. M.; Record, L.; Sluka, J. P.; Weber, G. M.; Nystrom, N. A.; Silverstein, J. C.; Blood, P. D.; Ropelewski, A. J.; Shirey, W. E.; Scibek, R. M.; Mabee, P.; Lenhardt, W. C.; Robasky, K.; Michailidis, S.; Satija, R.; Marioni, J.; Regev, A.; Butler, A.; Stuart, T.; Fisher, E.; Ghazanfar, S.; Rood, J.; Gaffney, L.; Eraslan, G.; Biancalani, T.; Vaishnav, E. D.; Conroy, R.; Procaccini, D.; Roy, A.; Pillai, A.; Brown, M.; Galis, Z.; Srinivas, P.; Pawlyk, A.; Sechi, S.; Wilder, E.; Anderson, J.; Hu, B. C.; Writing, G.; Caltech, U. T.; Stanford-Wash, U. T.; Ucsd, T. M. C.; University of Florida, T. M. C.; Vanderbilt University,

T. M. C.; California Institute of Technology, T. T. D.; Harvard, T. T. D.; Purdue, T. T. D.; Stanford, T. T. D.;
HuBmap Integration, V.; Engagement Collaboratory: Carnegie Mellon, T. C.; Harvard Medical School, T.
C.; Indiana University Bloomington, M. C.; Pittsburgh Supercomputing, C.; University of Pittsburgh, I.;
Engagement, C.; University of South Dakota, C. C.; New York Genome Center, M. C.; Group, N. I. H. H. W.,
The human body at cellular resolution: the NIH Human Biomolecular Atlas Program. *Nature* **2019,** *574*
(7777), 187-192.
36.    Neumann, E. K.; allen, J.; Brewer, M.; Anderson, D. M.; De Caestecker, M. P.; Gutierrez, D.;
Spraggins, J. M., VU Biomolecular Multimodal Imaging Center (BIOMIC) kidney characterization pipeline
for tissues collected through the Cooperative Human Tissue Network (CHTN). *protocols.io* **2020**.
37.    Basharat, A. R.; Zang, Y.; Sun, L.; Liu, X., TopFD: A Proteoform Feature Detection Tool for Top–
Down Proteomics. *Analytical Chemistry* **2023,** *95* (21), 8189-8196.
38.    Kou, Q.; Xun, L.; Liu, X., TopPIC: a software tool for top-down mass spectrometry-based
proteoform identification and characterization. *Bioinformatics* **2016,** *32* (22), 3495-3497.
39.    Toby, T. K.; Fornelli, L.; Srzentić, K.; DeHart, C. J.; Levitsky, J.; Friedewald, J.; Kelleher, N. L., A
comprehensive pipeline for translational top-down proteomics from a single blood draw. *Nature Protocols*
**2019,** *14* (1), 119-152.
40.    Martin, E. A.; Fulcher, J. M.; Zhou, M.; Monroe, M. E.; Petyuk, V. A., TopPICR: A Companion R
Package for Top-Down Proteomics Data Analysis. *Journal of Proteome Research* **2023,** *22* (2), 399-409.
41.    Park, J.; Piehowski, P. D.; Wilkins, C.; Zhou, M.; Mendoza, J.; Fujimoto, G. M.; Gibbons, B. C.;
Shaw, J. B.; Shen, Y.; Shukla, A. K.; Moore, R. J.; Liu, T.; Petyuk, V. A.; Tolić, N.; Paša-Tolić, L.; Smith, R.
D.; Payne, S. H.; Kim, S., Informed-Proteomics: open-source software package for top-down proteomics.
*Nature Methods* **2017,** *14* (9), 909-914.
42.    Belov, M. E.; Ellis, S. R.; Dilillo, M.; Paine, M. R. L.; Danielson, W. F.; Anderson, G. A.; de Graaf,
E. L.; Eijkel, G. B.; Heeren, R. M. A.; McDonnell, L. A., Design and Performance of a Novel Interface for
Combined Matrix-Assisted Laser Desorption Ionization at Elevated Pressure and Electrospray Ionization
with Orbitrap Mass Spectrometry. *Analytical Chemistry* **2017,** *89* (14), 7493-7501.
43.    Xi, Y.; Sohn, A. L.; Joignant, A. N.; Cologna, S. M.; Prentice, B. M.; Muddiman, D. C., SMART: A
data reporting standard for mass spectrometry imaging. *Journal of Mass Spectrometry* **2023,** *58* (2),
e4904.
44.    Degnan, D. J.; Zemaitis, K. J.; Lewis, L. A.; McCue, L. A.; Bramer, L. M.; Fulcher, J. M.; Veličković,
D.; Paša-Tolić, L.; Zhou, M., IsoMatchMS: Open-Source Software for Automated Annotation and
Visualization of High Resolution MALDI-MS Spectra. *Journal of the American Society for Mass*
*Spectrometry* **2023**.
45.    Bhargava, P.; Schnellmann, R. G., Mitochondrial energetics in the kidney. *Nature Reviews*
*Nephrology* **2017,** *13* (10), 629-646.
46.    Tabb, D. L.; Jeong, K.; Druart, K.; Gant, M. S.; Brown, K. A.; Nicora, C.; Zhou, M.; Couvillion, S.;
Nakayasu, E.; Williams, J. E.; Peterson, H. K.; McGuire, M. K.; McGuire, M. A.; Metz, T. O.; Chamot-
Rooke, J., Comparing Top-Down Proteoform Identification: Deconvolution, PrSM Overlap, and PTM
Detection. *Journal of Proteome Research* **2023,** *22* (7), 2199-2217.
47.    Spraggins, J. M.; Rizzo, D. G.; Moore, J. L.; Rose, K. L.; Hammer, N. D.; Skaar, E. P.; Caprioli, R.
M., MALDI FTICR IMS of Intact Proteins: Using Mass Accuracy to Link Protein Images with Proteomics Data.
*Journal of the American Society for Mass Spectrometry* **2015,** *26* (6), 974-985.
48.    Neumann, E. K.; Patterson, N. H.; Rivera, E. S.; Allen, J. L.; Brewer, M.; deCaestecker, M. P.;
Caprioli, R. M.; Fogo, A. B.; Spraggins, J. M., Highly multiplexed immunofluorescence of the human kidney
using co-detection by indexing. *Kidney Int* **2022,** *101* (1), 137-143.
49.    Smith, L. M.; Thomas, P. M.; Shortreed, M. R.; Schaffer, L. V.; Fellers, R. T.; LeDuc, R. D.;
Tucholski, T.; Ge, Y.; Agar, J. N.; Anderson, L. C.; Chamot-Rooke, J.; Gault, J.; Loo, J. A.; Paša-Tolić, L.;

Robinson, C. V.; Schlüter, H.; Tsybin, Y. O.; Vilaseca, M.; Vizcaíno, J. A.; Danis, P. O.; Kelleher, N. L., A five-level classification system for proteoform identifications. *Nature Methods* **2019,** *16* (10), 939-940.

50.      Zemaitis, K. J.; Zhou, M.; Kew, W.; Paša-Tolić, L., 193 nm Ultraviolet Photodissociation for the Characterization of Singly Charged Proteoforms Generated by MALDI. *Journal of the American Society for Mass Spectrometry* **2023,** *34* (2), 328-332.

51.      Tripathi, A.; Saini, V.; Marchese, A.; Volkman, B. F.; Tang, W.-J.; Majetschak, M., Modulation of the CXC Chemokine Receptor 4 Agonist Activity of Ubiquitin through C-Terminal Protein Modification. *Biochemistry* **2013,** *52* (24), 4184-4192.

52.      Bianchi, M. E.; Mezzapelle, R., The Chemokine Receptor CXCR4 in Cell Proliferation and Tissue Regeneration. *Frontiers in Immunology* **2020,** *11*.

53.      Ralat, L. A.; Kalas, V.; Zheng, Z.; Goldman, R. D.; Sosnick, T. R.; Tang, W.-J., Ubiquitin Is a Novel Substrate for Human Insulin-Degrading Enzyme. *Journal of Molecular Biology* **2011,** *406* (3), 454-466.

54.      Rabkin, R.; Ryan, M. P.; Duckworth, W. C., The renal metabolism of insulin. *Diabetologia* **1984,** *27* (3), 351-7.

55.      Piehowski, P. D.; Zhu, Y.; Bramer, L. M.; Stratton, K. G.; Zhao, R.; Orton, D. J.; Moore, R. J.; Yuan, J.; Mitchell, H. D.; Gao, Y.; Webb-Robertson, B.-J. M.; Dey, S. K.; Kelly, R. T.; Burnum-Johnson, K. E., Automated mass spectrometry imaging of over 2000 proteins from tissue sections at 100-µm spatial resolution. *Nature Communications* **2020,** *11* (1), 8.

56.      Yang, M.; Hu, H.; Su, P.; Thomas, P. M.; Camarillo, J. M.; Greer, J. B.; Early, B. P.; Fellers, R. T.; Kelleher, N. L.; Laskin, J., Proteoform-Selective Imaging of Tissues Using Mass Spectrometry**. *Angewandte Chemie International Edition* **2022,** *61* (29), e202200721.

57.      Sigdel, T. K.; Piehowski, P. D.; Roy, S.; Liberto, J.; Hansen, J. R.; Swensen, A. C.; Zhao, R.; Zhu, Y.; Rashmi, P.; Schroeder, A.; Damm, I.; Sur, S.; Luo, J.; Yang, Y.; Qian, W. J.; Sarwal, M. M., Near-Single-Cell Proteomics Profiling of the Proximal Tubular and Glomerulus of the Normal Human Kidney. *Front Med (Lausanne)* **2020,** *7*, 499.

58.      Burnum-Johnson, K. E.; Conrads, T. P.; Drake, R. R.; Herr, A. E.; Iyengar, R.; Kelly, R. T.; Lundberg, E.; MacCoss, M. J.; Naba, A.; Nolan, G. P.; Pevzner, P. A.; Rodland, K. D.; Sechi, S.; Slavov, N.; Spraggins, J. M.; Van Eyk, J. E.; Vidal, M.; Vogel, C.; Walt, D. R.; Kelleher, N. L., New Views of Old Proteins: Clarifying the Enigmatic Proteome. *Molecular & Cellular Proteomics* **2022,** *21* (7), 100254.

59.      Ardini-Poleske, M. E.; Clark, R. F.; Ansong, C.; Carson, J. P.; Corley, R. A.; Deutsch, G. H.; Hagood, J. S.; Kaminski, N.; Mariani, T. J.; Potter, S. S.; Pryhuber, G. S.; Warburton, D.; Whitsett, J. A.; Palmer, S. M.; Ambalavanan, N.; Consortium, T. L., LungMAP: The Molecular Atlas of Lung Development Program. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **2017,** *313* (5), L733-L740.

60.      Lee, P. J.; Benz, C. C.; Blood, P.; Börner, K.; Campisi, J.; Chen, F.; Daldrup-Link, H.; De Jager, P.; Ding, L.; Duncan, F. E.; Eickelberg, O.; Fan, R.; Finkel, T.; Furman, D.; Garovic, V.; Gehlenborg, N.; Glass, C.; Heckenbach, I.; Joseph, Z.-B.; Katiyar, P.; Kim, S.-J.; Königshoff, M.; Kuchel, G. A.; Lee, H.; Lee, J. H.; Ma, J.; Ma, Q.; Melov, S.; Metis, K.; Mora, A. L.; Musi, N.; Neretti, N.; Passos, J. F.; Rahman, I.; Rivera-Mulia, J. C.; Robson, P.; Rojas, M.; Roy, A. L.; Scheibye-Knudsen, M.; Schilling, B.; Shi, P.; Silverstein, J. C.; Suryadevara, V.; Xie, J.; Wang, J.; Wong, A. I.; Niedernhofer, L. J.; Wang, S.; Anvari, H.; Balough, J.; Benz, C.; Bons, J.; Brenerman, B.; Evans, W.; Gerencser, A.; Gregory, H.; Hansen, M.; Justice, J.; Kapahi, P.; Murad, N.; O'Broin, A.; Pavone, M. E.; Powell, M.; Scott, G.; Shanes, E.; Shankaran, M.; Verdin, E.; Winer, D.; Wu, F.; Adams, A.; Blood, P. D.; Bueckle, A.; Cao-Berg, I.; Chen, H.; Davis, M.; Filus, S.; Hao, Y.; Hartman, A.; Hasanaj, E.; Helfer, J.; Herr, B.; Joseph, Z. B.; Molla, G.; Mou, G.; Puerto, J.; Quardokus, E. M.; Ropelewski, A. J.; Ruffalo, M.; Satija, R.; Schwenk, M.; Scibek, R.; Shirey, W.; Sibilla, M.; Welling, J.; Yuan, Z.; Bonneau, R.; Christiano, A.; Izar, B.; Menon, V.; Owens, D. M.; Phatnani, H.; Smith, C.; Suh, Y.; Teich, A. F.; Bekker, V.; Chan, C.; Coutavas, E.; Hartwig, M. G.; Ji, Z.; Nixon, A. B.; Dou, Z.; Rajagopal, J.; Slavov, N.; Holmes, D.; Jurk, D.; Kirkland, J. L.; Lagnado, A.; Tchkonia, T.; Abraham, K.; Dibattista, A.; Fridell, Y.-W.; Howcroft, T. K.; Jhappan, C.; Montes, V. P.; Prabhudas, M.; Resat, H.;

Taylor, V.; Kumar, M.; Suryadevara, V.; Cigarroa, F.; Cohn, R.; Cortes, T. M.; Courtois, E.; Chuang, J.;
Davé, M.; Domanskyi, S.; Enninga, E. A. L.; Eryilmaz, G. N.; Espinoza, S. E.; Gelfond, J.; Kirkland, J.;
Kuchel, G. A.; Kuo, C.-L.; Lehman, J. S.; Aguayo-Mazzucato, C.; Meves, A.; Rani, M.; Sanders, S.;
Thibodeau, A.; Tullius, S. G.; Ucar, D.; White, B.; Wu, Q.; Xu, M.; Yamaguchi, S.; Assarzadegan, N.; Cho,
C.-S.; Hwang, I.; Hwang, Y.; Xi, J.; Adeyi, O. A.; Aliferis, C. F.; Bartolomucci, A.; Dong, X.; DuFresne-To,
M. J.; Ikramuddin, S.; Johnson, S. G.; Nelson, A. C.; Niedernhofer, L. J.; Revelo, X. S.; Trevilla-Garcia, C.;
Sedivy, J. M.; Thompson, E. L.; Robbins, P. D.; Wang, J.; Aird, K. M.; Alder, J. K.; Beaulieu, D.; Bueno,
M.; Calyeca, J.; Chamucero-Millaris, J. A.; Chan, S. Y.; Chung, D.; Corbett, A.; Gorbunova, V.; Gowdy, K.
M.; Gurkar, A.; Horowitz, J. C.; Hu, Q.; Kaur, G.; Khaliullin, T. O.; Lafyatis, R.; Lanna, S.; Li, D.; Ma, A.;
Morris, A.; Muthumalage, T. M.; Peters, V.; Pryhuber, G. S.; Reader, B. F.; Rosas, L.; Sembrat, J. C.;
Shaikh, S.; Shi, H.; Stacey, S. D.; Croix, C. S.; Wang, C.; Wang, Q.; Watts, A.; Gu, L.; Lin, Y.; Rabinovitch,
P. S.; Sweetwyne, M. T.; Artyomov, M. N.; Ballentine, S. J.; Chheda, M. G.; Davies, S. R.; DiPersio, J. F.;
Fields, R. C.; Fitzpatrick, J. A. J.; Fulton, R. S.; Imai, S.-i.; Jain, S.; Ju, T.; Kushnir, V. M.; Link, D. C.; Ben
Major, M.; Oh, S. T.; Rapp, D.; Rettig, M. P.; Stewart, S. A.; Veis, D. J.; Vij, K. R.; Wendl, M. C.;
Wyczalkowski, M. A.; Craft, J. E.; Enninful, A.; Farzad, N.; Gershkovich, P.; Halene, S.; Kluger, Y.;
VanOudenhove, J.; Xu, M.; Yang, J.; Yang, M.; SenNet, C.; Writing, G.; Brown University, T. D. A.; Buck
Institute for Research on Aging, T. T.; Consortium, O.; Data Coordinating, C.; Columbia, T. M. C.; Duke
University, T. M. C.; Massachusetts General Hospital, T. D. A.; Mayo Clinic, T. D. A.; National Institute of,
H.; Stanford, T. D. A.; University of Connecticut, T. M. C.; University of Michigan, T. D. A.; University of
Minnesota, T. M. C.; University of Pittsburgh, T. M. C.; University of Washington, T. D. A.; Washington
University, T. M. C.; Yale, T. M. C., NIH SenNet Consortium to map senescent cells throughout the human
lifespan to understand physiological health. *Nature Aging* **2022,** *2* (12), 1090-1100.
61.      Börner, K.; Teichmann, S. A.; Quardokus, E. M.; Gee, J. C.; Browne, K.; Osumi-Sutherland, D.;
Herr, B. W.; Bueckle, A.; Paul, H.; Haniffa, M.; Jardine, L.; Bernard, A.; Ding, S.-L.; Miller, J. A.; Lin, S.;
Halushka, M. K.; Boppana, A.; Longacre, T. A.; Hickey, J.; Lin, Y.; Valerius, M. T.; He, Y.; Pryhuber, G.;
Sun, X.; Jorgensen, M.; Radtke, A. J.; Wasserfall, C.; Ginty, F.; Ho, J.; Sunshine, J.; Beuschel, R. T.;
Brusko, M.; Lee, S.; Malhotra, R.; Jain, S.; Weber, G., Anatomical structures, cell types and biomarkers of
the Human Reference Atlas. *Nature Cell Biology* **2021,** *23* (11), 1117-1128.
62.      Uhlén, M.; Fagerberg, L.; Hallström, B. M.; Lindskog, C.; Oksvold, P.; Mardinoglu, A.; Sivertsson,
Å.; Kampf, C.; Sjöstedt, E.; Asplund, A., Tissue-based map of the human proteome. *Science* **2015,** *347*
(6220), 1260419.
63.      Rešetar Maslov, D.; Svirkova, A.; Allmaier, G.; Marchetti-Deschamann, M.; Kraljević Pavelić, S.,
Optimization of MALDI-TOF mass spectrometry imaging for the visualization and comparison of peptide
distributions in dry-cured ham muscle fibers. *Food Chem* **2019,** *283*, 275-286.
64.      Xiang, P.; Liyu, A.; Kwon, Y.; Hu, D.; Williams, S. M.; Veličković, D.; Markillie, L. M.; Chrisler, W.
B.; Paša-Tolić, L.; Zhu, Y., Spatial Proteomics toward Subcellular Resolution by Coupling Deep Ultraviolet
Laser Ablation with Nanodroplet Sample Preparation. *ACS Measurement Science Au* **2023**.
65.      Strasser, L.; Füssl, F.; Morgan, T. E.; Carillo, S.; Bones, J., Exploring Charge-Detection Mass
Spectrometry on Chromatographic Time Scales. *Analytical Chemistry* **2023,** *95* (40), 15118-15124.
66.      Papanastasiou, D.; Kounadis, D.; Lekkas, A.; Orfanopoulos, I.; Mpozatzidis, A.; Smyrnakis, A.;
Panagiotopoulos, E.; Kosmopoulou, M.; Reinhardt-Szyba, M.; Fort, K.; Makarov, A.; Zubarev, R. A., The
Omnitrap Platform: A Versatile Segmented Linear Ion Trap for Multidimensional Multiple-Stage Tandem
Mass Spectrometry. *Journal of the American Society for Mass Spectrometry* **2022,** *33* (10), 1990-2007.

850