Enhanced Fc Receptor expression and distinct immunomodulatory properties of neutrophils during HIV-1 infection and therapy

Soledad Marsile-Medun[1], Manon Souchard[1], Daouda Abba Moussa[1], Valérie Lorin[2], Hugo Mouquet[2], Elisa Reynaud[3], Rayane Dibsy[4], Edouard Tuaillon[3], Delphine Muriaux[4,5], Giang Ngo[6], Martine Pugnière[6], Mar Naranjo-Gomez[1*], Mireia Pelegrin[1*]

Addresses:

[1] IRMB, Univ Montpellier, INSERM, CNRS, Montpellier, France

[2] Humoral Immunology Unit, Institut Pasteur, Université Paris Cité, INSERM U1222, 75015 Paris, France

[3] Laboratoire de Virologie, Centre Hospitalier Universitaire de Montpellier, France

[4] IRIM, Université de Montpellier, UMR 9004 CNRS, Montpellier, France.

[5] CEMIPAI, University of Montpellier, UAR3725 CNRS, Montpellier, France.

[6] IRCM, Univ Montpellier, ICM, INSERM, Montpellier, France

[*] equally contributed

*Corresponding authors:

Mar Naranjo-Gomez and Mireia Pelegrin

Institute of Regenerative Medicine and Biotherapy of Montpellier, UMR1183

80, Avenue Agustin Fliche

34293 Montpellier Cedex 5

France

Phone number: + 33 4 67 33 52 67

maria.naranjo@inserm.fr; mireia.pelegrin@inserm.fr

Keywords

HIV-1, neutrophils, immunomodulation, immune complexes, broadly neutralizing antibodies.

Abstract

Neutrophils are innate immune cells that display multiple immunomodulatory properties. Using a mouse model of retroviral infection, we previously uncovered their key immunomodulatory function in promoting protective immunity during antiviral antibody therapy, which occurs through Fc-FcγR interactions. Here, we investigated the immunomodulatory properties of neutrophils in the context of HIV-1 infection and therapy. We conducted a thorough examination of the functional activation and modulation of Fcγ receptors (FcγRs) expression on neutrophils isolated from healthy donors (HD) and people living with HIV-1 (PLWH). Upon stimulation with virions, either free or in the form of immune complexes (ICs) formed with broadly neutralizing antibodies (bNAbs), neutrophils from HD secreted cytokines and chemokines in modest quantities. Interestingly, priming HD neutrophils with pro-inflammatory cytokines augmented their chemokine production upon exposure to HIV-1 and ICs, underscoring the pivotal role of the inflammatory milieu in eliciting a functional response from neutrophils to these stimuli. Noteworthy, under inflammatory conditions, IC-activated neutrophils showed higher secretion of CXCL1 and CCL4 than HIV-1-stimulated neutrophils. Highlighting the critical influence of the inflammatory environment on their functional properties, neutrophils from PLWH displayed increased release of cytokines (IFNγ) and chemokines (CXCL1, CCL2, CCL3 and CCL4) when stimulated with danger signals associated with HIV-1 infection. In addition, PLWH's neutrophils exhibited higher expression of two activating FcγRs (FcγRI and FcγRIII) than neutrophils from HD. Our study provides new insights into the immunomodulatory properties of neutrophils during HIV-1 infection and therapy. These findings hold potential for optimizing antiviral antibody therapies by harnessing the immunomodulatory properties of neutrophils.

Introduction

Neutrophils are the most abundant population of circulating white blood cells and are massively and rapidly recruited following infection. Through diverse mechanisms such as phagocytosis, releasing of reactive-oxygen species (ROS), trapping of the pathogenic agent by neutrophil extracellular traps (NET) formation (known as NETosis) (1), neutrophils can limit and counteract invading pathogens. Moreover, immunomodulatory properties of neutrophils have been described in recent years, showing their ability to shape the adaptive immune response (2,3). Neutrophils migrate from the site of infection to draining lymph nodes where they interact with multiple immune cells through the production of several cytokines and chemokines (3), making neutrophils key cells in the orchestration of innate and adaptive immunity (3,4). However, the immunomodulatory role of neutrophils in antiviral immunity is still ill-understood, including in the context of persistent viral infections such as HIV-1, as well as in the frame of antiviral monoclonal antibody (mAb)-based immunotherapies. Noteworthy, it has been reported that the functional activation of neutrophils is compromised upon HIV-1 infection which leads to impaired effector functions (i.e. phagocytosis, ROS release, …) and increased apoptosis (5,6). However, much less is known on the effect of HIV-1 infection on their cytokine and chemokine secretion profile as well as on the modulation of Fcγ receptors (FcγRs) expressed at their surface, which is a key point in the context of antibody therapy.

The discovery of potent broadly neutralizing antibodies (bNAbs) against HIV-1 has made passive immunotherapy a potential strategy for the prevention and treatment of HIV-1 infection (7). Importantly, recent clinical data have demonstrated the efficacy of several anti-HIV-1 bNAbs to control viremia when administered to people living with HIV-1 (PLWH) as well as to induce adaptive immune responses (so-called "vaccinal effect") (reviewed in 8,9). This has given strong support to the idea that bNAbs could broaden the therapeutic arsenal against HIV-1 infection (10–14). Beyond their neutralization capacity through the binding of their Fab fragment to viral antigens, the biological activity of mAbs is also mediated by the Fc moiety upon interaction with the complement system and FcγRs expressed by many cells of the immune system (15,16). This allows to clear free virions from the blood as well as to guide host immune effector cells to kill infected cells by several Fc-mediated mechanisms (i.e. antibody-dependent cellular phagocytosis (ADCP), antibody-dependent cell-mediated cytotoxicity (ADCC), …). Furthermore, mAbs can form immune complexes (ICs) *via* binding with different viral determinants (virions or infected cells) and the subsequent engagement of FcγRs by ICs has immunomodulatory effects leading to the induction of protective immunity.

We have previously identified different FcγR-expressing immune cells involved in the vaccinal

response during mAb-based therapies (8,9). By using a mouse model of retroviral infection, we

have shown a key role for ICs in enhancing antiviral immune responses by multiple innate

immune cells such as dendritic cells (DC) (17,18), neutrophils (19,20), monocytes (20) and NK

cells (21) in a FcγR-dependent manner. Notably, our findings revealed a key

immunomodulatory role for murine neutrophils in the induction of protective immunity upon

mAb-therapy (19,20). This occurs through the secretion of multiple cytokines and chemokines

upon activation by free virions and ICs. With the aim of extending our observations to the

context of HIV-1 infection, here we addressed the immunomodulatory properties of human

neutrophils upon activation by different stimuli/danger signals associated with HIV-1 infection

and mAb-based therapy.

Using neutrophils isolated from healthy donors (HD), we report here that their activation by

HIV-1 virions, free or in the form of ICs, induced the secretion of very limited amounts of

cytokines and chemokines, both in terms of diversity and quantity. However, priming of HD's

neutrophils with TNF-α or IFN-γ enhanced their chemokine production upon activation by

either free HIV-1 or ICs. Noteworthy, the inflammatory environment increased more the

chemokines secretion in IC-activated neutrophils than in HIV-stimulated neutrophils.

Furthermore, our findings reveal that neutrophils isolated from PLWH exhibited heightened

release of cytokines and chemokines compared to neutrophils from healthy HD when exposed

to danger signals associated with HIV-1 infection. We also report that PLWH's neutrophils

display higher expression several immune receptors. Our study provides new insights into the

immunomodulatory properties of neutrophils during HIV-1 infection and antibody treatment.

It also highlights that the inflammatory environment is a key factor for neutrophils to show a

functional response to HIV-1/ICs. These findings could have implications in improving the

treatment of HIV-1 infection with bNAbs alone or in combinatorial therapeutic approaches.

Results

Different cytokine/chemokine secretion profile and FcγR cell surface levels in neutrophils stimulated by TLR agonists or pro-inflammatory cytokines

First, we assessed the effect of TLR agonists and inflammatory cytokines (TNF-α and IFN-γ) on the phenotypic and functional activation of human neutrophils isolated from HD. The rationale behind this was to determine whether and how the microenvironment resulting from infection might be determinant for modulating the immune response, both in terms of chemokine/cytokine secretion profile and of FcγR expression pattern, as observed in our mouse studies (20). In this regard, virus-driven inflammation (including TNF-α and IFN-γ signature) has been associated with the development of bNAbs in spontaneous controllers of HIV (22). As for TLR agonists, we mainly focused on TLR4 and TLR7/8 agonists (lipopolysaccharide (LPS) and Resiquimod (R848), respectively), due to their role in immune activation associated with viral sensing (TLR7/8) and to the bacterial translocation observed in PLWH (i.e. LPS binding to TLR4). To assess the phenotypic and functional activation of neutrophils we used the following read-out: (i) the upregulation of the cell surface markers CD11b and CD66b, (ii) the modulation of the expression of the main FcγRs constitutively expressed on neutrophils: FcγRIIa (CD32a) and FcγRIIIb (CD16b), and (iii) the cytokine and chemokine secretion profile (**Figure 1**). Flow cytometry analysis was performed on the viable fraction of CD15$^+$/CD11b$^+$ cells (**Figure 1A**). While life span of neutrophils has been reported to be short (only few hours), specific culture conditions allow extending it. Upon TLR4 and TLR7/8 agonist stimulation during 16-20h, neutrophil survival ranged between 70-80% in average (**Figure 1B**). TLR4 and TLR7/8 agonists led to significant upregulation of CD11b and CD66b as compared to unstimulated neutrophils, with no significant differences between TLR agonists (**Figure 1C**). As for FcγR expression, neutrophil stimulation by either TLR agonist induced a strong shedding of the CD16b receptor, in particular TLR4, as well as a significant decrease in CD32a expression as compared to unstimulated neutrophils (**Figure 1C**). The assessment of neutrophil secretome revealed that both stimuli triggered the secretion of IL-8 and several chemokines (i.e. CCL3, CCL4, CXCL1, CXCL10 and CCL2) albeit to a different extent (**Figure 1D**). Due to high donor-dependent variability of chemokine secretion, no statistical differences were observed between TLR4 and TLR7/8 activation (**Figure 1E**). However, TLR4 stimulation led to the secretion of IFN-λ2/3 in 3 out of 7 samples, while TLR7/8 stimulation did not.

We also assessed the viability as well as the phenotypic and functional activation of human neutrophils stimulated with TNF-α and IFN-γ (**Figure 2**). At the concentrations used, TNF-α led to a lower neutrophil survival than IFN-γ (68% versus 85%, respectively) (**Figure 2A**). TNF-α led to a significant upregulation of CD11b and CD66b, a strong shedding of CD16b receptor as well as a significant decrease in CD32a expression while IFN-γ did not significantly affect the phenotypic activation of neutrophils (**Figure 2B**). Thus, significant differences between IFN-γ and TNF-α activation were observed for all the cell surface markers evaluated. The assessment of neutrophil's secretome revealed that TNF-α activation induced the secretion of several cytokines (IL-8, IL-1ß, IFN-λ2/3, IFN-λ1 and IFN-γ) and chemokines (CCL3, CCL4 and CXCL1) (**Figure 2C**). The secretion of IL-8, CCL3, CCL4 and CXCL1 was significantly higher than for unstimulated neutrophils, while increased secretion of IL-1ß, IFN-λ2/3 and IFN-γ was observed for half of the samples tested (**Figure 2D**). In contrast, IFN-γ activation only resulted in a very low secretion of IL-8 (**Figure 2C**). Overall, these results highlight a stimulus-specific effect on neutrophil activation, both phenotypically (FcγR and activation markers levels) and functionally (secretome). Furthermore, by studying multiple parameters, these results enabled to establish a comprehensive and comparative landscape of neutrophil activation by these stimuli (some of them used as a surrogate of viral activation) as a basis for comparison with neutrophil activation by HIV-1 and ICs. This is important to highlight as neutrophil activation by TLR agonists and cytokines has been partially addressed in previous studies (23–27), but the data are scattered and based on few analysed parameters.

Mild effect of neutrophil activation by HIV-1 virions, free or as ICs, on the expression of FcγRs

Considering that neutrophil activation by TLR agonists and TNF-α led to decreased levels of FcγRs, we next assessed the modulation of the expression of FcγRs in presence of HIV-1 virions, free or in the form of ICs. This is important to assess, as in the context of HIV-1 infection and antibody therapy, the expression of FcγRs might affect the activation of neutrophils by therapeutic antibodies. ICs were made using the anti-HIV-1 bNAb 10-1074. This bNAb is currently being tested in clinical trials and has been shown to induce vaccinal effects (28,29). The 10-1074 bNAb was chosen due to its higher affinity to the FcγRs CD16b and CD32a as compared to other bNAbs (**Supplementary Figure 1**). Activation by HIV-1 virions, ICs and free bNAb had a slight effect on neutrophil survival, which was maintained at around 85%. (**Figure 3A**). The stimulation with HIV-1 virions and ICs led to mild but significant

upregulation of CD11b and CD66b as compared to unstimulated neutrophils (**Supplementary Figure 2**) or neutrophils cultured in the presence of free bNAb, notably for CD11b (**Figure 3B**). No significant differences in the expression of these two cell surface receptors were observed between HIV-1- and IC-activated neutrophils (**Figure 3B**). Despite being similarly activated by both stimuli (as observed by the upregulation of activation markers), we observed a mild but significant reduced expression of CD16b in HIV-1-stimulated neutrophils but not in IC-stimulated cells (**Supplementary Figure 2**). However, a significant decrease in CD32a expression was observed in both HIV-1- and IC-activated neutrophils (**Supplementary Figure 2**), that was more marked in IC-activated cells (**Figure 3B**). Together, these results show a very mild effect of the activation of neutrophils by HIV-1/IC on FcγR expression. Worthy of note, the decrease in FcγRs expression was much less pronounced than that observed upon activation with the TLR7/8 agonist mimicking a viral stimulus (**Figure 1**). This is important to highlight as TLR7/8 is commonly used as surrogate of virus activation, instead of *bona fide* virions.

Neutrophils activated by HIV-1 and ICs secrete low levels of cytokines and chemokines

In parallel, we assessed the activation of neutrophils isolated from HD by HIV-1 virions, free or in the form of ICs. Free 10-1074 bNAb was used as control and showed no effect on either the expression of cell surface markers or the secretome of neutrophils. The study of the secretome of neutrophils activated by these stimuli revealed a weak secretion of several cytokines and chemokines (**Figure 3C**). Among them, IL-8, CXCL1 and CCL4 were significantly increased upon HIV-1 and IC-activation as compared to unstimulated neutrophils (**Figure 3D**). In addition, other cytokines and chemokines were detected in about half of samples tested (i.e. IL-1ß, IFN-λ1, IFN-λ2/3, CCL2 and CCL3). However, no significant differences in the secretome were observed between HIV-1- and IC-activated neutrophils (**Figure 3D**). Collectively, these results show a low activation of neutrophils by these stimuli. This contrasted to the strong secretion of multiple cytokines and chemokines by monocytes activated by HIV-1- and ICs (**Supplementary Figure 3**). This showed that these stimuli are able to induce the secretion of these soluble immune mediators by different myeloid cells but at different extents depending on the cell type.

Inflammatory conditions potentiate the activation of HD's neutrophils by HIV-1 and ICs

As the inflammatory microenvironment resulting from the viral infection and bNAbs therapy might affect the antiviral immune response, we next examined the phenotypic and functional activation of HD's neutrophils by HIV-1 free or in the form of ICs in an inflammatory

environment (i.e. priming with TNF-α and IFN-γ). Neutrophil priming with TNF-α or IFN-γ (for 10 or 30 min, respectively, **Figure 4A)** enhanced chemokine production (i.e. CXCL1, CCL2, CCL3 and CCL4) upon activation of neutrophils by either free HIV-1 or ICs. Interestingly, under these conditions, IC-activated neutrophils showed a significantly greater secretion of several chemokines than free HIV-stimulated neutrophils. Notably, TNF-α enhanced the secretion of CXCL1 in IC-stimulated neutrophils as compared to free virions stimulation (**Figure 4B**), while IFN-γ led to an enhanced secretion of CXCL1 and CCL4 (**Figure 4C**) as well as a trend to an increased secretion of CCL2. The enhanced secretion of chemokines by neutrophils upon cytokine priming indicates that the inflammatory environment (i) is key for the functional response of neutrophils to HIV-1 virions, and (ii) potentiates neutrophil chemokine secretion upon activation by ICs.

Neutrophil priming with TNF-α or IFN-γ leads to a dramatically different transcriptomic profile

To identify the potential molecular mechanisms involved in the enhancement of chemokine secretion by IC-activated neutrophils upon pro-inflammatory cytokine priming, we next studied the modulation of gene expression in neutrophils briefly conditioned by TNF-α or IFN-γ using the NanoString technology. This transcriptomic analysis allowed us to identify differentially expressed genes upon TNF-α or IFN-γ priming (**Figure 5**). Heatmap representation of upregulated and downregulated genes (**Figure 5A)** shows that IFN-γ and TNF-α activation resulted in changes in the expression pattern of a different set of genes. Volcano plots provided in **Figure 5B** show the differential gene expression by the stimulation by either cytokine. Only three genes (CD69, PTX3 and IL-1β) were upregulated by both cytokines (**Figure 5C**). IFN-γ induced a set of genes related to innate immune responses, viral infections, cytokine signaling, regulation of lymphocyte activation, antigen processing/presentation, MAPK cascade and PD1/PD-L1 axis. In contrast, TNF-α induced a set of genes related to IL-10/IL-18 signaling, AP1/NF-κB signaling, regulation of cell proliferation and apoptosis responses as well as nitric oxide biosynthesis (**Figure 5D**). The list of genes upregulated or downregulated and their fold increase is shown in **Supplementary Tables 1 and 2**. Worthy of note, IFN-γ led to the upregulation of the inducible FcγRI (CD64) (confirmed at the protein level, **Supplementary Figure 4**), while no effect of TNF-α activation on FcγR expression was observed. In addition, IFN-γ induced the expression of the TLR1 and TLR8 (involved in bacterial lipoproteins and viral sensing, respectively), in contrast to TNF-α that induced the expression of TLR5 (involved

in flagellin sensing) (**Figure 5A**). These results show a very distinct transcriptomic regulation of neutrophils by TNF-α and IFN-γ, with IFN-γ priming leading to the upregulation of FcγRI and TLR8 genes that could potentiate Fc-mediated antibody signaling and viral sensing, respectively.

Neutrophils from PLWH exhibit increased release of specific cytokines and chemokines upon stimulation with danger signals associated with HIV-1 infection

Neutrophil functions have been shown to be compromised upon HIV-1 infection. Notably, it has been reported that HIV-1 infection impaired the effector functions of neutrophils as well as increased their apoptosis (5,6). However, the effect of HIV-1 infection on their cytokine and chemokine secretion profiles is much less studied and mostly assessed in the AIDS stage (30). Thus, it is necessary to investigate how HIV-1 infection and ART treatment affect the functional activation of neutrophils, particularly their immunomodulatory potential. To this end, we assessed the activation of neutrophils isolated from the blood of ART-treated PLWH (**Supplementary 5A-D**) activated by TLR agonists, pro-inflammatory cytokines, HIV-1 or ICs as compared to HD. Importantly, no significant differences in the frequency of neutrophils in the blood of PLWH were observed as compared to HD (**Supplementary Figure 5E**). Furthermore, no significant differences in neutrophil frequencies were neither observed in PLWH regardless of the time since diagnosis (**Supplementary Figure 5F**). Neutrophils from PLWH had similar secretion profiles than HD upon activation with either TLR4 or TLR7/8, both qualitatively (type of cytokine/chemokine) and quantitatively (amount secreted) (**Figure 6A**), with several noticeable exceptions. In particular, neutrophils from PLWH showed an enhanced secretion of CCL3 and CXCL1 upon TLR4 stimulation and CCL3 and IFN-γ upon TLR7/8 activation (**Figure 6B**). The secretion of other chemokines (CCL4) and cytokines (IFN-λ2/3, IL-1ß) showed a trend of increase in neutrophils isolated from PLWH, but the difference did not reach statistical significance (**Figure 6B**). We assessed in parallel the response of PLWH's neutrophils to inflammatory cytokines (TNF-α and IFN-γ) (**Figure 6C**). Few significant differences were observed in the secretome of neutrophils from PLWH versus HD upon activation by either cytokine. Upon TNF-α activation we observed an increased secretion of CCL2, and on the contrary, a decreased secretion of IFN-λs and TNF-α (**Figure 6D**). Neutrophils from PLWH also had similar secretion profiles than those from HD upon stimulation with free HIV-1 virions or in the form of ICs (bNAb alone, was used as a control) (**Figure 7**). Overall, these results show that neutrophils from PLWH under ART not only

maintained their capacity to secrete cytokines and chemokines in response to danger signals associated with HIV-1 infection, but also showed increased release of specific soluble mediators (i.e. CXCL1, CCL2, CCL3 and IFN-γ) compared to neutrophils isolated from healthy donors (HD), particularly under certain stimulatory conditions and/or in specific donor samples.

Neutrophils from PLWH show stronger CD16 shedding in response to danger signals than neutrophils from HD

We analyzed in parallel the expression of FcγRs in neutrophils from PLWH upon activation by different danger signals associated with HIV-1 infection, as this is an important issue in the context of bNAb-based therapy. Upon TLR4 activation, CD16b and CD32a cell surface levels on neutrophils from PLWH were significantly lower than those observed in HD (**Figure 8A**). In contrast, TLR7/8 activation resulted in no significant difference in the expression of either FcγR as compared to HD (**Figure 8A**). Similar to TLR4 activation, TNF-α stimulation led to a significant lower expression of CD16b and CD32a on neutrophils from PLWH, as compared to HD (**Figure 8B**). In contrast, IFN-γ activation only led to significantly lower levels of CD16b (**Figure 8B**). No difference in expression was observed for CD16b or CD32a upon activation by free HIV-1 virions (**Figure 8C**) as with TLR7/8 activation. These results highlight that CD16b shedding can be affected by HIV-1 infection as neutrophils from PLWH showed a stronger CD16 shedding as compared to HD upon activation by TLR4 and inflammatory cytokines. However, such enhanced CD16 shedding was not observed upon activation by viral stimuli (i.e. TLR7/8, HIV-1, free or in the form of ICs).

Neutrophils from PLWH display enhanced expression of FcγRs, MHC-II molecules, CXCR4, and PD-L1

We then performed a phenotypic study of neutrophils isolated from PLWH and HD to further assess to what extent HIV-1 infection affected neutrophil properties. To this end, we measured the expression of 18 cell surface markers in the absence of stimulation. The panel was chosen to follow the expression of markers of interest in HIV-1 infection (CXCR4), neutrophil activation (CD11b, CD66b, CD63, CD62L), neutrophil maturity (CD10), neutrophil immunosuppressive properties (PD-L1) and Fc receptors (CD16, CD32, CD64 and CD89), among others. Within the 18 markers, we found a significantly higher expression on PLWH's neutrophils of two activating FcγRs (CD16b and CD64) and 3 other cell surface molecules, namely CXCR4, PD-L1 and HLA-DR. We also detected a decreased expression of the adhesion

318   molecule CD172α in neutrophils from PLWH (**Figure 9**). Thus, despite similar secretion

319   profiles when activated by HIV-1 virions and/or ICs, neutrophils from PLWH show phenotypic

320   differences as compared to HD. These differences might reflect different functional properties.

Discussion

We have previously shown in a mouse model of retroviral infection that neutrophils have a key immunomodulatory role in the induction of a protective immunity (19,20). Here, we provide new insights into the immunomodulatory properties of neutrophils in the context of HIV-1 infection and antiviral mAb-therapy. Our study dissects the secretome and expression of neutrophil immune receptors in a stimulus-specific manner both in HD and PLWH. Noteworthy, most studies on immunomodulatory functions of neutrophils are based on bacterial activation (i.e. TLR4 activation), while the effect of virus-dependent activation is understudied. The latter has been mostly investigated using TLRs as surrogates, and there are few data on whether and how HIV-1 and/or HIV-ICs affect their immunomodulatory properties. Our work highlights that neutrophil from ART-treated PLWH retain the ability to secrete cytokines/chemokines in response to danger signals and even display a stronger secretion of some of these soluble mediators compared to neutrophils isolated from HD. In addition, notable differences in phenotypic features distinguish them from neutrophils of HD origin. Our research also shows that inflammatory environment shapes neutrophils immunomodulatory properties and allows them to better respond to HIV-1 viral determinants, free or in the form of ICs.

Based on the observed upregulation of CD11b and CD66b cell surface activation markers, our data show that both TLR4 and TLR7/8 agonists similarly activate neutrophils. However, TLR4 agonist led to a significantly decrease expression of two FcγRs (CD16b and CD32a) as compared to the TLR7/8 agonist. This suggests a stronger effect of bacterial-associated danger signals on FcγRs modulation than virus-associated danger signals, notably on the shedding of CD16b. Similarly, our data also show that free HIV-1 and ICs had either a mild effect or no effect, respectively, on the modulation of CD16b cell surface levels, despite that both stimuli significantly increased the expression of CD11b and CD66b. However, TNF-α, but not IFN-γ, led to a strong CD16b shedding as well as a decrease in the CD32a cell surface levels (although the latter to a much lesser extent). This shows a clear effect of inflammatory conditions in promoting CD16b shedding that might have biological consequences. On one hand, soluble CD16b might have immunomodulatory functions as it has been shown to play a dominant role in clearing ICs (31) and to bind complement receptors (32) leading to an increased cell activation. On the other hand, it has been suggested that CD16 shedding might grant access to cell-activating CD32a (33). In addition, CD16b shedding can be affected by pathological conditions as observed for neutrophils from PLWH that showed stronger CD16 shedding upon activation than HD's neutrophils. This stimulus-dependent and pathology-dependent

modulation of FcγRs might have to be taken into account in the context of HIV-1 infection and

bNAb-based immunotherapies alone or in combinatorial therapeutic approaches involving TLR

agonists (14,34–38) as it might affect their therapeutic efficacy. This might also be relevant to

consider it in the context of bacterial co-infections in PLWH (39).

Our work also allowed us to dissect, at the protein level, the chemokine/cytokine secretion

profiles of neutrophils in a stimulus-dependent manner under both physiological (HD) and

pathological (PLWH) conditions. This is important to point out as multiple studies assessing

the secretion profile of neutrophils upon different stimuli were based on transcriptomic

approaches instead of *bona fide* cytokine/chemokine secretion. In agreement with previous

studies, we have shown that neutrophils isolated from HD and activated by LPS, R848 or TNF-

α secreted IL-8, CXCL1, CCL3 and CCL4 (23–25,27), while IFN-γ only induced the secretion

of IL-8. We also observed the secretion of IFN-λ2/3 by LPS-activated neutrophils in half of the

samples tested, which has not been described thus far. Finally, our work finely describes the

secretome of HIV-1-and IC-activated neutrophils, thus far unreported. Importantly, the

cytokines/chemokines secreted upon activation with HIV-1 and ICs are involved in neutrophil

activation (IL-8) and recruitment (CXCL1, CCL3) and/or act as chemoattractant for NK cells,

monocytes and various other immune cells in the site of inflamed or damaged tissue (CCL4)

(40). Interestingly, although it is only detected in a proportion of samples, the secretion of IFN-

λ by neutrophils observed upon HIV-1 stimulation has not been described to date. This family

of cytokines have been shown to modulate antiviral immune response (41), but its role in HIV-

1 infection is still unclear. Further investigations are necessary to ascertain the secretion of IFN-

λ by neutrophils activated by the different stimuli and to determine the biological consequences

of its production in the context on neutrophil biology and HIV-1 infection. In this regard, it has

been shown that IFN-λ (i) diminishes the production of reactive oxygen species and

degranulation in neutrophils (42) and (ii) IFN-λ3 inhibits HIV-1 infection of macrophages (43).

In the context of HIV-1 infection, a key question not yet fully addressed is the capacity of

neutrophils to respond, in terms of cytokines/chemokines secretion, to pathogen-associated

danger signals and cytokine stimulation. Our data show that neutrophils isolated from ART-

treated PLWH respond to TLR agonists, pro-inflammatory cytokines, HIV-1 and ICs in a

similar way than neutrophils from HD, if not better. This is the case for the enhanced secretion

of CCL2 upon TNFα activation as well as CCL3/CXCL1 and IFN-γ upon TLR activation.

These observations indicate that (i) neutrophils retain their capacity to secrete
cytokines/chemokines in the immune context associated with HIV-1 infection, and (ii) such
immune context may even potentiate the secretion of some specific chemokines. Our results
contrast with previous studies reporting impaired cytokine secretion by LPS-activated
neutrophils isolated from PLWH on ART for less than 2 years and having developed AIDS
(30). Thus, in advanced stages of disease, the ability of neutrophils to respond to danger signals
is apparently altered, which can contribute to the disease evolution. This does not seem to be
the case for neutrophils isolated from aviremic, ART-treated PLWH as shown here,
highlighting the importance of ART in maintaining not only their effector functions (6,44), but
also their cytokine/chemokine secretion capacity.

Our work also sheds new light on the phenotypic characteristics of neutrophils in PLWH. We
found that several cell surface molecules were significantly increased in PLWH. First, we
observed an increased expression of CD16b and CD64, while CD32a levels were not affected.
However, the upregulation of these FcγRs in PLWH did not result in an increased secretion of
cytokines and chemokines upon IC-activation of neutrophils as compared to HD. Indeed, while
increased FcγR-mediated phagocytic activity may directly influence immune activation, the
effect of specific FcγRs in modulating cytokines/chemokines secretion in neutrophils is still ill-
understood. Further investigations are required to determine the potential FcγR-dependent
enhancement of cytokines/chemokine by ICs as well as the FcγRs involved. Second, our results
show the upregulation of HLA-DR in PLWH neutrophils. This observation is also in agreement
with the increased frequency of HLA-DR[high] neutrophils in SIV-infected macaques as described
by Lemaitre et al (45). On the contrary, Bowers *et al* (46) reported that the expression of HLA-
DR was unmodified in PLWH neutrophils. However, experiments were performed using whole
blood and not purified neutrophils as we did. These controversial observations could therefore
result from the presence of several neutrophil subsets in whole blood as previously described
(47–49). Third, our results also show an enhanced expression of CXCR4 and PD-L1.
Interestingly, the upregulation of these two cell surface molecules has recently been associated
with neutrophil aging in PLWH (50). However, in contrast to our study, the assessment of
CXCR4 and PD-L1 expression was made using samples from ART-naive PLWH. Finally, our
findings provide further evidence of the upregulation of PD-L1 in PLWH neutrophils. PD-L1
expression has been previously described in different populations of neutrophils (46,47,50) and
irrespective of ART (46,50) in PLWH. It has been associated with T-cell immunosuppressive

effects that can be counteracted by blocking the PD-1/PD-L1 axis (46,50). Thus, the
upregulation of PD-L1 we found in PLWH suggests a potential immunosuppressive role of
PLWH neutrophils.
424
Another important observation of our study is the effect of the inflammatory environment on
the modulation of neutrophil properties. Thus, the stimulation with inflammatory cytokines
potentiated the release of several chemokines by HIV-1 and IC-activated neutrophils.
Interestingly, TNF-α and IFN-γ priming led to a significant increased secretion of CXCL1 and
CCL4 by IC-activated neutrophils as compared to virus-activated cells. This is in agreement
with our previous results using mouse neutrophils activated with IC-made with a murine
retrovirus (20). Our work also allowed us to identify a different effect of TNF-α and IFN-γ on
the enhancement of the functional properties of IC-activated human neutrophils. Notably, IFN-
γ but not TNF-α, led to enhanced secretion of CCL4 and CCL2 although to a lesser extent for
the latter. It is worth mentioning that the enhanced secretion of CCL2 by HIV-1 and HIV-IC-
activated neutrophils by IFN-γ priming is in agreement with the IFN-γ-mediated upregulation
of CCL2 production previously described in a different experimental setting (51). This finding
suggests that in a Th1 microenvironment, neutrophil might be involved in the recruitment of
monocytes through the secretion of CCL2. Worthy of note, IFN-γ priming led to the
upregulation of FcγRI and TLR8 (potentially involved in IC-mediated signaling and HIV-1
sensing, respectively). Interestingly, this observation is in agreement with the recent study by
Kumola *et al*, showing an upregulation of CD64 and TLR8 upon IFN-γ stimulation as well as
an enhanced expression of PD-L1 (52). This suggests that the enhanced expression of CD64
and PD-L1 we observed on the neutrophils from PLWH could be due to IFN-γ-priming *in vivo*.
However, our results showing that neutrophils from PLWH respond to HIV-1 or ICs as weakly
as those from HD, also raise the possibility that they may not be adequately primed *in vivo*, or
not to an efficient degree. Thus, the question of whether PLWH's neutrophils are primed or not
remains open and deserves further investigation.
448
In conclusion, our work dissects the phenotypic and functional properties of PLWH neutrophils.
It highlights that they maintain their cytokine and chemokine secretion capacity, while
displaying some immunosuppressive features. It also provides new insights on the modulation
of FcγR expression in the context of danger signals associated with HIV-1 infection which need
to be taken into account in the context of antibody therapy. These findings might be exploited

to improve HIV-1 antiviral therapies in PLWH. This could be achieved by tailoring therapeutic

interventions aiming at harnessing the immunomodulatory properties of these cells. Strategies

could involve leveraging the immunomodulatory properties of neutrophils (through the use of

cell-type specific, immunostimulatory compounds) and/or counteracting their

immunosuppressive functions (through the design of combinatorial approaches involving the

blocking of the PD-1/PD-L1 axis) (53,54).

Material & Methods

Ethics

All human blood samples were fully anonymized and given with previous written consent of the donor. Samples were given under ethical approval of appropriate institutions.

Human blood samples

Blood samples from anonymized healthy donors (HD) were provided by the French establishment for blood donations (Etablissement Français du Sang, EFS, Montpellier, France) according to the agreement between this establishment and INSERM (21PLER2018-0069). EFS provided 7 to 8mL of fresh whole blood in EDTA for each healthy donor.

The use of blood samples from anonymized people living with HIV-1 (PLWH) under antiretroviral therapies (ART) was approved by the Centre for Biological Resources (DRI#305-2019-NC and amendment DRI_2022-69-SR). The samples were collected and provided by the Virology laboratory of the Centre Hospitalier Universitaire (CHU) Lapeyronie of Montpellier (France). Median age was 57 years old ([22;76]) for 31 PLWH donors (19M; 13F). All donors were under successful ART with no detectable viral load. 2 to 3mL of whole blood in EDTA were provided for each PLWH donors.

Primary cells purification

Peripheral blood mononuclear cells (PBMC) and granulocytes were isolated using a Histopaque-1119 density gradient. Briefly, whole blood sample was diluted to three times its volume with a solution of PBS 1X (Gibco, cat #10010-015) supplemented with EDTA at 0,2 mM (Invitrogen, cat #15575-038) and loaded on Histopaque-1119 (Sigma-Aldrich, cat #11191). Density gradient was performed by 800g centrifugation with slow acceleration and no brake (acceleration 4, brake 0) for 20 minutes. Then, PBMCs ring was collected as well as the granulocytes- containing fraction underneath.

Monocytes and neutrophils were then purified from PBMC or granulocytes, respectively, using magnetic-based CD14- or CD15-positive selection kits (STEMCELL TECHNOLOGIES, CD15+ positive selection REF#18651, CD14+ positive selection REF#117858) according to the manufacturer's protocol. Purity assessed by flow cytometry was > 98% for both cell type.

Cell culture

Purified neutrophils were plated at a concentration of 2 million cells/mL in 96 U-bottom wells plate and cultured in RPMI 1640 supplemented with GlutaMAX^{TM} medium (Gibco, cat #61870-010), 10% fetal bovine serum (FBS, Eurobio, REF#CVFSVF00-01) and 1% penicillin/streptomycin (Gibco, cat #15140-122) with 10 μg/mL of G-CSF (R&D, cat #214-CS-025).

Purified monocytes were plated at a concentration of 1 million cells/mL in 96 U-bottom wells and cultured in RPMI 1640 medium supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin with 10 μg/mL of GM-CSF (Peprotech, cat #300-03). Cells were then cultured for 16-20h with or without stimulation.

In vitro cell stimulation

For TLR or cytokine stimulations, cells were stimulated overnight with TLR4 agonist (LPS ; Sigma-Aldrich, cat #L4524), TLR7/8 agonist (R848 ; Invivogen, cat #tlrl-r848) at 0,5 μg/mL each, with IFN-γ (R&D, cat #285-IF-100) at 1000 IU/mL or with TNF-α (Peprotech, cat #AF-300-01A) at 1ng/mL. Otherwise, cells were stimulated with either free HIV-1 virions (Ad8 strain, MOI of 3.10^4), immune complexes made with HIV-1 Ad8 and the 10-1074 bNAb or the bNAb alone.

Neutrophils priming

For primed conditions, neutrophils were resuspended in culture medium after purification and stimulated at 37°C 5%CO$_2$ with IFN-γ at 2000 IU/mL for 30 minutes or with TNF-α at 5ng/mL for 10 minutes. Cells were washed then cultured overnight with either free virus, immune complex or bNAb alone.

Virus production, purification and titration:

2.5 million human embryonic kidney cells (293THEK cell) were seeded in 10 ml of growth media 1 day before transfection. At 50-70% confluence, cells were transfected with calcium phosphate precipitate method, with 10 μg total quantity of HIV-1 molecular clone expressing plasmids, as in (55). At 12 hours post transfection, cells were washed and fresh media supplemented with 20mM Hepes was added on the cells. Cell culture supernatant containing viral particles was collected 36 hours post-transfection, pooled and filtrated through 0.45 μm and then purified by ultracentrifugation on a cushion of 25% sucrose in TNE buffer (10 mM Tris-HCl [pH 7.4], 100 mM NaCl, 1 mM EDTA) at 100000xg, for 1 hour 30 minutes at 4°C,

in a SW32Ti Beckman Coulter rotor. Dry pellet was resuspended with RPMI without serum, at
4°C overnight. Viral titer was quantified by alphaLISA immunoassay (Perkin Elmer) using
anti-HIV-1 p24. Recombinant HIV-1 p24 protein is used for titer standard range. Viral particles
were quantified using Videodrop system (Myriade France).

Recombinant HIV-1 *bNAb production*
Recombinant human HIV-1 IgG1 antibodies, anti-CD4bs 3BNC117 (56) and anti-V3-glycan
10-1074 (57), were produced by co-transfection of Freestyle 293-F cells (Thermo Fisher
Scientific) using the polyethyleneimine (PEI)-based precipitation method as previously
described (58), and purified by affinity chromatography using protein G Sepharose 4 fast flow
beads according to the manufacturer's instructions (Cytiva). Purified IgG antibodies were
dialyzed against PBS using Slide-A-Lyzer Cassettes (10K MWCO; Thermo Fisher Scientific),
and quantified using NanoDrop™ ONE instrument (Thermo Fisher Scientific). Both bNAbs
are of IgG1 isotype and were used at a concentration of 1μg/mL to form ICs.

Immune complexes formation
Viruses (3.10^4 viral particles) and bNAb (1μg/mL) were diluted in culture medium and mixed
together on a 1,5mL Eppendorf, then mixed on a rotative agitator for 20 minutes. ICs were
gently plated with cells in 96 U-bottom wells plate.

Flow cytometry immunophenotyping of neutrophils
Monitoring of phenotypic activation was done by flow cytometry using FACSCantoII, BD
LSRFortessa (BD Bioscience) or Novocyte (Agilent Technologies) flow cytometers. Cells were
stained at room temperature with a viability dye (Zombie NIR™, Biolegend #423105 or
ZombieRed™, Biolegend #423109) for 15 minutes then washed and stained for 20 minutes at
room temperature with fluorochrome-conjugated antibodies. Antibodies for flow cytometry
were PerCPCy5-5-CD15 (HI98, Biolegend #301922), AF488-CD11b (ICRF44, Biolegend
#301317), AF647-CD66b (G10F5, Biolegend #305110), PC7-CD16a/b (3G8, Biolegend
#302016), PE-CD32a/b (FUN-2, Biolegend #303206), PE-CD64 (10.1, Biolegend #305008),
FITC-CD14 (MoP9, STEMCELL Technologies #60124FI).
For immunophenotyping of HD's and PLWH's neutrophils the following antibodies were used:
PerCPCy5-5-CD15 (HI98, Biolegend #301922), PE-LOX-1 (15C4, Biolegend #358604),
BV510-CD11b (ICRF44, Biolegend, #301334), AF647-CD66b (G10F5, Biolegend, #305110),
BV605-CD62L (DREG-52, Biolegend, #304834), BV650-HLA-DR (L243, Biolegend,

#307650), BV711-CD63 (H5C6, Biolegend, #353042), PC7-CD16a/b (3G8, Biolegend,

#302016), PE-CF594-CD64 (10.1, Biolegend, #565389), APC-Fire750-CD32a/b (FUN-2,

Biolegend, #303220), PE-PD-1 (3.13, Miltenyi, #130-177-384), APC-CD177 (MEM-166,

Biolegend, #315808), BV605-CD49d (9F10, Biolegend, #304324), BV711-CD31 (WM59,

Biolegend, #303136), PC7-PD-L1 (MIH1, BD, #558017), PE-CF594-CD10 (HI10a,

Biolegend, #312227), PE-CD89 (Biolegend, #555686), APC- CD172α (15-414, eBiosciences

#17-1729-42), PC7-CXCR4 (12G5, Biolegend, #306514).

Forward scatter area and forward scatter height as well as side scatter area and height were used

to remove doublets from flow cytometry analyses. Gating is done on live cells as assessed by

viability dye and on CD15-positive cells. Data were analyzed using FlowJo software version

10.5.3 (TreeStar).

Cytokines and chemokines secretion quantification

Secretion of soluble cytokines and chemokines were assayed from cell-free culture supernatants

of *in vitro* stimulated neutrophils and monocytes using bead-based immunoassays (LegendPlex,

Biolegend). Two assays were used allowing the quantification of IL-1β, Il-6, TNF-α, CXCL10,

IFN-λ1, IL-8, IL-12p70, IFN-α2, IFN-λ2/3, GM-CSF, IFN-β, IL-10, IFN-γ (LegendPlex

Human anti-virus response panel, Biolegend #740390) and IL-8, CXCL10, CCL11, CCL17,

CCL2, CCL5, CCL3, CXCL9, CXCL5, CCL20, CXCL1, CXCL11, CCL4 (LegendPlex

Human proinflammatory chemokines panel, Biolegend #740985). Immunoassays plates were

read on a Novocyte (Agilent Technologies) flow cytometer and data were analyzed with the

appropriate Biolegend LEGENDplex^{TM} software (https://legendplex.qognit.com).

Transcriptomic studies

RNA was extracted from neutrophils obtained from 4 healthy blood donors. Total RNA was

then purified from 500,000 to 1 million cells using the miRNeasy Micro kit (Qiagen GmbH)

following the manufacturer's instructions. The quantification of all RNA samples was

performed using the Nanodrop One (Thermo Fisher Scientific), and RNA integrity was

evaluated using the Agilent 2100 Bioanalyzer system. RNA expression profiling was performed

using Myeloid Innate Immunity Panel 2.0 (770 genes in 19 different pathways). Gene

expression analysis was conducted with the NanoString technology. Briefly, 5 μL/sample

containing of 50 ng of total RNA was combined with the nCounter® reporter CodeSet (3 μL)

and nCounter® capture ProbeSet (2 μL) along with hybridization buffer (5 μL) for an overnight

hybridization reaction at 65 °C. The reaction was then cooled to 4 °C, and the samples were

purified, immobilized on a cartridge, and the data were assessed using the nCounter SPRINT.
All expression data was reviewed using NanoString® nSolver™ Analysis software 4.0. To
identify main biological processes modulated by cytokine priming, genes upregulated and
downregulated were analyzed using Metascape (59).

Surface Plasmon resonance (SPR)

Affinity of bNAbs for Fcγ Receptors was assessed by SPR experiments performed on a Biacore
T200 (GE Healthcare). SPR experiments were performed on a T200 apparatus at 25°C in PBS
containing 0,05 % P20 surfactant (Cytiva). Anti-histidine antibody (R&D Systems) was
covalently immobilized on a CM5-S sensor chip flowcell (Fc2) by amine coupling according
to the manufacturer's instructions (Cytiva). A control reference surface (flowcell Fc1) was
prepared using the same chemical treatment but without anti-His antibody. All kinetic
measurements in Fc1 and Fc2 were performed by single-cycle titration at 100µl/min. Each
human FcγR (R&D Systems) was captured on immobilized anti-His antibodies at 100-200 RU
level. Five increasing concentrations (3,6, 11, 33, 100, 300 nM) of antibody were injected
(injection time = 60s) at 100µl/min on captured receptors. After a dissociation step of 600s in
running buffer, sensor surfaces were regenerated using 10µl of glycine-HCl pH1.5. All the
sensorgrams were corrected by subtracting the low signal from the control reference surface
and buffer blank injections. Kinetic parameters were evaluated from the sensorgrams using a
two-states or a steady-state models from the T200 evaluation software.

Statistics

Statistical analyses were performed using Prism software version 9.5.1 (GraphPad). Simple
group comparisons were performed using the Wilcoxon signed rank test for paired data or the
Mann–Whitney test for unpaired data. Multiple group comparisons were performed using
Friedmann test for paired data or using Kruskal-Wallis test for unpaired data with additional
multiple comparisons Dunn test. Significance was assigned as follows: $*p < 0.05$, $**p < 0.01$,
$***p < 0.001$, and $****p < 0.0001$.

Acknowledgements

This work was supported by grants from Sidaction (A014-2-AEQ-08-01; BI25-1-02278, 20-1-AEQ-12652), ANRS (France Recherche Nord&sud Sida-hiv Hépatites; ECTZ46143; ECTZ47079, ECTZ205786), and INSERM state funding granted to M. Naranjo-Gomez (U1183NAR). This work was also supported by the consortium ACT4COVIDCellnex (220110FF, RAK200019FFA). R.Dibsy was a recipient of a Sidaction PhD fellowship. S. Marsile-Medun, M. Naranjo-Gomez, Manon Souchard, Daouda Abba Moussa, Martine Pugnière and M. Pelegrin are members of the "MabImprove Labex", a public grant overseen by the French National Research Agency (ANR) as part of the "Investments for the future" program (reference: ANR-10-LABX -53-01) that also supported this work. We thank the imaging facility MRI, which is part of the UAR BioCampus Montpellier and a member of the national infrastructure France-BioImaging, supported by the French National Research Agency (ANR-10-INBS-04, "Investments for the future"). We thank the BSL3 research laboratory CEMIPAI (CNRS UAR3725) for providing help and advice, and the Centre de Ressources Biologiques (CRB) for providing blood samples from PLWH. We thank the Transcriptomic Core Facility of the Institute in Regenerative Medicine and Biotherapy, CHU-INSERM-UM Montpellier where the applied Biosystem-Affymetrix microarrays were processed. We thank Cecile Monzo and Carole Crozet that are in charge of the Nanostring Sprint Profiler platform in the Institute for Neurosciences Montpellier (INM), for providing assistance in the transcriptomic studies. We thank the Montpellier Proteomics Platform (PPM-PP2I, BioCampus Montpellier) facilities where the SPR experiments were carried out. We are grateful to Dr. Paris for critical reading of the manuscript.

Authors contribution

Soledad Marsile-Medun (SMM), Mar Naranjo-Gomez (MN-G) and Mireia Pelegrin (MP) defined the research program; SMM, Manon Souchard (MS), Daouda Abba Mousssa (DAM), Giang Ngo (GN), Martine Pugnière (MPu,) and MN-G, performed the experiments; Valerie Lorin (VL) and Hugo Mouquet (HM) provided advice on and generated bNAbs; Elisa Reynaud (ER) and Edouard Tuaillon (ET) managed and provided PLWH blood samples. Rayane Dibsy (RD) and Delphine Muriaux DM) provided advice on and performed HIV-1 production, purification and quantification. SMM, MS, MN-G and MP carried out the data analyses. SMM, MN-G and MP wrote the manuscript. Grants to SMM, MN-G and MP funded the study. All authors discussed, commented on and approved the manuscript in its final form.

659 **Conflict of interests**

660 None

REFERENCES

1. Gabriel C, Her Z, Ng LFP. Neutrophils: Neglected Players in Viral Diseases. DNA and Cell Biology. 2013;32(12):665-75.

2. Mantovani A, Cassatella MA, Costantini C, Jaillon S. Neutrophils in the activation and regulation of innate and adaptive immunity. Nat Rev Immunol. 2011;11:519-31.

3. Leliefeld PHC, Koenderman L, Pillay J. How Neutrophils Shape Adaptive Immune Responses. Front Immunol. 2015;6:471.

4. Ley K, Hoffman HM, Kubes P, Cassatella MA, Zychlinsky A, Hedrick CC, et al. Neutrophils: New insights and open questions. Sci Immunol. 2018 ;3(30):eaat4579.

5. Hensley-McBain T, Klatt NR. The Dual Role of Neutrophils in HIV Infection. Curr HIV/AIDS Rep. 2018;15(1):1-10.

6. Casulli S, Elbim C. Interactions between human immunodeficiency virus type 1 and polymorphonuclear neutrophils. J Innate Immun. 2014;6:13-20.

7. Frattari GS, Caskey M, Søgaard OS. Broadly neutralizing antibodies for HIV treatment and cure approaches. Curr Opin in HIV and AIDS. 2023;18(4):157-63.

8. Pelegrin M, Naranjo-Gomez M, Piechaczyk M. Antiviral Monoclonal Antibodies: Can They Be More Than Simple Neutralizing Agents? Trends Microbiol. 2015;23(10):653-65.

9. Pelegrin M, Marsile-Medun S, Abba-Moussa D, Souchard M, Naranjo-Gomez M. Fc-Dependent Immunomodulation Induced by Antiviral Therapeutic Antibodies: New Perspectives for Eliciting Protective Immune Responses. Antibodies (Basel). 2022;11(3):50.

10. Julg B, Barouch D. Broadly neutralizing antibodies for HIV-1 prevention and therapy. Semin Immunol. 2021;51:101475.

11. Caskey M, Klein F, Nussenzweig MC. Broadly neutralizing anti-HIV-1 monoclonal antibodies in the clinic. Nat Medicine. 2019;25(4):547-53.

12. Spencer DA, Shapiro MB, Haigwood NL, Hessell AJ. Advancing HIV Broadly Neutralizing Antibodies: From Discovery to the Clinic. Front Public Health. 2021;9:690017.

13. Walsh SR, Seaman MS. Broadly Neutralizing Antibodies for HIV-1 Prevention. Front Immunol. 2021;12:712122.

14. Mahomed S, Garrett N, Baxter C, Abdool Karim Q, Abdool Karim SS. Clinical Trials of Broadly Neutralizing Monoclonal Antibodies for Human Immunodeficiency Virus Prevention: A Review. J Infect Dis. 2021;223(3):370-80.

15. Bruhns P, Jönsson F. Mouse and human FcR effector functions. Immunol Rev. 2015;268(1):25-51.

16. Gogesch P, Dudek S, van Zandbergen G, Waibler Z, Anzaghe M. The Role of Fc Receptors

on the Effectiveness of Therapeutic Monoclonal Antibodies. Int J Mol Sci. 2021;22(16):8947.

17. Lambour J, Naranjo-Gomez M, Piechaczyk M, Pelegrin M. Converting monoclonal

antibody-based immunotherapies from passive to active: bringing immune complexes into play.

Emerg Microbes Infect. 2016;5(8):e92.

18. Michaud HA, Gomard T, Gros L, Thiolon K, Nasser R, Jacquet C, et al. A crucial role for

infected-cell/antibody immune complexes in the enhancement of endogenous antiviral

immunity by short passive immunotherapy. PLoS Pathog. 2010;6:e1000948.

19. Naranjo-Gomez M, Lambour J, Piechaczyk M, Pelegrin M. Neutrophils are essential for

induction of vaccine-like effects by antiviral monoclonal antibody immunotherapies. JCI

Insight. 2018;3(9).

20. Lambour J, Naranjo-Gomez M, Boyer-Clavel M, Pelegrin M. Differential and sequential

immunomodulatory role of neutrophils and Ly6Chi inflammatory monocytes during antiviral

antibody therapy. Emerg Microbes Infect. 2021;10(1):964-81.

21. Naranjo-Gomez M, Cahen M, Lambour J, Boyer-Clavel M, Pelegrin M.

Immunomodulatory Role of NK Cells during Antiviral Antibody Therapy. Vaccines.

2021;9(2):137.

22. Dugast AS, Arnold K, Lofano G, Moore S, Hoffner M, Simek M, et al. Virus-driven

Inflammation Is Associated With the Development of bNAbs in Spontaneous Controllers of

HIV. Clin Infect Dis. 2017;64(8):1098-104.

23. Tamassia N, Le Moigne V, Calzetti F, Donini M, Gasperini S, Ear T, et al. The MyD88-

independent pathway is not mobilized in human neutrophils stimulated via TLR4. J Immunol.

2007;178(11):7344-56.

24. Cassatella MA. Neutrophil-derived proteins: selling cytokines by the pound. Adv Immunol.

1999;73:369-509.

25. Kasama T, Strieter RM, Standiford TJ, Burdick MD, Kunkel SL. Expression and regulation

of human neutrophil-derived macrophage inflammatory protein 1 alpha. J Exp Med.

1993;178(1):63-72.

26. Yeaman GR, Collins JE, Currie JK, Guyre PM, Wira CR, Fanger MW. IFN-gamma is

produced by polymorphonuclear neutrophils in human uterine endometrium and by cultured

peripheral blood polymorphonuclear neutrophils. J Immunol. 1998;160(10):5145-53.

27. Tecchio C, Cassatella MA. Neutrophil-derived chemokines on the road to immunity. Semin

Immunol. 2016;28(2):119-28.

28. Niessl J, Baxter AE, Mendoza P, Jankovic M, Cohen YZ, Butler AL, et al. Combination

anti-HIV-1 antibody therapy is associated with increased virus-specific T cell immunity. Nat

Med. 2020;26(2):222-7.

29. Naranjo-Gomez M, Pelegrin M. Vaccinal effect of HIV-1 antibody therapy: dream or

reality? Curr Opin in HIV and AIDS. 2023; 18(4):209-216.

30. Gasperini S, Zambello R, Agostini C, Trentin L, Tassinari C, Cadrobbi P, et al. Impaired

cytokine production by neutrophils isolated from patients with AIDS. AIDS. 1998;12(4):373-9.

31. Williams TE, Nagarajan S, Selvaraj P, Zhu C. Concurrent and independent binding of

Fcgamma receptors IIa and IIIb to surface-bound IgG. Biophys J. 2000;79(4):1867-75.

32. Galon J, Gauchat JF, Mazières N, Spagnoli R, Storkus W, Lötze M, et al. Soluble Fcgamma

receptor type III (FcgammaRIII, CD16) triggers cell activation through interaction with

complement receptors. J Immunol. 1996;157(3):1184-92.

33. Selvaraj P, Fifadara N, Nagarajan S, Cimino A, Wang G. Functional Regulation of Human

Neutrophil Fc γ Receptors. Immunology Research. 2004;29(1-3):219-30.

34. Hvilsom CT, Søgaard OS. TLR-Agonist Mediated Enhancement of Antibody-Dependent

Effector Functions as Strategy For an HIV-1 Cure. Front Immunol. 2021 ;12:704617.

35. Vibholm LK, Konrad CV, Schleimann MH, Frattari G, Winckelmann A, Klastrup V, et al.

Effects of 24-week Toll-like receptor 9 agonist treatment in HIV type 1+ individuals. AIDS.

2019;33(8):1315-25.

36. Moldt B, Chandrashekar A, Borducchi EN, Nkolola JP, Stephenson H, Nagel M, et al. HIV

envelope antibodies and TLR7 agonist partially prevent viral rebound in chronically SHIV-

infected monkeys. PLoS Pathog. 2022;18(4):e1010467.

37. Borducchi EN, Liu J, Nkolola JP, Cadena AM, Yu WH, Fischinger S, et al. Antibody and

TLR7 agonist delay viral rebound in SHIV-infected monkeys. Nature. 2018;563(7731):360-4.

38. Hsu DC, Schuetz A, Imerbsin R, Silsorn D, Pegu A, Inthawong D, et al. TLR7 agonist, N6-

LS and PGT121 delayed viral rebound in SHIV-infected macaques after antiretroviral therapy

interruption. PLoS Pathog. 2021;17(2):e1009339.

39. Bell LCK, Noursadeghi M. Pathogenesis of HIV-1 and Mycobacterium tuberculosis co-

infection. Nat Rev Microbiol. 2018;16(2):80-90.

40. Menten P, Wuyts A, Van Damme J. Macrophage inflammatory protein-1. Cytokine Growth

Factor Rev. 2002;13(6):455-81.

41. Syedbasha M, Egli A. Interferon Lambda: Modulating Immunity in Infectious Diseases.

Front Immunol. 2017;8.

42. Broggi A, Tan Y, Granucci F, Zanoni I. IFN-λ suppresses intestinal inflammation by non-

translational regulation of neutrophil function. Nat Immunol. 2017;18(10):1084-93.

43. Liu MQ, Zhou DJ, Wang X, Zhou W, Ye L, Li JL, et al. IFN-λ3 Inhibits HIV Infection of

Macrophages through the JAK-STAT Pathway. PLoS ONE. 2012;7(4):e35902.

44. Michailidis C, Giannopoulos G, Vigklis V, Armenis K, Tsakris A, Gargalianos P. Impaired

phagocytosis among patients infected by the human immunodeficiency virus: implication for a

role of highly active anti-retroviral therapy. Clin Exp Immunol. 2012;167(3):499-504.

45. Lemaitre J, Desjardins D, Gallouët AS, Gomez-Pacheco M, Bourgeois C, Favier B, et al.

Expansion of Immature Neutrophils During SIV Infection Is Associated With Their Capacity

to Modulate T-Cell Function. Front Immunol. 2022;13:781356.

46. Bowers NL, Helton ES, Huijbregts RP, Goepfert PA, Heath SL, Hel Z. Immune suppression

by neutrophils in HIV-1 infection: role of PD-L1/PD-1 pathway. PLoS Pathog.

2014;10:e1003993.

47. Marsile-Medun S, Souchard M, Abba Moussa D, Reynaud É, Tuaillon E, Naranjo-Gomez

M, et al. Fc receptors are key discriminatory markers of granulocytes subsets in people living

with HIV-1. Front Immunol. 2024;15:1345422.

48. Cloke T, Munder M, Taylor G, Müller I, Kropf P. Characterization of a novel population

of low-density granulocytes associated with disease severity in HIV-1 infection. PLoS One.

2012;7(11):e48939.

49. Cloke T, Munder M, Bergin P, Herath S, Modolell M, Taylor G, et al. Phenotypic alteration

of neutrophils in the blood of HIV seropositive patients. PLoS One. 2013;8(9):e72034.

50. Liu K, Huang HH, Yang T, Jiao YM, Zhang C, Song JW, et al. Increased Neutrophil Aging

Contributes to T Cell Immune Suppression by PD-L1 and Arginase-1 in HIV-1 Treatment

Naïve Patients. Front Immunol. 2021;12:670616.

51. Yoshimura T, Takahashi M. IFN-γ-Mediated Survival Enables Human Neutrophils to

Produce MCP-1/CCL2 in Response to Activation by TLR Ligands. J. Immunol.

2007;179(3):1942-9.

52. Kummola L, Salomaa T, Ortutay Z, Savan R, Young HA, Junttila IS. IL-4, IL-13 and IFN-

γ -induced genes in highly purified human neutrophils. Cytokine. 2023;164:156159.

53. Guihot A, Marcelin AG, Massiani MA, Samri A, Soulié C, Autran B, et al. Drastic decrease

of the HIV reservoir in a patient treated with nivolumab for lung cancer. Ann Oncol.

2018;29(2):517-8.

54. Gay CL, Bosch RJ, Ritz J, Hataye JM, Aga E, Tressler RL, et al. Clinical Trial of the Anti-

PD-L1 Antibody BMS-936559 in HIV-1 Infected Participants on Suppressive Antiretroviral

Therapy. J Infect Dis. 2017;215(11):1725-33.

55. Inamdar K, Tsai FC, Dibsy R, de Poret A, Manzi J, Merida P, et al. Full assembly of HIV-

1 particles requires assistance of the membrane curvature factor IRSp53. Elife.

2021;10:e67321.

56. Scheid JF, Mouquet H, Ueberheide B, Diskin R, Klein F, Oliveira TYK, et al. Sequence

and structural convergence of broad and potent HIV antibodies that mimic CD4 binding.

Science. 2011;333(6049):1633-7.

57. Mouquet H, Scharf L, Euler Z, Liu Y, Eden C, Scheid JF, et al. Complex-type N-glycan

recognition by potent broadly neutralizing HIV antibodies. Proc Natl Acad Sci U S A.

2012;109(47):E3268-3277.

58. Lorin V, Mouquet H. Efficient generation of human IgA monoclonal antibodies. J Immunol

Methods. 2015;422:102-10.

59. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape

provides a biologist-oriented resource for the analysis of systems-level datasets. Nat Commun.

2019;10(1):1523.

Figure Legends

Figure 1. Activation of human neutrophils stimulated *in vitro* by Toll-like receptors (TLRs) agonists.

Human neutrophils from healthy donors were activated by TLRs agonists, TLR4 (LPS), TLR7/8 (R848). (A). Gating strategy used for all experiments. (B-C) Viability (B) and phenotypic activation (C) assessed by monitoring cell surface markers. Fold change is measured versus unstimulated cells (dashed lines). It is represented as the ratio MFI stimulated cells/MFI unstimulated cells. (D-E) Cytokines and chemokines secretion profile, represented as heatmap (D) and histograms for each cytokine/chemokine assessed (E). Values are mean +/- SD of at least 25 (B-C) and 8 (D-E) independent measurements. Significance was assigned as follows: ∗∗p <0,01 and ∗∗∗p < 0.001. NS: non-stimulated.

Figure 2. Activation of human neutrophils stimulated *in vitro* by TNF-α or IFN-γ.

Human neutrophils from healthy donors were activated overnight by TNF-α at 1 ng/mL or IFN-γ at 1000 IU/mL. (A-B) Viability (A) and phenotypic (B) activation assessed by monitoring cell surface markers. Fold change is measured versus unstimulated cells (dashed lines). It is represented as the ratio MFI stimulated cells/MFI unstimulated cells. (C-D) Cytokines and chemokines secretion profile, represented as a heatmap (C) and histograms (D) for each cytokine/chemokine assessed. Values are mean +/- SD of at least 19 (A) and 8 (B-D) independent measurements. Significance was assigned as follows: ∗p < 0.05, and ∗∗p < 0.01. NS: non-stimulated.

Figure 3. Activation of human neutrophils stimulated *in vitro* by HIV-1 virions, ICs or free bNAb.

Human neutrophils from healthy donors were activated by HIV-1 virions, ICs or free bNAb. (A-B) Viability (A) and phenotypic activation (B) assessed by monitoring cell surface markers. Fold change is measured versus unstimulated cells (dashed lines). It is represented as the ratio MFI stimulated cells/MFI unstimulated cells. (C-D) Cytokines and chemokines secretion profile, represented as a heatmap (C) and histograms (D) for each cytokine/chemokine assessed. Values are mean +/- SD of at least 21 (A-B) and 17 (C-D) independent measurements. Significance was assigned as follows: ∗p < 0.05, ∗∗p < 0.01, ∗∗∗p < 0.001, and ∗∗∗∗p < 0.0001. NS: non-stimulated.

Figure 4. Secretion profile of human neutrophils primed with TNF-α (B) or IFN-γ (C) and then stimulated *in vitro* by HIV-1 virions, ICs or free bNAb.

(A) Protocol for priming of neutrophils. Dashed lines indicate the level of secretion induced by either TNF-α alone (B) or IFN-γ alone (C). Values are of 10 (B) and 6 (C) independent measurements. Significance was assigned as follows: ∗p < 0.05 and ∗∗p < 0.01

Figure 5. Transcriptomic analysis of neutrophils primed with TNF-α or IFN-γ.

Heatmap (A) and Volcano plot (B) representation of upregulated and downregulated genes by TNF-α or IFN-γ. (C) Venn diagram of regulated genes after cytokine stimulation. (D) Genes upregulated and downregulated were analyzed using Metascape (https://metascape.org/gp/index.html#/main/step1) to identify main biological processes modulated by cytokine priming. Data from 4 independent samples.

Figure 6. Activation of human neutrophils isolated from HD or PLWH stimulated *in vitro* by TLR agonists LPS (TLR4) and R848 (TLR7/8) or inflammatory cytokines (TNF-α and IFN-γ).

(A-B) Cytokines and chemokines secretion profile upon activation by TLR agonists, represented as a heatmap (A) and histograms (B) for each cytokine/chemokine assessed. (C-D) Cytokines and chemokines secretion profile upon activation by inflammatory cytokines, represented as a heatmap (C) and histograms (D) for each cytokine/chemokine assessed. Values are mean +/- SD of at least 9 (A-B) and 4 (C-D) independent measurements.

Figure 7. Activation of human neutrophils isolated from HD or PLWH stimulated *in vitro* by HIV-1 virions, ICs or free bNAb.
(A-B) Cytokines and chemokines secretion profile, represented as a heatmap (A) and histograms (B) for each cytokine/chemokine assessed. Values are mean +/- SD of 6 independent measurements.

Figure 8. Modulation of the expression of FcγR on human neutrophils isolated from HD or PLWH stimulated *in vitro* by TLR agonist, inflammatory cytokines, HIV-1 virions, ICs or free bNAb.
(A-C) Modulation of the cell surface levels of CD16 and CD32 upon activation by TLR agonists (A), cytokines (B), and HIV-1 virions, ICs or free bNAb (C). Fold change is measured versus unstimulated cells (dashed lines). It is represented as the ratio MFI stimulated cells/MFI unstimulated cells. Values are mean +/- SD of at least 14 (A), 8 (B) and 6 (C) independent measurements. Significance was assigned as follows: $*p < 0.05$ and $**p < 0.01$

Figure 9. Phenotypic characterization of neutrophils isolated from PLWH and HD blood samples.
Purified neutrophils were immunophenotyped by assessing the expression of 18 cell surface markers. Data include at least 19 PLWH blood samples and 4 HD samples. Statistical analyses are unpaired Mann-Whitney test. Significance was assessed as follows: $*p < 0.05$, $***p < 0.001$, and $****p < 0.0001$.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9