1

Title. *Structure, function and assembly of soybean primary cell wall cellulose synthases*

Authors: Ruoya Ho[1,#], Pallinti Purushotham[1,#+], Louis F. L. Wilson[1.2], Yueping Wan[1,], Jochen Zimmer[1,2]*

Affiliations:

[1] Department of Molecular Physiology and Biological Physics, University of Virginia School of Medicine, Charlottesville, VA 22903.

[2] Howard Hughes Medical Institute.

[#] Equal contribution

[+] Present address: Department of Life Sciences, GITAM University, Bengaluru, Karnataka 561203, India

* For correspondence: jz3x@virginia.edu

Abstract

Plant cell walls contain a meshwork of cellulose fibers embedded into a matrix of other

carbohydrate and non-carbohydrate-based biopolymers. This composite material exhibits

extraordinary properties, from stretchable and pliable cell boundaries to solid protective shells.

Cellulose, a linear glucose polymer, is synthesized and secreted across the plasma membrane by

cellulose synthase (CesA), of which plants express multiple isoforms. Different subsets of CesA

isoforms are necessary for primary and secondary cell wall biogenesis. Here, we structurally and

functionally characterize the *Glycine max* (soybean) primary cell wall CesAs CesA1, CesA3, and

CesA6. The CesA isoforms exhibit robust *in vitro* catalytic activity. Cryo-electron microscopy

analyses reveal their assembly into homotrimeric complexes *in vitro* in which each CesA

protomer forms a cellulose-conducting transmembrane channel with a large lateral opening.

Biochemical and co-purification analyses demonstrate that different CesA isoforms interact *in*

vitro, leading to synergistic cellulose biosynthesis. Interactions between CesA trimers are only

observed between different CesA isoforms and require the class-specific region (CSR). The CSR

forms a hook-shaped extension of CesA's catalytic domain at the cytosolic water-lipid interface.

Negative stain and cryo-electron microscopy analyses of mixtures of different CesA isoform

trimers reveal their side-by-side arrangement into loose clusters. Our data suggest a model by

which CesA homotrimers of different isoforms assemble into cellulose synthase complexes to

synthesize and secrete multiple cellulose chains for microfibril formation. Inter-trimer

interactions are mediated by fuzzy interactions between their CSR extension.

Introduction

Cellulose is a versatile biopolymer and a fundamental building block of plant cell walls. It is an

amphipathic linear β-1,4 linked glucose polymer that can be assembled into fibrillar structures.

As the load-bearing wall component of vascular plants, cellulose microfibrils are spun around the

cell and integrated with a variety of other biopolymers (Turner and Kumar, 2018). Cellulose is

synthesized by cellulose synthase (CesA), a membrane-integrated processive family-2

glycosyltransferase (GT) (Lombard et al., 2014). The enzyme synthesizes cellulose from UDP-

activated glucose (UDP-Glc) and translocates the polymer across the plasma membrane through

a channel formed by its own membrane-spanning segment (McNamara et al., 2015; Morgan et

al., 2013). While bacterial CesAs primarily function as monomeric enzymes (Abidi et al., 2021;

Acheson et al., 2021; Du et al., 2016; Morgan *et al.*, 2013), structural analyses of plant CesAs

revealed their assembly into triangular-shaped trimeric complexes of three catalytically active

subunits (Massenburg et al., 2024; Purushotham et al., 2020; Zhang et al., 2021a)

The overall CesA architecture and the mechanism of cellulose biosynthesis is evolutionarily

conserved from bacteria to land plants. However, plant CesAs contain specific domains absent in

most bacterial homologs. These include an extended cytosolic N-terminus beginning with a

RING-like region, a plant conserved region (PCR), as well as a class-specific region (CSR)

(Carpita, 2011; Purushotham *et al.*, 2020; Wilson et al., 2021). The CSR and PCR are inserted

into the cytosolic catalytic domain. While the PCR is a trimerization domain that connects three

CesA promoters in a trimeric complex, the function and structure of the CSR remains unknown.

Further, plants express different CesA isoforms at different developmental stages, of which

certain subsets are necessary for primary and secondary cell wall formation (Persson et al., 2007;

Taylor et al., 2003; Turner and Somerville, 1997). Based on the Arabidopsis nomenclature,

isoforms associated with primary cell wall formation include CesA1, CesA3 and CesA6, while

secondary cell wall CesAs are CesA4, CesA7 and CesA8. The isoenzymes vary the most within

the CSR and the N-terminal domain.

Genetic analyses demonstrated the importance of different CesA isoforms for primary and

secondary cell wall formation (Fagard et al., 2000; Turner and Somerville, 1997; Persson *et al.*,

2007; Sampathkumar et al., 2019) Further, co-immunoprecipitation analyses indicated direct

interactions between primary or secondary cell wall CesA isoforms (Gonneau et al., 2014;

Taylor *et al.*, 2003; Timmers et al., 2009). However, the specific functions of the different CesA

isoforms during *in vivo* cellulose biosynthesis remain unknown.

Plants organize cellulose into micro and macro-fibrils (Kubicki et al., 2018; Turner and Kumar,

2018). Cellulose microfibrils likely originate from supramolecular CesA complexes (CSCs)

observed in various species. In land plants, CSCs appear primarily as pseudo sixfold symmetric

membrane-integrated clusters by freeze fracture electron microscopy analyses (Herth and Weber,

1984; Kimura et al., 1999; Nixon et al., 2016). The CesA trimer likely represents the CSC repeat

unit, thereby accounting for 18 CesAs per CSC and, accordingly, 18 cellulose polymers in a

CSC-synthesized microfibril (Cosgrove et al., 2024; Nixon *et al.*, 2016; Turner and Kumar,
2018).

To analyze the oligomerization and function of primary cell wall CesAs, we recombinantly
expressed and purified *Glycine max* (soybean) CesA1, CesA3 and CesA6. Cryo-EM analyses of
all three CesA isoforms reveal the formation of homotrimeric complexes, similar to the
secondary cell wall CesAs. The CSR is resolved at the corners of the CesA trimer as a disordered
but hook-shaped domain that runs at the cytosolic water-lipid interface. *In vitro* co-purification
and electron microscopy studies demonstrate that homotrimers of different CesA isoforms
interact. This interaction requires the CSR and leads to synergistic cellulose biosynthesis. Our
results support a model by which CSCs are formed from homotrimers of different CesA isoforms.

Results

Primary cell wall CesAs purify as high and low molecular weight species
We selected a set of *G. max* (*Gm*) CesA isoforms that phylogenetically cluster with *Arabidopsis*
thaliana CesA1, CesA3 and CesA6, respectively, and that are widely and strongly expressed in
unlignified soybean tissues (***Figure 1*** and ***Figure supplement 1A*** *and* ***B***). Existing co-expression
data from ATTED-II (https://atted.jp/ (Obayashi et al., 2022) indicate that the enzymes are co-
expressed with other primary cell wall genes implicated in pectin, arabinogalactan, and
galactoglucomannan biosynthesis (***Figure 1***). We therefore conclude that the selected CesAs
indeed represent primary cell wall CesAs.
To biochemically and structurally characterize the CesAs, we followed a similar heterologous
expression protocol as established previously for hybrid aspen CesA8 (Purushotham *et al.*, 2020).
In short, the CesAs were expressed with N-terminal poly-histidine tags in Sf9 insect cells and
purified by metal affinity and size exclusion chromatography in the detergent glyco-diosgenin
(GDN) (Materials and Methods). Size exclusion chromatography separated all CesA isoforms
into high and low molecular weight species (***Figure 2***A-C). Cryogenic and negative stain EM
analyses identified these species as CesA trimers and monomers, respectively (see below). Of
note, compared to CesA1 and CesA3, the yield of trimeric CesA6 was more variable, with some

preparations producing primarily monomeric species. This suggests that CesA6 is less stable in a
detergent-solubilized state compared with the other isoforms.

To test whether the detergent-solubilized CesA trimers dissociate into monomers over time, the
purified CesA trimers were reinjected onto the size exclusion chromatography column after an
overnight incubation on ice. For all species, the reinjected material eluted as a trimeric complex,
indicating that assembled trimers are stable and do not interconvert with monomers within this
timeframe (*Figure 2***A-C**). The co-purifying monomers likely arise from incompletely assembled
trimers or oligomer dissociation during purification.

***In vitro* cellulose biosynthesis**
Cellulose biosynthetic activity of the purified CesAs was quantified by measuring the
incorporation of ^{3}H-labeled glucose into insoluble cellulose, followed by scintillation counting,
as previously described (Purushotham et al., 2016). As shown in *Figure 2***D**, the relative
activities of the monomeric and trimeric CesA fractions are comparable for each isoform,
demonstrating that both species are catalytically active *in vitro*. Between the different isoforms,
CesA1 exhibits greatest product accumulation (*Figure 2***D**). In all cases, the *in vitro* synthesized
polymer is readily degraded by a cellulase, indicating the formation of authentic cellulose. No
product was obtained in the presence of EDTA, in agreement with previous observations
(Purushotham *et al.*, 2016).

To further assess catalytic differences between the CesA isoforms, we determined their pH
optima for catalytic activity (*Figure 2***E**). All CesA isoforms show greatest catalytic activity at
neutral to mild-alkaline pH. CesA3 exhibits an activity optimum at pH 7 with a sharp decline at
pH 8 and 9. In contrast, the activities of CesA1 and CesA6 peak at pH 8, with a slight decline at
pH 9 (*Figure 2***E**). Quantifying the release of UDP during biosynthesis reactions using a 'UDP-
Glo' glycosyltransferase assay (Das et al., 2016) reveals Michaelis–Menten constants for CesA1
and CesA3 with respect to UDP-Glc of 0.44 mM and 0.18 mM, respectively. The apparent V_{max}
value is ~8-fold higher for *Gm*CesA1 compared to *Gm*CesA3 (*Figure supplement 1C*). The
activity of CesA6 was too weak to be analyzed by this method. Further, we analyzed inhibition
of the isoforms by UDP, which competitively inhibits CesA8 and related GT-2 enzymes (Gow

and Selitrennikoff, 1984; Omadjela et al., 2013; Purushotham et al., 2016; Tlapak-Simmons et al.,

2004). CesA1's apparent IC_{50} for UDP is about 0.8 mM, whereas this concentration is increased

to about 1.3 to 1.5 mM for CesA6 and CesA3, respectively (***Figure 2F***).

Homotrimer assembly

Cryo-EM analyses of the high molecular weight CesA fractions (***Figure 2*A-C**) revealed their

organization into homotrimeric complexes (***Figure 3*A - *Figure supplements 2 and 3 and***

Supplement table 1). For all isoforms, two-dimensional classification identified trimeric particles

similar to hybrid aspen CesA8 and cotton CesA7 (Purushotham *et al.*, 2020; Zhang *et al.*, 2021a).

Particle classification in 3 dimensions followed by non-uniform refinement with applied C3

symmetry and local refinement generated cryo-EM maps ranging in resolution from about 3.0 to

3.3 Å.

Overall, the CesAs contain a cytosolic catalytic domain that interacts with the channel-forming

TM region via three amphipathic IF helices (***Figure 3*B**). The helices surround the entrance to

the TM channel with the Trp residue of the QxxRW motif at its portal. As previously described

for hybrid aspen CesA8 and cotton CesA7, the CesA trimers are stabilized by the PCR domain

that is inserted into CesA's catalytic domain. (Purushotham *et al.*, 2020; Zhang *et al.*, 2021a).

The triangular PCR arrangement in a CesA complex positions the side chains of conserved Lys

and Arg residues towards the 3-fold symmetry axis. These residues include Arg449, Lys452, and

Arg453 in CesA6 and coordinate unidentified ligand(s) on the membrane distal and proximal

side of the PCR triangle. For all three CesA isoforms, the ligands' shapes are similar, suggesting

that they represent the same small molecule (***Figure supplement 3*B**). On the membrane

proximal side, the density extends by about 9 Å towards the membrane, perhaps representing a

nucleotide bound in different poses, as previously suggested for CesA8 (Purushotham *et al.*,

2020).

A transmembrane channel with a large later opening

CesAs contain seven TM helices of which helices 1-6 create a cellulose conducting channel. In

the previously described CesA8 and CesA7 complexes, TM helix 7 of one protomer packs

against TM helices 5 and 6 of a neighboring CesA subunit (***Figure 3*C**). In our new soybean

CesA structures, however, this helix is more flexible, as evidenced by weaker map quality (*Figure supplement 3***A**), and shifted to the periphery of the trimer. In the new position, the helix primarily mediates contacts with TM helix 5 and the C-terminal segment of TM helix 3 of the same CesA protomer (*Figure 3***B** *and* **C**). The helix has been modeled for CesA1 and CesA6, while its density is detectable at a similar position but too discontinuous for modeling in the CesA3 map.

The displacement of TM helix 7 away from the TM channel of the neighboring subunit opens a lateral lipid-exposed window in the neighboring subunit's channel architecture (*Figure 3***C**). The window is formed by the N-terminal region of IF helix 3 and TM helices 4 and 6. About midway across the membrane, the opening is roughly 6 Å wide, for example between the side chains of Ile901 in TM helix 4 and Trp1018 in TM helix 6 (*Figure 3***C**). The lateral window likely exposes the translocating nascent cellulose polymer to the hydrophobicity of the lipid bilayer. A similar lipid exposed polysaccharide translocation pathway has recently been described for hyaluronan synthase (Maloney et al., 2022).

The CSR forms a hook-shaped extension of the catalytic domain

Plant CesA's contain two structurally and functionally unresolved domains, which are the N-terminal domain (NTD) and the CSR. The NTD has been resolved at lower resolution for hybrid aspen CesA8, where it forms a helical stalk extending from the catalytic domains into the cytosol (Purushotham *et al.*, 2020). We observe a similar stalk-like extension in some of the trimeric *Gm*CesA3 particles (*Figure supplement 2*). However, the NTD is only resolved for a small subset of *Gm*CesA3 particles and not resolved at all for the other CesAs. This suggests that it can adopt multiple conformations, with the stalk being one of them.

The CSR has been proposed to be intrinsically disordered (Scavuzzo-Duggan et al., 2018) and only its short N- and C-terminal helical segments are visible in the cryo-EM maps. For *Gm*CesA1, however, at lower contour levels, additional CSR density is evident, extending from corners of the catalytic domains of the CesA1 trimer (*Figure 3D*). Viewed from the cytosol, the extra density resembles a hook extending by about 20 Å clockwise and tangentially along the trimer's corners at the water-lipid interface. At this position, the CSR's N-terminal conserved

cysteine residue(s) postulated to be acylated (Kumar et al., 2016) reside near the membrane
interface (*Figure 3D and E*).

Homotrimers of different CesA isoforms interact

We next tested whether CesA homotrimers of different isoforms would interact *in vitro*. To this
end, we individually expressed and purified trimers of poly-His tagged CesA1 and CesA6 (His-
CesA1 and His-CesA6) and TwinStrep-tagged CesA1 and CesA3 (Strep-CesA1 and Strep-
CesA3) (*Figure 4*). Cross-isoform interactions were tested by tandem purifications over Ni-NTA
and Strep-Tactin affinity matrices. His-CesA1 can be distinguished from Strep-CesA3 by
Coomassie stained SDS-PAGE due to size differences, whereas all other species comigrate.
Therefore, Western blotting together with Coomassie stained SDS-PAGE was performed to
evaluate the co-purification results.

An equimolar mixture (based on UV absorbance) of His-CesA1 and Strep-CesA3 was
incubated for 180 min on ice and sequentially purified using (1) Ni-NTA resin and (2) Strep-
Tactin beads (*Figure 4A*). The Coomassie stained SDS-PAGE resolves both CesA species in the
initial mixture, after elution from the Ni-NTA resin, as well as upon elution from the Strep-
Tactin beads. The identity of the bands as His-CesA1 and Strep-CesA3 was confirmed by
Western blotting. We observed no non-specific binding of His-CesA1 to Strep-Tactin beads or
Strep-CesA3 to Ni-NTA resin (*Figure 4E*).

Similar experiments with combinations of Strep-CesA3 and His-CesA6 (*Figure 4B*) and
Strep-CesA1 and His-CesA6 (*Figure 4C*) yielded comparable results, although the co-eluting
species cannot be distinguished by Coomassie staining alone, due to comigration. None of the
species showed detectable non-specific binding to the affinity resins in the absence of the
corresponding tags, (*Figure 4E-G*).

As an additional control, we analyzed whether differently tagged homotrimers of the
same isoform also interact with each other. To this end, a mixture of His-CesA1 and Strep-
CesA1 was subjected to tandem affinity purification as described above (*Figure 4D*). We failed
to detect any co-purification of Strep-CesA1 when applying Ni-NTA as the first affinity
chromatography step. Similar results were obtained for Strep- and His-tagged combinations of
homotrimers of CesA3 or CesA6 (*Figure supplements 4A-D*). Qualitatively, our interaction data

are consistent with previously published co-immunoprecipitations of primary and secondary
CesA isoforms (Gonneau *et al.*, 2014; Timmers *et al.*, 2009).

Similar tandem purification experiments were also performed with the monomeric CesA
fractions obtained from size exclusion chromatography (*Figure 2***A-C**). As observed for the
homotrimeric complexes, monomeric His-CesA1 co-purifies with monomeric Strep-CesA3,
demonstrating that trimeric assemblies are not necessary for the observed interactions (***Figure***
supplements 4E *and* **F**).

Only the CSR is required for isoform interaction
The biological functions of CesA's NTD and CSR are currently unknown. The NTD's RING-
like domain has been shown to form dimers and trimers *in vitro* (Kurek et al., 2002;
Purushotham *et al.*, 2020), raising the possibility that it could form inter-trimer complexes
accounting for the observed isoform interactions.

To test this hypothesis, N-terminally truncated constructs of CesA1 and CesA3 were
expressed and purified as described for the full-length variants. The constructs lack the first 259
(CesA1) and 242 (CesA3) residues yet purify as trimers (besides monomers) and exhibit *in vitro*
catalytic activity similar to the full-length constructs (*Figure 5***A**). A tandem affinity purification
of a mixture of the truncated His-CesA1 and Strep-CesA3 isoforms demonstrates their
interaction in the absence of the NTD, similar to the full-length enzymes (*Figure 5***B** *and* **C** *and*
*Figure 4***A**).

To test whether the CSR is involved in isoform interaction, the domain was replaced in
the N-terminally truncated *Gm*CesA1 and *Gm*CesA3 constructs with a flexible loop of 20
residues (see Methods), thereby generating CesA constructs devoid of the NTD as well as the
CSR (**Figure 5D**). Negative stain EM of the purified truncated CesA1 particles demonstrates
their trimeric assembly, as observed for the full-length and NTD truncated versions (*Figure 5D*).
Upon deletion of the CSR, the CesA1 and CesA3 isoforms no longer co-purify *in vitro*,
suggesting that the region is indeed required for inter-isoform interaction (*Figure 5E*). Further,
an N-terminally and CSR truncated CesA3 construct shows only minor, most likely non-specific,
interaction with full-length CesA1, suggesting that CesA isoforms interact primarily via their
CSR domains (*Figure 5F*).

Clustering of homotrimers of different CesA isoforms

Negative stain electron microscopy analyses of the individual CesA homotrimers revealed monodisperse particle distributions (*Figure 6***A-C**). The size and shape of the particles is consistent with the CesA homotrimers observed by cryo-EM and remains unchanged over a course of a week when incubated on ice.

To visualize the interaction of the isoforms, all three homotrimeric isoforms were combined and subjected to size exclusion chromatography after incubation overnight on ice. Negative stain EM analysis of high molecular weight fractions eluting after the void volume revealed CesA clusters of varying stoichiometries, ranging from 5 to >10 particles (*Figure 6***D**). The clusters likely arise from trimer-trimer interactions in different orientations and vary in diameter from about 50 to 100 nm. No clustering was observed for any of the individual CesA isoforms alone, even after prolonged incubations on ice.

Further, cryo-EM was used to analyze a mixture of individually purified CesA1 and CesA3 trimers. The trimers were combined and incubated on ice for 60 min prior to cryo grid preparation. Focusing on particles larger than a CesA trimer, 2-dimensional classification of the obtained particles revealed shapes resembling arrangements of two CesA trimers. The identified particles are consistent with side-by-side arrangements of two CesA trimers in opposite orientations. By this organization, the catalytic domains of the trimers are co-planar and the micelle-embedded TM segments are above and below the plane (*Figure 6***E**).

To support this interpretation, two cryo-EM volumes of a CesA1 trimer were manually arranged side-by-side but in inverted orientations. The individual volumes were placed such that the CSR densities of two CesA subunits of each trimer would contact each other. The obtained dimer-of-trimer volume was then used to calculate 2-dimensional projections for comparison with the experimentally obtained 2D class averages. Indeed, the generated 'upside down and side-by-side' arrangement of CesA trimer volumes resembles the experimental class averages (*Figure 6***F and G**). Steric interferences of the micelle-embedded TM regions likely prevent the physiological parallel arrangement of two CesA trimers under the experimental conditions.

Synergistic cellulose biosynthesis

We investigated whether the cross-isoform interaction of CesA trimers affects their *in vitro* catalytic activities. To this end, *in vitro* cellulose biosynthesis was quantified radiometrically

from reactions containing one CesA isoform at a constant concentration (20 µM) and increasing
concentrations of the other (1-20 µM). As a reference, cellulose biosynthetic activities were also
determined for each isoform alone at the concentrations used in the combined assays. As shown
in Figure 7A-F, for all isoform combinations, the measured activities exceed the theoretical
activities (calculated by adding the individually measured activities) at least one to two-fold,
depending on the isoform combination. This suggests synergistic cellulose biosynthesis in the
presence of two CesA isoforms. Performing the titration experiment with samples of the same
isoform does not reveal any synergy, consistent with the lack of interaction between trimers of
the same CesA isoform (*Figure 7**G** and **4D** - **Figure supplement 5***).

Additionally, comparing the measured and additive activities obtained after combining all
three CesA isoforms (at 6.6 µM each) reveals an experimental activity about 3-fold above the
theoretical value. This activity level may arise from different dimeric arrangements of CesA
trimers (1+3, 1+6, and 3+6) and/or the formation of larger complexes of different isoform
trimers (1+3+6) (*Figure 7**H***).

CesA1 exhibits higher *in vitro* catalytic activity compared to CesA3 and CesA6 (*Figure 2**D** and
*Figure supplement 1B***). To test whether the observed synergistic effects are due to altered
catalytic activity of only one isoform or both, one CesA isoform was inactivated after
purification by incubation with the oxidant sodium tetrathionate (Tie et al., 2004). While the
inactivated CesAs exhibit activity levels comparable to EDTA-treated negative controls, they
remain trimeric and interact with other isoforms as observed for the unmodified versions (*Figure
supplement 6***). Inactivation could be due to modification of a conserved cysteine residue in
CesA's catalytic pocket (such as Cys558 or Cys630 in CesA6).

Performing the above-described activity assays with pairs of inactive and active CesA
trimers demonstrates that all three isoforms exhibit increased catalytic activity when combined
with an inactive trimer of a different isoform (*Figure supplement 7***). This suggests that
intertrimer interactions impact the catalytic activity of all isoforms, perhaps by altering the
accessibility of the catalytic pocket (discussed below).

Discussion

A hallmark of plant cellulose biosynthesis is the deposition of microfibrils in the cell wall (Zhang et al., 2021b). Prevailing models of microfibril forming CSCs postulate that they contain at least three different CesA isoforms (Newman et al., 2013; Turner and Kumar, 2018). However, experimental evidence supporting the presence of different CesA isoforms in a CSC and its repeat unit is lacking.

Despite efficient co-expression of differently tagged CesA1, CesA3, and CesA6 in insect Sf9 cells, we failed to isolate hetero-oligomeric complexes suitable for structural analysis (*Figure supplement 8*). This failure does not exclude the formation of a small fraction of hetero-oligomeric CesA complexes; however, it indicates the preferred formation of homo-oligomeric CesA complexes. Therefore, we characterized the CesA isoforms individually. All three isoforms can be purified as catalytically active homotrimeric species. Trimerization is mediated by the cytosolic PCR, as observed in the secondary cell wall CesA8 and CesA7 isoforms. Because the PCR is highly conserved across the three *Gm*CesA isoforms, the apparent failure (or low efficiency) of heterotrimer formation may be due to subtle differences in sequence and shape complementarity between the isoforms.

The different arrangement of TM helix 7 in the *Gm*CesAs compared to CesA8 and CesA7 creates a large lateral window in the cellulose secretion channel. The window opens towards the trimer's 3-fold symmetry axis. While the biological function of this window or the flexibility of TM helix 7 are unclear, it could enable the lateral release of the glucan chains towards the center of the complex, thereby affecting protofibril formation (Kurek *et al.*, 2002; Purushotham *et al.*, 2020).

Our *Gm*CesA3 structure contrasts with the recently reported dimeric organization of a fragment of *A. thaliana's* (*At*) CesA3 catalytic domain, which self-associates via β-strand augmentation (Qiao et al., 2021). Our cryo-EM analysis of *Gm*CesA3 and the other isoforms only revealed monomeric and trimeric states, likely because the region involved in dimerization of the *At*CesA3 fragment is inaccessible in the full-length *Gm*CesA3 protein.

Our *in vitro* CesA interaction studies replicate previous *in vivo* co-immunoprecipitation results

on primary and secondary cell wall CesAs (Gonneau *et al.*, 2014; Timmers *et al.*, 2009). The

robust interaction of trimers of different CesA isoforms supports their physiological significance.

In a detergent solubilized state, the individual CesA trimers are not confined to the same plane,

as is the case in a biological membrane. Thus, the *in vitro* observed interactions lead to clustering

of the trimers, instead of their ordered close packing into symmetric particles.

364   Our data further demonstrate that the CSR, a region predicted to be intrinsically

disordered, is the primary mediator of trimer-trimer interactions (***Figure 6 and 8***), in agreement

with earlier suggestions (Sethaphong et al., 2013; Singh et al., 2020). The observation that CesA

trimers can assemble in a non-physiological upside-down orientation suggests that the CSRs in a

complex remain flexible, similar to previously described fuzzy complexes of intrinsically

disordered proteins (Fuxreiter, 2012). A fuzzy CSR-CSR interface may function like a magnet to

associate CesA trimers into a CSC, thereby bestowing flexibility on the CSC during microfibril

formation. The nascent cellulose fiber may further stabilize the complex.

We postulate the presence of distinct interaction hotspots for two nonidentical CesA isoforms

within a CSR. A dimeric complex of, for example CesA1 and CesA3, would then only be able to

interact with CesA6, thereby explaining the functional importance of three CesA isoforms. This

model can be modified by assuming multiple binding sites for the same 'non-like' isoform or

even a 'like' isoform to account for possible CSC configurations of two or one isoforms,

respectively. Because trimers of the same isoform do not interact *in vitro*, CSC models relying

on interactions between the same CesA isoforms across the repeat units are unlikely (***Figure 8***).

Due to its disordered nature (Scavuzzo-Duggan *et al.*, 2018) the CSR likely occupies a large

volume at the periphery of the catalytic domain. It is thus possible that the domain affects

nucleotide binding to or exchange at the active site. The CSR could be repositioned upon

complex formation with another isoform, which in turn could increase the catalytic activity and

account for the observed synergistic effects.

Lastly, the biological functions of the different CesA isoforms are currently unknown. Requiring

different isoforms to form a functional CSC could provide regulatory control over the cellulosic

material deposited in the cell wall. It is conceivable that CesA trimers function alongside fully

assembled CSCs in the plasma membrane, thereby producing proto- and microfibrils (from CesA

trimers and CSCs, respectively) that may interact. Accordingly, controlling the ability of the

CesAs to assemble into CSCs by regulating the isoform composition in the plasma membrane

may allow tailoring the fibril-to-protofibril ratio and thereby wall properties. Addressing these

questions will require detailed *in vivo* studies of the oligomerization and distribution of cellulose

depositing CesA complexes in the plasma membrane.

Materials and methods

Molecular phylogeny

A pre-computed cluster of loose CesA homologues (HOM05D000074) was downloaded from

Dicots PLAZA 5.0 (https://bioinformatics.psb.ugent.be/plaza/versions/plaza_v5_dicots/ (Van

Bel et al., 2022) , from which soy (*Glycine max*), Arabidopsis, cotton (*Gossypium hirsutum*),

poplar (*Populus trichocarpa*), tomato (*Solanum lycopersicum*), Physcomitrium, and Amborella

protein sequences were isolated. The sequence of CesA8 from *Populus*

tremula x *tremuloides* (*Ptt*CesA8) was also added at this stage for later reference. A region

corresponding to the Pfam 'Cellulose synthase' domain HMM profile (PF03552) was then

extracted using HMMER (Eddy, 2011) with an *E*-value cutoff of 1×10^{-80} and a bespoke Python

script. The extracted sequences were aligned using MAFFT v7.490 (Katoh and Standley, 2013)

before an initial near-maximum likelihood phylogeny using FastTree under default settings. This

tree was used to extract *bona fide* CesA sequences from Csl sequences using Figtree. The

corresponding rows of the alignment were then extracted alongside *A. thaliana* CslD5, which

was used as an outgroup in the following analysis. Model selection was carried out using

ProtTest version 3.4.2 (Darriba et al., 2011) before calculating the final maximum-

likelihood phylogeny using RAxML version 8.2.12 (Stamatakis, 2014) under a JTT+I+Γ model

with 100 rapid bootstrap pseudo-replicates. The tree was rendered in Figtree with subsequent

labelling in Inkscape.

Cloning

The primary cell wall CesA1, CesA3 and CesA6 genes from soybean (Glyma.06G069600,

Glyma.12G237000, and Glyma.02G080900) were synthesized (Gene Universal) with an N-

terminal 12x His-tag coding sequence and cloned into *Not*I and *Hind*III restriction sites in the

pACEBac1vector. The N-terminally TwinStrep-tagged CesA1 and CesA3 constructs were

generated by QuikChange mutagenesis from the pACEBac1-12xHisCesA vectors. The N-

terminally deleted GmCesA constructs (ΔNCesAs) were generated by PIPE cloning from the

full-length constructs, resulting in plasmids pACEBac1-ΔNCesA1R260, pACEBac1-

ΔNCesA3V243, and pACEBac1-ΔNCesA6M248. The CSR regions of CesA1 (residues 654-713)

and CesA3 (residues 648-709) were replaced with the sequence

ASGAGGSEGGGSEGGTSGAT(Baytshtok et al., 2017) in the -ΔNCesA1R260 and -

ΔNCesA3V243 backgrounds by gene synthesis.

A multi CesA expression cassette was generated for the co-expression of 12xHis-GmCesA1,

TwinStrep-GmCesA3 and 3xFlag-GmCesA6 according to the protocol detailed in the

MultiBac™ Multi-Protein Expression in Insect Cells manual (Geneva Biotech). The assembly

was done by homing endonuclease/BstXI multiplication. Briefly, the TwinStrep-GmCesA3-

pACEBac1vector (acceptor) was single restriction digested with BstXI. The His-GmCesA1-

pACEBac1vector was double restriction digested with I-CeuI and BstXI to excise the promoter

and terminator containing His-GmCesA1 (donor). The resulting fragment was inserted into the

BstXI digested TwinStrep-GmCesA3-pACEBac1vector to generate the TwinStrep-

GmCesA3_His-GmCesA1-pACEBac1 construct. The above cloning procedure was repeated one

more time to insert the 3xFlag-GmCesA6 (donor) into the GmCesA1 and GmCesA3 containing

vector (acceptor) to generate the multi CesA expression construct TwinStrep-CesA3_His-

GmCesA1_3xFlag-GmCesA6-pACEBac1.

Virus generation

An aliquot of 3 μL of 100 ng/μL pACEBac1-CesA plasmid was used for transformation into 50

μL chemically-competent DH10MultiBacTurbo *E. coli* cells. Bacmids were isolated from white

colonies on a Bluo-Gal agar plate and transfected into *Spodoptera frugiperda* 9 (SF9) cells. P0,

P1 and P2 baculovirus was generated according to the Joint Centre for Innovative Membrane

Protein Technologies (JCIMPT) protocol.

Protein expression and purification

For *Gm*CesA expression, Sf9 insect cells were infected with 15 mL P2 baculovirus at a density

of 3×10^{-6} cells per mL and grown at 27 °C for 48–72 h in an orbital shaker. Cells were then

harvested by centrifugation at 5,000×*g* for 10 min at 4 °C. Cell pellets were resuspended in

buffer A (20 mM Tris-HCl, pH 7.5, 100 mM NaCl, 5 mM sodium phosphate, 5 mM sodium

citrate, and 1 mM TCEP) supplemented with 1% lauryl maltose nopentyl glycol (LMNG,

Anatrace), 0.2% cholesteryl hemisuccinate (CHS, Anatrace), and protease inhibitors PIC (0.4

mM AEBSF, 2 μM aprotinin, 30 μM pepstatin, 7.5 μM pepstatin, 40 μM bestatin, 35 μM E-64,

and 4.5 mM benzamidine hydrochloride) and lysed in a glass dounce homogenizer ☐(~ 30

strokes). The lysate was solubilized at 4 °C for 1 h on a rocker. After separation of insoluble

material by centrifugation at 204,709×*g* for 45 min, 5 mL of Ni-NTA resin (HisPur Ni-NTA

Resin, Thermo scientific) and 20mM imidazole was added to the supernatant and incubated for 1

h at 4 °C on a rocker. After batch binding, the resin was packed into a gravity flow column and

then, sequentially washed twice with ten column volumes each of buffer A containing 40 mM

imidazole, 0.02% glyco-diosgenin (GDN, Anatrace) and PIC, followed by ten column volumes

of buffer A containing 1 M NaCl and 0.02% GDN (wash 3) and PIC. The final wash step (wash

4) was with ten column volumes of buffer A containing 0.02% GDN, PIC and 60 mM imidazole.

The *Gm*CesAs were eluted with six column volumes of buffer A containing 0.02% GDN, PIC

and 400 mM imidazole.

The TwinStrep-tagged GmCesAs were affinity purified by incubating the membrane extract with

5 ml Strep-Tactin sepharose at 4 °C for 1 h on a rocker. After batch binding, the resin was

packed into a gravity flow column and washed as described above for Ni-NTA affinity column

except the buffers lacked imidazole. The TwinStrep-tagged CesA was eluted using 6 column

volumes of buffer A containing 0.02% GDN and 5 mM desthiobiotin.

His- and TwinStrep-tagged *Gm*CesAs were further purified by size-exclusion chromatography

(SEC) using Superose 6 Increase 10/300 GL column (Cytiva) equilibrated in buffer A containing

0.02% GDN without any protease inhibitors. The purified *Gm*CesA trimers and monomers were

immediately used for activity assays and pulldown experiments or flash-frozen in liquid nitrogen

and stored at -80°C.

CesA-CesA pull-down assays

After SEC, either the trimeric or monomeric fractions of His-CesA and TwinStrep-CesA were

pooled and used for tandem affinity chromatography using Ni-NTA and Strep-Tactin sepharose

beads. His-CesA1 and TwinStrep-CesA3 trimers were mixed at 150 μg/mL concentration and

incubated at 4 °C for 3 h. The mixture was first loaded onto 200 μL bed volume Ni-NTA beads

for 1 h at 4 °C in the presence of 20 mM imidazole. After collecting the flowthrough, the beads

were washed with 10 bed volumes of buffer A containing 0.02% GDN and 20 mM imidazole for

3 washes followed by elution in 5 bed volumes of buffer A containing 0.02% GDN and 400 mM

imidazole.

The Ni-NTA eluted material was next bound to 200 μL Strep-Tactin sepharose beads for 1 h at

4 °C in a rotator shaker. After batch binding, the beads were packed into a gravity flow column.

The flow-through was collected and the column was subsequently washed 3-times with 10

column volumes of buffer A containing 0.02% GDN. Bound proteins were eluted with buffer A

containing 0.02% GDN and 5 mM desthiobiotin. All fractions were analyzed by SDS-

polyacrylamide gel electrophoresis (SDS-PAGE) and Western blotting using anti-His (Qiagen)

and anti-strep (MilliporeSigma) primary antibodies and an IRDye800-coupled anti-mouse

secondary antibody (Rockland) for detection.

The same protocol was used to study the interactions of CesA1 and CesA6 or CesA3 and CesA6

using Strep-tagged CesA1 and His-tagged CesA6, Strep-tagged CesA3 and His-tagged CesA6,

respectively. Control binding experiments were performed by loading the His-tagged CesA

species onto Strep-Tactin sepharose beads and the Strep-tagged CesAs onto the Ni-NTA beads.

In each case, the beads were washed as described above.

Inactivation of CesA

CesA was inactivated by treating CesA trimers with 10 mM sodium tetrathionate overnight at

room temperature (24 °C). The treated sample was purified over a size exclusion column to

remove the excess sodium tetrathionate. Binding experiments with inactivated and untreated

CesAs were performed using Ni-NTA beads as described above.

GmCesA activity assay

Freshly purified or aliquots of flash-frozen enzyme thawed on ice were used for activity assays.

In general, activity assays were performed as described earlier (Carpita, 2011; Purushotham *et al.*,

2020; Wilson *et al.*, 2021). Radiometric quantification of *in vitro* synthesized cellulose was

performed by combining 5 μM GmCesA, 5 mM UDP-glucose (UDP-Glc), and 0.34 μM / 1.25

Ci/L UDP-[^{3}H]-Glc in buffer containing 20 mM Tris-HCl, pH 7.5, 100 mM NaCl, 20 mM MgCl$_2$,

5 mM sodium phosphate, 5 mM sodium citrate, and 1 mM TCEP. The reaction mixtures were

incubated for 45 (3h?) min at 37°C. After incubation, the entire reaction mixture was spotted on
Whatman Grade 3MM chromatography paper. Free substrate was removed by ascending-
descending paper chromatography in 60% ethanol. The radioactivity retained at the origin was
quantified by scintillation counting.

The pH optima of GmCesAs were determined by incubating the CesAs in MMT buffer,
consisting of DL-malic acid, MES and Tris base in the molar ratios 1:2:2-DL-malic
acid:MES:Tris base. The desired pH was adjusted with NaOH or HCl. Activity assays were
performed as mentioned above in technical triplicate from two biological replicates.

Cellulase digestions were performed by adding 5 U of endo-β-1,4-glucanase (*Trichoderma*
longibrachiatum; Megazyme: E-CELTR) directly to the reaction mixture. Following the *in vitro*
synthesis reaction, cellulase treatment was performed for 3 h at 37°C. Cellulose quantification by
scintillation counting was performed as mentioned above to quantify the product.

Steady-state kinetic analyses were performed in triplicate using the UDP-Glo glycosyltransferase
Assay kit (Promega) to monitor the released UDP according to the manufacturer's instructions.
For measuring enzyme kinetics, the reaction mixtures containing 0.33 and 3.3 nM trimeric
*Gm*CesA1 or GmCesA3, respectively, and 0–1 mM UltraPure UDP-Glc (Promega) were added
to 20 mM Tris-HCl buffer, pH 7.5, 100 mM NaCl, 5 mM sodium phosphate, 5 mM sodium
citrate, 20 mM $MgCl_2$, 1 mM TCEP, and 0.02 % GDN and incubated in a final volume of 25 μl
for 1h at 30°C. Afterwards, the reaction mixture was mixed with an equal amount of UDP-Glo
reagent (Promega) in a 96-well Nunclon Delta-Treated flat-bottom microplate (Thermo
Scientific) and incubated for 1 h at room temperature before measuring luminescence using a
GloMax Explorer plate reader (Promega). A standard curve was used for quantification of the
UDP produced. Kinetic values were obtained using the nonlinear regression function in
GraphPad Prism.

_Gm_CesA UDP inhibition assays
UDP inhibition was analyzed using radiometric quantification of *in vitro* synthesized cellulose,
as described above by titrating 0.01-7.5 mM UDP in the reaction. Substrate concentrations for

the individual reactions were 1.4 mM, 0.5 mM, and 2.3 mM for CesA1, CesA3, and CesA6,

respectively. Inhibition constants (IC_{50}) for each CesA were obtained by data analysis in

GraphPad Prism.

Synergistic cellulose biosynthesis

558   Activity synergism between different CesA isoform trimers were studied by mixing two CesA

559   isoforms, one at a constant concentration of 20 μM and one at increasing concentrations from 1-

560   20 μM. The activities of the individual isoforms at the respective concentrations were also

561   measured. *In vitro* synthesized cellulose was quantified by scintillation counting, as described

562   above.

EM grid preparation and data collection

After size exclusion chromatography, the freshly purified protein fractions were pooled. The

protein quality was monitored by negative stain EM. The proteins were diluted to 0.01 mg/mL

and 4 μL was applied to a glow discharged Formvar/Carbon grid (Electron Microscopy Sciences)

for 30 s, followed by 2x washes with 4 μL H_2O. The grid was negatively stained with 4 μL 0.75%

Uranyl Formate (UF) in H_2O for 30 s. Excess UF was removed by blotting with filter paper and

the grid was air dried. Images were taken on a Tecnai F20 at the Macromolecular Electron

Microscopy Core (MEMC) facility at the University of Virginia.

For cryo grid preparation, the protein samples were concentrated to ~3 mg/mL (based on UV

absorbance). Samples of 2.5 µL were applied to a C-flat 300 mesh 1.2/1.3 copper grids (Electron

Microscopy Sciences), glow-discharged in the presence of amylamine at 25 mA for 45 s, and

blotted with a Vitrobot Mark IV (FEI, Thermo Fisher Scientific) with force 4 for 6 s at 4°C, 100%

humidity, and frozen in liquid ethane.

Cryo-EM data were collected at the MEMC at the University of Virginia on a Titan Krios

microscope operated at 300 keV and equipped with a Gatan K3 direct electron detector

positioned post a Gatan Quantum energy filter. On average, a total of ~6000 movies were

collected from one or two grids in counting mode at a magnification of 81K, pixel size of 1.08

A□, and defocus range from -2.2 to -1.2 µm with step size of 0.2 µm. The total dose was 50 e⁻

/Å^2. Movies with 40 frames were collected at 5.17 s/movie rate.

Cryo-EM data processing

Cryo-EM data processing was done in cryoSPARC v4 (Punjani et al., 2017). The general

workflow is described as supplemental information, Fig. S2. Based on improved map qualities,

all volumes were generated imposing threefold symmetry (C3). Maps generated without

symmetry assignment (C1) were used to visualize the unidentified ligand coordinated by the

PCR domains at the particle's symmetry axis. Initial protein models were generated in

AlphaFold and manually adjusted in Chimera and Coot (Emsley and Cowtan, 2004; Pettersen et

al., 2004). Models were refined in Phenix.refine (Afonine et al., 2012). The following regions

were omitted from the constructs due to weak or missing map density: CesA1:1-260 (NTD),

654-717 (CSR), 954-978 (gating loop), 1064-1078 (C term); CesA3: 1-251 (NTD), 648-712

(CSR), 948-974 (gating loop), 1029-1079 (C term and TM 7); and CesA6: 1-248 (NTD), 646-

712 (CSR), 952-971 (gating loop), 1060-1078 (C term). Cellobiose was modeled at the acceptor

positions of CesA3 and CesA6. The corresponding density in the CesA1 map was too weak for

interpretation. Structural representations were generated in ChimeraX or Pymol (Pettersen et al.,

2021; Pymol).

Data availability

PDB coordinates and cryo-EM maps have been deposited at the PDB with accessing codes

8VHZ, 8VHT, and 8VIO.

Funding

This work was supported in part by BASF (CesA cloning, expression, and structure

determination) and the Center for LignoCellulose Structure and Formation (CLSF, biochemical

and interaction studies). CLSF is an Energy Frontier Research Center funded by the U.S.

Department of Energy, Office of Science, Basic Energy Sciences (award DESC0001090). R.H.

and J.Z. are supported by NIH grant R35GM144130 awarded to J.Z. L.W. is supported by the

Howard Hughes Medical Institute of which J.Z. is an investigator.

Author contribution

R.H. prepared all EM grids, performed all EM experiments, performed the CesA kinetic analysis,

and analyzed the interaction between CSR deleted CesA constructs. P.P. purified all CesAs and

performed most biochemical assays. J.Z assisted in data processing and model building. R.H.

performed substrate titration assays and pull-down experiments with the CSR deleted CesA

constructs. L.F.L.W. generated the phylogenetic tree and co-expression analysis and assisted in

substrate titration assays. Y.W. assisted with baculovirus propagation and protein expression. J.Z.

generated the first draft and all authors contributed to manuscript preparation.

Acknowledgements

We are grateful to Kelly Dryden and Michael Purdy of UVA's Macromolecular Electron

Microscopy Core (MEMC) facility for support during EM data collection.

626
627 **Competing interest**

628 The authors declare no conflict of interests exist.

629
630

References

631 **Abidi, W., Zouhir, S., Caleechurn, M., Roche, S., and Krasteva, P.V.** (2021). Architecture and regulation of an enterobacterial cellulose secretion system. Sci Adv **7**10.1126/sciadv.abd8049.

Acheson, J.F., Ho, R., Goulart, N., Cegelski, L., and Zimmer, J. (2021). Molecular organization of the E. coli cellulose synthase macrocomplex. Nat Struct Mol Biol **28**:310-318.

Afonine, P.V., Grosse-Kunstleve, R.W., Echols, N., Headd, J.J., Moriarty, N.W., Mustyakimov, M., Terwilliger, T.C., Urzhumtsev, A., Zwart, P.H., and Adams, P.D. (2012). Towards automated crystallographic structure refinement with phenix.refine. Acta Crystallogr D Biol Crystallogr **68**:352-367. 10.1107/S0907444912001308.

Baytshtok, V., Chen, J.J., Glynn, S.E., Nager, A.R., Grant, R.A., Baker, T.A., and Sauer, R.T. (2017). Covalently linked HslU hexamers support a probabilistic mechanism that links ATP hydrolysis to protein unfolding and translocation. Journal of Biological Chemistry **292**:5695-5704. 10.1074/jbc.M116.768978.

Carpita, N.C. (2011). Update on mechanisms of plant cell wall biosynthesis: how plants make cellulose and other (1->4)-beta-D-glycans. Plant Physiol **155**:171-184. 10.1104/pp.110.163360.

Cosgrove, D.J., Dupree, P., Gomez, E.D., Haigler, C.H., Kubicki, J.D., and Zimmer, J. (2024). How Many Glucan Chains Form Plant Cellulose Microfibrils? A Mini Review. Biomacromolecules **25**:6357-6366. 10.1021/acs.biomac.4c00995.

Darriba, D., Taboada, G.L., Doallo, R., and Posada, D. (2011). ProtTest 3: fast selection of best-fit models of protein evolution. Bioinformatics **27**:1164-1165. 10.1093/bioinformatics/btr088.

Das, D., Walvoort, M.T.C., Lukose, V., and Imperiali, B. (2016). A Rapid and Efficient Luminescence-based Method for Assaying Phosphoglycosyltransferase Enzymes. Scientific Reports **6**ARTN 33412 10.1038/srep33412.

Du, J., Vepachedu, V., Cho, S.H., Kumar, M., and Nixon, B.T. (2016). Structure of the Cellulose Synthase Complex of Gluconacetobacter hansenii at 23.4 A Resolution. PloS one **11**:e0155886.

Eddy, S.R. (2011). Accelerated Profile HMM Searches. Plos Computational Biology **7**ARTN e1002195 10.1371/journal.pcbi.1002195.

Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. Acta Crystallogr D Biol Crystallogr **60**:2126-2132. 10.1107/S0907444904019158.

Fagard, M., Desnos, T., Desprez, T., Goubet, F., Refregier, G., Mouille, G., McCann, M., Rayon, C., Vernhettes, S., and Höfte, H. (2000). PROCUSTE1 encodes a cellulose synthase required for normal cell elongation specifically in roots and dark-grown hypocotyls of Arabidopsis. Plant Cell **12**:2409-2423. DOI 10.1105/tpc.12.12.2409.

Fuxreiter, M. (2012). Fuzziness: linking regulation to protein dynamics. Mol Biosyst **8**:168-177. 10.1039/c1mb05234a.

Gonneau, M., Desprez, T., Guillot, A., Vernhettes, S., and Hofte, H. (2014). Catalytic subunit stoichiometry within the cellulose synthase complex. Plant Physiol **166**:1709-1712. 10.1104/pp.114.250159.

Gow, L.A., and Selitrennikoff, C.P. (1984). Chitin Synthetase of Neurospora-Crassa - Inhibition by Nikkomycin, Polyoxin-B, and Udp. Curr Microbiol **11**:211-216. Doi 10.1007/Bf01567162.

Herth, W., and Weber, G. (1984). Occurrence of the Putative Cellulose-Synthesizing Rosettes in the Plasma-Membrane of Glycine-Max Suspension-Culture Cells. Naturwissenschaften **71**:153-154. Doi 10.1007/Bf01137780.

Katoh, K., and Standley, D.M. (2013). MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Molecular Biology and Evolution **30**:772-780.
10.1093/molbev/mst010.
Kimura, S., Laosinchai, W., Itoh, T., Cui, X., Linder, C.R., and Brown, R.M.J. (1999). Immunogold
labeling of rosette terminal cellulose-synthesizing complexes in the vascular plant Vigna
angularis. Plant Cell **11**:2075-2086. 10.1105/tpc.11.11.2075.
Kubicki, J.D., Yang, H., Sawada, D., O'Neill, H., Oehme, D., and Cosgrove, D. (2018). The Shape
of Native Plant Cellulose Microfibrils. Sci Rep **8**:13983. 10.1038/s41598-018-32211-w.
Kumar, M., Wightman, R., Atanassov, I., Gupta, A., Hurst, C.H., Hemsley, P.A., and Turner, S.
(2016). S-Acylation of the cellulose synthase complex is essential for its plasma membrane
localization. Science **353**:166-169. 10.1126/science.aaf4009.
Kurek, I., Kawagoe, Y., Jacob-Wilk, D., Doblin, M., and Delmer, D. (2002). Dimerization of
cotton fiber cellulose synthase catalytic subunits occurs via oxidation of the zinc-binding
domains. Proc Natl Acad Sci U S A **99**:11109-11114. 10.1073/pnas.162077099.
Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H.,
Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., et al. (2007). Clustal W and Clustal X version 2.0.
Bioinformatics **23**:2947-2948. 10.1093/bioinformatics/btm404.
Lombard, V., Ramulu, H.G., Drula, E., Coutinho, P.M., and Henrissat, B. (2014). The
carbohydrate-active enzymes database (CAZy) in 2013. Nucleic Acids Res **42**:D490-D495.
10.1093/nar/gkt1178.
Maloney, F.P., Kuklewicz, J., Corey, R.A., Bi, Y.C., Ho, R.Y., Mateusiak, L., Pardon, E., Steyaert,
J., Stansfeld, P.J., and Zimmer, J. (2022). Structure, substrate recognition and initiation of
hyaluronan synthase. Nature **604**:195-201. 10.1038/s41586-022-04534-2.
Massenburg, L., Nixon, B.T., O''Neill, H., and Willaims, A.N. (2024). Understanding the
Structural Ballet of Cellulose Formation – Using CryoEM to Reveal the Role of Protein Flexibility
on Higher-order Oligomer Formation, Glucan Catalysis, and
Cellulose Microfibril Extrusion. Micros Microanal **30**:714-718.
McNamara, J.T., Morgan, J.L.W., and Zimmer, J. (2015). A molecular description of cellulose
biosynthesis. Annu Rev Biochem **84**:17.11-17.27.
Morgan, J., Strumillo, J., and Zimmer, J. (2013). Crystallographic snapshot of cellulose synthesis
and membrane translocation. Nature **493**:181-186.
Newman, R.H., Hill, S.J., and Harris, P.J. (2013). Wide-angle x-ray scattering and solid-state
nuclear magnetic resonance data combined to test models for cellulose microfibrils in mung
bean cell walls. Plant Physiol **163**:1558-1567. 10.1104/pp.113.228262.
Nixon, B.T., Mansouri, K., Singh, A., Du, J., Davis, J.K., Lee, J.G., Slabaugh, E., Vandavasi, V.G.,
O'Neill, H., Roberts, E.M., et al. (2016). Comparative Structural and Computational Analysis
Supports Eighteen Cellulose Synthases in the Plant Cellulose Synthesis Complex. Sci Rep
6:28696. 10.1038/srep28696.
Obayashi, T., Hibara, H., Kagaya, Y., Aoki, Y., and Kinoshita, K. (2022). ATTED-II v11: A Plant
Gene Coexpression Database Using a Sample Balancing Technique by Subagging of Principal
Components. Plant and Cell Physiology **63**:869-881. 10.1093/pcp/pcac041.
Omadjela, O., Narahari, A., Strumillo, J., Mélida, H., Mazur, O., Bulone, V., and Zimmer, J.
(2013). BcsA and BcsB form the catalytically active core of bacterial cellulose synthase sufficient

for in vitro cellulose synthesis. Proc Natl Acad Sci U S A **110**:17856-17861.
10.1073/pnas.1314063110.
Persson, S., Paredez, A., Carroll, A., Palsdottir, H., Doblin, M., Poindexter, P., Khitrov, N., Auer,
M., and Somerville, C.R. (2007). Genetic evidence for three unique components in primary cell-
wall cellulose synthase complexes in Arabidopsis. Proc Natl Acad Sci USA **104**:15566-15571. DOI
10.1073/pnas.0706592104.
Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and
Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory research and analysis. J
Comput Chem **25**:1605-1612. 10.1002/jcc.20084.
Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H.,
and Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. Protein science : a publication of the Protein Society **30**:70-82. 10.1002/pro.3943.
Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. Nat Methods **14**:290-296.
10.1038/nmeth.4169.
Purushotham, P., Ho, R., and Zimmer, J. (2020). Architecture of a catalytically active
homotrimeric plant cellulose synthase complex. Science **369**:1089-1094.
Purushotham, P., Cho, S.H., Diaz-Moreno, S.M., Kumar, M., Nixon, B.T., Bulone, V., and
Zimmer, J. (2016). A single heterologously expressed plant cellulose synthase isoform is
sufficient for cellulose microfibril formation in vitro. Proc Natl Acad Sci U S A **113**:11360-11365.
10.1073/pnas.1606210113.
Pymol The PYMOL Molecular Graphics System.
Qiao, Z., Lampugnani, E.R., Yan, X.F., Khan, G.A., Saw, W.G., Hannah, P., Qian, F., Calabria, J.,
Miao, Y., Gruber, G., et al. (2021). Structure of Arabidopsis CESA3 catalytic domain with its
substrate UDP-glucose provides insight into the mechanism of cellulose synthesis. Proc Natl
Acad Sci U S A **118**10.1073/pnas.2024015118.
Sampathkumar, A., Peaucelle, A., Fujita, M., Schuster, C., Persson, S., Wasteneys, G.O., and
Meyerowitz, E.M. (2019). Primary wall cellulose synthase regulates shoot apical meristem
mechanics and growth. Development **146**10.1242/dev.179036.
Scavuzzo-Duggan, T.R., Chaves, A.M., Singh, A., Sethaphong, L., Slabaugh, E., Yingling, Y.G.,
Haigler, C.H., and Roberts, A.W. (2018). Cellulose synthase "class specific regions' are
intrinsically disordered and functionally undifferentiated. J Integr Plant Biol **60**:481-497.
10.1111/jipb.12637.
Sethaphong, L., Haigler, C., Kubicki, J., Zimmer, J., Bonetta, D., DeBolt, S., and Yingling, Y.
(2013). Tertiary model of a plant cellulose synthase. Proc Natl Acad Sci U S A **110**:7512-7517.
10.1073/pnas.1301027110.
Singh, A., Kwansa, A.L., Kim, H.S., Williams, J.T., Yang, H., Li, N.K., Kubicki, J.D., Roberts, A.W.,
Haigler, C.H., and Yingling, Y.G. (2020). In silico structure prediction of full-length cotton
cellulose synthase protein (GhCESA1) and its hierarchical complexes. Cellulose **27**:5597-5616.
Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics **30**:1312-1313. 10.1093/bioinformatics/btu033.
Taylor, N.G., Howells, R.M., Huttly, A.K., Vickers, K., and Turner, S.R. (2003). Interactions
among three distinct CesA proteins essential for cellulose synthesis. Proc Natl Acad Sci USA
100:1450-1455. 10.1073/pnas.0337628100.

Tie, J.K., Jin, D.Y., Loiselle, D.R., Pope, R.M., Straight, D.L., and Stafford, D.W. (2004). Chemical
modification of cysteine residues is a misleading indicator of their status as active site residues
in the vitamin K-dependent γ-glutamyl carboxylation reaction. J Biol Chem **279**:54079-54087.
10.1074/jbc.M408945200.
Timmers, J., Vernhettes, S., Desprez, T., Vincken, J.P., Visser, R.G., and Trindade, L.M. (2009).
Interactions between membrane-bound cellulose synthases involved in the synthesis of the
secondary cell wall. FEBS Lett **583**:978-982. 10.1016/j.febslet.2009.02.035.
Tlapak-Simmons, V.L., Baron, C.A., and Weigel, P.H. (2004). Characterization of the purified
hyaluronan synthase from Streptococcus equisimilis. Biochemistry **43**:9234-9242.
10.1021/bi049468v.
Turner, S., and Kumar, M. (2018). Cellulose synthase complex organization and cellulose
microfibril structure. Philos T R Soc A **376**ARTN 20170048 10.1098/rsta.2017.0048.
Turner, S.R., and Somerville, C.R. (1997). Collapsed xylem phenotype of Arabidopsis identifies
mutants deficient in cellulose deposition in the secondary cell wall. Plant Cell **9**:689-701. DOI
10.1105/tpc.9.5.689.
Van Bel, M., Silvestri, F., Weitz, E.M., Kreft, L., Botzki, A., Coppens, F., and Vandepoele, K.
(2022). PLAZA 5.0: extending the scope and power of comparative and functional genomics in
plants. Nucleic Acids Research **50**:D1468-D1474. 10.1093/nar/gkab1024.
Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M., and Barton, G.J. (2009). Jalview
Version 2--a multiple sequence alignment editor and analysis workbench. Bioinform **25**:1189-
1191. 10.1093/bioinformatics/btp033.
Wilson, T.H., Kumar, M., and Turner, S.R. (2021). The molecular basis of plant cellulose
synthase complex organisation and assembly. Biochem Soc T **49**:379-391.
10.1042/Bst20200697.
Zhang, X.N., Xue, Y., Guan, Z.Y., Zhou, C., Nie, Y.F., Men, S., Wang, Q., Shen, C.C., Zhang, D.L.,
Jin, S.X., et al. (2021a). Structural insights into homotrimeric assembly of cellulose synthase
CesA7 from Gossypium hirsutum. Plant Biotechnol J **19**:1579-1587. 10.1111/pbi.13571
Zhang, Y., Yu, J.Y., Wang, X., Durachko, D.M., Zhang, S.L., and Cosgrove, D.J. (2021b).
Molecular insights into the complex mechanics of plant epidermal cell walls. Science **372**:706-+.
10.1126/science.abf2824.
791
792

Figures and figure legends

Fig. 1: GmCesA Phylogeny and co-expression analysis

Left: Maximum-likelihood phylogeny of CesA protein sequences from soy, Arabidopsis, cotton, poplar, tomato, Physcomitrium and Amborella. For alignment, the Pfam-defined 'Cellulose synthase' domain was extracted from each sequence using HMMER. The final phylogeny was calculated using RAxML with 100 rapid bootstrap pseudo-replicates. Arrowheads mark the positions of *Gm*CesA1, *Gm*CesA3, and *Gm*CesA6 within the tree; structurally characterized proteins (*Ptt*CesA8 and *Gh*CesA7) and Arabidopsis sequences are also labelled. Branch lengths correspond to average number of substitutions per site (relative to scale bar); branch labels report bootstrap successes for each split. Right: Co-expressed gene networks for *Gm*CesA1, *Gm*CesA3, and *Gm*CesA6 from ATTED-II v11 (https://atted.jp). Relevant functional annotations for co-expressed cell wall genes are labelled by color (cellulose synthesis: yellow; pectin synthesis: orange; β-galactoglucomannan synthesis: violet; arabinogalactan proteins: purple).

811
812

Figure 2. Functional characterization of *Glycine max* primary cell wall CesAs. From (**A-C**) analytical size exclusion chromatography (Superose 6 Increase) of *Gm*CesA1 (A), *Gm*CesA3 (B), and *Gm*CesA6 (C). Void volume (*Vo*) and trimer and monomer peaks are marked. A rerun of the trimer fraction for each species is shown as a dashed profile. Inset: Coomassie-stained SDS-polyacrylamide gel electrophoresis of the indicated elution volumes. **D)** Catalytic activity of the purified CesAs. [3]H-labeled cellulose synthesized by trimeric and monomeric species was degraded with cellulase, followed by quantification by scintillation counting. (DS) and (PS) indicate cellulase treatments during and after the synthesis reaction, respectively. DPM: disintegrations per minute. **E)** pH optima for catalytic activity of CesA1, CesA3 and CesA6.

822   Activities are normalized to the highest activity for each isoform. **F)** UDP inhibits CesAs.

823   Cellulose biosynthesis was performed in the presence of 1.4, 0.5, and 2.3 mM UDP-Glc for

824   GmCesA1, CesA3, and CesA6, respectively, as well as the indicated increasing concentrations of

825   UDP. Product yields in the absence of UDP were set as 100 %. Error bars in panels D-F

826   represent deviations from the means of at least three replicas.

Figure 3. Soybean primary cell wall CesAs assemble into homotrimers.

A) CryoEM maps of the CesA homotrimers contoured at 4.5-5.6 σ. One subunit is shown in color, the others are shown in light and dark gray. The gray background indicates the estimated membrane boundaries. **B)** Cartoon representation of a CesA6 protomer. The transmembrane region is shown in green and dark pink, interface helices (IF) are shown in orange, and the catalytic domain is colored gray. The PCR and CSR regions are shown in blue and yellow, respectively. **C)** Comparison of CesA6 and CesA8. GmCesA6 is shown as a cartoon that is

overlaid with a semitransparent surface of hybrid aspen CesA8 (surface, PDB: 6WLB).

Transmembrane helix 7 is colored light and dark pink for CesA8 and CesA6, respectively. The

black triangle indicates the 3-fold symmetry axis of the homotrimer. Zoom views: Surface

representations of CesA6 (left) and CesA8 (right) highlighting the lateral window. $TM7_b$ refers

to TM helix 7 of another protomer. The view is from the 3-fold symmetry axis towards a CesA

protomer. **D)** CryoEM map of the CesA1 trimer shown at a low contour level (1.4 σ). The CesA1

structure is shown as a cartoon with one protomer colored yellow. The resolved CSR N- and C-

terminal regions are colored blue and red, respectively. **E)** Sequence alignment of the CSR

regions of CesA1, CesA3 and CesA6 generated in Clustal Omega (Larkin et al., 2007).

Figure 4. *In vitro* **interactions between different CesA isoforms.**

Tandem pull-down experiments using Ni-NTA and Strep-Tactin resin. Experiments were

performed with homotrimers of the indicated CesA isoforms tagged N-terminally either with

His- or TwinStrep-tags. Material eluted from the Ni-NTA resin was loaded onto the Strep-Tactin

beads. Top panels: Coomassie stained SDS-PAGE, bottom panels: Western blots using anti

penta-His or anti-Strep primary antibodies. **A-C)** Trimer-trimer interaction between CesA1 and

CesA3, CesA6 and CesA3, and CesA6 and CesA1, respectively. **D)** Differently tagged

homotrimers of the same isoform do not interact. Tandem purification of a mixture of His- and

TwinStrep-tagged CesA1. **E-H)** Control binding of His-tagged CesAs to StrepTactin beads and

Strep-tagged CesAs to Ni-NTA resin. F, W, E: Flow through, wash, and eluted fractions.

Figure 5. The CSR mediates trimer-trimer interactions.

Isoform interactions are independent of the NTD. **A)** Activity comparison of full-length and N-terminally truncated CesA isoforms. DPM: disintegrations per minute. **B)** Tandem pull-down experiments as in *Figure 4* but with N-terminally truncated homotrimers of CesA1 and CesA3. Top panel: SDS-PAGE, bottom panel: Western blots using anti-His and anti-Strep primary antibodies. **C)** Control binding of His-ΔNCesA1 to Strep-Tactin beads and Strep-ΔNCesA3 to Ni-NTA resin. L, Ft, W, E: Load, Flow through, Wash, and Eluted fractions. **D)** AlphaFold predicted model of N-terminally truncated CesA1 with the CSR replaced by a loop shown as an orange backbone (left), and negative stain images of the NTD and CSR truncated CesA1 trimer. E) Tandem purification of NTD and CSR truncated CesA1 and CesA3. **F)** Affinity purification of His-tagged full-length CesA1 and Strep-tagged NTD and CSR truncated CesA3. Shown is an SDS-PAGE after Coomassie staining.

Figure 6. Clustering of CesA homotrimers and a dimer of trimer.

A-C) Uranyl formate-stained EM images of homotrimers of purified GmCesA1 (A), GmCesA3 (B) and GmCesA6 (C). The proteins were incubated overnight on ice prior to grid preparation. **D)** The same for an equimolar mixture of all three CesA isoforms, incubated overnight, separated from individual trimers by size exclusion chromatography, and imaged by negative stain EM. Selected clusters are encircled. Scale bar: 100 nm. **E)** Purified trimers of CesA1 and CesA3 were combined and used for cryo-EM analysis. Shown are 2D class averages of dimers of trimers. **F and G)** Manually assembled CesA1 trimer volumes (F) were used to calculate 2D class average templates (G) for comparison with the experimentally obtained dimers of trimers shown in panel (E).

Figure 7. Synergistic catalytic activity.

Cellulose biosynthesis from mixtures of CesA isoform homotrimers. The formation of ^{3}H-

labeled cellulose was quantified by scintillation counting for reaction mixtures containing one

CesA isoform at 20 µM concentration and the other isoform at the indicated increasing

concentrations. Blue, magenta and green columns represent activities measured for the individual

single isoforms alone. Gray columns represent the calculated theoretical activities for the isoform

mixtures by adding the individually determined activities. Red columns represent the

experimentally determined activities for the isoform mixtures. **A and B)** CesA1$_{20\ \mu M}$ + CesA3$_{1-\ 20}$

$_{\mu M}$ and CesA3$_{20\ \mu M}$ + CesA1$_{1-\ 20\ \mu M}$; **C and D)** CesA3$_{20\ \mu M}$ + CesA6$_{1-\ 20\ \mu M}$ and CesA6$_{20\ \mu M}$ +

CesA3$_{1-\ 20\ \mu M}$; and **E and F**) CesA1$_{20\ \mu M}$ + CesA6$_{1-\ 20\ \mu M}$ and CesA6$_{20\ \mu M}$ + CesA1$_{1-\ 20\ \mu M}$,

respectively. **G)** The same as for panel (a) but titrating the same CesA isoform (CesA1$_{20\ \mu M}$ +

CesA1$_{1-\ 20\ \mu M}$). **H)** The same as for panels a-f but for a combination of all three CesA isoforms.

Individual and combined activities were determined at a concentration of 6.6 µM for each CesA

isoform. DPM, disintegrations per minute.

Figure 8. CSC models consisting of different CesA isoforms.

A) Association of homotrimers of three different CesA isoforms. Isoforms are indicated by different colors. The shapes represent the cytosolic CesA domains. **B and C)** Alternative models of CSC assembly from heterotrimeric CesA complexes that have not been detected *in vitro*. Model (B) would require interactions between the same CesA isoforms at the center, also not observed *in vitro*. PCR: Plant conserved region, GT: Glycosyltransferase, CSR: Class specific region.

937   **Supplemental Figure 1-8**

938   **Supplemental Table 1**

944
945

Figure 1. Sequence alignment of soybean CesA1, CesA3, and CesA6 and substrate turnover

kinetics. A) Sequences were aligned in Clustal Omega (Larkin *et al.*, 2007) and visualized in

Jalview (Waterhouse et al., 2009) showing sequence identity from dark blue (high) to white

(low). **B**) Gene expression profiles of soybean GmCesA1: Glyma.06G069600, GmCesA3:

Glyma.12G237000, and GmCesA6: Glyma.02G080900 from the soybean eFP Browser

(https://bar.utoronto.ca/efpsoybean/cgi-bin/efpWeb.cgi). **C**) In vitro Michaelis Menten kinetic

analysis of CesA1 and CesA3 by titrating UDP-Glc and quantifying the generated UDP using an

UDP-Glo assay kit. The data is normalized to the highest activity of GmCesA1. The activity of

GmCesA6 was too weak to be analyzed by this method. Error bars represent the standard

deviations from the means of three technical replicas.

958
959 **Figure 2. Cryo-EM data processing workflows**.

960 All steps were performed in CryoSparc v4 (Punjani *et al.*, 2017).

961
962 **Figure 3. Cryo-EM map quality examples.**

963 **A)** Selected helical segments of CesA1, CesA3, and CesA6. TM helix 7 was excluded from the

964 GmCesA3 model due to limiting map quality. **B)** Close-up views of the unidentified ligands

965 coordinated by the PCR domains of the homotrimers.

Figure 4. Interactions of CesA homotrimers of the same isoforms.

A-D) Tandem purifications of combinations of His- and Strep-tagged CesA3 or CesA6 (A and B) together with the non-specific binding controls (C and D). (E and F) Interactions of monomers of different CesA isoforms. **E)** Monomeric versions of His-tagged CesA1 and Strep-tagged CesA3 were subjected to tandem purification starting either with Ni-NTA resin or Strep-Tactin beads. **F)** Controls to account for non-specific interactions.

978
979
Figure 5. Control synergistic activity assays by titrating the same CesA isoforms.

A) CesA3$_{20\,\mu M}$ + CesA3$_{1-20\,\mu M}$, **B)** CesA6$_{20\,\mu M}$ + CesA6$_{1-20\,\mu M}$. Gray columns indicate calculated theoretical activities from individual measurements, red columns are experimentally determined activities. DPM: Disintegrations per minute.

Figure 6. Tetrathionate-inactivated CesAs interact with other CesA isoforms.

A-C) Co-purification of catalytically active CesA homotrimers with tetrathionate inactivated CesA homotrimers (indicated by a .T extension). The indicated combinations of CesAs were purified over Ni-NTA beads. Results were analyzed by Coomassie stained SDS-PAGE (top panels) and Western blotting (bottom panels). **D-F)** Control binding experiments of Strep-tagged CesAs and Ni-NTA resin. **G)** Catalytic activity of tetrathionate treated GmCesA3.

Figure 7. Synergistic cellulose biosynthesis with tetrathionate inactivated CesA trimers.

A-F) Inactivated CesA trimers (indicated by a '.T' extension) at a constant concentration were incubated with increasing concentrations of wild type CesAs of a different isoform. Background activities of the inactivated CesAs range from ~200-400 DPM. Calculated total theoretical activities are shown as gray columns, experimentally determined activities are shown as red columns. DPM: disintegrations per minute.

1001
1002

Figure 8. Attempt to purify hetero-oligomeric CesA oligomers. GmCesA1, GmCesA3, and

GmCesA6 were co-expressed in Sf9 cells and purified by sequential affinity chromatography. **A**)

Material eluted (Elu) from a Ni-NTA column was purified over Strep-Taction affinity resin. **B**)

The Elu fraction from the Strep-Tactin resin was loaded onto a Superose-6 size exclusion

chromatography column (SEC). The inset shows a Coomassie stained SDS-PAGE of the

indicated fractions. **C**) The peak fraction from (B) was loaded onto an anti-Flag affinity matrix,

washed (W), and eluted. **D**) Western blot analysis of eluting (Elu), flow through (Ft), and SEC (4,

5) fractions from all steps shows the presence of all CesA species, alongside significant loss in

all Ft fractions.

1012

Supplemental Table 1. Cryo-EM data collection, refinement, and validation statistic.

	GmCesA1 (EMDB-43244) (PDB: 8VHZ)	GmCesA3 (EMDB-43241) (PDB: 8VHT)	GmCesA6 (EMDB-43245) (PDB 8VI0)
Data collection and processing			
Microscope	Titan Krios	Titan Krios	Titan Krios
Camera	GIF K3	GIF K3	GIF K3
Magnification	81,000	81,000	81,000
Voltage (kV)	300	300	300
Electron exposure (e–/Å^2)	50	50	50
Defocus range (μm)	-1.1 to -2.2	-1.1 to -2.2	-1.1 to -2.2
Pixel size (Å)	1.08	1.08	1.08
Symmetry imposed	C3	C3	C3
Initial particle images (no.)			
Final particle images (no.)	181,101	237,568	192,666
Map resolution (Å)	3.3	3.2	3.0
FSC threshold	0.143	0.143	0.143
Map resolution range (Å)	3.3 -	3.2 -	3.0 -
Refinement			
Initial model used	AlphaFold generated	AlphaFold generated	AlphaFold generated
Map sharpening B factor (Å^2)	-117.5	-105.1	-92.4
Model composition			
Non-hydrogen atoms	17190	16644	17694
Protein residues	2139	2064	2172
Ligand		BGC	BGC
Mean B factors (Å^2)			
Protein	39.47	56.40	38.03
Ligand		85.27	72.40
R.m.s. deviations			
Bond lengths (Å)	0.003	0.002	0.002
Bond angles (°)	0.603	0.53	0.499
Ramachandran plot			
Favored (%)	96.88	97.05	97.7
Allowed (%)	3.12	2.95	2.3
Disallowed (%)	0.00	0.00	0.00
Validation			
MolProbity score	1.52	1.42	1.58
Clashscore	6.32	4.91	6.87
Poor rotamers (%)	0.16	0.00	0.00
Overall correlation coefficients			
CC (mask)	0.76	0.82	0.85
CC (peaks)	0.62	0.79	0.74
CC (volume)	0.74	0.66	0.83