Different subregions of monkey lateral prefrontal cortex respond to abstract

sequences and their components

Authors: Nadira Yusif Rodriguez[1,^], Aarit Ahuja[1], Debaleena Basu[1,2], Theresa H. McKim[3], and

Theresa M. Desrochers*[1,4,5]

Affiliations:

1. Department of Neuroscience, Brown University

2. Department of Biosciences and Bioengineering, IIT Bombay, Mumbai, Maharashtra, India

3. Department of Biology & Institute for Neuroscience, University of Nevada, Reno

4. Department of Psychiatry and Human Behavior, Brown University

5. Robert J. and Nancy D. Carney Institute for Brain Sciences, Brown University

*Corresponding Author: Theresa M. Desrochers, theresa_desrochers@brown.edu

^N.Y.R.'s present address: Massachusetts General Hospital, Athinoula A. Martinos Center for

Biomedical Imaging, Charlestown, MA.

Abbreviated title: Monkey LPFC regions respond to sequence components

Counts: 5 figures, 11 tables

Words: Abstract, 248; Introduction, 952; Discussion, 1,474

Conflict of Interest Statement: The authors declare no competing financial interests.

Acknowledgements

This study was supported by the National Science Foundation (NSF) Established Program to Stimulate Competitive Research (EPSCoR) Neural Basis of Attention Grant 1632738 (N.Y.R. and T.M.D.), the National Institute of General Medical Sciences (NIGMS)-National Institutes of Health (NIH) Initiative to Maximize Student Development Grant IMSD R25GM083270 (N.Y.R.), the NIH-NIGMS Grant COBRE P20GM103645 (T.M.D), NIH National Institutes of Mental Health (NIMH) R21MH125010 (T.M.D.), NSF Faculty Early Career Development (CAREER) Program Award BCS-2143656 (T.M.D.), NIMH Research Project Grant R01MH131615 (T.M.D.), and the Carney Institute for Brain Science Innovation Award (T.M.D.). Part of this research was conducted using computational resources and services at the Center for Computation and Visualization, Brown University (NIH Grant S10OD025181).

We thank Matthew Maestri for his assistance with animal training and data collection. We thank Dr. Michael Worden, Lynn Fanella, Fabienne McEleney, and Brown University's MRI Facilities staff for their support and guidance throughout this project. We thank Dr. Lucija Jankovic-Rapan, Dr. Nicola Palomero-Gallagher and Dr. Seán Froudist-Walsh for their assistance and sharing of the MEBRAINS atlas and regions of interest used for analysis. We also thank Dr. David Sheinberg, Dr. Amitai Shenhav, Dr. Katherine Conen, and Hannah Doyle for their continued support throughout this project and preparation of this publication as well as members of both the Sheinberg and the Desrochers Labs for many helpful discussions and contributions.

Author Contributions

Nadira Yusif Rodriguez: Conceptualization, Investigation, Formal analysis, Writing - Original

Draft, Writing - Review & Editing, Visualization. Aarit Ahuja: Investigation, Writing - Review

& Editing. Debaleena Basu: Conceptualization, Investigation, Writing - Review & Editing.

Theresa H. McKim: Conceptualization, Investigation, Writing - Review & Editing. Theresa M.

Desrochers: Conceptualization, Methodology, Investigation, Writing - Original Draft, Writing -

Review & Editing, Supervision, Resources, Funding acquisition.

<u>Abstract</u>

Sequential information permeates daily activities, such as when watching for the correct series of

buildings to determine when to get off the bus or train. These sequences include periodicity (the

spacing of the buildings), the identity of the stimuli (the kind of house), and higher-order more

abstract rules that may not depend on the exact stimulus (e.g. house, house, house, business).

Previously, we found that the posterior fundus of area 46 in the monkey lateral prefrontal cortex

(LPFC) responds to rule changes in such abstract visual sequences. However, it is unknown if

this region responds to other components of the sequence, i.e., image periodicity and identity, in

isolation. Further, it is unknown if this region dissociates from other, more ventral LPFC

subregions that have been associated with sequences and their components. To address these

questions, we used awake functional magnetic resonance imaging in three male macaque

monkeys during two no-report visual tasks. One task contained abstract visual sequences, and the

other contained no visual sequences but maintained the same image periodicity and identities.

We found the fundus of area 46 responded only to abstract sequence rule violations. In contrast,

the ventral bank of area 46 responded to changes in image periodicity and identity, but not

changes in the abstract sequence. These results suggest a functional specialization within

anatomical substructures of LPFC to signal different kinds of stimulus regularities. This

specialization may provide key scaffolding to identify abstract patterns and construct complex

models of the world for daily living.

Significance Statement

Daily tasks, such as a bus commute, require tracking or monitoring your place (same, same,

same, different building) until your stop. Sequence components such as rule, periodicity (timing),

and item identity are involved in this process. While prior work located responses to sequence

rule changes to area 46 of monkey lateral prefrontal cortex (LPFC) using awake monkey fMRI,

less was known about other components. We found that LPFC subregions differentiated between

sequence components. Area 46 posterior fundus responded to abstract visual sequence rule

changes, but not to changes in image periodicity or identity. The converse was true for the more

ventral, adjacent shoulder region. These results suggest that interactions between adjacent LPFC

subregions provide key scaffolding for complex daily behaviors.

Introduction

When commuting to work, such as when taking a bus, you may internally track house, house, house, store (same, same, same, different building) until you arrive at your stop. This recognition illustrates an essential process: the monitoring of abstract sequences. These sequences are abstract because they do not depend on the identity of the individual stimuli (e.g., changing the color of the house). The same system that facilitates sequential tracking also enables the detection of changes or deviations to an existing sequence. Sequences are defined by multiple elements that we specifically refer to as components. These sequential components include item identity, periodicity (temporal structure), and rule. Abstract sequence rule deviations may encompass changes in these individual sequence components. How does the brain track changes to sequences and their components?

We identified a brain region that responds to abstract sequence deviations, but whether the same or other regions also monitor changes in sequential components has not been tested. Previously, we identified a specific subregion within the fundus of posterior area 46 of the lateral prefrontal cortex (LPFC) as uniquely responding to sequential changes using awake monkey functional magnetic resonance imaging (fMRI) (Yusif Rodriguez et al., 2023). Beyond the rule that these abstract sequences followed, they had two (previously mentioned) main components: image identity and periodicity. These components were controlled for when determining responses to infrequent changes in the abstract visual sequence. However, LPFC is also known to respond to changes in image identity or periodicity. Responses to infrequent (sometimes referred to as "oddball") stimuli have been reported in monkey LPFC using an array of techniques (Chao et al., 2018; Camalier et al., 2019; Grohn et al., 2020). Changes in stimulus periodicity could also elicit responses in LPFC because responses in LPFC are modulated by the duration preceding the

auditory or visual stimulus (Onoe et al., 2001; Genovesio et al., 2006; Chiba et al., 2021).

Therefore, the question arises as to whether the same, abstract sequence coding, or other LPFC

subregions respond to changes in image periodicity or identity alone.

The ventral LPFC (VLPFC) is a prime candidate as another subregion within the LPFC

that could respond to sequential components. FMRI studies in macaques have shown VLPFC

activity during auditory sequential tasks and sequence deviants (Wang et al., 2015; Vergnieux

and Vogels, 2020). Studies using electrophysiology provided evidence for the representation of

generalizable sequential structures and changes to these structures in neuronal population

responses within VLPFC (Esmailpour et al., 2023; Bellet et al., 2024). VLPFC also responds to

non-sequential information that shares similar features with sequential tasks, such as prediction

error (Uhrig et al., 2014; Chao et al., 2018) and responses to infrequent ("oddball") items (Uhrig

et al., 2014; Suda et al., 2022). VLPFC is often observed as more directly representing sensory

visual information (compared to DLPFC, which can be more spatial or action oriented).

Responses in VLPFC for non-spatial object-based features include color, shape, and object type

(Meyer et al., 2011; Yamagata et al., 2012; Tang et al., 2021; Xu et al., 2022). Thus, there is

evidence to suggest that VLPFC could respond to sequential components and abstract sequences,

underscoring the importance for dissociating between LPFC subregions.

To dissociate between LPFC subregions, we defined a more ventral, yet adjacent brain

area. Historically, many conventions have been applied to naming subregions within the LPFC

(Walker, 1940; Petrides and Pandya, 1999; Rapan et al., 2023). While some studies refer to

VLPFC as the region ventral to the arcuate sulcus, primarily Brodmann area 44 and potentially

parts of 6VR (Rapan et al., 2023), others refer to the region ventral to the principal sulcus and

dorsal to the arcuate sulcus that can include Brodmann areas 46, 9/46, 45A, and 45B along with

more anterior territory such as Brodmann areas 47 and 12 (Rapan et al., 2023). Our purpose here

is not to adjudicate among the definitions, but take advantage of a commonality that, in general,

more ventral regions of the LPFC are more biased towards sensory, object-based, or non-spatial

information (Meyer et al., 2011; Yamagata et al., 2012; Tang et al., 2021; Xu et al., 2022).

Therefore, an area that is more ventral, yet adjacent to the subregion in area 46 where we

previously observed sequential responses is an ideal candidate for comparison. A region defined

as thus would not make assumptions about naming conventions and still be clearly within the

LPFC. Using this region, we determined if there were differences in abstract sequence or

component representation between the two LPFC sub-regions.

To address the questions of whether LPFC subregions dissociate based on 1) responses to

abstract sequences, and 2) responses to sequence components (image identity and periodicity),

we conducted an awake monkey fMRI experiment. Monkeys performed two no-report tasks, one

that contained abstract visual sequences (previously reported on in Yusif Rodriguez et al.

(2023)), and another that did not contain abstract visual sequences but maintained the image

identities and timing structure of the sequence task. We defined two LPFC subregions using a

parcellation of the PFC (Rapan et al., 2023): the posterior fundus of area 46 (p46f) that

overlapped with the previously identified sequence responsive subregion (Yusif Rodriguez et al.,

2023), and the adjacent posterior ventral shoulder of area 46 (p46v). Building on our previous

observations, we hypothesized that p46f would show responses unique to changes in abstract

visual sequences and not to their image identity and periodicity components. We hypothesized

the contrary for p46v: responses to changes in sequence components but not sequences

themselves. Our results broadly supported these hypotheses, with sequence responses in p46f and

not p46v, and responses to image identity and periodicity in p46v. These results further our

understanding of the representation of abstract visual sequences in adjacent subregions in the

LPFC.

Materials and Methods

Participants

165   We tested three adult male rhesus macaques (ages spanning 6-12 years during data collection, 9-

166   14 kg). All procedures followed the NIH Guide for Care and Use of Laboratory Animals and

167   were approved by the Institutional Animal Care and Use Committee (IACUC) at Brown

168   University.

Task Design and Procedure

All visual stimuli used in this study were displayed using an OpenGL-based software system

developed by Dr. David Sheinberg at Brown University. The experimental task was controlled

by a QNX real-time operating system using a state machine. Eye position was monitored using

video eye tracking (Eyelink 1000, SR Research). Stimuli were displayed at the scanner on a 24-

inch BOLDscreen flat-panel display (Cambridge Systems).

Each image presentation consisted of fractal stimulus (approximately 8° visual angle) with

varying colors and features. Fractals were generated using MATLAB for each scanning session

using custom scripts based on stimuli from Kim and Hikosaka (2013) following the instructions

outlined in Miyashita et al. (1991). For each scan session, new, luminance matched fractal sets

were generated. All stimuli were presented on a gray background, with a fixation spot that was

always present on the screen superimposed on the images. To provide behavioral feedback, the

fixation spot was yellow when the monkey was successfully maintaining fixation and red if the

monkey was not fixating. Stimuli were displayed for 0.1, 0.2, or 0.3 s each, depending on the

task, sequence type, and timing template.

The timing of liquid rewards was the same across tasks and not contingent on image

presentations, only on the monkey maintaining fixation. Rewards were delivered on a graduated

schedule such that the longer the monkey maintained fixation, the more frequent rewards were

administered (Leite et al., 2002). The first reward was given after 4 s of continuous fixation.

After two consecutive rewards of the same fixation duration, the fixation duration required to

obtain reward was decreased by 0.5 s. The minimum duration between rewards that the monkey

could obtain was 0.5 s. Fixation had to be maintained within a small window (typically 3° of

visual angle) around the fixation spot to not break fixation. The only exception was a brief time

window (0.32 s) provided for blinks. If the monkey's eyes left the fixation window and returned

within that time window, it would not trigger a fixation break. If fixation was broken, the reward

schedule would restart at the maximum 4 s duration required to obtain reward.

Tasks were organized into runs. Runs typically lasted approximately 10 min and only one task

was shown for each run. The order of tasks was pseudo-randomized within a scanning session

(one day) to balance the overall number of runs for each task and their presentation order.

Monkeys completed approximately 10 runs in a session.

Runs were initiated according to the monkey's fixation behavior to ensure that the monkey was

not moving and engaged in the task before acquiring functional images. During this pre-scan

period, a fixation spot was presented. Once the monkey successfully acquired this fixation spot

and received approximately four liquid rewards (12 – 16 s), functional image acquisition and the

first habituation block were initiated. Monkeys maintained fixation for the duration of the run.

Abstract Sequence Viewing (SEQ) Task

The details of the abstract sequence viewing task have been previously described (Yusif

Rodriguez et al., 2023) and are briefly summarized here. There were a total of five sequence

types and nine timing templates (**Figure 1**). The inter-sequence interval was jittered to

decorrelate across timing templates (mean 2 s, 0.25-8 s).

Habituation Sequences

Habituation sequences used images drawn from a pool of four fractals [A, B, C, D] and were

arranged to follow one of two possible rules: three the same, one different, and four the same. All

four-image sequences used one of six possible timing templates (**Figure 1C**).

Deviant Sequences

Deviant sequences used images drawn from a different pool of three fractals [E, F, G]. All

deviant images were displayed for 0.2 s and used the same general timing template (adjusted for

the number of items in the sequence). There were four deviant types, as follows:

● *New Items, Same Rule (NISR)*: four-image sequences that follow the same rule as the

habituation sequences.

● *Rule Deviants*: four-image sequences that follow the alternate rule not used in the

habituation sequences.

● *Number Deviants*: two- or six-image sequences that follow the same rule as the

habituation sequences.

● *Double Deviants*: combine the rule and number deviant types and contain two- or six-

image sequences that follow the alternate rule not used in the habituation sequences.

Block Structure

All blocks contained 30 sequences and an equal number of the six possible timing templates for

habituation sequences. Sequences could not start with the same image as the final fractal of the

previous sequence. In deviant blocks, six of the 30 sequences were deviant sequences. Deviant

sequences did not occur in the first six sequences (to avoid block initiation) or consecutively.

Blocks with two- and six-image sequences contained an equal number of both.

Run Structure

Each run contained five sequence blocks interleaved with 14 s fixation blocks, during which only

a fixation spot was present with no additional visual stimuli (**Figure 1D**). Monkeys maintained

fixation throughout the sequence and fixation blocks. The first sequence block was always all

habituation sequences. The four subsequent sequence blocks each contained one type of deviant

sequence. The sequential rule used for each run was counterbalanced across runs and sessions to

have an equal number of each. Monkeys typically completed 4-8 runs of this task (among other

tasks) in a session.

No Sequence (NoSEQ) Task

The main difference between NoSEQ and SEQ was that images were not arranged according to a

sequential rule (as in SEQ) and instead displayed in pseudorandom order such that there were no

consecutively repeated images. Images were still displayed as grouped into four-, two-, or six-

image sets, depending on the block. All the remaining basic structure of NoSEQ was the same as

SEQ. We adjusted the terminology to reflect this fact and more clearly dissociate between the

tasks. There were the same 9 timing templates in NoSEQ as in SEQ. However, for NoSEQ,

rather than referring to them as 6 habituation and 3 deviant timing templates as in SEQ, we refer

to them as 6 standard and 3 nonstandard timing templates in NoSEQ. Similarly, the same two

pools of fractal images referred to as habituation and deviant in SEQ are referred to as the 4

standard and 3 nonstandard images in NoSEQ. We underscore that the timing templates and

image pools are the same between SEQ and NoSEQ within a single scanning session, despite the

difference in naming.

Block Types and Structure

We define trials as a series of grouped images with the same timing structure. Most trials

contained four images, with some containing two or six (described further below). All blocks

contained 30 trials (120 images total). We note that even though images are grouped into trials,

reward is on an independent schedule based on the duration of fixation, as in the SEQ task. The

first six trials in a block did not contain nonstandard timing templates or nonstandard images.

Each block contained key differences with respect to the composition of the timing templates and

images used. There were four possible block types (**Figure 1E**), as follows:

● *All Standard Timing*: Each four-image trial used one of the 6 possible standard timing

templates (5 of each). Images were drawn only from the standard pool. This condition is

the same structure as habituation timing in the SEQ task.

● *Four-Image Nonstandard Timing (4NST)*: Six trials had four-image nonstandard timing

and the remaining 24 trials had standard timing. This timing structure matched the NISR

and rule deviant blocks in the SEQ task. The relative fraction of nonstandard images

matched the SEQ task (20%, 24 individual images), but they were randomly intermixed

with images from the standard pool.

● *Two- and Six-Image Nonstandard Timing (2/6NST)*: Six trials had two- or six-image

nonstandard timing (three of each) and the remaining 24 trials had standard timing. This

timing structure matched the number deviant blocks in the SEQ task. As in Four-Image

Nonstandard Timing blocks, 20% of images were drawn from the nonstandard image

pool and the remainder from the standard image pool. All images were displayed in

random order.

● *Novel*: As in the Standard Timing block, each four-image trial used one of the 6 possible

standard timing templates (5 of each). However, the images came from a novel pool of

four images that had not been used in either the standard or nonstandard image pools.

Run Structure

Each run was composed of four image blocks, interleaved with 14 s fixation blocks. As in the

SEQ task, fixation blocks consisted of only a fixation spot present and no additional visual

stimuli where the monkey had to maintain fixation. The first block of each run was always an All

Standard Timing block. The two subsequent blocks were either a Four-Image Nonstandard

Timing block or a Two- and Six-Image Nonstandard Timing block, with their order

counterbalanced across runs. The last block was always a Novel block. Runs lasted

approximately 10 min. Monkeys typically completed 2-4 runs of this task (among other tasks) in

a single scanning session.

Data Acquisition

<u>FMRI Data Acquisition</u>

Methods are as described in (Yusif Rodriguez et al., 2023) and briefly summarized here.

Monkeys sat in the "sphinx" position in an MR-safe primate chair (Applied Prototype, Franklin,

MA or custom-made by Brown University) with their head restrained using a plastic "post"

(PEEK, Applied Prototype, Franklin, MA) affixed to the monkeys' head and the primate chair.

Monkeys wore earplugs during MRI scanning (Mack's Soft Moldable Silicone Putty Ear Plugs,

"kid's" size). Monkeys were habituated to all scanning procedures prior to the MRI scanning

sessions.

Approximately 30-60 min prior to each scanning session, monkeys were intravenously injected

with a contrast agent: monocrystalline iron oxide nanoparticle (MION, Feraheme (ferumoxytol),

AMAG Pharmaceuticals, Inc., Waltham, MA, 30 mg per mL or BioPal Molday ION, Biophysics

Assay Lab Inc., Worcester, MA, 30 mg per mL). MION was injected into the saphenous vein

below the knee (7 mg/kg), then flushed with a volume of sterile saline approximately double the

volume of the MION injected. No additional MION was added during scanning.

A Siemens 3T PRISMA MRI system with a custom six-channel surface coil (ScanMed, Omaha,

NE) at the Brown University MRI Research Facility was used for whole-brain imaging.

Anatomical scans consisted of a T1-MPRAGE (repetition time, TR, 2700 ms; echo time, TE,

3.16 ms; flip angle, 9°; 208 sagittal slices; 0.5 x 0.5 x 0.5 mm), a T2 anatomical (TR, 3200 ms;

TE 410 ms; variable flip angle; 192 interleaved transversal slices; 0.4 x 0.4 x 0.4 mm), and an

additional high resolution T2 anatomical (TR, 8020 ms; TE 44 ms; flip angle, 122°; 30

interleaved transversal slices; 0.4 x 0.4 x 1.2 mm). Functional images were acquired using a fat-

saturated gradient-echoplanar sequence (TR, 1.8 s; TE, 15 ms; flip angle, 80°; 40 interleaved

axial slices; 1.1 x 1.1 x 1.1 mm).

Data Analysis

All preprocessing and data inclusion criteria are the same as in (Yusif Rodriguez et al., 2023).

Most analyses were performed in Matlab using SPM 12 (https://www.fil.ion.ucl.ac.uk/spm).

Prior to analysis, data were preprocessed using the following steps: reorienting (to ensure proper

assignment of the x,y,z planes), motion correction (realignment), normalization, and spatial

smoothing (2 mm isotropic Gaussian kernel separately for gray matter and white matter). All

steps were performed on individual runs separately. The T1-MPRAGE anatomical image was

skull stripped using FSL BET brain extraction tool (http://www.fmrib.ox.ac.uk/fsl/) to facilitate

normalization. All images were normalized to the 112-RM SL macaque atlas (McLaren et al.,

2009).

Runs were included for analysis only if they met the following criteria: the monkey had to be

performing well and a sufficient number of acquisition volumes within the run had to pass data

quality checks. The monkey's performance was evaluated by calculating the percentage of time

within a run that fixation was maintained. Runs were excluded if the monkey was fixating < 80%

of the time. We used the ART toolbox (Artifact Detection Tools,

https://www.nitrc.org/projects/artifact_detect) to detect outlier volumes (standard global mean;

global signal detection outlier detection threshold = 4.5; motion threshold = 1.1mm; scan to scan

motion and global signal change for outlier detection). Any run with greater than 12% of

volumes excluded was excluded from analysis (**Table 1**).

FMRI Models

For all models, data were binned to evenly distribute included runs from the SEQ and NoSEQ

tasks (**Table 1**) into pseudo-subject bins. Each bin contained data from only one monkey and

distributed runs from the SEQ and NoSEQ tasks as evenly as possible. Each bin contained

approximately 20 SEQ and 10 NoSEQ runs. Runs from earlier and later scanning sessions were

pseudorandomly distributed across bins. For the SEQ task, both rule types (AAAA and AAAB)

were evenly distributed in each bin. This binning procedure resulted in 11 total pseudo-subject

bins. Of the 11 pseudo-subject bins, 5 were monkey W, 4 were monkey J, and 2 were monkey B.

Within-subject statistical models were constructed under the assumptions of the general linear

model (GLM) in SPM 12 for each pseudo-subject bin. Condition regressors were all convolved

with a gamma function (shape parameter = 1.55, scale parameter = 0.022727) to model the

MION hemodynamic response function (Vanduffel and Farivar, 2014). The first six image

groups (24 images) and reward times were included as nuisance conditions. Additional nuisance

regressors were included for the six motion estimate parameters (translation and rotation) and

image variability (standard deviation of within-run image movement variability, calculated using

the ART toolbox). Outlier volumes determined with the ART toolbox in preprocessing were

"scrubbed" by adding an additional regressors, each with a "1" only at the volume to be

excluded. The equation for the GLM is below (Poline and Brett, 2012):

$$Y = X\beta + \varepsilon$$

Where Y is the (n,1) time series data (n = number of time points or scans), X is (n, p) design

matrix of regressors, β is the vector of parameters, and ε is the error vector. Regressors included

are the 20 listed in **Table 2** plus the nine nuisance regressors listed above (total of 29 regressors)

and any additional columns required for "scrubbing" (one regressor per volume scrubbed). The

baseline used for comparisons was implicit in that it included unmodeled time for which there

were no explicitly defined condition, nuisance, or "scrubbed" regressors (i.e., fixation only time

where there were no images displayed during fixation only blocks and inter-sequence intervals).

Regressors were estimated using a bin-specific fixed-effects model. Whole-brain estimates of

bin-specific effects were entered into second-level analyses that treated bin as a random effect.

One-sample t-tests (contrast value vs zero, $p < 0.005$) were used to assess significance. These

effects were corrected for multiple comparisons when examining whole-brain group voxelwise

effects using extent thresholds at the cluster level to yield false discovery rate (FDR) error

correction ($p < 0.05$).

To assess the univariate effects of deviant sequences, we constructed a general linear model

(GLM) using instantaneous stimulus onset regressors. Both tasks were modeled simultaneously,

with runs from both tasks included in each pseudo-subject bin. For the SEQ task, onsets were

modeled similarly as described in Yusif Rodriguez et al. (2023). Onsets were modeled at the first

item in each sequence type. Habituation and deviant sequences were modeled separately.

Habituation sequences were divided by timing template (short, medium, and long) and whether

they came from the first block containing only habituation sequences or a subsequent block that

contained deviant and habituation images, yielding six total habituation sequence regressors.

Deviant sequences were modeled separately according to their type: NISR, rule deviants, number

deviants (two- and six-image), and double deviants (two- and six-image), yielding six total

deviant sequence regressors. In total, the SEQ task contained 12 condition regressors (**Table 2**).

For the NoSEQ task, onsets were modeled for the first item in each group of images (a single

timing template). Standard and nonstandard timing templates were modeled separately. As in the

SEQ task, standard timing templates were divided by those occurring in the first block (where

there were no nonstandard timing templates or images) and those occurring in subsequent blocks

that contained nonstandard images and timing templates. Standard timing templates were again

divided by short, medium, and long yielding a total of six standard timing template regressors.

Nonstandard timing templates were modeled separately as four-image and two- and six-image,

yielding three total nonstandard timing template regressors. Nonstandard images that were

randomly interspersed in blocks that contained nonstandard timing templates were modeled

separately at the onset of each individual nonstandard image. The novel image block was also

separately modeled and divided by the three standard timing templates (short, medium, long);

however, these were not included in analyses. In summary, the NoSEQ task contained six

standard time, three nonstandard time, one nonstandard image, and three novel image regressors

for a total of 13 regressors (**Table 2**).

<u>ROI Analysis</u>

Individual area 46 subregion ROI images were directly acquired from the MEBRAINS

Multilevel Macaque Atlas (Balan et al., 2024) (https://www.ebrains.eu/tools/monkey-brain-

atlas). Individual subregion image warps were created from their native space to 112RM-SL

space using Rhemap (Sirmpilatze and Klink, 2020) (https://github.com/PRIME-RE/RheMAP).

Individual warps were then applied to create images used in ROI analysis for p46v (**Figure 2**).

Because the ROI used in Yusif Rodriguez et al. (2023) spanned subregions p46df and p46vf and

responses in these subregions were not distinct, for simplicity, we combined subregions in the

fundus of area 46 to create p46f (p46df + p46vf).

To compare activation within and across ROIs in a manner that controlled for variance, we

extracted average t-values for each ROI from whole-brain t-maps using the Marsbar toolbox

(Jean-Baptiste Poline, 2002). T-maps were the result of voxel-wise t-tests of parameter weights

for the conditions of interest scaled by the residual error in the model (GLM) compared to a null

hypothesis of zero, i.e., the resulting t-values from the condition > baseline contrast (see Poline

& Brett (2012) for further details). This procedure resulted in a total of 11 t-values for each

condition (one for each pseudo-subject bin: n = 11 bins) that were entered into repeated measures

analyses of variance (RM-ANOVAs) with the identity of the monkey entered as a covariate.

Results

We show awake fMRI results from three male monkeys (Macaca mulatta) during two no-report

(only central fixation was required) viewing tasks. One was an abstract sequence viewing task

(abbreviated SEQ hereafter) that contained a visual sequence rule and structured timing (as

reported in Yusif Rodriguez, et al., 2023). The second task did not contain abstract visual

sequences (abbreviated NoSEQ hereafter) but maintained the same stimulus frequencies and

periodicity structure as SEQ (**Figure 1**). Our main goals were to 1) test if sequential responses

differ between the fundus and more ventral LPFC subregions, and 2) test if and how these

subregions respond to changes in the components of abstract visual sequences.

The fundus of area 46 differentially represents changes to abstract visual sequences

To address our first question, we first tested if sequential responses differed between LPFC

subregions: one in the fundus and one adjacent and more ventral. To accomplish this goal, we

first needed to define two things: the precise anatomical locations of the regions of interest

(ROIs) and sequential responses.

To define the ROIs, we used a parcellation of PFC in conjunction with our previously defined

ROI in Yusif Rodriguez (2023). The previous ROI was a 3 mm radius (895 voxels) sphere in the

right hemisphere (named R46) based on coordinates chosen for their functional connectivity

similarity to human sequence responsive areas of the lateral frontal cortex (Sallet et al., 2013).

This ROI was not conducive to comparing directly adjacent areas because of its spherical shape

and because it necessarily included some white matter. Therefore, we created ROIs using the

MEBRAINS Multilevel Macaque Brain Atlas (Rapan et al., 2023; Balan et al., 2024) that

parcellated PFC according to cytoarchitectonics, functional connectivity, and neurochemical

data. This atlas divides area 46 into eight distinct regions (four anterior and four posterior) that

are then divided into dorsal and ventral shoulder and fundus regions (**Figure 2A**, **B**). Of the area

46 subdivisions, the posterior fundus (p46df and p46vf) regions showed the greatest overlap with

the previous R46 ROI constructed from functional connectivity seed coordinates (Sallet et al.,

2013; Yusif Rodriguez et al., 2023) (**Figure 2C**). Of the 895 voxels in the previous R46 ROI,

40.5% (420) overlapped with cortical gray matter, and all those voxels overlapped with p46df

and p46vf combined. We therefore combined the fundus regions and focused our analyses on the

posterior fundus (p46f). We compared p46f (1036 voxels) to an adjacent, more ventral

subregion: posterior ventral (p46v, 1060 voxels). We focused on ROIs in the right hemisphere

because previous results were observed on the right (though the right hemisphere was not

statistically different from the left) (Yusif Rodriguez et al., 2023).

To define sequential responses, we used deviations from an established (habituated) abstract

visual sequence in the SEQ task (as in Yusif Rodriguez et al. (2023)). The general logic was that

regions that responded to changes in the abstract visual sequence may play a role in tracking that

information. To create these changes, monkeys were first habituated to four-item sequences of

images that followed a particular rule, e.g., same, same, same, different. Images were drawn

from a pool of habituation images. In subsequent blocks, some of the sequences (6 out of 30)

were deviant sequences that drew images from a separate deviant pool and differed from

habituation sequences in one of four ways: new items, same rule (NISR), rule deviants, number

deviants, or double deviants (included to counterbalance the design but not included for

analysis). All comparisons to determine abstract sequence responses in the SEQ task were

between NISR and rule or number deviants. This comparison controlled for the use of less

frequent deviant images and any changes observed would be due to changes in the abstract

sequence.

For all analyses, we measured the cerebral blood volume (CBV) of a contrast agent,

monocrystalline iron oxide nanoparticle (MION) activity as our indicator of neural activity. We

created a single model for both the SEQ and NoSEQ tasks (see Methods for details). Statistical

testing was performed on approximately 20-run bins (n = 11), each consisting of data from a

single monkey. For each condition, t-values were extracted for that condition compared to

baseline (e.g., Rule Deviant > Baseline) to account for potential differences in variance across

conditions. These values were used to examine ROI activity throughout, and we refer to

comparisons by the condition of interest (i.e., without listing the contrast over baseline, e.g., Rule

Deviant). All statistical tests on ROIs were performed on binned data and included a covariate

for monkey identity (n = 3). While we report the effect of variation between monkeys in the

following analyses, the main focus of the study was not on individual differences, and our

discussion focuses on condition effects.

We first tested the hypothesis that sequential responses differ between right p46f and p46v. As in

our previous study, we compared the sequence deviants Rule and Number to NISR, with

increased activity for deviants indicating sequential processing. Replicating previous results

(with the newly defined ROI), both deviant responses were significantly greater than NISR in

right p46f (Rule > NISR: p = 0.01; Number > NISR: p = 0.04, **Figure 3A,B**, **Table 3**). In

contrast, responses in p46v did not differ between deviants and NISR, resulting in a significant

interaction between the two areas for number deviants compared to NISR (p = 0.02, **Figure 3B**,

Table 4) and a marginal interaction for rule deviants compared to NISR (p = 0.09, **Figure 3A**,

Table 4). These ROI results were supported by whole brain contrasts of Rule Deviants > NISR

(**Figure 3C**) and Number Deviants > NISR (**Figure 3D**) in the SEQ task (**Table 5**). Both deviant

types showed significant clusters of activation in right p46f. Therefore, these results support the

hypothesis that sequential responses differ between p46f, which responds to changes in abstract

visual sequences, and p46v, which does not.

The ventral shoulder of area 46 differentially represents changes to abstract sequence components

We next addressed the second question of if and how these regions respond to changes in sequential components in the absence of an abstract visual sequence. We examined two main features in the NoSEQ task: image periodicity (timing template) and identity. These two image features are components of the SEQ task, exist in parallel in the NoSEQ task, and will be described further below.

To test for responses in the DLPFC related to image periodicity, we compared standard to nonstandard timing templates in the NoSEQ task. Previous studies showed that regions of the frontal cortex process different types of timing structures (Onoe et al., 2001; Genovesio et al., 2006; Chiba et al., 2021), raising the possibility that a difference in the timing structure alone could be a component of sequence responses. In the context of this experiment, image periodicity refers to the timing template used. In the SEQ and NoSEQ tasks, most sequences/groups had one of six possible standard timing templates (referred to as habituation in the SEQ task, **Figure 1**). A unique timing template (0.2 s image duration for medium, 1.7 s, total duration) was used infrequently for deviants in the SEQ task (after the first block, 6 out of 30 sequences). In the NoSEQ task that structure was mirrored: after the first block, 6 out of the 30 stimulus groupings used a nonstandard timing template. These blocks either contained six 4-image nonstandard timing (4NST) or three each of 2- and 6-image nonstandard timing (2/6NST). Importantly, even though the timing template was the same as for SEQ deviants (just termed differently for the NoSEQ task), the images in NoSEQ were pseudorandomly presented and were not composed of entirely nonstandard images. To determine if brain areas responded to changes in timing

template alone, we compared responses to these nonstandard timing templates to the six other

standard timing templates in the NoSEQ task.

We tested the hypothesis that p46v would show a greater difference in responses to image

periodicity than p46f in the NoSEQ task. In general, more ventral LPFC regions are thought to

have more object-based or visual responses (Meyer et al., 2011; Yamagata et al., 2012; Tang et

al., 2021; Xu et al., 2022). First, we found that changes in timing structure alone did not elicit

deviant responses in right p46f. There were no reliable differences between 4NST ($p = 0.86$,

Figure 4A, **Table 6**) or 2/6NST compared to standard timing templates ($p = 0.85$, **Figure 4B**,

Table 6). In contrast, right p46v showed reliable differences when comparing standard to 4NST

($p = 0.03$, **Figure 4A**, **Table 6**) and standard to 2/6NST ($p = 0.01$, **Figure 4B**, **Table 6**). When

directly comparing p46f and p46v, overall responses were greater in p46v than p46f ($p = 0.01$,

Table 7) and there was a significant interaction such that the difference between nonstandard and

standard was significantly different by ROI for 2/6NST and marginal for 4NST (ROI x

condition, 2/6NST: $p = 0.05$, 4NST: $p = 0.09$; **Table 7**). These results were supported by whole

brain contrasts of 4NST > Standard Timing and 2/6NST > Standard Timing showing no

significant clusters of activation in p46f with other significant activation in distinct visual and

association areas (**Figure 4C,D**, **Table 8**). Together, these results support the hypothesis that

p46v responds to image periodicity and dissociates from responses in p46f. Further, p46v may be

part of a network of other brain areas that is specialized to detect periodicity differences,

independent of abstract sequential structure.

To test for responses in the LPFC related to image identity, we compared standard to

nonstandard images in the NoSEQ task. Deviant/nonstandard images in the SEQ task were less

frequent. It is unlikely that the deviant responses observed in the SEQ task were driven by these

images, because deviant comparisons were all made across conditions that contained images

from the deviant pool (e.g., rule deviant vs. NISR, **Figure 1**). However, infrequent or surprising

images have been shown to drive responses in LPFC (Chao et al., 2018; Camalier et al., 2019;

Grohn et al., 2020). Therefore, we aimed to determine if responses in area 46 could be driven by

less frequent image presentations, independent of sequential context. To examine this sequence

component, we again used conditions that were separate from an abstract visual sequence, i.e., in

the NoSEQ task. We compared responses to the randomly interspersed nonstandard images to

standard images to ensure other aspects of the task were held constant.

We tested the hypothesis that p46v would show a greater difference in responses to image

identity than p46f in the NoSEQ task. First, we found that nonstandard responses were not

reliably different from standard image responses in right p46f (p = 0.67, **Figure 5A**, **Table 9**). In

contrast, p46v differentiated between standard and nonstandard images, but with reliably greater

responses for standard images (p = 0.01, **Figure 5A**, **Table 9**). We directly compared responses

in p46f to p46v and found that responses to nonstandard compared to standard images were

reliably different (ROI x condition: p = 0.03, **Table 10**). These ROI results showing greater

responses to standard than nonstandard images were also supported by whole brain contrasts

(**Figure 5B**). Significant clusters of activation were observed for Standard > Nonstandard images

in right p46v but no other area 46 subregions (**Table 11**). In Nonstandard > Standard image there

were no significant clusters in the frontal cortex. These results support the hypothesis that p46v

differentially represents standard and nonstandard images, and p46f does not.

Discussion

We had two main goals in this experiment: 1) to test if SEQ deviant responses previously

observed in p46f were different from those in p46v, and 2) to test if and how p46f and p46v

responded to changes in abstract visual sequence components, image identity and periodicity,

alone in a NoSEQ task. We hypothesized that p46v would not respond to changes in abstract

visual sequences as a whole, but to the defined components. The results generally supported the

hypotheses, with p46f responding more strongly than p46v to abstract sequence deviants in the

SEQ task but not to differences in sequential components (standard vs nonstandard) in NoSEQ,

further strengthening its role as an area that uniquely represents abstract sequential changes.

P46v instead differentiated between standard and nonstandard sequence components, image

identity and periodicity, in NoSEQ. This result supported our predictions that this region was

more influenced by visual sensory inputs and not necessarily the higher order structures in

abstract sequences. These results provide important knowledge of the functional subdivisions

within area 46 of the LPFC and scaffold future understanding of this important area for

cognition.

The observed differences between p46f and p46v may expand on the notion of "dorsal" and

"ventral" distinctions within the LPFC. Classic anatomical definitions of dorsal and ventral have

used the principal sulcus as a dividing line, bisecting the fundus between the two (e.g., Petrides

and Pandya (1999)). Extracellular electrophysiology experiments often focus on the cortical

surface of LPFC, understandably due to the ability to visualize locations and position electrodes

and arrays. Therefore, it is less clear how the fundus region itself may or may not fit into

functional generalizations of LPFC. Anatomy and the present experiment suggest a more distinct

functional role for p46f, at least within more posterior LPFC regions. The present results

strikingly align with a multimodal parcellation of macaque area 46 showing that p46f

hierarchically clustered with the most rostral regions of LPFC whereas p46v clustered with more

caudal sensory and motor areas (Rapan et al., 2023). Current schema of the functional

organization of LPFC suggest that more spatial and action-oriented responses are localized more

dorsally and more non-spatial and object-oriented responses are localized more ventrally (Meyer

et al., 2011; Yamagata et al., 2012; Tang et al., 2021; Xu et al., 2022). These results are also

broadly consistent with a proposed distinction of dorsal 'How' (perception to action

transformation) and ventral 'What' (identity) information in the LPFC (O'Reilly, 2010), if it is

considered as a gradient and "actions" could be considered non-motor. Further, we did not

observe responses to changes in abstract visual sequences in p46v, as others have observed in

what was termed VLPFC (Wang et al., 2015; Bellet et al., 2024). It is unclear if p46v and

VLPFC are anatomically overlapping. More experiments are needed before strong conclusions

can be drawn, but the results here suggest that the subregions within what has been referred to as

DLPFC and VLPFC should be carefully considered, and the fundus may need separate

classification. Future experiments could benefit from recent developments in whole brain

imaging technology including functional contrast agents, PET-MR, and high field fMRI to

localize anatomical and functional brain regions both independently and pre/post-

electrophysiological recordings.

We did not observe responses in p46f to changes in the periodicity alone in NoSEQ. In contrast,

right p46v showed differences in responding to standard and nonstandard timing templates, with

increased responses to the more frequent standard timing presentations. These observations are

generally consistent with previous observations of timing-related activity in monkey LPFC (Niki

and Watanabe, 1979; Onoe et al., 2001; Genovesio et al., 2006; Cueva et al., 2020; Chiba et al.,

2021), although the precise anatomical location was not specified. Human LPFC responses in

temporal expectation tasks (Coull and Nobre, 2008) were also similar. Outside of area 46 we

observed some of the same regions that have been observed for duration perception in monkeys,

such as putamen, cerebellum, and V2 (Onoe et al., 2001). We also observed regions similar to

those observed in humans related to temporal expectation such as the basal ganglia, temporal

cortex, and cerebellum (Coull and Nobre, 2008). Together these results illustrate the specificity

of subregions within area 46 and suggest that adjacent subregions code for different stimulus

properties.

Area p46v responses differentiated between standard and nonstandard images in NoSEQ while

p46f did not. Responses in p46v were greater to standard images than nonstandard images. These

results suggest that this response was not a typical 'surprise' response. In our task, standard

images were presented with greater frequency, and closer in time to each other, than the isolated

individual appearances of the nonstandard images. Therefore, such responses could be partially

driven by the increased frequency, and thus a greater summed BOLD response, of standard

compared to nonstandard images and. This increased image frequency may have been the

primary driver of responses in regions such as visual cortex, which showed relatively large

significant clusters in Standard > Nonstandard NoSEQ images. However, if frequency was the

only driver of such responses, then we would have likely observed these responses throughout

the whole brain rather than in a specific set of regions. Outside of the visual cortex there were

roughly equal numbers of areas that showed responses to the reverse contrast (Nonstandard

Image > Standard Image), but these responses were not located in area 46. An intriguing

possibility is that a greater response to the standard images in p46v is due to the previous

association that those images have to abstract visual sequences, or their greater familiarity

(Rainer et al., 1999; Stern et al., 2001; Leaver et al., 2009). Further investigation will be

necessary to determine if that association is a component of the response in p46v along with

regions observed for Standard Image > Nonstandard Image in the whole brain. There were a

small number of regions observed outside of the frontal cortex that showed significantly greater

responses to nonstandard images. These regions were not necessarily overlapping with areas

typically associated with 'surprise' or prediction error (Grohn et al., 2020), again raising the

prospect that a form of association may govern these responses as well. Further research will be

needed to discern the underlying driving forces, but the fact remains that sensory related

responses localize to p46v and not adjacent p46f, again illustrating the specificity of responses

within area 46.

This study's approach was limited in the following ways. First, though p46f and p46v showed

significant differences, some effect sizes remained small. These results could be related to

limitations of whole-brain event-related monkey fMRI: the spatial resolution, signal-to-noise,

and inherent smoothness of the data. There could be differences in alignment of the voxels with

the regions and partial volume effects that would be difficult to resolve without fundamentally

changing the experiment by scanning a small volume at higher resolution, using a greater field

strength (which may not be available), or greatly increasing the sample size (introducing other

limitations). However, these experiments provide an ideal foundation for techniques with higher

spatial resolution such as electrophysiological recordings. Second, while the no-report paradigm

confers the advantage of eliminating possible confounds due to executing responses, it does not

allow for direct comparisons with behavior performance. In other words, even though we have

observed that regions within area 46 respond to changes in stimuli, we do not know how such

information may contribute to decisions or the production of actions. Third, we have specifically

focused on two subregions within area 46, which is itself only one of several areas defined as

belonging to LPFC. Other regions within and beyond LPFC warrant further investigation, and

the results here potentially contextualize further differences within and among subregions.

Though some of the preceding items are limitations of the chosen task and technique, we hope

that this experiment and others like it highlights the utility of different data acquisition modalities

and opens important avenues of future research.

In conclusion, we provide unique evidence for the anatomical and functional specificity of

abstract visual sequence deviant responses in a specific subregion of LPFC, p46f. In tandem, we

provide evidence that the adjacent region p46v, differentiates image identity and periodicity

components. These results reinforce the potential parallel with findings in human brain areas.

Rostrolateral PFC in humans is necessary for abstract task sequences and is most analogous to

p46f in monkeys (Desrochers et al., 2015; Yusif Rodriguez et al., 2023). Further, these results

illustrate the utility in using fMRI to isolate components to cognitive processes, in this case,

sequential components. The LPFC, which may have in the past appeared to be a more

homogenous region, may in fact be even more distinct in its subdivisions and functional

mapping. This study lays the foundation for an approach to functionally dissociating subregions

in the cortical structures that underlie many complex and abstract daily functions, such as

cooking a meal or appreciating a piece of music.

Tables

Table 1. Data excluded and included for analysis.

Percent Excluded Fixation

	Monkey B	Monkey J	Monkey W
SEQ	6.89%	10.46%	3.03%
NoSEQ	6.96%	10.13%	5.06%

Percent Excluded Motion

	Monkey B	Monkey J	Monkey W
SEQ	15.15%	0%	0.6%
NoSEQ	16.5%	0.63%	1.89%

Total Included Runs

	Monkey B	Monkey J	Monkey W	Total Runs
SEQ	70	65	97	232
NoSEQ	17	38	43	98

686   **Table 2.** Regressors used in SEQ and NoSEQ model. Both tasks are modeled together.

SEQ Task Regressors	
First Block	**Subsequent Blocks**
Habituation Short	Habituation Short
Habituation Medium	Habituation Medium
Habituation Long	Habituation Long
	New Item Same Rule (NISR)
	Rule Deviant
	Number Deviant – 2 items
	Number Deviant – 6 items
	Double Deviant – 2 items
	Double Deviant – 6 items
NoSEQ Task Regressors	
First Block	**Subsequent Blocks**
Standard timing - Short	Standard timing - Short
Standard timing - Medium	Standard timing - Medium
Standard timing - Long	Standard timing - Long
	Nonstandard image
	4-image nonstandard timing
	2-image nonstandard timing
	6-image nonstandard timing
	Novel image, standard timing - Short
	Novel image, standard timing - Medium
	Novel image, standard timing - Long

689   **Table 3.** Activity during SEQ task deviants compared to NISR in right area 46 using repeated measures ANOVAs.

690   P-values in bold are conditions of interest.

Rule Deviant

	Factor	DFs	F	p	η_p^2
	monkey	2,8	1.2	0.36	0.22
46f	condition	1,8	12.3	**0.01**	0.61
	monkey:condition	2,8	4.6	0.05	0.54
	monkey	2,8	4.5	0.05	0.53
46v	condition	1,8	0	0.93	0
	monkey:condition	2,8	0.2	0.82	0.05

Number Deviant

	Factor	DFs	F	p	η_p^2
	monkey	2,8	0.6	0.57	0.13
46f	condition	1,8	5.8	**0.04**	0.42
	monkey:condition	2,8	5.5	0.03	0.58
	monkey	2,8	2.8	0.12	0.41
46v	condition	1,8	0.3	0.58	0.04
	monkey:condition	2,8	0.1	0.88	0.03

692   **Table 4.** Comparisons of activity in right p46f and p46v during deviants compared to NISR in the SEQ task using

693   repeated measures ANOVAs. P-values in bold are conditions of interest.

Rule Deviant

Factor	DFs	F	p	η_p^2
monkey	2,18	0.2	0.82	0.02
ROI	1,18	5	0.04	0.22
condition	1,18	2.9	0.11	0.14
monkey:condition	2,18	1.9	0.17	0.18
ROI:condition	1,18	3.3	**0.09**	0.15

Number Deviant

Factor	DFs	F	p	η_p^2
monkey	2,18	0.6	0.57	0.06
ROI	1,18	3.7	0.07	0.17
condition	1,18	1	0.32	0.05
monkey:condition	2,18	2.5	0.11	0.22
ROI:condition	1,18	6	**0.02**	0.25

696 **Table 5.** Coordinates of activity clusters in Rule and Number deviant > NISR contrasts in the SEQ task.

Contrast Location	Extent (vox)	Peak T-val	X	Y	Z
Rule Deviant > NISR					
Rostral Medial Frontal Pole	105	6.25	0.5	45.5	14.5
Dorsal Area 46	381	7.53	14	35.5	25.5
Ventral Area 46	100	5.53	-15	35.5	22
Medial Agranular Insular Region	159	7.53	8	30.5	15
Orbital Area 12	148	5.59	-18	30.5	12.5
Area F5 of Ventral Pre-motor Cortex	173	6.06	-19	13	9.5
Granular Layer of Dentate Gyrus	137	5.66	-9.5	2.5	12.5
Cerebellum	229	4.85	6	-5.5	5.5
Visual Area 2	123	5.65	12	-12	20
	115	7.21	3	-22	23
Number Deviant > NISR					
Dorsal Area 46	166	5.73	11	36.5	25.5
Orbital Area 12	133	5.17	-18	31	10.5
Medial Area 13	162	5.21	7.5	27.5	17.5
Visual Area 2	332	7.99	5	-17	18

699 **Table 6.** Activity during NoSEQ task 4NST and 2/6NST compared to standard timing in right area 46 using

700 repeated measures ANOVAs. P-values in bold are conditions of interest.

4NST	Factor	DFs	F	p	eta2p
	monkey	2,8	0.8	0.5	0.16
46f	condition	1,8	0	0.86	0
	monkey:condition	2,8	0.1	0.95	0.01
	monkey	2,8	9.6	0.01	0.71
46v	condition	1,8	6.8	**0.03**	0.46
	monkey:condition	2,8	5.1	0.04	0.56
2/6NST	Factor	DFs	F	p	eta2p
	monkey	2,8	0.1	0.95	0.01
46f	condition	1,8	0	0.85	0
	monkey:condition	2,8	1.1	0.37	0.22
	monkey	2,8	14.5	0	0.78
46v	condition	1,8	10	**0.01**	0.55
	monkey:condition	2,8	4.4	0.05	0.52

Table 7. Comparisons of activity in right p46f and p46v during nonstandard compared to standard timing in the

NoSEQ task using repeated measures ANOVAs. P-values in bold are conditions of interest.

4NST

Factor	DFs	F	p	η_p^2
monkey	2,18	6.3	0.01	0.41
ROI	1,18	8.4	**0.01**	0.32
condition	1,18	2.8	0.11	0.14
monkey:condition	2,18	1.9	0.19	0.17
ROI:condition	1,18	3.1	**0.09**	0.15

2/6NST

Factor	DFs	F	p	η_p^2
monkey	2,18	3.4	0.06	0.27
ROI	1,18	8.2	**0.01**	0.31
condition	1,18	5.3	0.03	0.23
monkey:condition	2,18	5	0.02	0.36
ROI:condition	1,18	4.3	**0.05**	0.19

708 **Table 8.** Coordinates of activity clusters in 4NST and 2/6NST > Standard timing contrasts in the NoSEQ task.

Contrast Location	Extent (vox)	Peak T-val	X	Y	Z
4NST > Standard timing					
Putamen	135	6.8	15.5	22.5	10.5
Temporal Parietooccipital Associated Area	113	5.71	-16	22.5	2
Secondary Somatosensory Cortex	92	6.31	25	18	14
Amygdala	85	5.03	8.5	17.5	1.5
Putamen	137	5.33	-15	15.5	13
Areas 1 and 2	84	4.69	-24	12.5	21
Visual Area 2	192	5.77	-12	-5.5	14
Cerebellum	112	6.1	-8	-9.5	10
Visual Area 2	150	5.26	-14	-18	10.5
Primary Visual Cortex	184	6.63	-6	-22	18.5
Primary Visual Cortex	197	5.44	-16	-22	18.5
Standard timing > 4NST					
Lateral Area 13	204	6.61	-15	33.5	18
Putamen	227	5.45	12.5	21	19
Area 29	494	7.43	16	0	12
2/6NST > Standard timing					
Dorsal Area 46	142	5.75	-7.5	45	22.5
Intermediate Agranular Insula Area	144	6.37	-11	32	9
Area F5 of Ventral Premotor Cortex	94	7.27	26	27	17.5
Agranular and Dysgranular Insula	177	6.52	21	22	15
Area TEm	222	11	-25	8	7
		3.92	-20	4.5	2
Ventral Intraparietal Area	119	4.76	8.5	5	26
Area Pga	244	6.73	-18	4	16.5
Medial Superior Temporal Area	252	6.09	-14	-1	22
Posterior Intraparietal Area	604	6.9	9.5	-5	22.5
Standard timing > 2/6NST					
Area F5 of Ventral Premotor Cortex	123	4.65	-21	29	17.5
Putamen	137	4.57	12	27.5	10.5
Putamen	110	5.16	-17	8.5	12
Temporal Parietooccipital Associated Area	141	6.34	22.5	5	14
Dorsal Visual Area 4	171	7.22	-28	-0.5	17.5
Dorsal Visual Area 4	183	5.95	-28	-4.5	21.5
Lateral Reticular Nucleus	104	4.58	5.5	-7	-4
Primary Visual Cortex	504	6.36	25.5	-7.5	16.5
Primary Visual Cortex	215	5.22	21.5	-12	20.5
Primary Visual Cortex	192	5.29	-20	-13	26

711 **Table 9.** Activity during NoSEQ task nonstandard images compared to standard images in right area 46 using

712 repeated measures ANOVAs. P-values in bold are conditions of interest.

	Factor	DFs	F	p	η_p^2
	monkey	2,8	1.3	0.32	0.25
46f	condition	1,8	0.2	0.67	0.02
	monkey:condition	2,8	3.3	0.09	0.45
	monkey	2,8	9.7	0.01	0.71
46v	condition	1,8	11.8	**0.01**	0.6
	monkey:condition	2,8	10.2	0.01	0.72

715 **Table 10.** Comparisons of activity in right p46f and p46v during standard compared to nonstandard images in the

716 NoSEQ task using repeated measures ANOVAs. P-values in bold are conditions of interest.

Factor	DFs	F	p	η_p^2
monkey	2,18	1.8	0.2	0.17
ROI	1,18	8.7	0.01	0.33
condition	1,18	7	0.02	0.28
monkey:condition	2,18	12	0	0.57
ROI:condition	1,18	6	**0.03**	0.25

719 **Table 11.** Coordinates of activity clusters in Nonstandard Image > Standard Image and Standard Image >

720 Nonstandard Image contrasts in the NoSEQ task.

Contrast Location	Extent (vox)	Peak T-val	X	Y	Z
Nonstandard Image > Standard Image					
Hypothalamus	139	5.12	2	15.5	8.5
Middle Temporal Area	374	7.21	10	3	18
		4.83	17	-2.5	20.5
Ventral Visual Area 4	134	7.09	20	-7.5	8.5
Ventral Visual Area 6A	172	5.83	-3	-14	32
Visual Area 2	101	6.3	-7.5	-15	16.5
		4.06	-3	-20	12
Standard Image > Nonstandard Image					
Caudal Medial Frontal Pole	302	5.93	0	40	10.5
Dorsal Area 46	294	6.05	7	37.5	19
		4.84	7.5	45.5	21.5
Medial Area 13	217	7.08	6.5	34.5	12.5
Ventral Area 46	268	6.27	17.5	34.5	23.5
Area 45b	105	4.08	-20	29	22
Area F5 of Ventral Premotor Cortex	146	5.82	22	27.5	12
STS Part of Temporal Pole	339	5.97	15	24.5	-1.5
		3.25	18	17	-3
Anterior Ventral Area TE	472	7.53	-19	21	-2.5
Putamen	568	6.67	13	19	17
		3.41	8	14.5	21.5
Caudal Dorsal Premotor Cortex	187	6.79	17.5	18	31
Area TEm	344	6.17	23.5	9	9
		5.39	18	11.5	3.5
Temporal Parietooccipital Associated Area	171	5.01	-25	5.5	15.5
Medial Pulvinar Nucleus	93	5.09	-5.5	4.5	17.5
Primary Motor Cortex	88	5.35	11	1.5	31
Ventral Visual Area 4	304	5.32	18.5	-0.5	12
Posterior Intraparietal Areas	150	5.39	-11	-4.5	20.5
Visual Area 2	25046	43.43	19	-11	24
		30.66	23.5	-14	17.5
		21.88	28	0.5	21
		17.48	15.5	-15	30.5
		16.25	25	-5	14
		11.1	25.5	-6	25
		10.18	28	3	13
		8.78	11.5	-3	22.5
		6.57	16.5	-19	12.5
		6.35	18.5	-19	22.5
		5.11	13	-13	16.5

			4.97	3.5	-4.5	24
			4.51	16	-5	15.5
			4.16	26.5	6.5	5.5
Visual Area 2		21128	33.77	-20	-11	30
			24.6	-26	-4	20.5
			18.87	-21	-8.5	14
			17.13	-20	-15	21.5
			14.89	-13	-14	32.5
			6.23	-29	0.5	13.5
			6.15	-20	-20	12.5
			4.56	-14	-19	25.5
Cerebellum		205	5.82	-13	-11	-1.5

 Figure Legends

Figure 1. Sequence viewing task (SEQ) and No Sequence (NoSEQ) task structure. Both tasks are no-report. The

monkey maintains fixation at the central fixation spot throughout both tasks. **A**. Example partial habituation block

from SEQ task for sequence rule three same, one different (AAAB) and habituation timing templates. **B**. Example

partial standard block from NoSEQ illustrating non-sequential image order and standard timing templates. **C**.

Example stimuli pools (top) show a set of images that would be used in a single scanning session for both tasks (but

termed differently depending on the task). NoSEQ additionally contains a novel images category, with different

images not exemplified here. New images are used each session. Six possible habituation/standard event timing

templates (bottom, left) and deviant/nonstandard event timing templates (bottom, right) illustrated with gray

rectangles indicating individual image presentations. Total sequence/grouping durations are listed for each template

type. **D**. Example SEQ run, with each bar indicating one multi-image sequence: four images in habituation, new

items same rule (NISR), and rule deviants; two or six images in number and double deviants. Each block contains 30

sequences. The first block contains only habituation sequences and subsequent blocks (order counterbalanced)

contain only one of the four deviant types in six out of the 30 sequences. **E.** Example NoSEQ run where there is no

sequential order to the displayed images. Each bar indicates a multi-image set grouped by timing template and each

block contains 30 image groupings. To parallel SEQ structure, the first block in NoSEQ contains only standard

images and timing templates. In the two following blocks (order counterbalanced), six out of the 30 groupings have

nonstandard timing templates. One block has 4-image nonstandard timing (4NST) that parallels the rule deviant

block in SEQ, and the other block has 2- and 6-image nonstandard timing (2/6NST) together to parallel the number

deviant block in SEQ. Also, in the nonstandard blocks, 20% of the fractal images shown (in pseudorandom order)

are nonstandard (indicated by miniaturized nonstandard fractals), and the rest standard, again to mirror the

proportions in SEQ. Task relevant blocks alternate with fixation blocks for both SEQ and NoSEQ tasks. In fixation

blocks, monkeys maintained fixation on the fixation spot while no other images were displayed. Blue water droplets

schematize reward delivery, which is decoupled from sequence viewing and delivered on a graduated schedule

based on the duration the monkey has maintained fixation.

Figure 2. Schematic of anatomical subdivisions of area 46. A. Schematic of the area 46 subregions in the LPFC based on the atlas by Rapan et al. (2023) depicted on the right lateral surface of the macaque brain. **B.** Same cortical subregions illustrated in **A** with areas of comparison p46f and p46v in green and blue. Area p46d (light blue) shown only for context, indicated by gray lettering. **C.** Coronal slices displaying the area 46 ROI sphere used in Yusif Rodriguez et al. 2023 (red, outlined in black) superimposed on area 46 subregions. ROIs used for analysis in this study were green and blue, corresponding to regions illustrated in **A**). Light blue p46d only included for comparison and not used in analyses. Yellow voxels indicate overlap between the previous red sphere and current p46f ROI (green).

Figure 3. Right p46f and not p46v showed deviant responses in SEQ task. (A and B) T-values for the condition of interest > baseline are shown. Error bars are 95% confidence intervals (1.96 x standard error of the within-bin mean). **A**. Responses in rule deviants compared to new items, same rule (NISR) were different between p46f and p46v such that there was a significant main effect of ROI, and a marginal interaction between ROI and condition (indicated with ~). **B**. Number deviants compared to NISR between p46f and p46v showed a marginal main effect of ROI and significant interaction between ROI and condition. **C**. Voxel-wise contrast of Rule Deviants > NISR, false discovery rate (FDR) error cluster corrected for multiple comparisons (FDRc < 0.05, height p < 0.005 unc., extent = 100) are shown. **D.** Voxel-wise contrast of Number Deviants > NISR (FDRc < 0.05, height p < 0.005 unc., extent = 133) is shown. Color bar indicates T-values in **C** and **D**.

Figure 4. Right p46v and not p46f showed different responses to timing templates in NoSEQ. (A and B) T-values for the condition of interest > baseline are shown. Error bars are 95% confidence intervals (1.96 x standard error of the within-bin mean). **A.** 4NST compared to standard timing between p46f and p46v shows a significant main effect of ROI and a marginal interaction between ROI and condition (indicated with ~). **B**. 2/6NST compared to standard between p46f and p46v shows a significant main effect of ROI and interaction between ROI and condition (indicated with *). **C.** Voxel wise contrasts of 4NST > Standard timing (Hot colors) false discovery rate (FDR) error cluster corrected for multiple comparisons (FDRc < 0.05, height p < 0.005 unc., extent = 84), overlaid

with Standard timing > 4NST (Cold colors) false discovery rate (FDR) error cluster corrected for multiple

comparisons (FDRc < 0.05, height p < 0.005 unc., extent = 204) are shown. **D.** Voxel wise contrasts of 2/6NST >

Standard timing (hot colors; FDRc < 0.05, height p < 0.005 unc., extent = 94), overlaid with Standard timing >

2/6NST (cold colors; FDRc < 0.05, height p < 0.005 unc., extent = 104) are shown. Color bars indicate T-values in

C and **D**.

Figure 5. Right p46v and not p46f shows different responses to standard and nonstandard images in the

NoSEQ task. T-values for the condition of interest > baseline are shown. Error bars are 95% confidence intervals

(1.96 x standard error of the within-bin mean). **A.** Nonstandard compared to standard images showing reliable

differences between p46f and p46v showing a significant main effect of ROI with a significant interaction of ROI

and condition (indicated with *). **B.** Voxel wise contrasts of Nonstandard > Standard images false discovery rate

(FDR) error cluster corrected for multiple comparisons (hot colors, FDRc < 0.05, height p < 0.005 unc., extent =

101), overlaid with Standard > Nonstandard image (cold colors, FDRc < 0.05, height p < 0.005 unc., extent = 88)

are shown. Color bar indicates T-values.

<u>References</u>

Balan PF, Zhu Q, Li X, Niu M, Rapan L, Funck T, Wang H, Bakker R, Palomero-Gallagher N, Vanduffel W (2024) MEBRAINS 1.0: A new population-based macaque atlas. Imaging Neuroscience 2:1–26.

Bellet ME, Gay M, Bellet J, Jarraya B, Dehaene S, van Kerkoerle T, Panagiotaropoulos TI (2024) Spontaneously emerging internal models of visual sequences combine abstract and event-specific information in the prefrontal cortex. Cell Rep 43:113952.

Camalier CR, Scarim K, Mishkin M, Averbeck BB (2019) A Comparison of Auditory Oddball Responses in Dorsolateral Prefrontal Cortex, Basolateral Amygdala, and Auditory Cortex of Macaque. Journal of Cognitive Neuroscience 31:1054–1064.

Chao ZC, Takaura K, Wang L, Fujii N, Dehaene S (2018) Large-Scale Cortical Networks for Hierarchical Prediction and Prediction Error in the Primate Brain. Neuron 100:1252-1266.e3.

Chiba A, Morita K, Oshio K, Inase M (2021) Neuronal activity in the monkey prefrontal cortex during a duration discrimination task with visual and auditory cues. Sci Rep 11:17520.

Coull JT, Nobre AC (2008) Dissociating explicit timing from temporal expectation with fMRI. Current Opinion in Neurobiology 18:137–144.

Cueva CJ, Saez A, Marcos E, Genovesio A, Jazayeri M, Romo R, Salzman CD, Shadlen MN, Fusi S (2020) Low-dimensional dynamics for working memory and time encoding. Proc Natl Acad Sci USA.

Desrochers TM, Chatham CH, Badre D (2015) The necessity of rostrolateral prefrontal cortex for higher-level sequential behavior. Neuron 87:1357–1368.

Esmailpour H, Raman R, Vogels R (2023) Inferior temporal cortex leads prefrontal cortex in response to a violation of a learned sequence. Cereb Cortex 33:3124–3141.

Genovesio A, Tsujimoto S, Wise SP (2006) Neuronal Activity Related to Elapsed Time in Prefrontal Cortex. Journal of Neurophysiology 95:3281–3285.

Grohn J, Schüffelgen U, Neubert F-X, Bongioanni A, Verhagen L, Sallet J, Kolling N, Rushworth MFS (2020) Multiple systems in macaques for tracking prediction errors and other types of surprise. PLoS Biol 18:e3000899.

Jean-Baptiste Poline MB (2002) Region of interest analysis using an SPM toolbox.

Kim HF, Hikosaka O (2013) Distinct basal ganglia circuits controlling behaviors guided by flexible and stable values. Neuron 79:1001–1010.

Leaver AM, Lare JV, Zielinski B, Halpern AR, Rauschecker JP (2009) Brain Activation during
Anticipation of Sound Sequences. J Neurosci 29:2477–2485.

Leite FP, Tsao D, Vanduffel W, Fize D, Sasaki Y, Wald LL, Dale AM, Kwong KK, Orban G a,
Rosen BR, Tootell RBH, Mandeville JB (2002) Repeated fMRI using iron oxide contrast
agent in awake, behaving macaques at 3 Tesla. NeuroImage 16:283–294.

McLaren DG, Kosmatka KJ, Oakes TR, Kroenke CD, Kohama SG, Matochik JA, Ingram DK,
Johnson SC (2009) A population-average MRI-based atlas collection of the rhesus
macaque. NeuroImage 45:52–59.

Meyer T, Qi X-L, Stanford TR, Constantinidis C (2011) Stimulus Selectivity in Dorsal and
Ventral Prefrontal Cortex after Training in Working Memory Tasks. J Neurosci 31:6266–
6276.

Miyashita Y, Higuchi SI, Sakai K, Masui N (1991) Generation of fractal patterns for probing the
visual memory. Neuroscience Research 12:307–311.

Niki H, Watanabe M (1979) Prefrontal and cingulate unit activity during timing behavior in the
monkey. Brain Res 171:213–224.

Onoe H, Komori M, Onoe K, Takechi H, Tsukada H, Watanabe Y (2001) Cortical Networks
Recruited for Time Perception: A Monkey Positron Emission Tomography (PET) Study.
NeuroImage 13:37–45.

O'Reilly RC (2010) The What and How of prefrontal cortical organization. Trends in
Neurosciences 33:355–361.

Petrides M, Pandya DN (1999) Dorsolateral prefrontal cortex: comparative cytoarchitectonic
analysis in the human and the macaque brain and corticocortical connection patterns. Eur
J Neurosci 11:1011–1036.

Poline J-B, Brett M (2012) The general linear model and fMRI: Does love last forever?
NeuroImage 62:871–880.

Rainer G, Rao SC, Miller EK (1999) Prospective Coding for Objects in Primate Prefrontal
Cortex. J Neurosci 19:5493–5505.

Rapan L, Froudist-Walsh S, Niu M, Xu T, Zhao L, Funck T, Wang X-J, Amunts K, Palomero-
Gallagher N (2023) Cytoarchitectonic, receptor distribution and functional connectivity
analyses of the macaque frontal lobe Badre D, Baker CI, Thiebaut de Schotten M, eds.
eLife 12:e82850.

Sallet J, Mars RB, Noonan MP, Neubert F-X, Jbabdi S, O'Reilly JX, Filippini N, Thomas AG,
Rushworth MF (2013) The organization of dorsal frontal cortex in humans and
macaques. J Neurosci 33:12255–12274.

Sirmpilatze N, Klink PC (2020) RheMAP: Non-linear warps between common rhesus macaque
brain templates. Available at: https://zenodo.org/records/3668510 [Accessed July 10,
2024].

Stern CE, Sherman SJ, Kirchhoff BA, Hasselmo ME (2001) Medial temporal and prefrontal
contributions to working memory tasks with novel and familiar stimuli. Hippocampus
11:337–346.

Suda Y, Tada M, Matsuo T, Kawasaki K, Saigusa T, Ishida M, Mitsui T, Kumano H, Kirihara K,
Suzuki T, Matsumoto K, Hasegawa I, Kasai K, Uka T (2022) Prediction-Related Frontal-
Temporal Network for Omission Mismatch Activity in the Macaque Monkey. Front
Psychiatry 13:557954.

Tang H, Bartolo R, Averbeck BB (2021) Reward-related choices determine information timing
and flow across macaque lateral prefrontal cortex. Nat Commun 12:894.

Uhrig L, Dehaene S, Jarraya B (2014) A Hierarchy of Responses to Auditory Regularities in the
Macaque Brain. J Neurosci 34:1127–1132.

Vanduffel W, Farivar R (2014) Functional MRI of Awake Behaving Macaques Using Standard
Equipment. In: Advanced Brain Neuroimaging Topics in Health and Disease
(Papageorgiou TD, Christopoulos GI, Smirnakis SM, eds), pp Ch. 6. Rijeka: IntechOpen.
Available at: https://doi.org/10.5772/58281 [Accessed March 9, 2023].

Vergnieux V, Vogels R (2020) Statistical Learning Signals for Complex Visual Images in
Macaque Early Visual Cortex. Front Neurosci 14:789.

Walker AE (1940) A cytoarchitectural study of the prefrontal area of the macaque monkey.
Journal of Comparative Neurology 73:59–86.

Wang L, Uhrig L, Jarraya B, Dehaene S (2015) Representation of numerical and sequential
patterns in macaque and human brains. Current Biology 25:1966–1974.

Xu R, Bichot NP, Takahashi A, Desimone R (2022) The cortical connectome of primate lateral
prefrontal cortex. Neuron 110:312-327.e7.

Yamagata T, Nakayama Y, Tanji J, Hoshi E (2012) Distinct information representation and
processing for goal-directed behavior in the dorsolateral and ventrolateral prefrontal
cortex and the dorsal premotor cortex. J Neurosci 32:12934–12949.

Yusif Rodriguez N, McKim TH, Basu D, Ahuja A, Desrochers TM (2023) Monkey Dorsolateral
Prefrontal Cortex Represents Abstract Visual Sequences during a No-Report Task. J
Neurosci 43:2741–2755.

A Sequence Viewing Task (SEQ)

A A A B   D D D A

B No Sequence Task (NoSEQ)

A D A B   C D B A

C Example Stimuli Pools and Timing Templates

Habituation/Standard

A B C D

Deviants/Nonstandard

E F G

Short: 1.1 s

Medium: 1.7 s

Long: 2.3 s

Image Duration
☐ 0.1 s
☐ 0.2 s
☐ 0.3 s

Number / Two and Six Nonstandard Time:

2-item: 1.1 s

6-item: 2.3 s

Rule / Four-Image Nonstandard

4-item: 1.7 s

D SEQ Example Run

▯ = 1 multi-item sequence, bar color = image + time template type   ▮ Fixation Block

Habituation   NISR   Rule Deviant   Number Deviant   Double Deviant

E NoSEQ Example Run

▯ = 1 multi-item group, bar color = time template type   ▮ Fixation Block

Standard Timing, Nonstandard Image Block

Standard Timing   Four-Image Nonstandard   Two- and Six-Image Nonstandard

Habituation Images   Novel Images

A

B

p46d	a46d
p46f	a46f
p46v	a46v

C

y = 31   33   35

R

A

B
Nonstandard > Standard (Hot)
Standard > Nonstandard (Cold)