Title: Repeated behavioral evolution is associated with targeted convergence of gene expression in cavity-nesting songbirds

Authors: Sara E Lipshutz[1,2,3], Mark S Hibbins[1,4,5], Alexandra B Bentz[1,6], Aaron M Buechlein[7], Tara A Empson[1,8], Elizabeth M George[1,9], Mark E Hauber[10,11,12], Doug B Rusch[7], Wendy M Schelsky[11], Quinn K. Thomas[2], Samuel J. Torneo[1], Abbigail M Turner[1,10], Sarah E Wolf[1,13], Mary J Woodruff[1], Matthew W Hahn[1,14], Kimberly A Rosvall[1]

*correspondence to: sara.lipshutz@duke.edu or krosvall@iu.edu

Affiliations:
1: Department of Biology, Indiana University, Bloomington, IN, USA
2. Department of Biology, Loyola University Chicago, IL, USA
3: Department of Biology, Duke University, Durham, NC, USA
4: Department of Ecology & Evolutionary Biology, University of Toronto, Canada
5: Department of Biology, University of Rochester, NY, USA
6: School of Biological Sciences, University of Oklahoma, OK, USA
7: Center for Genomics and Bioinformatics, Indiana University, Bloomington, IN, USA
8: Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN, USA
9: Department of Evolution, Ecology, and Organismal Biology, The Ohio State University, OH, USA
10: Department of Evolution, Ecology, and Behavior, University of Illinois, Urbana-Champaign, IL, USA
11: Illinois Natural History Survey, Prairie Research Institute, University of Illinois, Urbana-Champaign, IL, USA
12: Advanced Science Research Center and Program in Psychology, Graduate Center of the City University of New York, New York, NY, USA
13: School of Biological Sciences, University of Edinburgh, Scotland
14: Department of Computer Science, Indiana University, Bloomington, IN, USA

Author Contributions:

S.E.L. and K.A.R. conceived of the study; S.E.L. led the data collection, supported by A.B.B., A.J.T., E.M.G, M.E.H., M.J.W., S.J.T, S.E.W., T.A.E., and W.M.S. S.E.L., M.S.H., A.B.B., A.M.B., D.B.R., Q.K.T., M.W.H. and K.A.R. analyzed and interpreted the data; S.E.L. M.S.H. and K.A.R. wrote the paper, and all authors contributed to revisions. The manuscript reflects the contributions and ideas of all authors.

Competing Interests: Authors declare no competing interests.

Classification: Biological Sciences, Evolution

Keywords: Competition, Behavioral Physiology, Molecular Evolution, Transcriptomics, Songbird, Aggression

This PDF file includes: Main Text, Figures 1 to 4

Abstract

Uncovering the genomic bases of phenotypic adaptation is a major goal in biology, but this has been hard to achieve for complex behavioral traits. Here, we leverage the repeated, independent evolution of obligate cavity-nesting in birds to test the hypothesis that pressure to compete for a limited breeding resource has facilitated convergent evolution in behavior, hormones, and gene expression. We used an integrative approach, combining aggression assays in the field, testosterone measures, and transcriptome-wide analyses of the brain in wild-captured females and males. Our experimental design compared species pairs across five avian families, each including one obligate cavity-nesting species and a related species with a more flexible nest strategy. We find behavioral convergence, with higher levels of territorial aggression in obligate cavity-nesters, particularly among females. Across species, levels of testosterone in circulation were not associated with nest strategy, nor aggression. Phylogenetic analyses of individual genes and co-regulated gene networks revealed more shared patterns of brain gene expression than expected by drift, but the scope of convergent gene expression evolution was limited to a small percent of the genome. When comparing our results to other studies that did not use phylogenetic methods, we suggest that accounting for shared evolutionary history may reduce the number of genes inferred as convergently evolving. Altogether, we find that behavioral convergence in response to shared ecological pressures is associated with largely independent gene expression evolution across different avian families, punctuated by a narrow set of convergently evolving genes.

Main

Biologists have long been fascinated by phenotypic convergence as a window into the predictability of evolution (1-3). However, our understanding of the molecular mechanisms of behavioral convergence lags behind that of morphological or physiological traits (4-6). Some studies of behavioral convergence find that evolution repeatedly uses the same (or similar) changes in particular genes or pathways (7-9). Alternatively, the building blocks of behavioral convergence may be independent, with lineage-specific mechanisms across replicated evolutionary events (10, 11). Understanding the relative contributions of these two processes requires a comparative approach that embraces the likely polygenic nature of complex behaviors (12-16). However, efforts to connect behavioral evolution to transcriptomics rarely use phylogenetic methods (6), despite the potential for changes in gene expression to shape phenotypic diversification. Likewise, genetic drift may shape gene expression independently of adaptation, and phylogenetic methods are necessary to disentangle these processes.

Territorial aggression is a widespread behavioral trait that is well-suited for evaluating these processes. In vertebrates, aggression is mediated by diverse neuroendocrine mechanisms (17), including the hormone testosterone (18) and its metabolite 17β-estradiol, both of which can act on sex steroid receptors in the brain to promote aggression (19, 20). Aggression is also linked to brain metabolic pathways (7) and G-protein coupled receptor signaling of dopamine, serotonin, and glutamate (21, 22). Many of these candidates are shared across distantly related species (7, 21), demonstrating the potential for conserved or repeatedly evolved mechanisms in the evolution of territorial aggression. However, there is also interspecific variation in the abundance and distribution of these endocrine-molecular building blocks (23, 24), indicating that mechanisms of aggression can diverge over time.

Mechanisms of aggression can also differ within a species, including between males and females (25), both of whom exhibit territorial aggression in many species (26, 27). This provides

a unique opportunity to include both sexes in analyses of behavioral evolution. After all, females
and males share the majority of their genome, yet sex-specific selective pressures can alter
gene expression (28) and hormone secretion (29), which together may shape aggression or the
mechanisms that promote the expression of aggression, the latter of which is still remarkably
understudied in females (30, 31). Critically, in both sexes, aggression shapes access to
resources and mates, and the degree of such competition varies among species (26, 27).
99
Competition for breeding territories is thought to be especially intense for obligate secondary
cavity-nesting birds, which must secure an excavated space inside a tree or other substrate.
They need a cavity to breed, they cannot excavate one themselves, and physical conflicts over
cavities can lead to serious injury or death (32-34). For many cavity-nesting species, both
females and males compete to acquire and defend nesting territories (35-39), and our earlier
case study showed elevated aggression in two species of obligate cavity-nesters, though
without quantitative phylogenetic methods (40). There have been multiple evolutionary
transitions in and out of secondary cavity-nesting across passerines (41, 42), providing a strong
foundation to evaluate the degree of mechanistic convergence in behavioral evolution.
109
Here, we test the hypothesis of evolutionary convergence using a phylogenetic approach that
integrates behavioral, hormonal, and neurogenomic data. We compare species pairs from five
avian families - Hirundinidae (Swallows), Parulidae (Woodwarblers), Passeridae (Sparrows),
Turdidae (Thrushes), and Troglodytidae (Wrens) - which each represent independent origins of
obligate secondary cavity-nesting (Figure 1A; further details in SI §1) (42). Each pair includes
one obligate secondary cavity-nesting species, which regularly nests in artificial nest boxes, and
one species with a more flexible nest strategy, either open cup-nesting or facultative cavity-
nesting. Each species pair is similar in other aspects of its ecology and life history (e.g. foraging
ecology, degree of biparental care). Across species, we compared territorial aggression,
testosterone in circulation, and brain gene expression. By applying quantitative phylogenetic
approaches across these datasets, we evaluate molecular convergence against a null
hypothesis of shared evolutionary history and genetic drift.

122
Figure 1: (A) *Left*: Consensus phylogeny of 10 species from five families, which diverged from a common

ancestor ~44 mya (36-50 mya). Species pairs diverged ~9-20 mya (43). Nestboxes indicate obligate
cavity-nesters, nests represent open nesters, and both symbols together represent facultative cavity-
nesters. *Right*: Sex- and species-level aggression towards a conspecific decoy, measured by the
proportion of 5-sec intervals that contained physical contact during a 5-min aggression assay. Species
listed in descending order: Swallows: tree swallows (*Tachycineta bicolor*), barn swallows (*Hirundo
rustica*), Woodwarblers: prothonotary warblers (*Protonotaria citrea*), yellow warblers (*Setophaga
petechia*), Sparrows: Eurasian tree sparrows (*Passer montanus*), house sparrows (*Passer domesticus*),
Thrushes: Eastern bluebirds (*Sialia sialis*), American robins (*Turdus migratorius*), and Wrens: house
wrens (*Troglodytes aedon*), Carolina wrens (*Thryothorus ludovicianus*). **(B)** Aggression, grouped by nest
strategy. Each point is one assay on a unique free-living individual; box plots convey interquartile range.
Obligate cavity-nesting females were significantly more aggressive than females with more flexible nest
strategies. Illustrations reproduced with permission from Lynx Edicions.

Results

Behavioral convergence among obligate cavity-nesters
Our study occurred during territorial establishment at the beginning of the breeding season,
when birds are defending territories and acquiring mates, but before egg laying. We assayed
aggression in 304 free-living female and male birds in Indiana, Illinois, or Kentucky, USA (SI §1,
Table 1). Each aggression assay lasted 5 minutes and included a conspecific, taxidermic decoy
of either female or male sex, coupled with playback of female or male vocalizations that occur
during natural aggressive interactions (details in SI §2). We measured a variety of aggressive
behaviors and focused on physical attacks (i.e. contact with the decoy). To evaluate the role of
nest strategy as a driver of territorial aggression, we used phylogenetic generalized linear mixed
models (PGLMMs). Briefly, these models correct for covariation in traits among species due to
common ancestry by incorporating phylogenetic relatedness as a random effect. Along with sex
of the territory holder and sex of the decoy, we included nest strategy as a fixed effect with two
levels: obligate cavity-nesting vs. more flexible strategies (including both facultative cavity- and
open-nesting). We also ran three-level models differentiating obligate cavity, facultative cavity,
and open nest strategies (SI §9-11).
152
153 **Table 1:** *Sample sizes for behavior, testosterone, and gene expression in the ventromedial
154 telencephalon (VmT).*

Family	Species	Nest Strategy	Aggression		Testosterone		VmT RNASeq	
			F	M	F	M	F	M
Swallows *Hirundinidae*	Tree Swallow	Obligate cavity	14	24	14	8	7	6
	Barn Swallow	Open cup	10	11	13	14	6	6
Woodwarblers *Parulidae*	Prothonotary Warbler	Obligate cavity	12	19	9	9	6	6
	Yellow Warbler	Open cup	10	20	8	8	6	6
Sparrows *Passeridae*	Eurasian Tree Sparrow	Obligate cavity	8	13	7	8	6	6
	House Sparrow	Facultative	14	17	8	10	6	6
Thrushes *Turidae*	Eastern Bluebird	Obligate cavity	17	20	15	15	6	6
	American Robin	Open cup	9	11	7	12	6	6
Wrens *Troglodytidae*	House Wren	Obligate cavity	16	22	6	11	6	6
	Carolina Wren	Facultative	15	22	6	8	6	6

We found significant effects of nest strategy, sex of the territory holder, sex of the decoy, and
their interaction on *aggression*. Obligate cavity-nesters spent more time attacking the decoy,
compared to species with more flexible nest strategies (PGLMM: p = 0.0011; Figure 1A; Table
S1). This difference in aggression was larger between obligate cavity-nesters and open nesters
relative to facultative cavity-nesters (Figure 1A). Males attacked marginally more than females
(PGLMM: p = 0.058). Notably, sex interacted with nest strategy (PGLMM: p = 0.0064), a pattern
driven by elevated levels of territorial aggression in obligate cavity-nesting females (Figure 1B).
We also found a significant interaction between sex of the territory holder and sex of the decoy
(PGLMM: p = 0.011), such that male territory holders were more aggressive towards male
decoys, whereas female territory holders attacked female and male decoys with similar levels of
aggression (Figure S1; Table S1).
167
Additionally, we measured the average distance from the focal individual to the decoy, to
confirm each subject was present and engaged during a trial, regardless of whether they were
aggressively attacking. *Distance from the decoy* was not related to nest strategy (PGLMM: p =
0.22), sex (p = 0.70), nor their interaction (p = 0.39) (Table S2; Figure S2). This result indicates
that decoy placement was a salient stimulus, in that all species were in audio-visual proximity to
the simulated intruder (average distance = 5.0m), but obligate cavity-nesters spent more time
attacking the decoy compared to their close relatives with more flexible nesting strategies
(Figure 1, Table S1).
176
Testosterone levels are not associated with nest strategy, nor aggression, across species
Next, we sought to measure levels of testosterone and gene expression as potential
physiological drivers of behavioral convergence. We focused our analyses on constitutive
hormonal and neurogenomic states that are representative of unprovoked, free-living animals (n
= 196 hormone samples, n = 121 brain gene expression samples; Table 1). We followed the
assumption that social elevation of testosterone peaks around 30-45 min after HPG axis
activation (44, 45), and that socially responsive genes likewise show peak transcriptional
responses 30-60 minutes after a stimulus (46). To maximize our number of aggression assays
while meeting these assumptions, our sample collection took three approaches: a) *passive*
collection, in which we set up a mist-net or trap in the target territory and waited to capture the
focal individual without any stimulation (testosterone: n = 108, brain: n = 51), b) *immediate*
collection, in which we sampled individuals immediately after a short aggression assay
(testosterone: n = 55, brain: n = 43), and c) *delayed* collection in which we sampled individuals
several days after a short aggression assay (testosterone: n = 33, brain: n = 27). We found no
differences in testosterone among immediate, passive, or delayed sample collection
approaches (ANOVA: $F_{2, 184}$ = 0.011, p = 0.90; Figure S3A), and therefore combined these
samples.
194
We previously identified a positive correlation between territorial aggression and testosterone
within female tree swallows (S = 2, p = 0.0028, rho = 0.96) (40), which are included in this
study, but we did not find this relationship in male tree swallows. To explore whether similar
patterns apply to other species sampled here, we conducted Pearson's correlations across
species for each sex, and Spearman's correlations within each family for each sex, given the
reduced sample size. We did not find any relationship between testosterone and aggression in
Eastern bluebirds, house wrens, Carolina wrens, prothonotary warblers, nor yellow warblers (p
> 0.13), and we did not have enough data on testosterone and aggression from the same
individuals for the other species. Considered collectively for all samples, we did not find a
significant relationship between territorial aggression and testosterone for females (t = 1.87, df =
25, p = 0.073, r = 0.35), nor for males (t = 1.17, df = 35, p = 0.25, r = 0.19). Testosterone was

not related to nest strategy (PGLMM: p = 0.80) nor the interaction between nest strategy and
sex (PGLMM: p = 0.85; Figure S4, Table S3). As expected, we found that males had
significantly higher levels of testosterone in circulation than females (PGLMM: p < 0.0001).
209
Neurogenomic mechanisms of behavioral convergence
Finally, we examined convergent evolution in brain gene expression. Using RNA-seq, we
measured mRNA abundance for 10,672 orthologous genes expressed in all 10 focal species in
the ventromedial telencephalon (Table 1, Figure S5); this region contains core nodes of the
vertebrate social behavior network, which regulates behaviors including aggression (47). The
number of differentially expressed genes generally increased with divergence time between
species pairs within each family (p = 0.08, R^2 = 0.83; SI §5, Figure S8, Table S4), underscoring
the need for phylogenetic methods. Using these expression data, we took three analytical
approaches to understand the mechanisms underlying convergent behavioral evolution.
219
Gene expression is highly concordant between pairs of families, but involves different sets of genes
across the phylogeny
We used the Rank Rank Hypergeometric Overlap (RRHO) approach to compare the magnitude
and direction of gene expression differences among species pairs. If the same genes are
repeatedly targeted by selection for the obligate cavity-nesting strategy, these genes should be
differentially expressed in the same direction for cavity-nesters across multiple family
comparisons. As a null hypothesis, we also generated permuted differential expression datasets
by randomly sampling the log2FoldChange and associated p-values from the observed values
with replacement for each gene. This was done independently for each family. The permuted
datasets were then given to the same RRHO pipeline used for the empirical datasets.
230
In contrast to the largely random distributions of overlap from the randomly permuted dataset
(Figure S9), RRHO for our empirical dataset revealed significantly concordant patterns of gene
expression (all p-values < 0.01, permutation test, Figure 2, Table S5). Genes that were more
highly expressed in the obligate cavity-nester of one species pair were also more highly
expressed in the obligate cavity-nester of another species pair, and vice-versa for genes with
lower expression (Data S4), evidenced by clustering in the upper-right and lower-left quadrants
of the RRHO heatmaps (Figure 2C). This concordant pattern applied to an average of 1390
genes per family comparison (Figure S12), amounting to ~13.0% of genes studied, suggesting a
large number of genes evolving in a similar direction of expression *within each family.* In
comparison, the randomly permuted dataset had an average of 562 concordant genes per
family comparison (~5.3% of total genes). The empirical dataset had more than twice as many
concordant genes as expected by random chance from the permuted dataset. Therefore, our
empirical RRHO data for each family comparison are overwhelmingly concordant, beyond the
null expectation.
245
This high level of concordance in gene expression did not persist across multiple family
comparisons, indicating a lack of convergence in the *identity* of genes differentially expressed
between species pairs. The number of shared genes decreased rapidly with increasing inclusion
of additional family comparisons (Figure 2A). Specifically, about 1,000 genes (~10% of
orthologs) displayed concordant expression patterns among two or three families, and 497
genes (~5%) were shared by at least 5 out of 10 family comparisons at a time (Figure 2A).
Relative to the empirical data, the randomly permuted data had more shared genes across 2
family comparisons, and fewer shared genes across 5+ family comparisons (Figure S9). Gene
Ontology (GO) analyses of the 497 shared genes (Data S4, S5) revealed no significant
functional enrichment. This pattern indicates that – despite a significant degree of concordance

between pairs of species – concordantly expressed genes are not repeatedly recruited from any
particular set of known biological processes. However, we must recognize that GO databases
are biased towards model organisms, and a lack of GO enrichment does not necessarily
indicate an absence of a shared biological function.
260
Across all ten family comparisons, only 11 genes were shared, meaning that 0.1% of 10,672
orthologs exhibited complete concordance across five independent origins of obligate cavity-
nesting (Table S6). This number was still larger than the genes shared across families using the
randomly permuted dataset, which was only 1 gene. Some of these 11 globally concordant
genes have established connections with behavior or brain function (48). For instance, *EMC3,*
NPTX1, and RGS19 are involved in neurotransmitter reception, *ATP6V0E1* and *DDX56* have
ATPase activity, and most of the completely concordant genes have established connections
with Alzheimer's (*ARRDC4, CTNND1, NPTX1, WIPF2*), addiction (*RGS19*) or other neurological
disorders (*AFAP1L1, PQBP1*) (48). s genes could be functionally validated in relation to cavity
nesting or aggression in the future. Overall, these results from RRHO indicate a high degree of
independent gene expression evolution in association with the convergent evolution of obligate
cavity nesting.
273
We repeated these RRHO analyses with females and males separately, because sex-specific
selection pressures could generate concordance that would be masked by analyzing both sexes
together. We found similarly high concordance in these sex-specific analyses, in that significant
gene overlap was concentrated primarily in the upper-right and lower-left quadrants. However,
the degree (indicated by the scale of the heatmap) and direction (concentration of significant
overlap in the upper-right vs. lower-left) of concordance differed between sexes (Figure 2A,
Figure S10, S11). Males had more cases of higher expression in the obligate cavity nester of
both families (upper-right), whereas females had more cases of lower expression (lower-left).
282

283
Figure 2: Concordance in differential expression based on rank-rank hypergeometric overlap (RRHO).
Heatmap colors reflect adjusted hypergeometric -log(p-value). Larger values indicate more concordance
in expression between families. Nestboxes indicate obligate cavity-nesters, nests represent open nesters,
and both together represent facultative cavity nesters. **(A)** Number of concordantly expressed genes
shared across family comparisons (i.e. across heatmaps) for males (blue), females (pink) and both sexes
combined (green). Solid lines indicate higher expression in obligate cavity-nesters, dashed lines indicate
lower expression **(B)** Key for interpreting individual heatmaps. Each quadrant of the key corresponds to a
quadrant of an individual heatmap, shown in **(C)** Each pixel within the heatmaps contains two sets of
approximately 100 genes being compared between the two families; the color legend indicates the log p-
value of the overlap between these gene sets, with a higher value indicating stronger overlap.
294
Nest strategy and aggression are associated with convergently evolving genes
Next, we used PGLMMs to model expression of all ~10k orthologs as a function of nest
strategy, sex, aggression, sex-by-nest-strategy interactions, and aggression-by-nest-strategy
interactions. To identify convergent expression evolution, these phylogenetic models explicitly
account for shared expression due to evolutionary history. We expect some degree of
convergence in expression evolution due to random chance (i.e. genetic drift along the
phylogeny), so we employed a false discovery rate (FDR) correction to test this null hypothesis.
Many genes showed some relationship with obligate cavity nesting, aggression, or a relevant

interaction, supporting a hypothesis of convergent expression evolution. Specifically, there were
234 genes associated with obligate cavity-nesting, 79 genes associated with aggression, and
five genes overlapped among these. Of these five genes (*RNASEH2B, ERN1, UGGT2, TAF1B,*
and HIGD1A), the former four relate to protein or RNA processing, the latter four relate to stress
responses, and all have connections with neurodegenerative disorders (48-52). An additional 62
genes exhibited an interaction between nest strategy and aggression. 76 genes exhibited an
interaction between sex and obligate cavity-nesting, with approximately equal numbers of genes
that had a stronger association in females vs. males (Table S7, Data S6). Sex had the largest
influence on expression variation and was associated with 510 genes (Table S7), the majority of
which had higher expression in males than females. Genes with sex-biased expression were
significantly enriched for cellular metabolic process.
314
Because some nest-strategy and aggression-associated patterns may be driven by expression
in one or two species, we developed additional criteria to pinpoint a robust set of convergently
evolving genes. We retained only those that were (i) significantly different in expression (by *t*-
test) between species in at least 3 out of 5 family comparisons; and (ii) different in a consistent
direction (i.e. all higher or all lower expression) with respect to nest strategy in all families with a
significant difference. 168 of 234 genes met these criteria of convergence for nest strategy
(Data S8). Further restricting our analyses to all 5 family comparisons, we identified 40 genes
(0.4% of orthologues) associated with nest strategy. Several convergent genes associated with
nest strategy have some connection to ATP and mitochondrial function (*ATP1B1, PITRM1,*
SLC25A24), as well as behavior or psychiatric risk (*TRMT1L, NT5C2, BCHE*) (48).
325
326 We also used PGLMMs to examine the expression of a subset of candidate genes related to
327 testosterone and steroid hormone signaling in our transcriptome dataset, but we did not find a
328 relationship with obligate cavity nesting (SI §9).
329
Nest strategy and female aggression are associated with convergent gene networks
Complex phenotypes are often regulated by subtle but coordinated changes in gene networks,
which may also be targets of selection (53), and approaches that analyze genes individually
may miss important emergent properties. Therefore, we used weighted gene co-expression
network analyses (WGCNA) to estimate co-expression across all 10 species (SI §8), yielding 93
networks of correlated genes (Data S10). Using each network's eigengene (akin to a principal
component of expression), we ran PGLMM with fixed effects of nest strategy, aggression, sex,
and their interactions. After accounting for phylogeny, and with FDR correction, two networks
showed significant associations that are unlikely to occur by drift alone. Expression in the red
network was higher in males than in females (Figure S13a), and 147 of 193 genes (76%) were
on the zebra finch Z chromosome. Expression in the tan4 network was associated with nest
strategy (Data S9, Table S8), with lower eigenvalues in each of the five independent origins of
obligate cavity-nesting (Figure 3A). Though not significantly enriched for any GO terms (Table
S8), several genes in the tan4 network related to mitochondrial function and energy metabolism,
cognition, and stress response (Figure 3C).
345
346

Figure 3. (A) Weighted gene co-expression network analysis (WGCNA) eigengene values for the tan4 network, which included 44 genes whose expression was significantly associated with obligate cavity-nesting relative to non-obligate nesting strategies. Individual points represent the eigengene value for a sampled individual, colored by sex. Nestboxes indicate obligate cavity-nesters, nests represent open nesters, and both together represent facultative cavity nesters. **(B)** Heatmap of log-scaled genes in tan4 network. Columns represent individuals, grouped by sex and nest strategy. Color scale indicates log of gene expression. **(C)** Network depicts genes with network membership |>| 0.6. Illustrations reproduced with permission from Lynx Edicions.

Because sexes have the potential to differ in mechanisms of behavior, we repeated this analysis for each sex separately (SI §10). In this sex-specific analysis, we identified two female-specific gene networks correlated with species-level aggression – pink4 and sienna4 (Pearson's correlation: r |>| 0.78, p < 0.0001, Figure 4, Table S9). These networks contained genes with established connections to aggression, including DRD3, a dopamine receptor in the pink4 network, and GRIA2, a glutamate receptor in the sienna4 network. Finding aggression-associated networks specific to females further suggests that there are multiple gene regulatory routes to aggression, including those that may differ between sexes.

Figure 4. For females only, WGCNA network eigengenes for the **(A)** pink4 network and **(B)** sienna

network were associated with average, species-level aggression towards a conspecific decoy, measured

by the proportion of 5-sec intervals that contained physical contact during a 5-min aggression assay.

Networks for **(C)** pink4 and **(D)** sienna4 depict genes with network membership |>| 0.6.

Gene expression provides many unique and some shared routes to convergent behavioral

evolution

To identify candidate genes that are robustly (i.e. independently across multiple analyses)

associated with nest strategy and/or aggression, we integrated results across differentially

expressed genes, individual gene- and network level PGLMMs, and concordant genes from at

least 5 out of 10 RRHO comparisons (Table S11). For nest strategy-associated candidates, 27

out of 44 genes (61%) in the tan4 network recapitulated convergent individual genes in the

PGLMM analysis, a strong signal that this gene set is consistently associated with the

convergent evolution of obligate cavity-nesting. Out of the 221 differentially expressed genes

shared among species pairs, 13 genes overlapped with nest-strategy associated individual

PGLMM analyses and the tan4 network. We also identified 6 aggression-associated genes that

overlapped with nest-strategy associated analyses, including 3 with concordant RRHO genes, 2
with individual PGLMM analyses, and 1 gene, *TAF1B*, which overlapped with both the tan4
network and individual PGLMM analyses. Altogether, these results suggest that repeated use of
a small core set of genes may have contributed to the evolution of obligate cavity nesting and
associated changes in behavior.
387
To place our results in a broader context, we compared our list of aggression-associated genes
to prior studies (Data S12), including those on the brain's response to an aggressive challenge
(7, 54) and those that are known to mediate aggression in diverse species, including humans
(22, 55). Using a permutation analysis, we found significant overlap between our 79 aggression-
associated genes and another study's network of 290 genes that were sensitive to experimental
competition in the tree swallow brain (SI §11, Data S12, Table S12, Figure S19; (54)). Of the
seven genes in common between these datasets (Table S13), *DPF3*, *PLXND1*, and *ZIC4* are
involved in brain development, *CHN2* is associated with schizophrenia, *SPHKAP* is associated
with neuroblastoma and apoptosis, and *GDF10* promotes neural repair after stroke (48), again
linking aggression to key elements of brain function.

Discussion

Striking phenotypic similarities among distantly related organisms exposed to similar ecological
pressures yields critical insights into the mechanisms of evolution. However, investigations of
the proximate mechanisms by which natural selection generates evolutionary change are often
limited to traits with a simple and well-understood genetic basis (2, 3, 56), and less is known
about the underpinnings of convergence in complex behavioral traits. A particularly unresolved
question is the relative contributions of repeated vs. independent molecular changes to
convergent behavioral evolution. Across five avian families with independent origins of obligate
secondary cavity-nesting, we find behavioral convergence in territorial aggression, but limited
transcriptomic convergence in the brain involving a small, core set of genes. These results
highlight how complex behaviors can arise from largely independent evolution, punctuated by a
few specific cases of repeated molecular changes in response to shared ecological pressures.
410
Across ten species, those with obligate cavity-nesting strategies exhibited greater physical
aggression, though all species were present and responsive to the simulated intruder. Among
females, aggression was particularly high for obligate cavity nesters, consistent with the
hypothesis that female aggression is adaptive during competition for nesting territories (27, 37,
40, 57). Females were aggressive towards decoys of both sexes, whereas males were more
aggressive towards their own sex. Conspecific aggression may serve multiple functions, most
likely nest site defense or mate guarding (27, 58), considering the timing of stimulated intrusions
during the early spring period of territorial establishment before egg laying. Our results
complement a macroevolutionary study finding that cavity-nesters display more territorial
behaviors against heterospecifics (59), underscoring multiple contexts linking territorial
aggression with a cavity-nesting strategy.
422
Despite behavioral convergence, we did not observe higher levels of testosterone in obligate
cavity-nesters. Testosterone-focused hypotheses have dominated aggression-related research
for decades (18), yet we find that testosterone does not explain species-level differences in
female or male territorial aggression, at least when birds are sampled at a constitutive state,
without a prolonged social challenge (40). Recent phylogenetic analyses have found that

breeding season length and mating system predict testosterone levels across species (60), but
these life history traits are similar across our focal species. Interspecific variation in testosterone
does not track interspecific variation in territorial aggression, for either sex. This finding aligns
with recent perspectives that variation in testosterone is context-dependent (61, 62), and that
testosterone is but one of many potential mechanisms regulating aggression (63, 64).

Using multiple phylogenetically informed approaches, we find that the convergent evolution of
obligate cavity-nesting is associated with a small set of convergently expressed genes in the
brain, alongside a larger set of lineage-specific genes shared only by some families or species.
With these quantitative phylogenetic approaches, we find changes that occur more than is
expected due to shared evolutionary history and random chance. These patterns could be
driven by expression evolution in either the obligate-cavity or open nesters, but we are currently
unable to resolve this directionality, given that we did not explicitly construct ancestral states for
expression values. RRHO analyses revealed striking patterns of expression concordance
between family comparisons, but only 0.1% of orthologs (11 genes) were associated with
obligate cavity nesting across all comparisons. Single-gene and network PGLM analyses
likewise revealed a small set of convergently evolving genes (0.4%; ~40 genes). These
proportions are similar to a recent phylogenetic study on muscle gene expression and courtship
behavior in birds (14). However, they represent less convergence than most other studies of
brain gene expression and behavioral evolution (9, 13, 65), which report that 4-6% of expressed
orthologs are convergently evolving alongside behavior. Though our study differs by the number
of taxa and/or degree of evolutionary divergence, another key difference is our explicit
phylogenetic approach to identify convergently evolving genes, which statistically eliminates
shared patterns of gene expression that stem from common ancestry and more directly tests for
convergence. Thus, ours and other phylogenetic approaches may reduce the number of genes
inferred as convergently evolving. Altogether, our results suggest that the convergent evolution
of complex behavioral phenotypes can be underlain by mostly independent, lineage-specific
changes in gene expression, with some convergent expression in a small, core set of genes
shared across species.

This constellation of changes in gene expression may be associated with obligate cavity-nesting
or aggression, and our PGLMMs disentangle these effects directly. Genes associated with nest
strategy or aggression exhibited uncoupled regulatory evolution, with differing degrees of
convergence and low overlap in gene identity. These patterns suggest that there are diverse
neuromolecular correlates of aggression, including many that differ among species. Notably, we
also found significant gene overlap with our aggression-associated genes and a previously
published socially-sensitive gene network (54), indicating concordance between short-term
functional responses to competition and long-term evolutionary change in competitive
environments. Though our study focused on genes constitutively expressed during territorial
establishment, future work should examine convergence in genes whose expression is
activated or suppressed by acute social challenges, particularly since such environmentally
sensitive genes may also play a key role in behavioral evolution (66).

By including both females and males, our study also assesses key elements of sex-specific
evolution. Across species, we find sex-biased gene expression, for which males had higher
expression that females. In birds, males carry two copies of the Z chromosome, and dosage
compensation of the Z chromosome may be incomplete or absent (67), leading to higher

expression of Z-linked genes in males than females. This mechanism likely explains our results
for the 'red' network of co-regulated genes, which had higher expression in males than in
females. Higher expression in males could also stem from stronger sexual selection (68), or
greater constraint on the evolution of reduced expression (i.e. floor effects). Our sex-specific
analyses further evaluate the potential for sex-specific behavioral and mechanistic convergence.
Consistent with this hypothesis, we found that female obligate cavity nesters are more
aggressive than females of both facultative and open-cup nesting species – a pattern that was
not recapitulated in males. We also identified two female-specific gene networks that correlated
with aggression. These sex-specific patterns suggest that female aggression has evolved
differently from male aggression in obligate cavity nesting species, again underscoring multiple
regulatory routes to behavioral convergence.
486
In summary, convergent evolution provides a natural experiment to identify the genomic
underpinnings of complex phenotypes, and our work highlights the power of comparative
approaches to uncover these mechanistic bases of convergence (13, 14, 21). Though future
experiments are needed to functionally validate the nesting- and aggression-associated patterns
we identify, our study is among the first to use phylogenetically explicit models of brain gene
expression to understand the evolution of a complex and continuously varying behavioral trait.
Our findings support the hypothesis that there are largely independent evolutionary routes to
building an aggressive bird, layered atop a small set of convergently evolving genes.
495

Methods

497 This work was approved by Indiana University Bloomington IACUC #18-004 and #21-003, and
498 relevant federal (MB59069B) and state permits (IL: W20.6355, KY: SC1911001, IN: 18-030, 19-
499 272, 2545). Unless otherwise noted, analyses used R version 4 (R-Core-Team 2019).
500
Assay of aggression
For cavity-nesting species, we placed the conspecific decoy on the nestbox and hung a Bluetooth
speaker nearby. For non-cavity-nesters, we located nest sites or observed individuals for at least an
hour to determine where they spent their time. For facultative cavity nesting species, including Carolina
wrens and house sparrows, some aggression assays were conducted on individuals for which the nest
could not be located, and therefore the nest type could not be confirmed. We compared levels of
physical aggression and distance from decoy between individuals with a known nestbox vs. these other
individuals and found no significant differences between groups for either species (p > 0.19). Additional
details on audio stimuli and decoys can be found in SI §2. We played a conspecific vocal lure to
capture the attention of the focal individual and waited 30 s before beginning the 5-min aggression
assay. We measured a suite of aggressive behaviors (see SI §2) and focused on the proportion of the
trial spent physically attacking the decoy. We calculated a maximum attack score of 60, based on the
number of 5-second (sec) intervals that contained any physical contact. To visualize this behavior, we
converted attack scores to a proportion of the trial spent attacking (# intervals including attack/ total #
intervals x 100). We also measured distance from the focal individual to the decoy to confirm that all
focal territory holders were present and engaged with the simulated intruder. We evaluated the effects
of nest strategy, sex, decoy sex, and their interaction on physical attacks and distance from the decoy
using phylogenetic linear mixed models (details below and in SI §9). For downstream analyses, we
include attack averages among species and sex categories as a fixed effect, referred to simply as
'aggression.'
521
Sample collection

For brain gene expression, we euthanized individuals with an anaesthetic overdose of isoflurane and
rapid decapitation. We collected trunk blood into heparinized BD Microtainers (#365965). Using tools
cleaned with RNAse-away and 95% ethanol, we immediately dissected out whole brains and flash froze
on powdered dry ice within 8 min 26 sec ± 12 sec of euthanasia followed by storage at -80°C. Some
additional hormonal samples for were obtained from barn swallows via brachial venipuncture (n = 4
females, n = 8 males), and sex was confirmed from blood via PCR (69). Blood was stored on ice packs
in the field and later centrifuged for 10 min at 10,000 rpm. Plasma was separated and stored at -20°C.
See SI §3 for details on enzyme immunoassays.
531
Immediate collections occurred an average of 18 min and 25 sec after the assay started. There was no
relationship between latency to sampling and testosterone (Pearson's correlation: r = 0.031, p = 0.84;
Figure S3B), or brain gene expression (r |<| 0.32, p > 0.13). For the facultative cavity-nesters, we also
found no difference in testosterone between individuals nesting in nestboxes compared with individuals
whose nest type was unknown, for either species (p > 0.51), and no difference in brain gene expression
for the tan4 eigengenes (p > 0.88).
538
Brain dissection and RNA isolation
For 6 to 7 individuals per sex per species, we dissected whole brains into functional regions following
(70). We removed the cerebellum, hindbrain, optic chiasm, optic tecta, and hypothalamus to the depth
of the anterior commissure. Gene expression analyses focused on the ventromedial telencephalon
(VmT), which contains nodes in the vertebrate social behavior network (47) including the extended
medial amygdala, bed nucleus of the stria terminalis, and lateral septum (Figure S5). We collected this
region by removing ~1mm of the ventromedial portion of the caudal telencephalon. We extracted total
RNA using Trizol (Invitrogen), and resuspended RNA in UltraPure water. Tape Station Bioanalyzer
(Agilent) on 121 samples showed RIN = 8.56 ± 0.05 (mean ± SE).
548
Sequencing, alignment, and mapping
We submitted total RNA to the Indiana University Center for Genomics and Bioinformatics,
using paired end sequencing (Illumina NextSeq500, 75 cycle sequencing module) to generate
an average of ~28.6 million reads per sample (Data S1). We used Trinity version 2.13.2 (71) to
assemble transcriptomes per species, which we aligned to the zebra finch (*Taeniopygia guttata*)
proteome (NCBI assembly GCF_003957565.2_bTaeGut1.4.pri_protein.faa) (72) using tblastn v.
2.2.9. See SI §4 for additional bioinformatics details on alignment, mapping, and downstream
analyses. We identified 10,672 orthologous genes with high confidence in all 10 species. . We
normalized counts (Data S2) and performed differential gene expression using DESeq2 (73) (SI
§5).
559
Rank-Rank Hypergeometric Overlap
To identify shared transcriptomic patterns in obligate cavity-nesting species across pairs of families, we
used the R package Rank-Rank Hypergeometric Overlap (RRHO) (74) (details in SI §8). RRHO uses
normalized counts to rank genes based on the direction and magnitude of expression difference
between groups, in our case each obligate cavity-nester and its species pair within the same family.
Hypergeometric tests evaluate overlap among these ranked lists, visualized as heatmaps of p-values
from 100 permutations. The pipeline also identifies the set of genes with the highest degree of
concordance for each comparison.
568
Weighted Gene Co-Expression Network Analysis
We constructed gene networks for both sexes combined using WGCNA (75). We used the normalized
counts from DESeq2 and filtered out genes with <15 norm counts in 90% of the samples. We
generated a signed hybrid network by selecting a soft threshold power (β) = 6, in accordance with

scale-free topology. We calculated a minimum network size of 30 and used a biweight midcorrelation
(bicor) function. We merged modules in Dynamic Tree Cut using a threshold of 0.25. Genes with an
absolute network membership value >0.6, which indicates high network connectivity, were assessed for
enrichment of biological processes in PantherGO (76), using our list of 10,672 orthologs as a reference
set. Networks of interest were visualized in Cytoscape v3.10.1 (77). See SI §8 for gene network
analyses run separately on females and males.
579
Phylogenetic Generalized Linear Mixed Models
Phylogenetic generalized linear mixed models (PGLMM) used MCMCglmm (78), with the phylogenetic
covariance matrix from the consensus tree as a random effect (SI §9). Our consensus tree (Figure 1A)
made use of the BirdTree.org (79) resource to obtain 1000 ultrametric trees from Ericson All Species.
We constructed a majority rule consensus topology from this tree set using the *ape* (80) function
'consensus', and inferred branch lengths using the *phytools* (81) function 'consensus.edges' with the
least squares method. For each PGLMM, we evaluated MCMC convergence using Heidelberger and
Welch's convergence diagnostic as implemented by the heidel.diag function in the R package coda.
588
To evaluate whether nest strategy predicted aggression, average distance from the decoy, and
testosterone, we fit models with nest type, sex, and their interaction as fixed effects. Our
aggression dataset contained many individuals that never attacked, and so we used a zero-
inflated binomial distribution, with random effect and residual variance priors fixed at 1, the
default burn-in period of 3000, and 2,000,000 iterations. For this aggression analysis, technical
limitations in using this specific model with a zero-inflated binomial distribution restricted our
ability to specify obligate cavity-nesting as the baseline factor level; therefore, we combined
facultative nesting and open nesting into a single category to contrast with obligate cavity-
nesting. For all other PGLMMs, we fit models containing either two (obligate vs. non-obligate) or
three (obligate vs. facultative vs. open) factor levels (SI §9). For these other models, we fit a
standard Gaussian model with no priors, the default burn-in, and 100,000 iterations.
600
To predict nest-strategy and aggression associated genes from our set of ~10k orthologs, we
included species/sex-level aggression sex, nest strategy, nest strategy by sex interaction and
nest strategy by aggression interaction as fixed effects. After fitting a model for the log-
expression of each gene, we applied a Benjamini-Hochberg false discovery rate correction in
the *statsmodels* package (82) in Python version 3.7.11. Since our PGLMMs could not estimate
p-values with precision lower than 1×10^{-4} (due to computational constraints), we set all raw p-
values reported as less than this value to exactly 1×10^{-4}. Since the Benjamini-Hochberg
procedure relies on the relative ranks of raw p-values, rather than their magnitudes, this should
not significantly affect our FDR correction.
610
611   To identify gene networks associated with nest strategy, sex, and their interaction we ran a
612   similar phylogenetic analysis using the first eigengene value drawn from a weighted gene co-
613   expression network analysis (SI §10).
614
615   Our PGLMM-related scripts are available at
616   https://github.com/slipshut/CavityNesting/tree/main/PGLMM
617
618   *Functional interpretation*
619   We inferred gene function based on genecards.org (48) and GO analyses. We inferred GO terms from
620   *Homo sapiens* because its ontologies are orthologous to, but more complete than avian references.

621 We conducted an overrepresentation analysis of non-redundant biological process GO terms in
622 PantherGO (76), using our list of orthologs as a reference set.
623
624 We also conducted a custom permutation analysis to evaluate the degree of overlap between our
625 aggression-related genes and published genomic studies of aggression (7, 22, 54, 55) (Data S12; SI
626 §11). Our enrichment scripts are available at
627 https://github.com/slipshut/CavityNesting/tree/main/Enrichment

Acknowledgements

For facilitating fieldwork, we thank the Indiana Department of Natural Resources, IU's Research
and Teaching Preserve, Thomas Rothfus and Chris Hagie at Therkilsden Field station, Auriel
Fournier and Steve Havera at Forbes Field Station, Jeff Hoover, Nick Antonson, Hannah Scharf,
and Angela Funk. For vocal stimuli, we thank Matt Wilkins, Dustin Reichard, Cara Krieg, and
Elsa Chen. Thanks to Emmi Mueller for help with 3D printing; Sumitha Nallu and Jie Huang for
TapeStation and library preparation; Emma Dossey Curole and Jace Kuske for assistance in the
lab; Becca Young for assistance with data analysis; Amanda Clark for assistance with data
visualization; and our reviewers for their feedback.

Funding

638 US National Science Foundation (NSF) grant DBI-1907134 (to S.E.L.), DBI-2146866 (to
639 M.W.H.), IOS-1953226 (to M.E.H.), and CAREER grant IOS-1942192 (to K.A.R.), as well as
640 support from IU Biology, Loyola Biology, and Duke Biology.

Data Accessibility

642 Raw sequence reads and count data can be obtained from the Gene Expression Omnibus
643 database (GEO accession number GSE244480). Supplementary information, including R
644 scripts and trait data can be obtained from Github: https://github.com/slipshut/CavityNesting.
645
646
647 <div align="center">References Cited</div>
648

1. C. R. Darwin, *On the origin of species* (Harvard University Press, Cambridge, MA, 1859).
2. E. B. Rosenblum, C. E. Parent, E. E. Brandt, The Molecular Basis of Phenotypic
Convergence. *Annual Review of Ecology, Evolution, and Systematics* **45**, 203-226 (2014).
3. J. Arendt, D. Reznick, Convergence and parallelism reconsidered: what have we learned
about the genetics of adaptation? *Trends Ecol Evol* **23**, 26-32 (2008).
4. J. R. Gallant, L. A. O'Connell, Studying convergent evolution to relate genotype to
behavioral phenotype. *J Exp Biol* **223** (2020).
5. N. Jourjine, H. E. Hoekstra, Expanding evolutionary neuroscience: insights from
comparing variation in behavior. *Neuron* **109**, 1084-1099 (2021).
6. K. A. Rosvall, Evolutionary endocrinology and the problem of Darwin's tangled bank.
Horm Behav **146**, 105246 (2022).
7. C. C. Rittschof *et al.*, Neuromolecular responses to social challenge: common
mechanisms across mouse, stickleback fish, and honey bee. *Proc Natl Acad Sci U S A*
111, 17929-17934 (2014).

8.   A. L. Toth *et al.*, Brain transcriptomic analysis in paper wasps identifies genes associated
with behaviour across social insect lineages. *Proc Biol Sci* **277**, 2139-2148 (2010).
9.   A. R. Pfenning *et al.*, Convergent transcriptional specializations in the brains of humans
and song-learning birds. *Science* **346**, 1256846 (2014).
10.   A. S. Mikheyev, T. A. Linksvayer, Genes associated with ant social behavior show distinct
transcriptional and evolutionary patterns. *Elife* **4**, e04775 (2015).
11.   M. R. Warner, L. Qiu, M. J. Holmes, A. S. Mikheyev, T. A. Linksvayer, Convergent eusocial
evolution is based on a shared reproductive groundplan plus lineage-specific plastic
genes. *Nat Commun* **10**, 2651 (2019).
12.   E. A. Boyle, Y. I. Li, J. K. Pritchard, An Expanded View of Complex Traits: From Polygenic
to Omnigenic. *Cell* **169**, 1177-1186 (2017).
13.   R. L. Young *et al.*, Conserved transcriptomic profiles underpin monogamy across
vertebrates. *Proc Natl Acad Sci U S A* **116**, 1331-1336 (2019).
14.   J. B. Pease *et al.*, Layered evolution of gene expression in "superfast" muscles for
courtship. *Proc Natl Acad Sci U S A* **119**, e2119671119 (2022).
15.   M. C. Harrison *et al.*, Hemimetabolous genomes reveal molecular basis of termite
eusociality. *Nat Ecol Evol* **2**, 557-566 (2018).
16.   Z. V. Johnson *et al.*, Cellular profiling of a recently-evolved social behavior in cichlid
fishes. *Nature Communications* **14**, 4891 (2023).
17.   R. J. Nelson, B. C. Trainor, Neural mechanisms of aggression. *Nat Rev Neurosci* **8**, 536-
546 (2007).
18.   J. C. Wingfield, R. E. Hegner, J. Dufty, Alfred M. , G. F. Ball, The "Challenge Hypothesis":
Theoretical Implications for Patterns of Testosterone Secretion, Mating Systems, and
Breeding Strategies. *The American Naturalist* **136**, 829-846 (1990).
19.   B. M. Horton *et al.*, Estrogen receptor alpha polymorphism in a species with alternative
behavioral phenotypes. *Proc Natl Acad Sci U S A* **111**, 1443-1448 (2014).
20.   K. A. Rosvall *et al.*, Neural sensitivity to sex steroids predicts individual differences in
aggression: implications for behavioural evolution. *Proc Biol Sci* **279**, 3547-3555 (2012).
21.   M. C. Saul *et al.*, Cross-species systems analysis of evolutionary toolkits of neurogenomic
response to social challenge. *Genes Brain Behav* **18**, e12502 (2019).
22.   Y. Zhang-James *et al.*, An integrated analysis of genes and functional pathways for
aggression in human and rodent models. *Mol Psychiatry* **24**, 1655-1667 (2019).
23.   A. M. Kelly, L. C. Wilson, Aggression: Perspectives from social and systems neuroscience.
Horm Behav **123**, 104523 (2020).
24.   L. A. O'Connell, H. A. Hofmann, Evolution of a vertebrate social decision-making
network. *Science* **336** (2012).
25.   K. A. Rosvall, Proximate perspectives on the evolution of female aggression: good for
the gander, good for the goose? *Philos Trans R Soc Lond B Biol Sci* **368**, 20130083 (2013).
26.   T. Clutton-Brock, Sexual Selection in Males and Females. *Science* **318**, 1882-1885 (2007).
27.   K. A. Rosvall, Intrasexual competition in females: evidence for sexual selection? *Behav*
Ecol **22**, 1131-1140 (2011).
28.   A. E. Wright, J. E. Mank, The scope and strength of sex-specific selection in genome
evolution. *J Evol Biol* **26**, 1841-1853 (2013).

29.   W. Goymann, J. C. Wingfield, Male-to-female testosterone ratios, dimorphism, and life
history—what does it really tell us? *Behavioral Ecology* **25**, 685-699 (2014).

30.   K. A. Rosvall, A. B. Bentz, E. M. George, How research on female vertebrates contributes
to an expanded challenge hypothesis. *Horm Behav* **123**, 104565 (2020).

31.   K. O. Smiley *et al.*, Beyond a biased binary: A perspective on the misconceptions,
challenges, and implications of studying females in avian behavioral endocrinology.
Front Physiol **13**, 970603 (2022).

32.   R. A. Duckworth, Adaptive dispersal strategies and the dynamics of a range expansion.
Am Nat **172 Suppl 1**, S4-17 (2008).

33.   D. Leffelaar, R. J. Robertson, Nest usurpation and female competition for breeding
opportunities by Tree Swallows. *The Wilson Bulletin* **97**, 221-224 (1985).

34.   J. Merilä , D. A. Wiggins, Interspecific Competition for Nest Holes Causes Adult Mortality
in the Collared Flycatcher. *The Condor* **97**, 445-450 (1995).

35.   A. N. Albers, J. A. Jones, L. Siefferman, Behavioral Differences among Eastern Bluebird
Populations Could Be a Consequence of Tree Swallow Presence: A Pilot Study. *Frontiers*
in Ecology and Evolution **5** (2017).

36.   C. A. Krieg, T. Getty, Fitness benefits to intrasexual aggression in female house wrens,
Troglodytes aedon. *Animal Behaviour* **160**, 79-90 (2020).

37.   K. A. Rosvall, Sexual selection on aggressiveness in females: evidence from an
experimental test with tree swallows. *Animal Behaviour* **75**, 1603-1610 (2008).

38.   D. R. Rubenstein, I. J. Lovette, Reproductive skew and selection on female
ornamentation in social species. *Nature* **462**, 786-789 (2009).

39.   E. R. Schuppe, G. D. Sanin, M. J. Fuxjager, The social context of a territorial dispute
differentially influences the way individuals in breeding pairs coordinate their aggressive
tactics. *Behavioral Ecology and Sociobiology* **70**, 673-682 (2016).

40.   S. E. Lipshutz, K. A. Rosvall, Nesting strategy shapes territorial aggression but not
testosterone: A comparative approach in female and male birds. *Horm Behav* **133**,
104995 (2021).

41.   Y. T. Fang, M. N. Tuanmu, C. M. Hung, Asynchronous evolution of interdependent nest
characters across the avian phylogeny. *Nat Commun* **9**, 1863 (2018).

42.   R. Zenil-Ferguson, J. P. McEntee, J. G. Burleigh, R. A. Duckworth, Linking Ecological
Specialization to Its Macroevolutionary Consequences: An Example with Passerine Nest
Type. *Syst Biol* **72**, 294-306 (2023).

43.   S. Kumar, G. Stecher, M. Suleski, S. B. Hedges, TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. *Mol Biol Evol* **34**, 1812-1819 (2017).

44.   J. M. Jawor *et al.*, Seasonal and individual variation in response to GnRH challenge in
male dark-eyed juncos (Junco hyemalis). *Gen Comp Endocrinol* **149**, 182-189 (2006).

45.   K. A. Rosvall, C. M. Bergeon Burns, S. P. Jayaratna, E. D. Ketterson, Divergence along the
gonadal steroidogenic pathway: Implications for hormone-mediated phenotypic
evolution. *Horm Behav* **84**, 1-8 (2016).

46.   S. A. Bukhari *et al.*, Temporal dynamics of neurogenomic plasticity in response to social
interactions in male threespined sticklebacks. *PLoS Genet* **13**, e1006840 (2017).

47.   J. L. Goodson, The vertebrate social behavior network: evolutionary themes and
variations. *Horm Behav* **48**, 11-22 (2005).

48. G. Stelzer *et al.*, The GeneCards Suite: From Gene Data Mining to Disease Genome
Sequence Analyses. *Current Protocols in Bioinformatics* **54**, 1.30.31-31.30.33 (2016).
49. H. H. Hung *et al.*, Selective involvement of UGGT variant: UGGT2 in protecting mouse
embryonic fibroblasts from saturated lipid-induced ER stress. *Proc Natl Acad Sci U S A*
119, e2214957119 (2022).
50. C. Yan *et al.*, Correction: IRE1 promotes neurodegeneration through autophagy-
dependent neuron death in the Drosophila model of Parkinson's disease. *Cell Death Dis*
11, 150 (2020).
51. J. Y. Zhu, M. Chen, W. J. Mu, H. Y. Luo, L. Guo, The functional role of Higd1a in
mitochondrial homeostasis and in multiple disease processes. *Genes Dis* **10**, 1833-1845
(2023).
52. E. M. Crombie, K. Cleverley, H. T. M. Timmers, E. M. C. Fisher, The roles of TAF1 in
neuroscience and beyond. *R Soc Open Sci* **11**, 240790 (2024).
53. N. M. Baran, P. T. McGrath, J. T. Streelman, Applying gene regulatory network logic to
the evolution of social behavior. *Proc Natl Acad Sci U S A* **114**, 5886-5893 (2017).
54. A. B. Bentz *et al.*, Experimental competition induces immediate and lasting effects on
the neurogenome in free-living female birds. *Proc Natl Acad Sci U S A* **118** (2021).
55. A. L. Filby, G. C. Paull, T. F. Hickmore, C. R. Tyler, Unravelling the neurophysiological
basis of aggression in a fish model. *BMC Genomics* **11**, 498 (2010).
56. J. F. Storz, Causes of molecular convergence and parallelism in protein evolution. *Nat*
Rev Genet **17**, 239-250 (2016).
57. M. J. West-Eberhard, Sexual selection, social competition, and speciation. *The Quarterly*
Review of Biology **58**, 155-183 (1983).
58. A. M. Harts, I. Booksmythe, M. D. Jennions, Mate guarding and frequent copulation in
birds: A meta-analysis of their relationship to paternity and male phenotype. *Evolution*
70, 2789-2808 (2016).
59. J. P. Drury, M. C. Cowen, G. F. Grether, Competition and hybridization drive interspecific
territoriality in birds. *Proc Natl Acad Sci U S A* **117**, 12923-12930 (2020).
60. J. F. Husak *et al.*, Life history and environment predict variation in testosterone across
vertebrates. *Evolution* **75**, 1003-1010 (2021).
61. W. Goymann, I. T. Moore, R. F. Oliveira, Challenge Hypothesis 2.0: A Fresh Look at an
Established Idea. *BioScience* **69**, 432-442 (2019).
62. J. C. Wingfield, M. Ramenofsky, R. E. Hegner, G. F. Ball, Whither the challenge
hypothesis? *Horm Behav* **123**, 104588 (2020).
63. J. E. Lischinsky, D. Lin, Neural mechanisms of aggression across species. *Nat Neurosci* **23**,
1317-1328 (2020).
64. S. E. Lipshutz, E. M. George, A. B. Bentz, K. A. Rosvall, Evaluating testosterone as a
phenotypic integrator: From tissues to individuals to species. *Mol Cell Endocrinol* **496**,
110531 (2019).
65. S. C. P. Renn *et al.*, Gene expression signatures of mating system evolution. *Genome* **61**,
287-297 (2018).
66. C. Alaux *et al.*, Honey bee aggression supports a link between gene regulation and
behavioral evolution. *Proc Natl Acad Sci U S A* **106**, 15400-15405 (2009).

67.   C. Mullon, A. E. Wright, M. Reuter, A. Pomiankowski, J. E. Mank, Evolution of dosage
compensation under sexual selection differs between X and Z chromosomes. *Nat*
Commun **6**, 7720 (2015).
68.   S. Grath, J. Parsch, Sex-Biased Gene Expression. *Annu Rev Genet* **50**, 29-44 (2016).
69.   A.-K. Fridolfsson, H. Ellegren, A simple and universal method for molecular sexing of
non-ratite birds. *Journal of Avian Biology* **30**, 116-121 (1999).
70.   K. K. Soma, B. A. Schlinger, J. C. Wingfield, C. J. Saldanha, Brain aromatase, 5 alpha-
reductase, and 5 beta-reductase change seasonally in wild male song sparrows:
relationship to aggressive and sexual behavior. *J Neurobiol* **56**, 209-221 (2003).
71.   M. G. Grabherr *et al.*, Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nature Biotechnology* **29**, 644-652 (2011).
72.   A. Rhie *et al.*, Towards complete and error-free genome assemblies of all vertebrate
species. *Nature* **592**, 737-746 (2021).
73.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).
74.   S. B. Plaisier, R. Taschereau, J. A. Wong, T. G. Graeber, Rank–rank hypergeometric
overlap: identification of statistically significant overlap between gene-expression
signatures. *Nucleic Acids Research* **38**, e169-e169 (2010).
75.   P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**, 559 (2008).
76.   H. Mi, A. Muruganujan, D. Ebert, X. Huang, P. D. Thomas, PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools.
Nucleic Acids Research **47**, D419-D426 (2018).
77.   P. Shannon *et al.*, Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res* **13**, 2498-2504 (2003).
78.   J. D. Hadfield, MCMC Methods for Multi-Response Generalized Linear Mixed Models:
The MCMCglmm R Package. *Journal of Statistical Software* **33**, 1-22 (2010).
79.   W. Jetz, G. H. Thomas, J. B. Joy, K. Hartmann, A. O. Mooers, The global diversity of birds
in space and time. *Nature* **491**, 444-448 (2012).
80.   E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**, 526-528 (2019).
81.   L. J. Revell, phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* **3**, 217-223 (2011).
82.   S. Seabold, J. Perktold (2010) Statsmodels: Econometric and Statistical Modeling with
Python. in *SciPy Conferences*.
828