1

2
3
4 **Unique growth and morphology properties of Clade 5 *Clostridioides difficile* strains**
5 **revealed by single-cell time-lapse microscopy**
6

John W. Ribis[1,2], César Nieto[3], Nicholas V. DiBenedetto[1,2], Anchal Mehra,[4] Qiwen Dong[1], Irene

Nagawa,[1] Imane El Meouche,[5] Bree B. Aldridge,[1,6] Mary J. Dunlop,[7] Rita Tamayo,[4] Abhyudai

Singh,[3] Aimee Shen[1*]

Affiliations

[1]Department of Molecular Biology and Microbiology, Tufts University School of Medicine,

Boston, Massachusetts, USA, [2] Graduate School of Biomedical Sciences, Tufts University

School of Medicine, Boston, Massachusetts, USA. [3]Department of Electrical and Computer

Engineering, University of Delaware, Newark, DE 19716, USA. [4]Department of Microbiology

and Immunology, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill,

NC, USA, [5]INSERM, Université Paris Cité, Université Sorbonne Paris Nord, Inserm, IAME, F-

75018, Paris, France, [6]Department of Biomedical Engineering, Tufts University School of

Engineering, Medford, Massachusetts, USA, [7]Biomedical Engineering, Boston University,

Boston, MA 02215.

[*]Address correspondence to
Aimee Shen, aimee.shen@tufts.edu
Phone number: (617)636-3792

Abstract

Clostridioides difficile is a major One Health threat as a gastrointestinal pathogen of both

humans and agricultural animals. The *C. difficile* species consists of five main clades, with Clade

5 currently undergoing speciation from Clades 1-4. Notably, Clade 5 strains are highly prevalent

in agricultural animals and can cause zoonotic infections, suggesting that Clade 5 strains have

evolved phenotypes that distinguish them from Clade 1-4 strains. Here, we compare the growth

properties of Clade 5 strains to Clade 1, 2, and 4 strains using anaerobic time-lapse microscopy

coupled with automated image analysis. These analyses reveal that Clade 5 strains grow faster

than Clade 1, 2, and 4 strains and are more likely to form long chains of cells. Comparative

genomic analyses revealed that the orientation of the invertible *cmr* switch sequence likely drives

the cell chaining phenotype of Clade 5 strains during growth in rich media. Interestingly, during

murine infection, Clade 5 strains with a high frequency of the *cmr* switch in the ON orientation

shifted to a largely *cmr* OFF state, suggesting that the *cmr* OFF state may be positively selected

during infection. Taken together, our data suggest that Clade 5 strains have distinct growth

properties that allow them to inhabit diverse ecological niches.

Importance

The Clade 5 strains of the *Clostridioides difficile* species are so phylogenetically divergent that

they almost meet the threshold of being a distinct species. Although these strains are

ubiquitously isolated from agricultural and environmental settings and an important source of

zoonotic and community-acquired infections, it is unclear whether they have distinct phenotypic

properties that allow them to efficiently colonize diverse hosts or persist in the environment. By

combining a novel anaerobic time-lapse microscopy method with automated image analysis, we

discovered that Clade 5 strains grow faster than strains from other *C. difficile* clades and that

they frequently form long chains. These chaining properties appear to be driven by the

environmentally responsive expression of a non-canonical signal transduction system, which we

also found is likely selected against during growth in mice. Collectively, our analyses reveal that

Clade 5 strains have distinct growth properties that may promote their persistence in the

environment.

Introduction

       Clostridioides difficile is a leading cause of nosocomial infection in the United States, with approximately 500,000 new infections and 14,000 deaths attributed to this organism annually (1, 2). As an obligate anaerobe, *C. difficile* relies on its ability to form hardy, metabolically dormant spores to survive outside the gut environment and transmit disease to new hosts (3). When *C. difficile* spores are ingested by susceptible hosts (4, 5), they germinate and outgrow into vegetative cells that subsequently colonize the colon. The vegetative cells secrete toxins that damage gut epithelial tissue (6), which triggers a significant inflammatory response that can cause disease pathologies ranging from mild diarrhea to pseudomembranous colitis and even death (2, 4). *C. difficile* also causes recurrent infections (~20% of infections), which can lead to more severe disease symptoms (2, 7, 8).

       C. difficile's success as a pathogen may be related to its tremendous genetic diversity (9-11). Genomic analyses indicate that *C. difficile*'s core genome represents only ~10-20% of its pan-genome (9, 10). The plasticity of its "open" pan-genome likely helps *C. difficile* colonize the gastrointestinal tract of diverse animals, from mammals to invertebrates, and persist in environmental reservoirs like sewage and compost (12). Multi-locus sequence typing (MLST) analyses have revealed that the *C. difficile* species comprises five distinct phylogenetic clades that can be further subdivided into different ribotypes (RTs) or sequence types (STs) (10, 13).

       Between these five clades, there are notable differences in geographic and host distributions. Clade 1 is the largest and most heterogeneous clade, containing over 200 STs that have the broadest geographic distribution (14, 15). It includes both toxin-producing and non-toxigenic strains, including the well-characterized, genetically tractable toxigenic strain 630 (9). Clade 2 harbors epidemic-associated strains found within the ribotype 027 (RT027) lineage. Strains from this ribotype have been associated with outbreaks in hospitals, particularly in North America, due to their frequent resistance to fluoroquinolones (16-18). These epidemic strains can also cause severe disease symptoms in part due to their production of three toxins: TcdA, TcdB, and CDT (binary toxin) (6). Although RT027 strains have frequently been associated with "hypervirulence," there is considerable phenotypic diversity within this lineage with respect to virulence, toxin production levels, and sporulation (19-21).

Clade 3 strains are relatively uncommon and harbor a unique cell surface due to their

lack of CwpV production (22), but phenotypic analyses of biofilm formation and motility

suggest that they share similarities with Clade 2 strains like R20291 (23). Clade 4 contains

RT017 (ST37) strains that only encode a single toxin, TcdB (i.e., TcdA⁻CDT⁻), and are often

clindamycin- and fluoroquinolone-resistant. RT017 strains have been associated with outbreaks

in Europe and North America and are the most common strains found in Asia (24). Clade 5 is

the most genetically distant from the other 4 main *C. difficile* clades (94) and is thought to have

emerged before Clades 1-4 (25). While the average nucleotide identity (ANI) for Clades 1-4

ranges between 97.1 - 99.8%, Clade 5 strains exhibit ANI values around 96%, which is close to

the ANI value demarcation used by NCBI to define organisms of the same species (10, 26).

Thus, Clade 5 strains appear to be actively diverging from Clades 1-4 (10).

Clade 5 strains are an increasing problem in healthcare and agricultural settings because

they can cause severe disease in humans and are commonly found in livestock, particularly pigs

(12, 27). While other *C. difficile* strains have been known to infect both humans and animals,

only Clade 5 strains have been associated with zoonotic transmission from both animal-to-

human and human-to-animal (27, 28). The mechanisms underlying this bidirectional zoonotic

transmission are poorly understood, but the increased carriage of antimicrobial resistance genes

by Clade 5 strains may contribute to their ability to persist in agricultural and community

settings (27, 29). Thus, Clade 5 strains are of particular relevance from a One Health

perspective (12, 30), especially since they frequently cause community-acquired infections (29)

and are often detected in retail foods (31). These observations highlight the importance of

understanding the unique properties of this group of strains. Indeed, a recent genomic analysis

suggests that RT078/ST11 strains within Clade 5 frequently carry zinc acquisition and

homeostasis genes (11).

Despite numerous genomic analyses revealing the remarkable genetic diversity of *C.*

difficile strains, relatively few studies have investigated the phenotypic diversity between strains

from different clades. Clade-specific differences in colony morphology between Clade 5 strains

relative to Clade 1-4 strains have been described in a limited set of analyses (25, 32), suggesting

that differences in growth and/or cellular morphology may exist within clades. While

differences in bulk growth rates between *C. difficile* strains have been reported (33), most

phenotypic analyses have been conducted on a limited subset of strains within a given clade.

Furthermore, systematic comparisons of the growth properties of the different clades or

comparisons of cell morphology have not been performed to date.

Here, we compare the growth properties of multiple strains derived from the four

predominant main phylogenetic clades of *C. difficile* (Clade 1, 2, 4, and 5) using anaerobic

time-lapse microscopy. These analyses unexpectedly revealed striking differences in the growth

and cell morphology of the Clade 5 lineage relative to strains from Clades 1, 2, and 4.

Specifically, we found that Clade 5 strains grow faster and frequently form long chains, in

contrast with Clade 1, 2, and 4 strains. Genomic comparisons of the Clade 5 strains tested

strongly suggest that the presence of the *cmr* switch in the ON orientation drives their chaining

phenotype. While the orientation of this switch has previously been associated with cell

chaining (32), we found that Clade 5 strains with a strong *cmr* ON bias mostly reverted to a *cmr*

OFF phenotype during murine infection. Taken together, our data reveal that Clade 5 strains

have unique growth properties relative to Clade 1, 2, and 4 strains that may contribute to the

improved ability of Clade 5 strains to colonize and persist in a wider range of animal hosts.

Results

Development of a simple method for time-lapse imaging under anaerobic conditions

Time-lapse imaging of single cells has been widely used in bacteria due to the growing

appreciation that clonal bacterial populations can exhibit considerable phenotypic heterogeneity,

which can impact antibiotic resistance and virulence (32, 34-37). However, live single-cell

analyses in *C. difficile* have been complicated by its inability to grow in the presence of

atmospheric oxygen (38). While there have been limited reports of live cell time-lapse

microscopy of *C. difficile*, these analyses have required custom growth chambers to maintain

anaerobic conditions (39), which may limit the accessibility of these experimental systems to

most investigators.

To overcome these limitations, we established a simple system that relies solely upon

commercially available reagents and materials to grow *C. difficile* cells under anaerobic

conditions. This system uses gas-tight, adhesive Gene Frames™, which have been used

extensively in imaging applications for bacteria (40). Notably, the gas-impermeability of these

commercial seals allows anaerobic conditions to be maintained when agarose pads made with

growth media are prepared in the anaerobic chamber (**Figure 1**). Gene Frames also generate

thick agarose pads, which are critical for *C. difficile* to grow in a sealed system under ambient

conditions. Upon preparing agarose pads in the anaerobic chamber, *C. difficile* cultures are

inoculated onto the pads, and the pads are sealed a coverslip. At this point, the growth chamber is

removed from the chamber and imaged on a heated stage under ambient conditions for up to 6

hours. Although the growth chamber maintains its anaerobic status during this time, movies were

generally limited to 6 hours because *C. difficile* stopped growing exclusively as a monolayer.

159
160

Figure 1. Schematic of the anaerobic single-cell imaging set-up. Exponentially growing *C. difficile*
cells in TY medium supplemented with cysteine (TYC) are spotted onto 1.5% agarose pads formed
within gas-tight adhesive Gene Frames inside the anaerobic chamber. Up to 6 strains can be spotted
onto a pad. The pad is sealed with a coverslip, and the imaging chamber is removed from the anaerobic

chamber and transferred to a heated (37˚C) microscope stage. Time-lapse microscopy is used to
visualize the growth of individual bacterial cells for 2-6 hrs. The output data is segmented and tracked
with the DeLTA Python package (191). An example filmstrip of strain 630 grown on TYC medium
over time is shown (Bottom).
169

Time-lapse microscopy reveals clade-specific differences in elongation rate and cell length

With the establishment of our time-lapse imaging setup, we compared the single-cell

growth properties of representative *C. difficile* strains from Clade 1 (630, ribotype (RT) 012),

Clade 2 (R20291, RT027), Clade 4 (M68, RT017), and Clade 5 (M120, RT078) (**Figure 2,**

Table 1). We did not have access to Clade 3 strains, so these were not included in this study.

The four "representative" strains were all isolated from patients with *C. difficile*-associated

disease and are frequently used as reference genomes for their clades and ribotype groups.

Notably, RT027 (ST1), RT017 (ST45), and RT078 (ST11) strains are from ribotypes/ multi-

locus sequencing types that are frequently isolated from patients with *C. difficile* infection

(CDI) (12, 16, 19, 24).

To quantify the growth properties of single cells visualized by time-lapse microscopy,

we used the Deep Learning for Time-lapse Analysis (DeLTA) software to rapidly and

accurately segment and track bacteria growing in two dimensions on agarose pads (41, 42).

This software uses deep convolutional neural networks to analyze time-lapse microscopy

images, allowing us to obtain information on the growth properties of individual cells growing

in microcolonies on agarose pads. The segmentation and tracking of *C. difficile* cells were

highly accurate (**Figure 1**), and minimal user input or post-image processing was needed to

obtain growth property measurements.

Robust growth was observed for all strains using our system. We quantified growth

using the elongation rate defined as doublings/hr. Notably, the Clade 5 strain M120 elongated

the fastest (2.1 doublings/hr), followed by Clade 2 strain R20291 (1.8 doublings/hr), and then

Clade 1 strain 630 and Clade 4 strain M68 (1.6-1.7 doublings/hr) (**Figure 2A**). The elongation

rate was defined as doublings/hr to indicate the number of times that a cell's size doubles in one

hour. Notably, the elongation rate (doublings/hr) should not be confused with doubling time, or

generation time, which represent the length of *time* that it takes before a bacterium divides.

Instead, the elongation rate reflects how fast the cell is increasing in length over time.

198
199
Figure 2. Clade 5 strain M120 elongates more quickly and exhibits cell chaining. (A) Violin plot of
the elongation rates measured during time-lapse microscopy analyses of strains 630 (Clade 1), R20291
(Clade 2), M68 (Clade 4), M120 (Clade 5) grown on TY supplemented with cysteine (TYC) agar. Data
are from three biological replicates, with the mean of each replicate shown as a point on the violin. (B)
Violin plot of the cell or chain length measured during time-lapse microscopy for strains 630 (Clade 1),
R20291 (Clade 2), M68 (Clade 4), M120 (Clade 5). Each replicate mean is shown as a point on the
violin. Statistical significance for A and B was determined by comparing the mean of the three replicates
of Clade 1, 2, and 4 strains relative to the Clade 5 M120 strain using a Kruskal-Wallis test * p < 0.05.
(C) Phase-contrast image from time-lapse microscopy movies. Scale bar is 10 μm. D. Optical density-
based analyses of bulk population growth of the indicated strains in TYC or BHIS medium. The number
in brackets indicates the clade to which a given strain belongs.
211

Importantly, the differences in single-cell elongation rates measured for the four strains

using DeLTA were also observed in bulk population analyses of their growth using optical

density in TYC and BHIS media (**Figure 2D**). These analyses confirmed that the Clade 5 strain

M120 grows faster (using optical density as a proxy for bacterial growth) than the Clade 1, 2,

and 4 strains analyzed. In contrast, negligible differences in bulk growth rates were observed

between Clade 1, 2, and 4 strains (**Figure 2D**), consistent with the relatively subtle difference

observed in the single-cell elongation rates measured for the Clade 2 R20291 strain relative to

the Clades 1 and 4 strains (**Figure 2A**).

The Clade 5 strain M120 exhibited another distinct growth property from the strains

from Clades 1, 2, and 4. While strains from Clades 1, 2, and 4 produced cells of similar length

prior to cell division, with an average length of ~13 µm, cells of Clade 5 strain M120 were

significantly longer, appearing as chains with an average length of 22 µm; M120 chains of ~50

µm were also readily observed (**Figure 2B**), and chains appeared to bend readily (**Figure 2C**).

Notably, to fully visualize the chains made by M120, it was necessary to stitch together several

fields of view. This modification revealed that some M120 chains approached several hundred

microns and even up to ~1 mm in length (**Figure 3**). By incorporating the lipophilic fluorescent

dye FM4-64 into the agarose pads to visualize cell membranes (43), we determined that strain

M120 forms chains rather than filaments based on the presence of division septa throughout the

length of the cell (**Figure 3, inset**). Notably, the spacing between division septa was relatively

similar to the other *C. difficile* strains analyzed, suggesting that Clade 5 strain M120 undergoes

cell separation less efficiently than strains from the other clades tested.

235
236 **Figure 3. *C. difficile* Clade 5 can form large heterogenous chains.** Large mosaic phase-contrast
237 image of the Clade 5 strain M120. Zoomed image shows chains revealed by staining with the
238 membrane dye FM4-64. Image was stitched from 8 individual fields of view at 63x magnification.
239 Scale bar is 100 μm.
240

Clade 5 clinical isolates typically form long chains and grow more quickly than strains

from other clades

Since prior work indicated that Clade 5 strains produce colony morphologies distinct from

Clade 1, 2, and 4 strains (25), we sought to determine whether the striking cell chaining

phenotype and faster growth rate observed in Clade 5 strain M120 were properties shared by

other Clade 5 strains. Thus, we compared the single-cell growth properties of five additional

Clade 5 clinical isolates obtained from three separate hospitals on TYC agarose (**Table 1**) using

time-lapse microscopy analyses (**Figure 2**). Clade 1 strain 630 was included as a control since it

does not form chains under these conditions.

These analyses revealed that all but one of the Clade 5 strains tested formed long chains, with

TAL29600 forming the longest chains (29 μm on average, **Figure 4A-C**). In contrast, strain

TAL29996 formed shorter chains that were comparable in length to those observed for strains

from Clades 1, 2, and 4 (12-13 μm, **Figures 2B, 4C**). Notably, DeLTA segmented most Clade 5

strains as single cells because it was unable to identify division septa (based on FM4-64

staining) within a given chain because cell separation had not yet been initiated. To overcome

this limitation and accurately quantify cell length within long chains, namely the distance

between division septa, we modified our image processing pipeline to use a thresholding

method to detect division septa. After generating masks in DeLTA to segment the chains, we

modified the mask so that only the interior of the contour was analyzed. We then applied an

adaptive thresholding method to identify division septa based on their elevated fluorescence

relative to the long axis of the cell; the thresholds were defined using a Gaussian-weighted

method (**Figure 4A**, red dots, **Figure 4B**). These analyses revealed that the average length of

cells made by Clade 5 strains that form long chains is only slightly longer than the average

length of cells made by strains that do not form chains. Specifically, the Clade 5 strains that

produced the longest chains (TAL29600 and TAL30550) had average cell lengths of 14 and 12

μm, respectively, that were 30-50% longer than strains that do not form chains like strains 630

(Clade 1, 8.5 μm) and TAL29996 (Clade 5, 9.4 μm) (**Table 2**). Notably, even though the Clade

5 strain TAL29996 did not form chains, it still elongated at rates similar to those of other Clade

5 strains (2.1 doublings/hr vs. 1.7 doublings/hr for Clade 1 strain 630**, Figure 4D**).

272
273

Figure 4. Faster growth is a common feature of Clade 5 strains, but the chaining phenotype is not fully
penetrant in Clade 5 strains. (A) Phase-contrast microscopy images from time-lapse microscopy studies of
Clade 5 strains; Clade 1 strain 630 is included for comparison. Fluorescence microscopy was used to
visualize the FM4-64 stain incorporated into the agarose pads. Masks generated using DeLTA are shown,
with the bottom panel showing division septa identified with our adaptive thresholding approach (red dots).
(B) Example image showing the parameters identified using DeLTA combined with our adaptive threshold
method for detecting division septa within a chain of cells. (C) Cell or chain length measured using
automated DeLTA analyses (pink violin plot) or DeLTA combined with adaptive threshold analyses (purple
violin plot). The former method is more likely to measure chain length, while the latter method accurately

measures cell length. (D) Violin plot of the elongation rates measured based on three biological replicates.
Each replicate mean is shown as a point on the violin; statistical significance was determined by comparing
the mean of the three replicates using a Kruskal-Wallis test, * p < 0.05.
286

Indeed, all the Clade 5 strains tested grew faster in bulk analyses in broth culture.

Specifically, in optical density-based analyses, the Clade 5 strains grew faster than Clade 1, 2,

and 4 strains irrespective of their ability to form chains or the growth medium used (**Figure S1**).

To assess whether these conclusions could be generalized to additional Clade 5 strains vs. Clade

1, 2, and 4 strains, we analyzed the growth of additional strains from all four clades in broth

culture. These analyses confirmed that Clade 1, 2, and 4 strains grow at similar rates, which are

slower than those measured for the nine Clade 5 strains analyzed in both TYC and BHIS media

(**Figure S1**). Taken together, these analyses strongly suggest that Clade 5 strains grow faster

than Clade 1, 2, and 4 strains and are more likely to form chains presumably because their cell

separation mechanisms are less efficient.

Cell chaining in Clade 5 strains is not dependent on growth on a solid medium.

C. difficile has previously been shown to promote cell elongation and chain formation

upon induction of the *cmrRST* locus (32), which encodes a non-canonical signal transduction

system. Expression of this locus is also responsive to c-di-GMP levels (44), which increases in

cells grown on solid surfaces such as in a biofilm or on an agar plate (44). To determine

whether the chaining phenotype observed in most Clade 5 strains is induced by growth on a

surface, we assessed whether cell chaining would be observed in cells grown in broth culture.

The Clade 5 strains, along with strains from other clades, were grown to log-phase in rich

media, and their ability to form chains was assessed by visualizing division septa with the

fluorescent D-amino acid label, HADA. Chains of cells were still observed in the Clade 5

strains during broth culture growth, although the chains formed in broth culture were not as

long as those observed during growth on the agarose pads (**Figure 5, S2**). Chains were also

observed in four additional Clade 5 strains analyzed during exponential-phase growth in broth

culture. Interestingly, two of the Clade 5 strains, V48 and WU66, failed to efficiently

incorporate the HADA label for unknown reasons, and the cells appeared lighter in phase-

contrast images despite undergoing the fixation process shortly after HADA addition (**Figure

S2). Regardless, these results indicate that many Clade 5 strains undergo cell separation less

efficiently than strains from other clades irrespective of whether they are grown on a solid

surface or in broth culture.

319
320
Figure 5. Clade 5 strains form chains during logarithmic growth in broth culture. Representative
micrographs showing phase-contrast (top) and peptidoglycan labeling with the fluorescent D-amino acid,
HADA, (bottom) following growth in rich broth (BHIS) to logarithmic phase. All strains shown with the
exception of 630 (Clade 1) are Clade 5 strains. Scale bar, 10 μm. Data are representative of multiple
independent experiments. **ADJ** indicates that the brightness of the image was enhanced to detect HADA
labeling in V48.

327
Cell length does not correlate with the propensity to sporulate.

We next wondered whether the propensity to form chains impacts the ability of Clade 5

strains to sporulate because analyses in *Bacillus subtilis* suggest that smaller cells, such as those

formed during stationary phase growth (45), may concentrate proteins involved in the

phosphorelay that induces sporulation (46, 47). Specifically, in *B. subtilis*, the decrease in cell

length helps the kinase KinA reach the threshold concentration needed to trigger sporulation

initiation. Although *C. difficile* lacks homologs of KinA and other components of the

phosphorelay system (48), it is possible that the longer cells generated by Clade 5 strains would

be less likely to induce sporulation due to dilution of a currently unknown sporulation regulator.

To test this hypothesis, we analyzed the propensity of Clade 5 strains to form spores when

plated on 70:30 sporulation medium using phase-contrast microscopy and heat resistance

assays. These analyses revealed that Clade 5 strains exhibit striking differences in sporulation

frequency, with some strains exhibiting close to 100% sporulation levels, and others exhibiting

levels closer to 20% (**Figure 6A, S3**). Strikingly, strain TAL29600 exhibited extremely low

levels of sporulation (0.003%). Despite this strain producing the longest chains, no correlation

between sporulation and cell length or chaining among the strains was observed (**Figure 6B,**

S3). For example, M120 forms long chains yet readily forms spores (~100% sporulation

frequency). The low sporulation levels of TAL29600 were rather surprising since this strain was

isolated from a clinical setting, so it likely forms spores more efficiently in conditions

encountered during infection. Interestingly, Clade 5 strains did not form long chains when

grown on 70:30 sporulation medium, and the spores produced by these isolates exhibited

similar lengths and proportions relative to the Clade 1 strain 630 (**Figure S4**). Thus, our data

suggest that Clade 5 strains alter their cell length and propensity to form chains depending on

the growth conditions encountered.

353
354
Figure 6. Sporulation levels in Clade 5 isolates grown on 70:30 medium. Phase-contrast microscopy
of the indicated strains ~24 hrs after sporulation induction. All strains shown with the exception of 630
(Clade 1) are Clade 5 strains. Heat resistance efficiencies, indicated below the respective images, are
based on 20-24 hr sporulating cultures and represent the mean and standard deviation for a given strain
relative to strain 630 based on a minimum of three biological replicates. Statistical significance relative
to strain 630 was determined using a one-way ANOVA and Tukey's test. The scale bar represents 5 μm.
361

Genomic analyses of Clade 5 strains to identify putative loci controlling the chaining

phenotype

Since our phenotypic analyses revealed that TAL29996 differs from the 8 other Clade 5

strains analyzed, we sought to gain insight into the molecular basis for this difference by

comparing their genomes. To this end, we sequenced 5 of the Clade 5 strains, including

TAL29996, and determined the average nucleotide identity (ANI) for orthologous genes. These

analyses revealed that the ANI between the strains ranging from 99.83-99.99% (**Table S1**). Since

the pan-genome between the 5 strains is 12%, they are quite closely related. A comparison of

their pan-genomes revealed that, relative to the other four strains, TAL29996 is missing one

duplication of *blaR1*, which encodes an integral membrane protein that senses beta-lactams, and

a gene region predicted to be involved in nicotinate metabolism. We also compared the genomes

using breseq (49) to identify SNPs that might distinguish TAL29996 from the other strains. We

used strain M120 as the reference genome, since this strain has been the Clade 5 strain

traditionally characterized (33, 50). Around 10 SNPs were unique to TAL29996, but none were

clearly implicated in regulating cell separation or peptidoglycan synthesis (**Table S2**).

Since none of the SNPs or absent genes led to hypotheses for why TAL29996 undergoes

cell separation more efficiently than the other Clade 5 strains, we took a candidate approach and

analyzed the orientation of the *cmr* switch, also known as the Cdi6 DNA invertible element (32,

51). The orientation of the *cmr* switch affects the expression of the adjacent *cmrRST* operon,

resulting in phase-variable production of the non-canonical CmrRST signal transduction system.

This system promotes cell chaining when produced. In addition, the orientation of the *cmr* switch

determines whether *C. difficile* produces rough or smooth colony morphologies (32, 44, 51).

Specifically, cells from rough colonies are highly biased to the ON orientation of the *cmr* switch

and exhibit a chaining phenotype, whereas cells from smooth colonies are highly biased to the

OFF orientation and do not form chains (32). Consistent with the breseq analyses, manual

inspection of the *cmrRST* upstream region revealed that this region is identical between all the

Clade 5 strains analyzed (100% identity), including strain TAL29996.

However, when we specifically assessed the orientation of the *cmr* switch during growth

in broth culture using qPCR, we found that Clade 5 strains form chains with a *cmr*-ON frequency

of between 70-96% on average (**Figure 7A**), whereas the *cmr*-ON frequency of the non-chaining

Clade 5 strain TAL29996 was greatly reduced (~20%, **Figure 7**). Furthermore, when we

extended these analyses to the four additional Clade 5 strains characterized that also produce

chains, we found that they exhibit a *cmr*-ON frequency of between 80-93%. Collectively, these

results suggest that a *cmr*-ON switch orientation correlates with the chaining phenotype observed

in the majority of Clade 5 strains even during growth in broth culture.

Surface motility in Clade 5 strains relative to strains of other clades

Since the *cmr*-ON state has also been correlated with increased surface motility, we analyzed the

surface motility of our Clade 5 strains. After a 5-day incubation, *cmr*-ON strains spread further

on agar plates than the TAL29996 *cmr*-OFF strain (**Figure 7B**). Consistent with prior reports

(32), Clade 5 strains whose *cmr* switch is predominantly in the ON orientation exhibited greater

and more uniform surface motility (**Figure 7**). In contrast, the surface motility of TAL29996 was

considerably reduced and more asymmetric, with fractal-like extensions extending from a couple

of sites. The surface motility of TAL29996 resembles the phenotype previously reported for

Clade 2 strain R20291, whose *cmr* switch is also predominantly in the OFF position in liquid

cultures (32).

However, growth of R20291 on agar medium selects for bacteria with the *cmr* switch in

the ON orientation, which leads to asymmetric surface motility on plates (32). These

observations suggest that, even though TAL29996 is biased to the *cmr*-OFF orientation during

broth culture growth, a subset of TAL29996 cells switch to the *cmr*-ON orientation during

growth on BHIS agar, leading to the asymmetric spreading phenotype.

Figure 7. Orientation of the *cmr* switch correlates with chaining phenotype and surface motility in Clade 5 strains. (A) Orientation-specific qPCR for detecting the orientation of the *cmr* switch in the indicated strains. The mean and standard deviation based on one to three biological replicates are shown. (B) Representative images of surface motility 5 days after exponentially growing liquid cultures were spotted onto BHIS agar plates. All strains shown with the exception of 630 (Clade 1) are Clade 5 strains.

We next analyzed the propensity of Clade 1, 2, and 4 strains to exhibit surface motility.

These analyses revealed that asymmetric motility was more frequently observed in Clade 2

strains. Clade 1 strains exhibited a range of surface motility, from high surface motility with

strain 630 to lower surface motility with strain WU38 (**Figure S5**). While analyzing a larger

number of Clade 1, 2, and 4 strains would reveal whether there are clade-specific differences in

surface motility, the data suggest that *cmr* switching in Clade 2 strains is more variable between

strains during growth on agar medium, with strain Wup14 exhibiting the least surface motility of

the Clade 2 strains tested. Collectively, the data imply that the ON orientation of the *cmr* switch

during broth culture growth contributes to the chaining phenotype observed in most Clade 5

strains.

Colonization and virulence properties of Clade 5 strains

The *cmrRST* system impacts the virulence of the Clade 2 strain, R20291, in a hamster

model of infection, with loss of *cmrR* reducing R20291's ability to cause disease and the *cmr-*

OFF orientation correlating with less severe disease in hamsters (32). Since chaining in *Bacillus*

anthracis strains promotes virulence (52), while chaining in *Enterococcus faecalis* promotes

colonization (53), we compared the ability of Clade 5 strains to colonize mice and cause disease.

Specifically, we analyzed the ability of several Clade 5 strains and the Clade 1 strain 630 to

colonize and cause disease in antibiotic-treated mice. We also analyzed the orientation of the *cmr*

switch over the course of the 14-day infection for strains TAL29600 and TAL29996, which

exhibited the highest and lowest *cmr*-ON orientations, respectively, during growth in rich media

(**Figure 7A**).

All Clade 5 strains tested colonized to relatively similar levels for throughout the 14 days

of the infection. Strain 630 also colonized mice to similar levels in the first two days of infection

and then maintained colonization, albeit at 1-2 logs lower than the Clade 5 strains (**Figure 8A**).

Interestingly, only the Clade 5 strain TAL29600 caused significant weight loss relative to the

other strains on Days 2 through 4, while the Clade 1 strain 630 caused some weight loss on Day

3 (**Figure 8B**). This latter phenotype is consistent with prior reports of strain 630 causing only

mild disease symptoms in the cefoperazone model of murine infection (54, 55). Analyses of the

cmr orientation revealed that the *cmr*-OFF orientation appeared to be selected for over the course

of the infection. While the TAL29600 spore inoculum started off with ~30% *cmr*-ON frequency, the frequency of TAL29600 cells detected in the *cmr*-ON orientation decreased rapidly to < 5% *cmr*-ON by 24 hrs post-inoculation (**Figure 8C**). As the infection progressed, two of the 8 mice tested exhibited an increase in TAL29600 cells with the *cmr*-ON orientation (6-30% *cmr*-ON) (**Figure S6**). Conversely, the TAL29996 strain retained the ~1% *cmr*-ON frequency of the inoculum for the greater part of the 14-day infection (**Figure 8C**). Taken together, these analyses reveal that Clade 5 RT078 strains efficiently colonize mice but vary in their ability to cause disease. Furthermore, the ability of the Clade 5 strains to colonize or cause disease did not correlate with their ability to chain in rich media.

Figure 8. Infection and colonization dynamics of Clade 5 strains in mice. (A) Fecal colony-forming units measured by selective plating and (B) Percentage of weight loss to baseline of infected mice on Days 1-4, 7, and 14. The mean and standard deviation are shown based on the results of two experiments consisting of four mice each (n = 8). (C) Percentage of *cmr* ON switch orientation measured in fecal pellets on the indicated days by qPCR. For the strain TAL29996 inoculum, 1% of the spores had the *cmr*-

470   ON switch orientation, while the TAL29600 inoculum consisted of 32% *cmr*-ON spores. The mean and
471   standard deviation based on analyses of eight mice are shown, although fecal pellets could not be
472   collected from some of the mice on Day 2.
473

Discussion

While Clade 5 strains are genetically distinct from Clade 1-4 strains (10, 25) and more

prevalent in animals (12, 28, 29), the phenotypic impact of their genetic divergence has been

poorly characterized. By phenotypically characterizing *C. difficile* strains from multiple clades

using time-lapse microscopy, we discovered that Clade 5 strains have distinct growth properties

from the Clade 1, 2, and 4 strains tested. Specifically, Clade 5 strains elongate more quickly

(**Figures 2, 4, S1**) and form long chains more readily than strains from Clades 1, 2, and 4,

irrespective of media type or growth on a surface (**Figures 2, 4, S2**). In contrast, long chains

were not observed in any of the Clade 1, 2, and 4 strains, regardless of whether the cells were

grown on agarose pads or in broth culture. Thus, Clade 5 strains undergo cell separation far less

efficiently than strains from other clades during growth in rich media. While our analyses of

Clade 5 strains were largely limited to the RT078 ribotype, we note that a prior report described

an RT126 strain isolated from a patient experiencing multiple recurrences that also formed long

chains (56). Since the RT126 ribotype is closely related to the RT078 ribotype (12), the

chaining phenotype will likely be observed in other Clade 5 ribotypes.

Our data strongly suggest that the chaining phenotype of Clade 5 strains relates to their

propensity to have the *cmr* switch (32, 44) in the ON orientation during growth in rich media

(**Figure 7**) because the non-chaining TAL29996 strain exhibits a *cmr*-ON orientation frequency

of 20%, whereas the 8 other Clade 5 strains tested exhibit *cmr*-ON orientation frequencies of

~70-95% (**Figure 7**). This conclusion is further supported by our finding that Clade 5 strains

form longer chains more frequently during growth on agarose pads compared to broth culture

(**Figures 4, 5**), which is similar to the finding that the *cmrRST* operon is induced by elevated c-

di-GMP levels encountered during growth on agar plates in the Clade 2 strain R20291 (44). In

addition, Clade 5 strains also exhibited high levels of surface motility, which is a *cmrRST*-

induced phenotype in strain R20291 (32). Interestingly, *cmrRST* expression is negatively

correlated with flagellar motility in strain R20291, and growth conditions favoring flagellar

motility select against *cmr*-ON bacteria in this strain (32). Since Clade 5 strains lack flagellar

motility (12), it is tempting to hypothesize that they are "primed" to exhibit surface motility.

Analyzing the regulation of c-di-GMP in different growth conditions in Clade 5 strains,

particularly for TAL29996 relative to the other strains, will likely provide insight into the

mechanisms that drive *cmrRST* expression, chaining, and surface motility in Clade 5 strains.

Such analyses would be greatly facilitated by methods for genetically manipulating Clade 5

strains.

While the propensity of Clade 5 strains to form chains and carry the *cmr* switch in the

ON orientation might be expected to promote colonization or disease in mice based on studies

of other Gram-positive pathogens (52, 53), we observed that the frequency of *cmr*-ON cells

decreases during the first 4 days of infection for strain TAL 29600 (**Figure 8**). This suggests

that there may be a selection against *cmr*-ON cells during the initial stages of murine

colonization. Indeed, a similar selection for the *cmr*-OFF orientation was observed during

infection of hamsters with R20291 (32). However, since the frequency of *cmr*-ON TAL29600

cells increased in some mice at later stages of colonization (**Figure 8**), our data imply that the

invertibility of the *cmrRST* switch region likely promotes *C. difficile*'s ability to adapt to

different growth conditions. While it is possible that the operon is expressed during infection

even if the orientation of the *cmr* switch is in the "OFF" position due to the induction of the

cmrRST operon by c-di-GMP (44), assessing whether Clade 5 strains form long chains during

murine infection, for example, using fluorescence *in situ* hybridization (57, 58) or using

transcriptional reporters to visualize the expression of the *cmrRST* operon at the single-cell level

during infection would allow these questions to be addressed.

It was also somewhat surprising that the five Clade 5 strains analyzed during murine

infection caused minimal to no disease (**Figure 8**) given that all of these strains were isolated

from human patients experiencing CDI-related disease symptoms (**Table 1**). To our knowledge,

very few Clade 5 strains have been analyzed during murine infection, with one study observing

minor disease symptoms in mice for two Clade 5 strains three days post-infection, despite one

of the strains causing severe disease in humans (20). Interestingly, RT078 Clade 5 strains

frequently cause asymptomatic infections in agricultural animals (31, 59) and mice may be

important vectors of transmission in these settings (60, 61). Thus, it is intriguing that our

preliminary analyses suggest that Clade 5 strains persist at high levels in the murine gut over

time compared to the Clade 1 strain 630 (**Figure 8**, $p < 0.005$, except TAL29996). Identifying

factors that allow Clade 5 strains to grow more quickly would provide critical insight into

whether this faster growth rate contributes to the persistence of these strains in mice.

Another unexpected finding was that the chain length of Clade 5 strains does not

correlate with their propensity to sporulate (**Figure 6**) because the strains tested varied wildly in

their sporulation frequencies. Overall, Clade 5 strains sporulated at high levels (>30%), with the

exception of strain TAL29600, which sporulates ~5,000-fold less efficiently than the other

Clade 5 strains, **Figure 6**). While little is known about the mechanisms regulating sporulation

initiation outside of strains 630 and R20291, future analyses of strain TAL29600, which

sporulates poorly under laboratory conditions (**Figure 6**) but not during murine infection

(**Figure 8**), could provide insight into the molecular determinants of sporulation initiation in

Clade 5 strains. For example, differences in gene presence or polymorphisms in several c-di-

GMP-related genes were observed in TAL29600 relative to the other Clade 5 strains (**Table

S2**), and c-di-GMP has also been implicated in regulating sporulation initiation events through

an unknown mechanism (62, 63).

Finally, a key finding of our time-lapse microscopy analyses is that *C. difficile* strains

generally undergo cell separation inefficiently after cell division, leading to the formation of

chains of two cells (Clades 1, 2, and 4) or long chains (most Clade 5 strains). While there was a

significant delay between cell division and cell separation in the latter case, the Clade 1 strain

630 and Clade 5 strain TAL29996, formed chains of two cells that were typically detected by

DeLTA as a single cell because cell separation had not yet initiated (**Figure 4**, red dot). Thus,

cell division and cell separation in *C. difficile* are not as tightly correlated as they are in many

bacteria.

The lack of coordination likely relates to the absence of FtsEX homologs in *C. difficile*,

which couple septal PG synthesis with PG hydrolases that mediate cell separation (64-67).

Indeed, recent work has identified novel factors that control cell chaining in *C. difficile*. The

CwlA peptidoglycan hydrolase is required for cell separation (68), and its activity is controlled

by the Ser/Thr kinase PrkC: phosphorylation inhibits cell separation by preventing CwlA export

(68). Since the septum-localizing MldA or MldB proteins also promote chaining in *C. difficile*

through unknown mechanisms (69), future work could address whether the inefficient cell

separation phenotype of Clade 5 strains is due to decreased CwlA export or MldA/MldB levels.

Importantly, our insights into the basic physiology of *C. difficile* were enabled by our

development of a facile method for conducting time-lapse microscopy under anaerobic

conditions. We anticipate that the methods used in this manuscript can be applied by many

anaerobic microbiologists to study the growth properties of their organisms(s) of interest, under

different conditions and mutant backgrounds. This is because the growth chamber set-up

involves only commercially available GeneFrames and open-source software for rapidly

conducting automated image analyses of time-lapse microscopy data (41, 42). Our anaerobic

set-up could be further coupled with recently developed, fluorogen-activated anaerobic imaging

tags (70) to facilitate single-cell analyses of gene-specific transcription during anaerobic growth

and dynamic protein localization studies (71). Thus, there are many potential applications for

the simple methods described here for studying the growth of anaerobes over time at the single-

cell level.

575
576   **Materials and Methods**

577
Bacterial strains and growth conditions
All *C. difficile* strains were grown on brain heart infusion (BHIS) medium supplemented
with 0.5% w/v yeast extract and 0.1% w/v L-cysteine with taurocholate (TCA; 0.1% w/v; 1.9
mM). Strains were sub-cultured into tryptone yeast extract (TY) broth supplemented with 0.1%
w/v L-cysteine (TYC medium) prior to inoculation onto the time-lapse microscopy agarose
pads. All strains were grown at 37°C under anaerobic conditions using a gas mixture of 85%
hydrogen, 5% CO_2, and 10% H_2. For time-lapse experiments, 1.5% agarose pads supplemented
with TYC medium were used as described above. Sporulation analyses were carried out on
70:30 medium (70% BHIS and 30% SMC) for 24h as described previously (72).

587
Anaerobic time-lapse imaging of C. difficile growth
All imaging was carried out on a Leica DMi8 inverted microscope with a HC plan
apochromat 63x 1.4 NA oil immersion phase contrast objective. Fluorescent membrane staining
experiments were done with a Lumencor Spectra X light source, coupled with an XLED-QP
quadruple-band dichroic beam-splitter (Leica) (transmission: 415, 470, 570, and 660 nm) along
with an external emission filter wheel (Leica). FM4-64 was excited using a 470nm LED
through a 470/20nm excitation filter and emitted light was filtered through a 590/50nm
emission filter and captured with a Leica DFC9000GTC sCMOS camera. All experiments were
carried out at 37°C using a microscope incubation system (Pecon), Leica Adaptive Focus
Control hardware autofocus, and a high precision stage (Pecon) were used for all imaging
experiments.
For time-lapse imaging of *C. difficile* growth, all bacterial strains were grown in 2 mL
liquid TY medium to a turbid OD600 > 2-3; after 2 hours of growth, bacteria were diluted 1:100
for Clade 5 strains and all other strains were diluted 1:50 in fresh media and grown to mid-log
phase (OD600 0.4-0.7).

An imaging chamber with a gas-tight seal was constructed using a 125 µL Gene
Frame (Thermo Fisher) adhered to a glass slide generating a well for growth medium. The slide
was then transferred to the anaerobic chamber. In the anaerobic chamber, the gene frame was
filled with 500µl 1.5% Top vision low melting point agarose and tryptone yeast extract media
containing 0.1% w/v L-cysteine to scavenge oxygen and maintain anaerobic conditions. While
the agarose was molten, a second clean slide was placed over the top and the agar pad was
placed on a frozen small freezer block (for holding PCR strip tubes) for 10-30 minutes until the
agarose-media mixture was solid. For experiments using FM4-64, agarose pads were made the
same way, with the addition of FM4-64 to a final concentration of 1 µg/mL directly to the
agarose/media solution prior to making the agar pad.
The agar pad was dried for 5-10 minutes until visible liquid on the surface of the pad
was evaporated. 1 µL of mid-log cells were spotted on the pad, dried, and a #1.5 coverslip
(VWR) was adhered to the Gene Frame. The cells were imaged at 37°C until they reached
confluency in the field of view. This was anywhere from 2.5 hours for Clade 5 strains to 6 hours
for Clades 1-4 for all experiments with images taken at 5-minute intervals.
618
Image analysis, computing hardware, and statistical analysis.
All movie frames were trimmed to the point when cells were not overlapping and out of
focus regions were cropped. The resulting images were analyzed using the Python library
DeLTA 2.0 (41, 42). All image and data analyses were done on a PC running Windows 10
equipped with an AMD Ryzen 5900HX 8-core CPU, 32GB DDR4 RAM, 2 1TB NVME SSDs,
and an NVIDIA RTX3080 GPU with 16GB VRAM. Analysis of the output data and data
visualizations were done in Python using Matplotlib/Seaborn, Pandas, Numpy, Scipy, and the
Statannotations library.
627
Septum detection during live-cell microscopy
The image processing starts with the masks generated using DeLTA 2.0. First, we performed
erosion on the mask with a disk of radius of 1 pixel to avoid effects of membrane fluorescence.
The photo was cropped following the eroded mask contour. Hence, only the interior of the
contour was considered. Pixel values were rescaled such that the minimum pixel value of inner
pixels was mapped to 0 and the maximum was mapped to 255. We set a threshold intensity as
the pixel value of the 95% quantile of pixel intensities. The image was slightly blurred by
convolution with a Gaussian filter with standard deviation of 2 pixels. The septum corresponds
to the contours found after performing an adaptive thresholding based on thresholds defined
using a Gaussian-weighted method. This thresholding was performed using the open cv2 library
in Python. The resultant contours were manually verified from videos generated using the
tracking from the outputs from DeLTA. The manual validation mainly involved filling in the
time points where the algorithm missed a ring that was visible in the video. We did not assume
the existence of a ring if the algorithm did not detect it in a previous frame.
642
Cell length estimation
We measured the cell projected area from the DeLTA contours of the images as the pixel amount
of the contours. However, estimating cell length was challenging because some cells were very
long and bent. To overcome this problem, we selected 30 images of three different cells (from
strains 630, TAL3050 and TAL28131) that were straight and had different lengths. For these
cells, we calculated the cell length as the longest side of the minimum bounding rectangle of the
contour. From these lengths of straight cells, we also estimated the best cell width as the average

of the projected area divided by the length. Considering the extreme cell length, the effects of the
rounded tips were negligible and the rectangle shape adequately approximated length. We
verified that this mean value showed low variability for the three strains. Then, we used this
width value to estimate the length of all the cells, including the bent ones, by dividing their
projected area by the width. This way, we obtained a consistent measure of cell length that was
independent of bending.
656
Elongation rate estimation
We tracked cell size over time and identified the division points as the ones where the cell size
(projected area) dropped by more than 30% compared to the current cell size value. We fitted an
exponential function (with base e) of time to the data points between two divisions and estimated
the elongation rate from the exponent of the best fit. We expressed the elongation rate in
doublings/hr, which means how many times the cell size doubles in one hour. For example, an
elongation rate of 2 doubling/hr means that the cell size doubles two times in 1 hr, which
corresponds to an exponent of $2\ln(2)$ 1/hr. For the statistics, we only included the elongation
rates that had a high quality of fit, with an R^2 coefficient greater than 0.9.
666
Bulk growth measurements
Starter cultures were grown until early stationary phase in BHIS (or TYC medium as indicated)
then diluted 1:50 into BHIS (or TYC medium). When the cultures reached an OD_{600} of 0.5, they
were diluted 1:25 into 200 µL of either BHIS or TYC in a flat 96 well polystyrene plate
(CellTreat). The OD_{600} was analyzed every 15 min for 24 hrs in a BioTek Epoch plate reader
with shaking. Bulk growth measurements are based on a minimum of three independent
replicates across a minimum of 2 experiments.
674
Cell wall labeling
HADA (Tocris Bioscience) was added to exponentially growing cell culture to a final
concentration of 50-100 µM and incubated for ~2 mins before cell fixation. Cells were fixed as
previously described (73). Briefly, 400 µL of cell suspension was added to 100 µL of a 5X
fixation solution containing paraformaldehyde and $NaPO_4$ buffer. Samples were mixed and
incubated in the dark for 30 min at room temperature, followed by 30 min on ice. Fixed cells
were washed three times in phosphate-buffered saline (PBS) and resuspended in ~50 µL of PBS.
Cells were imaged within 72 hours after fixation.
683
Sporulation assays.
Starter cultures were grown until early stationary phase in BHIS then diluted 1:50 into BHIS.
When the cultures reached an OD_{600} between 0.35 and 0.75, 120 µL of the culture was spread
onto 70:30 (70% SMC media and 30% BHIS media) agar plates (40 ml media per plate) and then
incubated for 20-24 hrs before the sporulating cells were scraped from the plate into phosphate-
buffered saline (PBS). Sporulation levels were visualized by phase-contrast microscopy as
previously described (74).
691
692   *Heat resistance assay.*
693   Heat-resistant spore formation was measured 20-24 hrs after sporulation was induced on 70:30
694   agar plates as previously described (72). The percent sporulation of given culture represents the
695   ratio of heat-resistant colony-forming units (CFUs) to total CFUs. Percent sporulation was
696   determined from a minimum of 3 biological replicates.

Spore purification.

Spores were purified as previously described (75) by scraping up sporulating cells incubated on
70:30 medium for 3 days into ice-cold H_2O. The cells were washed several times in ice-water
over the course of a day and incubated on ice overnight. The following morning, the sample was
pelleted, and cells were suspended in 1 X DNAse buffer (New England Biolabs) and then treated
with DNAse (New England Biolabs) for 30 min at 37˚C. The samples were washed one more
time before being resuspended in 20% Histodenz and then layered onto a 50% Histodenz layer.
The resulting mixture was pelleted, and the supernatant was aspirated off using a vacuum
aspirator. The pelleted spores were washed in ice-cold water 2-3 times and the optical density of
the purified spores was measured.

Genomic DNA preparation.

Starter cultures were grown until early stationary phase in BHIS then back-diluted 1:50 into
BHIS and grown until an OD_{600} of around 0.7-0.8 was reached. 10 mL of the culture was
pelleted and then frozen at -80˚C. After thawing the sample, it was resuspended in a 25%
sucrose TE buffer (10 mM Tris, 1mM EDTA), incubated with 100 mg/mL lysozyme for 37˚C
for 1 hr. After the cultures tarted to lyse, proteinase K, RNAse A, EDTA, Sarkosyl, and NaCl
was added. Phenol:Chloroform:IAA (25:24:1) was added to extract proteins, gently mixed, and
then the sample was pelleted to separate the phenol and aqueous layer. The aqueous layer was
then added to Chloroform:IAA (24:1), mixed gently, then centrifuged. The aqueous layer was
then precipitated using isopropanol and incubated at -20˚C for a minimum of 15 min. The
precipitated DNA was pelleted and then washed with 70% ethanol. The pellet was air dried and
then gently resuspended in 10 mM Tris pH 8.0 elution buffer.

Genomic analyses.

Genomic DNA was sequenced by MiGS at the University of Pittsburgh (now SeqCenter)
according to their standard protocol. Libraries were sequenced on an Illumina NextSeq 500
platform to generate paired-end 150 bp reads. Illumina reads of RT078 genomes were assembled
into contigs using SPAdes (v3.13.0),(76) and genes were called and annotated using Prokka
(v1.11) (77). Assembled and annotated contigs of five RT078 strains (TAL28131, TAL29600,
TAL29996, TAL30550, TAL30574) were applied for pangenomic analysis. Default settings
were used based on the Anvi'o workflow for microbial pangenomcis with adjustments for minbit
as 0.5 and mcl-inflation as 10(78-80). For SNPs analyses, reads of five RT078 genomes were
aligned to the reference M120 and variants were called by breseq (v. 0.38.1) by default
settings.(81)

qPCR analyses.

735 Each genomic DNA sample was analyzed by qPCR with primers that amplify the *cmr*-ON
736 sequence orientation, the *cmr*-OFF orientation, or the reference gene *rpoA* (32) . Each 20-µL
737 qPCR reaction consisted of 100 ng genomic DNA, 100 nM primers, and SensiMix™ SYBR®
738 reagents (Bioline). The reactions were run on a LightCycler® 96 (Roche Diagnostics), and *cmr*
739 switch orientation frequencies were calculated as described previously (44).

Surface motility assays

742 Starter cultures were grown until early stationary phase in BHIS then back-diluted 1:50 into
743 BHIS and grown until an OD_{600} of 0.5 was reached. 10 µL of the exponential-phase cultures
744 were then spotted onto BHIS plates and incubated at 37˚C for 5 days after which the plates were
745 scanned using a flatbed scanner.
746
Mouse infection experiments
Conventional 7-week-old C57BL/6 female mice from Jackson Laboratories were housed in a
sterile (autoclaved cage and bedding) large cage (24"x17") with autoclaved water and irradiated
food (Teklad 2918) for 10 days to allow for normalization of microbiota across mice through
coprophagy. After the 10-day normalization period, mice were started on cefoperazone, which
was added to their water at a concentration of 0.5mg/ml. Mice were allowed to drink the
cefoperazone water ad libitum for 10 days, after which they were placed back on sterile water
without antibiotic. After a 2-day period of being on normal sterile water, mice were weighed and
given a single dose of clindamycin (10mg/kg) via intraperitoneal injection. Immediately after IP
injection, mice were moved to standard-size autoclaved mouse cages (4 mice per cage) with
sterile food and water. 24 hours following the clindamycin injection, mice were inoculated with
1×10^5 spores of *C. difficile* (in 1xPBS) via oral gavage using a metal, reusable needle. Mice
were weighed by being placed in a plastic Nalgene cup on top of a scale. Fecal pellets were
collected just prior to oral gavage to ensure no prior *C. difficile* colonization and to note a
baseline weight. Following inoculation, mice were weighed to monitor % weight change over
time on days 1-4, 7, and 14. Fecal pellets were collected in duplicate on the same days for *C.*
difficile CFU enumeration and qPCR for detection of *cmr* switch orientation. On day 14, a
terminal weight was taken, and a fecal pellet was collected, followed by sacrifice via CO_2
inhalation with cervical dislocation as secondary method of euthanasia. Two experimental
replicates were completed using 4 mice per group, resulting in a total of 8 mice per condition.
767
C. difficile CFU enumeration from mouse fecal pellets
C. difficile engraftment in mice was monitored over a 14-day period. Fecal pellets were collected
from mice on days 0-4, 7, and 14. Each pellet was weighed and then suspended in 1x PBS. 10 µL
of the suspension was then serially diluted 1:10 in 1x PBS in a 96 well plate, and 5 µL was
spotted onto TCCFA agar to select for *C. difficile*, such that the dilutions plated were $2 \times 10^3 - 2$
$\times 10^7$. *C. difficile* colonies were counted 24 hours after plating to allow for sufficient growth. *C.*
difficile CFUs were normalized by gram of fecal material.
775
776 *Data accessibility statement*
777   The data set and data processing scripts are publicly available
778 at https://doi.org/10.5281/zenodo.13352469 (82).
779
Author contribution statement
J.W.R. conducted all live-cell microscopy experiments and analyzed the single-cell data.
C.N. generated data visualizations from the time-lapse microscopy data, developed the method
for identifying septa within chains that had not initiated cell separation, and calculated
elongation rates and cell lengths. A. Singh advised C.N. in these quantitative analyses. I.N. and
A.S. conducted sporulation analyses and liquid culture experiments. J.W.R., R.T., and A.S.
conceptualized all experiments with input from A.Singh. Q.D. and A.S. extracted genomic
DNA and visualized cells at different growth phases. Q.D. conducted all the genomic analyses

and genome comparisons. N.D. and I.N. conducted spore purification analyses and imaging. N.
B.A. and I.E.M. developed and optimized methods for constructing agarose pads that could
support anaerobic growth in an ambient atmosphere. M.J.D. provided input on the DeLTA
analysis pipeline. N.D. conducted the animal experiments. A.M. conducted the qPCRs to
determine *cmr* switch orientation in Clade 5 isolates. The manuscript was written by J.W.R and
A.S with input from all authors.
794
795
796 **Table 1.** *Clostridioides difficile* **clinical isolates used in this study**
797

Strain Name	Clade	Ribotype	ST group	Source	Reference
630	1	012	2	Zurich, 1982 (Sanger Institute)	(83)
R20291	2	027	1	London, 2006 (Sanger Institute)	(84)
M68	4	017	81	Dublin, 2006 (Sanger Institute)	(50)
M120	5	078	11	UK, 2007 (Sanger Institute)	(50)
TAL28131	5	078	11	Tufts Medical Center	This paper
TAL29600	5	078	11	Tufts Medical Center	This paper
TAL29996	5	078	11	Tufts Medical Center	This paper
TAL30550	5	078	11	Tufts Medical Center	This paper
TAL30574	5	078	11	Tufts Medical Center	This paper
V48	5	078	11	Brigham & Women's Hospital	This paper
139b	5	078	11	Memorial Sloan Kettering	(20)
WU66	5	078	11	Barnes-Jewish Hospital	(20)
BBL2	1	012	2	Memorial Sloan Kettering	(20)
WU38	1	012	2	Barnes-Jewish Hospital	(20)
190B	1	087	46	Memorial Sloan Kettering	(20)
Wup14	2	027	1	Barnes-Jewish Hospital	(20)
BBL4	2	027	1	Memorial Sloan Kettering	(20)
186A	2	027	1	Memorial Sloan Kettering	(20)

798
799 Memorial Sloan Kettering Cancer Center (MSK), Barnes-Jewish Hospital (BJH)
800
801
802 **Table 2: Cell size and growth rate statistics for the studied strains.**
803

Strain Name	Clade	Mean Growth rate	Mean Cell Length (um)	S.D. Cell Length	Mean Chain Length (um)	S.D. Chain Length
630	1	1.63±0.02	8.51±0.06	2.24±0.04	12.53±0.11	3.46±0.07
R20291	2	1.88±0.01	not measured	not measured	11.72±0.08	3.15±0.05
M68	4	1.64±0.01	not measured	not measured	13.69±0.08	4.33±0.06
M120	5	2.15±0.01	not measured	not measured	21.8±0.2	10.2±0.2
TAL28131	5	2.17±0.02	11.15±0.06	3.34±0.04	19.7±0.2	6.9±0.1
TAL29600	5	2.11±0.02	14.0±0.1	4.44±0.08	29.1±0.4	13.1±0.3
TAL29996	5	2.15±0.02	9.44±0.04	2.70±0.03	13.65±0.06	3.87±0.04
TAL30550	5	2.05±0.04	11.82±0.08	3.34±0.05	27.3±0.3	9.8±0.3

V48	5	2.10±0.02	10.80±0.08	3.75±0.06	19.4±0.2	8.5±0.2

804
805 S.D. stands for standard deviation. The ± interval represents the 95% confidence interval of the
806 estimated metric.
807
808 **References**
809

1. Desai K, Gupta SB, Dubberke ER, Prabhu VS, Browne C, Mast TC. 2016.
Epidemiological and economic burden of *Clostridium difficile* in the United States:
estimates from a modeling approach. BMC Infect Dis 16:303.
2. Guh AY, Mu Y, Winston LG, Johnston H, Olson D, Farley MM, Wilson LE, Holzbauer
SM, Phipps EC, Dumyati GK, Beldavs ZG, Kainer MA, Karlsson M, Gerding DN,
McDonald LC, Emerging Infections Program *Clostridioides difficile* Infection Working
G. 2020. Trends in U.S. Burden of *Clostridioides difficile* Infection and Outcomes. N
Engl J Med 382:1320-1330.
3. Shen A. 2020. *Clostridioides difficile* Spore Formation and Germination: New Insights
and Opportunities for Intervention. Annu Rev Microbiol 74:545-566.
4. Abt MC, McKenney PT, Pamer EG. 2016. *Clostridium difficile* colitis: pathogenesis and
host defence. Nat Rev Microbiol doi:10.1038/nrmicro.2016.108.
5. Theriot CM, Young VB. 2015. Interactions Between the Gastrointestinal Microbiome
and *Clostridium difficile*. Annu Rev Microbiol 69:445-61.
6. Kordus SL, Thomas AK, Lacy DB. 2022. *Clostridioides difficile* toxins: mechanisms of
action and antitoxin therapeutics. Nat Rev Microbiol 20:285-298.
7. Maroo S, Lamont J. 2006. Recurrent *Clostridium difficile*. Gastroenterology 130:1311-
1316.
8. Zanella Terrier MC, Simonet ML, Bichard P, Frossard JL. 2014. Recurrent *Clostridium*
difficile infections: the importance of the intestinal microbiota. World J Gastroenterol
20:7416-23.
9. Knight DR, Elliott B, Chang BJ, Perkins TT, Riley TV. 2015. Diversity and Evolution in
the Genome of *Clostridium difficile*. Clin Microbiol Rev 28:721-41.
10. Knight DR, Imwattana K, Kullin B, Guerrero-Araya E, Paredes-Sabja D, Didelot X,
Dingle KE, Eyre DW, Rodriguez C, Riley TV. 2021. Major genetic discontinuity and
novel toxigenic species in *Clostridioides difficile* taxonomy. eLife 10.
11. Norsigian CJ, Danhof HA, Brand CK, Midani FS, Broddrick JT, Savidge TC, Britton
RA, Palsson BO, Spinler JK, Monk JM. 2022. Systems biology approach to functionally
assess the *Clostridioides difficile* pangenome reveals genetic diversity with
discriminatory power. Proc Natl Acad Sci U S A 119:e2119396119.
12. Knight DR, Kullin B, Androga GO, Barbut F, Eckert C, Johnson S, Spigaglia P, Tateda
K, Tsai PJ, Riley TV. 2019. Evolutionary and Genomic Insights into *Clostridioides*
difficile Sequence Type 11: a Diverse Zoonotic and Antimicrobial-Resistant Lineage of
Global One Health Importance. mBio 10.
13. Seth-Smith HMB, Biggel M, Roloff T, Hinic V, Bodmer T, Risch M, Casanova C,
Widmer A, Sommerstein R, Marschall J, Tschudin-Sutter S, Egli A. 2021. Transition
From PCR-Ribotyping to Whole Genome Sequencing Based Typing of *Clostridioides*
difficile. Front Cell Infect Microbiol 11:681518.

14.   Dingle KE, Elliott B, Robinson E, Griffiths D, Eyre DW, Stoesser N, Vaughan A,
849   Golubchik T, Fawley WN, Wilcox MH, Peto TE, Walker AS, Riley TV, Crook DW,
850   Didelot X. 2014. Evolutionary history of the *Clostridium difficile* pathogenicity locus.
851   Genome Biol Evol 6:36-52.

852   15.   Mengoli M, Barone M, Fabbrini M, D'Amico F, Brigidi P, Turroni S. 2022. Make It Less
853   difficile: Understanding Genetic Evolution and Global Spread of *Clostridioides difficile*.
854   Genes (Basel) 13.

855   16.   Clements AC, Magalhaes RJ, Tatem AJ, Paterson DL, Riley TV. 2010. *Clostridium
856   difficile* PCR ribotype 027: assessing the risks of further worldwide spread. Lancet Infect
857   Dis 10:395-404.

858   17.   Endres BT, Begum K, Sun H, Walk ST, Memariani A, Lancaster C, Gonzales-Luna AJ,
859   Dotson KM, Basseres E, Offiong C, Tupy S, Kuper K, Septimus E, Arafat R, Alam MJ,
860   Zhao Z, Hurdle JG, Savidge TC, Garey KW. 2019. Epidemic *Clostridioides difficile*
861   Ribotype 027 Lineages: Comparisons of Texas Versus Worldwide Strains. Open Forum
862   Infect Dis 6:ofz013.

863   18.   McDonald LC, Killgore GE, Thompson A, Owens RC, Jr., Kazakova SV, Sambol SP,
864   Johnson S, Gerding DN. 2005. An epidemic, toxin gene-variant strain of *Clostridium
865   difficile*. N Engl J Med 353:2433-41.

866   19.   Dong Q, Lin H, Allen MM, Garneau JR, Sia JK, Smith RC, Haro F, McMillen T, Pope
867   RL, Metcalfe C, Burgo V, Woodson C, Dylla N, Kohout C, Sundararajan A, Snitkin ES,
868   Young VB, Fortier LC, Kamboj M, Pamer EG. 2023. Virulence and genomic diversity
869   among clinical isolates of ST1 (BI/NAP1/027) *Clostridioides difficile*. Cell Rep
870   42:112861.

871   20.   Lewis BB, Carter RA, Ling L, Leiner I, Taur Y, Kamboj M, Dubberke ER, Xavier J,
872   Pamer EG. 2017. Pathogenicity Locus, Core Genome, and Accessory Gene Contributions
873   to *Clostridium difficile* Virulence. mBio 8.

874   21.   Merrigan M, Venugopal A, Mallozzi M, Roxas B, Viswanathan VK, Johnson S, Gerding
875   DN, Vedantam G. 2010. Human hypervirulent *Clostridium difficile* strains exhibit
876   increased sporulation as well as robust toxin production. J Bacteriol 192:4904-11.

877   22.   Shaw HA, Preston MD, Vendrik KEW, Cairns MD, Browne HP, Stabler RA, Crobach
878   MJT, Corver J, Pituch H, Ingebretsen A, Pirmohamed M, Faulds-Pain A, Valiente E,
879   Lawley TD, Fairweather NF, Kuijper EJ, Wren BW. 2020. The recent emergence of a
880   highly related virulent *Clostridium difficile* clade with unique characteristics. Clin
881   Microbiol Infect 26:492-498.

882   23.   Li C, Harmanus C, Zhu D, Meng X, Wang S, Duan J, Liu S, Fu C, Zhou P, Liu R, Wu A,
883   Kuijper EJ, Smits WK, Fu L, Sun X. 2018. Characterization of the virulence of a non-
884   RT027, non-RT078 and binary toxin-positive *Clostridium difficile* strain associated with
885   severe diarrhea. Emerg Microbes Infect 7:211.

886   24.   Imwattana K, Rodriguez C, Riley TV, Knight DR. 2021. A species-wide genetic atlas of
887   antimicrobial resistance in *Clostridioides difficile*. Microb Genom 7.

888   25.   Kumar N, Browne HP, Viciani E, Forster SC, Clare S, Harcourt K, Stares MD, Dougan
889   G, Fairley DJ, Roberts P, Pirmohamed M, Clokie MRJ, Jensen MBF, Hargreaves KR, Ip
890   M, Wieler LH, Seyboldt C, Noren T, Riley TV, Kuijper EJ, Wren BW, Lawley TD. 2019.
891   Adaptation of host transmission cycle during *Clostridium difficile* speciation. Nat Genet
892   51:1315-1320.

26.   Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput
ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun
9:5114.

27.   Knetsch CW, Kumar N, Forster SC, Connor TR, Browne HP, Harmanus C, Sanders IM,
Harris SR, Turner L, Morris T, Perry M, Miyajima F, Roberts P, Pirmohamed M, Songer
JG, Weese JS, Indra A, Corver J, Rupnik M, Wren BW, Riley TV, Kuijper EJ, Lawley
TD. 2018. Zoonotic Transfer of *Clostridium difficile* Harboring Antimicrobial Resistance
between Farm Animals and Humans. J Clin Microbiol 56.

28.   Hensgens MP, Keessen EC, Squire MM, Riley TV, Koene MG, de Boer E, Lipman LJ,
Kuijper EJ, European Society of Clinical M, Infectious Diseases Study Group for
Clostridium d. 2012. *Clostridium difficile* infection in the community: a zoonotic disease?
Clin Microbiol Infect 18:635-45.

29.   Goorhuis A, Bakker D, Corver J, Debast SB, Harmanus C, Notermans DW, Bergwerff
AA, Dekker FW, Kuijper EJ. 2008. Emergence of *Clostridium difficile* infection due to a
new hypervirulent strain, polymerase chain reaction ribotype 078. Clin Infect Dis
47:1162-70.

30.   Lim SC, Knight DR, Riley TV. 2020. *Clostridium difficile* and One Health. Clin
Microbiol Infect 26:857-863.

31.   Bolton D, Marcos P. 2023. The Environment, Farm Animals and Foods as Sources of
Clostridioides difficile Infection in Humans. Foods 12.

32.   Garrett EM, Sekulovic O, Wetzel D, Jones JB, Edwards AN, Vargas-Cuebas G, McBride
SM, Tamayo R. 2019. Phase variation of a signal transduction system controls
Clostridioides difficile colony morphology, motility, and virulence. PLoS Biol
17:e3000379.

33.   Thanissery R, Winston JA, Theriot CM. 2017. Inhibition of spore germination, growth,
and toxin activity of clinically relevant C. difficile strains by gut microbiota derived
secondary bile acids. Anaerobe 45:86-100.

34.   Ackermann M. 2015. A functional perspective on phenotypic heterogeneity in
microorganisms. Nat Rev Microbiol 13:497-508.

35.   Aldridge BB, Fernandez-Suarez M, Heller D, Ambravaneswaran V, Irimia D, Toner M,
Fortune SM. 2012. Asymmetry and aging of mycobacterial cells lead to variable growth
and antibiotic susceptibility. Science 335:100-4.

36.   Anjuwon-Foster BR, Tamayo R. 2017. A genetic switch controls the production of
flagella and toxins in *Clostridium difficile*. PLoS Genet 13:e1006701.

37.   Anjuwon-Foster BR, Tamayo R. 2018. Phase variation of *Clostridium difficile* virulence
factors. Gut Microbes 9:76-83.

38.   Kint N, Janoir C, Monot M, Hoys S, Soutourina O, Dupuy B, Martin-Verstraete I. 2017.
The alternative sigma factor sigma(B) plays a crucial role in adaptive strategies of
Clostridium difficile during gut infection. Environ Microbiol 19:1933-1958.

39.   Courson DS, Pokhrel A, Scott C, Madrill M, Rinehold AJ, Tamayo R, Cheney RE,
Purcell EB. 2019. Single cell analysis of nutrient regulation of Clostridioides
(Clostridium) difficile motility. Anaerobe 59:205-211.

40.   de Jong IG, Beilharz K, Kuipers OP, Veening JW. 2011. Live Cell Imaging of Bacillus
subtilis and Streptococcus pneumoniae using Automated Time-lapse Microscopy. J Vis
Exp doi:10.3791/3145.

41. Lugagne JB, Lin H, Dunlop MJ. 2020. DeLTA: Automated cell segmentation, tracking, and lineage reconstruction using deep learning. PLoS Comput Biol 16:e1007673.

42. O'Connor OM, Alnahhas RN, Lugagne JB, Dunlop MJ. 2022. DeLTA 2.0: A deep learning pipeline for quantifying single-cell spatial and temporal dynamics. PLoS Comput Biol 18:e1009797.

43. Pogliano J, Osborne N, Sharp MD, Abanes-De Mello A, Perez A, Sun YL, Pogliano K. 1999. A vital stain for studying membrane dynamics in bacteria: a novel mechanism controlling septation during Bacillus subtilis sporulation. Mol Microbiol 31:1149-59.

44. Garrett EM, Mehra A, Sekulovic O, Tamayo R. 2021. Multiple Regulatory Mechanisms Control the Production of CmrRST, an Atypical Signal Transduction System in *Clostridioides difficile*. mBio 13:e0296921.

45. Sauls JT, Cox SE, Do Q, Castillo V, Ghulam-Jelani Z, Jun S. 2019. Control of Bacillus subtilis Replication Initiation during Physiological Transitions and Perturbations. mBio 10.

46. Eswaramoorthy P, Duan D, Dinh J, Dravis A, Devi SN, Fujita M. 2010. The threshold level of the sensor histidine kinase KinA governs entry into sporulation in Bacillus subtilis. J Bacteriol 192:3870-82.

47. Narula J, Kuchina A, Zhang F, Fujita M, Suel GM, Igoshin OA. 2016. Slowdown of growth controls cellular differentiation. Mol Syst Biol 12:871.

48. Mehdizadeh Gohari I, Edwards AN, McBride SM, McClane BA. 2024. The impact of orphan histidine kinases and phosphotransfer proteins on the regulation of clostridial sporulation initiation. mBio 15:e0224823.

49. Deatherage DE, Traverse CC, Wolf LN, Barrick JE. 2014. Detecting rare structural variation in evolving microbial populations from new sequence junctions using breseq. Front Genet 5:468.

50. He M, Sebaihia M, Lawley T, Stabler R, Dawson L, Martin M, Holt K, Seth-Smith H, Quail M, Rance R, Brooks K, Churcher C, Harris D, Bentley S, Burrows C, Clark L, Corton C, Murray V, Rose G, Thurston S, van Tonder A, Walker D, Wren B, Dougan G, Parkhill J. 2010. Evolutionary dynamics of *Clostridium difficile* over short and long time scales. Proceedings of the National Academy of Sciences of the United States of America 107:7527-7532.

51. Sekulovic O, Mathias Garrett E, Bourgeois J, Tamayo R, Shen A, Camilli A. 2018. Genome-wide detection of conservative site-specific recombination in bacteria. PLoS Genet 14:e1007332.

52. Glomski IJ, Piris-Gimenez A, Huerre M, Mock M, Goossens PL. 2007. Primary involvement of pharynx and peyer's patch in inhalational and intestinal anthrax. PLoS Pathog 3:e76.

53. McKenney PT, Yan J, Vaubourgeix J, Becattini S, Lampen N, Motzer A, Larson PJ, Dannaoui D, Fujisawa S, Xavier JB, Pamer EG. 2019. Intestinal Bile Acids Induce a Morphotype Switch in Vancomycin-Resistant Enterococcus that Facilitates Intestinal Colonization. Cell Host Microbe 25:695-705 e5.

54. Leslie JL, Jenior ML, Vendrov KC, Standke AK, Barron MR, O'Brien TJ, Unverdorben L, Thaprawat P, Bergin IL, Schloss PD, Young VB. 2021. Protection from Lethal *Clostridioides difficile* Infection via Intraspecies Competition for Cogerminant. mBio 12.

982   55.   Theriot CM, Koumpouras CC, Carlson PE, Bergin, II, Aronoff DM, Young VB. 2011.
983   Cefoperazone-treated mice as an experimental platform to assess differential virulence of
984   *Clostridium difficile* strains. Gut Microbes 2:326-34.

985   56.   Sachsenheimer FE, Yang I, Zimmermann O, Wrede C, Muller LV, Gunka K, Gross U,
986   Suerbaum S. 2018. Genomic and phenotypic diversity of *Clostridium difficile* during
987   long-term sequential recurrences of infection. Int J Med Microbiol 308:364-377.

988   57.   Semenyuk EG, Poroyko VA, Johnston PF, Jones SE, Knight KL, Gerding DN, Driks A.
989   2015. Analysis of Bacterial Communities during *Clostridium difficile* Infection in the
990   Mouse. Infect Immun 83:4383-91.

991   58.   Smith AB, Jenior ML, Keenan O, Hart JL, Specker J, Abbas A, Rangel PC, Di C, Green
992   J, Bustin KA, Gaddy JA, Nicholson MR, Laut C, Kelly BJ, Matthews ML, Evans DR,
993   Van Tyne D, Furth EE, Papin JA, Bushman FD, Erlichman J, Baldassano RN, Silverman
994   MA, Dunny GM, Prentice BM, Skaar EP, Zackular JP. 2022. Enterococci enhance
995   *Clostridioides difficile* pathogenesis. Nature 611:780-786.

996   59.   Kim HY, Cho A, Kim JW, Kim H, Kim B. 2018. High prevalence of *Clostridium difficile*
997   PCR ribotype 078 in pigs in Korea. Anaerobe 51:42-46.

998   60.   Moono P, Foster NF, Hampson DJ, Knight DR, Bloomfield LE, Riley TV. 2016.
999   *Clostridium difficile* Infection in Production Animals and Avian Species: A Review.
Foodborne Pathog Dis 13:647-655.

61.   Burt SA, Siemeling L, Kuijper EJ, Lipman LJ. 2012. Vermin on pig farms are vectors for
Clostridium difficile PCR ribotypes 078 and 045. Vet Microbiol 160:256-8.

62.   Dhungel BA, Govind R. 2021. Phase-variable expression of pdcB, a phosphodiesterase,
influences sporulation in *Clostridioides difficile*. Mol Microbiol 116:1347-1360.

63.   Edwards AN, Willams CL, Pareek N, McBride SM, Tamayo R. 2021. c-di-GMP Inhibits
Early Sporulation in *Clostridioides difficile*. mSphere 6:e0091921.

64.   Du S, Henke W, Pichoff S, Lutkenhaus J. 2019. How FtsEX localizes to the Z ring and
interacts with FtsA to regulate cell division. Mol Microbiol 112:881-895.

65.   Li J, Xu X, Shi J, Hermoso JA, Sham LT, Luo M. 2023. Regulation of the cell division
hydrolase RipC by the FtsEX system in Mycobacterium tuberculosis. Nat Commun
14:7999.

66.   Pichoff S, Du S, Lutkenhaus J. 2019. Roles of FtsEX in cell division. Res Microbiol
170:374-380.

67.   Xu X, Li J, Chua WZ, Pages MA, Shi J, Hermoso JA, Bernhardt T, Sham LT, Luo M.
2023. Mechanistic insights into the regulation of cell wall hydrolysis by FtsEX and EnvC
at the bacterial division site. Proc Natl Acad Sci U S A 120:e2301897120.

68.   Garcia-Garcia T, Poncet S, Cuenot E, Douche T, Giai Gianetto Q, Peltier J, Courtin P,
Chapot-Chartier MP, Matondo M, Dupuy B, Candela T, Martin-Verstraete I. 2021.
Ser/Thr Kinase-Dependent Phosphorylation of the Peptidoglycan Hydrolase CwlA
Controls Its Export and Modulates Cell Division in *Clostridioides difficile*. mBio 12.

69.   Ransom EM, Williams KB, Weiss DS, Ellermeier CD. 2014. Identification and
characterization of a gene cluster required for proper rod shape, cell division, and
pathogenesis in *Clostridium difficile*. J Bacteriol 196:2290-300.

70.   Streett H, Charubin K, Papoutsakis ET. 2021. Anaerobic fluorescent reporters for cell
identification, microbial cell biology and high-throughput screening of microbiota and
genomic libraries. Curr Opin Biotechnol 71:151-163.

71. Anjou C, Lotoux A, Zhukova A, Royer M, Caulat LC, Capuzzo E, Morvan C, Martin-
Verstraete I. 2024. The multiplicity of thioredoxin systems meets the specific lifestyles of
Clostridia. PLoS Pathog 20:e1012001.
72. Shen A, Fimlaid KA, Pishdadian K. 2016. Inducing and Quantifying *Clostridium difficile*
Spore Formation. Methods Mol Biol 1476:129-42.
73. Ransom EM, Weiss DS, Ellermeier CD. 2016. Use of mCherryOpt Fluorescent Protein in
Clostridium difficile. Methods Mol Biol 1476:53-67.
74. Pishdadian K, Fimlaid KA, Shen A. 2015. SpoIIID-mediated regulation of sigma(K)
function during *Clostridium difficile* sporulation. Mol Microbiol 95:189-208.
75. Fimlaid KA, Jensen O, Donnelly ML, Francis MB, Sorg JA, Shen A. 2015. Identification
of a Novel Lipoprotein Regulator of *Clostridium difficile* Spore Germination. PLoS
Pathogens doi:10.1371/journal.ppat.1005239.
76. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes
De Novo Assembler. Curr Protoc Bioinformatics 70:e102.
77. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics
30:2068-9.
78. Benedict MN, Henriksen JR, Metcalf WW, Whitaker RJ, Price ND. 2014. ITEP: an
integrated toolkit for exploration of microbial pan-genomes. BMC Genomics 15:8.
79. Eren AM, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, Fink I, Pan JN, Yousef
M, Fogarty EC, Trigodet F, Watson AR, Esen OC, Moore RM, Clayssen Q, Lee MD,
Kivenson V, Graham ED, Merrill BD, Karkman A, Blankenberg D, Eppley JM, Sjodin
A, Scott JJ, Vazquez-Campos X, McKay LJ, McDaniel EA, Stevens SLR, Anderson RE,
Fuessel J, Fernandez-Guerra A, Maignien L, Delmont TO, Willis AD. 2021. Community-
led, integrated, reproducible multi-omics with anvi'o. Nat Microbiol 6:3-6.
80. van Dongen S, Abreu-Goodger C. 2012. Using MCL to extract clusters from networks.
Methods Mol Biol 804:281-95.
81. Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using breseq. Methods Mol Biol
1151:165-88.
82. César N. 2024. Dataset for the article: Unique growth and morphology properties of
Clade 5 *Clostridioides difficile* strains revealed by single-cell time-lapse microscopy. .
Zenodo doi:https://doi.org/10.5281/zenodo.13352469.
83. Sebaihia M, Wren B, Mullany P, Fairweather N, Minton N, Stabler R, Thomson N,
Roberts A, Cerdeño-Tárraga A, Wang H, Holden M, Wright A, Churcher C, Quail M,
Baker S, Bason N, Brooks K, Chillingworth T, Cronin A, Davis P, Dowd L, Fraser A,
Feltwell T, Hance Z, Holroyd S, Jagels K, Moule S, Mungall K, Price C, Rabbinowitsch
E, Sharp S, Simmonds M, Stevens K, Unwin L, Whithead S, Dupuy B, Dougan G,
Barrell B, Parkhill J. 2006. The multidrug-resistant human pathogen *Clostridium difficile*
has a highly mobile, mosaic genome. Nature genetics 38:779-786.
84. Stabler R, He M, Dawson L, Martin M, Valiente E, Corton C, Lawley T, Sebaihia M,
Quail M, Rose G, Gerding D, Gibert M, Popoff M, Parkhill J, Dougan G, Wren B. 2009.
Comparative genome and phenotypic analysis of *Clostridium difficile* 027 strains
provides insight into the evolution of a hypervirulent bacterium. Genome biology 10.
1070