1

Unique growth and morphology properties of Clade 5 *Clostridioides difficile* strains

revealed by single-cell time-lapse microscopy

John W. Ribis[1,2], César A. Nieto-Acuña[3], Qiwen Dong[1], Nicholas V. DiBenedetto[1,2], Anchal

Mehra,[4] Irene Nagawa,[1] Imane El Meouche,[5] Bree B. Aldridge,[1] Mary J. Dunlop,[6] Rita Tamayo,[4]

Abyudhai Singh,[3] Aimee Shen[1*]

Affiliations

[1]Department of Molecular Biology and Microbiology, Tufts University School of Medicine,

Boston, Massachusetts, USA, [2] Graduate School of Biomedical Sciences, Tufts University

School of Medicine, Boston, Massachusetts, USA. [3]Department of Electrical and Computer

Engineering, University of Delaware, Newark, DE 19716, USA. [4]Department of Microbiology

and Immunology, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill,

NC, USA, [5]INSERM, Université de Paris Cité, Université Sorbonne Paris, Nord, IAME, F-

75018, Paris, France, [6]Biomedical Engineering, Boston University, Boston, MA 02215.

[*]Address correspondence to
Aimee Shen, aimee.shen@tufts.edu
Phone number: (617)636-3792

Abstract

Clostridioides difficile is a major One Health threat as an important gastrointestinal pathogen of
both humans and agricultural animals. The *C. difficile* species can be subdivided into 5 main
clades, with Clade 5 currently undergoing speciation from Clades 1-4. Since Clade 5 strains are
found more frequently in agricultural animals and can cause zoonotic infections, Clade 5 strains
likely have evolved phenotypes that distinguish them from Clade 1-4 strains. Here, we compare
the growth properties of Clade 5 strains to Clade 1, 2, and 4 strains using an anaerobic time-lapse
microscopy system coupled with automated image analysis. These analyses revealed that Clade 5
strains grow faster than Clade 1, 2, and 4 strains and are more likely to form long chains of cells.
Notably, the chaining phenotype was not shared among all Clade 5 strains examined, since 1 of
the 9 strains did not form chains. Genomic analyses of the Clade 5 strains revealed that the
orientation of the *cmr* switch, an invertible DNA element controlling the expression of a signal
transduction system that regulates chaining, correlates with the propensity of a given Clade 5
strain to form chains. Taken together, Clade 5 strains appear to have distinct growth properties
that allow them to inhabit more diverse ecological niches.

Introduction

Clostridioides difficile is a leading cause of nosocomial infection in the United States, with approximately 500,000 new infections and 14,000 deaths attributed to this organism annually (1, 2). As an obligate anaerobe, *C. difficile* relies on its ability to form hardy, metabolically dormant spores to survive outside the gut environment and transmit disease (3). When *C. difficile* spores are ingested by susceptible hosts (4, 5), they encounter bile salt signals in the small intestine that trigger their germination. These stimuli allow them to exit dormancy and outgrow into vegetative cells, which colonize the colon and secrete toxins that damage gut epithelial tissue (6). This damage triggers a significant inflammatory response that can cause disease pathologies ranging from mild diarrhea to pseudomembranous colitis, and even death (2, 4). Beyond these acute complications, *C. difficile* frequently causes recurrent infections (~20% of infections), which can lead to more severe disease symptoms (14–16) (2, 7, 8).

Notably, as a species, *C. difficile* exhibits tremendous genetic diversity (9-11). Genomic analyses indicate that *C. difficile*'s core genome represents only ~10-20% of its pan-genome (9, 10). The extreme plasticity of the "open" pan-genome of *C. difficile* likely helps the bacterium inhabit the gastrointestinal tract of diverse animals, from mammals to invertebrates, and environmental reservoirs like sewage and compost (12). Multi-locus sequence typing (MLST) analyses have revealed that the *C. difficile* species comprises 5 distinct phylogenetic clades that can be further subdivided into different ribotypes (RTs) or sequence types (STs) (10, 13). While whole-genome sequencing analyses identified three additional cryptic clades, subsequent analyses have revealed that the genomic divergence between the 5 main clades and three additional cryptic clades is so significant that the cryptic clade species fall below the threshold determined by average nucleotide identity (ANI) to qualify as part of the *C. difficile* species (10, 14).

Between the five remaining phylogenetic clades, there are notable differences in geographic and host distributions. Clade 1 is the largest and most heterogeneous (and likely ancient) clade, containing over 200 STs that have the broadest geographic distribution (15, 16). It includes both toxin-producing and non-toxigenic strains, including the well-characterized, genetically tractable toxigenic strain 630 (9). Clade 2 harbors epidemic-associated strains found within the ribotype 027 (RT027) lineage. Strains from this ribotype have been associated with

outbreaks in hospitals, particularly in North America, due to their frequent resistance to

fluoroquinolones (17-19). These epidemic strains can also cause severe disease symptoms in

part due to their production of three toxins: TcdA, TcdB, and CDT (binary toxin) (6). Although

RT027 strains have frequently been associated with "hypervirulence," there is considerable

phenotypic diversity within this lineage with respect to virulence, toxin production levels, and

sporulation (20-22).

Clade 3 strains are relatively uncommon and harbor a unique cell surface due to their

lack of CwpV production (23), but phenotypic analyses of biofilm formation and motility

suggest that they share similarities with Clade 2 strains like R20291 (24). Clade 4 contains

RT017 (ST37) strains that only encode a single toxin, TcdB (i.e., TcdA–CDT–), and are often

clindamycin- and fluoroquinolone-resistant. RT017 strains have been associated with outbreaks

in Europe and North America and are the most common strains found in Asia (25). Clade 5 is

the most genetically distant from the other 4 main *C. difficile* clades (94) and is thought to have

emerged before Clades 1-4 (26). While the ANI for Clades 1-4 ranges between 97.1 - 99.8%,

Clade 5 strains exhibit ANI values around 96%, which is close to the ANI value demarcation

used by NCBI to define organisms of the same species (10, 14). Thus, Clade 5 strains appear to

be actively diverging from Clades 1-4 (10).

Clade 5 strains are an increasing problem in healthcare and agricultural settings because

they can cause severe disease in humans and are commonly found in livestock, particularly pigs

(12, 27). While other *C. difficile* strains have been known to infect both humans and animals,

only Clade 5 strains have been associated with zoonotic transmission from both animal-to-

human and human-to-animal. The mechanisms underlying this bidirectional zoonotic

transmission are poorly understood, but the increased carriage of antimicrobial resistance genes

by Clade 5 strains may contribute to their ability to persist in livestock and humans. The

prevalence of antibiotic resistance genes in Clade 5 strains has major implications for One

Health, particularly with respect to antibiotic use in agriculture and humans (12, 27, 28).

Furthermore, Clade 5 strains frequently cause community-acquired infections, which likely

reflects wide environmental distribution and highlights the importance of understanding the

unique properties of this group of strains. Indeed, a recent genomic analysis suggests that

RT078/ST11 strains within Clade 5 frequently carry zinc acquisition and homeostasis genes

(11).

Despite numerous genomic analyses revealing the remarkable genetic diversity of *C.*
difficile strains, relatively few studies have investigated the phenotypic diversity between strains
from different clades. Clade-specific differences in colony morphology between Clade 5 strains
relative to Clade 1-4 strains have been described in a limited set of analyses (26, 29), suggesting
that differences in growth and/or cellular morphology may exist within clades. While
differences in bulk growth rates between *C. difficile* strains have been reported (30), most
phenotypic analyses have been conducted on a limited subset of strains within a given clade.
Indeed, no systematic comparisons of the growth properties of the different clades or
comparisons of cell morphology have been performed to date.
Here, we compare the growth properties of multiple strains derived from the four
predominant main phylogenetic clades of *C. difficile* (Clade 1, 2, 4, and 5) using anaerobic
time-lapse microscopy. These analyses unexpectedly revealed striking differences in the growth
and cell morphology of the Clade 5 lineage relative to strains from Clades 1, 2, and 4. These
morphological differences correlate with the orientation of the *cmr* switch, which has previously
been implicated in modulating cell and colony morphology and cell chaining (31). Thus, the
cmr switch appears to be an important contributor to the chaining phenotype frequently
observed in Clade 5 strains.
122
123   **Results**
124
125   **Development of a simple method for time-lapse imaging under anaerobic conditions**
126
Time-lapse imaging of single cells has been widely used in bacteria due to the growing
appreciation that clonal bacterial populations can exhibit considerable phenotypic heterogeneity,
with implications for antibiotic resistance and virulence (31-35). However, live single-cell
analyses in *C. difficile* have been complicated by its inability to grow in the presence of
atmospheric oxygen (36). While there have been limited reports of live cell time-lapse
microscopy of *C. difficile*, these analyses have required custom growth chambers to maintain
anaerobic conditions (37), which may limit the accessibility of these experimental systems to
most investigators.

To overcome these limitations, we established a simple system that relies solely upon

commercially available reagents and materials to grow *C. difficile* cells under anaerobic

conditions. This system uses gas-tight, adhesive Gene Frames™, which have been used

extensively in imaging applications for bacteria (38). Notably, the gas-impermeability of these

commercial seals allows maintenance of anaerobic conditions when the agarose pads, which

contain growth media, are prepared in the anaerobic chamber (**Figure 1**).

142
143
Figure 1. Schematic of the anaerobic single-cell imaging set-up. Exponentially growing *C. difficile*
cells in TY medium supplemented with cysteine (TYC) are spotted onto a 1.5% agarose pads formed
within gas-tight adhesive Gene Frames inside the anaerobic chamber. Up to 6 strains can be spotted
onto a pad. The pad is sealed with a coverslip, and the imaging chamber is removed from the anaerobic
chamber and transferred to a heated (37˚C) microscope stage. Time-lapse microscopy is used to
visualize the growth of individual bacterial cells for 2-6 hrs. The output data is segmented and tracked
with the DeLTA 2.0 Python package (191). An example filmstrip of strain 630 grown on TYC medium
over time (Bottom).

Gene Frames also allow highly uniform, thick agarose pads to be generated, which are properties

critical for conducting automated time-lapse microscopy analyses and enabling *C. difficile* to

grow in a sealed system under ambient conditions.

Time-lapse microscopy reveals clade-specific differences in elongation rate and cell length

Following the establishment of our time-lapse imaging setup, we compared the single-

cell growth properties of representative *C. difficile* strains from Clade 1 (630, ribotype (RT)

012), Clade 2 (R20291, RT027), Clade 4 (M68, RT017), and Clade 5 (M120, RT078) (**Figure**

2, Table 1). We did not have access to Clade 3 strains, so these were not included in this study.

The four "representative" strains were all isolated from patients with *C. difficile*-associated

disease and are frequently used as reference genomes for their clades and ribotype groups.

Notably, RT027 (ST1), RT017 (ST45), and RT078 (ST11) strains are from ribotypes/

multilocus sequencing types that are frequently isolated from patients with *C. difficile* infection

(CDI) (12, 19, 22, 25).

To quantify the growth properties of single cells visualized by time-lapse microscopy,

we used DeLTA 2.0, a Deep Learning for Time-lapse Analysis (DeLTA) software that allows

rapid and accurate segmentation and tracking of bacteria growing in two dimensions on agarose

pads (39, 40). This software uses deep convolutional neural networks to obtain information on

the growth properties of individual cells growing in microcolonies on agarose pads. The

segmentation and tracking of *C. difficile* cells was highly accurate (**Figure 1**), and minimal user

input or post-image processing was needed to obtain growth property measurements.

Robust growth was observed for all strains using our system, with Clade 5 strain M120

elongating the fastest (2.1 doublings/hr), followed by Clade 2 strain R20291 (1.8 doublings/hr),

and then Clade 1 strain 630 and Clade 4 strain M68 (1.6-1.7 doublings/hr) (**Fig 2A**). This

elongation rate was defined as doublings/hr to indicate the number of times that a cell's size

doubles in one hour. For example, an elongation rate of 2 doublings/hr means that the cell size

doubled twice in 1 hr, which corresponds to an exponent of 2ln(2) 1/hr. Notably, the elongation

rate (doublings/hr) should not be confused with doubling time, or generation time, which are

defined as the length of *time* that it takes before a bacterium divides. Instead, the elongation rate

reflects how fast the cell is increasing in length over time. Nevertheless, elongation rate and

doubling time are related because most cells undergo a division event when they reach twice

their original size.

Figure 2. Clade 5 strain M120 elongates more quickly and exhibits cell chaining. A. Violin plot of
the elongation rates measured during time-lapse microscopy analyses of strains 630 (Clade 1), R20291
(Clade 2), M68 (Clade 4), M120 (Clade 5) grown on TY supplemented with cysteine (TYC) agar. Data
are from three biological replicates, with the mean of each replicate shown as a point on the violin. B.
Violin plot of the cell length measured during time-lapse microscopy for strains 630 (Clade 1), R20291
(Clade 2), M68 (Clade 4), M120 (Clade 5). Each replicate mean is shown as a point on the violin.
Statistical significance for A and B was determined by comparing the mean of the three replicates of
Clade 1, 2, and 4 strains to the Clade 5 M120 strain using a Kruskal-Wallis test * $p < 0.05$. C. Phase-
contrast image from time-lapse microscopy movies. Scale bar is 10 μm. D. Optical density-based
analyses of bulk population growth of the indicated strains in TYC or BHIS medium. The number in
brackets indicates the clade to which a given strain belongs.

199   Importantly, the differences in single-cell elongation rates measured for the four strains

using DeLTA were also observed in bulk population analyses of their growth using optical

density in TYC and BHIS media (**Figure 2D**). These analyses confirmed that the Clade 5 strain

M120 grows faster (using optical density as a proxy for bacterial growth) than the Clade 1, 2,

and 4 strains analyzed. In contrast, negligible differences in bulk growth rates for Clade 1, 2,

and 4 strains were observed (**Figure 2A**), consistent with the relatively subtle difference

observed in the single-cell elongation rates measured for the Clade 2 R20291 strain relative to

the Clades 1 and 4 strains.

The Clade 5 strain M120 exhibited another distinct growth property from the strains

from Clades 1, 2, and 4. While strains from Clades 1, 2, and 4 produced cells of similar length

prior to division, with an average cell size of ~13 μm prior to division, cells of Clade 5 strain

M120 were significantly longer, appearing as long filaments (or chains) with an average length

of 22 μm but chains close to 50 μm readily being measured (**Figure 2B**). These filaments or

chains also appeared to bend readily (**Figure 2C**). Notably, to visualize an entire filament (or

chain), it was necessary to stitch together several fields of view, revealing that the bacterial

filaments (or chains) approached several hundred microns and even up to ~1 mm in length

(**Figure 3**). To determine whether strain M120 forms filaments, i.e. long cells that lack division

septa, or chains of cells, i.e. cells that remain attached despite forming division septa, we

incorporated the lipophilic fluorescent dye FM4-64 which visualizes membranes (Pogliano

1999), directly into the agarose pads. This stain revealed division septa throughout the length of

the cell, indicating that the extremely long cells were not filaments but instead extended chains

of cells (**Figure 3, inset**). Since the spacing between division septa generated cells that were

relatively similar in length to the other *C. difficile* strains analyzed, Clade 5 strain M120 is

distinguished by a decreased efficiency in cell separation.

Clade 5 clinical isolates typically form long chains and grow more quickly than strains

from other clades

Since prior work indicated that Clade 5 strains produce colony morphologies distinct from

Clade 1, 2, and 4 strains (26), we sought to determine whether the (i) striking cell chaining

phenotype and (ii) faster growth rate observed in Clade 5 strain M120 is observed in other

Clade 5 strains. To address these questions, we compared the single-cell growth properties on

TYC agarose of 5 additional Clade 5 clinical isolates obtained from three separate hospitals

(**Table 1**) using time-lapse microscopy analyses (**Figure 2**). Clade 1 strain 630 was included as

a control since it does not form chains under these conditions.

235
236
237 **Figure 3. *C. difficile* Clade 5 can form large heterogenous chains.** Large mosaic phase-contrast
238 image of the Clade 5 *C. difficile* strain M120. Zoomed image shows chains revealed by staining with
239 the membrane dye FM4-64. Image was stitched from 8 individual fields of view at 63x magnification.
240 Scale bar is 100 μm.
241

These analyses revealed that all but one of the Clade 5 strains tested formed long chains,

with TAL29600 forming the longest chains (29 μm on average, **Figure 4A-C**). In contrast,

strain TAL29996 formed shorter chains that were comparable in length to those observed for

strains from Clades 1, 2, and 4 (12-13 μm, **Figures 2B, 4C**). Notably, DeLTA over-estimated

cell length in Clade 5 strains that form chains because it could not identify cells that had formed

division septa (based on FM4-64 staining) but had not yet initiated cell separation.

To overcome this limitation and accurately quantify cell length within long chains, namely the

distance between division septa, we modified our image processing pipeline to use a

thresholding method to detect division septa. After generating masks in DeLTA to segment the

chains, we modified the mask so that only the interior of the contour was analyzed. We then

applied an adaptive thresholding method to identify division septa based on their elevated long

chains do not produce cells that are dramatically longer than strains that do not form chains;

cells within the Clade 5 strains that produced the longest chains had average cell lengths

(TAL29600 and TAL30550, 14 and 12 μm, respectively) that were 30-50% longer than strains

that do not form chains like strains 630 (Clade 1, 8.5 μm) and TAL29996 (Clade 5, 9.4 μm).

Notably, even though TAL29996 did not form long chains like the other Clade 5 strains

analyzed, it elongated at rates similar to those of other Clade 5 strains (2.1 doublings/hr vs. 1.7

260
261
Figure 4. Faster growth is a common feature of Clade 5 strains, but the chaining phenotype is not fully
penetrant in Clade 5 strains. (A) Phase-contrast microscopy images from time-lapse microscopy studies of
Clade 5 strains; Clade 1 strain 630 is included for comparison. Fluorescence microscopy was used to
visualize the FM4-64 stain incorporated into the agarose pads. Masks generated using DeLTA 2.0 are shown,
with the bottom panel showing division septa identified with our adaptive thresholding approach. (B)
Example image showing the parameters identified using DeLTA 2.0 combined with our adaptive threshold
method for detecting division septa within a chain of cells. (C) Cell length measured based on automated
DeLTA 2.0 analyses (pink violin plot) or DeLTA 2.0 combined with adaptive threshold analyses (purple
violin plot). The former method is more likely to measure chain length, while the latter method more
accurately measures cell length. (D) Violin plot of the elongation rates measured for the indicated strains
based on three biological replicates. For all the violin plots shown, each replicate mean is shown as a point on
the violin, and statistical significance was determined by comparing the mean of the three replicates using a
Kruskal-Wallis test * $p < 0.05$.
275

doublings/hr for Clade 1 strain 630**, Figure 4D**). Taken together, these analyses strongly

suggest that Clade 5 strains grow faster than Clade 1, 2, and 4 strains and are more likely to

form chains presumably because their cell separation mechanisms are less efficient.

While the long chains formed by Clade 5 strains limited the number of divisions that

could be analyzed within a given field using time-lapse microscopy, the faster elongation rates

measured on the agarose pads were also observed in bulk analyses of growth in broth culture.

These optical density-based analyses revealed that Clade 5 strains all grow faster than Clade 1,

2, and 4 strains irrespective of their ability to form chains or the growth media used (**Figure**

S1).

To assess whether these conclusions could be generalized to additional Clade 5 strains

vs. Clade 1, 2, and 4 strains, we analyzed the growth of additional strains from all four clades in

broth culture. These analyses confirmed that Clade 1, 2, and 4 strains grow at similar rates,

which are slower than those measured for the nine Clade 5 strains analyzed in both TYC and

BHIS media (**Figure S1**).

Cell chaining in Clade 5 strains is not dependent on growth on a solid medium.

C. difficile has previously been shown to promote cell elongation and chain formation

upon induction of the *cmrRST* locus (31), which encodes a signal transduction system.

Expression of this locus is also responsive to c-di-GMP levels (29), which increases in cells

grown on solid surfaces such as in a biofilm or on an agar plate (29). To determine whether the

chaining phenotype observed in most Clade 5 strains is induced by growth on a surface, we

assessed whether cell chaining would be observed in cells grown in broth culture. The Clade 5

strains along with strains from other clades were grown to log-phase in rich media, and their

ability to form chains was assessed by visualizing division septa with the fluorescent D-amino

acid label, HADA. Chains of cells were still observed in the Clade 5 strains by time-lapse

microscopy analyses, although the chains formed in broth culture were not as long as those

observed during growth on the agarose pads (**Figure 5, S2**). Chains were also observed in 4

additional Clade 5 strains analyzed during exponential-phase growth in broth culture.

Interestingly, two of the Clade 5 strains, V48 and WU66, did not appear to incorporate the

HADA label efficiently, and the cells appeared lighter in phase-contrast images despite
undergoing the fixation process shortly after HADA addition. It is unclear why the HADA was
poorly incorporated by these strains across multiple independent experiments. Regardless, these
results indicate that many Clade 5 strains undergo cell separation less efficiently than strains
from other clades irrespective of whether they are grown on a solid surface or in broth culture.

313
Figure 5. Clade 5 strains form chains during logarithmic growth in broth culture. Representative
micrographs showing phase-contrast (top) and peptidoglycan labeling with the fluorescent D-amino
acid, HADA, (bottom) following growth in rich broth (BHIS) to logarithmic phase. Scale bar, 10 μm.
Data are representative of multiple independent experiments. **ADJ** indicates that the brightness of the
image was enhanced to detect HADA labeling in V48.

Cell length does not correlate with the propensity to sporulate.

We next wondered whether the propensity to form chains impacts the ability of Clade 5
strains to sporulate because analyses in *Bacillus subtilis* suggest that smaller cells, such as those
formed during stationary phase growth (41), may concentrate proteins involved in the
phosphorelay that induces sporulation. In *B. subtilis*, the decrease in cell length helps the kinase
KinA reach the threshold concentration needed to trigger sporulation initiation (42, 43).
Although *C. difficile* lacks homologs of KinA and other components of the phosphorelay
system, it is possible that the longer cell length observed in Clade 5 strains may similarly reduce
their ability to induce sporulation by diluting the levels of a currently unknown sporulation
regulator. To test this hypothesis, we analyzed the propensity of Clade 5 strains to form spores
when plated on 70:30 sporulation medium using phase-contrast microscopy and heat resistance
assays.
Striking differences in the frequency of spore formation were observed across the Clade
5 strains, with several strains exhibiting close to 100% sporulation levels (**Figure 6A, S3**).

Although TAL29600, which forms the longest chains of the strains tested, exhibited extremely

low levels of sporulation (0.003%), we did not observe a correlation between sporulation and

cell length among the strains (**Figure 6B, S3**). For example, M120 forms long chains yet readily

forms spores (~100% sporulation frequency). The low sporulation levels of TAL29600 were

rather surprising since this strain was isolated from a clinical setting, so it likely forms spores

more efficiently in conditions encountered during infection. Notably, when grown on 70:30

medium, Clade 5 strains did not appear to form the long chains observed during broth culture

and grown on agarose pads. In addition, the spores produced by all *C. difficile* isolates analyzed

exhibited similar lengths and proportions despite arising from vegetative cells that varied

widely in their cell length (**Figure S4**). Thus, our data suggest that Clade 5 strains alter their cell

length and propensity to form chains depending on the media encountered.

347
348
Figure 6. Sporulation levels in Clade 5 isolates grown on 70:30 medium. Phase-contrast microscopy
of the indicated strains ~24 hrs after sporulation induction. Heat resistance efficiencies, indicated below
the respective images, are based on 20-24 hr sporulating cultures and represent the mean and standard
deviation for a given strain relative to wild type based on a minimum of three biological replicates.
Statistical significance relative to strain 630 was determined using a one-way ANOVA and Tukey's test.
The scale bar represents 5 μm.
355
356

Genomic comparisons of Clade 5 strains

Since our phenotypic analyses revealed that TAL29996 differs from the 8 additional

Clade 5 strains analyzed, we sought to gain insight into the molecular basis for this difference by

comparing their genomes. To this end, we sequenced 5 of the Clade 5 strains, including

TAL29996, and determined that the average nucleotide identity (ANI) for orthologous genes

between the strains ranged from 99.83-99.99% (**Table S1**). Since the pan-genome between the 5

strains is 12%, they are quite closely related. A comparison of their pan-genomes revealed that

TAL29996 is missing one duplication of *blaR1*, which encodes an integral membrane protein

that senses beta-lactams, and a gene region predicted to be involved in nicotinate metabolism

compared to the other 4 strains. We also compared the genomes using breseq (44) to identify

SNPs that might distinguish TAL29996 from the other strains using strain M120 as the

reference, since this strain has been the Clade 5 strain traditionally characterized (30, 45).

Around 10 SNPs were unique to TAL29996, but none were clearly implicated in regulating cell

separation or peptidoglycan synthesis (**Tables S2, S3**).

Since testing whether one or some of these SNPs or absent genes help TAL29996

undergo cell separation more efficiently would require genetic manipulation of TAL29996, we

instead analyzed the orientation of the *cmr* switch, also known as the Cdi6 DNA invertible

element (31, 46). The orientation of the *cmr* switch affects expression of the adjacent *cmrRST*

operon, which encodes a non-canonical two-component system, resulting in phase variation of

CmrRST. Because CmrRST promotes cell chaining, the orientation of the *cmr* switch determines

whether *C. difficile* produces rough or smooth colony morphologies (29, 31, 46). Specifically,

cells from rough colonies are highly biased to the ON orientation of the *cmr* switch and exhibit a

chaining phenotype, whereas cells from smooth colonies are highly biased to the OFF orientation

and do not form chains (31). Consistent with the breseq analyses, manual inspection of the

cmrRST upstream region revealed that this region is identical between all the Clade 5 strains

analyzed (100% identity), including strain TAL29996. In the Illumina whole genome sequencing

performed on the subset of 5 Clade 5 strains, the *cmr* switch is in the OFF orientation in the

assembled contigs.

However, in contrast with the Illumina analyses, qPCR analyses of the *cmr* switch region

in the Clade 5 isolates revealed that TAL29996 exhibits a *cmr*-ON frequency of ~20%, whereas

the four remaining Clade 5 strains exhibit a *cmr*-ON frequency of between 70-96% on average

(**Fig 7**). Furthermore, when we extended these analyses to the four additional Clade 5 strains

characterized that also produce chains, we found that they exhibit a *cmr*-ON frequency of

between 80-93%. Collectively, these results suggest that the orientation of the *cmr* switch

orientation correlates with the propensity to form chains in Clade 5 strains and that most Clade 5

strains are biased to the *cmr*-ON orientation even during growth in broth culture.

Since the *cmr*-ON state has also been correlated with increased surface motility, we next

analyzed the surface motility of Clade 5 strains. After incubation for 5 days, *cmr*-ON strains

spread further on agar plates than the TAL29996 *cmr*-OFF strain (**Fig 7B**). Consistent with prior

reports, Clade 5 strains whose *cmr* switch is predominantly in the ON orientation exhibited

greater and more uniform surface motility. In contrast, TAL29996 exhibited less and more

asymmetric surface motility, with fractal-like extensions extending from a couple of sites. The

surface motility of TAL29996 resembles the phenotype previously reported for Clade 2 strain

R20291, whose *cmr* switch is also predominantly in the OFF position in liquid cultures.

However, growth of R20291 on agar medium selects for bacteria with the *cmr* switch in the ON

orientation, which leads to asymmetric surface motility on plates (31). These observations

suggest that even though TAL29996 is biased to the *cmr*-OFF orientation during broth culture

growth, during growth on BHIS agar, a subset of cells switch to the *cmr*-ON orientation, leading

to the asymmetric spreading phenotype.

408
409
410 **Figure 7. Orientation of the *cmr* switch correlates with chaining phenotype and surface motility in**
411 **Clade 5 strains.** A. Orientation-specific PCR for detecting the orientation of the *cmr* switch in the

412 indicated strains. B. Representative images of surface motility 5 days after exponentially growing liquid
413 cultures were spotted onto BHIS agar plates.
414
Analyses of the surface motility of strains from Clade 1, 2, and 4 revealed that

asymmetric motility was more frequently observed in Clade 2 strains. Clade 1 strains exhibited a

range of surface motility, from high surface motility with strain 630 to lower surface motility

with strain WU38 (**Figure S5**). While analyzing a larger number of Clade 1, 2, and 4 strains

would reveal whether there are clade-specific differences in surface motility, the data suggest

that Clade 2 strains may exhibit more variability in *cmr* switching within a population during

growth on plates, with strain Wup14 exhibiting the least surface motility of the Clade 2 strains

tested. Collectively, the data imply that the ON orientation of the *cmr* switch during broth culture

growth contributes to the chaining phenotype observed in most Clade 5 strains.

424
425
Discussion

While the genetically distinct nature of Clade 5 strains from Clade 1-4 strains is well-

established (10, 26), the phenotypic impacts of this genetic diversity are poorly understood,

with only one study to our knowledge systematically comparing Clade 5 strains to Clade 1-4

strains (10, 26). By phenotypically characterizing *C. difficile* strains from multiple clades using

time-lapse microscopy, we discovered that Clade 5 strains have distinct growth properties from

the Clade 1, 2, and 4 strains tested. Our analyses of multiple strains from each of Clades 1, 2, 4,

and 5 revealed that Clade 5 strains grow more quickly (**Figures 2, 4, S1**) and are more likely to

form chains than strains from Clades 1, 2, and 4, irrespective of media type or growth on a

surface (**Figures 2, 4, S2**). Notably, these analyses represent the first report of the single-cell

growth properties of *C. difficile* to our knowledge and were enabled by our development of a

simple method for conducting anaerobic time-lapse microscopy analyses (**Figure 1**) and recent

developments in automated image analyses of time-lapse microscopy data (39, 40).

Using this time-lapse microscopy method, we discovered that many Clade 5 strains

produce extremely long chains of cells (>100 μm in length, **Figures 3, 4**). This chaining

phenotype was observed in 8 of 9 Clade 5 strains versus none of the Clade 1, 2, and 4 strains,

regardless of whether the cells were grown on agarose pads or in broth culture. Thus, Clade 5

strains appear to undergo cell separation less frequently than strains from other Clades. The

exception to this conclusion is Clade 5 strain TAL29996, which did not form chains but

nevertheless grew at similarly fast rates as the other Clade 5 strains (**Figure 4**). Taken together,

these data suggest that faster growth is likely a characteristic that distinguishes Clade 5 strains

from Clade 1, 2, and 4 strains.

Our genomic analyses suggest that a driving factor of the chaining phenotype of Clade 5

strains is their propensity to have the *cmr* switch in the ON orientation (**Figure 7**), which

increases the expression of an operon encoding a signal transduction system that induces

chaining (31). This conclusion is based on our finding that the non-chaining TAL29996 strain

exhibits a *cmr*-ON orientation frequency of 20%, whereas the 8 additional Clade 5 strains tested

exhibited *cmr*-ON orientation frequencies of between ~70-95% (**Figure 7**). In addition, strain

TAL29996 exhibited less surface motility during growth on agar plates, with only a fraction of

the population spreading on the surface relative to the more uniform spreading observed for the

other 8 Clade 5 strains (**Figure 7**).

Previous work in Clade 2 strain R20291 has shown that growth on agar plates promotes

selection of *cmr*-ON bacteria, and increased c-di-GMP during surface growth may also promote

cmrRST expression (29), so Clade 5 strain TAL29996 may exhibit a similar surface-inducible

expression of the *cmrRST* operon. Indeed, the chaining phenotype of Clade 5 strains was more

frequent and extensive during time-lapse microscopy analyses on agarose pads compared to

broth culture-grown cells (**Figures 4, 5**). Thus, the chaining phenotype of Clade 5 strains may

be similarly responsive to c-di-GMP levels during growth on a surface (29). While testing this

hypothesis will depend on the ability to genetically manipulate Clade 5 strains, it is tempting to

speculate that *cmrRST* expression is an important determinant of the growth properties of Clade

5 strains. Expression of this operon in Clade 2 strain R20291 not only promotes chaining and

surface motility in a Type IV pilus-independent manner, it also negatively regulates biofilm

formation and flagellar motility (31). In addition, growth conditions favoring flagellar motility

select against *cmr*-ON bacteria. Since Clade 5 strains lack flagellar motility (12), they may be

"primed" to exhibit surface motility in the absence of selection against the *cmr*-ON state.

Regardless, analyzing the regulation of c-di-GMP in different growth conditions in Clade 5

strains, particularly for TAL29996 relative to the other strains, will likely provide insight into

the mechanisms that drive the higher incidence of *cmr*-ON cells in most Clade 5 strains.

Even though Clade 5 strains form long chains and produce longer cells than strains from

the other clades, neither cell nor chain length was predictive of the propensity to sporulate

(**Figure 6**). Clade 5 strains exhibited wildly different sporulation frequencies, with TAL29600

forming spores inefficiently (~5,000-fold less efficiently than most Clade 5 strains). While little

is known about the mechanisms regulating sporulation initiation outside of strains 630 and

R20291, future analyses of strain TAL29600, which sporulates poorly under the laboratory

media tested (**Figure 6**), could provide insight into the molecular determinants of sporulation in

Clade 5 strains. For example, differences in gene presence or polymorphisms in several c-di-

GMP-related genes were observed in TAL29600 relative to the other Clade 5 strains (**Table

S2**), and c-di-GMP has also been implicated in regulating sporulation initiation events through

an unknown mechanism (47, 48). Thus, analyzing the contribution of these genes in future work

could provide important insight into c-di-GMP-mediated regulation of sporulation.

The variable sporulation levels observed for the nine Clade 5 strains analyzed in this

manuscript contrast with the findings of a previous study, which reported that Clade 5 strains

sporulate at lower rates in the presence of glucose and fructose (26). Kumar *et al*. previously

correlated phenotypic differences in sporulation and mouse colonization between Clade 5

strains vs. Clade 1-4 strains to their finding that genes involved in sporulation and the

metabolism of simple sugars are under positive selection in Clade 1-4 strains (26). However, the

five Clade 5 strains analyzed by Kumar *et al*. all exhibited very low sporulation frequencies

under the conditions tested. While we used different conditions to induce sporulation in the

present study than those by Kumar *et al*., our analyses of nine Clade 5 strains indicate that there

are marked phenotypic differences in sporulation frequency within this clade. Thus, analyzing

more strains within a clade may reveal fewer differences between clades. A caveat to our

analyses is that *C. difficile* strains exhibit considerable genetic diversity even within a ribotype

(20, 22), so our conclusions may only reflect the behavior of RT078 Clade 5 strains. It is also

possible that analyzing more strains within this ribotype will reveal a broader spectrum of

phenotypes than the somewhat bimodal ones we have reported with respect to cell length and

chaining propensity (**Figures 2, 3**). Indeed, a recent analysis of a large number of RT027 strains

indicates that strains within this ribotype can exhibit marked differences in virulence in mice

(22) despite these strains traditionally being associated with "epidemic" outbreaks in hospital

setting.

While addressing these questions will depend on analyzing a larger number of strains

from each clade, an important outcome of our study is its development of a facile method for

conducting time-lapse microscopy under anaerobic conditions. Although this method is a closed

system, i.e. no additional reagents can be introduced once the bacteria are inoculated onto the

pad in the anaerobic chamber, it could be coupled to recently developed anaerobic imaging tags

based on fluorogen-activated proteins (49). Thus, there are many potential applications for this

simple method for studying the growth of anaerobes over time at the single-cell level.

Materials and Methods

Bacterial strains and growth conditions

All *C. difficile* strains were grown on brain heart infusion (BHIS) medium supplemented

with 0.5% w/v yeast extract and 0.1% w/v L-cysteine with taurocholate (TCA; 0.1% w/v; 1.9

mM). Strains were sub-cultured into tryptone yeast extract (TY) broth supplemented with 0.1%

w/v L-cysteine (TYC medium) prior to inoculation onto the time-lapse microscopy agarose

pads. All strains were grown at 37°C under anaerobic conditions using a gas mixture of 85%

hydrogen, 5% CO_2, and 10% H2. For time-lapse experiments, 1.5% agarose pads were used

supplemented with TYC medium as described above. Sporulation analyses were carried out on

70:30 medium (70% BHIS and 30% SMC) for 24h as described previously (50).

Anaerobic time-lapse imaging of C. difficile growth

All imaging was carried out on a Leica DMi8 inverted microscope with a HC plan

apochromat 63x 1.4 NA oil immersion phase contrast objective. Fluorescent membrane staining

experiments were done with a Lumencor Spectra X light source, coupled with an XLED-QP

quadruple-band dichroic beam-splitter (Leica) (transmission: 415, 470, 570, and 660 nm) along

with an external emission filter wheel (Leica). FM4-64 was excited using a 470nm LED

through a 470/20nm excitation filter and emitted light was filtered through 590/50nm emission

filter and captured with a Leica DFC9000GTC sCMOS camera. All experiments were carried

out at 37°C using a microscope incubation system (Pecon), Leica Adaptive Focus Control

hardware autofocus, and a high precision stage (Pecon) were used for all imaging experiments.

For time-lapse imaging of *C. difficile* growth all bacterial strains were grown in 2 mL

liquid TY medium to a turbid OD600 > 2-3; after 2 hours of growth, bacteria were diluted 1:100

for Clade 5 strains and all other strains were diluted 1:50 in fresh media and grown to mid-log

stage (OD600 0.4-0.7).

An imaging chamber with a gas-tight seal was constructed using a 125 μL Gene

Frame (Thermo Fisher) adhered to a glass slide generating a well for growth medium. The slide

was then transferred to the anaerobic chamber. In the anaerobic chamber, the gene frame was

filled with 500μl 1.5% Top vision low melting point agarose and tryptone yeast extract media

containing 0.1% w/v L-cysteine to scavenge oxygen and maintain anaerobic conditions. While

the agarose was molten, a second clean slide was placed over the top and the agar pad was

placed on a frozen small freezer block (for holding PCR strip tubes) for 10-30 minutes until the

agarose-media mixture was solid. For experiments where FM4-64 was used, agarose pads were

made the same way, with the addition of FM4-64 to a final concentration of 1 μg/mL directly to
the agarose/media solution prior to making the agar pad.
The agar pad was dried for 5-10 minutes until visible liquid on the surface of the pad
was evaporated. 1 μL of mid-log cells were spotted on the pad, dried, and a #1.5 coverslip
(VWR) was adhered to the Gene Frame. The cells were imaged at 37°C until they reached
confluency in the field of view. This was anywhere from 2.5 hours for Clade 5 strains to 6 hours
for Clades 1-4 for all experiments with images taken at 5-minute intervals.
556
Image analysis, computing hardware, and statistical analysis.
All movie frames were trimmed to the point when cells were not overlapping and out of
focus regions were cropped. The resulting images were analyzed using the Python library
DeLTA 2.0 (39, 40). All image and data analyses were done on a PC running
Windows 10 equipped with an AMD Ryzen 5900HX 8-core CPU, 32GB DDR4 RAM, 2 1TB
NVME SSDs, and an NVIDIA RTX3080 GPU with 16GB VRAM. Analysis of the output data
and data visualizations were done in Python using Matplotlib/Seaborn,
Pandas, Numpy, Scipy, and the Statannotations library.
565
Septum detection
The image processing starts with the masks generated using DeLTA 2.0. First, we performed
erosion on the mask with a disk of radius of 1 pixel to avoid effects of the membrane fluorescence.
The photo was cropped following the eroded mask contour. Hence, only the interior of the contour
was considered. Pixel values were rescaled such as the minimum pixel value of inner pixels was
mapped to 0 and the maximum is mapped to 255. We set a threshold intensity as the pixel value
of the 95% quantile of pixel intensities. The image was slightly blurred by convolution with a
Gaussian filter with standard deviation of 2 pixels. The septum corresponds to the contours found
after performing an adaptive thresholding based on thresholds defined using a Gaussian-weighted
method. This thresholding was performed using the open cv2 library in Python. The resultant
contours were manually verified from videos generated using the tracking from the outputs from
DeLTA. The manual validation mainly involved filling in the time points where the algorithm
missed a ring that was visible in the video. We did not assume the existence of a ring if the
algorithm did not detect it in a previous ,m,frame.
580
Cell length estimation
We measured the cell projected area from the DeLTA contours of the images as the pixel amount
of the contours. However, estimating the cell length was challenging because some cells were very
long and bent. To overcome this problem, we selected 30 images of three different cells (from
strains 630, TAL3050 and TAL28131) that were straight and had different lengths. For these cells,
we calculated the cell length as the longest side of the minimum bounding rectangle of the contour.
From these lengths of straight cells, we also estimated the best cell width as the average of the
projected area divided by the length. Considering the extreme cell length, the effects of the rounded
tips were negligible and the rectangle shape adequately approximated length. We verifed that this
mean value showed low variability for the three strains. Then, we used this width value to estimate
the length of all the cells, including the bent ones, by dividing their projected area by the width.
This way, we obtained a consistent measure of cell length that was independent of bending.
593
Elongation rate estimation

We tracked the cell size over time and identified the division points as the ones where the cell size
(projected area) dropped by more than 30% compared to the current cell size value. We fitted an
exponential function (with base e) of time to the data points between two divisions and estimated
the elongation rate from the exponent of the best fit. We expressed the elongation rate in
doublings/hr, which means how many times the cell size doubles in one hour. For example, an
elongation rate of 2 doubling/hr means that the cell size doubles two times in 1 hr, which
corresponds to an exponent of 2ln(2) 1/hr. For the statistics, we only included the elongation rates
that had a high quality of fit, with an R^2 coefficient greater than 0.9.
603
Bulk growth measurements. Starter cultures were grown until early stationary phase in BHIS
(or TYC medium as indicated) then back-diluted 1:50 into BHIS (or TYC medium). When the
cultures reached an OD_{600} of 0.5, they were back-diluted 1:25 into 200 µL of either BHIS or
TYC in a flat 96 well polystyrene plate (CellTreat). The OD_{600} was analyzed every 15 min for 24
hrs in a BioTek Epoch plate reader with shaking. Bulk growth measurements are based on a
minimum of three independent replicates across a minimum of 2 experiments.
610
Sporulation assays. Starter cultures were grown until early stationary phase in BHIS then back-
diluted 1:50 into BHIS. When the cultures reached an OD_{600} between 0.35 and 0.75, 120 µL of
the culture was spread onto 70:30 (70% SMC media and 30% BHIS media) agar plates (40 ml
media per plate) and then incubated for 20-24 hrs before the sporulating cells were scraped from
the plate into phosphate-buffered saline (PBS). Sporulation levels were visualized by phase-
contrast microscopy as previously described (51).
617
618 **Heat resistance assay**. Heat-resistant spore formation was measured 20-24 hrs after sporulation
619 was induced on 70:30 agar plates as previously described (50). The percent sporulation of given
620 culture represents the ratio of heat-resistant colony-forming units (CFUs) to total CFUs. Percent
621 sporulation was determined from a minimum of 3 biological replicates.
622
Spore purification. Spores were purified as previously described (52) by scraping up
sporulating cells incubated on 70:30 medium for 3 days into ice-cold H_2O. The cells were
washed several times in ice-water over the course of a day and incubated on ice overnight. The
following morning, the sample was pelleted, and cells were resuspended in 1 X DNAse buffer
(New England Biolabs) and then treated with DNAse (New England Biolabs) for 30 min at 37˚C.
The samples was washed one more time before being resuspended in 20% Histodenz and then
layered onto a 50% Histodenz layer. The resulting mixture was pelleted and the supernatant was
aspirated off using a vacuum aspirator. The pelleted spores were washed in ice-cold water 2-3
times and the optical density of the purified spores was measured.
632
Genomic DNA preparation. Starter cultures were grown until early stationary phase in BHIS
then back-diluted 1:50 into BHIS and grown until an OD_{600} of around 0.7-0.8 was reached. 10
mL of the culture was pelleted and then frozen at -80˚C. After thawing the sample, it was
resuspended in a 25% sucrose TE buffer (10 mM Tris, 1mM EDTA), incubated with 100 mg/mL
lysozyme for 37˚C for 1 hr. After the cultures tarted to lyse, proteinase K, RNAse A, EDTA,
Sarkosyl, and NaCl was added. Phenol:Chloroform:IAA (25:24:1) was added to extract proteins,
gently mixed, and then the sample was pelleted to separate the phenol and aqueous layer. The
aqueous layer was then added to Chloroform:IAA (24:1), mixed gently, then centrifuged. The
aqueous layer was then precipitated using isopropanol and incubated at -20˚C for a minimum of

15 min. The precipitated DNA was pelleted and then washed with 70% ethanol. The pellet was
air dried and then gently resuspended in 10 mM Tris pH 8.0 elution buffer.
644
Genomic analyses. Genomic DNA was sequenced by MiGS at the University of Pittsburgh
(now SeqCenter) according to their standard protocol. Libraries were sequenced on an Illumina
NextSeq 500 platform to generate paired-end 150 bp reads. Illumina reads of RT078 genomes
were assembled into contigs using SPAdes (v3.13.0),(53) and genes were called and annotated
using Prokka (v1.11).(54) Assembled and annotated contigs of five RT078 strains (TAL28131,
TAL29600, TAL29996, TAL30550, TAL30574) were applied for pangenomic analysis. Default
settings were used based on the Anvi'o workflow for microbial pangenomcis with adjustments
for minbit as 0.5 and mcl-inflation as 10.(55-57) For SNPs analyses, reads of five RT078
genomes were aligned to the reference M120 and variants were called by breseq (v. 0.38.1) by
default settings.(58)
655
656   **qPCR analyses**. Each genomic DNA sample was analyzed by qPCR with primers that amplify
657   the *cmr*-ON sequence orientation, the *cmr*-OFF orientation, or the reference gene *rpoA* (31) .
658   Each 20-µL qPCR reaction consisted of 100 ng genomic DNA, 100 nM primers, and SensiMix™
659   SYBR® reagents (Bioline). The reactions were run on a LightCycler® 96 (Roche Diagnostics),
660   and *cmr* switch orientation frequencies were calculated as described previously (29).
661
662   **Surface motility assays**
663   Starter cultures were grown until early stationary phase in BHIS then back-diluted 1:50 into
664   BHIS and grown until an OD_{600} of 0.5 was reached. 10 µL of the exponential-phase cultures
665   were then spotted onto BHIS plates and incubated at 37˚C for 5 days after which the plates were
666   scanned using a flatbed scanner.
667
Author contribution statement
J.W.R. conducted all live-cell microscopy experiments and analyzed the single-cell data.
C.A.N.A. generated data visualizations from the time-lapse microscopy data, developed the
method for identifying septa within chains that had not initiated cell separation, and calculated
elongation rates and cell lengths. I.N. and A.S. conducted sporulation analyses and liquid
culture experiments. J.W.R., R.T., and A.S conceptualized all experiments with input from .
Q.D. and A.S extracted genomic DNA and visualized cells at different growth phases. Q.D.
conducted all the genomic analyses and genome comparisons. N.D. and I.N. conducted spore
purification analyses and imaging. B.A. and IE developed and optimized methods for
constructing agarose pads that could support anaerobic growth in an ambient atmosphere.
M.J.D. developed the DeLTA 2.0 analysis pipeline. A.M. conducted qPCRs to determine *cmr*
switch orientation in Clade 5 isolates. Writing was by J.W.R and A.S with input from all the
authors.
681
682
683   **Table 1. *Clostridioides difficile* clinical isolates used in this study**
684

Strain Name	Clade	Ribotype	ST group	Source	Reference
630	1	012	2	Zurich, 1982 (Sanger Institute)	(59)
R20291	2	027	1	London, 2006 (Sanger Institute)	(60)

M68	4	017	81	Dublin, 2006 (Sanger Institute)	(45)
M120	5	078	11	UK, 2007 (Sanger Institute)	(45)
TAL28131	5	078	11	Tufts Medical Center	This paper
TAL29600	5	078	11	Tufts Medical Center	This paper
TAL29996	5	078	11	Tufts Medical Center	This paper
TAL30550	5	078	11	Tufts Medical Center	This paper
TAL30574	5	078	11	Tufts Medical Center	This paper
V48	5	078	11	Brigham & Women's Hospital	This paper
139b	5	078	11	Memorial Sloan Kettering	(20)
WU66	5	078	11	Barnes-Jewish Hospital	(20)
BBL2	1	012	2	Memorial Sloan Kettering	(20)
WU38	1	012	2	Barnes-Jewish Hospital	(20)
190B	1	087	46	Memorial Sloan Kettering	(20)
Wup14	2	027	1	Barnes-Jewish Hospital	(20)
BBL4	2	027	1	Memorial Sloan Kettering	(20)
186A	2	027	1	Memorial Sloan Kettering	(20)

685
686 Memorial Sloan Kettering Cancer Center (MSK), Barnes-Jewish Hospital (BJH)
687

References

1. Desai K, Gupta SB, Dubberke ER, Prabhu VS, Browne C, Mast TC. 2016.
Epidemiological and economic burden of Clostridium difficile in the United States:
estimates from a modeling approach. BMC Infect Dis 16:303.
2. Guh AY, Mu Y, Winston LG, Johnston H, Olson D, Farley MM, Wilson LE, Holzbauer
SM, Phipps EC, Dumyati GK, Beldavs ZG, Kainer MA, Karlsson M, Gerding DN,
McDonald LC, Emerging Infections Program Clostridioides difficile Infection Working
G. 2020. Trends in U.S. Burden of Clostridioides difficile Infection and Outcomes. N
Engl J Med 382:1320-1330.
3. Shen A. 2020. Clostridioides difficile Spore Formation and Germination: New Insights
and Opportunities for Intervention. Annu Rev Microbiol 74:545-566.
4. Abt MC, McKenney PT, Pamer EG. 2016. Clostridium difficile colitis: pathogenesis and
host defence. Nat Rev Microbiol doi:10.1038/nrmicro.2016.108.
5. Theriot CM, Young VB. 2015. Interactions Between the Gastrointestinal Microbiome
and Clostridium difficile. Annu Rev Microbiol 69:445-61.
6. Kordus SL, Thomas AK, Lacy DB. 2022. Clostridioides difficile toxins: mechanisms of
action and antitoxin therapeutics. Nat Rev Microbiol 20:285-298.
7. Maroo S, Lamont J. 2006. Recurrent Clostridium difficile. Gastroenterology 130:1311-
1316.
8. Zanella Terrier MC, Simonet ML, Bichard P, Frossard JL. 2014. Recurrent Clostridium
difficile infections: the importance of the intestinal microbiota. World J Gastroenterol
20:7416-23.
9. Knight DR, Elliott B, Chang BJ, Perkins TT, Riley TV. 2015. Diversity and Evolution in
the Genome of Clostridium difficile. Clin Microbiol Rev 28:721-41.

10.   Knight DR, Imwattana K, Kullin B, Guerrero-Araya E, Paredes-Sabja D, Didelot X,
714   Dingle KE, Eyre DW, Rodriguez C, Riley TV. 2021. Major genetic discontinuity and
715   novel toxigenic species in Clostridioides difficile taxonomy. Elife 10.

716   11.   Norsigian CJ, Danhof HA, Brand CK, Midani FS, Broddrick JT, Savidge TC, Britton
717   RA, Palsson BO, Spinler JK, Monk JM. 2022. Systems biology approach to functionally
718   assess the Clostridioides difficile pangenome reveals genetic diversity with
719   discriminatory power. Proc Natl Acad Sci U S A 119:e2119396119.

720   12.   Knight DR, Kullin B, Androga GO, Barbut F, Eckert C, Johnson S, Spigaglia P, Tateda
721   K, Tsai PJ, Riley TV. 2019. Evolutionary and Genomic Insights into Clostridioides
722   difficile Sequence Type 11: a Diverse Zoonotic and Antimicrobial-Resistant Lineage of
723   Global One Health Importance. mBio 10.

724   13.   Seth-Smith HMB, Biggel M, Roloff T, Hinic V, Bodmer T, Risch M, Casanova C,
725   Widmer A, Sommerstein R, Marschall J, Tschudin-Sutter S, Egli A. 2021. Transition
726   From PCR-Ribotyping to Whole Genome Sequencing Based Typing of Clostridioides
727   difficile. Front Cell Infect Microbiol 11:681518.

728   14.   Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput
729   ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun
730   9:5114.

731   15.   Dingle KE, Elliott B, Robinson E, Griffiths D, Eyre DW, Stoesser N, Vaughan A,
732   Golubchik T, Fawley WN, Wilcox MH, Peto TE, Walker AS, Riley TV, Crook DW,
733   Didelot X. 2014. Evolutionary history of the Clostridium difficile pathogenicity locus.
734   Genome Biol Evol 6:36-52.

735   16.   Mengoli M, Barone M, Fabbrini M, D'Amico F, Brigidi P, Turroni S. 2022. Make It Less
736   difficile: Understanding Genetic Evolution and Global Spread of Clostridioides difficile.
737   Genes (Basel) 13.

738   17.   Endres BT, Begum K, Sun H, Walk ST, Memariani A, Lancaster C, Gonzales-Luna AJ,
739   Dotson KM, Basseres E, Offiong C, Tupy S, Kuper K, Septimus E, Arafat R, Alam MJ,
740   Zhao Z, Hurdle JG, Savidge TC, Garey KW. 2019. Epidemic Clostridioides difficile
741   Ribotype 027 Lineages: Comparisons of Texas Versus Worldwide Strains. Open Forum
742   Infect Dis 6:ofz013.

743   18.   McDonald LC, Killgore GE, Thompson A, Owens RC, Jr., Kazakova SV, Sambol SP,
744   Johnson S, Gerding DN. 2005. An epidemic, toxin gene-variant strain of Clostridium
745   difficile. N Engl J Med 353:2433-41.

746   19.   Clements AC, Magalhaes RJ, Tatem AJ, Paterson DL, Riley TV. 2010. Clostridium
747   difficile PCR ribotype 027: assessing the risks of further worldwide spread. Lancet Infect
748   Dis 10:395-404.

749   20.   Lewis BB, Carter RA, Ling L, Leiner I, Taur Y, Kamboj M, Dubberke ER, Xavier J,
750   Pamer EG. 2017. Pathogenicity Locus, Core Genome, and Accessory Gene Contributions
751   to Clostridium difficile Virulence. mBio 8.

752   21.   Merrigan M, Venugopal A, Mallozzi M, Roxas B, Viswanathan VK, Johnson S, Gerding
753   DN, Vedantam G. 2010. Human hypervirulent Clostridium difficile strains exhibit
754   increased sporulation as well as robust toxin production. J Bacteriol 192:4904-11.

755   22.   Dong Q, Lin H, Allen MM, Garneau JR, Sia JK, Smith RC, Haro F, McMillen T, Pope
756   RL, Metcalfe C, Burgo V, Woodson C, Dylla N, Kohout C, Sundararajan A, Snitkin ES,
757   Young VB, Fortier LC, Kamboj M, Pamer EG. 2023. Virulence and genomic diversity

among clinical isolates of ST1 (BI/NAP1/027) Clostridioides difficile. Cell Rep
42:112861.

23.   Shaw HA, Preston MD, Vendrik KEW, Cairns MD, Browne HP, Stabler RA, Crobach
MJT, Corver J, Pituch H, Ingebretsen A, Pirmohamed M, Faulds-Pain A, Valiente E,
Lawley TD, Fairweather NF, Kuijper EJ, Wren BW. 2020. The recent emergence of a
highly related virulent Clostridium difficile clade with unique characteristics. Clin
Microbiol Infect 26:492-498.

24.   Li C, Harmanus C, Zhu D, Meng X, Wang S, Duan J, Liu S, Fu C, Zhou P, Liu R, Wu A,
Kuijper EJ, Smits WK, Fu L, Sun X. 2018. Characterization of the virulence of a non-
RT027, non-RT078 and binary toxin-positive Clostridium difficile strain associated with
severe diarrhea. Emerg Microbes Infect 7:211.

25.   Imwattana K, Rodriguez C, Riley TV, Knight DR. 2021. A species-wide genetic atlas of
antimicrobial resistance in Clostridioides difficile. Microb Genom 7.

26.   Kumar N, Browne HP, Viciani E, Forster SC, Clare S, Harcourt K, Stares MD, Dougan
G, Fairley DJ, Roberts P, Pirmohamed M, Clokie MRJ, Jensen MBF, Hargreaves KR, Ip
M, Wieler LH, Seyboldt C, Noren T, Riley TV, Kuijper EJ, Wren BW, Lawley TD. 2019.
Adaptation of host transmission cycle during Clostridium difficile speciation. Nat Genet
51:1315-1320.

27.   Knetsch CW, Kumar N, Forster SC, Connor TR, Browne HP, Harmanus C, Sanders IM,
Harris SR, Turner L, Morris T, Perry M, Miyajima F, Roberts P, Pirmohamed M, Songer
JG, Weese JS, Indra A, Corver J, Rupnik M, Wren BW, Riley TV, Kuijper EJ, Lawley
TD. 2018. Zoonotic Transfer of Clostridium difficile Harboring Antimicrobial Resistance
between Farm Animals and Humans. J Clin Microbiol 56.

28.   Lim SC, Knight DR, Riley TV. 2020. Clostridium difficile and One Health. Clin
Microbiol Infect 26:857-863.

29.   Garrett EM, Mehra A, Sekulovic O, Tamayo R. 2021. Multiple Regulatory Mechanisms
Control the Production of CmrRST, an Atypical Signal Transduction System in
Clostridioides difficile. mBio 13:e0296921.

30.   Thanissery R, Winston JA, Theriot CM. 2017. Inhibition of spore germination, growth,
and toxin activity of clinically relevant C. difficile strains by gut microbiota derived
secondary bile acids. Anaerobe 45:86-100.

31.   Garrett EM, Sekulovic O, Wetzel D, Jones JB, Edwards AN, Vargas-Cuebas G, McBride
SM, Tamayo R. 2019. Phase variation of a signal transduction system controls
Clostridioides difficile colony morphology, motility, and virulence. PLoS Biol
17:e3000379.

32.   Ackermann M. 2015. A functional perspective on phenotypic heterogeneity in
microorganisms. Nat Rev Microbiol 13:497-508.

33.   Aldridge BB, Fernandez-Suarez M, Heller D, Ambravaneswaran V, Irimia D, Toner M,
Fortune SM. 2012. Asymmetry and aging of mycobacterial cells lead to variable growth
and antibiotic susceptibility. Science 335:100-4.

34.   Anjuwon-Foster BR, Tamayo R. 2017. A genetic switch controls the production of
flagella and toxins in Clostridium difficile. PLoS Genet 13:e1006701.

35.   Anjuwon-Foster BR, Tamayo R. 2018. Phase variation of Clostridium difficile virulence
factors. Gut Microbes 9:76-83.

36.   Kint N, Janoir C, Monot M, Hoys S, Soutourina O, Dupuy B, Martin-Verstraete I. 2017.
The alternative sigma factor sigma(B) plays a crucial role in adaptive strategies of
Clostridium difficile during gut infection. Environ Microbiol 19:1933-1958.
37.   Courson DS, Pokhrel A, Scott C, Madrill M, Rinehold AJ, Tamayo R, Cheney RE,
Purcell EB. 2019. Single cell analysis of nutrient regulation of Clostridioides
(Clostridium) difficile motility. Anaerobe 59:205-211.
38.   de Jong IG, Beilharz K, Kuipers OP, Veening JW. 2011. Live Cell Imaging of Bacillus
subtilis and Streptococcus pneumoniae using Automated Time-lapse Microscopy. J Vis
Exp doi:10.3791/3145.
39.   Lugagne JB, Lin H, Dunlop MJ. 2020. DeLTA: Automated cell segmentation, tracking,
and lineage reconstruction using deep learning. PLoS Comput Biol 16:e1007673.
40.   O'Connor OM, Alnahhas RN, Lugagne JB, Dunlop MJ. 2022. DeLTA 2.0: A deep
learning pipeline for quantifying single-cell spatial and temporal dynamics. PLoS
Comput Biol 18:e1009797.
41.   Sauls JT, Cox SE, Do Q, Castillo V, Ghulam-Jelani Z, Jun S. 2019. Control of Bacillus
subtilis Replication Initiation during Physiological Transitions and Perturbations. mBio
10.
42.   Narula J, Kuchina A, Zhang F, Fujita M, Suel GM, Igoshin OA. 2016. Slowdown of
growth controls cellular differentiation. Mol Syst Biol 12:871.
43.   Eswaramoorthy P, Duan D, Dinh J, Dravis A, Devi SN, Fujita M. 2010. The threshold
level of the sensor histidine kinase KinA governs entry into sporulation in Bacillus
subtilis. J Bacteriol 192:3870-82.
44.   Deatherage DE, Traverse CC, Wolf LN, Barrick JE. 2014. Detecting rare structural
variation in evolving microbial populations from new sequence junctions using breseq.
Front Genet 5:468.
45.   He M, Sebaihia M, Lawley T, Stabler R, Dawson L, Martin M, Holt K, Seth-Smith H,
Quail M, Rance R, Brooks K, Churcher C, Harris D, Bentley S, Burrows C, Clark L,
Corton C, Murray V, Rose G, Thurston S, van Tonder A, Walker D, Wren B, Dougan G,
Parkhill J. 2010. Evolutionary dynamics of Clostridium difficile over short and long time
scales. Proceedings of the National Academy of Sciences of the United States of America
107:7527-7532.
46.   Sekulovic O, Mathias Garrett E, Bourgeois J, Tamayo R, Shen A, Camilli A. 2018.
Genome-wide detection of conservative site-specific recombination in bacteria. PLoS
Genet 14:e1007332.
47.   Dhungel BA, Govind R. 2021. Phase-variable expression of pdcB, a phosphodiesterase,
influences sporulation in Clostridioides difficile. Mol Microbiol 116:1347-1360.
48.   Edwards AN, Willams CL, Pareek N, McBride SM, Tamayo R. 2021. c-di-GMP Inhibits
Early Sporulation in Clostridioides difficile. mSphere 6:e0091921.
49.   Streett H, Charubin K, Papoutsakis ET. 2021. Anaerobic fluorescent reporters for cell
identification, microbial cell biology and high-throughput screening of microbiota and
genomic libraries. Curr Opin Biotechnol 71:151-163.
50.   Shen A, Fimlaid KA, Pishdadian K. 2016. Inducing and Quantifying Clostridium difficile
Spore Formation. Methods Mol Biol 1476:129-42.
51.   Pishdadian K, Fimlaid KA, Shen A. 2015. SpoIIID-mediated regulation of sigma(K)
function during Clostridium difficile sporulation. Mol Microbiol 95:189-208.

52. Fimlaid KA, Jensen O, Donnelly ML, Francis MB, Sorg JA, Shen A. 2015. Identification
of a Novel Lipoprotein Regulator of Clostridium difficile Spore Germination. PLoS
Pathogens doi:10.1371/journal.ppat.1005239.
53. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes
De Novo Assembler. Curr Protoc Bioinformatics 70:e102.
54. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics
30:2068-9.
55. Benedict MN, Henriksen JR, Metcalf WW, Whitaker RJ, Price ND. 2014. ITEP: an
integrated toolkit for exploration of microbial pan-genomes. BMC Genomics 15:8.
56. Eren AM, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, Fink I, Pan JN, Yousef
M, Fogarty EC, Trigodet F, Watson AR, Esen OC, Moore RM, Clayssen Q, Lee MD,
Kivenson V, Graham ED, Merrill BD, Karkman A, Blankenberg D, Eppley JM, Sjodin
A, Scott JJ, Vazquez-Campos X, McKay LJ, McDaniel EA, Stevens SLR, Anderson RE,
Fuessel J, Fernandez-Guerra A, Maignien L, Delmont TO, Willis AD. 2021. Community-
led, integrated, reproducible multi-omics with anvi'o. Nat Microbiol 6:3-6.
57. van Dongen S, Abreu-Goodger C. 2012. Using MCL to extract clusters from networks.
Methods Mol Biol 804:281-95.
58. Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using breseq. Methods Mol Biol
1151:165-88.
59. Sebaihia M, Wren B, Mullany P, Fairweather N, Minton N, Stabler R, Thomson N,
Roberts A, Cerdeño-Târraga A, Wang H, Holden M, Wright A, Churcher C, Quail M,
Baker S, Bason N, Brooks K, Chillingworth T, Cronin A, Davis P, Dowd L, Fraser A,
Feltwell T, Hance Z, Holroyd S, Jagels K, Moule S, Mungall K, Price C, Rabbinowitsch
E, Sharp S, Simmonds M, Stevens K, Unwin L, Whithead S, Dupuy B, Dougan G,
Barrell B, Parkhill J. 2006. The multidrug-resistant human pathogen Clostridium difficile
has a highly mobile, mosaic genome. Nature genetics 38:779-786.
60. Stabler R, He M, Dawson L, Martin M, Valiente E, Corton C, Lawley T, Sebaihia M,
Quail M, Rose G, Gerding D, Gibert M, Popoff M, Parkhill J, Dougan G, Wren B. 2009.
Comparative genome and phenotypic analysis of Clostridium difficile 027 strains
provides insight into the evolution of a hypervirulent bacterium. Genome biology 10.
878