1

Unique growth and morphology properties of Clade 5 *Clostridioides difficile* strains

revealed by single-cell time-lapse microscopy

John W. Ribis[1,2*‡], César Nieto[3*], Nicholas V. DiBenedetto[1,2], Anchal Mehra,[4] Pola Kuhn[1,2],

Qiwen Dong[1], Irene Nagawa,[1] Imane El Meouche,[5] Bree B. Aldridge,[1,6] Mary J. Dunlop,[7] Rita

Tamayo,[4] Abhyudai Singh,[3] Aimee Shen[1†]

Affiliations

[1]Department of Molecular Biology and Microbiology, Tufts University School of Medicine,

Boston, Massachusetts, USA, [2] Graduate School of Biomedical Sciences, Tufts University

School of Medicine, Boston, Massachusetts, USA. [3]Department of Electrical and Computer

Engineering, University of Delaware, Newark, DE 19716, USA. [4]Department of Microbiology

and Immunology, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill,

NC, USA, [5]INSERM, Université Paris Cité, Université Sorbonne Paris Nord, Inserm, IAME, F-

75018, Paris, France, [6]Department of Biomedical Engineering, Tufts University School of

Engineering, Medford, Massachusetts, USA, [7]Biomedical Engineering, Boston University,

Boston, MA 02215.

*These authors contributed equally to this work.

‡Current address: Nikon Instruments

†Address correspondence to
Aimee Shen, aimee.shen@tufts.edu
Phone number: (617)636-3792

Abstract

Clostridioides difficile is a gastrointestinal pathogen of both humans and agricultural animals and

thus a major One Health threat. The *C. difficile* species consists of five main clades, with Clade 5

currently undergoing speciation from Clades 1-4. Clade 5 strains are highly prevalent in

agricultural animals and can cause zoonotic infections, suggesting that these strains have evolved

phenotypes that distinguish them from Clade 1-4 strains. Here, we compare the growth

properties of Clade 5 strains to those of Clade 1-4 strains using anaerobic time-lapse microscopy

coupled with automated image analysis. Our analyses indicate that Clade 5 strains grow faster

and are more likely to form long chains of cells than Clade 1-4 strains. Using comparative

genomic and CRISPRi analyses, we show that the chaining phenotype of Clade 5 strains is

driven by the orientation of the invertible *cmr* switch sequence, with chaining strains exhibiting a

bias to the *cmr*-ON state. Interestingly, Clade 5 strains with a bias towards the *cmr*-ON state

shifted to a largely *cmr*-OFF state during murine infection, suggesting that the *cmr*-OFF state is

under positive selection during infection. Collectively, our data reveal that Clade 5 strains have

distinct growth properties, which may allow them to inhabit diverse ecological niches.

Author Summary

The Clade 5 strains of the *Clostridioides difficile* species are so phylogenetically divergent that they almost meet the threshold of being a distinct species. Although these strains are ubiquitously isolated from agricultural and environmental settings and an important source of zoonotic and community-acquired infections, it is unclear whether they have distinct phenotypic properties that allow them to colonize diverse hosts or persist in the environment. By combining a novel anaerobic time-lapse microscopy method with automated image analysis, we discovered that Clade 5 strains grow faster than strains from other *C. difficile* clades and that they frequently form long chains. These chaining properties are driven by the environmentally responsive expression of a non-canonical signal transduction system, which our analyses suggest is selected against during murine infection. Collectively, our analyses reveal that Clade 5 strains have distinct growth properties that may promote their persistence in the environment.

Introduction

Clostridioides difficile is a leading cause of nosocomial infections in the United States, with approximately 500,000 new infections and 14,000 deaths being attributed to this organism annually [1, 2]. As an obligate anaerobe, C. difficile relies on its hardy, metabolically dormant spore form to survive outside the host and transmit disease to new hosts [3]. When C. difficile spores are ingested by susceptible hosts [4, 5], they germinate and outgrow into vegetative cells that subsequently colonize the colon. The vegetative cells secrete toxins that damage gut epithelial tissue [6], which triggers an inflammatory response that can cause disease pathologies ranging from mild diarrhea to pseudomembranous colitis and even death [2, 4]. C. difficile also causes recurrent infections in ~20% of infections, which can lead to more severe disease symptoms [2, 7, 8].

C. difficile's success as a pathogen may be related to its tremendous genetic diversity [9-11], with genomic analyses indicating that C. difficile's core genome represents only ~10-20% of its pan-genome [9, 10]. The plasticity of its "open" pan-genome likely helps C. difficile colonize the gastrointestinal tract of diverse animals, from mammals to invertebrates, and persist in environmental reservoirs like sewage and compost [12]. Indeed, the C. difficile species is so genetically diverse that it has been divided into five distinct phylogenetic clades based on multi-locus sequence typing (MLST) analyses, and the clades have been further subdivided into different ribotypes (RTs) or sequence types (STs) [10, 13].

Between these five clades, there are notable differences in geographic and host distributions. Clade 1 is the largest, most heterogeneous clade with the broadest geographic distribution [14, 15]. It includes over 200 STs, which can contain both toxin-producing and non-toxigenic strains, including the well-characterized, genetically tractable toxigenic strain 630 [9]. Clade 2 harbors epidemic-associated strains found within the ribotype 027 (RT027) lineage. Strains from this ribotype have been associated with outbreaks in hospitals, particularly in North America, due to their frequent resistance to fluoroquinolones [16-18]. These epidemic strains can also cause severe disease symptoms in part due to their production of three toxins: TcdA, TcdB, and CDT (binary toxin) [6]. Although RT027 strains have frequently been associated with "hypervirulence," there is considerable phenotypic diversity within this lineage with respect to virulence, toxin production levels, flagellar motility, and sporulation [19-22].

Clade 3 strains are relatively uncommon and harbor a unique cell surface due to their lack of

CwpV production [23], but phenotypic analyses of biofilm formation and motility suggest that

they share similarities with Clade 2 strains like R20291 [24]. Clade 4 contains RT017 (ST37)

strains that only encode a single toxin, TcdB (i.e., $TcdA^-CDT^-$), and are often clindamycin- and

fluoroquinolone-resistant. RT017 strains have been associated with outbreaks in Europe and

North America and are the most common strains found in Asia [25]. Clade 5 is the most

genetically distant from the other 4 main *C. difficile* clades and is thought to have emerged

before Clades 1-4 [26]. While the average nucleotide identity (ANI) for Clades 1-4 ranges

between 97.1 - 99.8%, Clade 5 strains exhibit ANI values around 96%, which is close to the

ANI value demarcation used by NCBI to define organisms of the same species [10, 27]. Thus,

Clade 5 strains appear to be actively diverging from Clades 1-4 [10].

Clade 5 strains are an increasing problem in healthcare and agricultural settings because

they can cause severe disease in humans and are commonly found in livestock, particularly pigs

[12, 28]. While other *C. difficile* strains have been known to infect both humans and animals,

only Clade 5 strains have been associated with zoonotic transmission from both animal-to-

human and human-to-animal [28, 29]. The mechanisms underlying this bidirectional zoonotic

transmission are poorly understood, but the increased carriage of antimicrobial resistance genes

by Clade 5 strains may contribute to their ability to persist in agricultural and community

settings [28, 30]. Thus, Clade 5 strains are of particular relevance from a One Health

perspective [12, 31], especially since they frequently cause community-acquired infections [30]

and are often detected in retail foods [32]. These observations highlight the importance of

understanding the unique properties of this group of strains. Indeed, a recent genomic analysis

suggests that RT078/ST11 strains within Clade 5 frequently carry zinc acquisition and

homeostasis genes [11].

Despite numerous genomic analyses revealing the remarkable genetic diversity of *C.*

difficile strains, relatively few studies have investigated the phenotypic diversity between strains

from different clades. Clade-specific differences in colony morphology between Clade 5 strains

relative to Clade 1-4 strains have been described in a limited set of analyses [26, 33], suggesting

that differences in growth and/or cellular morphology may exist within clades. While

differences in bulk growth rates between *C. difficile* strains have been reported [34], most

phenotypic analyses have been conducted on a limited subset of strains within a given clade.

Furthermore, systematic comparisons of the growth properties of different clades have only

recently been described [35], while comparisons of their cell morphology have not been

performed to date.

Here, we compare the growth properties of multiple strains derived from all five

phylogenetic clades of *C. difficile* using anaerobic time-lapse microscopy. These analyses

unexpectedly reveal striking differences in the growth and cell morphology of the Clade 5

lineage relative to strains from Clades 1-4. Specifically, we found that Clade 5 strains grow

faster and frequently form long chains, in contrast with strains from Clade 1-4 strains. Genomic

comparisons and genetic analyses indicate that the chaining phenotype of Clade 5 strains is

driven by the phase-variable expression of the *cmrRST* operon by the invertible *cmr* switch [33].

Interestingly, we found that Clade 5 strains with a strong *cmr*-ON bias mostly reverted to a *cmr*-

OFF phenotype during murine infection. Taken together, our data reveal that Clade 5 strains

have unique growth properties relative to Clade 1-4 strains that may contribute to the

widespread distribution of Clade 5 strain among diverse animal hosts.

Results

Development of a simple method for time-lapse imaging under anaerobic conditions

Time-lapse imaging of single cells has been widely used to study phenotypic

heterogeneity in bacteria, which can impact important traits like antibiotic resistance and

virulence [33, 36-39]. However, live single-cell analyses in *C. difficile* have been complicated by

its inability to grow in the presence of atmospheric oxygen [40]. While time-lapse microscopy

analyses of *C. difficile* have previously been reported, they require custom growth chambers to

maintain anaerobic conditions [41] and thus may limit the accessibility of these experimental

systems to a broad range of investigators.

To overcome these limitations, we established a simple system that relies solely upon

commercially available reagents and materials to grow *C. difficile* cells under anaerobic

conditions. This system uses gas-tight, adhesive Gene Frames, which have been used extensively

in imaging applications for bacteria [42]. Notably, the gas-impermeability of these commercial

seals allows anaerobic conditions to be maintained when agarose pads made with growth media

are prepared in the anaerobic chamber (**Figure 1**). Gene Frames also generate thick agarose pads,

which are critical for *C. difficile* to grow in a sealed system under ambient conditions. After

agarose pads are prepared in the anaerobic chamber, *C. difficile* cultures are inoculated onto the

pads, and the pads are sealed with a coverslip. The growth chamber is then removed from the

chamber and imaged on a heated microscope stage under ambient conditions for up to 6 hours or

until *C. difficile* stops growing as a monolayer.

Time-lapse microscopy reveals clade-specific differences in elongation rate and cell length

Having established an anaerobic time-lapse imaging setup, we compared the single-cell

growth properties of representative *C. difficile* strains from Clade 1 (630, ribotype (RT) 012),

Clade 2 (R20291, RT027), Clade 3 (E15, RT075), Clade 4 (M68, RT017), and Clade 5 (M120,

RT078) (**Figure 2, Table 1**). The five "representative" strains were all isolated from patients

with *C. difficile*-associated disease and are frequently used as reference genomes for their

clades and ribotype groups. Notably, RT027 (ST1), RT017 (ST45), and RT078 (ST11) strains

are from ribotypes/ multi-locus sequencing types that are frequently isolated from patients with

C. difficile infection (CDI) [12, 16, 19, 25]. In contrast, Clade 3 strains are more rare and the

least characterized of *C. difficile* strains [23].

The growth properties of single cells visualized by time-lapse microscopy were

quantified using Deep Learning for Time-lapse Analysis (DeLTA) software, which rapidly and

accurately segments and tracks bacteria growing in two dimensions on agarose pads [43, 44].

This software uses deep convolutional neural networks to analyze time-lapse microscopy

images, allowing the growth properties of individual cells growing in microcolonies on agarose

pads to be determined. The segmentation and tracking of *C. difficile* cells were highly accurate

(**Figure 1**), and minimal user input or post-image processing was needed to obtain growth

property measurements.

Robust growth was observed for all strains using our system. Growth was quantified by

measuring the elongation rate, which was defined as doublings/hr to indicate the number of

times that a cell's length doubles in one hour. The elongation rate (doublings/hr) is distinct from

the doubling time, or generation time, which represents the length of *time* that it takes before a

bacterium divides. Instead, the elongation rate reflects how fast the cell is increasing in length

over time. Notably, the Clade 5 strain M120 elongated the fastest (2.1 doublings/hr, p < 0.05),

followed by Clade 3 E15 (2.0 doublings/hr) and Clade 2 strain R20291 (1.8 doublings/hr), and

then Clade 1 strain 630 and Clade 4 strain M68 (1.6-1.7 doublings/hr) (**Figure 2A**).

Importantly, the differences in single-cell elongation rates measured for the five strains

were also observed in bulk population analyses of their growth using optical density in TYC

and BHIS media (**Figure 2B** & **S1**). These analyses confirmed that the Clade 5 strain M120

grew faster (based on optical density-based analyses) in these media than the Clade 1-4 strains

(p < 0.001). In contrast, negligible differences in bulk growth rates were observed between

Clade 1-4 strains in BHIS media, although, in TYC medium, the Clade 3 strain grew faster than

Clades 1, 2, and 4 strains (**Figures 2B** & **S1**).

The Clade 5 strain M120 exhibited another distinct growth property from the Clade 1-4

strains. While strains from Clades 1-4 produced cells of similar length prior to cell division,

with an average apparent length of ~13 µm, cells of Clade 5 strain M120 were significantly

longer, with an average apparent length of ~22 µm. Indeed, cells ~50 µm were readily observed

for the Clade 5 strain M120 (**Figure 2C**), and these cells appeared to bend readily (**Figure 2D**).

By incorporating the FM4-64 membrane stain into the agarose pads to visualize division septa

[45], we assessed whether the Clade 5 strain M120 forms chains vs. filaments, These analyses

revealed that septa were readily observed in Clade 5 strain M120 across the length of a given

cell (**Figure 3, inset**), Since the spacing between division septa was relatively consistent, the

Clade 5 strain M120 appear to undergo cell separation less efficiently than strains from the

other clades tested. Indeed, cell separation in strain M120 was so inefficient that it was often

necessary to stitch together several fields of view to fully visualize M120 chains, which

approached several hundred microns and even up to ~1 mm in length (**Figure 3**).

Clade 5 clinical isolates typically form long chains and grow more quickly than strains

from other clades

Since prior work indicated that Clade 5 strains produce colony morphologies distinct from

Clade 1, 2, and 4 strains [26], we sought to determine whether the striking cell chaining

phenotype and faster growth rate observed in Clade 5 strain M120 were properties shared by

other Clade 5 strains. Thus, we compared the single-cell growth properties of five additional

Clade 5 clinical isolates obtained from several hospitals around the country on TYC agarose

(**Table 1**) using time-lapse microscopy analyses (**Figure 2**). Clade 1 strain 630 was included as

a control since it does not form chains in any of the conditions we have tested.

These analyses revealed that all but one of the Clade 5 strains tested formed long chains,

with TAL29600 forming the longest chains (29 µm on average, **Figure 4A-C**). In contrast,

strain TAL29996 formed shorter chains that were comparable in length to those observed for

strains from Clades 1-4 (12-13 µm, **Figures 2B** & **4C**). Notably, DeLTA segmented many

Clade 5 chains as single cells because cell separation (i.e., invagination) had not yet initiated at

division septa visualized via FM4-64 staining. To overcome this limitation and accurately

quantify cell length within long chains, namely the distance between division septa, we

modified our image processing pipeline to use a thresholding method to detect division septa.

After generating masks in DeLTA to segment the chains, we modified the mask so that only the

interior of the contour was analyzed. We then applied an adaptive thresholding method to

identify division septa based on their elevated fluorescence relative to the long axis of the cell;

the thresholds were defined using a Gaussian-weighted method (**Figure 4A**, red dots, **Figure**

4B). Septa detected with this automated method were also manually inspected. These analyses

revealed that the average length of cells within the chains of cells made by Clade 5 strains is

only slightly longer than the average length of cells made by strains that do not form chains,

namely Clade 1 strain 630 and Clade 5 strain TAL29996 (**Figure 4C**, **Table 2**). For example,

the Clade 5 strains that produced ”he l’ngest chains (TAL29600 and TAL30550) had average

cell lengths of 14 and 12 µm, respectively, which is only 30-50% longer than the non-chaining

strains 630 (Clade 1, 8.5 µm) and TAL29996 (Clade 5, 9.4 µm) (**Table 2**). Notably, even

though the Clade 5 strain TAL29996 did not form chains, it still exhibited higher elongation

rates, which were similar to those measured for other Clade 5 strains (2.1 doublings/hr vs. 1.7

doublings/hr for Clade 1 strain 630**, Figure 4D**).

Importantly, the additional Clade 5 strains tested also grew faster in bulk optical density-

based analyses in broth culture than Clade 1-4 strains irrespective of their ability to form chains

(**Figure S2**). To assess whether these findings would extend to additional Clade 5 strains vs.

Clade 1-4 strains, we analyzed the growth of additional strains from all five clades in broth

culture. These analyses confirmed that Clade 1-4 strains grow at similar rates in BHIS media,

which are slower than those observed for the nine Clade 5 strains analyzed in this media

(**Figure S2**). However, Clade 3 strains grew relatively faster than the Clade 1 630 strain in TYC

medium, but their growth was still slower than the Clade 5 strain M120 (**Figure S2**). Taken

together, these analyses strongly suggest that Clade 5 strains grow faster than Clade 1-4 strains

and are more likely to form chains, presumably because their cell separation mechanisms are

less efficient.

Cell chaining in Clade 5 strains is not dependent on growth on a solid medium.

C. difficile has previously been shown to promote cell elongation and chain formation

upon induction of the *cmrRST* locus [33], which encodes a non-canonical signal transduction

system. Expression of this locus is also responsive to c-di-GMP levels [46], which increases in

cells grown on solid surfaces such as in a biofilm or on an agar plate [46]. To test whether the

chaining phenotype observed in Clade 5 strains is induced by growth on a surface, we assessed

the chaining properties of Clade 5 strains during logarithmic growth in rich medium broth

culture using the fluorescent D-amino acid label, HADA, to stain septa. These analyses revealed

that Clade 5 strains still form chains during broth culture growth, although the chains are not as

long as those observed during growth on the agarose pads (**Figures 5 and S3**).

Since Clade 5 strains grow rapidly in rich media, we considered the possibility that the

chaining phenotype might be mitigated by allowing more time for cell separation to occur after

cell division. To test this possibility, we grew Clade 5 strains in CDDM minimal medium and

analyzed their chaining properties [47]. While Clade 5 strains grew slower in CDDM medium

relative to richer media (compare **Figure S4A** to **Figures 2 & S2**), the Clade 5 strains

nevertheless formed chains in minimal medium (**Figure S4B**), with the exception of strain

TAL29996 strain. Taken together, our results reveal that Clade 5 strains undergo cell separation

less efficiently in a range of growth conditions relative to strains from other clades.

Cell length does not correlate with the propensity to sporulate.

We next wondered whether the propensity to form chains impacts the ability of Clade 5

strains to sporulate. Analyses in *Bacillus subtilis* suggest that smaller cells, such as those

formed during stationary phase growth [48], are more likely to sporulate likely because they

concentrate proteins involved in the phosphorelay that induces sporulation [49, 50]. For

example, a decrease in cell length in *B. subtilis* helps the kinase KinA reach the threshold

concentration needed to trigger sporulation initiation. Although *C. difficile* lacks homologs of

KinA and other components of the phosphorelay system [51], the longer cells generated by

Clade 5 strains may be less likely to induce sporulation due to dilution of a currently unknown

sporulation regulator. To test this hypothesis, we analyzed the propensity of Clade 5 strains to

form spores when plated on 70:30 sporulation medium using phase-contrast microscopy and

heat resistance assays. These analyses revealed that Clade 5 strains exhibit striking differences

in sporulation frequency, with some strains exhibiting close to 100% sporulation levels, and

others exhibiting levels closer to 20% (**Figures 6 & S5**). Interestingly, strain TAL29600

exhibited extremely low levels of sporulation (0.003%), and it continued to form chains during

growth on 70:30 medium. In contrast, the other Clade 5 strains analyzed did not form long

chains when grown on 70:30 sporulation medium, and the spores produced by these isolates

exhibited similar lengths and proportions relative to the Clade 1 strain 630 (**Figure S6**). These

data suggest that Clade 5 strains alter their cell length and propensity to form chains depending

on the growth conditions encountered.

Consistent with this hypothesis, even though strain M120 forms long chains during

growth in or on rich medium (**Figures 3** & **S3**), it readily formed spores during growth on 70:30

medium (~100% sporulation frequency, **Figure S5**). Notably, the average cell length for visibly

sporulating Clade 5 cells was ~5-6 µm (**Figure S8**) compared to the average ~11 µm cell length

measured for the Clade 5 strains in rich broth culture. This reduced cell length was observed

even for Clade 5 strain TAL29600 (**Figure S8**), which sporulates poorly on 70:30 medium,

indicating that there was little correlation between cell length and propensity to sporulate for the

Clade 5 strains analyzed.

Comparative genomics reveals that the chaining phenotype of Clade 5 strains is driven by

***cmrRST* operon expression.**

Given the phenotypic difference in chaining observed for the TAL29996 strain relative to

the 8 other Clade 5 strains analyzed, we sought to gain insight into the mechanism driving this

difference by comparing the genomes of five of the Clade 5 strains, including TAL29996. These

analyses revealed that the average nucleotide identity (ANI) for orthologous genes ranged

between 99.83-99.99% (**Table S1**) and that the pan-genome between the five strains is 12%.

Thus, all five strains are quite closely related. The pan-genome analysis revealed that, relative to

the other four strains, TAL29996 is missing one duplication of *blaR1*, which encodes an integral

membrane protein that senses beta-lactams, and a gene region predicted to be involved in

nicotinate metabolism. To identify SNPs that might distinguish TAL29996 from the other strains,

we used breseq [52]; the Clade 5 strain M120 genome sequence was used as the reference

genome because it is the Clade 5 strain traditionally characterized [34, 53]. These analyses

identified 10 SNPs that were unique to TAL29996, but none were obviously involved in

regulating cell separation or peptidoglycan synthesis (**Table S2**).

We next took a candidate approach to gain insight into why TAL29996 mediates cell

separation more efficiently than the other Clade 5 strains by analyzing the orientation of the *cmr*

switch, also known as the Cdi6 DNA invertible element [33, 54]. This phase-variable element

affects the expression of the adjacent *cmrRST* operon, which encodes a non-canonical CmrRST

signal transduction system that regulates cell chaining and colony morphology [33, 46, 54]. Cells

from rough colonies form chains and are highly biased to the ON orientation of the *cmr* switch,

whereas cells from smooth colonies do not form chains and are highly biased to the OFF

orientation [33]. Although the sequence of the *cmr* switch is identical between the Clade 5 strains

analyzed, including strain TAL29996, we found that the orientation of the *cmr* switch during

growth in broth culture was markedly different for Clade 5 strain TAL29996. Specifically, qPCR

analyses revealed that Clade 5 strains that form chains are biased towards the *cmr*-ON

orientation (between 70-96% ON) (**Figure 7A**), whereas the *cmr*-ON orientation was markedly

less frequent in the non-chaining strain TAL29996 (~20%, **Figure 7**).

Since these analyses correlated the *cmr*-ON switch orientation to the chaining phenotype

of Clade 5 strains, we tested whether knocking down the expression of the *cmrRST* operon using

CRISPRi in strains that are biased towards the *cmr*-ON state would reduce their chaining

properties. A plasmid targeting the *cmrR* gene using CRISPRi was introduced into the Clade 5

strains TAL28131, TAL30550, and TAL30574, which typically produce chains and are found

predominantly in the *cmr*-ON state in rich broth culture (70-96%). Knocking down the

expression of the *cmrRST* operon in all three strain backgrounds reduced their chaining

phenotypes under these conditions (**Figures 7B** & **S8**), indicating that the expression of the

cmrRST operon in Clade 5 strains drives their propensity to form chains.

Surface motility in Clade 5 strains relative to strains of other clades

342   Since the *cmr*-ON state has also been correlated with increased surface motility [33], we

analyzed the surface motility of our Clade 5 strains. Consistent with prior reports [33], the

primarily *cmr*-ON state strains exhibited greater and more uniform surface motility (**Figure 7C**),

whereas the predominantly *cmr*-OFF state TAL29996 strain exhibited less and more asymmetric

surface motility, with fractal-like extensions emerging from only a few sites. This asymmetric

phenotype has previously been reported for the Clade 2 strain R20291, whose *cmr* switch is

predominantly in the OFF position in liquid cultures but converts to the ON orientation during

growth on plates [33]. These observations suggest that, even though TAL29996 is biased to the

cmr-OFF orientation during broth culture growth, a subset of TAL29996 cells switch to the *cmr*-

ON orientation during growth on BHIS agar, leading to the asymmetric spreading phenotype.

352   We next assessed whether additional Clade 1-4 strains exhibit surface motility. These

analyses revealed that asymmetric motility was more frequently observed in Clade 2 strains,

although Clade 2 strain Wup14 exhibited little surface motility (**Figure S9**). Clade 1 strains

exhibited a range of surface motility, from high surface motility with strain 630 to lower surface

motility with strain WU38 (**Figure S9**). While the data suggest that *cmr* switching in Clade 2

strains varies between strains during growth on agar medium, Clade 5 strains biased towards the

cmr-ON state are more likely to exhibit surface motility. However, it is important to note that

additional factors contribute to surface motility beyond expression of the *cmrRST* operon [33],

since loss of pili can also decrease surface motility on plates [55]. Regardless, the data imply that

the *cmr*-ON state promotes surface motility in Clade 5 strains.

Colonization and virulence properties of Clade 5 strains

365   Beyond the effects of the CmrRST system on cell chaining and surface motility, this

system has also been shown to impact the virulence of the Clade 2 strain, R20291, in a hamster

model of infection, with loss of *cmrR* reducing R20291's ability to cause disease and the *cmr*-

OFF orientation correlating with less severe disease in hamsters [33]. Since chaining in *Bacillus*

anthracis strains promotes virulence [56], while chaining in *Enterococcus faecalis* promotes

colonization [57], we compared the ability of Clade 5 strains to colonize mice and cause disease.

Mice were infected with 10^5 spores of several Clade 5 strains and the Clade 1 strain 630 and the

weight loss induced by these strains, their colonization levels, and orientation of the *cmr* switch

over the course of the 14-day infection were assessed. For this latter analysis, we focused on

strains TAL29600 and TAL29996 because they exhibited the highest and lowest *cmr*ON

orientations, respectively, during growth in rich media (**Figure 7A**).

All Clade 5 strains tested colonized to relatively similar levels throughout the 14 days of

the infection. Strain 630 also colonized mice to similar levels in the first two days of infection

and then maintained colonization, albeit at 1-2 logs lower than the Clade 5 strains (**Figure 8A**).

Interestingly, only the Clade 5 strain TAL29600 caused significant weight loss relative to the

other strains on Days 2 through 4, although the Clade 1 strain 630 caused some weight loss on

Day 3 (**Figure 8B**). This latter phenotype is consistent with prior reports of strain 630 causing

only mild disease symptoms in the cefoperazone model of murine infection [58, 59]. Analyses of

the *cmr* orientation revealed that the *cmr*-OFF orientation appeared to be selected for over the

course of the infection. While the TAL29600 spore inoculum started off with ~30% *cmr*-ON

frequency, the frequency of TAL29600 cells detected in the *cmr*-ON orientation decreased

rapidly to < 5% *cmr*-ON by 24 hrs post-inoculation (**Figure 8C**). As the infection progressed,

two of the 8 mice tested exhibited an increase in TAL29600 cells with the *cmr*-ON orientation

(6-30% *cmr*-ON) (**Figure S10**). Conversely, the TAL29996 strain retained the ~1% *cmr*-ON

frequency of the inoculum for the greater part of the 14-day infection (**Figure 8C**). Taken

together, these analyses reveal that Clade 5 RT078 strains efficiently colonize mice but vary in

their ability to cause disease. Furthermore, the ability of the Clade 5 strains to colonize or cause

disease did not strongly correlate with their ability to form chains in broth culture.

Discussion

While Clade 5 strains are genetically distinct [10, 26] and more prevalent in animals

than Clade 1-4 strains [12, 29, 30], the phenotypes that distinguish Clade 5 strains from Clade

1-4 strains are not well understood. By phenotypically characterizing *C. difficile* strains from

multiple clades using time-lapse microscopy, we discovered that Clade 5 strains have distinct growth properties from Clade 1-4 strains. Specifically, Clade 5 strains elongate more quickly (**Figures 2, 4, S1**) and form long chains more readily than strains from Clades 1-4, irrespective of media type or growth on a surface (**Figures 2, 4, S2**, and **S4**). In contrast, long chains were not observed in any of the Clade 1-4 strains, regardless of whether the cells were grown on agarose pads or in broth culture. Thus, Clade 5 strains undergo cell separation far less efficiently than strains from other clades during growth in rich media. While our analyses of Clade 5 strains were largely limited to the RT078 ribotype, we note that a prior report described an RT126 strain isolated from a patient experiencing multiple recurrences that also formed long chains [60]. Since the RT126 ribotype is closely related to the RT078 ribotype [12], it is likely that the chaining phenotype will be observed in other Clade 5 ribotypes.

Our data indicate that the chaining phenotype of Clade 5 strains relates to their propensity to express the *cmrRST* operon [33, 46] because they favor the *cmr*-ON state (**Figure 7**). The non-chaining TAL29996 strain has a *cmr*-ON orientation frequency of 20%, whereas the 8 other Clade 5 strains have a strong bias for the *cmr*-ON orientation (~70-96%, **Figure 7**). The chaining phenotype of Clade 5 strains prone to the *cmr*-ON state was enhanced during growth on a solid surface compared to broth culture (**Figures 4, 5**), consistent with prior analyses of the Clade 2 strain R20291, which exhibits greater *cmrRST* expression during growth on agar plates due to elevated c-di-GMP levels [46]. In addition, Clade 5 strains exhibit high levels of surface motility on agar plates, which is a *cmrRST*-induced phenotype in strain R20291 [33]. Finally, knocking-down the expression of the *cmrRST* operon in three Clade 5 strains biased to the *cmr*-ON state reduced their propensity to form chains (**Figures 7** & **S8**).

These findings lead to the question of what benefit *cmrRST* expression might confer to Clade 5 strains. In the Clade 2 R20291 strain, *cmrRST* expression is negatively correlated with flagellar motility, and growth conditions that favor flagellar motility select against the *cmr*-ON state [33]. Since Clade 5 strains lack flagellar motility [12], it is tempting to hypothesize that they are "primed" to form chains as a method to promote motility. Analyzing the regulation of c-di-GMP in different growth conditions in Clade 5 strains, particularly for TAL29996 relative to the other strains, will likely provide insight into the mechanisms that drive *cmrRST* expression, chaining, and surface motility in Clade 5 strains and the importance of these properties to these strains.

While the propensity of Clade 5 strains to form chains in a *cmrRST*-dependent manner

might be expected to promote colonization or disease in mice based on studies of other Gram-

positive pathogens [56, 57], we found that the frequency of *cmr*-ON cells decreases during the

first 4 days of infection for strain TAL29600 (**Figure 8**). This suggests that there may be a

selection against *cmr*-ON cells during the initial stages of murine colonization. Consistent with

this hypothesis, a decrease in *cmr*-ON orientation was observed for Clade 2 strain R20291

during infection of hamsters [33]. However, since the frequency of *cmr*-ON TAL29600 cells

increased in some mice at later stages of colonization (**Figure 8**), our data imply that the

invertibility of the *cmrRST* switch region may promote *C. difficile*'s ability to adapt to different

growth conditions. While it is possible that high levels of c-di-GMP during infection induce the

expression of the *cmrRST* operon expression during murine infection [46], assessing whether

Clade 5 strains form long chains during murine infection, for example, using fluorescence *in*

situ hybridization [61, 62] or using transcriptional reporters to visualize the expression of the

cmrRST operon at the single-cell level during infection would provide insight into these

questions.

Indeed, the avirulence of most Clade 5 strains analyzed during murine infection (**Figure**

8) was somewhat surprising given that all of these strains were isolated from human patients

experiencing CDI-related disease symptoms (**Table 1**). To our knowledge, very few Clade 5

strains have been analyzed during murine infection, with one study observing minor disease

symptoms in mice for two Clade 5 strains three days post-infection, despite one of the strains

causing severe disease in humans [20]. Interestingly, RT078 Clade 5 strains frequently cause

asymptomatic infections in agricultural animals [32, 63] and mice may be important vectors of

transmission in these settings [64, 65]. Thus, it is possible that Clade 5 strains are adapted for

colonization rather than virulence in non-human systems. Consistent with this hypothesis, we

found that the Clade 5 strains persist at high levels in the murine gut over time compared to the

Clade 1 strain 630 (**Figure 8**, $p < 0.005$). Identifying factors that allow Clade 5 strains to grow

more quickly would provide insight into whether their faster growth rate promotes their

persistence in mice. It is also possible that increasing the inoculum could have allowed for a

greater degree of disease severity to be observed and differences in virulence between the

strains to be detected.

Interestingly, the chain length of Clade 5 strains did not correlate with their propensity

to sporulate (**Figure 6**). The Clade 5 strains tested varied markedly in their sporulation

frequencies, with most strains forming spores at frequencies >30% unlike strain TAL29600,

which sporulates ~5,000-fold less efficiently than the other Clade 5 strains (**Figure 6**). While

little is known about the mechanisms regulating sporulation initiation outside of strains 630 and

R20291, analyses of strain TAL29600, which sporulates poorly under laboratory conditions

(**Figure 6**), could provide insight into the molecular determinants of sporulation initiation in

Clade 5 strains. For example, differences in gene presence or polymorphisms in several c-di-

GMP-related genes were observed in TAL29600 relative to the other Clade 5 strains (**Table S2**),

and c-di-GMP has been implicated in regulating sporulation initiation events through unknown

mechanisms [66, 67].

Our time-lapse microscopy analyses further revealed that a notable delay between cell

division and cell separation is common to *C. difficile* strains, irrespective of their propensity to

form long chains, because chains of two cells were frequently observed for Clades 1-4 and

Clade 5 strain TAL29996 (**Figures S3** & **4**). Notably, these two-cell chains were typically

segmented by DeLTA as a single cell because cell separation had not initiated, i.e. no

invagination was detected (**Figure 4**, red dot). The loose coordination between cell division and

cell separation in *C. difficile* relative to other bacteria likely relates to the absence of FtsEX

homologs in *C. difficile*. In diverse bacteria, the FtsEX complex couples septal PG synthesis

with PG hydrolases that mediate cell separation to result in a fast splitting of recently divided

cells [68-71]. While it remains unclear whether coordination between cell division and cell

separation exists in *C. difficile*, recent work has identified novel factors that control chaining in

C. difficile. The CwlA peptidoglycan hydrolase mediates cell separation in *C. difficile* [72], and

its export and thus activity is controlled by the Ser/Thr kinase PrkC [72]. The septum-localizing

MldA or MldB proteins also promote chaining in *C. difficile* through unknown mechanisms

[73], so future work could address whether the inefficient cell separation phenotype of Clade 5

strains is due to decreased CwlA export or MldA/MldB levels.

Importantly, these insights into the basic physiology of *C. difficile* were enabled by our

development of a facile method for conducting time-lapse microscopy under anaerobic

conditions. Since the growth chamber set-up involves commercially available GeneFrames and

open-source software for conducting automated image analyses of time-lapse microscopy data

[43, 44], the methods described in this manuscript can be applied to many anaerobic systems for

studying the growth properties of diverse organisms and the impact of different growth

conditions and mutant backgrounds on these properties. Our anaerobic set-up could be further

coupled with recently developed, fluorogen-activated anaerobic imaging tags [74] to facilitate

single-cell analyses of gene-specific transcription during anaerobic growth and dynamic protein

localization studies [75]. Thus, there are many potential applications for the simple methods

described here for studying the growth of anaerobes over time at the single-cell level.

Materials and Methods

Bacterial strains and growth conditions

All *C. difficile* strains were grown on brain heart infusion (BHIS) medium supplemented with 0.5% w/v yeast extract and 0.1% w/v L-cysteine with taurocholate (TCA; 0.1% w/v; 1.9 mM). Strains were sub-cultured into tryptone yeast extract (TY) broth supplemented with 0.1% w/v L-cysteine (TYC medium) prior to inoculation onto the time-lapse microscopy agarose pads. All strains were grown at 37°C under anaerobic conditions using a gas mixture of 85% hydrogen, 5% CO_2, and 10% H_2. For time-lapse experiments, 1.5% agarose pads supplemented with TYC medium were used as described above. Sporulation analyses were carried out on 70:30 medium (70% BHIS and 30% SMC) for 24h as described previously [76].

Anaerobic time-lapse imaging of C. difficile growth

All imaging was carried out on a Leica DMi8 inverted microscope with a HC plan apochromat 63x 1.4 NA oil immersion phase contrast objective. Fluorescent membrane staining experiments were done with a Lumencor Spectra X light source, coupled with an XLED-QP quadruple-band dichroic beam-splitter (Leica) (transmission: 415, 470, 570, and 660□nm) along with an external emission filter wheel (Leica). FM4-64 was excited using a 470nm LED through a 470/20nm excitation filter and emitted light was filtered through a 590/50nm emission filter and captured with a Leica DFC9000GTC sCMOS camera. All experiments were carried out at 37°C using a microscope incubation system (Pecon), Leica Adaptive Focus Control hardware autofocus, and a high precision stage (Pecon) were used for all imaging experiments.

For time-lapse imaging of *C. difficile* growth, all bacterial strains were grown in 2 mL liquid TY medium to a turbid OD600 > 2-3; after 2 hours of growth, bacteria were diluted 1:100 for Clade 5 strains and all other strains were diluted 1:50 in fresh media and grown to mid-log phase (OD600 0.4-0.7).

An imaging chamber with a gas-tight seal was constructed using a 125 μL Gene Frame (Thermo Fisher) adhered to a glass slide generating a well for growth medium. The slide was then transferred to the anaerobic chamber. In the anaerobic chamber, the gene frame was filled with 500μl 1.5% Top vision low melting point agarose and tryptone yeast extract media containing 0.1% w/v L-cysteine to scavenge oxygen and maintain anaerobic conditions. While the agarose was molten, a second clean slide was placed over the top and the agar pad was placed on a frozen small freezer block (for holding PCR strip tubes) for 10-30 minutes until the agarose-media mixture was solid. For experiments using FM4-64, agarose pads were made the same way, with the addition of FM4-64 to a final concentration of 1 μg/mL directly to the agarose/media solution prior to making the agar pad.

The agar pad was dried for 5-10 minutes until visible liquid on the surface of the pad was evaporated. 1 μL of mid-log cells were spotted on the pad, dried, and a #1.5 coverslip (VWR) was adhered to the Gene Frame. The cells were imaged at 37°C until they reached confluency in the field of view. This was anywhere from 2.5 hours for Clade 5 strains to 6 hours for Clades 1-4 for all experiments with images taken at 5-minute intervals.

Image analysis, computing hardware, and statistical analysis.

All movie frames were trimmed to the point when cells were not overlapping and out of focus regions were cropped. The resulting images were analyzed using the Python library

547 DeLTA 2.0 [43, 44]. All image and data analyses were done on a PC running Windows 10
548 equipped with an AMD Ryzen 5900HX 8-core CPU, 32GB DDR4 RAM, 2 1TB NVME SSDs,
549 and an NVIDIA RTX3080 GPU with 16GB VRAM. Analysis of the output data and data
550 visualizations were done in Python using Matplotlib/Seaborn, Pandas, Numpy, Scipy, and the
551 Statannotations library.
552
Septum detection during live-cell microscopy
The image processing starts with the masks generated using DeLTA. First, we performed erosion
on the mask with a disk of radius of 1 pixel to avoid effects of membrane fluorescence. The
photo was cropped following the eroded mask contour. Hence, only the interior of the contour
was considered. Pixel values were rescaled such that the minimum pixel value of inner pixels
was mapped to 0 and the maximum was mapped to 255. We set a threshold intensity as the pixel
value of the 95% quantile of pixel intensities. The image was slightly blurred by convolution
with a Gaussian filter with standard deviation of 2 pixels. The septum corresponds to the
contours found after performing an adaptive thresholding based on thresholds defined using a
Gaussian-weighted method. This thresholding was performed using the open cv2 library in
Python. The resultant contours were manually verified from videos generated using the tracking
from the outputs from DeLTA. The manual validation mainly involved filling in the time points
where the algorithm missed a ring that was visible in the video. We did not assume the existence
of a ring if the algorithm did not detect it in a previous frame.
567
Cell length estimation
We measured the cell projected area from the DeLTA contours of the images as the pixel amount
of the contours. However, estimating cell length was challenging because some cells were very
long and bent. To overcome this problem, we selected 30 images of three different cells (from
strains 630, TAL3050 and TAL28131) that were straight and had different lengths. For these
cells, we calculated the cell length as the longest side of the minimum bounding rectangle of the
contour. From these lengths of straight cells, we also estimated the best cell width as the average
of the projected area divided by the length. Considering the extreme cell length, the effects of the
rounded tips were negligible and the rectangle shape adequately approximated length. We
verified that this mean value showed low variability for the three strains. Then, we used this
width value to estimate the length of all the cells, including the bent ones, by dividing their
projected area by the width. This way, we obtained a consistent measure of cell length that was
independent of bending.
581
Elongation rate estimation
We tracked cell size over time and identified the division points as the ones where the cell size
(projected area) dropped by more than 30% compared to the current cell size value. We fitted an
exponential function (with base e) of time to the data points between two divisions and estimated
the elongation rate from the exponent of the best fit. We expressed the elongation rate in
doublings/hr, which means how many times the cell size doubles in one hour. For example, an
elongation rate of 2 doubling/hr means that the cell size doubles two times in 1 hr, which
corresponds to an exponent of 2ln(2) 1/hr. For the statistics, we only included the elongation
rates that had a high quality of fit, with an R^2 coefficient greater than 0.9.
591
Bulk growth measurements

Starter cultures were grown until early stationary phase in BHIS (or TYC medium as indicated)
then diluted 1:50 into BHIS (or TYC medium). For the CDDM growth analyses, starter cultures
were prepared in CDDM medium at a relatively high density and then back-diluted 1:25 into
CDDM. When the cultures (for all three media conditions) reached an OD_{600} of 0.5, they were
diluted 1:50 into 200 µL of either BHIS, TYC, or CDDM in a flat 96 well polystyrene plate
(CellTreat). The OD_{600} was analyzed every 15 min for 24 hrs in a BioTek Epoch plate reader
with shaking. Bulk growth measurements are based on a minimum of three independent
replicates across a minimum of 2 experiments. The growth rate was calculated from the linear
range of the growth curves, between 105 min to 180 min of growth.

Cell wall labeling
HADA (Tocris Bioscience) was added to exponentially growing cell culture to a final
concentration of 50-100 µM and incubated for ~2 mins before cell fixation. Cells were fixed as
previously described [77]. Briefly, 500 µL of cell suspension was added to 120 µL of a 5X
fixation solution containing paraformaldehyde and $NaPO_4$ buffer. Samples were mixed and
incubated in the dark for 30 min at room temperature, followed by 30 min on ice. Fixed cells
were washed three times in phosphate-buffered saline (PBS) and resuspended in ~50 µL of PBS.
Cells were imaged within 72 hours after fixation.

Sporulation assays.
Starter cultures were grown until early stationary phase in BHIS then diluted 1:50 into BHIS.
When the cultures reached an OD_{600} between 0.35 and 0.75, 120 µL of the culture was spread
onto 70:30 (70% SMC media and 30% BHIS media) agar plates (40 ml media per plate) and
then incubated for 20-24 hrs before the sporulating cells were scraped from the plate into
phosphate-buffered saline (PBS). Sporulation levels were visualized by phase-contrast
microscopy as previously described [78].

Heat resistance assay.
Heat-resistant spore formation was measured 20-24 hrs after sporulation was induced on 70:30
agar plates as previously described [76]. The percent sporulation of given culture represents the
ratio of heat-resistant colony-forming units (CFUs) to total CFUs. Percent sporulation was
determined from a minimum of 3 biological replicates.

Spore purification.
Spores were purified as previously described [79] by scraping up sporulating cells incubated on
70:30 medium for 3 days into ice-cold H_2O. The cells were washed several times in ice-water
over the course of a day and incubated on ice overnight. The following morning, the sample was
pelleted, and cells were suspended in 1 X DNAse buffer (New England Biolabs) and then treated
with DNAse (New England Biolabs) for 30 min at 37˚C. The samples were washed one more
time before being resuspended in 20% Histodenz and then layered onto a 50% Histodenz layer.
The resulting mixture was pelleted, and the supernatant was aspirated off using a vacuum
aspirator. The pelleted spores were washed in ice-cold water 2-3 times and the optical density of
the purified spores was measured.

Genomic DNA preparation.
Starter cultures were grown until early stationary phase in BHIS then back-diluted 1:50 into
BHIS and grown until an OD_{600} of around 0.7-0.8 was reached. 10 mL of the culture was

pelleted and then frozen at -80˚C. After thawing the sample, it was resuspended in a 25%
sucrose TE buffer (10 mM Tris, 1mM EDTA), incubated with 100 mg/mL lysozyme for 37˚C
for 1 hr. After the cultures tarted to lyse, proteinase K, RNAse A, EDTA, Sarkosyl, and NaCl
was added. Phenol:Chloroform:IAA (25:24:1) was added to extract proteins, gently mixed, and
then the sample was pelleted to separate the phenol and aqueous layer. The aqueous layer was
then added to Chloroform:IAA (24:1), mixed gently, then centrifuged. The aqueous layer was
then precipitated using isopropanol and incubated at -20˚C for a minimum of 15 min. The
precipitated DNA was pelleted and then washed with 70% ethanol. The pellet was air dried and
then gently resuspended in 10 mM Tris pH 8.0 elution buffer.
649
Genomic analyses.
Genomic DNA was sequenced by MiGS at the University of Pittsburgh (now SeqCenter)
according to their standard protocol. Libraries were sequenced on an Illumina NextSeq 500
platform to generate paired-end 150 bp reads. Illumina reads of RT078 genomes were assembled
into contigs using SPAdes (v3.13.0),[80] and genes were called and annotated using Prokka
(v1.11) [81]. Assembled and annotated contigs of five RT078 strains (TAL28131, TAL29600,
TAL29996, TAL30550, TAL30574) were applied for pangenomic analysis. Default settings
were used based on the Anvi'o workflow for microbial pangenomcis with adjustments for minbit
as 0.5 and mcl-inflation as 10[82-84]. For SNPs analyses, reads of five RT078 genomes were
aligned to the reference M120 and variants were called by breseq (v. 0.38.1) by default
settings.[85]
661
qPCR analyses.
Each genomic DNA sample was analyzed by qPCR with primers that amplify the *cmr*-ON
sequence orientation, the *cmr*-OFF orientation, or the reference gene *rpoA* [33] . Each 20-µL
qPCR reaction consisted of 100 ng genomic DNA, 100 nM primers, and SensiMix™ SYBR
reagents (Bioline). The reactions were run on a LightCycler® 96 (Roche Diagnostics), and *cmr*
switch orientation frequencies were calculated as described previously [46].
668
669   *Surface motility assays*
670   Starter cultures were grown until early stationary phase in BHIS then back-diluted 1:50 into
671   BHIS and grown until an OD_{600} of 0.5 was reached. 10 µL of the exponential-phase cultures
672   were then spotted onto BHIS plates and incubated at 37˚C for 5 days after which the plates were
673   scanned using a flatbed scanner.
674
Mouse infection experiments
Mouse experiments were performed under the guidance of veterinary staff within the Tufts
Comparative Medicine Services (TCMS) core. All animal studies were done with prior approval
from the Tufts Institutional Animal Care and Use Committee (IACUC protocol #B2024-30).
Conventional 7-week-old C57BL/6 female mice from Jackson Laboratories were housed in a
sterile (autoclaved cage and bedding) large cage (24"x17") with autoclaved water and irradiated
food (Teklad 2918) for 10 days to allow for normalization of microbiota across mice through
coprophagy. After the 10-day normalization period, mice were started on cefoperazone, which
was added to their water at a concentration of 0.5mg/ml. Mice were allowed to drink the
cefoperazone water ad libitum for 10 days, after which they were placed back on sterile water
without antibiotic. After a 2-day period of being on normal sterile water, mice were weighed and

given a single dose of clindamycin (10mg/kg) via intraperitoneal injection. Immediately after IP injection, mice were moved to standard-size autoclaved mouse cages (4 mice per cage) with sterile food and water. 24 hours following the clindamycin injection, mice were inoculated with 1 x 10^5 spores of *C. difficile* (in 1xPBS) via oral gavage using a metal, reusable needle. Mice were weighed by being placed in a plastic Nalgene cup on top of a scale. Fecal pellets were collected just prior to oral gavage to ensure no prior *C. difficile* colonization and to note a baseline weight. Following inoculation, mice were weighed to monitor % weight change over time on days 1-4, 7, and 14. Fecal pellets were collected in duplicate on the same days for *C. difficile* CFU enumeration and qPCR for detection of *cmr* switch orientation. On day 14, a terminal weight was taken, and a fecal pellet was collected, followed by sacrifice via CO_2 inhalation with cervical dislocation as secondary method of euthanasia. Two experimental replicates were completed using 4 mice per group, resulting in a total of 8 mice per condition.

C. difficile CFU enumeration from mouse fecal pellets
C. difficile engraftment in mice was monitored over a 14-day period. Fecal pellets were collected from mice on days 0-4, 7, and 14. Each pellet was weighed and then suspended in 1x PBS. 10 µL of the suspension was then serially diluted 1:10 in 1x PBS in a 96 well plate, and 5 µL was spotted onto TCCFA agar to select for *C. difficile*, such that the dilutions plated were 2 x 10^3 – 2 x 10^7. *C. difficile* colonies were counted 24 hours after plating to allow for sufficient growth. *C. difficile* CFUs were normalized by gram of fecal material.

Data accessibility statement
The data set and data processing scripts are publicly available at https://doi.org/10.5281/zenodo.13352469 [86]. Genomes are available from NCBI under the BioProject ID PRJNA1238863.

Acknowledgements
We are grateful to Dr. Cheleste Thorpe, Dr. Eric Pamer, Dr. Bruno Dupuy, Dr. Lynn Bry, and Dr. Trevor Lawley for providing the strains used in this study.

Funding
This work was supported by NIAID grants R21 AI153853 (A Shen, BBA, and MJD), R01 AI122232 (A Shen), R01 AI143638 (RT), and T32 AI007422 (NVD) and NIGMS grants R35 GM148351 (A Singh), and T32 GM007310 (JWR). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

723 **Table 1.** *Clostridioides difficile* **clinical isolates used in this study**
724

Strain Name	Clade	Ribotype	ST group	Source	Reference
630	1	012	2	Zurich, 1982 (Sanger Institute)	[87]
BBL2	1	012	2	Memorial Sloan Kettering	[20]
WU38	1	012	2	Barnes-Jewish Hospital	[20]
190B	1	087	46	Memorial Sloan Kettering	[20]
R20291	2	027	1	London, 2006 (Sanger Institute)	[88]
Wup14	2	027	1	Barnes-Jewish Hospital	[20]
BBL4	2	027	1	Memorial Sloan Kettering	[20]
186A	2	027	1	Memorial Sloan Kettering	[20]
E15	3	075	unknown	France (Bruno Dupuy/ Lynn Bry)	[89, 90]
BI1	3	unknown	unknown	Bruno Dupuy/ Lynn Bry	This study
95-978	3	unknown	unknown	Bruno Dupuy/ Lynn Bry	This study
M68	4	017	81	Dublin, 2006 (Sanger Institute)	[53]
1002	4	unknown	39	Memorial Sloan Kettering	[20]
M120	5	078	11	UK, 2007 (Sanger Institute)	[53]
TAL28131	5	078	11	NY Presbyterian/Weill Cornell Medical Center	[91]
TAL29600	5	078	11	RM Alden Research Lab	[91]
TAL29996	5	078	11	Vines VA Hospital	[91]
TAL30550	5	078	11	Mayo Clinic	[91]
TAL30574	5	078	11	Tufts Medical Center	[91]
V48	5	078	11	Brigham & Women's Hospital	This study
139b	5	078	11	Memorial Sloan Kettering	[20]
WU66	5	078	11	Barnes-Jewish Hospital	[20]

725
726 Memorial Sloan Kettering Cancer Center (MSK), Barnes-Jewish Hospital (BJH)
727
728
729 **Table 2: Cell size and growth rate statistics for the studied strains.**
730

Strain Name	Clade	Mean Growth rate	Mean Cell Length (um)	Mean Chain Length (um)
630	1	1.63 ± 0.02	8.5 ± 2.2	12.5 ± 3.5
R20291	2	1.88 ± 0.01	not measured	11.7 ± 3.2
E15	3	1.92 ± 0.01	not measured	13.6 ± 4.2
M68	4	1.64 ± 0.01	not measured	13.7 ± 4.3
M120	5	2.15 ± 0.01	not measured	21.8 ± 10.2
TAL29996	5	2.15 ± 0.02	9.4 ± 2.7	13.7 ± 3.9
TAL28131	5	2.17 ± 0.02	11.2 ± 3.3	19.7 ± 6.9
V48	5	2.10 ±0.02	10.8 ± 3.7	19.4 ± 8.5
TAL30550	5	2.05 ± 0.04	11.8 ±.3.3	27.3 ± 9.8

| TAL29600 | 5 | 2.11 ± 0.02 | 14.0 ± 4.4 | 29.1 ± 13.1 |

731
732 The $\pm$ interval represents the 95% confidence interval of the mean growth rate. The standard
733 deviation (S.D) is provided.
734
735
736

References

1.   Desai K, Gupta SB, Dubberke ER, Prabhu VS, Browne C, Mast TC. Epidemiological and economic burden of *Clostridium difficile* in the United States: estimates from a modeling approach. BMC Infect Dis. 2016;16:303. Epub 20160618. doi: 10.1186/s12879-016-1610-3. PubMed PMID: 27316794; PubMed Central PMCID: PMCPMC4912810.

2.   Guh AY, Mu Y, Winston LG, Johnston H, Olson D, Farley MM, et al. Trends in U.S. Burden of *Clostridioides difficile* Infection and Outcomes. N Engl J Med. 2020;382(14):1320-30. doi: 10.1056/NEJMoa1910215. PubMed PMID: 32242357; PubMed Central PMCID: PMCPMC7861882.

3.   Shen A. *Clostridioides difficile* Spore Formation and Germination: New Insights and Opportunities for Intervention. Annu Rev Microbiol. 2020;74:545-66. doi: 10.1146/annurev-micro-011320-011321. PubMed PMID: 32905755.

4.   Abt MC, McKenney PT, Pamer EG. *Clostridium difficile* colitis: pathogenesis and host defence. Nat Rev Microbiol. 2016. doi: 10.1038/nrmicro.2016.108. PubMed PMID: 27573580.

5.   Theriot CM, Young VB. Interactions Between the Gastrointestinal Microbiome and *Clostridium difficile*. Annu Rev Microbiol. 2015;69:445-61. doi: 10.1146/annurev-micro-091014-104115. PubMed PMID: 26488281.

6.   Kordus SL, Thomas AK, Lacy DB. *Clostridioides difficile* toxins: mechanisms of action and antitoxin therapeutics. Nat Rev Microbiol. 2022;20(5):285-98. Epub 20211126. doi: 10.1038/s41579-021-00660-2. PubMed PMID: 34837014; PubMed Central PMCID: PMCPMC9018519.

7.   Maroo S, Lamont J. Recurrent *Clostridium difficile*. Gastroenterology. 2006;130(4):1311-6. doi: 10.1053/j.gastro.2006.02.044.

8.   Zanella Terrier MC, Simonet ML, Bichard P, Frossard JL. Recurrent *Clostridium difficile* infections: the importance of the intestinal microbiota. World journal of gastroenterology : WJG. 2014;20(23):7416-23. doi: 10.3748/wjg.v20.i23.7416. PubMed PMID: 24966611; PubMed Central PMCID: PMC4064086.

9.   Knight DR, Elliott B, Chang BJ, Perkins TT, Riley TV. Diversity and Evolution in the Genome of *Clostridium difficile*. Clin Microbiol Rev. 2015;28(3):721-41. doi: 10.1128/CMR.00127-14. PubMed PMID: 26085550; PubMed Central PMCID: PMCPMC4475645.

10.   Knight DR, Imwattana K, Kullin B, Guerrero-Araya E, Paredes-Sabja D, Didelot X, et al. Major genetic discontinuity and novel toxigenic species in *Clostridioides difficile* taxonomy. Elife. 2021;10. Epub 20210611. doi: 10.7554/eLife.64325. PubMed PMID: 34114561; PubMed Central PMCID: PMCPMC8241443.

11.   Norsigian CJ, Danhof HA, Brand CK, Midani FS, Broddrick JT, Savidge TC, et al. Systems biology approach to functionally assess the *Clostridioides difficile* pangenome reveals genetic diversity with discriminatory power. Proc Natl Acad Sci U S A. 2022;119(18):e2119396119. Epub 20220427. doi: 10.1073/pnas.2119396119. PubMed PMID: 35476524; PubMed Central PMCID: PMCPMC9170149.

12.   Knight DR, Kullin B, Androga GO, Barbut F, Eckert C, Johnson S, et al. Evolutionary and Genomic Insights into *Clostridioides difficile* Sequence Type 11: a Diverse Zoonotic and Antimicrobial-Resistant Lineage of Global One Health Importance. mBio. 2019;10(2). Epub 20190416. doi: 10.1128/mBio.00446-19. PubMed PMID: 30992351; PubMed Central PMCID: PMCPMC6469969.

13.   Seth-Smith HMB, Biggel M, Roloff T, Hinic V, Bodmer T, Risch M, et al. Transition
From PCR-Ribotyping to Whole Genome Sequencing Based Typing of *Clostridioides difficile*.
Front Cell Infect Microbiol. 2021;11:681518. Epub 20210601. doi: 10.3389/fcimb.2021.681518.
PubMed PMID: 34141631; PubMed Central PMCID: PMCPMC8204696.
14.   Dingle KE, Elliott B, Robinson E, Griffiths D, Eyre DW, Stoesser N, et al. Evolutionary
history of the *Clostridium difficile* pathogenicity locus. Genome Biol Evol. 2014;6(1):36-52. doi:
10.1093/gbe/evt204. PubMed PMID: 24336451; PubMed Central PMCID: PMCPMC3914685.
15.   Mengoli M, Barone M, Fabbrini M, D'Amico F, Brigidi P, Turroni S. Make It Less
difficile: Understanding Genetic Evolution and Global Spread of *Clostridioides difficile*. Genes
(Basel). 2022;13(12). Epub 20221124. doi: 10.3390/genes13122200. PubMed PMID: 36553467;
PubMed Central PMCID: PMCPMC9778335.
16.   Clements AC, Magalhaes RJ, Tatem AJ, Paterson DL, Riley TV. *Clostridium difficile*
PCR ribotype 027: assessing the risks of further worldwide spread. Lancet Infect Dis.
2010;10(6):395-404. doi: 10.1016/S1473-3099(10)70080-3. PubMed PMID: 20510280; PubMed
Central PMCID: PMCPMC7185771.
17.   Endres BT, Begum K, Sun H, Walk ST, Memariani A, Lancaster C, et al. Epidemic
Clostridioides difficile Ribotype 027 Lineages: Comparisons of Texas Versus Worldwide Strains.
Open Forum Infect Dis. 2019;6(2):ofz013. Epub 20190209. doi: 10.1093/ofid/ofz013. PubMed
PMID: 30793006; PubMed Central PMCID: PMCPMC6368847.
18.   McDonald LC, Killgore GE, Thompson A, Owens RC, Jr., Kazakova SV, Sambol SP, et
al. An epidemic, toxin gene-variant strain of *Clostridium difficile*. N Engl J Med.
2005;353(23):2433-41. Epub 20051201. doi: 10.1056/NEJMoa051590. PubMed PMID:
16322603.
19.   Dong Q, Lin H, Allen MM, Garneau JR, Sia JK, Smith RC, et al. Virulence and genomic
diversity among clinical isolates of ST1 (BI/NAP1/027) *Clostridioides difficile*. Cell Rep.
2023;42(8):112861. Epub 20230730. doi: 10.1016/j.celrep.2023.112861. PubMed PMID:
37523264; PubMed Central PMCID: PMCPMC10627504.
20.   Lewis BB, Carter RA, Ling L, Leiner I, Taur Y, Kamboj M, et al. Pathogenicity Locus,
Core Genome, and Accessory Gene Contributions to *Clostridium difficile* Virulence. mBio.
2017;8(4). Epub 20170808. doi: 10.1128/mBio.00885-17. PubMed PMID: 28790208; PubMed
Central PMCID: PMCPMC5550754.
21.   Merrigan M, Venugopal A, Mallozzi M, Roxas B, Viswanathan VK, Johnson S, et al.
Human hypervirulent *Clostridium difficile* strains exhibit increased sporulation as well as robust
toxin production. J Bacteriol. 2010;192(19):4904-11. Epub 20100730. doi: 10.1128/JB.00445-10.
PubMed PMID: 20675495; PubMed Central PMCID: PMCPMC2944552.
22.   Nhu NTQ, Lin H, Pigli Y, Sia JK, Kuhn P, Snitkin ES, et al. Flagellar switch inverted
repeat impacts flagellar invertibility and varies *Clostridioides difficile* RT027/MLST1 virulence.
bioRxiv. 2024. Epub 20240924. doi: 10.1101/2023.06.22.546185. PubMed PMID: 39386689;
PubMed Central PMCID: PMCPMC11463649.
23.   Shaw HA, Preston MD, Vendrik KEW, Cairns MD, Browne HP, Stabler RA, et al. The
recent emergence of a highly related virulent *Clostridium difficile* clade with unique
characteristics. Clin Microbiol Infect. 2020;26(4):492-8. Epub 20190913. doi:
10.1016/j.cmi.2019.09.004. PubMed PMID: 31525517; PubMed Central PMCID:
PMCPMC7167513.
24.   Li C, Harmanus C, Zhu D, Meng X, Wang S, Duan J, et al. Characterization of the
virulence of a non-RT027, non-RT078 and binary toxin-positive *Clostridium difficile* strain

associated with severe diarrhea. Emerg Microbes Infect. 2018;7(1):211. Epub 20181212. doi:
10.1038/s41426-018-0211-1. PubMed PMID: 30542069; PubMed Central PMCID:
PMCPMC6291415.
25.   Imwattana K, Rodriguez C, Riley TV, Knight DR. A species-wide genetic atlas of
antimicrobial resistance in *Clostridioides difficile*. Microb Genom. 2021;7(11). doi:
10.1099/mgen.0.000696. PubMed PMID: 34793295; PubMed Central PMCID:
PMCPMC8743556.
26.   Kumar N, Browne HP, Viciani E, Forster SC, Clare S, Harcourt K, et al. Adaptation of
host transmission cycle during *Clostridium difficile* speciation. Nat Genet. 2019;51(9):1315-20.
Epub 20190812. doi: 10.1038/s41588-019-0478-8. PubMed PMID: 31406348.
27.   Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI
analysis of 90K prokaryotic genomes reveals clear species boundaries. Nature Comm.
2018;9(1):5114. Epub 20181130. doi: 10.1038/s41467-018-07641-9. PubMed PMID: 30504855;
PubMed Central PMCID: PMCPMC6269478.
28.   Knetsch CW, Kumar N, Forster SC, Connor TR, Browne HP, Harmanus C, et al.
Zoonotic Transfer of *Clostridium difficile* Harboring Antimicrobial Resistance between Farm
Animals and Humans. J Clin Microbiol. 2018;56(3). Epub 20180222. doi: 10.1128/JCM.01384-
17. PubMed PMID: 29237792; PubMed Central PMCID: PMCPMC5824051.
29.   Hensgens MP, Keessen EC, Squire MM, Riley TV, Koene MG, de Boer E, et al.
Clostridium difficile infection in the community: a zoonotic disease? Clin Microbiol Infect.
2012;18(7):635-45. Epub 20120427. doi: 10.1111/j.1469-0691.2012.03853.x. PubMed PMID:
22536816.
30.   Goorhuis A, Bakker D, Corver J, Debast SB, Harmanus C, Notermans DW, et al.
Emergence of *Clostridium difficile* infection due to a new hypervirulent strain, polymerase chain
reaction ribotype 078. Clin Infect Dis. 2008;47(9):1162-70. doi: 10.1086/592257. PubMed
PMID: 18808358.
31.   Lim SC, Knight DR, Riley TV. *Clostridium difficile* and One Health. Clin Microbiol
Infect. 2020;26(7):857-63. Epub 20191101. doi: 10.1016/j.cmi.2019.10.023. PubMed PMID:
31682985.
32.   Bolton D, Marcos P. The Environment, Farm Animals and Foods as Sources of
Clostridioides difficile Infection in Humans. Foods. 2023;12(5). Epub 20230304. doi:
10.3390/foods12051094. PubMed PMID: 36900611; PubMed Central PMCID:
PMCPMC10000743.
33.   Garrett EM, Sekulovic O, Wetzel D, Jones JB, Edwards AN, Vargas-Cuebas G, et al.
Phase variation of a signal transduction system controls *Clostridioides difficile* colony
morphology, motility, and virulence. PLoS Biol. 2019;17(10):e3000379. Epub 20191028. doi:
10.1371/journal.pbio.3000379. PubMed PMID: 31658249; PubMed Central PMCID:
PMCPMC6837544.
34.   Thanissery R, Winston JA, Theriot CM. Inhibition of spore germination, growth, and
toxin activity of clinically relevant *C. difficile* strains by gut microbiota derived secondary bile
acids. Anaerobe. 2017;45:86-100. Epub 20170306. doi: 10.1016/j.anaerobe.2017.03.004.
PubMed PMID: 28279860; PubMed Central PMCID: PMCPMC5466893.
35.   Beebe MA, Paredes-Sabja D, Kociolek LK, Rodriguez C, Sorg JA. Phenotypic analysis
of various *Clostridioides difficile* ribotypes reveals consistency among core processes. bioRxiv.
2025. Epub 20250110. doi: 10.1101/2025.01.10.632434. PubMed PMID: 39829883; PubMed
Central PMCID: PMCPMC11741275.

36. Ackermann M. A functional perspective on phenotypic heterogeneity in microorganisms.
Nat Rev Microbiol. 2015;13(8):497-508. Epub 20150706. doi: 10.1038/nrmicro3491. PubMed
PMID: 26145732.
37. Aldridge BB, Fernandez-Suarez M, Heller D, Ambravaneswaran V, Irimia D, Toner M,
et al. Asymmetry and aging of mycobacterial cells lead to variable growth and antibiotic
susceptibility. Science. 2012;335(6064):100-4. Epub 20111215. doi: 10.1126/science.1216166.
PubMed PMID: 22174129; PubMed Central PMCID: PMCPMC3397429.
38. Anjuwon-Foster BR, Tamayo R. A genetic switch controls the production of flagella and
toxins in *Clostridium difficile*. PLoS Genet. 2017;13(3):e1006701. Epub 20170327. doi:
10.1371/journal.pgen.1006701. PubMed PMID: 28346491; PubMed Central PMCID:
PMCPMC5386303.
39. Anjuwon-Foster BR, Tamayo R. Phase variation of *Clostridium difficile* virulence factors.
Gut microbes. 2018;9(1):76-83. Epub 20170921. doi: 10.1080/19490976.2017.1362526.
PubMed PMID: 28806147; PubMed Central PMCID: PMCPMC5914908.
40. Kint N, Janoir C, Monot M, Hoys S, Soutourina O, Dupuy B, et al. The alternative sigma
factor sigma(B) plays a crucial role in adaptive strategies of *Clostridium difficile* during gut
infection. Environ Microbiol. 2017;19(5):1933-58. Epub 20170321. doi: 10.1111/1462-
2920.13696. PubMed PMID: 28198085.
41. Courson DS, Pokhrel A, Scott C, Madrill M, Rinehold AJ, Tamayo R, et al. Single cell
analysis of nutrient regulation of *Clostridioides* (*Clostridium*) *difficile* motility. Anaerobe.
2019;59:205-11. Epub 20190803. doi: 10.1016/j.anaerobe.2019.102080. PubMed PMID:
31386902; PubMed Central PMCID: PMCPMC6785396.
42. de Jong IG, Beilharz K, Kuipers OP, Veening JW. Live Cell Imaging of *Bacillus subtilis*
and Streptococcus pneumoniae using Automated Time-lapse Microscopy. J Vis Exp. 2011;(53).
Epub 20110728. doi: 10.3791/3145. PubMed PMID: 21841760; PubMed Central PMCID:
PMCPMC3197447.
43. Lugagne JB, Lin H, Dunlop MJ. DeLTA: Automated cell segmentation, tracking, and
lineage reconstruction using deep learning. PLoS Comput Biol. 2020;16(4):e1007673. Epub
20200413. doi: 10.1371/journal.pcbi.1007673. PubMed PMID: 32282792; PubMed Central
PMCID: PMCPMC7153852.
44. O'Connor OM, Alnahhas RN, Lugagne JB, Dunlop MJ. DeLTA 2.0: A deep learning
pipeline for quantifying single-cell spatial and temporal dynamics. PLoS Comput Biol.
2022;18(1):e1009797. Epub 20220118. doi: 10.1371/journal.pcbi.1009797. PubMed PMID:
35041653; PubMed Central PMCID: PMCPMC8797229.
45. Pogliano J, Osborne N, Sharp MD, Abanes-De Mello A, Perez A, Sun YL, et al. A vital
stain for studying membrane dynamics in bacteria: a novel mechanism controlling septation
during *Bacillus subtilis* sporulation. Mol Microbiol. 1999;31(4):1149-59. PubMed PMID:
10096082; PubMed Central PMCID: PMC2885269.
46. Garrett EM, Mehra A, Sekulovic O, Tamayo R. Multiple Regulatory Mechanisms
Control the Production of CmrRST, an Atypical Signal Transduction System in *Clostridioides*
difficile. mBio. 2021;13(1):e0296921. Epub 20220215. doi: 10.1128/mbio.02969-21. PubMed
PMID: 35164558; PubMed Central PMCID: PMCPMC8844915.
47. Karasawa T, Ikoma S, Yamakawa K, Nakamura S. A defined growth medium for
Clostridium difficile. Microbiology (Reading). 1995;141 (Pt 2):371-5. doi: 10.1099/13500872-
141-2-371. PubMed PMID: 7704267.

48.   Sauls JT, Cox SE, Do Q, Castillo V, Ghulam-Jelani Z, Jun S. Control of *Bacillus subtilis*
Replication Initiation during Physiological Transitions and Perturbations. mBio. 2019;10(6).
Epub 20191217. doi: 10.1128/mBio.02205-19. PubMed PMID: 31848269; PubMed Central
PMCID: PMCPMC6918070.
49.   Eswaramoorthy P, Duan D, Dinh J, Dravis A, Devi SN, Fujita M. The threshold level of
the sensor histidine kinase KinA governs entry into sporulation in *Bacillus subtilis*. J Bacteriol.
2010;192(15):3870-82. Epub 20100528. doi: 10.1128/JB.00466-10. PubMed PMID: 20511506;
PubMed Central PMCID: PMCPMC2916370.
50.   Narula J, Kuchina A, Zhang F, Fujita M, Suel GM, Igoshin OA. Slowdown of growth
controls cellular differentiation. Mol Syst Biol. 2016;12(5):871. Epub 20160523. doi:
10.15252/msb.20156691. PubMed PMID: 27216630; PubMed Central PMCID:
PMCPMC5289222.
51.   Mehdizadeh Gohari I, Edwards AN, McBride SM, McClane BA. The impact of orphan
histidine kinases and phosphotransfer proteins on the regulation of clostridial sporulation
initiation. mBio. 2024;15(4):e0224823. Epub 20240313. doi: 10.1128/mbio.02248-23. PubMed
PMID: 38477571; PubMed Central PMCID: PMCPMC11210211.
52.   Deatherage DE, Traverse CC, Wolf LN, Barrick JE. Detecting rare structural variation in
evolving microbial populations from new sequence junctions using breseq. Front Genet.
2014;5:468. Epub 20150121. doi: 10.3389/fgene.2014.00468. PubMed PMID: 25653667;
PubMed Central PMCID: PMCPMC4301190.
53.   He M, Sebaihia M, Lawley T, Stabler R, Dawson L, Martin M, et al. Evolutionary
dynamics of *Clostridium difficile* over short and long time scales. Proceedings of the National
Academy of Sciences of the United States of America. 2010;107(16):7527-32. doi:
10.1073/pnas.0914322107.
54.   Sekulovic O, Mathias Garrett E, Bourgeois J, Tamayo R, Shen A, Camilli A. Genome-
wide detection of conservative site-specific recombination in bacteria. PLoS Genet.
2018;14(4):e1007332. Epub 20180405. doi: 10.1371/journal.pgen.1007332. PubMed PMID:
29621238; PubMed Central PMCID: PMCPMC5903667.
55.   Purcell EB, McKee RW, Bordeleau E, Burrus V, Tamayo R. Regulation of Type IV Pili
Contributes to Surface Behaviors of Historical and Epidemic Strains of *Clostridium difficile*. J
Bacteriol. 2016;198(3):565-77. Epub 20151123. doi: 10.1128/JB.00816-15. PubMed PMID:
26598364; PubMed Central PMCID: PMCPMC4719444.
56.   Glomski IJ, Piris-Gimenez A, Huerre M, Mock M, Goossens PL. Primary involvement of
pharynx and peyer's patch in inhalational and intestinal anthrax. PLoS Pathog. 2007;3(6):e76. doi:
10.1371/journal.ppat.0030076. PubMed PMID: 17542645; PubMed Central PMCID:
PMCPMC1885272.
57.   McKenney PT, Yan J, Vaubourgeix J, Becattini S, Lampen N, Motzer A, et al. Intestinal
Bile Acids Induce a Morphotype Switch in Vancomycin-Resistant Enterococcus that Facilitates
Intestinal Colonization. Cell Host Microbe. 2019;25(5):695-705 e5. Epub 20190425. doi:
10.1016/j.chom.2019.03.008. PubMed PMID: 31031170; PubMed Central PMCID:
PMCPMC6939634.
58.   Leslie JL, Jenior ML, Vendrov KC, Standke AK, Barron MR, O'Brien TJ, et al.
Protection from Lethal *Clostridioides difficile* Infection via Intraspecies Competition for
Cogerminant. mBio. 2021;12(2). Epub 20210330. doi: 10.1128/mBio.00522-21. PubMed PMID:
33785619; PubMed Central PMCID: PMCPMC8092246.

59.   Theriot CM, Koumpouras CC, Carlson PE, Bergin, II, Aronoff DM, Young VB.
Cefoperazone-treated mice as an experimental platform to assess differential virulence of
Clostridium difficile strains. Gut microbes. 2011;2(6):326-34. Epub 20111101. doi:
10.4161/gmic.19142. PubMed PMID: 22198617; PubMed Central PMCID: PMCPMC3337121.
60.   Sachsenheimer FE, Yang I, Zimmermann O, Wrede C, Muller LV, Gunka K, et al.
Genomic and phenotypic diversity of *Clostridium difficile* during long-term sequential
recurrences of infection. Int J Med Microbiol. 2018;308(3):364-77. Epub 20180221. doi:
10.1016/j.ijmm.2018.02.002. PubMed PMID: 29490877.
61.   Semenyuk EG, Poroyko VA, Johnston PF, Jones SE, Knight KL, Gerding DN, et al.
Analysis of Bacterial Communities during *Clostridium difficile* Infection in the Mouse. Infect
Immun. 2015;83(11):4383-91. Epub 20150831. doi: 10.1128/IAI.00145-15. PubMed PMID:
26324536; PubMed Central PMCID: PMCPMC4598419.
62.   Smith AB, Jenior ML, Keenan O, Hart JL, Specker J, Abbas A, et al. Enterococci
enhance *Clostridioides difficile* pathogenesis. Nature. 2022;611(7937):780-6. Epub 20221116.
doi: 10.1038/s41586-022-05438-x. PubMed PMID: 36385534; PubMed Central PMCID:
PMCPMC9691601.
63.   Kim HY, Cho A, Kim JW, Kim H, Kim B. High prevalence of *Clostridium difficile* PCR
ribotype 078 in pigs in Korea. Anaerobe. 2018;51:42-6. Epub 20180328. doi:
10.1016/j.anaerobe.2018.03.012. PubMed PMID: 29604338.
64.   Moono P, Foster NF, Hampson DJ, Knight DR, Bloomfield LE, Riley TV. *Clostridium
difficile* Infection in Production Animals and Avian Species: A Review. Foodborne Pathog Dis.
2016;13(12):647-55. Epub 20160907. doi: 10.1089/fpd.2016.2181. PubMed PMID: 27602596.
65.   Burt SA, Siemeling L, Kuijper EJ, Lipman LJ. Vermin on pig farms are vectors for
Clostridium difficile PCR ribotypes 078 and 045. Vet Microbiol. 2012;160(1-2):256-8. Epub
20120518. doi: 10.1016/j.vetmic.2012.05.014. PubMed PMID: 22682200.
66.   Dhungel BA, Govind R. Phase-variable expression of pdcB, a phosphodiesterase,
influences sporulation in *Clostridioides difficile*. Mol Microbiol. 2021;116(5):1347-60. Epub
20211018. doi: 10.1111/mmi.14828. PubMed PMID: 34606654; PubMed Central PMCID:
PMCPMC8876291.
67.   Edwards AN, Willams CL, Pareek N, McBride SM, Tamayo R. c-di-GMP Inhibits Early
Sporulation in *Clostridioides difficile*. mSphere. 2021;6(6):e0091921. Epub 20211208. doi:
10.1128/msphere.00919-21. PubMed PMID: 34878288; PubMed Central PMCID:
PMCPMC8653836.
68.   Du S, Henke W, Pichoff S, Lutkenhaus J. How FtsEX localizes to the Z ring and interacts
with FtsA to regulate cell division. Mol Microbiol. 2019;112(3):881-95. Epub 20190620. doi:
10.1111/mmi.14324. PubMed PMID: 31175681; PubMed Central PMCID: PMCPMC6831102.
69.   Li J, Xu X, Shi J, Hermoso JA, Sham LT, Luo M. Regulation of the cell division
hydrolase RipC by the FtsEX system in *Mycobacterium tuberculosis*. Nature communications.
2023;14(1):7999. Epub 20231204. doi: 10.1038/s41467-023-43770-6. PubMed PMID: 38044344;
PubMed Central PMCID: PMCPMC10694151.
70.   Pichoff S, Du S, Lutkenhaus J. Roles of FtsEX in cell division. Res Microbiol.
2019;170(8):374-80. Epub 20190801. doi: 10.1016/j.resmic.2019.07.003. PubMed PMID:
31376483; PubMed Central PMCID: PMCPMC6899183.
71.   Xu X, Li J, Chua WZ, Pages MA, Shi J, Hermoso JA, et al. Mechanistic insights into the
regulation of cell wall hydrolysis by FtsEX and EnvC at the bacterial division site. Proc Natl

Acad Sci U S A. 2023;120(21):e2301897120. Epub 20230515. doi: 10.1073/pnas.2301897120. PubMed PMID: 37186861; PubMed Central PMCID: PMCPMC10214136.

72.   Garcia-Garcia T, Poncet S, Cuenot E, Douche T, Giai Gianetto Q, Peltier J, et al. Ser/Thr Kinase-Dependent Phosphorylation of the Peptidoglycan Hydrolase CwlA Controls Its Export and Modulates Cell Division in *Clostridioides difficile*. mBio. 2021;12(3). Epub 20210518. doi: 10.1128/mBio.00519-21. PubMed PMID: 34006648; PubMed Central PMCID: PMCPMC8262956.

73.   Ransom EM, Williams KB, Weiss DS, Ellermeier CD. Identification and characterization of a gene cluster required for proper rod shape, cell division, and pathogenesis in *Clostridium difficile*. J Bacteriol. 2014;196(12):2290-300. doi: 10.1128/JB.00038-14. PubMed PMID: 24727226; PubMed Central PMCID: PMC4054185.

74.   Streett H, Charubin K, Papoutsakis ET. Anaerobic fluorescent reporters for cell identification, microbial cell biology and high-throughput screening of microbiota and genomic libraries. Curr Opin Biotechnol. 2021;71:151-63. Epub 20210807. doi: 10.1016/j.copbio.2021.07.005. PubMed PMID: 34375813.

75.   Anjou C, Lotoux A, Zhukova A, Royer M, Caulat LC, Capuzzo E, et al. The multiplicity of thioredoxin systems meets the specific lifestyles of Clostridia. PLoS Pathog. 2024;20(2):e1012001. Epub 20240208. doi: 10.1371/journal.ppat.1012001. PubMed PMID: 38330058; PubMed Central PMCID: PMCPMC10880999.

76.   Shen A, Fimlaid KA, Pishdadian K. Inducing and Quantifying *Clostridium difficile* Spore Formation. Methods Mol Biol. 2016;1476:129-42. doi: 10.1007/978-1-4939-6361-4_10. PubMed PMID: 27507338.

77.   Ransom EM, Weiss DS, Ellermeier CD. Use of mCherryOpt Fluorescent Protein in *Clostridium difficile*. Methods Mol Biol. 2016;1476:53-67. Epub 2016/08/11. doi: 10.1007/978-1-4939-6361-4_5. PubMed PMID: 27507333.

78.   Pishdadian K, Fimlaid KA, Shen A. SpoIIID-mediated regulation of sigma(K) function during *Clostridium difficile* sporulation. Mol Microbiol. 2015;95(2):189-208. doi: 10.1111/mmi.12856. PubMed PMID: 25393584.

79.   Fimlaid KA, Jensen O, Donnelly ML, Francis MB, Sorg JA, Shen A. Identification of a Novel Lipoprotein Regulator of *Clostridium difficile* Spore Germination. PLoS Pathogens. 2015. Epub October 23, 2015. doi: 10.1371/journal.ppat.1005239.

80.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De Novo Assembler. Current protocols in bioinformatics / editoral board, Andreas D Baxevanis [et al]. 2020;70(1):e102. doi: 10.1002/cpbi.102. PubMed PMID: 32559359.

81.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30(14):2068-9. Epub 20140318. doi: 10.1093/bioinformatics/btu153. PubMed PMID: 24642063.

82.   Benedict MN, Henriksen JR, Metcalf WW, Whitaker RJ, Price ND. ITEP: an integrated toolkit for exploration of microbial pan-genomes. BMC Genomics. 2014;15:8. Epub 20140103. doi: 10.1186/1471-2164-15-8. PubMed PMID: 24387194; PubMed Central PMCID: PMCPMC3890548.

83.   Eren AM, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, et al. Community-led, integrated, reproducible multi-omics with anvi'o. Nat Microbiol. 2021;6(1):3-6. doi: 10.1038/s41564-020-00834-3. PubMed PMID: 33349678; PubMed Central PMCID: PMCPMC8116326.

84.   van Dongen S, Abreu-Goodger C. Using MCL to extract clusters from networks.
Methods Mol Biol. 2012;804:281-95. doi: 10.1007/978-1-61779-361-5_15. PubMed PMID:
22144159.
85.   Deatherage DE, Barrick JE. Identification of mutations in laboratory-evolved microbes
from next-generation sequencing data using breseq. Methods Mol Biol. 2014;1151:165-88. doi:
10.1007/978-1-4939-0554-6_12. PubMed PMID: 24838886; PubMed Central PMCID:
PMCPMC4239701.
86.   César N. Dataset for the article: Unique growth and morphology properties of Clade 5
Clostridioides difficile strains revealed by single-cell time-lapse microscopy. Zenodo. 2024. doi:
https://doi.org/10.5281/zenodo.13352469.
87.   Sebaihia M, Wren B, Mullany P, Fairweather N, Minton N, Stabler R, et al. The
multidrug-resistant human pathogen *Clostridium difficile* has a highly mobile, mosaic genome.
Nature genetics. 2006;38(7):779-86. doi: 10.1038/ng1830.
88.   Stabler R, He M, Dawson L, Martin M, Valiente E, Corton C, et al. Comparative genome
and phenotypic analysis of *Clostridium difficile* 027 strains provides insight into the evolution of
a hypervirulent bacterium. Genome biology. 2009;10(9). doi: 10.1186/gb-2009-10-9-r102.
89.   Kurka H, Ehrenreich A, Ludwig W, Monot M, Rupnik M, Barbut F, et al. Sequence
similarity of *Clostridium difficile* strains by analysis of conserved genes and genome content is
reflected by their ribotype affiliation. PLoS One. 2014;9(1):e86535. Epub 20140123. doi:
10.1371/journal.pone.0086535. PubMed PMID: 24482682; PubMed Central PMCID:
PMCPMC3902958.
90.   Moura I, Monot M, Tani C, Spigaglia P, Barbanti F, Norais N, et al. Multidisciplinary
analysis of a nontoxigenic *Clostridium difficile* strain with stable resistance to metronidazole.
Antimicrobial agents and chemotherapy. 2014;58(8):4957-60. Epub 20140609. doi:
10.1128/AAC.02350-14. PubMed PMID: 24913157; PubMed Central PMCID:
PMCPMC4135993.
91.   Thorpe CM, McDermott LA, Tran MK, Chang J, Jenkins SG, Goldstein EJC, et al. U.S.-
Based National Surveillance for Fidaxomicin Susceptibility of *Clostridioides difficile*-Associated
Diarrheal Isolates from 2013 to 2016. Antimicrobial agents and chemotherapy. 2019;63(7). Epub
20190624. doi: 10.1128/AAC.00391-19. PubMed PMID: 31085514; PubMed Central PMCID:
PMCPMC6591623.
1086
1087

	Figure Legends

	Figure 1. Schematic of the anaerobic single-cell imaging set-up. Exponentially growing *C.*

	difficile cells in TY medium supplemented with cysteine (TYC) are spotted onto 1.5% agarose

	pads formed within gas-tight adhesive Gene Frames inside the anaerobic chamber. Up to 6

	strains can be spotted onto a pad. The pad is sealed with a coverslip, and the imaging chamber

	is removed from the anaerobic chamber and transferred to a heated (37˚C) microscope stage.

	Time-lapse microscopy is used to visualize the growth of individual bacterial cells for 2-6 hrs.

	The output data is segmented and tracked with the DeLTA Python package (191). An example

	filmstrip of strain 630 grown on TYC medium over time is shown (Bottom).

	Figure 2. Clade 5 strain M120 elongates more quickly and exhibits cell chaining. (A)

	Violin plot of the elongation rates measured during time-lapse microscopy analyses of strains

	630 (Clade 1), R20291 (Clade 2), E15 (Clade 3), M68 (Clade 4), and M120 (Clade 5) grown on

	TY supplemented with cysteine (TYC) agar. Data are from three biological replicates, with the

	mean of each replicate shown as a point on the violin. (B) Optical density-based analyses of

	bulk population growth of the indicated strains in TYC or BHIS media. The number in brackets

	indicates the clade to which a given strain belongs. (C) Violin plot of the cell or chain length

	measured during time-lapse microscopy for the strains shown in **A**. Each replicate mean is

	shown as a point on the violin. Statistical significance for A and B was determined by

	comparing the mean of the three replicates of strains from Clades 1-4 strains relative to the

	Clade 5 M120 strain using a Kruskal-Wallis test * $p < 0.05$. (B) Phase-contrast image from

	time-lapse microscopy movies. Scale bar is 10 μm.

	Figure 3. *C. difficile* Clade 5 can form large heterogenous chains. Large mosaic phase-

	contrast image of the Clade 5 strain M120. Inset shows chains revealed by staining with the

	membrane dye FM4-64; septa are highlighted with yellow arrows. The image was stitched

	from 8 individual fields of view at 63X magnification. Scale bar is 100 μm.

	Figure 4. Faster growth is a common feature of Clade 5 strains, but the chaining

	phenotype is not fully penetrant in Clade 5 strains. (A) Phase-contrast microscopy images

	from time-lapse microscopy studies of Clade 5 strains; Clade 1 strain 630 is included for

comparison. Fluorescence microscopy was used to visualize the FM4-64 stain incorporated into

the agarose pads. Masks generated using DeLTA are shown, with the bottom panel showing

division septa identified with our adaptive thresholding approach (red dots). (B) Example image

showing the parameters identified using DeLTA combined with our adaptive threshold method

for detecting division septa (red dot) within a chain of cells. Following the automated

thresholding analysis for detecting septa, the images were manually inspected to ensure that all

septa were properly identified. (C) Cell or chain length measured using automated DeLTA

analyses (pink violin plot) or DeLTA combined with the adaptive threshold-manual inspection

analyses (purple violin plot). The former method is more likely to measure chain length, while

the latter method accurately measures cell length. (D) Violin plot of the elongation rates

measured based on three biological replicates. Each replicate mean is shown as a point on the

violin; statistical significance was determined by comparing the mean of the three replicates

using a Kruskal-Wallis test, * $p < 0.05$.

1133

Figure 5. Clade 5 strains form chains during logarithmic growth in broth culture.

Representative micrographs showing phase-contrast (top) and peptidoglycan labeling with the

fluorescent D-amino acid, HADA, (bottom) following growth in rich broth (BHIS) to mid-

logarithmic phase. All strains shown, with the exception of Clade 1 strain 630, are Clade 5

strains. Scale bar, 10 µm. Data are representative of three independent experiments. **ADJ**

indicates that the brightness of the image was enhanced to detect HADA labeling in V48.

1140

Figure 6. Sporulation levels in Clade 5 isolates grown on 70:30 medium. Phase-contrast

microscopy of the indicated strains ~24 hrs after sporulation induction. All strains shown with

the exception of 630 (Clade 1) are Clade 5 strains. The percent heat-resistant spores is indicated

below the respective images. The percentage was determined from 20-24 hr sporulating cultures

and represent the mean and standard deviation for a given strain based on a minimum of three

biological replicates. Statistical significance relative to strain 630 was determined using a one-

way ANOVA and Tukey's test. The scale bar represents 5 µm.

1148

Figure 7. The orientation of the *cmr* switch promotes Clade 5 strain cell chaining and

surface motility. (A) Orientation-specific qPCR for detecting the orientation of the *cmr* switch

in the indicated strains. The mean and standard deviation based on one to three biological

replicates are shown. Statistical significance relative to strain TAL29996 was determined using a

one-way ANOVA and Tukey's test for strains where data from three independent experiments

was obtained. **** $p < 0.0001$. (B) Representative micrographs of *cmrRST* CRISPRi knock-

down strains compared to a no target control. Phase-contrast (top) and peptidoglycan labeling

with the fluorescent D-amino acid, HADA, (bottom) images following growth in rich broth

(BHIS) to mid-logarithmic phase. Scale bar, 5 µm. (C) Representative images of surface

motility 5 days after exponentially growing liquid cultures of the indicated strains were spotted

onto BHIS agar plates.

1160

Figure 8. Infection and colonization dynamics of Clade 5 strains in mice. (A) Fecal colony-

forming units measured by selective plating and (B) Percentage of weight loss to baseline of

infected mice on Days 1-4, 7, and 14. The mean and standard deviation are shown based on the

results of two experiments consisting of four mice each (n = 8). (C) Percentage of *cmr*-ON

switch orientation measured in fecal pellets on the indicated days by qPCR. For the strain

TAL29996 inoculum, 1% of the spores had the *cmr*-ON switch orientation, while the TAL29600

inoculum consisted of 32% *cmr*-ON spores. The mean and standard deviation based on analyses

of eight mice are shown, although fecal pellets could not be collected from some of the mice on

Day 2. The same mouse exhibited a higher *cmr*-ON frequency for TAL29600 over time (15%,

Day 7 and 27%, Day 14).

1171
1172

Supporting Information

S1 Fig. Growth rates measured from optical density-based analyses of bulk population growth in TYC and BHIS media.

S2 Fig. Optical density-based analyses of bulk population growth of the indicated strains during growth in TYC and BHIS media.

S3 Fig. Clade 5, but not Clade 1 or 4, strains form chains during logarithmic growth in BHIS broth culture.

S4 Fig. Clade 5 strains form chains even when growth is slowed during growth in minimal medium broth culture.

S5 Fig. Sporulation levels in clinical isolates grown on 70:30 medium.

S6 Fig. Spores purified from Clade 5 strains.

S7 Fig. Expression of the *cmrRST* operon promotes cell chaining.

S8 Fig. Cell length during growth on 70:30 sporulation medium.

S9 Fig. Surface motility in Clade 1-4 strains.

S10 Fig. Infection and *cmr* switch orientation dynamics during TAL29600 infection.

S1 Table. Average nucleotide identity for orthologous genes for Clade 5 strains

S2 Table. Breseq analyses of Clade 5 strain genomes.

S3 Table. Anvi'o analysis of the accessory genome identified for the Clade 5 strains sequenced.

1. Grow Bacteria to Mid-log

2. Assemble and Inoculate Pad in Anaerobic Chamber

1 µL Culture

Coverglass

Adhesive "Gene Frame"

1.5% Agarose Medium with reducing agent

3. Time-lapse Imaging @ 37°C

4. Cell Segmentation using DeLTA

| 0 min | 10 min | 20 min | 30 min | 40 min | 50 min |

Bright-field

10µm

Segmentation

630	TAL29996	TAL28131	V48	TAL30550	TAL29600
8 ± 4%	40 ± 14% p < 0.01	163 ± 78% p < 0.0001	26 ± 7%	31 ± 21%	0.003 ± 0.001% p < 0.0001

A.

% *cmr*-ON orientation

B.

C.

| TAL29996 | TAL28131 | V48 | TAL30550 | TAL29600 |

| TAL30574 | 139b | WU66 | M120 |

A.

Colonization levels

B.

% Weight Change

— 630   — TAL29996   — TAL28131   — V48   — TAL30550   — TAL29600

C.

***% cmr*-ON**

● TAL29996
● TAL29600