Immune escape and replicative capacity of Omicron lineages BA.1, BA.2, BA.5.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1

Meriem Bekliz[#1,3], Manel Essaidi-Laziosi[#1,3], Kenneth Adea[2], Krisztina Hosszu-Fellous[2,3], Catia Alvarez[1,2], Mathilde Bellon[1,2], Pascale Sattonnet-Roche[2], Olha Puhach[1,2], Damien Dbeissi[1,2], Maria Eugenia Zaballa[4], Silvia Stringhini[4,5,6], Idris Guessous[4,5], Pauline Vetter[2,3], Christiane S Eberhardt[7,8,9], Laurent Kaiser[2,3], Isabella Eckerle[1,2,3*]

[#]contributed equally

[1]*Department of Medicine, University of Geneva, Geneva, Switzerland.*

[2]*Geneva Centre for Emerging Viral Diseases, University Hospitals of Geneva and University of Geneva, Geneva, Switzerland.*

[3]*Division of Infectious Diseases, Geneva University Hospitals, Geneva, Switzerland*

[4]*Unit of Population Epidemiology, Division of Primary Care Medicine, Geneva University Hospitals, Geneva, Switzerland.*

[5]*Department of Health and Community Medicine, Faculty of Medicine, University of Geneva, Geneva, Switzerland.*

[6]*University Center for General Medicine and Public Health, University of Lausanne, Lausanne, Switzerland.*

[7]*Center for Vaccinology and Neonatal Immunology, Department of Pathology and Immunology, University of Geneva, Geneva, Switzerland.*

[8]*Division of General Pediatrics, Department of Woman, Child and Adolescent Medicine, Faculty of Medicine, University of Geneva, Geneva, Switzerland.*

[9]*Center of Vaccinology, University Hospitals of Geneva, Geneva, Switzerland*

Corresponding author:

Isabella Eckerle

Geneva Centre for Emerging Viral Diseases, Geneva University Hospitals, Geneva, Switzerland. Geneva University Hospitals, Rue Gabrielle-Perret-Gentil 4, 1205 Geneva, Switzerland.

Tel: +41223729820;

Email: isabella.eckerle@unige.ch

Abstract

In the 5th year after the emergence of SARS-CoV-2, Omicron lineages continue to evolve and cause infections. Here, we used eight authentic SARS-CoV-2 isolates to assess their capacity to escape immunity of different exposure histories and their replicative capacity in polarized human airway epithelial cells (HAE) derived from the nasal and bronchial epithelium.

Using live-virus neutralization assays of 108 human sera or plasma of different immunological backgrounds, progressive immune escape was observed from B.1 (ancestral virus) to EG.5.1, but no significant difference between EG.5.1 and JN.1.1. Vaccinated individuals without natural infection and individuals with a single infection, but no vaccination showed markedly reduced or completely lost neutralization against the latest variants, while in those with hybrid immunity almost all sera showed some neutralization capacity. Furthermore, although absolute titers differed between groups, the pattern of immune escape between the variants remains comparable with strongest loss of neutralization observed for the latest variants.

In vitro studies with HAE at 33°C and 37°C showed some, but minor differences in virus replication and innate immune responses upon infection. Notably, infection with XBB.1.5, EG.5.1 and JN.1.1 showed slightly increased viral growth in nasal HAE at 33°C.

Altogether, these data underscore increasing immune escape across heterogeneous immunological backgrounds with gradually increasing antibody escape of evolving Omicron lineages until variant EG.5.1, but not any further for the latest dominant lineage JN.1.1. They also suggest that viral dynamics within Omicron lineages are driven by a combination of immune evasion and increase in viral replication.

Keywords: SARS-CoV-2, omicron variants, viral replication, immune escape, neutralisation.

Introduction

The emergence of the SARS-CoV-2 in late 2019 has been followed by emergence of numerous virus variants [1]. Two years later, the emergence of the Omicron variant of concern (VOC) marked a significant shift, enhancing immune evasion due to extensive mutations in its Spike region. This has led to widespread infections by overcoming immunity from vaccines and prior natural infection. Omicron's distinct genetic profile, particularly in the Spike region, positions it as a potential new serotype of SARS-CoV-2 [2, 3].

Global observations have documented successive waves of infections since the initial emergence of Omicron, with a huge number of evolving Omicron lineages. Consequently, a complex immunological landscape now exists in the population, where immunity is derived from vaccines, one or more infections due to pre-Omicron or Omicron variants, or a combination therefore, known as hybrid immunity. In most individuals in high income countries, the immunological background of SARS-CoV-2 today consists of multiple vaccinations and multiple exposures to SARS-CoV-2 through natural infection. On the other hand, some individuals have never received a vaccine and were only infected after the circulation of Omicron, in particular younger children, for which there is no vaccine recommendation in many countries. This leads to a heterogeneous immunological background in the population, ranging from both extremes of the exposure spectrum [2-9]. Despite background immunity in the population, SARS-CoV-2 continues to circulate. Multiple new variants have arisen within the Omicron clade, with XBB.1.5, XBB.1.16, EG.5, BA.2.86 and JN.1, being the latest variants of interest (VOIs) designated by WHO. BA.2.86, first identified in August 2023, has a remarkable number of mutations, known to allow antibody evasion, compared to earlier Omicron variants [10]. BA.2.86 did not show strong epidemiological signs of spread, but its decedent did, after BA.2.86 acquired an additional mutation S:L455S in the

Spike, which became JN.1. This mutation was shown to be associated with significantly

enhanced immune evasion capabilities but also increased transmissibility [5, 11-15]. In early

2024, JN.1 and its descendant lineages showed a strong increase globally, outcompeting earlier

variants, and remains dominant as of mid 2024.

With reduced testing and surveillance and a highly variable immunological background in the

population, obtaining data from epidemiology and/or clinical specimen collections has become

more challenging. Therefore, an experimental assessment of immune escape and replicative

capacity of emerging Omicron lineages *in vitro* is interesting, although it does not fully reflect

the *in vivo* situation. Here, we investigated these aspects in SARS-CoV-2 Omicron lineages:

BA.1, BA.2, BA.5.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1 compared to the ancestral SARS-

CoV-2 B.1, using live viruses as a widely acknowledged gold standard, first, to compare

immune escape capacities using sera from individuals with different immunological

backgrounds by neutralization assays and second, to study the infection in relevant primary

polarized airway epithelial culture models of the upper and lower respiratory tract.

Methods

Viruses and cells

Vero-E6 (ATCC CRL-1586) and Vero-E6-TMPRSS (Vero-E6 overexpressing TMPRSS2

protease, provided by National Institute for Biological Standards and Controls, NIBSC, Cat.

Nr. 100978) cells were cultured as previously described [2, 16, 17]. All SARS-CoV-2 viruses

used in this study were isolated from anonymized nasopharyngeal swabs collected at

University Hospitals of Geneva (HUG) under an approval that allows the usage of anonymized

left-over materials for virus culture. For this study, the following virus isolates were used

(according to Pango lineages designation [18]): B.1 (ancestral SARS-CoV-2) and BQ.1,

isolated and propagated on Vero-E6; BA.1 and BA.5.1, isolated on Vero-TMPRSS, then

propagated to Vero-E6; BA.2, XBB.1.5, EG.5.1 and JN.1.1, isolated and propagated on Vero-

TMPRSS. Both the initial clinical specimen and the obtained virus isolates were fully

sequenced (**Table S1**). All virus stocks were titrated on the same cell line on which the virus

stock was produced (either Vero-E6 or Vero-TMPRSS).

SARS-CoV-2 infections in HAE

Infections with SARS-CoV-2 Omicron lineages BA.1, BA.2, BA.5.1, BQ.1, XBB.1.5, EG.5.1

and JN.1.1 were performed at 37°C or 33°C at 5% CO_2 at a multiplicity of infection of

approximately 0.1 in commercially available polarized tissues "MucilAir™" (Epithelix

SARL), *in vitro* reconstituted from human nasal or bronchial (3 donors from each group)

epithelial cells of adult healthy donors cultured in an air-liquid interface (ALI) system, as

previously described [16, 19, 20]. Viral replication was assessed at 24, 48, 72 and 96 hours

post infection (hpi), as previously described [16, 19, 20].

Assessment of host gene response

Induction of interferons IFN-α and IFN-β IFN-λ, ISG15 (Interferon stimulated gene 15),

angiotensin-converting enzyme 2 (ACE-2) and Transmembrane Serine Protease 2 (TMPRSS2)

was assessed by semi-quantitative real-time PCR for intracellular RNA collected at 96 hpi, as

previously described [16, 19, 20].

Human serum and plasma samples

Immunocompetent and healthy individual samples consisted of serum or plasma samples

collected after vaccination, infection or a combination of both (hybrid immunity). Plasma or

serum samples from vaccinated healthy individuals, vaccinated either with two or three doses

(boosted) of BNT162b2 (Pfizer/BioNTech) or mRNA-1273 (Moderna) were available from a

prospective observational studies (Ethics approval number: CCER 2021-00430 and CCER

2020-02323). Asymptomatic or undetected infections of the vaccinated-only group were excluded in those samples by testing all specimens for antibodies against SARS-CoV-2 nucleocapsid (Roche Elecsys anti-SARS-CoV-2 N assay). Specimens of individuals with hybrid immunity were collected from adults vaccinated with BNT162b2 or mRNA-1273 vaccines and with one or more documented infections. Breakthrough infection samples were collected from individuals with either 2x or 3x mRNA vaccination, followed by an Omicron BA.1 or BA.2 breakthrough infection, respectively (Ethics approval number: CCER 2020-02323). Serum samples from individuals with XBB breakthrough infection had received either 2x (n=2), 3x (n=7) or 4x (n=2) mRNA vaccination, followed by breakthrough infection with one of the XBB sublineages that occurred between March-June 2023 (specimens were left-over samples from a prospective observational study, ethics approval number: CCER 2022-01722). Four of these individuals had a subsequent infection prior to XBB infection. One of the individuals vaccinated with 4x mRNA had received a bivalent vaccine. For individuals with infections from two different VOCs (e.g. Alpha and Omicron or Delta and Omicron) had received either 1x (n=6), 2x (n=10) or 3x (n=2) mRNA vaccine (Ethics approval number: CCER 2020-02323). Convalescent sera from unvaccinated adults and children with confirmed SARS-CoV-2 infection in early 2022 were also available (Ethics approval number: CCER 2020-00881). Based on the data from our Swiss national genomic surveillance, the vast majority of the variants circulating at that time was Omicron BA.1 and BA.2, with only few Delta sequences remaining in January 2022 [21]. The infecting variant of each episode was either determined by sequencing of the diagnostic samples or extrapolated by the time of infection according to the information that was self-reported by the participant and/or by their parent (for children), taking the knowledge on variant circulation generated by the Swiss national genomic surveillance program into account [22]. Data on collection time of specimens

after vaccination and/or infection are displayed in **Tables 1-5**. Only one serum per individual

of a single collection time point was used in this study.

A written informed consent was obtained from all adult participants, and from the legally

appointed representatives (parents) of all minor participants. All necessary approvals were

obtained from the Cantonal Ethical board of the Canton of Geneva, Switzerland (Commission

Cantonal d'Ethique de la Recherche, CCER). Since no differences are to be expected in

neutralizing capacity between plasma or serum, both sample types were used in parallel.

Focus reduction neutralization test (FRNT)

FRNT was used to determine the infectious titer after neutralization. Vero-TMPRSS cells were

seeded at a density of 4×10^5 cells/mL in 96-well cell culture plates. All sera/plasma and Vero-

TMPRSS cells infections were prepared as previously described [2]. After incubation for 16-

24 hours at 37 °C, 5% CO_2, the plates were fixed and stained for SARS-CoV-2 nucleocapsid

protein as described previously [17, 23]. The 90% reduction endpoint titers ($FRNT_{90}$) were

calculated as previously described [2]. For samples that did not reach 90% reduction at a 1:10

dilution, we extrapolated the titer until a dilution of 0.5. If the extrapolation reached a titer

below 0.5, the sample was given a value of 0.5. All samples with a titer below 1, i.e. undiluted

sample are considered negative.

Data was recorded in Excel 2019. Geometric means with 95% CI were used for the comparison

of $FRNT_{90}$ titers. Statistical analyses for $FRNT_{90}$ were conducted using GraphPad Prism

version 9.1.0 software and performed using repeated measures one-way ANOVA with

Dunnett's multiple comparisons test with $\log_{10}$ transformed $FRNT_{90}$ titers.

Results

1. Virus neutralization to Omicron lineages of sera or plasma after infection, vaccination, and hybrid immunity

A panel of sera/plasma was used from: (i) double or monovalent boosted vaccinated individuals with BNT162b2 or mRNA-1273 without prior or subsequent infection (**Table 1**); (ii) individuals with hybrid immunity after receiving two or three doses of mRNA vaccine followed by a break-through infection with either Omicron BA.1 or BA.2, respectively (**Table 2**); (iii) individuals with hybrid immunity due to XBB-variant breakthrough infection (**Table 3**); (iv) individuals with hybrid immunity receiving between one to three doses of mRNA vaccine and two different documented VOCs infections (either infection with Alpha or Delta, followed by infection with Omicron) (**Table 4**); and (v) a panel of convalescent sera from unvaccinated individuals, adults and children infected between January and March 2022, a period that was characterized by high circulation of Omicron BA.1 and BA.2 (**Table 5**).

2.1. *Neutralizing capacity in vaccinated but never-infected individuals against Omicron lineages*

We investigated a total of 33 individuals' specimens, either double-vaccinated (n=14) or monovalent boosted individuals that had never been infected with SARS-CoV-2 (n=19) for neutralization against ancestral SARS-CoV-2 lineage B.1 as well as SARS-CoV-2 Omicron lineages BA.1, BA.2, BA.5.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1. For double-vaccinated individuals, the highest neutralizing capacity was observed against the ancestral SARS-CoV-2 virus B.1 with geometric mean $FRNT_{90}$ titers of 251.7 (95%CI: 158.3-400.1) but very reduced titers were observed against all Omicron lineages with $FRNT_{90}$ titers of 5.7 (95%CI: 2.8-11.5) for BA.1, 10.4 (95%CI: 6.6-16.3) for BA.2, 2.9 (95%CI: 1.5-5.5) for BA.5.1, 1.5 (95%CI: 0.8-3.0) for BQ.1, 1.0 (95%CI: 0.6-1.4) for XBB.1.5 and 0.6 (95%CI: 0.5-0.8) for JN.1.1. None of

the samples neutralized EG.5.1. Although titers were very reduced compared to B.1, none of the samples failed to neutralize Omicron BA.2. An increasing number of samples with complete failure to neutralize was observed for Omicron BA.1, BA.5.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1 with 2/14, 4/14, 7/14, 7/14, 14/14 and 12/14, respectively (**Figure 1A**).

For boosted individuals, overall geometric mean $FRNT_{90}$ titers were higher against all viruses. Geometric mean $FRNT_{90}$ titers were 357.5 (95%CI: 255.3-500.6), 68.8 (95%CI: 39.7-119.2), 27.6 (95%CI: 15.5-49.1), 13.7 (95%CI: 10.7-17.4), 6.2 (95%CI: 3.1-12.3), 2.1 (95%CI: 1.0-4.1), 1.5, 0.8 (95%CI: 0.5-1.3) and 1.0 (95%CI: 0.6-1.6) against B.1, BA.1, BA.2 , BA.5.1, BQ.1, XBB, EG.5.1 and, JN.1.1, respectively. No complete loss of neutralization in this group was observed for variants B.1, BA.1, BA.2, and BA.5.1, while 3/19 samples were not neutralized for BQ.1, 9/19 for XBB.1.5, 16/19 for EG.5.1 and 12/19 for JN.1.1 (**Figure 1B**).

2.2. Neutralizing capacity of vaccinated individuals with breakthrough infection (hybrid immunity)

We investigated the impact of BA.1 breakthrough infection in individuals vaccinated with two doses of mRNA vaccine. Geometric mean $FRNT_{90}$ titers in the hybrid immunity group were higher than for vaccinated individuals. They were 865.4 (95%CI: 450.9-1661.0) against B.1, 416.5 (95%CI: 175.6-987.8) against BA.1, 56.7 (95%CI: 31.8-100.9) against BA.2, 65.4 (95%CI: 30.3-141.4) against BA.5.1, 17.4 (95%CI: 7.6-39.9) against BQ.1, 11.3 (95%CI: 5.1-24.9) against XBB.1.5, 8.4 (95%CI: 3.5-20.3) against EG.5.1 and 7.1 (95%CI: 3.6-14.0) against JN.1.1. Complete loss of neutralization was observed only for 1/11 sample each for XBB.1.5, EG.5.1 and JN.1.1 (**Figure 1C**).

For boosted individuals with BA.2 breakthrough infection, geometric mean $FRNT_{90}$ titers were 818.7 (95%CI: 541.5-1238.0) for B.1, 266.2 (95%CI: 134.2-528.0) for BA.1; 415.8 (95%CI: 250.5-690.1) for BA.2, 85.1 (95%CI: 44.7-161.9) for BA.5.1, 50.2 (95%CI: 24.2-104.0) for

BQ.1, 33.6 (95%CI: 16.7-67.4) for XBB.1.5, 7.4 (95%CI: 2.7-20.7) for EG.5.1 and 6.2

(95%CI: 2.8-13.5) for JN.1.1. No complete loss of neutralization in this group was observed

for variants B.1, BA.1, BA.2, BA.5.1, BQ.1 and XBB.1.5, while 2/11 samples were not

neutralized for EG.5.1 and 1/11 for JN.1.1 (**Figure 1D**).

Although titers for B.1 were highest in both groups, the second highest neutralization titers

were found against the infecting virus, *e.g.*: neutralization of BA.1 was higher in the BA.1-

infected group and neutralization of BA.2 was higher in the BA.2.-infected group. No

difference was seen between the groups in the neutralization titers for EG.5.1. and JN.1.1 which

were both comparably low independent of the infecting virus.

We then investigated the impact of breakthrough infections in individuals (n=11) who have

been infected with one of the Omicron XBB subvariants between March and June 2023. The

geometric mean $FRNT_{90}$ titers were 1984.0 (95%CI: 1109-3547) against B.1, 46.0 (95%CI:

18.2-116.4) against XBB.1.5, 31.8 (95%CI: 9.3-108.6) against EG.5.1 and 21.1 (95%CI: 9.1-

48.7) against JN.1.1. All sera were able to neutralize in this group (**Figure 2A**).

2.3. Neutralizing capacity in vaccinated individuals with breakthrough infections from two

antigenically different VOCs (multi-variant hybrid immunity) towards Omicron lineages

We investigated vaccinated individuals (n=18) who subsequently have been exposed to at least

two antigenically different variants through two independent infection episodes that included

a pre-Omicron VOC (Alpha, n=9 or Delta, n=9) and another infection episode with an Omicron

lineage. The geometric mean $FRNT_{90}$ titers were 226.2 (95%CI: 153.8-332.8) against B.1,

80.33 (95%CI: 57.3-112.6) against BA.1, 32.3 (95%CI: 20.5-50.7) against Omicron BA.2, 22.4

(95%CI: 17.7-28.4) against BA.5.1, 7.6 (95%CI: 3.6-16.1) against BQ.1, 8.1 (95%CI: 4.3-15.3)

against XBB.1.5, 2.4 (95%CI: 1.0-5.4) against EG.5.1 and 2.5 (95%CI: 1.2-4.8) against JN.1.1.

Loss of neutralization was observed for BQ.1, XBB.1.5, EG.5.1 and JN.1.1 in 2/18, 1/18, 7/18

and 6/18 samples, respectively (**Figure 2B**).

2.4. Neutralizing capacity from unvaccinated adults and children infected in early 2022

We also studied neutralization of Omicron variants in 24 sera (adults, n=12; children, n=12) of

unvaccinated individuals with a single infection between January and March 2022 (most likely

exposed to BA.1 or BA.2). Due to the limited volume of serum available for this group, we

only assessed neutralization towards Omicron variants BA.1, BQ.1, XBB.1.5, EG.5.1 and

JN.1.1 in this cohort. Geometric mean $FRNT_{90}$ titers in this group for adult individuals were

29.1 (95%CI: 13.0-65.1) against BA.1, 1.3 (95%CI: 0.6-2.6) against BQ.1; 0.8 (95%CI: 0.5-

1.4) against XBB.1.5 and none of the samples neutralized EG.5.1 and Omicron JN.1.1.

Complete loss of neutralization was observed for 7/12 samples for BQ.1, 8/12 samples for

XBB.1.5 and all samples for EG.5.1 and JN.1.1 (**Figure 3A**). For children, geometric mean

$FRNT_{90}$ titers were 43.3 (95%CI: 18.0-103.7) against BA.1, 2.3 (95%CI: 0.9-5.8) against BQ.1,

1.1 (95%CI: 0.5-2.4) against XBB.1.5, 1.2 (95%CI: 0.6-2.5) against EG.5.1 and 0.6 (95%CI:

0.4-0.8) against JN.1.1. Of note, complete loss of neutralization was observed for 5/12 samples

for BQ.1, 8/12 samples for XBB.1.5, 6/12 samples for EG.5.1 and 11/12 for JN.1.1 (**Figure

3B**).

2.5. Heatmap of neutralization data across different immunological backgrounds

To summarize the findings across the cohorts, we have displayed the fold change of geometric

mean $FRNT_{90}$ titers in comparison to the ancestral virus B.1 for all cohorts (**Figure 3**). Across

cohorts, a consecutive loss in neutralization was observed from B.1 to BA.1/BA.2. to BA.5.1

to BQ.1 to XBB.1.5. and to EG.5.1/JN.1. The effect was the strongest for sera from individuals

that were only vaccinated but never infected, and the least pronounced for individuals exposed

to more than one natural infection with different variants. The differences between EG.5.1 and

JN.1. were only subtle across groups, and JN.1.1 did not show an enhanced immune escape

compared to EG.5.1. In vaccinated but never infected individuals, there was even a tendency

for better neutralization of JN.1.1 compared to EG.5.1.

2. Replicative capacity and innate immune responses of Omicron sublineages in nasal and bronchial HAE

To understand if Omicron lineages differ in their ability to replicate, we infected polarized

HAE of nasal (3 donors) and bronchial (3 donors) origin at different temperatures (**Figure 1**)

with different Omicron lineages. Comparison across lineages overall revealed similar kinetics

and replication range, with a rapid increase in viral RNA, reaching peak viral loads at 24/48hpi

followed by a slight decline at 72 and 96hpi. An increase in viral RNA was observed at 24h for

almost all lineages under all conditions. In nasal HAE, virus replication was slightly lower in

the physiological conditions (33°C) than at 37°C, where the peak of replication, reached at 48h,

varied from 9.6 log10 RNAc/mL (for BA.2), to 10.9 log10RNA c/mL (for XBB). The most

recent subvariants XBB.1.5, EG.5.1 and JN.1.1 showed better replication efficacy (higher than

10.5 log10 RNAc/mL). At 37°C, the peak was reached at 1dpi, except for BA.2, BA.5.1 and

BQ.1. Similar patterns were found in bronchial HAE with comparable levels of replication at

both temperatures. BA.2 again showed the lowest level of replication (10.1 log10 RNAc/mL

at 33°C and 10.0 log10 RNAc/mL at 37°C). The three most recent subvariants all showed

earlier replication peaks with a stronger increase of viral RNA at 37°C.

To compare innate immune responses after infection between Omicron lineages, we studied

IFN-α and -β and IFN-λ interferon responses and the induction of downstream interferon-

stimulated genes 15 (ISG15) at the end of the infection experiments (96hpi) in the infected and

non-infected HAE cultures (**Figure 6**). IFN-α by all variant's induction was barely observed

(less than 1log increase versus non-infected cells) at 33°C in infected nasal and bronchial HAE

and at 37°C in only in bronchial tissues. For all IFNs and ISG15, the lowest and highest

inductions were found in infected nasal HAE at 33°C and bronchial HAE at 37°C, respectively.

Weak induction (less than 0.5 log log10FC) of IFN-α and -β and ISG15 was observed in

bronchial HAE at 37°C, except for the IFN-β induction by the recent subvariants XBB.1.5,

EG.5.1 and JN.1.1 (0.65, 0.48 and 0.84 log10 FC, respectively).

IFN-λ showed the highest level of induction with all variants (except BA.2 in nasal HAE at

33°C) and the most pronounced variability between subvariants. While low induction was

observed in BA.1-, BA.2- and BQ.1- infected nasal tissues at 33°C, XBB.1.5, EG.5.1 and

JN.1.1 induced higher induction levels (2.17, 1.57- and 1.92-log10 FC, respectively). Inversely,

the latter showed lower induction levels in bronchial HAE at 33°C (from 0.58 to 0.81 log10

FC versus from 1.19 to 2.34 log10 FC for BA.1, BA.2, BA.5.1 and BQ.1) and at 37°C from

0.76 to 0.95 $\log_{10}$ FC versus from 1.63 to 1.84 $\log_{10}$ FC for BA.1, BA.2, BA.5.1 and BQ.1).

Induction of the main entry host factors, ACE-2 and TMPRSS2, was not enhanced in all

conditions of Omicron lineages' infections (**Figure S2**).

Altogether, our data showed, regardless of inter-donor variability, comparable replicative

capacity in both tissue origins under both temperature conditions, although lower for BA.2 and

slightly better for the most recent subvariants in nasal HAE.

Omicron lineages showed low induction of host responses, with an overall only slight

difference between variants.

Discussion

With the emergence of Omicron in late 2021, the COVID-19 pandemic has entered a new

phase. It has previously been shown that Omicron, compared to earlier VOCs, could overcome

immunity from various exposures, including prior infections and/or vaccinations, and showed

a distinct phenotype in *ex-vivo* infections compared to previous variants [2]. In the continuous emergence of new Omicron lineages, intrinsic transmissibility and immune pressure are considered as the main drivers of viral evolution [24]. A range of studies have shown increasing escape from prior immunity for Omicron lineages, particularly those that are currently designated as VOIs such as XBB.1.5 like and BA.2.86.

Omicron BA.2.86, with its highly mutated spike carrying over 30 mutations compared to other Omicron lineages and a genetic distance that is comparable to that of the first Omicron lineages BA.1 to that of Delta, was initially suspected to have potentially enhanced immune escape properties [25]. Multiple neutralization studies showed similar or slightly diminished immunity evasion compared to other Omicron variants [10, 14, 26-33]. Despite its first detection in mid-2023, the prevalence of BA.2.86 remained low, possibly due to presumed lower viral fitness observed in cell culture studies and reduced pathogenicity in animal models compared to other Omicron lineages [14, 27, 34].

In line with clinical observations, previous studies including ours recently confirmed the faster but shorter replication of Omicron BA.1 compared to previous SARS-CoV-2 variants in nasal HAE [16]. We here confirmed this typical replication for BA.1 and extended this observation to more recent subvariants relative to Omicron, despite modest differences (especially with BA.2) in nasal and bronchial HAE models. A sustained phenotype of Omicron lineages has also been shown when looking at their intrinsic host response mainly involving IFN-λ. Even with an overall equal replicative capacity at upper and lower respiratory tract temperatures, as previously shown for Omicron but not the previous ancestral (B.1) and Alpha variants [35], little differences in IFN induction were observed at 37°C compared to 33°C. Higher replication efficiency in *in vitro* (Calu3) and *in vivo* (Balb/c mice) of BA.5, compared to BA.1, in lung tissues/cells have been reported [36].

More *ex-vivo* (explant) and *in vivo* (mice) studies suggested an association between the milder severity of Omicron lineages and their enhanced replication efficacy in the upper, compared to the lower, respiratory tract, in contrast to Delta variant [37-39]. One study found increased replicative capacity and infectivity of BA.5 in comparison to BA.1 and the ancestral virus in human nasal and airway organoids at 37°C [40]. In HAE, Zaderer *et al.* showed that, compared to the Deta variant, there was a decreased replication efficiency with Omicron subvariants BA.1, BA.2, BA.5 and BQ.1.1, and a superficial localization into the pseudostratified tissue with less pronounced anti-viral response [41]. The more recently emerging lineages, like the XBB, EG.5 and JN.1 benefit from an additional fitness advantage, as experimental data obtained with live virus in cell culture/primary cell cultures infection studies. We showed a small replication advantage and slightly higher IFN-λ response [19, 20] in nasal epithelia with these variants, in line with Planas *et al* [14].

In comparison with the parent lineage BA.2.86, JN.1 has an additional L455S substitution in the spike protein that was described to be associated with increased escape from humoral immunity as well as transmissibility [5, 11-15]. However, in contrast to its parental lineage BA.2.86, JN.1 rapidly outcompeted earlier variants in late 2023 and became domiant, associated with a wave of infections worldwide [11, 15]. Limited data on the immune escape assessment of JN.1 are available, but it demonstrated lower geometric mean neutralization titers and lower fold change values compared to earlier Omicron variants [15, 42].

The present study aimed at the assessment of emerging SARS-CoV-2 variants in an authentic virus neutralization assay against a panel of human sera and plasma. To add to the complexity of the underlying immunity, we used cohorts with immunity at both ends of the exposure spectrum, which reflects the complex situation in the population in the fifth year of the pandemic. Here we show that recently emerged SARS-CoV-2 Omicron variants, namely BQ.1, XBB.1.5, EG.5.1 and JN.1.1, display pronounced immune evasion to earlier Omicron variants,

but that JN.1.1, does not show additional immune escape compared to EG.5.1. We also showed

that earlier findings on hybrid immunity remain valid; specifically, hybrid immunity continues

to offer higher neutralization titers compared to monovalent vaccination or natural infection.

The use of authentic live virus isolates and $FRNT_{90}$ to assess the neutralization titers of a large

number of serum/plasma samples with a heterogeneous immunological background of the

population adds strength to our findings. Indeed, according to the reported data, results may be

different using a pseudovirus instead of a live virus, underscoring the importance of having

data with authentic live viruses. It should be noted that our study has some limitations. First,

we had a low number of sera that were available for the individual groups, especially for those

that were exposed to more than one variant. It is important to note that vaccine sera from

bivalent vaccines or updated vaccine formulations were not included in the analysis, which

could impact the comprehensiveness of the findings regarding vaccine efficacy against

emerging variants.

In summary, our data show that continuous assessment of newly evolving SARS-CoV-2, taking

the different groups within the population into account, remains crucial to understand viral

strategies to overcome existing immunity. In the case of JN.1, that showed a rapid global

increase but no enhanced immune escape, other factors than immune escape seem to be the

driving force behind this variant success. Collectively, this comparative study of the fitness and

the immune escape capacity of the most relevant SARS-CoV-2 Omicron lineages, using

pertinent cell models, authentic viruses and human specimens from immunized individuals,

highlights the role of both virus fitness and adaptive immune response pressure on the evolution

of Omicron lineages. It hence contributes to the better understanding of SARS-CoV-2

dynamics including its main driving forces as well as its phenotypical impact on viral

properties.

Acknowledgments

The authors acknowledge NIBSC for providing the VERO-E6-derived cells, Samuel Cordey for sequence analysis and Pascal Cherpillod for facilitating the work in the BSL-3 laboratory at HUG, staff of the laboratory of virology at the HUG for support, clinicians and technicians responsible for clinical cohorts, the population epidemiology of the HUG Unit for their help with participants' recruitment and sample collection and all the donors.

Declaration of Competing Interest

IE has received research funding for an unrelated research project (IIS) as well as speakers fees from Moderna. CSE received funding from Pfizer for an unrelated research project (IIS). All other authors: No conflict of interest.

Authorship contributions

M.Bek., and M.E.L performed the experiments and conducted the analysis; K.A., C.A., P.S.R, D.D. and M.Bel helped to carry out some experiments; M.Bek, K.H.F, C.S.E., M.E.Z, S.S., I.G., O.P., M.Bel., and L.K. conducted the clinical studies and/or helped with clinical sample collection; I.E., M.Bek designed and supervised the study; M.Bek., M.E.L and I.E wrote the main manuscript. All authors have contributed to the final version of the manuscript.

Funding

This work was supported by the HUG Private Foundation, the Pictet Foundation, the Swiss National Science Foundation (Nr. 215567) and by the European Union Horizon Europe program, under project "EU-Africa Concerted Action on SARS-CoV-2 Virus Variant and Immunological Surveillance" (CoVICIS, grant nr: 101046041). Views and opinions expressed

References

1. World Health Organization (WHO), 2024. Tracking SARS-CoV-2 variants: https://www.who.int/activities/tracking-SARS-CoV-2-variants

2. Bekliz M, Adea K, Vetter P, Eberhardt CS, Hosszu-Fellous K, Vu DL, Puhach O, Essaidi-Laziosi M, Waldvogel-Abramowski S, Stephan C *et al*: **Neutralization capacity of antibodies elicited through homologous or heterologous infection or vaccination against SARS-CoV-2 VOCs**. *Nat Commun* 2022, **13**(1):3840.

3. Simon-Loriere E, Schwartz O: **Towards SARS-CoV-2 serotypes?** *Nat Rev Microbiol* 2022, **20**(4):187-188.

4. Hoffmann M, Arora P, Nehlmeier I, Kempf A, Cossmann A, Schulz SR, Morillas Ramos G, Manthey LA, Jack HM, Behrens GMN *et al*: **Profound neutralization evasion and augmented host cell entry are hallmarks of the fast-spreading SARS-CoV-2 lineage XBB.1.5**. *Cell Mol Immunol* 2023, **20**(4):419-422.

5. Jeworowski LM, Muhlemann B, Walper F, Schmidt ML, Jansen J, Krumbholz A, Simon-Loriere E, Jones TC, Corman VM, Drosten C: **Humoral immune escape by current SARS-CoV-2 variants BA.2.86 and JN.1, December 2023**. *Euro Surveill* 2024, **29**(2).

6. Kurhade C, Zou J, Xia H, Liu M, Chang HC, Ren P, Xie X, Shi PY: **Low neutralization of SARS-CoV-2 Omicron BA.2.75.2, BQ.1.1 and XBB.1 by parental mRNA vaccine or a BA.5 bivalent booster**. *Nat Med* 2023, **29**(2):344-347.

7. Miller J, Hachmann NP, Collier AY, Lasrado N, Mazurek CR, Patio RC, Powers O, Surve N, Theiler J, Korber B *et al*: **Substantial Neutralization Escape by SARS-CoV-2 Omicron Variants BQ.1.1 and XBB.1**. *N Engl J Med* 2023, **388**(7):662-664.

8. Uraki R, Ito M, Kiso M, Yamayoshi S, Iwatsuki-Horimoto K, Furusawa Y, Sakai-Tagawa Y, Imai M, Koga M, Yamamoto S *et al*: **Antiviral and bivalent vaccine efficacy against an omicron XBB.1.5 isolate**. *Lancet Infect Dis* 2023, **23**(4):402-403.

9. Wang Q, Iketani S, Li Z, Liu L, Guo Y, Huang Y, Bowen AD, Liu M, Wang M, Yu J *et al*: **Alarming antibody evasion properties of rising SARS-CoV-2 BQ and XBB subvariants**. *Cell* 2023, **186**(2):279-286 e278.

10. Wang Q, Guo Y, Liu L, Schwanz LT, Li Z, Nair MS, Ho J, Zhang RM, Iketani S, Yu J *et al*: **Antigenicity and receptor affinity of SARS-CoV-2 BA.2.86 spike**. *Nature* 2023, **624**(7992):639-644.

11. Kaku Y, Okumura K, Padilla-Blanco M, Kosugi Y, Uriu K, Hinay AA, Jr., Chen L, Plianchaisuk A, Kobiyama K, Ishii KJ *et al*: **Virological characteristics of the SARS-CoV-2 JN.1 variant**. *Lancet Infect Dis* 2024, **24**(2):e82.

12. Najat Bdeir TL, Henrike Maaß, Stefan Schmelz, Henning Jacobsen, Kristin Metzdorf, Anne Cossmann, Metodi V. Stankov, Markus Hoffmann, Stefan Pöhlmann, Wulf Blankenfeldt, Alexandra Dopfner-Jablonka, Georg M.N. Behrens, Luka Čičin-Šain: **Reverse mutational scanning of spike BA.2.86 identifies the epitopes contributing to immune escape from polyclonal sera**. *medRxiv* 2024.

13. Naomi S. Coombes KRB, Yann Le Duff, Nassim Alami-Rahmouni, Kathryn A. Ryan, Sarah Kempster, Deborah Ferguson, Elizabeth R. Davies, Thomas M. Weldon, Eleanor S. Cross, Lauren Smith, Conner Norris, Karly Rai Rogers-Broadway, Kuiama Lewandowski, Samantha Treagus, Isobel Everall, Steven T. Pullan, Bassam Hallis, Sue Charlton, Yper Hall, Simon G. P. Funnell: **Evaluation of the cross reactivity of neutralising antibody response in vaccinated human and convalescent hamster sera against SARS-CoV-2 variants up to and including JN.1 using an authentic virus neutralisation assay**. *bioRxiv* 2024.

14. Planas D, Staropoli I, Michel V, Lemoine F, Donati F, Prot M, Porrot F, Guivel-Benhassine F, Jeyarajah B, Brisebarre A *et al*: **Distinct evolution of SARS-CoV-2 Omicron XBB and BA.2.86 lineages combining increased fitness and antibody evasion**. *bioRxiv* 2023.

15. Yang S, Yu Y, Xu Y, Jian F, Song W, Yisimayi A, Wang P, Wang J, Liu J, Yu L *et al*: **Fast evolution of SARS-CoV-2 BA.2.86 to JN.1 under heavy immune pressure**. *Lancet Infect Dis* 2024, **24**(2):e70-e72.

16. Essaidi-Laziosi M, Perez-Rodriguez FJ, Alvarez C, Sattonnet-Roche P, Torriani G, Bekliz M, Adea K, Lenk M, Suliman T, Preiser W *et al*: **Distinct phenotype of SARS-CoV-2 Omicron BA.1 in human primary cells but no increased host range in cell lines of putative mammalian reservoir species**. *Virus Res* 2024, **339**:199255.

17. Puhach O, Bellon M, Adea K, Bekliz M, Hosszu-Fellous K, Sattonnet P, Hulo N, Kaiser L, Eckerle I, Meyer B: **SARS-CoV-2 convalescence and hybrid immunity elicits mucosal immune responses**. *EBioMedicine* 2023, **98**:104893.

18. Pango lineages: Latest epidemiological lineages of SARS-CoV-2, 2024: https://cov-lineages.org

19. Essaidi-Laziosi M, Alvarez C, Puhach O, Sattonnet-Roche P, Torriani G, Tapparel C, Kaiser L, Eckerle I: **Sequential infections with rhinovirus and influenza modulate the replicative capacity of SARS-CoV-2 in the upper respiratory tract**. *Emerg Microbes Infect* 2022, **11**(1):412-423.

20. Essaidi-Laziosi M, Brito F, Benaoudia S, Royston L, Cagno V, Fernandes-Rocha M, Piuz I, Zdobnov E, Huang S, Constant S *et al*: **Propagation of respiratory viruses in human airway epithelia reveals persistent virus-specific signatures**. *J Allergy Clin Immunol* 2018, **141**(6):2074-2084.

21. SARS-CoV-2 genomic surveillance in Geneva, weekly update, January, 2022: https://www.hug.ch/sites/interhug/files/structures/laboratoire_de_virologie/documents/Rapport%20hebdomadaire%20variants%20GE/2022/sars-cov-2_variants_geneva_update_12.01.22.pdf

22. Cov-Spectrum collections, Switzerland: https://cov-spectrum.org/explore/Switzerland/AllSamples/Past6M

23. Puhach O, Adea K, Hulo N, Sattonnet P, Genecand C, Iten A, Jacquerioz F, Kaiser L, Vetter P, Eckerle I *et al*: **Infectious viral load in unvaccinated and vaccinated individuals infected with ancestral, Delta or Omicron SARS-CoV-2**. *Nat Med* 2022, **28**(7):1491-1500.

24. Markov PV, Ghafari M, Beer M, Lythgoe K, Simmonds P, Stilianakis NI, Katzourakis A: **The evolution of SARS-CoV-2**. *Nat Rev Microbiol* 2023, **21**(6):361-379.

25. World Health Organization (WHO); Initial Risk Evaluation of BA.2.86 and its sublineages, 2023: https://www.who.int/docs/default-source/coronaviruse/21112023_ba.2.86_ire.pdf

26. Hu Y, Zou J, Kurhade C, Deng X, Chang HC, Kim DK, Shi PY, Ren P, Xie X: **Less neutralization evasion of SARS-CoV-2 BA.2.86 than XBB sublineages and CH.1.1**. *Emerg Microbes Infect* 2023, **12**(2):2271089.

27. Khan K, Lustig G, Romer C, Reedoy K, Jule Z, Karim F, Ganga Y, Bernstein M, Baig Z, Jackson L *et al*: **Evolution and neutralization escape of the SARS-CoV-2 BA.2.86 subvariant**. *Nat Commun* 2023, **14**(1):8078.

28. Lasrado N, Collier AY, Hachmann NP, Miller J, Rowe M, Schonberg ED, Rodrigues SL, LaPiana A, Patio RC, Anand T *et al*: **Neutralization escape by SARS-CoV-2 Omicron subvariant BA.2.86**. *Vaccine* 2023, **41**(47):6904-6909.

29. Lassauniere R, Polacek C, Utko M, Sorensen KM, Baig S, Ellegaard K, Escobar-Herrera LA, Fomsgaard A, Spiess K, Gunalan V *et al*: **Virus isolation and neutralisation of SARS-CoV-2 variants BA.2.86 and EG.5.1**. *Lancet Infect Dis* 2023, **23**(12):e509-e510.

30. Qu P, Xu K, Faraone JN, Goodarzi N, Zheng YM, Carlin C, Bednash JS, Horowitz JC, Mallampalli RK, Saif LJ *et al*: **Immune Evasion, Infectivity, and Fusogenicity of SARS-CoV-2 Omicron BA.2.86 and FLip Variants**. *bioRxiv* 2023.

31. Sheward DJ, Yang Y, Westerberg M, Oling S, Muschiol S, Sato K, Peacock TP, Karlsson Hedestam GB, Albert J, Murrell B: **Sensitivity of the SARS-CoV-2 BA.2.86 variant to prevailing neutralising antibody responses**. *Lancet Infect Dis* 2023, **23**(11):e462-e463.

32.   Uriu K, Ito J, Kosugi Y, Tanaka YL, Mugita Y, Guo Z, Hinay AA, Jr., Putri O, Kim Y, Shimizu R *et al*: **Transmissibility, infectivity, and immune evasion of the SARS-CoV-2 BA.2.86 variant**. *Lancet Infect Dis* 2023, **23**(11):e460-e461.

33.   Yang S, Yu Y, Jian F, Song W, Yisimayi A, Chen X, Xu Y, Wang P, Wang J, Yu L *et al*: **Antigenicity and infectivity characterisation of SARS-CoV-2 BA.2.86**. *Lancet Infect Dis* 2023, **23**(11):e457-e459.

34.   Tamura T, Mizuma K, Nasser H, Deguchi S, Padilla-Blanco M, Oda Y, Uriu K, Tolentino JEM, Tsujino S, Suzuki R *et al*: **Virological characteristics of the SARS-CoV-2 BA.2.86 variant**. *Cell Host Microbe* 2024.

35.   Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, Bleicker T, Brunink S, Schneider J, Schmidt ML *et al*: **Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR**. *Euro Surveill* 2020, **25**(3).

36.   Hoffmann M, Wong LR, Arora P, Zhang L, Rocha C, Odle A, Nehlmeier I, Kempf A, Richter A, Halwe NJ *et al*: **Omicron subvariant BA.5 efficiently infects lung cells**. *Nat Commun* 2023, **14**(1):3500.

37.   Hui KPY, Ho JCW, Cheung MC, Ng KC, Ching RHH, Lai KL, Kam TT, Gu H, Sit KY, Hsin MKY *et al*: **SARS-CoV-2 Omicron variant replication in human bronchus and lung ex vivo**. *Nature* 2022, **603**(7902):715-720.

38.   Hui KPY, Ng KC, Ho JCW, Yeung HW, Ching RHH, Gu H, Chung JCK, Chow VLY, Sit KY, Hsin MKY *et al*: **Replication of SARS-CoV-2 Omicron BA.2 variant in ex vivo cultures of the human upper and lower respiratory tract**. *EBioMedicine* 2022, **83**:104232.

39.   Shuai H, Chan JF, Hu B, Chai Y, Yoon C, Liu H, Liu Y, Shi J, Zhu T, Hu JC *et al*: **The viral fitness and intrinsic pathogenicity of dominant SARS-CoV-2 Omicron sublineages BA.1, BA.2, and BA.5**. *EBioMedicine* 2023, **95**:104753.

40.   Li C, Huang J, Yu Y, Wan Z, Chiu MC, Liu X, Zhang S, Cai JP, Chu H, Li G *et al*: **Human airway and nasal organoids reveal escalating replicative fitness of SARS-CoV-2 emerging variants**. *Proc Natl Acad Sci U S A* 2023, **120**(17):e2300376120.

41.   Zaderer V, Abd El Halim H, Wyremblewsky AL, Lupoli G, Dachert C, Muenchhoff M, Graf A, Blum H, Lass-Florl C, Keppler OT *et al*: **Omicron subvariants illustrate reduced respiratory tissue penetration, cell damage and inflammatory responses in human airway epithelia**. *Front Immunol* 2023, **14**:1258268.

42.   Wang Q, Guo Y, Bowen A, Mellis IA, Valdez R, Gherasim C, Gordon A, Liu L, Ho DD: **XBB.1.5 monovalent mRNA vaccine booster elicits robust neutralizing antibodies against XBB subvariants and JN.1**. *Cell Host Microbe* 2024, **32**(3):315-321 e313.

Table 1. Characteristic of vaccinated individuals' samples.

Vaccine	Number of individuals	Sample type	Gender *(M/F)*	Age *mean*	WPLV *Mean weeks (range)*	Date of last vaccination
2x mRNA vaccine	14	Plasma	2/12	51 (36-62)	4 (4-5)	March-May 2021
3x mRNA vaccine	19	Serum	9/10	43 (26-63)	13 (3-22)	November 2021-January 2022

552 **Table 2.** Characteristic of hybrid immunity (breakthrough with BA.1 or BA.2) individuals' samples.

Vaccination/infection status	Number of individuals	Sample type	Gender *(M/F)*	Age *mean*	WPLV *Mean weeks (range)*	WPLP *Mean weeks (range)*	Interval vaccination – infection *Mean weeks (range)*	Variant Identification[1]
2x mRNA vaccine & BA.1 breakthrough	11	Serum	5/6	39 (25-55)	34 (12-52)	15 (2-15)	27 (8-41)	Sequencing (n=11)
3x mRNA vaccine & BA.2 breakthrough	11	Serum	5/6	39 (25-62)	22 (15-35)	7 (3-10)	15 (5-32)	Sequencing (n=11)

556 **Table 3.** Characteristic of hybrid immunity (breakthrough with XBB) individuals' samples.

Vaccination[2]/infection status[3]	Number of individuals	Sample type	Gender (M/F)	Age mean	Number of vaccinations	WPLV Mean weeks (range)	WPLP Mean weeks (range)	Variant[1] identification
Vaccination & XBB breakthrough infection	11	Serum	1/10	44 (24-61)	2, 3 or 4	73 (36-93)	10 (2-19)	Sequencing (n=11)

Table 4. Characteristic of multi-VOC infected individuals' samples.

Vaccination[4]/infection status	Number of individuals	Sample type	Gender (M/F)	Age mean	WPLV Mean weeks (range)	WPLD Mean weeks (range)	Interval vaccination – last infection Mean weeks (range)	Variant identification[1]	
								Alpha or Delta	Omicron
Vaccination + Alpha & Omicron breakthrough	9	Serum	5/4	41 (25-50)	42 (30-64)	17 (10-29)	25 (12-44)	Sequencing (n=3)	Sequencing (n=5)
Vaccination + Delta & Omicron breakthrough	9	Serum	3/6	40 (29-51)	53 (27-87)	21 (4-41)	31 (18-55)	Sequencing (n=6)	Sequencing (n=1)

565 **Table 5.** Characteristic of convalescent children and adults' individuals' samples.

Infecting virus[5]	Number of individuals	Sample type	Gender (M/F)	Age mean	WPLD Mean days (range)	Infection period	Variant Identification[1]
Omicron in adults	12	Serum	2/10	40 (28-46)	15 (9-21)	January-March 2022	Extrapolated (n=12)
Omicron in children	12	Serum	6/6	5 (2-7)	17 (12-21)	January-March 2022	Extrapolated (n=12)

WPLV weeks post last dose of vaccine; *WPLP* weeks post last positive RT-PCR; *WPLD* weeks post last positive diagnosis;

[1]The infecting variant of each episode was either determined by sequencing of the diagnostic samples or extrapolated according to the dominance of the variant
at that time based on the data from our Swiss national genomic surveillance (**Figure S1**).

[2]Vaccination consisted of either 2 (n=2), 3 (n=7) or 4 (n=2) doses of BNT162b2 or mRNA-1273 vaccine. One of the individuals vaccinated with 4 doses of
mRNA had received a bivalent vaccine.

[3]Infection consisted of either 1 (n=7) or 2 (n=4) infection including one of the Omicron XBB variants infection.

[4]Vaccination consisted of either 1 (n=6), 2 (n=10) or 3 (n=2) doses of BNT162b2 or mRNA-1273 vaccine.

[5]Individuals were infected between January and March 2022 when the predominant circulation of Omicron BA.1 and BA.2 were taking place (<90%).

Figures

Figure 1. Neutralization in vaccine and hybrid immunity-derived blood specimens against eight authentic isolates of SARS-CoV-2 variants (B.1 and Omicron lineages including BA.1, BA.2, BA.5.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1). Bars represent geometric mean titers (GMT) of 90% reduction endpoint titers ($FRNT_{90}$) with 95% confidence interval. **A–D** Cohorts of individuals with **A)** double-dose mRNA vaccination (n=14), **B)** boosted mRNA vaccination (n=19), **C)** BA.1 breakthrough infection following double mRNA vaccination (n=11) and **D)** BA.2 breakthrough infection following 3 mRNA vaccination (n=11). Coloured numbers below each bar represent the number of specimens with complete loss of neutralization ($FRNT_{90}$ titer < 1). Repeated measures one-way ANOVA with Dunnett's multiple comparisons test using $\log_{10}$ transformed $FRNT_{90}$ titers was performed to analyze the statistical significance.

Figure 2. A) Neutralization in XBB exposure-derived blood specimens against four authentic isolates of SARS-CoV-2 variants (B.1 and Omicron lineages including XBB.1.5, EG.5.1 and JN.1.1). Cohort of specimens after XBB-derived infections with one of Omicron XBB variant. Bars represent geometric mean titers (GMT) of 90% reduction endpoint titers ($FRNT_{90}$) with 95% confidence interval. Coloured numbers below each bar represent the number of specimens with complete loss of neutralization ($FRNT_{90}$ titer < 1). Repeated measures one-way ANOVA with Dunnett's multiple comparisons test using $\log_{10}$ transformed $FRNT_{90}$ titers was performed to analyze the statistical significance. **B)** Neutralization in hybrid immunity-derived blood specimens against eight authentic isolates of SARS-CoV-2 variants (B.1 and Omicron lineages including BA.1, BA.2, BA.5.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1). Bars represent geometric mean titers (GMT) of 90% reduction endpoint titers ($FRNT_{90}$) with 95% confidence interval. Cohort of vaccinated individuals with dual SARS-CoV-2 infections (e.g. Alpha and Omicron (n=9) or Delta and Omicron (n=9)). Coloured numbers below each bar represent the number of specimens with complete loss of neutralization ($FRNT_{90}$ titer < 1). Repeated measures one-way ANOVA with Dunnett's multiple comparisons test using $\log_{10}$ transformed $FRNT_{90}$ titers was performed to analyze the statistical significance.

Figure 3. Neutralization in infection-derived blood specimens against five authentic isolates of SARS-CoV-2 variants (omicron lineages including BA.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1). Bars represent geometric mean titers (GMT) of 90% reduction endpoint titers ($FRNT_{90}$) with 95% confidence interval. Cohorts of convalescent specimens that are derived from **A)** unvaccinated adult individuals (n=12) and **B)** unvaccinated children individuals

(n=12) with confirmed SARS-CoV-2 infection in early 2022 (thus, probably Omicron BA.1 or

BA.2). Coloured numbers below each bar represent the number of specimens with complete

loss of neutralization ($FRNT_{90}$ titer < 1). Repeated measures one-way ANOVA with Dunnett's

multiple comparisons test using log_{10} transformed $FRNT_{90}$ titers was performed to analyze the

statistical significance.

Figure 4. Heatmap of fold-reduction in neutralization based on $FRNT_{90}$ data. Values of fold-

reduction in neutralization ($FRNT_{90}$) of B.1 and Omicron sublineages including BA.1, BA.2,

BA.5.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1 are presented as heat maps with lighter colors

implying greater changes. The immune sera/plasma were organized into cohorts of individuals

with double-dose mRNA vaccination (n=14), boosted individuals with three doses of mRNA

vaccine (n=19), BA.1 breakthrough infection of double-vaccinated individuals (n=11), BA.2

breakthrough infection individuals following 3 doses of mRNA vaccine (n=11), vaccinated

individual with XBB breakthrough infection (n=11) and vaccinated Individuals with dual

SARS-CoV-2 Infections (n=18).

Figure 5. Replication of SARS-CoV-2 Omicron lineages in HAE**.** Nasal (**A** and **B)** and

Bronchial (**C** and **D**) HAE were infected with SARS-CoV-2 Omicron lineages BA.1, BA.2,

BA.5.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1 at 33°C (**A** and **C**) and 37°C (**B** and **D**). Viral

replication was assessed by the quantification of viral RNA from apical washes collected at

3hpi (baseline), 24hpi, 48hpi, 72hpi and 96hpi. For each cell origin, HAE from 3 donors have

been tested. Data are expressed as mean of the log of viral RNA copies/mL (log_{10} RNA/mL)

and SEM.

Figure 6. Induction of HAE intrinsic host response during by SARS-CoV-2 Omicron

lineages

Nasal and Bronchial HAE infected with SARS-CoV-2 Omicron lineages BA.1, BA.2, BA.5.1,

BQ.1, XBB.1.5, EG.5.1 and JN.1.1 at 33°C (the same shown **Figure 1**) were lysed at 96hpi.

Induction of IFN-α (A) IFN-β (B), IFN-λ (C) and ISG15 (D) was assessed by semi-quantitative

real time RT-PCR using intracellular RNA and expressed in fold change relative to non-

infected and normalized to RNAseP. Data are represented as mean and SEM (n=3 donors), as

for each cell origin.

Figure S1. Trend of SARS-CoV-2 variants in Switzerland over time. From December 2020 until September 2023: from left to right, Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617), BA.1, BA.2, BA.5, BQ.1, XBB.1.5, EG.5.1 and JN.1.1 SARS-CoV-2 variants (https://cov-spectrum.org/explore/Switzerland/AllSamples/Past6M).

Figure S2. Induction entry host factor in HAE during infection by SARS-CoV-2 Omicron lineages

Nasal and Bronchial HAE infected with SARS-CoV-2 Omicron lineages BA.1, BA.2, BA.5.1, BQ.1, XBB.1.5, EG.5.1 and JN.1.1. at 33°C (as in **Figures 5** and **6**) were lysed at 96hpi. Induction of host ACE-2 receptor (A) and TMPRSS2 protease (B), involved in SARS-CoV-2 entry during infection, was assessed by semi quantitative real time PCR using total RNA from the cell and expressed in fold change relative to non-infected and normalized to RNAseP. Data are represented as mean and SEM (n=3 epithelia from 3 different donors tested for each cell origin, nasal/bronchial).

Figure 1

Figure 2

A

XBB

	<0,0001
	<0,0001
	<0,0001

FRNT$_{90}$ titer

0/11 0/11 0/11 0/11

B.1 XBB.1.5 EG.5.1 JN.1.1

1984 46.0 31.8 21.1

FRNT$_{90}$ titer

B.1 XBB.1.5 EG.5.1 JN.1.1

B

Alpha or Delta   Omicron

<0,0001
<0,0001
<0,0001
<0,0001
<0,0001
<0,0001
<0,0001

FRNT$_{90}$ titer

0/18 0/18 0/18 0/18 2/18 1/18 7/18 6/18

B.1 BA.1 BA.2 BA.5.1 BQ.1 XBB.1.5 EG.5.1 JN.1.1

226.1 80.3 32.3 22.4 7.6 8.1 2.4 2.5

FRNT$_{90}$ titer

B.1 BA.1 BA.2 BA.5.1 BQ.1 XBB.1.5 EG.5.1 JN.1.1

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Sequences over time in Switzerland

Plot by cov-spectrum.org, enabled by data from GISAID

Figure S2

A

ACE-2 4dpi

B

TMPRSS2 4dpi