**Locomotion-dependent auditory gating to the parietal cortex mediates flexible multisensory decisions**

Ilsong Choi[1], Seung-Hee Lee[1,2*]

[1]Center for Synaptic Brain Dysfunctions, IBS, Daejeon, 34141, Republic of Korea

[2]Department of Biological Sciences, KAIST, Daejeon, 34141, Republic of Korea

*Correspondence and requests for materials should be addressed to S.-H.L. (e-mail: shlee1@kaist.ac.kr).

**Abstract**

Decision-making in mammals is fundamentally based on the integration of several senses. When different sensory inputs guide conflicting outcomes, animals make decisions following the prepotent sensory modality. However, the neural basis of state-dependent flexibility in sensory dominance and multisensory decisions remains obscure. Here, we found that locomotion switched auditory-dominant decisions to visual-dominant ones in mice with audiovisual conflicts. When mice made auditory-dominant decisions, the visual representation in the posterior parietal cortex (PPC) was weakened. Prolonged locomotion restored visual representation in the PPC and induced visually dominant decisions by inhibiting the auditory cortical neurons projecting to the PPC (AC_{PPC}). In contrast, it had no effect on the AC projection to the striatum, which mediated auditory decisions. Locomotion activated the AC-projecting secondary motor cortex neurons, which preferentially inhibited the AC_{PPC} neurons. Our findings demonstrate that animals make flexible multisensory decisions by gating auditory afferents to the association cortex depending on locomotor conditions.

**Main**

Multisensory integration in the brain, which combines neural information from different sensory modalities, is essential for goal-directed perceptual decisions in an ever-changing environment[1]. Animals can make faster[2,3] and more accurate[4,5] perceptual decisions when they integrate congruent (coherent) multisensory stimuli. On the other hand, animals frequently prioritize one modality over the other in response to conflicting multisensory stimuli and show sensory dominance in their perceptual decisions[6-13]. For instance, when head-fixed mice faced conflicting audiovisual cues (one instructed 'Go' responses while the other instructed 'No-go' responses), they often made decisions following auditory cues[12,13]. This sensory dominance in multisensory decisions cannot be fully explained by a simple additive integration of individual stimuli as explained by a classic Bayesian model[14-16]. Rather, it requires under-weighting one modality more than the other before the integration[10,11,17], potentially via crossmodal inhibition[12,18,19]. Interestingly, the prepotent modality in multisensory decisions not only varies individually but also switches flexibly from trial to trial[10,11,20,21]. Neural underpinnings of this flexibility in multisensory decisions, however, remain largely unexplored.

Perceptual decision-making occurs with sensory representation associated with motor actions across cortical areas[22,23]. The sensory representation is not stable and fluctuates depending on the behavioral states of the animals[24,25]. One key factor affecting the cortical representation of sensory stimuli is the locomotion. Locomotion increases arousal, which amplifies visual gain in the visual cortex (VC)[26-34] but suppresses auditory representations in the auditory cortex (AC)[35,36]. Furthermore, locomotion generates corollary discharge signals copied from motor commands, which also suppress auditory responses in the AC[37,38]. As a result of these representational changes, mice showed better detection of mild visual cues[27] and reduced detection of weak auditory cues[35,39]. Despite this, animals still can identify salient

auditory cues well and perform auditory-dependent perceptual tasks without disruption even

during locomotion[40,41]. These reports suggest that, even during locomotion, animals retain

salient auditory information in the brain, enabling them to make appropriate auditory decisions.

In a multisensory environment containing both visual and auditory stimuli, it is more

challenging to determine how locomotion influences the multisensory representation and

decision-making processes. Compared to head-fixed mice, freely roaming mice exhibited no

differences in unisensory decisions but only shifted prepotent modality from audition to vision

during multisensory decisions[13]. However, it is unclear how locomotion modulates neural

circuits representing multisensory stimuli in the brain to alter behavioral decisions in mice.

One of the key brain areas that mediate perceptual decision-making is the posterior

parietal cortex (PPC)[42-44]. The PPC encodes various task-relevant activities during visual

decision-making tasks, conveying information of the category of stimuli[45-48], accumulating

sensory evidence[49-51], history of the task variables[48,52], animal's choices[45,48,49,53], and body

movements[54]. Furthermore, inactivation of the PPC resulted in a substantial decline in an

animal's ability to make visual decisions[45,49,50]. Similar to the visual acitivites, the PPC also

showed task-relevant activities during auditory decision-making tasks[48,50,55,56]. Its inactivation,

however, did not impair perceptual decisions based on simple auditory feature detection or

discrimiation[50,51,56], suggesting that the PPC may work for auditory-dependent decision-

making in more complicated context[57,58]. The PPC receives converging inputs from visual,

auditory, and somatosensory cortices[59-61] and responds to multisensory stimuli in humans[62],

primates[19,63] and rodents[64,65]. However, its role in multisensory decisions is still unclear. PPC

inactivation altered multisensory decisions under audiovisual conflicts[12] (but see also[16]), but it

did not affect multisensory decisions in coherent conditions[16,50]. These reports suggest that the

PPC in multisensory decisions may depend on the task paradigms or multisensory contexts that

the animal encounters.

To understand how the PPC is involved in multisensory decisions in animals making

perceptual decisions across auditory, visual, and congruent or incongruent audiovisual stimuli,

at different behavioral states, we used *in vivo* calcium imaging and optogenetic manipulation

techniques in mice performing an audiovisual discrimination task on the treadmill. We

discovered that mice flexibly switched the dominant modality from audition to vision during

locomotion. The PPC neurons discriminated visual inputs better than auditory inputs, and the

visual representation was suppressed when animals made auditory-dominant decisions in

audiovisual conflicts. During locomotion, the AC output to the PPC (AC$_{PPC}$) was selectively

suppressed, while the AC output to the striatum was maintained. This circuit-specific

modulation preserved the visual representation in the PPC and allowed animals to make

visually dominant multisensory decisions when animals confronted conflicting audiovisual

stimuli during locomotion. The locomotion-induced suppression of AC$_{PPC}$ neurons was

mediated by top-down inputs from the secondary motor cortex (M2) to the AC. Our data

demonstrate a neural mechanism of flexible multisensory decisions in the face of crossmodal

conflicts, which require gating in the auditory-to-parietal circuit according to the animal's

behavioral states.

## Results

### Locomotion enhances visually dominant decisions in mice under audiovisual conflicts

To examine state-dependent multisensory decisions in mice, we trained head-fixed mice to discriminate auditory and visual stimuli and make Go/No-go decisions on a non-motorized treadmill (Fig. 1a,b; 10 kHz vs. 5 kHz pure-tones: auditory Go ($A_G$) vs. auditory No-go ($A_{NG}$); vertically vs. horizontally drifting gratings: visual Go ($V_G$) vs. visual No-go ($V_{NG}$)). After mice became experts in discriminating both auditory and visual stimuli at similar correct rates in the stationary (S) state (Fig. 1c), we compared their performance in the moving (M) state (Fig. 1d). During the test, either congruent ($A_G V_G$ or $A_{NG} V_{NG}$) or incongruent ($A_G V_{NG}$ or $A_{NG} V_G$) audiovisual stimuli were randomly presented in a quarter of trials to examine their licking decisions in such multisensory context (Fig. 1e). In the incongruent trials, mice made either auditory-dominant or visual-dominant decisions (Fig. 1e; Methods). Head-fixed mice showed better performance of discriminating congruent audiovisual stimuli compared to auditory- or visual-only stimuli in stationary conditions (Fig. 1f). In moving sessions, the same mice discriminated the unisensory (auditory or visual) stimuli better, often hit the maximum performances, and therefore showed no further improvement in the congruent trials (Fig. 1f). As reported previously[12], stationary mice showed auditory-dominant decisions in response to the incongruent audiovisual stimuli (80.7% auditory dominance rate; Fig. 1g). Interestingly, the moving mice made considerably fewer auditory-dominant decisions (45.6% of the incongruent trials; Fig. 1g).

We further examined the effects of different conditions of locomotion on switching auditory dominance to visual dominance. First, we examined the effect of locomotion duration, as it has been shown that the locomotion-induced changes in cortical states depend on its duration[36,38,66,67]. We compared mice's decisions in the incongruent trials when the stimuli were presented right after locomotion (start-moving, SM) with those when stimuli were presented

after prolonged (> 3s) locomotion (keep-moving, KM). Interestingly, mice showed stronger

visual dominance in the keep-moving trials than in the start-moving trials (Fig. 1h). Once they

kept moving at a certain speed (> 3cm/s), the pre-stimulus speed did not further affect their

decisions in the incongruent trials (Extended Data Fig. 1a). These data indicate that the duration,

not the speed, of locomotion is important for switching the dominant modality from audition

to vision in mice making decisions under audiovisual conflicts. The unisensory trial types and

outcomes of previous trials did not affect the mice's decision in the incongruent trials with or

without locomotion (Extended Data Fig. 1b). The level of arousal measured by pupil size was

also unrelated to the effect of locomotion on changes in dominant decisions (Extended Data

Fig. 1c,d). Therefore, the locomotion duration prior to the stimuli, compared with any other

internal factors, is the most critical factor determining which modality becomes dominant.

**Locomotion reduces auditory representation in the PPC**

As the PPC activity is important for mice to make perceptual decisions by integrating

auditory and visual inputs[12], we next examined how the PPC excitatory neurons represent task-

relevant sensory information in mice performing the audiovisual discrimination task by *in vivo*

calcium imaging using a miniature microscope through a prism-combined gradient reflective

index (GRIN) lens (Fig. 2a). We first classified neuronal types based on their responses to

different sensory stimuli in the unisensory trials accompanied by correct decisions (Fig. 2b,c).

To avoid any potent influence of the outcome-related activities, we calculated fluorescence

changes during the stimulus period before the outcomes (0 ~ 1s). In the stationary sessions,

many of the PPC excitatory neurons were identified as visual-only (41.7%) or multisensory

responsive neurons (36.7%), but relatively small portion of them were auditory-only

responsive neurons (10.6%) (Fig. 2c). In the moving sessions, the proportion of visual neurons

increased (54.7%), and many neurons lost their auditory responses (Fig. 2c,d and Extended

Data Fig. 2a). When we compared the response amplitudes of all PPC neurons, the auditory

responses were significantly reduced during locomotion. Still, the visual responses were

similar between the stationary and the moving states (Fig. 2d). This finding indicates that

locomotion selectively degrades auditory, but not visual responses in the PPC of task-

performing mice.

During the task, mice often started to lick within the stimulus period, and the stimulus-

evoked activity of the PPC neurons might be mixed with the licking-evoked activities during

the stimulus presentation period. To decompose the PPC population activity into specific

activities representing sensory stimulus, licking decisions, stimulus-decision interactions, or

other independent factors, we used demixed principal component analysis (dPCA)[68]. We fed

neural data from the unisensory trials (i.e., $A_G$, $A_{NG}$, $V_G$, $V_{NG}$ with licking or no-licking

decisions) for the dPCA (Fig. 2e). We confirmed that the decomposed PPC activity in the

stimulus PC axes reliably represented each stimulus information regardless of licking actions

after the stimuli (Extended Data Fig. 2b). On the other hand, dPCA properly dissociated

decision-relevant activity in the PPC neurons in other axes (Extended Data Fig. 2c,d).

Interestingly, the stimulus axis explained the PPC population activity much more than any

other axes in both stationary and moving conditions (Fig. 2f and Extended Data Fig. 2e),

indicating that the PPC neurons represent sensory information the most strongly during the

task.

We next examined how the PPC activity represented auditory and visual Go/No-go

stimuli in the stimulus subspace by comparing trial-averaged neuronal trajectories of each

unisensory stimulus in the top three PCs (Fig. 2g; mean % explanation by top three stimulus

PCs: 62.8%). First, we quantitatively measured the magnitude of sensory representations by

calculating the projection distance of each trajectory during the stimulus presentation period

(for 1 s after stimulus onset) (Fig. 2g; Methods). Auditory trajectories were much shorter than

visual trajectories and became even shorter as animals made longer locomotion (Fig. 2h). Next,

we calculated the Euclidean distance between a pair of trajectories to quantify similarities in

the representation of each stimulus type (Fig. 2i and Extended Data Fig. 2f). Auditory Go and

No-go trajectories were close to each other and less distinct during locomotion (Fig. 2i). In

contrast, visual Go and No-go trajectories were divergent and distinct from each other (Fig.

2i). The visual trajectories also differed from the auditory trajectories (Extended Data Fig. 2f).

Finally, to understand the trial-by-trial reliability of unisensory representation in the PPC, we

developed a classifier, which predicted the presented stimulus type in the current trial based on

the Euclidean distance between the current-trial trajectory and the trial-averaged trajectory

(Fig. 2j). Classification accuracies were significantly higher for the visual trials than the

auditory trials (Fig. 2k,l). Moreover, accuracies of auditory classification were significantly

lower in the moving sessions than in the stationary sessions (Fig. 2l). Collectively, the PPC

neurons processed task-relevant visual information more strongly than the auditory

information, showing robust visual but weak auditory Go/No-go discrimination. More

importantly, locomotion further dampened the auditory representation in the PPC.

**Multisensory representation in the PPC determines the dominant modality for perceptual**

**decisions under audiovisual conflict**

We next examined how the PPC neurons represent multisensory information by

projecting their activity into the stimulus subspace during the multisensory trials (Fig. 3a,b;

Methods). The neural trajectories representing the congruent multisensory stimuli were very

close to the visual trajectories but highly distinct from the auditory trajectories in both

stationary and moving conditions (Fig. 3a,b). Euclidean distances, projection distances, and

classifier accuracies between the congruent Go and No-go trajectories were not different from

those between the visual Go and No-go trajectories regardless of locomotion conditions

(Extended Data Fig. 3a-d). These findings suggest that PPC neurons represent congruent

multisensory inputs in a similar way that they do visual stimuli.

By projecting PPC activity trajectories during the incongruent multisensory trials, we

found that PPC neurons represented the same stimulus condition ($A_{NG}V_G$ or $A_GV_{NG}$) differently

depending on the impending decisions of mice (Fig. 3c; auditory dominance in grey trajectories

vs. visual dominance in black trajectories). We first found that the projection and the Euclidean

distances of the trajectories for the auditory-dominant incongruent trials were smaller than

those for the visual-dominant trials (Extended Data Fig. 3e,f). Furthermore, the trajectories

representing the incongruent stimuli were nearly identical to those representing visual stimuli

when mice made visually dominant decisions (Fig. 3d and Extended Data Fig. 3g). When mice

made auditory-dominant decisions, however, the neural trajectories for the incongruent stimuli

diverged from the visual trajectories, becoming closer to but still separate from the auditory

trajectories (Fig. 3d and Extended Data Fig. 3g).

We next classified the incongruent trials individually by calculating the Euclidean

distance between the single-trial trajectory and the mean trajectories of the four types of

multisensory trials and found that the classification accuracy was significantly higher in the

stimulus subspace than in other subspaces (Extended Data Fig. 3h). Therefore, the sensory

representation in the PPC is the most critical factor determining the multisensory decisions of

mice in the audiovisual conflict during the task. We further developed another classifier by

setting the boundary of the visual trajectories based on their trial-by-trial variations and

classified the individual incongruent trajectories as visually dominant ones if they were within

the boundary of the visual trajectories (Fig. 3e). Using this, we successfully classified the

dominant modality across the incongruent trials (Fig. 3f). Collectively, if the PPC neurons

dominantly represented visual information in the incongruent audiovisual stimuli, mice showed

visually dominant decisions in the incongruent trials. Conversely, if the visually dominant

representation weakened in the PPC, mice made auditory-dominant decisions in response to

the incongruent stimuli.

To understand if the stimulus representation in the PPC changes with locomotion, we

calculated the divergence between the incongruent multisensory trajectory and the visual

trajectory at different locomotion states (Fig. 3g). The two trajectories were closer as the animal

made longer locomotion, and this was correlated with the decrease in auditory dominance rates

in the animal's multisensory decisions (Fig. 3g). In correlation with such changes, visually

responsive neurons showed stronger suppression in stationary conditions than in moving

conditions, but only in the incongruent trials (Fig. 3h). In contrast, auditory-responsive neurons

showed significantly low auditory responses during locomotion across all types of trials (Fig.

3h). Therefore, locomotion increased the likelihood of maintaining the visual representation

while reducing the auditory representation of the incongruent multisensory stimuli in the PPC

and caused mice to make visually dominant multisensory decisions.

**Locomotion selectively suppresses auditory cortical outputs to the parietal cortex but not**

**to the striatum**

The robust reduction of auditory representation in the PPC during locomotion might

be due to direct suppression of the auditory neurons in the PPC or indirect suppression of the

AC output neurons that project to the PPC (i.e. AC$_{PPC}$ neurons). To examine which case it was,

we measured the spontaneous activity changes in the PPC auditory-responsive neurons during

locomotion. Neither the locomotion onset nor offset induced significant suppression in their

activities (Extended Data Fig. 4a-d). Therefore, the suppression of the auditory responses in

the PPC during locomotion is unlikely to result from direct inhibition of the PPC auditory

neurons. Instead, it might be due to the reduction of bottom-up auditory inputs to the PPC.

To examine this, we first traced the $AC_{PPC}$ neurons by injecting the retrograde adeno-

associated viruses (AAVrg) encoding tdTomato into the PPC (Fig. 4a). We co-traced the AC

neurons that project to the posterior striatum ($AC_{STR}$), which are known to play a critical role

in the auditory decision-making of mice[69-72], by injecting the AAVrg encoding eGFP into the

striatum (Fig. 4a). $AC_{PPC}$ neurons were mostly found in the anterior and dorsal AC (AuD),

whereas $AC_{STR}$ neurons were found across the anterior-to-posterior and dorsal-to-ventral AC

(Fig. 4b,c). Only 7% of the traced neurons in the AC were co-labeled, suggesting that the $AC_{PPC}$

and the $AC_{STR}$ neurons are distinct and non-overlapping neuronal populations in the AC (Fig.

4d).

We next measured the activity of $AC_{PPC}$ and $AC_{STR}$ neurons in mice running on the

treadmill (Fig. 4e,k). When animals started to move, the spontaneous activity of $AC_{PPC}$ neurons

was significantly reduced, and their calcium activities and locomotion onset speeds were

negatively correlated (Fig. 4f). $AC_{PPC}$ neurons were liberated from the inhibition at locomotion

offsets (Fig. 4g). Unlike the $AC_{PPC}$ neurons, $AC_{STR}$ neurons did not exhibit any activity changes

at the onset or offset of locomotion (Fig. 4l,m). These results demonstrate that locomotion

suppressed $AC_{PPC}$ neurons directly and selectively without inhibiting $AC_{STR}$ neurons.

**Locomotion reduces task-relevant auditory representation in the $AC_{PPC}$ neurons**

We next examined the stimulus-evoked activities of $AC_{PPC}$ and $AC_{STR}$ neurons during

the task. Both $AC_{PPC}$ and $AC_{STR}$ neurons showed significant auditory responses but no

profound visual responses during the task (Extended Data Fig. 5a-c and Extended Data Fig.

6a-c). Prolonged locomotion suppressed the auditory responses of the $AC_{PPC}$ neurons

(Extended Data Fig. 5c). In the incongruent trials, $AC_{PPC}$ neurons showed strong responses

when animals made auditory-dominant decisions but significantly weak responses during

visually-dominant decisions (Fig. 4h,i). Locomotion-induced suppression of the $AC_{PPC}$

neurons was correlated with reduced auditory dominance in multisensory decisions of mice

(Fig. 4j). However, there was no such suppression in the AC$_{STR}$ neurons during locomotion,

and the population of AC$_{STR}$ neurons showed auditory responses at similar levels across the

trials independent of locomotion conditions and dominant modalities (Fig. 4n-p and Extended

Data Fig. 6c).

We further analyzed AC population activity using the dPCA. As shown in the PPC

neurons, the stimulus axis explained the activities of both AC subpopulations more strongly

than other task-relevant axes (data not shown). During locomotion, the auditory trajectories

significantly shrank only in the AC$_{PPC}$ neurons (Extended Data Fig. 5d-f), but not in the AC$_{STR}$

neurons (Extended Data Fig. 6d-f). Furthermore, only AC$_{PPC}$ neurons showed decreased

classification accuracies for the auditory stimuli during locomotion (Extended Data Fig. 5g and

Extended Data Fig. 6g). During the incongruent trials, the response amplitudes and the neural

trajectories of the AC$_{PPC}$ neurons were significantly smaller in the visual-dominant trials than

in the auditory-dominant trials (Extended Data Fig. 5h-j). In contrast, AC$_{STR}$ neurons showed

consistent responses to the incongruent stimuli regardless of dominant modalities  (Extended

Data Fig. 6h-j). These data indicate that locomotion suppressed the AC$_{PPC}$ neurons more than

AC$_{STR}$ neurons, and this selective suppression switched mice to make visually dominant

decisions in the incongruent trials. In contrast to the vulnerability of the AC$_{PPC}$ neurons, AC$_{STR}$

neurons were unaffected by locomotion, suggesting that the AC$_{STR}$ neurons reliably encode

auditory information even during locomotion and are unrelated to multisensory decisions

influenced by the locomotion.

**Optogenetic double dissociation reveals distinct roles of the AC subpopulations in**

**perceptual decisions during the audiovisual discrimination task**

We wondered whether the two distinct AC outputs, $AC_{PPC}$ and $AC_{STR}$, play distinct

functional roles during the audiovisual discrimination task. We first examined whether the

activity in the $AC_{PPC}$ neurons was required for auditory dominance in multisensory decisions

of mice in the incongruent trials. We expressed halorhodopsin (eNpHR3.0) in $AC_{PPC}$ neurons

and optogenetically inhibited them in task-performing mice (Fig. 5a). Optogenetic inhibition

of the $AC_{PPC}$ neurons did not affect the mice's decisions in the auditory, visual, and congruent

audiovisual trials (Fig. 5b,d). However, it caused a significant reduction in auditory-dominant

decisions in stationary mice (Fig. 5c,e).

We next examined the role of $AC_{STR}$ neurons during the task by optogenetic inhibition

of those neurons (Fig. 5f). Similar to previous studies[69,70], inhibition of the $AC_{STR}$ neurons

significantly hampered auditory discrimination in stationary mice (Fig. 5g,i). At the same time,

it caused only a mild reduction of auditory-dominant decisions in the incongruent trials (Fig,

5h,j). Our findings indicate that the AC-to-STR circuit is important for perceptual decisions in

auditory-only conditions. In contrast, the AC-to-PPC circuit is necessary for animals to make

auditory-dominant decisions under audiovisual conflicts.

**Motor cortical projection to the AC suppresses $AC_{PPC}$ neurons during locomotion**

Previous research reported that the neural projection from the M2 to the AC suppressed

auditory responses of AC neurons[37,38]. Our data further showed that locomotion suppressed the

$AC_{PPC}$ neurons but not the $AC_{STR}$ neurons. To examine if the M2 mediates this selective

suppression of the $AC_{PPC}$ neurons, we first anatomically mapped the spread of fluorescence-

labeled M2 axons in the AC (Fig. 6a). Similar to the previous report[37], M2 axons primarily

innervated the superficial or deep layers of the AuD (Fig. 6b). When we compared the M2 axon

distribution with the distributions of the $AC_{PPC}$ and the $AC_{STR}$ neurons across AC areas, M2

axons were localized at a similar location where the $AC_{PPC}$ neurons were found (Fig. 6c). On

the other hand, majority number of the $AC_{STR}$ neurons did not overlap well with the M2 axons

(Fig. 6c). This anatomical feature of the overlap between the M2 axons and the $AC_{PPC}$ neurons

in the AuD suggests potential modulation of the $AC_{PPC}$ neurons by the M2 during locomotion.

We next investigated whether M2 neurons that project to the AC were activated during

locomotion. We measured the calcium activity of the AC-projecting M2 neurons ($M2_{AC}$) in

mice running on the treadmill (Fig. 6d). $M2_{AC}$ neurons showed robust activation at the

locomotion onset (Fig. 6e), and the calcium activities of these neurons were significantly higher

during the later period of locomotion than those right after the locomotion onset (Fig. 6f). This

result indicates that the M2 neurons copied and delivered the locomotion onset signal to the

AC. We then examined whether this M2 activity suppressed the $AC_{PPC}$ neurons directly without

affecting the $AC_{STR}$ neurons. Using the miniature fluorescence microscope conjugated with

optogenetic manipulation system (nVoke), we activated the M2 axon terminals in the AC while

imaging the calcium activity in the AC subpopulations (Fig. 6g). Optogenetic activation of M2

axons significantly dampened auditory responses of the $AC_{PPC}$ neurons (Fig. 6h,i). In contrast,

the $AC_{STR}$ neurons were unaffected by the M2 axon activation, showing consistent auditory

responses during the M2 axon activation (Fig. 6j,k). Lastly, we optogenetically inhibited the

M2 axon terminals in the AC in mice performing the task (Fig. 6l) and found that it did not

alter the animal's decisions in the unisensory or congruent trials (Fig. 6m). However, it

significantly enhanced auditory dominance in the incongruent trials during locomotion (Fig.

6n). These results strongly support the idea that locomotion raises activity of the M2-to-AC

circuit[38], and we further found that the locomotion-induced activation of this cortical circuit

selectively inhibits the AC outputs to the PPC. Consequently, PPC activity is dominated by

visual information during locomotion, leading to more visually dominant decisions in mice

facing audiovisual conflicts (Fig. 6o).

**Discussions**

In this study, we newly identified the neural mechanism for flexible multisensory decisions in mice under audiovisual conflicts. The auditory cortical outputs to the parietal cortex inhibited visual representation in the PPC and caused auditory-dominant decisions. These output neurons were suppressed by locomotion, and such suppression caused animals make more visually dominant decisions. However, locomotion did not suppress the auditory outputs to the STR, allowing animals to make precise auditory decisions even during locomotion. Our findings demonstrate that top-down projection from the M2 to the AC gates auditory outputs to the higher association cortex specifically and shapes flexible decisions under audiovisual conflicts without disrupting perceptual decisions in a simple auditory environment in behaving animals.

**Locomotion and multisensory decision-making**

As all animals naturally locomote while collecting sensory information in the environment to make decisions, understanding the role of locomotion in perceptual decision-making is ethologically critical. More recently, using small animals like mice, perceptual experiments for head-fixed mice on the treadmill were devised[31]. Using this system, previous studies have examined the effects of locomotion on the simple detection of a single type of sensory cue[27,35,39], but the effect of locomotion on the discrimination of different modality inputs and multisensory decision-making has never been tested. Our study designed a new behavioral paradigm to measure perceptual decision-making performances in the unisensory (visual or auditory) and multisensory (congruent or incongruent) conditions at different locomotion states within one session in the same animal. We found that locomotion can improve both visual and auditory perceptual decisions after discrimination of salient stimuli in the same animal. This might be because of the increased arousal during locomotion, which can enhance visual and auditory discriminability in the sensory cortices[34,73].

We further found that locomotion can switch multisensory decisions from auditory-

dominant to visual-dominant ones in mice. The effect of locomotion on multisensory

perception and decision has never been examined before. Our findings demonstrate that the

dominant modality can be flexibly switched from audition to vision depending on the

locomotion states. Sensory dominance in multisensory decisions can be beneficial for animals

to make rapid perceptual decisions[74], which is critical for the survival of animals in a

complicated world. Depending on the task design, specific sensory modality information

provides more reliable information than other modalities during multisensory integration[74]. A

more reliable sensory modality can provide a stronger effect on multisensory decisions in

animals performing the task. For example, visual information is more critical for animals to

discriminate spatial locations of the multisensory stimuli[9], while auditory information is more

important for temporal discrimination of the stimuli[75]. Our study suggests that the locomotion

condition of an animal can flexibly switch the appropriate modality even during the same task.

The determination of the dominant modality has been explained by Bayesian models, which

explain the optimal strategy of re-weighting different sensory inputs based on their reliability

(or variance)[10,11,14]. Our data further suggest that the locomotion generates a self-initiated

internal signal that can change the relative weights of the auditory inputs compared to the visual

inputs by lowering reliability of the auditory information in the parietal cortex, where the

dominant modality is determined.

**Locomotion-induced circuit-specific suppression of auditory outputs during multisensory**

**decisions**

Locomotion increases corollary discharge signals in the M2, potentially degrading

auditory representations in the AC[38,76]. We further found that locomotion-evoked suppression

of auditory responses only happened in the auditory-to-parietal projection, without impairing

auditory responses in the auditory-to-striatal projection (Fig. 4 and Extended Data Fig. 5,6).

Therefore, corollary discharge signals evoked by locomotion weaken a specific auditory flow

from the AC to the higher association cortex. This caused animals to rely less on auditory

stimuli for their decisions under crossmodal conflicts. Future studies are required to understand

how the top-down inhibitory synaptic connection from the M2 is only formed in the $AC_{PPC}$ but

not in the $AC_{STR}$ neurons.

Using circuit-specific calcium-imaging and optogenetic experiments, we found that

$AC_{PPC}$ neurons were mainly necessary for auditory-dominant decisions in the audiovisual

conflicts, but $AC_{STR}$ neurons were critical for auditory-based discrimination. The divided roles

of the AC outputs may allow the mammalian brain to take advantage of preserved auditory

processing during locomotion to make auditory-dependent decisions errorlessly while

switching dominant modality in multisensory decisions only when sensory inputs make

conflicts.

Similar to the locomotion effect on the auditory system, saccadic eye movement

suppresses visual representation in the lower visual areas of the animal[77,78]. However, we found

that locomotion did not weaken the visual representation in the PPC (Fig. 2), although some

individual neurons reshaped (either increased or decreased) their visual responsiveness during

locomotion. It is also known that locomotion evokes strong cholinergic activation, which

enhances visual representation in the VC[29]. Future studies are required to know whether

inhibitory modulation by the M2 activation during locomotion is specific in the AC and how

the locomotion-evoked sensory degradation is compensated by the enhanced arousal in the

visual system.

**Functional role of PPC in perceptual decision making**

Similar to the previous research[51], we found the PPC excitatory neurons represent

visual information more reliably than the auditory information during the task. Our data further

showed that strong auditory inputs to the PPC can disrupt the visual representation, facilitating

auditory-dominant multisensory decisions. Interestingly, the PPC excitatory neurons could not

discriminate auditory Go/No-go information during the auditory-based decisions, indicating

that the PPC itself is not involved in auditory decision-making. Rather, its function as a higher

visual area is integrating auditory inputs to determine the visual weights for the multisensory

decisions. We also found that the AC output to the STR, which is important for auditory

decision-making, showed stable auditory representation even in the visually dominant

decisions (Fig. 4 and Extended Data Fig. 6). Future studies are required to fully understand

whether the strong visual representation in the PPC is only required or whether any downstream

areas of the $AC_{STR}$ neurons need to be further suppressed to switch the auditory-dominant

decisions to the visually dominant ones.

Our data also indicate that the suppression of visual responses in the PPC only occurs

in the incongruent trials (not congruent trials) when animals make auditory-dominant decisions

(Fig. 3h). One potential explanation for this congruency-dependent crossmodal suppression

would be an antagonistic selectivity between Go and No-go stimuli in auditory-to-visual

inhibitory circuits. Supporting this idea, opponent inhibition between choice-selective neurons

was recently found in the PPC[79]. Our previous report showed that pavalbumin-expressing

(PV[+]) neurons in the PPC strongly received AC inputs and mediated auditory-dominant

behaviors[12]. Future studies are required to examine whether the selectivity of inhibitory

neurons emerges with learning in these inhibitory neurons in the PPC. Similar to the role of the

mouse PPC exerting crossmodal inhibition, the human prefrontal cotex[80] and the superior

temporal sulcus[17,81] also showed multisensory depression in resolution of conflicting

audiovisual speech stimuli. Therefore, multisensory depression might happen across

association areas when animals face multisensory conflicts, and our study demonstrates how

the brain flexibly controls the resolution of multisensory conflicts by gating bottom-up inputs.

Our task design used Go/No-go decisions of animals in each trial, which are binary

reports and hard to represent differences in the level of perceptual decisions as a continuum

across trials. Furthermore, the population of the PPC neurons did not show any significant

enhancement in the representation of the congruent audiovisual stimuli compared to the visual-

only stimuli. Therefore, other areas, instead of the PPC, may integrate auditory inputs

additively and represent congruent multisensory information more reliably to improve animals'

perceptual discriminability in the congruent trials. One potential downstream area might be the

frontal motor cortex, as the additive multisensory integration[16], as well as the auditory-based

decision-making, require M2 activity[82]. Future studies are required to determine whether these

circuits are further recruited beyond the PPC to make flexible multisensory decisions in mice.

**Acknowledgements**

We thank Dr. Alex C. Kwan for helpful discussions of imaging data analysis. We thank Dr. Jae-Hyun Kim for helping pupil data analysis.This work was supported by grants to S.-H.L. from the National Research Foundation of Korea funded by the Ministry of Science & ICT (NRF-2021R1A2C3012159) and the Institute for Basic Science (IBS-R002-A2).

**Author Contributions**

I.C. and S.-H.L. conceived and designed all experiments and wrote the manuscript. I.C. performed all experiments and analyzed the data.

**Competing interest declaration**

The authors declare no competing interests.

**Data availability**

All the representative data that support the findings will be publicly available on GitHub. Further requests for the data used in this study can be directed to the corresponding author (shlee1@kaist.ac.kr)

**Code availability**

All custom MATLAB codes to analyze and visualize the data will be publicly available on GitHub.

## Methods

### Animals and surgery

*Animals*

All experimental procedures were approved by the KAIST Institutional Animal Care and Use Committee (IACUC-18-100, IACUC-18-235). Both male and female mice (P45-P150) were used in this study. Animals were maintained *ad libitum* under light (8 am-8 pm) and dark cycle (8 pm-8 am) conditions and housed under single animal per cage conditions from the start until the last day of the experiment. We used C57BL/6J wild-type (WT) mice, B6.Cg-Tg(Camk2a-cre)T29-1Stl/J mice (CaMKIIα-Cre, stock no. 005359, Jackson Laboratory), and Ai162 mice (stock no. 031562, Jackson Laboratory).

*Common procedure of animal surgery*

Adult mice (P45-P80) were anesthetized with isoflurane (1.5%–2% in oxygen) and fixed their head on the stereotaxic apparatus. The body temperature was monitored and maintained at 37°C using a heating pad connected to a body temperature sensor. The scalp was alternately sterilized three times with 70% ethanol and povidone-iodine solution. After the sterilization of the scalp, surgical scissors was used to cut the scalp according to the midline (for virus injection) or remove a portion of the scalp (for the implantation of head plates, GRIN lens, and optic fiber). Periodsteum above the skull was removed, and povidone-iodine solution was used to sanitize the exposed skull and incision. Lastly, we cleaned the exposed skull with sterilized phosphate-buffered saline (PBS).

*Virus injection for retrograde tracing*

For the retrograde tracing of the AC neurons that project to the PPC and the STR, we injected 0.5 μl of retrograde AAV expressing tdTomato (AAVrg-CAG-tdTomato, Addgene)

and 0.5 µl of retrograde AAV expressing GFP (AAVrg-CAG-GFP, Addgene) into the target

areas respectively. After 2~3 weeks from the injections, we perfused mice to collect the brain

samples for histology.

*Head plate implantation surgery for mouse head-fixation*

To implant a stainless steel head-plate on the mouse head, three miniature stainless

steel screws (J.I. Morris) were implanted into the exposed skull using cyanoacrylate glue

(Loctite super glue 401, Henkel). A customized stainless steel head-plate was attached on the

skull with the cyanoacrylate glue, and the between the skull and head-plate was filled by dental

acrylic (Ortho-Jet, Lang Dental).

*Surgical procedures for in vivo calcium imaging*

To express GCaMP6s in the target neurons, we made a small craniotomy (diameter

~0.5 mm) above the targeted location. To conduct *in vivo* calcium imaging at the PPC, we

injected ~0.5 µl of Cre-recombinase dependent adeno-associated virus (AAV) expressing

GCaMP6s (AAV5-Syn-Flex-GCaMP6s-WPRE-SV40, UPENN) into a site 0.2mm posterior

from the right PPC (bregma -2.15mm, lateral -1.6mm, depth -0.35mm) of CaMKIIα-Cre mice

using a thin pulled glass pipette (20~25µm of tip diameter, item no. 504949, WPI) installed in

a Nanoliter 2010 injector (WPI) at a speed of 24nl/min. For *in vivo* calcium imaging of $AC_{PPC}$

neurons, we injected ~0.5 µl of retrograde AAV expressing GCaMP6s (AAVrg-hSyn1-

GCaMP6s-P2A-nls-dTomato, Addgene) into the right PPC (bregma -1.95mm, lateral -1.6mm,

depth -0.35mm) of the WT mice. For *in vivo* calcium imaging of $AC_{STR}$ neurons, we injected

~0.5 µl of retrograde AAV expressing GCaMP6s (AAVrg-hSyn1-GCaMP6s-P2A-nls-

dTomato, Addgene) into the right STR (bregma -1.5mm, lateral -3.45mm, depth -2.8mm) of

the WT mice. For *in vivo* calcium imaging of AC-projecting M2 neurons, we injected ~0.5 µl

of retrograde AAV expressing Cre-recombinase (AAVrg-hSyn-Cre, Addgene) into the right

AC (bregma -2.5mm, lateral 4.2mm, depth -0.8mm) of the Ai162 mice. For *in vivo* calcium

imaging of AC neruons with optogenetic of M2 axons with nVoke system (Inscopix), we

injected ~0.5 μl of retrograde AAV expressing Cre recombinase into the right PPC or STR,

and injected ~0.5 μl of AAV-hSyn-ChrimsonR-tdTomato (Addgene) into the right M2 at two

anterior-posterior (AP) coordinates (bregma +1.0mm & +1.7mm, lateral -0.75mm, depth -

0.6mm) of Ai162 mice. After the injection, the incision was sutured with a stitching fiber and

disinfected with a povidone-iodine solution.

After 10-14 days from the virus injection, gradient reflective index (GRIN) lens

implantation surgery was conducted as previously demonstrated[83]. Mice received

subcutaneously injection of carprofen (5mg/Kg) and dexamethasone (5mg/Kg) 1 hour before

the surgery. After anesthetization of a mouse, a 1mm$^2$ of square-shaped craniotomy was made

on the skull above the area of implantation. A prism-attached GRIN lens (1050-004601 or

1050-004606, Inscopix) was inserted 0.2mm away from the imaging area, where we made a

small linear incision of the cortical surface using a surgical knife (10055-12, Fine Science

Tools) coupled to the stereotaxic apparatus. The GRIN lens was gently inserted (-0.1mm/20 s,

depth -0.8mm for the PPC and the VC; -1.4mm for the AC; 1.1mm for the M2) in different

directions of the image plane based on the target areas (PPC and AC: along the medial-lateral

axis; VC and M2: along the anterior-posterior axis). The exposed brain tissue between the

GRIN lens and skull was covered with Kwik-Sil (WPI), and the inserted GRIN lens was

secured by applying Super-Bond (Super-Bond C&B Kit, Sun Medical Co.) and the

cyanoacrylate glue over the cured Kwik-Sil layer and connected skull. Mice were then

implanted with the head plates. Mice were subcutaneously injected with carprofen and

dexamethasone for three days following surgery. The field of view of each mouse was

examined on a regular basis, and if we detected obvious neuronal activity, we fastened a

baseplate (1050-004638, Inscopix) atop the GRIN lens using dental acrylic. To protect the lens

from dirt, the baseplate was covered with a baseplate cover (1050-004639, Inscopix).

*Surgical procedure for optogenetic manipulation during behavior task*

For optogenetic inhibition of projection-specific AC neurons, we bilaterally injected

retrograde AAV expressing Cre-recombinase (AAVrg-hSyn-Cre-WPRE-hGH, Addgene) into

the right PPC or bilateral STR (0.5 μl per each site) and 0.5 μl of Cre-recombinase-dependent

AAV expressing eNpHR3.0-mCherry (AAV2-EF1a-DIO-eNpHR3.0-mCherry, UNC Vector

Core) into the AC. For optogenetic inhibition of the M2-to-AC circuit, we injected the

excitatory-neuron-specific AAV expressing ChR2-eYFP (AAV2-CAMKIIa-ChR2-EYFP,

UNC Vector Core) or the AAV expressing eNpHR3.0-eYFP in neurons (AAV2-hSyn-

eNpHR3.0-eYFP, UNC Vector Core) into the right M2 at two anterior-posterior (AP)

coordinates of WT mice (0.5 μl per each coordinate). After the injection, mice were further

implanted with the head plates. Mice had 14 days of viral expression before starting water

restriction for the behavioral experiment. When mice were trained and ready for the

manipulation experiment (see Methods; **Behaviors**), we provided mice with sufficient water

for 2 days and implanted optic fiber on target sites for the laser light delivery. Mice were

anesthetized with 1.5%-2% isoflurane, and a 0.5mm-diameter craniotomy was made on the

right AC. An optic fiber with a 200μm diameter was implanted into the AC (0.6mm depth) and

glued with Kwik-Sil and cyanoacrylate glue. For the $AC_{STR}$ inhibition (Fig. 5f-j), we implanted

optic fibers into the AC bilaterally. We recovered mice for three days after the surgery and

restarted the water restriction and behavioral training.

**Behaviors**

*Hardware setups for behavior experiments*

Visual stimuli were generated with a custom MATLAB code. A gamma-corrected

LCD monitor (12cm x 9cm; minimal luminance: 0 lux; maximal luminance: 120 lux; refresh

rate: 60Hz) was placed 10 cm away from the left eye of mice at a $45°$ angle from the rostral-

caudal axis. For the binocular visual stimulation task, we put another monitor in front of the

right eye of mice as we put on the left side. Full-field drifting gratings (100% contrast, 2 Hz

temporal resolution, 0.04 cycles/° spatial resolution, 60Hz refresh rate) were used for the visual

discrimination task in two directions (rightward for Go; upward for No-go). Auditory stimuli

were generated with a custom MATLAB code (sampling rate 119,000Hz, 16bit). A speaker

(#60108, Avisoft) was placed 10cm away from the left ear of the mice. Pure tones with different

frequencies (10kHz for Go, 5kHz for No-go in the main task) were used for the auditory

discrimination task. All sound stimuli were calibrated to have a 76dB sound pressure level

(SPL) at the left ear of mice.

Licking behavior of task-performing mice was detected by a custom-made lickometer.

We delivered water reward and air punishment via solenoid valves (EV-2-24, Clippard). All

the hardware components were operated and controlled by Presentation (Neurobehavioral

Systems) with USB data acquisition devices (USB-201, Measurement Computing). An optical

mouse (G100s, Logitech) was placed beside a spinning disk of the treadmill. The rotation of

the disk during locomotion was converted to digitalized signals by the optical mouse, and real-

time locomotion speed was calculated by a customized LabView code (sampling rate: 40Hz).

If the speed exceeded the locomotion threshold (3 cm/s), a 5V-digital signal was sent to the

Presentation software in real time through a USB data acquisition device (USB-6001, National

Instruments) to present sensory stimuli during locomotion.

*Behavioral training and multisensory discrimination experiments*

Mice underwent water restriction (~1 ml of water per day) throughout the behavioral

experiment. The weight of mice under the water restriction was monitored throughout the

experiments and maintained at more than 75% of the initial body weight before the water

restriction. We calculated the hit rate, FA rate, auditory dominance rate, and correct rate as

follows:

*Correct rate (%) = (#Hit trials + #CR trials) / #Total trials X 100*

*Hit rate (%) = #Hit trials / (#Hit trials + #Miss trials) X 100*

*FA rate (%) = #FA trials / (#CR trials + #FA trials) X 100*

*Auditory dominance rate (%) =*

*(#$A_GV_{NG}$→A trials + #$A_{NG}V_G$→A trials) / #Incongruent trials X 100*

*Auditory dominance rate of $A_GV_{NG}$ trials (%) = #$A_GV_{NG}$→A trials / #$A_GV_{NG}$ trials X 100*

*Auditory dominance rate of $V_GA_{NG}$ trials (%) = #$A_{NG}V_G$→A trials / #$A_{NG}V_G$ trials X 100*

All mice were trained and tested through the following five steps:

1) reward conditioning, 2) stimulus conditioning, 3) audiovisual discrimination training, 4)

treadmill habituation; 5) audiovisual integration test

1) Reward conditioning: water-restricted mice received 0.8μl of water reward whenever they

licked the water port. If mice did not show any spontaneous licking behavior within the first 3

minutes after the beginning of a session, we manually delivered water through the water port

until mice started to lick the water port. If mice got more than 0.8ml of total water reward (>

1000 licks) within the first 20 minutes without manual water delivery, mice were moved to the

second step in the next session.

2) Stimulus conditioning: auditory go stimulus, visual go stimulus, and no stimulation trials

were pseudo-randomly presented in the proportion of 1:1:2. If mice licked the water port during

2 s of response window after the stimulus offset, 1μl of water reward and 2 s of reward

consumption time were presented (hit). If mice did not lick the water port until the offset of the

response window (miss), 1μl of water reward was delivered at the end of the response window

to encourage learning. If mice licked the water port during the no stimulation trial (false-alarm,

FA), 6 s of additional time-out period was given. We gave 4 seconds extra time for the inter-

trial interval before beginning the next trial. Task sessions automatically ended when mice

consecutively missed 30 auditory- or visual-Go stimuli. When mice showed more than 80% hit

rates for visual and auditory go stimuli, mice were moved on to the third step in the next session.

3) Audiovisual discrimination training: 4 types of stimuli (auditory Go, auditory No-go, visual

Go, visual No-go) were pseudo-randomly presented in equal proportion. There was no water

reward in the miss trials, and 400ms of air puff punishment was delivered to the right cheek of

mice with the time-out period (6 s) in the FA trials. Task sessions automatically ended when

mice consecutively missed 10 auditory- or visual-Go stimuli. If mice achieved correct rates

higher than 80% for both auditory and visual discrimination, we considered that the mice

learned the discrimination tasks.

4) Treadmill habituation: We first habituated mice to run on the treadmill during the task by

presenting the same auditory and visual stimuli when mice ran faster than a speed threshold

(3cm/s). If mice reached an expert level of discrimination (> 80% of correct rates) while

showing enough locomotion activity (> 400 trials within 1 hour), mice consecutively

performed the task in the stationary block on the immobilized treadmill and in the moving

block on the free-rotating treadmill during a single day in random order. If mice showed an

expert level of performance in both conditions, they were moved on to the final step. We

conducted optic fiber implantation surgery for the optogenetic manipulation experiments when

animals passed this step.

5) Audiovisual integration test: auditory, visual, congruent ($A_G V_G$, $A_{NG} V_{NG}$), and incongruent

($A_G V_{NG}$, $A_{NG} V_G$) stimuli were pseudo-randomly presented in the proportion of 3:3:1:1 or

3:3:2:2 for main behavioral test. 3:3:2:2 (for optogenetic experiments and *in vivo* calcium

imaging) or 3:3:1:3 (*in vivo* calcium imaging) ratio was used for the audiovisual integration

test to reserve enough trial numbers. There was no water reward and air puff punishment in the

incongruent trials. Mice conducted both stationary and moving blocks in random order on a

single day except for *in vivo* calcium imaging experiments (See the section on *in vivo* calcium

imaging experiments).

In moving sessions, we divided trials into two categories: start-moving trials and keep-

moving trials. If the average speed before the stimulus onset (-3 ~ -1.5 s) exceeded the speed

threshold (> 3 cm/s), that trial was classified as keep-moving. To evaluate the correlation

between locomotion speed and auditory dominance (Extended Data Fig. 1a), we categorized

keep-moving incongruent trials according to mean locomotion speed (3 ~ 9 cm/s, 9 ~ 15 cm/s,

and higher than 15 cm/s) before the stimulus onset (-1 ~ 0 s). We next determined the auditory

dominance rate for each category. To analyze the correlation between trial history and auditory

dominance (Extended Data Fig. 1b), we collected and calculated the auditory dominance rates

for the following trial sequences: 1) $V_G$ or $A_{NG}$ trial followed by $A_{NG}V_G$ trial, and 2) $A_G$ or $V_{NG}$

followed by $A_GV_{NG}$ trial during the moving sessions. We then estimated the auditory

dominance rate based on the modality of the previous trial. To find the relationship between

auditory dominance rate and pupil size, we averaged the maximum-normalized pupil size

before the stimulus onset (-1 ~ 0 s) during moving sessions. We then calculated the average

value of each variable according to the dominant modality.

*Optogenetic manipulation*

To inhibit $AC_{STR}$ neurons in the bilateral AC (Fig. 5), we delivered a continuous green

light from the laser (5mW, 532nm, Shanghai Laser & Optics Century) during the stimulus

period (-0.2 ~ 1s from the stimulus onsets) in 50% of trials. To suppress $M2_{\rightarrow AC}$ axon terminals

in the right AC, we delivered a continuous green light from the laser (5 ~ 8mW, 532nm,

Shanghai Laser & Optics Century) starting 200ms before the stimulus presentation (-0.2 ~ 1 s

from the stimulus onset). To control the effectiveness of laser light itself during optogenetic

inhibition of the target neurons, we conducted control experiments delivering laser lights onto

the dental cement.

**Video monitoring of orofacial and body movements of mice performing a task**

The data acquisition and analysis were performed as previously reported[84]. We used a

camera (G3-GM11-M2020-Genie, Teledyne DALSA) and lens (TEC-M55 2/3" 55mm

telecentric lens, C-Mount) to record left eye of task-performing animals at a high resolution

(1200 x 800 pixels) by illuminating the left side of the animal's face with infrared light source

(850 nm, S-IR5850, Skycares). We recorded images at 30 Hz using the Image Acquisition

Toolbox provided by MATLAB.

We calculated the pupil size of the left eye of the task-performing mice using the

following custom-written MATLAB code: 1) drawing an ROI that covers the left eye, 2) setting

the optimal brightness threshold to determine the pupil boundary, 3) converting the ROI image

to a binary image by filling in each pixel with the black and white values according to the

threshold (*imbinarize*, built-in MATLAB function), 4) filling holes within the pupil boundary

to avoid any artifacts from measuring pupil area due to light reflection on the cornea (*imfill*,

built-in MATLAB function), 5) identifying a convex hull that includes the inside part of the

pupil (*bwconvhull*, a built-in MATLAB function), and 6) counting the number of pixels within

the convex hull as the pupil size across frames. We normalized the pupil size using the

following equation, resulting in a number ranging from 0 (smallest size) to 1 (largest size):

*Normalized Pupil(t) = (Pupil(t) – min(Pupil))/max(Pupil)*

***In vivo* calcium imaging experiments**

*Data acquisition*

To avoid bleaching of calcium signals during the behavior task, we conducted an

imaging experiment in a stationary or moving block one day and tested the other block the next

day. Each imaging session was completed when mice showed three consecutive miss trials or

when the imaging time reached one hour. We analyzed the same neurons tracked over two

consecutive days in imaging data.

We performed an *in vivo* calcium imaging experiment using a miniature fluorescence

microscope (Inscopix). Images were acquired at 20 frames per second (10 frames per second

for AC$_{PPC}$ neurons) using nVista HD software (Inscopix). We cropped superfluous imaging

areas that contained no neural signal. The LED power (0.2 ~ 0.5mW) and gain value (1 ~ 4)

were adjusted based on the overall brightness of the GCaMP signals. For the simultaneous M2

axon activation and calcium imaging at the AC (Fig. 6g-k), we used the nVoke system

(Inscopix) to acquire images at 10 frames per second alternating with 0.4mW of EX-LED light

activation (red excitation light to activate ChrimsonR). The light delivery started 200ms before

the stimulus onset and ended at the stimulation offset (-0.2 ~ 1s).

*Preprocessing of imaging data*

The acquired image data were processed with Inscopix Data Processing Software

(IDPS, Inscopix). Images were downsampled by a factor of 4 and spatially filtered (highCutoff:

0.5; lowCutoff: 0.01 for PPC excitatory neurons; highCutoff: 0.5; lowCutoff: 0.005 for the

other groups). Filtered images were corrected for lateral motion (IDPS implementation).

Motion-corrected images were converted to ΔF/F images where F represents the average

intensity of each pixel during the whole imaging time. Principal component analysis -

independent component analysis (PCA-ICA) was used to identify cell boundaries on ΔF/F

images using IDPS implementation. We next extracted calcium signals by manually inspecting

cell boundaries obtained from PCA-ICA on motion-corrected images and exported average

intensities within cell boundaries. The same neurons were identified throughout several

imaging sessions using longitudinal registration (IDPS implementation), and only neurons that

appeared across imaging sessions were selected for further analysis.

*Calculation of response amplitudes and sorting neuronal type*

Exported calcium data were analyzed using custom MATLAB codes. The extracted

calcium response data was converted to ΔF/F for each neuron. ΔF/F was calculated as $\Delta F/F =$

$(F - F0)/F0$, where F0 represents the average fluorescence of a neuron across the imaging

time. All the analyses were conducted based on the ΔF/F value.

To determine if a neuron responded to distinct stimuli (Fig. 2c), we compared average

calcium activity during baseline (-0.2 ~ 0 s) and stimulation (0 ~ 1 s) periods using Wilcoxon

signed-rank test (*P<0.01*, MATLAB implementation). The response amplitude for each trial

was calculated by subtracting the average of baseline activity from the average of calcium

activity during the first 1 s after stimulus onset. To dissect neuronal type (Fig. 2c), we compared

Go and No-go responses if neurons showed significant responses to Go and No-go stimuli in

one modality and classified as Go or No-go neurons if they had a significantly stronger

response in the Go or No-go trials. Neurons that showed significant responses in one auditory

and one visual stimulus were classified as "Dual," while neurons with significant responses in

three unisensory trial types were classified as "Triple." "All" type neurons responded

significantly in all unisensory trials. To examine how locomotion affects unisensory responses

(Extended Data Fig. 2a), we compared the response amplitudes of each neuron from stationary

and moving sessions using the Mann-Whitney U-test (*P < 0.01*).

744             To calculate the change in spontaneous activity of imaged neurons by locomotion

onset and offset, we collected calcium activity and locomotion speed data at all locomotion

onsets or offsets (-2 ~ 2s) and concatenated them. We then calculated the Pearson correlation

coefficient between the two variables. We used the following conditions to define locomotion

onsets and offsets during an imaging session.

Locomotion onset at 0 s must satisfy all of the following conditions:

*1) Average locomotion speed (-1 ~ -0.5 s) < 0.5cm/s;*

*2) Average locomotion speed (-0.5 ~ 1 s) < 0.5cm/s;*

*3) Average locomotion speed (0 ~ 250 ms) > 1cm/s;*

*4) Average locomotion speed (250 ~ 1000 ms) > 2cm/s;*

*5) No stimulus during -2 ~ 2 s.*

Locomotion offset at 0 s must satisfy all of the following conditions:

*1) Average locomotion speed (-1 ~ -0.5 s) > 2 cm/s;*

*2) Average locomotion speed (-0.5 ~ 1 s) > 1cm/s;*

*3) Average locomotion speed (0 ~ 250 ms) < 0.5cm/s;*

*4) Average locomotion speed (250 ~ 1000 ms) < 0.5cm/s;*

*5) No stimulus during -2 ~ 2 s.*

761             To calculate the change in M2$_{AC}$ activity at locomotion onset (Fig. 6f), we subtracted

762 mean baseline activity (-1 ~ -0.5s from the locomotion onset) from mean calcium signal at

763 early (0 ~ 1.5s from the locomotion onset) or late (3 ~ -1 from the locomotion offset) periods.

765 *Demixed principal component analysis of imaging data*

766             To visualize populational activity during the behavioral task over time, we used

demixed principal component analysis (dPCA)[68]. The dPCA was performed using trial-

averaged calcium activity aligned at stimulus onset (-1 ~ 3 s) of all neurons across the eight

unisensory trial types (visual hit, visual miss, visual CR, visual FA, auditory hit, auditory miss,

auditory CR, and auditory FA). To denoise the calcium data, we temporally downsampled it

(factor 2) and applied Gaussian smoothing with a window size of three bins. We excluded

sessions that had fewer than three trials of lick or no-lick in the unisensory and incongruent

trials. The encoding PCs from the dPCA were classified into stimulus-related, decision-related,

stimulus-decision interaction-related, and independent axes. To show trajectories for each trial

type in Fig. 2g, 3a, 3c, and Extended Data Fig. 5d,6d, we used dPCA results derived from the

all imaged neurons. For all quantifications, we used the dPCA results from each imaging

session. We projected the activities of the multisensory trials (congruent and incongruent trials)

by using transformation vectors obtained from the dPCA with the unisensory trials.

To quantify the similarities between trial types, we calculated Euclidean distances

between coordinates after the stimulus onset (0 ~ 1 s) in a subspace containing the top 3 sorted

PCs (top 3 stimulus PCs explained the 62.8% variance of the data on average). To compare

neural trajectories based on locomotion state, trial-averaged activities of unisensory trials from

stationary, start-moving, and keep-moving trials were analyzed by dPCA. To distinguish

between start-moving and keep-moving data in the moving sessions, we projected trial data

into the encoding PC axes from the dPCA of each moving session. The Euclidean distance

between two trajectories was calculated 1 s after the stimulus onset. Projection distance was

measured as the Euclidean distance between 0 and 1 s following the stimulus onset.

*dPCA based classifiers*

For classifiers (Fig. 2 and Extended Data Fig. 3,5,6), we projected single-trial activity

onto the top encoding PC axes and computed Euclidean distances between single trial data

(only correct trials for unisensory classification) and trial-averaged trajectories of the

unisensory trials (for unisensory classification) or incongruent trials (for incongruent

classification) at each time points. Each test trial was classified into a trial type of the trial-

averaged trajectory that showed the smallest Euclidean distance from the test trial trajectory. If

the test trajectory showed a very short projection in the PC axes, its distance to the center of

the trajectories at the pre-stimulus time points was smaller than the distance to the trial-

averaged trajectories at each time point. We counted it as incorrect classification.

To determine the significance of similarity with visual trajectories in determining

dominant modality (Fig. 3e,f), we first projected the activities of individual unisensory trials

onto the top three stimulus encoding PC axes and calculated the Euclidean distance between

each trajectory and the trial-averaged target visual trajectory (target visual trajectory: $V_G$ for

$A_{NG}V_G$ classification; $V_{NG}$ for $A_GV_{NG}$ classification). To calculate the distribution of individual

visual trials, we used linear discriminant analysis (LDA) to determine an Euclidean distance

threshold that distinguishes visual trials from other visual and auditory trials. Finally, we

classified individual incongruent trials according to their Euclidean distance to the trial-

averaged target visual trajectory, with each trial classified as a visual-dominant trial if its

Euclidean distance was less than the threshold. To statistically compare classifier accuracies,

we calculated mean classifier accuracy between 0.5 and 1s following stimulus onset, when the

classifiers attained a peak of classification accuracy, and compared between groups.

**Histology and Immunohistochemistry**

We performed histology experiments to confirm imaging sites after *in vivo* calcium imaging,

injection and expression sites of viral vectors, and anatomical circuit tracing data. Mice were

anesthetized with avertin (2,2,2-Tribromoethanol (Sigma-Aldrich), 125-250mg/kg,

intraperitoneally) and transcardially perfused with 15ml of PBS, followed by 4%

paraformaldehyde (PFA, w/v in PBS). Brain samples were post-fixed for 4 hours in the PFA

at 4°C, then washed 3 times with PBS for 10 minutes each. After washing, the brain samples

were immersed in a 30% sucrose solution (w/v in PBS) for 2-5 days. When the brain samples

sank, they were embedded in optimal cutting temperature medium (Tissue-Tek O.C.T.

Compound, Sakura Finetek) and rapidly froze at -80°C. Frozen brain samples were sectioned

into 40μm or 20μm (for immunostaining) in a coronal direction using a cryocut (Leica). We

washed brain slices 3 times in PBS for 10 minutes before mounting them with an anti-bleaching

mounting medium containing 4′,6-diamidino-2-phenylindole (DAPI) (Vector Labs). To

immunostain the CaMKIIα, we permeabilized the brain sections for 30 min with 0.5% w/v

Triton X-100 in PBS, then blocked with 2% w/v normal goat serum (NGS) in PBS for 2 hours

at room temperature. The samples were incubated at 4°C for 48 hours with mouse anti-

CaMKIIα antibodies (ab22609, abcam, 1:250 dilution in 2% NGS). After treating the primary

antibodies, we washed the samples 3 times in PBS for 10 minutes. We used Alexa Fluor 594

goat anti-mouse IgG (A11005, Invitrogen, 1:250 dilution in PBS) to stain CaMKIIα for 2 hours

at room temperature. Following that, we washed the samples in PBS 3 times for 10 minutes

before mounting them with anti-bleaching medium with DAPI. Fluorescent images of brain

slices were taken using a slide scanner (Zeiss Axio Scan. Z1) or a confocal microscope (Nikon

A1 HD25).

**Statistical analysis**

We analyzed the data using custom-written codes in MATLAB (Mathworks). All data were

presented as mean ± SEM unless otherwise noted. 'N' denotes the number of mice or sessions,

whereas 'n' represents the number of neurons. All statistical analyses were described in the

figure legends. All the statistical significances of the datasets were indicated as NS (not

significant; $P \geq 0.05$), * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), and **** ($P < 0.0001$).

**References**

1     Stein, B. E. *The new handbook of multisensory processing*. (MIT Press, 2012).

2     Gielen, S. C., Schmidt, R. A. & Van den Heuvel, P. J. On the nature of intersensory
facilitation of reaction time. *Percept Psychophys* **34**, 161-168 (1983).
https://doi.org/10.3758/bf03211343

3     Diederich, A. & Colonius, H. Bimodal and trimodal multisensory enhancement: effects
of stimulus onset and intensity on reaction time. *Percept Psychophys* **66**, 1388-1404
(2004). https://doi.org/10.3758/bf03195006

4     Raposo, D., Sheppard, J. P., Schrater, P. R. & Churchland, A. K. Multisensory decision-
making in rats and humans. *J Neurosci* **32**, 3726-3735 (2012).
https://doi.org/10.1523/JNEUROSCI.4998-11.2012

5     Meijer, G. T., Pie, J. L., Dolman, T. L., Pennartz, C. M. A. & Lansink, C. S. Audiovisual
Integration Enhances Stimulus Detection Performance in Mice. *Front Behav Neurosci*
**12**, 231 (2018). https://doi.org/10.3389/fnbeh.2018.00231

6     Colavita, F. B. Human Sensory Dominance. *Perception & Psychophysics* **16**, 409-412
(1974). https://doi.org/Doi 10.3758/Bf03203962

7     McGurk, H. & MacDonald, J. Hearing lips and seeing voices. *Nature* **264**, 746-748
(1976). https://doi.org/10.1038/264746a0

8     Battaglia, P. W., Jacobs, R. A. & Aslin, R. N. Bayesian integration of visual and
auditory signals for spatial localization. *J Opt Soc Am A Opt Image Sci Vis* **20**, 1391-
1397 (2003). https://doi.org/10.1364/josaa.20.001391

9     Alais, D. & Burr, D. The ventriloquist effect results from near-optimal bimodal
integration. *Curr Biol* **14**, 257-262 (2004). https://doi.org/10.1016/j.cub.2004.01.029

10    Fetsch, C. R., Turner, A. H., DeAngelis, G. C. & Angelaki, D. E. Dynamic reweighting
of visual and vestibular cues during self-motion perception. *J Neurosci* **29**, 15601-
15612 (2009). https://doi.org/10.1523/JNEUROSCI.2574-09.2009

11    Fetsch, C. R., Pouget, A., DeAngelis, G. C. & Angelaki, D. E. Neural correlates of
reliability-based cue weighting during multisensory integration. *Nat Neurosci* **15**, 146-
154 (2011). https://doi.org/10.1038/nn.2983

12    Song, Y. H. *et al.* A Neural Circuit for Auditory Dominance over Visual Perception.
*Neuron* **93**, 940-+ (2017). https://doi.org/10.1016/j.neuron.2017.01.006

13    Choi, I. & Lee, S. H. Reduced activity of parvalbumin-positive interneurons in the
posterior parietal cortex causes visually dominant multisensory decisions in freely
navigating mice. *Mol Brain* **15**, 82 (2022). https://doi.org/10.1186/s13041-022-00968-
x

14    Ernst, M. O. & Banks, M. S. Humans integrate visual and haptic information in a
statistically optimal fashion. *Nature* **415**, 429-433 (2002).
https://doi.org/10.1038/415429a

15    Kording, K. P. & Wolpert, D. M. Bayesian integration in sensorimotor learning. *Nature*
**427**, 244-247 (2004). https://doi.org/10.1038/nature02169

16    Coen, P., Sit, T. P. H., Wells, M. J., Carandini, M. & Harris, K. D. Mouse frontal cortex
mediates additive multisensory decisions. *Neuron* **111**, 2432-2447 e2413 (2023).
https://doi.org/10.1016/j.neuron.2023.05.008

17    Keil, J., Muller, N., Ihssen, N. & Weisz, N. On the variability of the McGurk effect:
audiovisual integration depends on prestimulus brain states. *Cereb Cortex* **22**, 221-231
(2012). https://doi.org/10.1093/cercor/bhr125

18    Jiang, W., Jiang, H. & Stein, B. E. Two corticotectal areas facilitate multisensory
orientation behavior. *J Cogn Neurosci* **14**, 1240-1255 (2002).
https://doi.org/10.1162/089892902760807230

19    Avillac, M., Ben Hamed, S. & Duhamel, J. R. Multisensory integration in the ventral
intraparietal area of the macaque monkey. *J Neurosci* **27**, 1922-1932 (2007).
https://doi.org/10.1523/JNEUROSCI.2646-06.2007

20    Mallick, D. B., Magnotti, J. F. & Beauchamp, M. S. Variability and stability in the
McGurk effect: contributions of participants, stimuli, time, and response type. *Psychon*
*Bull Rev* **22**, 1299-1307 (2015). https://doi.org/10.3758/s13423-015-0817-4

21    Sinnett, S., Spence, C. & Soto-Faraco, S. Visual dominance and attention: the Colavita
effect revisited. *Percept Psychophys* **69**, 673-686 (2007).
https://doi.org/10.3758/bf03193770

22    Crochet, S., Lee, S. H. & Petersen, C. C. H. Neural Circuits for Goal-Directed
Sensorimotor Transformations. *Trends Neurosci* **42**, 66-77 (2019).
https://doi.org/10.1016/j.tins.2018.08.011

23    Esmaeili, V. *et al.* Rapid suppression and sustained activation of distinct cortical
regions for a delayed sensory-triggered motor response. *Neuron* **109**, 2183-2201 e2189
(2021). https://doi.org/10.1016/j.neuron.2021.05.005

24    Choi, I., Lee, J. Y. & Lee, S. H. Bottom-up and top-down modulation of multisensory
integration. *Curr Opin Neurobiol* **52**, 115-122 (2018).
https://doi.org/10.1016/j.conb.2018.05.002

25    McGinley, M. J. *et al.* Waking State: Rapid Variations Modulate Neural and Behavioral
Responses. *Neuron* **87**, 1143-1161 (2015).
https://doi.org/10.1016/j.neuron.2015.09.012

26    Ayaz, A., Saleem, A. B., Scholvinck, M. L. & Carandini, M. Locomotion controls
spatial integration in mouse visual cortex. *Curr Biol* **23**, 890-894 (2013).
https://doi.org/10.1016/j.cub.2013.04.012

27    Bennett, C., Arroyo, S. & Hestrin, S. Subthreshold Mechanisms Underlying State-
Dependent Modulation of Visual Responses. *Neuron* **80**, 350-357 (2013).
https://doi.org/10.1016/j.neuron.2013.08.007

28    Dipoppa, M. *et al.* Vision and Locomotion Shape the Interactions between Neuron
Types in Mouse Visual Cortex. *Neuron* **98**, 602-+ (2018).
https://doi.org/10.1016/j.neuron.2018.03.037

29    Fu, Y. *et al.* A Cortical Circuit for Gain Control by Behavioral State. *Cell* **156**, 1139-
1152 (2014). https://doi.org/10.1016/j.cell.2014.01.050

30    Lee, A. M. *et al.* Identification of a Brainstem Circuit Regulating Visual Cortical State in Parallel with Locomotion. *Neuron* **83**, 455-466 (2014). https://doi.org/10.1016/j.neuron.2014.06.031

31    Niell, C. M. & Stryker, M. P. Modulation of Visual Responses by Behavioral State in Mouse Visual Cortex. *Neuron* **65**, 472-479 (2010). https://doi.org/10.1016/j.neuron.2010.01.033

32    Paken, J. M. P. *et al.* Behavioral-state modulation of inhibition is context-dependent and cell type specific in mouse visual cortex. *Elife* **5** (2016). https://doi.org/ARTN e14985

10.7554/eLife.14985

Polack, P. O., Friedman, J. & Golshani, P. Cellular mechanisms of brain state-dependent gain modulation in visual cortex. *Nature Neuroscience* **16**, 1331-U1227 (2013). https://doi.org/10.1038/nn.3464

Vinck, M., Batista-Brito, R., Knoblich, U. & Cardin, J. A. Arousal and Locomotion Make Distinct Contributions to Cortical Activity Patterns and Visual Encoding. *Neuron* **86**, 740-754 (2015). https://doi.org/10.1016/j.neuron.2015.03.028

McGinley, M. J., David, S. V. & McCormick, D. A. Cortical Membrane Potential Signature of Optimal States for Sensory Signal Detection. *Neuron* **87**, 179-192 (2015). https://doi.org/10.1016/j.neuron.2015.05.038

Zhou, M. *et al.* Scaling down of balanced excitation and inhibition by active behavioral states in auditory cortex. *Nature Neuroscience* **17**, 841-850 (2014). https://doi.org/10.1038/nn.3701

Nelson, A. *et al.* A Circuit for Motor Cortical Modulation of Auditory Cortical Activity. *Journal of Neuroscience* **33**, 14342-+ (2013). https://doi.org/10.1523/Jneurosci.2275-13.2013

Schneider, D. M., Nelson, A. & Mooney, R. A synaptic and circuit basis for corollary discharge in the auditory cortex. *Nature* **513**, 189-+ (2014). https://doi.org/10.1038/nature13724

Schneider, D. M., Sundararajan, J. & Mooney, R. A cortical filter that learns to suppress the acoustic consequences of movement. *Nature* **561**, 391-+ (2018). https://doi.org/10.1038/s41586-018-0520-5

Runyan, C. A., Piasini, E., Panzeri, S. & Harvey, C. D. Distinct timescales of population coding across cortex. *Nature* **548**, 92-96 (2017). https://doi.org/10.1038/nature23020

Williamson, R. S., Hancock, K. E., Shinn-Cunningham, B. G. & Polley, D. B. Locomotion and Task Demands Differentially Modulate Thalamic Audiovisual Processing during Active Search. *Curr Biol* **25**, 1885-1891 (2015). https://doi.org/10.1016/j.cub.2015.05.045

Zhou, Y. & Freedman, D. J. Posterior parietal cortex plays a causal role in perceptual and categorical decisions. *Science* **365**, 180-185 (2019). https://doi.org/10.1126/science.aaw8347

Lyamzin, D. & Benucci, A. The mouse posterior parietal cortex: Anatomy and functions. *Neurosci Res* **140**, 14-22 (2019). https://doi.org/10.1016/j.neures.2018.10.008

Freedman, D. J. & Ibos, G. An Integrative Framework for Sensory, Motor, and Cognitive Functions of the Posterior Parietal Cortex. *Neuron* **97**, 1219-1234 (2018). https://doi.org/10.1016/j.neuron.2018.01.044

Goard, M. J., Pho, G. N., Woodson, J. & Sur, M. Distinct roles of visual, parietal, and frontal motor cortices in memory-guided sensorimotor decisions. *Elife* **5** (2016). https://doi.org/10.7554/eLife.13764

Pho, G. N., Goard, M. J., Woodson, J., Crawford, B. & Sur, M. Task-dependent representations of stimulus and choice in mouse parietal cortex. *Nat Commun* **9**, 2596 (2018). https://doi.org/10.1038/s41467-018-05012-y

Nikbakht, N., Tafreshiha, A., Zoccolan, D. & Diamond, M. E. Supralinear and Supramodal Integration of Visual and Tactile Signals in Rats: Psychophysics and Neuronal Mechanisms. *Neuron* **97**, 626-639 e628 (2018). https://doi.org/10.1016/j.neuron.2018.01.003

Oude Lohuis, M. N., Marchesi, P., Pennartz, C. M. A. & Olcese, U. Functional (ir)relevance of posterior parietal cortex during audiovisual change detection. *J Neurosci* **42**, 5229-5245 (2022). https://doi.org/10.1523/JNEUROSCI.2150-21.2022

Harvey, C. D., Coen, P. & Tank, D. W. Choice-specific sequences in parietal cortex during a virtual-navigation decision task. *Nature* **484**, 62-68 (2012). https://doi.org/10.1038/nature10918

Raposo, D., Kaufman, M. T. & Churchland, A. K. A category-free neural population supports evolving demands during decision-making. *Nature Neuroscience* **17**, 1784-1792 (2014). https://doi.org/10.1038/nn.3865

Licata, A. M. *et al.* Posterior Parietal Cortex Guides Visual Decisions in Rats. *J Neurosci* **37**, 4954-4966 (2017). https://doi.org/10.1523/JNEUROSCI.0105-17.2017

Hwang, E. J., Dahlen, J. E., Mukundan, M. & Komiyama, T. History-based action selection bias in posterior parietal cortex. *Nat Commun* **8**, 1242 (2017). https://doi.org/10.1038/s41467-017-01356-z

Driscoll, L. N., Pettit, N. L., Minderer, M., Chettih, S. N. & Harvey, C. D. Dynamic Reorganization of Neuronal Activity Patterns in Parietal Cortex. *Cell* **170**, 986-999 e916 (2017). https://doi.org/10.1016/j.cell.2017.07.021

Tseng, S. Y., Chettih, S. N., Arlt, C., Barroso-Luque, R. & Harvey, C. D. Shared and specialized coding across posterior cortical areas for dynamic navigation decisions. *Neuron* **110**, 2484-2502 e2416 (2022). https://doi.org/10.1016/j.neuron.2022.05.012

Akrami, A., Kopec, C. D., Diamond, M. E. & Brody, C. D. Posterior parietal cortex represents sensory history and mediates its effects on behaviour. *Nature* **554**, 368-372 (2018). https://doi.org/10.1038/nature25510

Zhong, L. *et al.* Causal contributions of parietal cortex to perceptual decision-making during stimulus categorization. *Nat Neurosci* **22**, 963-973 (2019). https://doi.org/10.1038/s41593-019-0383-6

Gallero-Salas, Y. *et al.* Sensory and Behavioral Components of Neocortical Signal Flow in Discrimination Tasks with Short-Term Memory. *Neuron* **109**, 135-148 e136 (2021). https://doi.org/10.1016/j.neuron.2020.10.017

Yao, J. D., Gimoto, J., Constantinople, C. M. & Sanes, D. H. Parietal Cortex Is Required for the Integration of Acoustic Evidence. *Curr Biol* **30**, 3293-3303 e3294 (2020). https://doi.org/10.1016/j.cub.2020.06.017

Zingg, B. *et al.* Neural networks of the mouse neocortex. *Cell* **156**, 1096-1111 (2014). https://doi.org/10.1016/j.cell.2014.02.023

Zhang, S. *et al.* Organization of long-range inputs and outputs of frontal cortex for top-down control. *Nat Neurosci* **19**, 1733-1742 (2016). https://doi.org/10.1038/nn.4417

Lewis, J. W. & Van Essen, D. C. Corticocortical connections of visual, sensorimotor, and multimodal processing areas in the parietal lobe of the macaque monkey. *J Comp Neurol* **428**, 112-137 (2000). https://doi.org/10.1002/1096-9861(20001204)428:1<112::aid-cne8>3.0.co;2-9

Driver, J. & Noesselt, T. Multisensory interplay reveals crossmodal influences on 'sensory-specific' brain regions, neural responses, and judgments. *Neuron* **57**, 11-23 (2008). https://doi.org/10.1016/j.neuron.2007.12.013

Schlack, A., Sterbing-D'Angelo, S. J., Hartung, K., Hoffmann, K. P. & Bremmer, F. Multisensory space representations in the macaque ventral intraparietal area. *J Neurosci* **25**, 4616-4625 (2005). https://doi.org/10.1523/JNEUROSCI.0455-05.2005

Olcese, U., Iurilli, G. & Medini, P. Cellular and synaptic architecture of multisensory integration in the mouse neocortex. *Neuron* **79**, 579-593 (2013). https://doi.org/10.1016/j.neuron.2013.06.010

Wallace, M. T., Ramachandran, R. & Stein, B. E. A revised view of sensory cortical parcellation. *Proc Natl Acad Sci U S A* **101**, 2167-2172 (2004). https://doi.org/10.1073/pnas.0305697101

Shimaoka, D., Harris, K. D. & Carandini, M. Effects of Arousal on Mouse Sensory Cortex Depend on Modality (vol 22, pg 3160, 2018). *Cell Reports* **25**, 3230-3230 (2018). https://doi.org/10.1016/j.celrep.2018.11.105

Schneider, D. M. & Mooney, R. Motor-related signals in the auditory system for listening and learning. *Curr Opin Neurobiol* **33**, 78-84 (2015). https://doi.org/10.1016/j.conb.2015.03.004

Kobak, D. *et al.* Demixed principal component analysis of neural population data. *Elife* **5** (2016). https://doi.org/10.7554/eLife.10989

Chen, L., Wang, X. X., Ge, S. Y. & Xiong, Q. J. Medial geniculate body and primary auditory cortex differentially contribute to striatal sound representations. *Nature Communications* **10** (2019). https://doi.org/ARTN 418
10.1038/s41467-019-08350-7

70    Guo, L., Walker, W. I., Ponvert, N. D., Penix, P. L. & Jaramillo, S. Stable representation of sounds in the posterior striatum during flexible auditory decisions. *Nature Communications* **9** (2018). https://doi.org/ARTN 1534
10.1038/s41467-018-03994-3

71    Xiong, Q. J., Znamenskiy, P. & Zador, A. M. Selective corticostriatal plasticity during acquisition of an auditory discrimination task. *Nature* **521**, 348-+ (2015). https://doi.org/10.1038/nature14225

72   Ponvert, N. D. & Jaramillo, S. Auditory Thalamostriatal and Corticostriatal Pathways
Convey Complementary Information about Sound Features. *Journal of Neuroscience*
**39**, 271-280 (2019). https://doi.org/10.1523/Jneurosci.1188-18.2018
73   Lin, P. A., Asinof, S. K., Edwards, N. J. & Isaacson, J. S. Arousal regulates frequency
tuning in primary auditory cortex. *Proc Natl Acad Sci U S A* **116**, 25304-25310 (2019).
https://doi.org/10.1073/pnas.1911383116
74   Welch, R. B. & Warren, D. H. Immediate perceptual response to intersensory
discrepancy. *Psychol Bull* **88**, 638-667 (1980).
75   Repp, B. H. & Penel, A. Auditory dominance in temporal processing: New evidence
from synchronization with simultaneous visual and auditory sequences. *J Exp Psychol*
*Human* **28**, 1085-1099 (2002). https://doi.org/10.1037//0096-1523.28.5.1085
76   Crapse, T. B. & Sommer, M. A. Corollary discharge across the animal kingdom. *Nat*
*Rev Neurosci* **9**, 587-600 (2008). https://doi.org/10.1038/nrn2457
77   Matin, E. Saccadic suppression: a review and an analysis. *Psychol Bull* **81**, 899-917
(1974). https://doi.org/10.1037/h0037368
78   Bremmer, F., Kubischik, M., Hoffmann, K. P. & Krekelberg, B. Neural dynamics of
saccadic     suppression.     *J     Neurosci*     **29**,     12374-12383     (2009).
https://doi.org/10.1523/JNEUROSCI.2908-09.2009
79   Kuan, A. T. *et al.* Synaptic wiring motifs in posterior parietal cortex support decision-
making.        *bioRxiv*,        2022.2004.2013.488176        (2022).
https://doi.org/10.1101/2022.04.13.488176
80   Romanski, L. M. Representation and integration of auditory and visual stimuli in the
primate ventral lateral prefrontal cortex. *Cereb Cortex* **17 Suppl 1**, i61-69 (2007).
https://doi.org/10.1093/cercor/bhm099
81   Calvert, G. A., Campbell, R. & Brammer, M. J. Evidence from functional magnetic
resonance imaging of crossmodal binding in the human heteromodal cortex. *Curr Biol*
**10**, 649-657 (2000). https://doi.org/10.1016/s0960-9822(00)00513-3
82   Siniscalchi, M. J., Phoumthipphavong, V., Ali, F., Lozano, M. & Kwan, A. C. Fast and
slow transitions in frontal ensemble activity during flexible sensorimotor behavior. *Nat*
*Neurosci* **19**, 1234-1242 (2016). https://doi.org/10.1038/nn.4342
83   Gulati, S., Cao, V. Y. & Otte, S. Multi-layer Cortical Ca2+ Imaging in Freely Moving
Mice with Prism Probes and Miniaturized Fluorescence Microscopy. *Jove-J Vis Exp*
(2017). https://doi.org/ARTN e55579
10.3791/55579
84   Kim, J. H., Ma, D. H., Jung, E., Choi, I. & Lee, S. H. Gated feedforward inhibition in
the frontal cortex releases goal-directed action. *Nat Neurosci* **24**, 1452-1464 (2021).
https://doi.org/10.1038/s41593-021-00910-9

1089

**Fig. 1 | Locomotion switches auditory-dominant decisions to visually dominant ones in mice under audiovisual conflicts**

**a**, Schematic illustration of the audiovisual discrimination task in a head-fixed mouse on the treadmill. G, go; NG, no-go; L, lick; NL, no lick; FA, false alarm; CR, correct rejection.

**b**, Stimuli used for behavioral training. Aud, auditory stimuli; Vis, visual stimuli.

**c**, Learning curves across mice trained with the task (N = 22 mice). Thin lines represent individual mice. The black dashed line indicates a criterion for expert performance (80% correct rate). Red, visual discrimination performance; Blue, auditory discrimination performance.

**d**, Schematics of stationary (top, S) and moving (bottom, M) conditions.

**e**, Audiovisual stimuli used for the multisensory test. Note that letter 'A' and 'V' after the arrow

marks indicates auditory dominance and visual dominance respectively.

**f**, Correct rates of mice in the auditory (Aud), visual (Vis), and congruent (Con) trials (N = 22

mice). Each circle indicates one mouse. S, stationary; M, moving. Two-way ANOVA with

Tukey's test. Locomotion, $P < 0.0001$; Stimulus, $P < 0.0001$; Interaction: $P < 0.0001$.

**g**, Auditory dominance rates in the incongruent trials (Incon) (N = 22 mice). Note that auditory

dominance rates were calculated as the fraction of incongruent trials when mice showed

decisions following auditory Go/No-go rules. Wilcoxon signed-rank test.

**h**, Auditory dominance rates in the incongruent trials at different locomotion states (N = 22

mice). S, stationary; SM, start-moving; KM, keep-moving. Gray lines represent individual

mouse data. Wilcoxon signed-rank test.

Data are presented as means ± SEM. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$;

1112

1113

Fig. 2 | Locomotion suppresses auditory representation while preserving visual representation in the PPC.

a, Top, schematics of *in vivo* calcium imaging of the PPC excitatory neurons in a task-performing mouse. Bottom left, lens-inserted sites across the imaged mice (scale bar: 1mm); bottom right, immunohistochemical confirmation of CaMKIIα expression in GCaMP6s-expressing neurons (scale bars: 20 μm). 10 imaging experiments (10 stationary + 10 moving sessions) from 7 mice.

b, Measured calcium activities of 4 example neurons (black lines) and the locomotion speed (gray line) of an example moving mouse.

**c**, Response types of the PPC neurons (n = 961 neurons) in the stationary (left) and moving (right) conditions. Top, percentages of neurons classified according to the response types. Bottom, percentages of unisensory and multisensory neurons.

**d**, Stimulus-evoked response amplitudes of the PPC excitatory neurons (n = 961) in the correct unisensory trials at different locomotion states. Dots indicate individual neurons. Red cross lines indicate means ± standard deviations. Red, paired-sample t-test between response amplitudes at stationary and moving conditions.

**e**, Schematic illustration of dPCA to decompose the PPC populational activity into demixed axes representing specific task-relevant variables. Stim, stimulus; Deci, decision.

**f**, Percentages of explained variances of top 20 demixed PCs calculated in 10 stationary and 10 moving sessions (N = 20). Lines with shades are means ± SEM. Asterisks indicate statistical comparisons between the stimulus PCs and other PCs. One-way ANOVA with Fisher's LSD test.

**g**, Stimulus PC trajectories averaged across unisensory correct trials from all stationary or moving sessions. Circles indicate 1 s after the stimulus onset. Projection distance indicates the Euclidean distance between the points at 0 s and 1 s after the stimulus onset. Go/No-go distance indicates the Euclidean distance between Go and No-go trajectories at 1 s after the stimulus onset.

**h**, Projection distances of the unisensory trajectories at different states (N = 10). Two-way ANOVA with Fisher's LSD test. Stimulus, $P < 0.0001$; Locomotion, $P = 0.00085$; Interaction: $P = 0.0029$.

**i**, Go/No-go distances in the auditory (Aud) or visual (Vis) trials at different states (N = 10). Two-way ANOVA with Fisher's LSD test. Stimulus, $P < 0.0001$; Locomotion, $P = 0.039$; Interaction, $P = 0.034$.

**j**, Schematic illustration of the dPCA classifier that predicts trial types based on the Euclidean

distance between a test trial trajectory and trial-averaged trajectories in the same subspace.

**k**, Accuracies of the dPCA classifier in sorting unisensory trial types. Solid lines, mean

accuracies in stationary sessions; Dashed lines, mean accuracies in moving sessions; Thin lines,

upper boundaries of 95% confidence intervals with shuffled trial labels. Colors, unisensory

trial types. Vertical gray lines, stimulus onset (solid) and offset (dashed).

**l**, Mean classifier accuracies during the $0.5 - 1s$ period after the stimulus onset (N = 10). Two-

way ANOVA with Fisher's LSD test. Stimulus, $P < 0.0001$; Locomotion, $P < 0.0001$;

Interaction, $P < 0.0001$.

Data are presented as means ± SEM except for **d**. NS (not significant); *$P < 0.05$, **$P < 0.01$,

***$P < 0.001$, ****$P < 0.0001$.

1158

1159

**Fig. 3 | The similarity between visual and multisensory representations in the PPC**

**predicts dominant modality in multisensory decisions.**

**a**, Trajectories of trial-averaged calcium traces across all sessions for congruent and unisensory

trials in the stimulus subspace from stationary (left) and moving (right) trials.

**b**, Mean Euclidean distances between congruent and unisensory trajectories for 1s during

stimulus presentation at different locomotion states (N = 10). Two-way ANOVA with Fisher's

LSD test. Pair type, $P < 0.0001$; Locomotion, $P = 0.022$; Interaction, $P = 0.0068$.

**c**, Same as **a**, but for incongruent and unisensory trials. Note that the incongruent trajectories

were similar to the visual trajectories when mice made visual-dominant decisions. Black lines,

trajectories for visual-dominant trials; Gray lines, trajectories for auditory-dominant trials.

Solid lines, for $A_{NG}V_G$ trials; Dashed lines, for $A_GV_{NG}$ trials.

**d**, Same as **b**, but between incongruent and unisensory trajectories (N = 10). Two-way ANOVA

with Fisher's LSD test. Pair type, $P < 0.0001$; Locomotion, $P = 0.014$; Interaction, $P = 0.64$.

**e**, Left, an example session data showing Euclidean distances between individual unisensory

trajectories and the trial-averaged $V_G$ trajectory at 1s after the stimulus onset. Note that the

LDA boundaries were set at the distance where $V_G$ trials were separated from other trials the

most. Right, Euclidean distances between individual $A_{NG}V_G$ trajectories and the trial-averaged

$V_G$ trajectory. LDA boundary was used to classify the dominant modality in $A_{NG}V_G$ trials. Dark

gray, visual-dominant trials; light gray, auditory-dominant trials.

**f**, Left, accuracies of the LDA classifier in predicting dominance of multisensory trials (N =

10). Solid lines for $A_{NG}V_G$ trials; Dashed lines for $A_GV_{NG}$ trials; Thin gray lines, upper

boundaries of 95% confidence intervals with shuffled trial labels. Vertical gray lines, stimulus

onset (solid) and offset (dashed). Right, mean classifier accuracy at 1s after the stimulus onset

(N = 10). S, stationary state; M, moving state. One-way ANOVA, $P = 0.93$.

**g**, Euclidean distances from the incongruent trials to the visual trajectory and the auditory

dominance rates of mice at different locomotion states (N = 10). Left, $A_{NG}V_G$ trials; right,

$A_GV_{NG}$ trials. Dots connected by gray lines, data from individual imaging. Red crosses, means

± standard deviations. Wilcoxon signed-rank test.

**h**, Response amplitudes of unisensory-responsive neurons to congruent and incongruent

stimuli at different locomotion states. Circles with error bars indicate response amplitudes of

the same neurons in unisensory visual (red) and auditory (blue) trials. Mann-Whitney U test

(between S and SM or KM) and Wilcoxon signed-rank test (between SM and KM) with

Bonferonni correction (multiplied by 3) on the multisensory responses.

Data are presented as means ± SEM except for **g**. NS (not significant); $\#P < 0.1$, $*P < 0.05$,

$**P < 0.01$, $***P < 0.001$, $****P < 0.0001$.

1195

**Fig. 4 | Locomotion preferentially suppresses auditory outputs to the parietal cortex.**

**a**, Schematic illustration (left) and injection sites (right) of retrograde AAVs for dual retrograde

tracing of the AC neurons that project to the PPC (red, $AC_{PPC}$) and to the posterior STR (green,

$AC_{STR}$). Scale bars, 1 mm.

**b**, Representative images showing $AC_{PPC}$ and $AC_{STR}$ neurons at the anterior (left, -2.3mm from

the bregma) and the posterior (right, -3.2mm from the bregma) AC. White lines, boundaries of

the AC; White dashed lines, boundaries between AC subareas. Scale bars, 500 μm.

**c**, Heatmaps representing percentages of labeled $AC_{PPC}$ neurons (left) or $AC_{STR}$ neurons (right)

in each subregion of the AC at different anterior-posterior coordinates.

**d**, Percentages of labeled neurons in the AC. Circles, data from individual brain samples.

**e**, Schematic illustration of *in vivo* calcium imaging of the $AC_{PPC}$ neurons. 127 neurons were

imaged from 6 mice.

**f**, Left, mean calcium traces of $AC_{PPC}$ neurons (green) and mean locomotion speeds of mice

(black) aligned to the locomotion onset. Right, distribution of Pearson correlation coefficients

between locomotion speeds and calcium activities of individual neurons at locomotion onsets.

The red triangle represents the mean correlation coefficients across all neurons. A vertical

dotted line indicates zero. Wilcoxon signed-rank test.

**g**, Same for **f**, but for the locomotion offset. Wilcoxon signed-rank test.

**h**, Calcium activities of $AC_{PPC}$ neurons in the incongruent trials at stationary state. Black line

and shades, for visual-dominat trials; Gray line and shades, for auditory-dominant trials.

Calcium activities between the two dominant trials were significantly different at every time

points ($P < 0.0001$). Wilcoxon signed-rank test.

**i**, Same for **h**, but for moving state. Calcium activities between the two dominant trials were

significantly different at every time points ($P < 0.0001$). Wilcoxon signed-rank test.

**j**, Mean response amplitudes of $AC_{PPC}$ neurons and auditory dominance rates of mice in $A_{NG}V_G$

(left) and $A_G V_{NG}$ (right) trials at different locomotion states (N = 6). Dots connected with gray

lines, individual mice data. Red crosses, means ± standard deviations. Wilcoxon signed-rank

test.

**k-p**, Same as **e-j**, but for the $AC_{STR}$ neurons (214 neurons from 6 mice).

Data are presented as means ± SEM except for **j,p**. NS (not significant), $*P < 0.05$, $****P <$

0.0001.

1227

**Fig. 5 | Double dissociation of AC_PPC and AC_STR neurons in perceptual decision-making.**

**a,** Schematic for optogenetic inhibition of the AC_PPC neurons in stationary mice performing audiovisual discrimination task.

**b,** Performance changes by optogenetic inactivation of the AC_PPC neurons in the auditory, visual, and congruent trials (N = 7 mice). Gray lines, individual mice. Inhibition, sessions with light on the AC; Control, sessions with light on the dental cement. Wilcoxon signed-rank test for each data.

**c,** Changes in auditory-dominant rates by AC_PPC inactivation in the incongruent trials. Wilcoxon signed-rank test for each data.

**d,** Changes in Hit and FA rates by AC_PPC inactivation in the auditory trials from the inhibition sessions. Off, laser off trial; On, laser on trial. Wilcoxon signed-rank test.

**e,** Changes in lick rates by AC_PPC inactivation in the incongruent trials from the inhibition sessions. Wilcoxon signed-rank test.

**f-j,** Same as **a-e**, but for optogenetic inactivation of the AC_STR neurons (N = 6 mice). Gray lines, individual mice; Bars, means ± SEM. Statistical comparisons with zero were indicated above individual bars. NS (non-significant), *$P < 0.05$.

1244

**Fig. 6 | M2-to-AC projection inhibits AC_PPC neurons and induces visually dominant decisions during locomotion.**

**a**, Schematic illustration of anterograde labeling M2 axons in the AC. Inset, an example coronal brain slice showing virus injection site at M2. Scale bar, 1 mm.

**b**, A representative image of virus-expressing M2 axons in the subregions of anterior AC. Scale bar, 500 μm.

**c**, Percentages of the number of labeled neurons (black) and the intensity of M2 axons (green) in the subregions of anterior AC. Solid line, AC_PPC neurons; dotted line, AC_STR neurons.

**d**, Schematic illustration of *in vivo* calcium imaging of M2_AC neurons.

**e**, Distribution of Pearson correlation coefficients between calcium activities and locomotion onsets across individual M2_AC neurons. 85 neurons from 5 mice. Wilcoxon signed-rank test.

**f**, Top, mean activity of $M2_{AC}$ neurons after locomotion onsets. A gray vertical line, locomotion onset; lighter gray shade, early locomotion period (0 – 1.5 s from the onset); darker gray shade, late locomotion period (> 3s from the onset). Bottom, mean activities of $M2_{AC}$ neurons in the early and late locomotion periods. Wilcoxon signed-rank test.

**g**, Schematic illustration of *in vivo* calcium imaging of the projection-specific AC neurons together with optogenetic activation of M2 axons in the AC using the nVoke system.

**h,i**, Average calcium traces (**h**) and mean response amplitudes (**i**) of the $AC_{PPC}$ neurons (42 neurons from 5 mice) in response to the 10kHz (left) and 5kHz (right) pure tones with (dotted) or without (solid) optogenetic activation of the M2 axons. Wilcoxon signed-rank test.

**j,k**, Same as **h,i**, but for the $AC_{STR}$ neurons (61 neurons from 5 mice). Wilcoxon signed-rank test.

**l**, Schematic illustration of optogenetic inhibition of the M2 axons in the AC in task-performing mice at moving states.

**m**, Performance changes by optogenetic inhibition of the M2 axon in the AC during the auditory, visual, and congruent trials (N = 6 mice). Gray lines, individual mice. Inhibition, sessions with light on the AC; Control, sessions with light on the dental cement. Wilcoxon signed-rank test for each data.

**n**, Changes in auditory dominance rates by the M2 axon inhibition in the AC during the incongruent trials. Wilcoxon signed-rank test for each data.

**o**, Summary illustration of the brain areas interacting with each other to resolve audiovisual conflicts and make multisensory decisions at different locomotion states.

Data are presented as means ± SEM. NS (not significant), $*P < 0.05$, $**P < 0.01$, $***P < 0.001$.

 **Extended data figures and legends**

1279

**Extended Data Fig. 1 | Analysis of the effects of pre-stimulus locomotion speeds, trial**

**history, and pupil size on the auditory dominance rates.**

**a**, Auditory dominance rates at different pre-stimulus locomotion speeds in $A_{NG}V_G$ (gray solid

line), $A_GV_{NG}$ (gray dashed line), and total incongruent trials (black) from the keep-moving state.

One-way ANOVA with Fisher's LSD test.

**b**, Auditory dominance rates in incongruent trials after different types of unisensory trials. Gray

lines indicate individual mice. Wilcoxon signed-rank test.

**c**, Example snapshot of a mouse head on the treadmill (top) and automatically estimated pupil

boundaries at three different time points (bottom).

**d**, Mean pre-stimulus pupil sizes in the incongruent trials when mice showed auditory (A) or

visual (V) dominance at different locomotion states (N = 6). Gray lines indicate individual mice.

Data are presented as means ± SEM. NS (not significant).

1292

**Extended Data Fig. 2 | Unisensory response changes during locomotion and additional**

**dPCA analysis of the PPC excitatory neurons.**

**a**, Changes in unisensory responses of the PPC excitatory neurons by locomotion.

**b**, Differences in the top 3 PC values in lick trials compared to no-lick trials in the stimulus PC

axes after dPCA (N = 20). Note that the PCs in the stimulus axes were not different between

lick and no-lick decisions of mice (Δ Score ≈ 0).

**c**, Comparisons of the top 3 PC values in the decision axes. Colors indicate unisensory types

as shown in **d**. Note that decision PCs clearly dissect lick (solid line) and no-lick (dotted line)

trials.

**d**, Same as **c**, but for the PCs in the interaction axes. Note that interaction PCs clearly dissect

correct and incorrect trials

**e**, Percentage of explained variances of top 20 demixed PCs from dPCA of the stationary (left,

N = 10) and the moving sessions (right, N = 10). Asterisks indicate statistical comparisons

between the stimulus PCs and other PCs. One-way ANOVA with Fisher's LSD test.

**f**, Mean Euclidean distances in a pair of the unisensory trajectories in the stimulus subspace

isolated from stationary (left, N = 10) and moving (right, N = 10) sessions. Note that the

Euclidean distances are larger between visual Go and No-go trajectories than those between

auditory Go and No-go trajectories in both conditions. One-way ANOVA with Fisher's LSD

test.

Data are presented as means ± SEM. NS (not significant), \*\*\*$P < 0.001$; \*\*\*\*$P < 0.001$.

1313

**Extended Data Fig. 3 | Additional dPCA results from the PPC neurons in the multisensory trials.**

**a**, Projection distances of neural trajectories in the stimulus subspace during congruent audiovisual and visual trials at different states (N = 10). One-way ANOVA, $P$ = 0.48.

**b**, Euclidean distances between stimulus trajectories of Go and No-go trials (Go/No-go distances) in the congruent (Con) and visual (Vis) trials at different states (N = 10). One-way ANOVA, $P$ = 0.97.

**c**, Classification accuracies of the visual Go and No-go trials and congruent Go and No-go trials in the stationary (left) and the moving (right) states (N = 10).

**d**, Mean Go/No-go classifier accuracies during the 0.5 – 1s period after the stimulus onset (N = 10). One-way ANOVA, $P$ = 0.53.

**e,f**, Same as **a,b**, but for the incongruent audiovisual trials (N = 10). Note that the incongruent trials were sorted into visual-dominant and auditory-dominant trials. Two-way ANOVA with Fisher's LSD test. **e**, Trial type, $P < 0.0001$; Locomotion, $P$ = 0.61; Interaction, $P$ = 0.28. **f**, Dominance, $P < 0.0001$; Locomotion, $P$ = 0.076; Interaction, $P$ = 0.39.

**g**, Euclidean distances between the incongruent audiovisual trajectories and the unisensory

visual or auditory trajectories of all PCs in the stimulus subspace (N = 10). Two-way ANOVA

with Fisher's LSD test. Pair type, $P < 0.0001$; Locomotion, $P = 0.033$; Interaction, $P = 0.68$.

**h**, Mean classifier accuracies (N = 20) in identifying the four types of incongruent trials (two

stimulus conditions with visual- or auditory-dominant decisions) by using neural trajectories

in the stimulus, decision, interaction, and independent subspaces. One-way ANOVA with

Fisher's LSD test.

Data are presented as means ± SEM. NS (not significant), $*P < 0.05$, $**P < 0.01$, $***P < 0.001$,

$****P < 0.0001$.

1338

**Extended Data Fig. 4 | Locomotion-related activities in the PPC excitatory neurons.**

**a**, Left, the mean calcium trace of the $A_G$ neurons (green for $A_G$ neurons that showed weaker

auditory responses during locomotion (as shown in Extended Data Fig. 2a); gray for all $A_G$

neurons) and the mean locomotion speed of mice (black) aligned to the locomotion onset. Right,

a histogram of Pearson correlation coefficients between the activity changes in those neurons

and the locomotion speeds during -2 ~ 2 s of the locomotion onset. Arrowheads, mean

correlation coefficients. Wilcoxon signed-rank test.

**b**, Same as **a**, but for the locomotion offsets. Wilcoxon signed-rank test.

**c,d**, Same as **a,b**, but for the $A_{NG}$ neurons. Wilcoxon signed-rank test.

Traces with shades are means ± SEM. NS (not significant), \*\*\**P* < 0.001.

1349

1350

**Extended Data Fig. 5 | Neural activities of the AC$_{PPC}$ neurons at different locomotion**

**states.**

**a**, Mean calcium traces of the AC$_{PPC}$ neurons (n = 127) in the trials with the A$_G$ stimulus (left)

and with the A$_{NG}$ stimulus (right) at different locomotion states. Note that the responses were

suppressed when mice showed longer locomotion.

**b**, Same as **a**, but for the AC$_{PPC}$ neurons in the trials with the V$_G$ (left) and V$_{NG}$ stimuli (right)

at different locomotion states.

**c**, Mean response amplitudes of the AC$_{PPC}$ neurons (n = 127) in the unisensory trials at different

locomotion states. Two-way ANOVA test with Fisher's LSD test. Stimulus, $P < 0.0001$;

Locomotion, $P < 0.0001$; Interaction, $P < 0.0001$;

**d**, Neural trajectories of the $AC_{PPC}$ neurons in three PCs of the stimulus subspace from the

dPCA. Four colors in trajectories represent unisensory trial types. Solid lines, stationary

conditions; dotted lines, moving conditions.

**e,f**, Projections distances of the unisensory trajectories (**e**) and distances between Go and No-

go trajectories (**f**) at different locomotion states (N = 6 for each data). Two-way ANOVA test

with Fisher's LSD test. **e**, Stimulus, $P < 0.0001$; Locomotion, $P < 0.0001$; Interaction, $P <$

$0.0001$. **f**, Stimulus, $P < 0.0001$; Locomotion, $P < 0.0001$; Interaction, $P < 0.0001$;

**g**, Mean classifier accuracies of auditory stimuli classification (N = 6). Two-way ANOVA with

Fisher's LSD test. Stimulus, $P < 0.0001$ ; Locomotion, $P < 0.0001$ ; Interaction, $P = 0.0009$.

**h-j**, Same as **c,e,f**, but for the incongruent trials (N = 6). Two-way ANOVA test with Fisher's

LSD test. **h**, Trial type, $P < 0.0001$; Locomotion, $P < 0.0001$; Interaction, $P < 0.0001$. **i**, Trial

type, $P < 0.0001$; Locomotion, $P < 0.0001$; Interaction, $P < 0.0001$. **j**, Dominance, $P < 0.0001$;

Locomotion, $P < 0.0001$; Interaction, $P < 0.0001$.

Gray scales in lines and bars represent locomotion states. Light gray, stationary (S); dark gray,

start-moving (SM); black, keep-moving (KM). Data are presented as means ± SEM. NS (not

significant), $\#P < 0.1$, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$.

**Extended Data Fig. 6 | Neural activities of the AC_STR neurons at different locomotion states.**

**a**-**j**, Same as Extended Data Fig. 5, but for the AC_STR neurons.

**a**, Mean calcium traces of the AC_STR neurons (n = 214) in the trials with the A_G stimulus (left) and with the A_NG stimulus (right) at different locomotion states.

**b**, Same as **a**, but for the AC_STR neurons in the trials with the V_G (left) and V_NG stimuli (right) at different locomotion states.

**c**, Mean response amplitudes of the AC_STR neurons (n = 214) in the unisensory trials at different locomotion states (n = 214 neurons). Two-way ANOVA test with Fisher's LSD test. Stimulus, $P < 0.0001$; Locomotion, $P = 0.87$; Interaction, $P = 0.99$;

**d**, Neural trajectories of the $AC_{STR}$ neurons in three PCs of the stimulus subspace from the dPCA. Four colors in trajectories represent unisensory trial types. Solid lines, stationary conditions; dotted lines, moving conditions.

**e,f**, Projections distances of the unisensory trajectories (**e**) and distances between Go and No-go trajectories (**f**) at different locomotion states (N = 6 for each data). Two-way ANOVA with Fisher's LSD test. **e**, Stimulus, $P < 0.0001$ ; Locomotion, $P = 0.49$; Interaction, $P = 0.99$. **f**, Stimulus, $P < 0.0001$ ; Locomotion, $P = 0.49$; Interaction, $P = 0.99$.

**g**, Mean classifier accuracies of auditory stimuli classification (N = 6). Two-way ANOVA with Fisher's LSD test. Stimulus, $P < 0.0001$ ; Locomotion, $P = 0.70$; Interaction, $P = 0.81$.

**h-j**, Same as **c,e,f**, but for the incongruent trials (N = 6). Two-way ANOVA test with Fisher's LSD test. **h**, Trial type, $P = 0.77$; Locomotion, $P = 0.97$; Interaction, $P = 0.98$. **i**, Trial type, $P < 0.0001$ ; Locomotion, $P = 0.94$; Interaction, $P = 0.99$. **j**, Dominance, $P = 0.99$ ; Locomotion, $P = 0.98$; Interaction, $P = 0.97$.

Gray scales in lines and bars represent locomotion states. Light gray, stationary (S); dark gray, start-moving (SM); black, keep-moving (KM). Data are presented as means ± SEM. NS (not significant), ****$P < 0.0001$.