Genome sequence of the ornamental plant *Digitalis purpurea* reveals the molecular basis of flower color and morphology variation

Jakob Maximilian Horz[1,2], Katharina Wolff[1], Ronja Friedhoff[2], Boas Pucker[1*]

[1] Plant Biotechnology and Bioinformatics, IZMB, University of Bonn, Kirschallee 1, 53115 Bonn, Germany

[2] Plant Biotechnology and Bioinformatics,TU Braunschweig, Mendelssohnstrasse 4, 38106 Braunschweig, Germany

* corresponding author: pucker@uni-bonn.de

ORCIDs:
JMH: 0009-0007-7367-9439
KW: 0000-0002-3616-0482
RF: 0009-0002-0270-4661
BP: 0000-0002-3321-7471

Abstract

Digitalis purpurea (foxglove) is a widely distributed ornamental plant and the producer of the biomedical compound digoxin. Here, we present a long read sequencing-based genome sequence of a magenta flowering *D. purpurea* plant and a corresponding prediction of gene models. The high assembly continuity is indicated by the N50 of 4.3 Mbp and the completeness is supported by discovery of about 96% complete BUSCO genes. This genomic resource paves the way for an in-depth investigation of the flower pigmentation of *D. purpurea*. Structural genes of the anthocyanin biosynthesis and the corresponding transcriptional regulators were identified. The comparison of magenta and white flowering plants revealed a large insertion in the anthocyanidin synthase gene in white flowering plants that most likely renders this gene non-functional and could explain the loss of anthocyanin pigmentation. Furthermore, we found a large insertion in the *DpTFL1/CEN* gene to be likely responsible for the development of large terminal flowers.

Keywords:

Genome sequence, foxglove, flavonoids, anthocyanins, pigmentation, ONT sequencing, MYB

Introduction

Digitalis purpurea, commonly known as foxglove, is a biennial plant in the Plantaginaceae family [1]. The species is native to Europe, but appears in many other parts of the world. D. purpurea has been used for centuries for its medicinal properties, particularly for treating heart conditions such as atrial fibrillation and congestive heart failure [2, 3]. The plant contains a variety of cardiac glycosides, with the most important being digoxin [2]. These compounds work by increasing the force and efficiency of heart contractions, which are used to treat heart failure and other cardiac conditions. D. purpurea and the close relative D. lanata are cultivated for the extraction of cardenolides [1]. Scientists have been studying the plant for many years to better understand the chemistry of the cardiac glycosides it produces and their effects on the human body. Many experiments explored the potential of tissue cultures to propagate plants and to improve the yield of medicinal compound production [1]. A cytochrome P450 CYP87A4 is crucial for the biosynthesis of digoxin [3]. The identification of all genes required for a biosynthesis process is the first step towards production in a heterologous system.

Besides its medicinal value, D. purpurea is also famous for beautiful flowers (**Fig. 1a**). It is a tall plant that can reach up to six feet in height and produces striking magenta flowers on tall spikes. Various Digitalis species display a plethora of diverse flower colors and shapes. While the plant is visually appealing, it is considered toxic due to the formation of specialized metabolites like the cardenolides. Nonetheless, D. purpurea is widespread as an ornamental plant. Contributing to the flower colors are anthocyanins, the product of one specific branch of the flavonoid biosynthesis (**Fig. 1b**). The substrate of the flavonoid biosynthesis is the aromatic amino acid phenylalanine, which is channeled through the general phenylpropanoid pathway and the flavonoid biosynthesis towards anthocyanins [4, 5]. Major branches of the flavonoid biosynthesis lead to colorful anthocyanins, colorless flavonols, colorless flavones, and proanthocyanidins which appear dark after oxidation. The anthocyanin biosynthesis shares several enzymatic steps with other pathways like the proanthocyanidin biosynthesis. Dihydroflavonol 4-reductase (DFR), anthocyanidin synthase (ANS), UDP-glucose dependent 3-O-anthocyanidin-glucosyltransferase (UFGT), and other decorating enzymes are often considered as the anthocyanin biosynthesis branch [4, 6] (**Fig. 1c**). A recent study revealed anthocyanin-related glutathione S-transferase (arGST) as an additional enzyme in the anthocyanin biosynthesis [7]. After biosynthesis of anthocyanins at the endoplasmic reticulum, a transport through the cytoplasm and import into the central vacuole is required for long term storage. This process involves several proteins including two tonoplast-localized transporters and a tonoplast-localized ATPase for generation of a proton gradient [5, 8, 9]. A previously postulated 'ligandin', that might protect anthocyanins during the transport through the cytoplasm [10], was recently characterized as an enzyme in the cyanidin biosynthesis, which makes an involvement in the anthocyanin transport less likely [7]. Besides direct transport through the cytoplasm, there are several studies that report an involvement of vesicles in the anthocyanin transport in different plant species [5, 11–14].

The pigmentation of a plant structure is not only determined by the activity of the anthocyanin biosynthesis, but also by the activity of pathways that compete for the same substrate. High activity of the flavonol biosynthesis can deplete the available substrate of the anthocyanin

biosynthesis branch [15]. Subtle differences in substrate specificity of the competing
enzymes ,e.g. flavonol synthase (FLS) and DFR, can determine the pattern of synthesized
flavonoids [16, 17]. Differences in expression of the first committed genes *DFR* and *FLS*,
respectively, appear as a decisive factor in controlling the activity of the anthocyanin and flavonol
biosynthesis branch [17].

Activity of different branches of the flavonoid biosynthesis is mostly controlled at the
transcriptional level through regulation of gene expression by transcription factors [18–21].
Decades of research turned the regulation of the flavonoid biosynthesis into a model system for
research on cis- and trans-regulatory elements in eukaryotes. The anthocyanin biosynthesis
genes are activated by a complex comprising three different proteins: a R2R3-MYB protein, a
bHLH protein, and a WD40 protein [21–23]. Different subgroup 6 MYBs and subgroup IIIf bHLHs
can be included in this complex leading to a diverse set of different complexes which differ in their
ability to activate certain target genes [24, 25]. The WD40 component is often TTG1, but LWD1
and LWD2 can also participate in complex formation [26]. An additional WRKY component (TTG2)
has been suggested to specifically increase the expression of genes associated with the
anthocyanin transport into the vacuole [21, 27]. Anthocyanin biosynthesis-regulating MYBs have
been reported in different MYB clades. PAP1 (MYB75)/PAP2 (MYB90)/PAP3 (MYB113)/PAP4
(MYB114) are well known anthocyanin regulators in *Arabidopsis thaliana* [23, 28, 29]. While most
anthocyanin regulating MYBs are activators, the incorporation of subgroup 4 MYBs into the MBW
complex can result in an inhibitory function [30].

Pigmentation differences between plants of the same species are often due to genetic variations
affecting the functionality or gene expression of the MYB and bHLH transcription factors [31–34].
This results in extremely low or no transcriptional activation of specific flavonoid biosynthesis
genes under certain conditions. However, activation of the structural genes in the context of a
different pathway is still possible [34]. Loss of function or gene loss affecting biosynthesis genes or
transporters involved in the biosynthesis pathway are rare, but can explain deeper loss of the
anthocyanin pigmentation in the Caryophyllales and might be irreversible [35]. Changes in flower
color can affect the attraction of pollinating animals and are thus subject to selection. For example,
the transition of purple blue to orange-red in *Ipomoea quamoclit* L. probably facilitated the shift
from bee pollination to hummingbird pollination [36]. A study in *Silene littorea* discovered that a
lack of anthocyanins exclusively in flower petals appeared as a stable polymorphism, but
complete loss of anthocyanins in all plant parts was rare suggesting anthocyanins are the target of
selection in photosynthetic tissues [37].

The flower pigmentation of *D. purpurea* is not uniform, but characterized by intensely pigmented
dark spots on the bottom petals (**Fig. 1c**). Petal spots have been reported in other plant species
including *Gorteria diffusa* [38], *Kohleria warszewiczii* [39], *Clarkia gracilis* [40], *Lilium leichtlinii*
[41], and *Mimulus guttatus* [42]. Pigmentation spots on petals can be caused by the accumulation
of high levels of anthocyanins (cyanidin derivatives) [40, 43]. These spots can be surrounded by a
white region characterized by high levels of flavonols, the products of a pathway competing with
anthocyanin biosynthesis [44]. Formation of intensely pigmented spots can be explained by a
system involving a local R2R3-MYB activator (NEGAN) and a lateral R3-MYB repressor (RTO)

[42]. NEGAN interacts with a bHLH and a WD40 partner (MBW complex) to activate *RTO*
expression and RTO represses *NEGAN* expression by sequestering bHLH proteins required for
the MBW complex [42]. RTO proteins appear to travel through the endosomal system to other
cells, where they act as repressors [42]. Another study in *C. gracilis* reported an anthocyanin
activating MYB (CgsMYB12) as the causal gene for the formation of a single large spot [45]
indicating that alternative mechanisms have evolved to control the formation of pigmentation
spots. Numerous functions have been described for such spots including an attraction and
guidance of pollinating insects to increase the number of visits and to make each event more
successful [39, 46–49].
132
133

134
Fig. 1: Pictures showing magenta (a) and white (b) flowers of *Digitalis purpurea* (foxglove). The
anthocyanin biosynthesis is displayed as the central branch of the flavonoid biosynthesis (c).
Abbreviations: CHS, chalcone synthase; CHI, chalcone isomerase; F3H, flavanone 3-
hydroxylase; DFR, dihydroflavonol 4-reductase; ANS, anthocyanidin synthase; arGST,
anthocyanin-related glutathione S-transferase; 3GT, UDP-glucose anthocyanidin 3-O-

glucosyltransferase; MYB, Myeloblastosis protein; bHLH, basic helix-loop-helix; WD40, proposed
scaffolding protein.

The inflorescence of *D. purpurea* is usually constructed in the form of an indeterminate spike with
zygomorphic tubular flowers on the side [50]. But in rather rare cases individuals develop large
terminal flowers at the top of the spike, rendering the inflorescence determinate. These
phenotypes show two types of flowers in the same plant: the normal zygomorphic flowers at the
lower part of the spike and one actinomorphic structured flower at the top (**Fig. 2**). Keeble *et al*.
investigated the heredity of this special trait and found it to be inherited recessively [51]. The
characterization of the genetics behind the development of a terminal flower in *D. purpurea* has
not been investigated yet. The occurrence of terminal flowers in plants with otherwise
indeterminate inflorescence architecture has been described in different species like *Antirrhinum*
[52] and Arabidopsis [53]. In *Antirrhinum*, a loss of function mutation in the *CEN*
(*CENTRORADIALIS*) gene leads to the development of a terminal flower [52]. The functional *CEN*
gene is expressed in the subapical meristem and disrupts the initiation of flower development [52].
Therefore, CEN is a flowering repressor. The *Arabidopsis* ortholog *TERMINAL FLOWERING 1*
(*TFL1*) plays a similar role [54]. The mutant *tfl1* no longer represses flowering in the apical
meristem, so it changes its fate and develops into flower primordia and subsequently into a flower
[53, 55]. Both genes encode a protein belonging to the family of phosphatidylethanolamine
binding proteins. Therefore, they are not conventional regulators of the transcription or translation
of any specific target genes [56]. They have been shown to interact directly with other proteins so
they might be part of a more elaborate signaling system [56].

163
164   **Fig. 2**: Three terminal flowers. (A): A purple terminal flower with purple spots originated from the
165   Botanical Garden of TU Braunschweig. (B and C): White terminal flowering plants.

166
Here, we reported about the genome sequence of *D. purpurea* and the corresponding annotation.
As a proof of concept, we demonstrate the power of this genome sequence for the discovery of
specialized metabolism pathways by investigating the flavonoid biosynthesis. A comparison of
white and magenta flowering individuals revealed molecular differences in the flavonoid
biosynthesis genes that could explain the pigmentation loss in the white flowering plants. The

molecular basis of large terminal flowers was discovered through comparative genomics based on
this genome sequence.
174
175
176

Materials & Methods

Plant material and DNA extraction

Digitalis purpurea is a native species in Germany that is widespread at the borders of forests
and in gardens. We collected seeds of a white and magenta flowering plant in a garden close to
these coordinates 51.92351 N, 8.84441 E. Plants were cultivated under long day conditions
(16h light, 8h darkness) at 20°C. LEDs (Niello®, 300W) were used as the sole light source.
Plants were incubated in the dark for two days immediately prior to the sampling of leaves for
the DNA extraction to reduce polysaccharide content. Young leaves were harvested for the
DNA extraction. The DNA extraction was performed with a CTAB-based protocol as previously
described [57]. Quality and purity of the extracted DNA were assessed through NanoDrop
measurement, agarose gel electrophoresis, and Qubit measurement. Short fragments (<25kb)
were depleted with the Short Read Eliminator kit (Pacific Biosciences) (see [58] for a detailed
workflow).
190

Nanopore sequencing

The library preparation for sequencing was started with 1µg of high molecular weight DNA. The
initial DNA repair was performed with two enzyme mixes of the NEBNext® Companion Module
Oxford Nanopore Technologies® (ONT) Ligation Sequencing (E7180L) following the suppliers'
instructions. All following steps were performed according to the SQK-LSK109 protocol (ONT).
Sequencing was performed with MinIONs using R9.4.1 flow cells. Upon blockage of a
substantial number of nanopores, a wash step was performed with the EXP-WSH004 following
ONT's instructions to recover the blocked nanopores. Afterwards, a fresh library was loaded
onto the flow cell. High accuracy basecalling was performed with guppy v6.4.6+ae70e8f (ONT)
and minimap2 v2.24-r1122 [59] on a graphical processor unit (GPU) in the de.NBI cloud.
201

De novo genome sequence assembly

A *de novo* genome sequence assembly was generated based on all long reads of a magenta
flowering plant. NextDenovo v2.5.0 [60], Shasta 0.10.0 [61], and Flye 2.9.1-b1780 [62] were
deployed to generate assemblies. Based on completeness, continuity, and assumed
correctness, an assembly produced by NextDenovo v2.5.0 with read_cutoff = 1k, genome_size
= 1g, and seed_depth = 30 was selected as the best representation of the genome. Additional
polishing of this assembly was performed with NextPolish v1.4.1 [63] based on read mappings

that were generated with minimap2 [59] and samtools v1.15.1 [64]. The genomic BUSCO
assessment command contained the arguments '--long --limit 10 -sp arabidopsis', 10^{-3} as e-
value cutoff for BLAST, and eudicotyledon_odb10 (embryophyta) as the reference. AUGUSTUS
v3.2.2 [65, 66] was used for the gene prediction as part of BUSCO.
213

Structural annotation

Paired-end RNA-seq reads (Additional file 1) were retrieved from the Sequence Read Archive
via fastq-dump [67] and aligned to the assembled *D. purpurea* genome sequence with STAR
v2.5.1b [68, 69] in 2-pass-mode using a minimal similarity of 95% and a minimal alignment
length of 90% as previously described [70] to provide hints for the gene prediction process.
Gene models were predicted with GeMoMa v1.9 [71, 72] using the following arguments: pc=true
pgr=true p=true o=true GeMoMa.c=0.4 GeMoMa.Score=ReAlign Extractor.r=true
GAF.f="start=='M' and stop=='*' and (isNaN(score) or score/aa>='0.75')". In addition to RNA-seq
hints, the annotation of the following datasets was supplied as hints: *Salvia hispanica*
(GCF_023119035.1), *Rehmannia glutinosa* (GCA_016081115.2), *Sesamum indicum*
(GCF_000512975.1), *Salvia miltiorrhiza* (GCF_028751815.1), *Penstemon davidsonii*
(GCA_034814905.1), and *Penstemon smallii* (GCA_046254845.1). Predicted gene models
were filtered via GeMoMa (f="start=='M' and stop=='*' and (isNaN(tie) or tie>0) and tpc>0 and
aa>50" atf="tie>0 and tpc>0") and renamed. BUSCO v3.0.2 [73, 74] was used to assess the
completeness of the genome sequence and the completeness of the gene prediction. The
completeness of predicted gene models was assessed with BUSCO v5.7.1 by running in
proteome mode with default parameters.
231

Functional annotation

Homologs in *Arabidopsis thaliana* were identified for the predicted peptide sequences to transfer
annotation terms from this model organism to the *D. purpurea* sequences. Previously developed
Python scripts [70, 75] were modified to enable the efficient identification of Reciprocal Best
BLAST hits (RBHs) supplemented with best BLAST hits for query sequences that are not
assigned to a RBH partner. BLASTp v2.13.0+ [76, 77] was run with an e-value cutoff of 0.0001.
This approach allows an efficient identification of putative orthologs and forms the basis for the
transfer of functional annotation details across species borders. *Arabidopsis thaliana*
TAIR10/Araport11 annotation terms were automatically transferred to the predicted *D. purpurea*
sequences. This workflow is implemented in the Python script construct_anno.py [78]. Enzyme
and transporter encoding genes of the flavonoid biosynthesis were identified using KIPEs v3.2.6
with the flavonoid biosynthesis data set v3.4 [6, 79] with additional parameters listed in
Additional file 3. Flavonoid biosynthesis controlling MYB transcription factors were annotated
with the MYB_annotator v1.0.3 [80]. The following parameters were used for the MYB
annotation: initial candidate identification via BLASTp v2.13.0+ [76, 77], alignment construction
with MAFFT v7.475 [81], phylogenetic tree construction with FastTree v2.1.10 [82], minimal
BLASTp alignment length of 75 amino acids, and 10 neighbors were used for the

ingroup/outgroup classification (see Additional file 4 for details). The bHLH transcription factors
were annotated using the bHLH_annotator v1.04 [83]. The following parameters were used for
the bHLH annotation: identification of initial candidates via BLASTp v2.13.0+ [76, 77],
alignment with muscle v5.1 [84], phylogenetic tree construction with FastTree v2 [82], a minimal
BLASTp hit similarity of 40%, minimal BLASTp hit alignment length of 80 amino acids, and
considering 10 neighbors for the ingroup/outgroup classification (see Additional file 5 for
details).
256

Analysis of gene expression

258   Fastq-dump [67] was applied to retrieve the FASTQ files of *D. purpurea* RNA-seq experiments
259   from the Sequence Read Archive (SRA). Gene expression was quantified using kallisto v0.44.0
260   [85] to process all RNA-seq data sets (Additional file 1) based on the predicted coding
261   sequences. Individual count tables were merged with a customized Python script and served as
262   the basis for a visualization of gene expression as previously described [17, 78].

Identification of sequence variants associated with anthocyanin loss

Reads of two magenta flowering plants and two white flowering plants (Additional file 5) were
aligned against the reference genome sequence of a magenta flowering plant with minimap
v2.24-r1122 [59]. The resulting BAM files were indexed with samtools v1.13 [64]. Candidate
genes associated with the flavonoid biosynthesis were manually inspected for systematic
sequence differences via Integrative Genomics Viewer (IGV) v2.15.4 [86]. The same method
was applied to one terminally flowering individual and three indeterminate flowering individuals

Genotyping of a *Digitalis purpurea* population

272   A population of 89 plants was subjected to genotyping via PCR for the discovered mutations
273   responsible for the white flower color or the changed flower morphology, respectively. The
274   plants were first grown under long day conditions (16h light, 8h darkness) at 20°C. LEDs
275   (Niello®, 300W) were used as the sole light source during that time. After four months the plants
276   were moved outside the plant cultivation chamber and placed into two garden beds.
277
278   DNA for subsequent genotyping via PCR was extracted based on the GABI-Kat CTAB DNA
279   preparation protocol [87]. This protocol, designed for *A. thaliana,* was adapted for the use on *D.*
280   *purpurea*: Only one young leaf was harvested from each plant. After extraction the DNA
281   samples were stored at -20 °C.
282
283   PCR was conducted with oligonucleotides flanking the *TFL1*/*CEN* locus and a 106 bp deletion
284   linked to the mutant *ans* allele, respectively (Additional file 9, Additional file 10). Sizes of the
285   PCR products were analyzed via agarose gel electrophoresis and revealed the genotypes.
286

287 Initial phenotypes of the plants resulting from the *ANS* mutation were derived from the visual
288 inspection of the leaves as *D. purpurea* with a functional *ANS* would show magenta colored
289 petioles (Additional file 11). These phenotypes were later validated by adding information about
290 the flower color.
291

Results

Genome sequence assembly and structural annotation

A magenta flowering plant (DR1) was selected for sequencing and construction of a reference
genome sequence [88]. The rationale behind this selection was the assumption that a magenta
flowering plant should have all genes required for the biosynthesis of anthocyanins which are
most likely responsible for the magenta flower pigmentation. For comparison, another magenta
flowering plant (DR2) and two white flowering plants (DW1, DW2) were subjected to nanopore
long read sequencing (Additional file 5).
300
301 Different assemblers were evaluated with respect to their performance in this study (**Table 1**).
302 The assembly produced by NextDenovo2 was selected as the representative *D. purpurea*
303 genome sequence due to the large assembly size of 940 Mbp, the high continuity
304 (N50=4.3Mbp), and the high completeness of 96% of complete BUSCOs.
305
306 **Table 1**: Comparison of assembler performance for the analysis of *Digitalis purpurea* long read
307 sequencing data of the magenta flowering plant DR1.

Assembly ID	**DR1_v1**	**DR1_v2**	**DR1_v3**
Assembler	NextDenovo (v2.5.0)	Shasta (v0.10.0)	Flye (2.9.1-b1780)
Size	940 Mbp	773 Mbp	1379 Mbp
# contigs	684	4883	3738
N50	4.3 Mbp	0.9 Mbp	1.9 Mbp
N90	464 kbp	71 kbp	132 kbp
BUSCO	C:95.7% [S:63.5%,D:32.2%], F:0.6%, M:3.7% n:2326	C:94.4% [S:68.7%,D:25.7%], F:0.6%, M:5.0%, n:2326	C:25.9% [S:23.3%,D:2.6%], F:6.0%, M:68.1% n:2326

308
Only the representative genome sequence DR1_v1 was subjected to a gene prediction process
that resulted in 35,916 protein encoding genes [89]. A completeness check revealed 96.3%
complete BUSCO genes in this annotation. The presence of 55.8% duplicated BUSCO genes
suggested that a large fraction of the assembly represents separate haplotypes. This was

supported by a coverage analysis that found an average coverage of 20x for large parts of the
assembly, while small parts showed a coverage of around 40x (Additional file 7).

315 Anthocyanin biosynthesis genes in *Digitalis purpurea*

Anthocyanins are well known pigments responsible for the coloration of many plant structures -
especially flowers. Therefore, the genes encoding enzymes of the anthocyanin biosynthesis
(*CHS*, *CHI*, *F3H*, *F3'H*, *DFR*, *ANS*, *arGST*, *3GT*) were identified (**Fig. 3**; Additional file 6).
Additionally, the transcription factors controlling the anthocyanin biosynthesis genes in *D.*
purpurea were investigated. Genes belonging to the MYB75/PAP1 (DP112203, DP103545,
DP109418) and MYB123/TT2 (DP102989) lineages were identified. In addition, a candidate for
the bHLH partner in the anthocyanin biosynthesis MBW regulatory complex (bHLH42/TT8) was
detected in *D. purpurea* (DP105472). An ortholog of the WD40 gene *TTG1* (DP111282) was
also identified in the phylogenetic analysis. In summary, all genes expected for a complete
anthocyanin biosynthesis pathway were successfully identified in the reference genome
sequence DR1_v1.
327
328

329
Fig. 3: *Digitalis purpurea* candidate genes associated with the anthocyanin biosynthesis
including the transcriptional regulators of the MBW complex. Yellow and magenta arrows
indicate that the biosynthesis of pelargonidin and cyanidin derivatives would be possible, while
the lack of a flavonoid 3',5'-dihydroxylase (*F3'5'H*) does not allow the synthesis of delphinidin
derivatives. *CHS*, naringenin-chalcone synthase; *CHI*, chalcone isomerase; *F3H*, flavanone 3-
hydroxylase; *F3'H*, flavonoid 3'-hydroxylase; *DFR*, dihydroflavonol 4-reductase; *ANS*,
anthocyanidin synthase; *arGST*, anthocyanin-related glutathione S-transferase; *3GT*,
anthocyanidin 3-O-glucosyltransferase; MYB, Myeloblastosis; bHLH, basic helix-loop-helix;
TTG1, TRANSPARENT TESTA GLABRA 1.

Additional support for candidate genes comes from a gene expression analysis that was
conducted across a wide range of different samples (**Fig. 4**, Additional file 1). Anthocyanin
biosynthesis genes are expected to be highly expressed in pigmented structures like flowers
and in particular within the intensely pigmented purple spots on the lower flower petal. A
consideration of the gene expression pattern enabled a reduction of the candidate gene set to
those gene copies which appear active in the anthocyanin-pigmented plant organs.

Genes encoding enzymes for all steps of the flavonoid and anthocyanin biosynthetic pathways
were found to contain at least one paralog that is transcriptionally active in petal tissues. Two
paralogous *F3'H* genes exhibited expression above baseline levels. *F3'Ha* (DP114378) was
strongly active in floral tissues, whereas *F3'Hb* (DP125257) showed elevated expression
primarily in root tissues (Additional File 12). Although roots typically do not accumulate
anthocyanins, *F3'Hb* is reported here for completeness.

The putative anthocyanin-related transcription factor MYB75 is represented by three paralogous
genes that displayed distinct expression profiles. *DpMYB75a* was active in the lower, upper, and
side petal tissues but not in the spot sample. In contrast, *DpMYB75b* was strongly active in spot
tissues and, consequently, also showed elevated expression in the lower petal samples, which
include the spot region. *DpMYB75c* exhibited a comparatively uniform expression pattern
across young floral tissues, with a modest increase in the spot samples.

Fig. 4: Expression plot showing the activity of anthocyanin biosynthesis associated genes across different petal tissues. The coloration indicates the z-score, the numbers inside the squares are the corresponding TPM-values. Abbreviations: *CHS*, chalcone synthase; *CHI*, chalcone isomerase; *F3H*, flavanone 3-hydroxylase; *F3'H*, flavonoid 3'-hydroxylase; *DFR*, dihydroflavonol 4-reductase; *ANS*, anthocyanidin synthase; *arGST*, anthocyanin-related glutathione S-transferase; *F3GT*, UDP-glucose flavonoid 3-O-glucosyltransferase; *A3GT*, UDP-

368 glucose anthocyanidin 3-O-glucosyltransferase; *A5GT*, UDP-glucose anthocyanidin 5-O-
369 glucosyltransferase; MYB, Myeloblastosis; bHLH, basic helix-loop-helix; *TTG1*,
370 *TRANSPARENT TESTA GLABRA 1*.

371 Sequence variants underlying pigmentation differences

All identified candidate genes associated with the anthocyanin biosynthesis were manually
inspected in a read mapping comparing magenta flowering and white flowering plants. The most
striking difference is an insertion of approximately 13.5 kb into the *ANS* gene (**Fig. 5**). This
insertion is homozygous in the two white flowering plants, while the magenta flowering plants
harbor an *ANS* allele without this insertion. Since the reference genome sequence DR1_v1
contains the 13.5 kb sequence, the automatic prediction of the *ANS* gene model is inaccurate.
The inserted sequence was misannotated as an intron of the *ANS* gene because no RNA-seq
reads were aligned to the inserted region. However, in the wildtype allele, the first exon extends
across the position of the insertion, indicating that the sequence cannot be removed by splicing
in the mutant allele. This strongly suggests that the insertion disrupts normal *ANS* transcript
structure rather than representing an intronic region (**Fig. 5**). The inserted sequence is most
likely a transposable element, as it occurs at multiple loci within the genome. In the region
containing the transposable element (TE) insertion, the coverage plots of magenta-flowering
plants do not display exactly half the read depth of adjacent genomic regions, as would be
expected for heterozygous individuals. Instead, the coverage is slightly elevated. Similarly, in
white-flowering plants, the coverage across the insertion site is also somewhat higher than that
of the surrounding sequences. These deviations from the expected pattern are likely caused by
misaligned reads originating from additional copies of the TE elsewhere in the genome.
390
Genotyping of 89 plants for the *ANS* mutation marker identified 14 plants with the homozygous
wildtype allele, 31 plants with a heterozygous genotype, and 44 plants with the homozygous
mutant allele. In the genotyped population, 61 plants have flowered. The flowering phenotype
was found to fit the genotype in 57 of these 61 plants, making up 93.4 %. Four plants showed a
phenotype mismatching the genotyping results. Two plants flowered white while being
heterogenous for the mutation and two plants showed magenta flowers but had a homozygous
mutant genotype.
398
399
400

401
Fig. 5: Comparison of the *ANS* locus in magenta and white flowering *D. purpurea* plants revealed
a large insertion that appears homozygous in the white flowering plants. The insertion is expected
to be a loss-of-function mutation that results in the loss of ANS enzymatic activity and thus a block
in the anthocyanin biosynthesis ultimately resulting in the white flower phenotype. A: Schematic of
the insertion. B: Screenshot from IGV with ONT read mappings of two magenta flowering plants
and two white flowering ones at the *ANS* locus. The red box highlights the inserted reads. The
green boxes highlight RNAseq read mapping coverage to corroborate the gene annotation.

409
410
411
412

Sequence variants leading to the formation of a large terminal flower

In the search for the genetic basis of the formation of large terminal flowers the sequencing
reads of one plant with a terminal flower were compared to three wildtype plants. Upon
investigation of the *Digitalis* orthologs of the *Arabidopsis TFL1* a 14337 bp insertion was found
starting only 4 base pairs after the start codon of the gene and leading to a premature stop
codon after 4 codons (**Fig. 7**). The insertion has a high similarity to the insertion in the *ans* gene
described above.
Out of 24 characterized plants 23 showed the phenotype fitting the predicted genotype. While
17 plants showed the wildtype floral architecture and contained at least one wildtype allele, 7
plants developed terminal flowers. Of those plants six had the homozygous mutant genotype
and one had a heterozygous genotype.

Fig. 7. Genomic region showing the insertion of 14 kb in the *DpFL1/CEN* (DP105791) in the reads of the plant with a terminal flower (red box). The plants with indeterminate flowering architecture show at least some reads without the insert.

Discussion

The genome sequence of *D. purpurea* and the corresponding annotation of protein encoding genes enabled the identification of genes associated with the anthocyanin biosynthesis. Long read sequencing of two white flowering individuals enabled the discovery of sequence variants that might explain the lack of pigmentation. *ANS* encodes the anthocyanidin synthase, a crucial enzyme for the anthocyanin biosynthesis, and shows a homozygous insertion of a large DNA fragment in both white flowering plants, while the two magenta flowering plants show at least one allele without this insertion. The consequence of this insertion is most likely a loss of ANS activity that disrupts the anthocyanin biosynthesis. A loss of ANS activity fits all observations regarding the presence/absence of flavonoids in *D. purpurea*.

We noticed that flavonols are produced in the white flowering plants (Additional file 8). There is an overlap regarding the enzymes required for flavonol and anthocyanin biosynthesis. CHS, CHI, F3H, and F3'H are shared between both pathways. The presence of flavonols indicates that the genes *CHS*, *CHI*, *F3H*, and *F3'H* are functional, because they are required for the flavonol biosynthesis [90–94]. This observation reduces the set of remaining structural genes in the anthocyanin pathways to *DFR*, *ANS*, *arGST*, and *UFGT*. While we cannot completely rule out that *DFR*, *arGST*, or *UFGT* contribute to the loss of anthocyanin pigmentation, we have not observed any evidence for systematic sequence variants in these genes. Several previous studies discovered loss-of-function mutations in *ANS* as the cause of anthocyanin pigmentation loss [95–97]. The above described *ans* mutation could be a universal explanation for the lack of pigments in white *D. purpurea* flowers. Because the tested population did not show a perfect correlation between *ANS* genotype and floral phenotype, it remains possible that additional mechanisms contribute to the absence of pigmentation in white-flowering *D. purpurea*

individuals. Another possibility is that the putative TE responsible for the mutation excised from
the insertion site after genotyping, resulting in a chimeric plant with distinct genotypes present in
different tissues. A comparable phenomenon is well known in maize and formed the basis of
Barbara McClintock's discovery of transposable elements [98]. In this system, the non-
autonomous element Ds inserts into mutable pigment-related genes such as *c1, bz1 or a1* [98,
99]*, thereby disrupting their function and giving rise to non pigmented kernels. Subsequent
excision of Ds during later developmental stages can restore gene activity in a subset of cells,
resulting in chimeric plants or in kernels exhibiting spotted or striped pigmentation, depending
on the developmental timing and cellular context of the excision event . A further plausible
explanation is that two plants grew in close proximity and physically merged, thereby
compromising the representativeness of the sampled tissue. As transcription factors are often
implicated in the loss of anthocyanin pigmentation [34], the identified candidate genes were
checked for sequence variants associated with the anthocyanin loss. However, no variants
fitting the phenotypes could be observed in these genes. Nevertheless, three paralogous
DpMYB75 candidates were found to have differential expression patterns in petal tissues.
DpMYB75b is strongly active in the dark purple spots which suggests its involvement in the
regulation of spot pigmentation DpMYB75a is expressed in the remaining floral samples but not
in the spot sample. This suggests the existence of two separate regulatory complexes involved
in the pigmentation of *Digitalis* flowers: One activating spot pigmentation (DpMYB75b) and one
activating background pigmentation (DpMYB75a).
Whether or not the absence of color and spot pigmentation in white flowering plants has any
effect on pollination efficiency remains to be tested. However, there are many examples of
plants where different visual cues in the flower lead to increased pollination efficiency [100–
102]. The visual cues can consist of flower forms indicating landing sites for insects, simply
pigmented flowers letting the flower stand out against the background or contrasting patterns
indicating landing-position and -direction for insects on the flower [103, 104]. This suggests that
the pigmentation patterns in *D. purpurea* flowers might contribute to increased attractiveness of
the flower for pollinators. The contrasting spots on the lower petal possibly indicate a landing
zone and the two UV fluorescent stripes at the base of the connation of the side petal with the
lower petal may act as nectar guides (Additional file 8).
483
484 The present study focuses on explaining the completely white flowering phenotype (**Fig. 1**), but
485 there are also cases of white flowers with magenta spots in *D. purpurea*. Similar patterns have
486 been described before in other plant species [42]. It seems plausible to assume that a similar
487 local activator/lateral repressor system comprising NEGAN and RTO also underlies the
488 pigmentation pattern in *D. purpurea*. Further investigations are required to test this hypothesis.
489
DpTFL1/CEN is an ortholog of *AmCEN* and *AtTFL1*, genes crucial for the development of an
indeterminate inflorescence architecture. Loss-of-function mutations of these genes have been
shown to lead to large terminal flowers [52, 54]. The system responsible for inflorescence
architecture seems to be highly conserved across plant species [56], suggesting a similar
system in *D. purpurea* as well. Genotyping of a *D. purpurea* population for the *DpTFL1/CEN*
locus revealed that only the homozygous mutant genotype leads to the development of a
determinate inflorescence architecture, corroborating the findings of Keeble *et al*. [51].

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All analyzed RNA-seq data sets are publicly available from the Sequence Read Archive (SRA) (Additional file 1). ONT sequencing data of *Digitalis purpurea* are available from the European Nucleotide Archive (ENA) under PRJEB63449 (Additional file 5).The assembled genome sequence of DR1 and the corresponding annotation are available via bonndata: https://doi.org/10.60507/FK2/4KUKXI [89].

Competing interests

The authors declare that they have no competing interests.

Funding

Not applicable.

Authors' contributions

BP planned the study. JH, KW, andRF performed the sequencing. JH and BP wrote the software and performed the bioinformatic analysis. JH performed genotyping experiments. JH, KW, RF, and BP interpreted the results, and wrote the manuscript. All authors have read the final version of the manuscript and approved its submission.

Acknowledgements

This work was supported by the de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI) and ELIXIR-DE (Forschungszentrum Jülich and W-de.NBI-001, W-de.NBI-004, W-de.NBI-008, W-de.NBI-010, W-de.NBI-013, W-de.NBI-014, W-de.NBI-016, W-de.NBI-022). We also thank all students participating in our 'Data Literacy in Genome Research' course and members of the research group Plant Biotechnology and Bioinformatics for discussion and support. We thank Hanna Marie Schilbert for spotting large terminal flowers on *Digitalis* and contributing to the brainstorming for experiments.

Additional files

Additional file 1: *Digitalis purpurea* RNA-seq data sets that delivered hints for the gene prediction process.

Additional file 2: *Digitalis purpurea* long read sequencing runs with corresponding accessions assigned by the European Nucleotide Archive (ENA).

Additional file 3: Documentation of all parameters and input files used for the annotation of the flavonoid biosynthesis genes with KIPEs3.

Additional file 4: Documentation of all parameters and input files used for the annotation of the MYB transcription factor gene family with the MYB_annotator.

Additional file 5: Documentation of all parameters and input files used for the annotation of the bHLH transcription factor gene family with the bHLH_annotator.

Additional file 6: *Digitalis purpurea* candidate genes associated with the anthocyanin biosynthesis.

Additional file 7: Histogram showing the read depth across the entire assembly. A coverage of 20x suggests a successful separation of haplophases, while a coverage of 40x suggests that haplophases have been merged.

Additional file 8: Purple flower (left) and white flower (right) under UV-illumination. The two upper petals were removed to open the flower and expose the inside of the remaining flower.

Additional file 9: Design of the genotyping PCRs for the *ANS* and *TFL1/CEN* locus.

Additional file 10: PCR cycle program for the *DpCEN* and *DpANS* genotyping.

Additional file 11: *D. purpurea* plants showing magenta colored petioles.

Additional file 12: Expression plot showing the activity of anthocyanin biosynthesis associated genes across different samples

References

1. Kreis W. The Foxgloves (Digitalis) Revisited. Planta Med. 2017;83:962–76. https://doi.org/10.1055/s-0043-111240.

2. Bucca A. Role of Digoxin in Heart Failure. In: Vasan RS, Sawyer DB, editors. Encyclopedia of Cardiovascular Research and Medicine. Oxford:

Elsevier; 2018. p. 323–6. https://doi.org/10.1016/B978-0-12-809657-4.99654-6.

3. Carroll E, Ravi Gopal B, Raghavan I, Mukherjee M, Wang ZQ. A cytochrome P450 CYP87A4 imparts sterol side-chain cleavage in digoxin biosynthesis. Nat Commun. 2023;14:4042. https://doi.org/10.1038/s41467-023-39719-4.

4. Winkel-Shirley B. Flavonoid Biosynthesis. A Colorful Model for Genetics, Biochemistry, Cell Biology, and Biotechnology. Plant Physiol. 2001;126:485–93. https://doi.org/10.1104/pp.126.2.485.

5. Pucker B, Selmar D. Biochemistry and Molecular Basis of Intracellular Flavonoid Transport in Plants. Plants Basel Switz. 2022;11:963. https://doi.org/10.3390/plants11070963.

6. Pucker B, Reiher F, Schilbert HM. Automatic Identification of Players in the Flavonoid Biosynthesis with Application on the Biomedicinal Plant Croton tiglium. Plants. 2020;9:1103. https://doi.org/10.3390/plants9091103.

7. Eichenberger M, Schwander T, Hüppi S, Kreuzer J, Mittl PRE, Peccati F, et al. The catalytic role of glutathione transferases in heterologous anthocyanin biosynthesis. Nat Catal. 2023;6:927–38. https://doi.org/10.1038/s41929-023-01018-y.

8. Goodman CD, Casati P, Walbot V. A Multidrug Resistance–Associated Protein Involved in Anthocyanin Transport in Zea mays. Plant Cell. 2004;16:1812–26. https://doi.org/10.1105/tpc.022574.

9. Zhao J, Dixon RA. MATE Transporters Facilitate Vacuolar Uptake of Epicatechin 3′-O-Glucoside for Proanthocyanidin Biosynthesis in Medicago truncatula and Arabidopsis. Plant Cell. 2009;21:2323–40. https://doi.org/10.1105/tpc.109.067819.

10. Mueller LA, Goodman CD, Silady RA, Walbot V. AN9, a Petunia Glutathione S-Transferase Required for Anthocyanin Sequestration, Is a Flavonoid-Binding Protein. Plant Physiol. 2000;123:1561–70. https://doi.org/10.1104/pp.123.4.1561.

11. Nozue M, Yasuda H. Occurrence of anthocyanoplasts in cell suspension cultures of sweet potato. Plant Cell Rep. 1985;4:252–5. https://doi.org/10.1007/BF00269370.

12. Grotewold E, Chamberlin M, Snook M, Siame B, Butler L, Swenson J,
et al. Engineering Secondary Metabolism in Maize Cells by Ectopic
Expression of Transcription Factors. Plant Cell. 1998;10:721–40.
https://doi.org/10.1105/tpc.10.5.721.

13. Zhang H, Wang L, Deroles S, Bennett R, Davies K. New insight into the
structures and formation of anthocyanic vacuolar inclusions in flower
petals. BMC Plant Biol. 2006;6:29. https://doi.org/10.1186/1471-2229-6-29.

14. Poustka F, Irani NG, Feller A, Lu Y, Pourcel L, Frame K, et al. A
Trafficking Pathway for Anthocyanins Overlaps with the Endoplasmic
Reticulum-to-Vacuole Protein-Sorting Route in Arabidopsis and Contributes
to the Formation of Vacuolar Inclusions. Plant Physiol. 2007;145:1323–35.
https://doi.org/10.1104/pp.107.105064.

15. Luo P, Ning G, Wang Z, Shen Y, Jin H, Li P, et al. Disequilibrium of
Flavonol Synthase and Dihydroflavonol-4-Reductase Expression
Associated Tightly to White vs. Red Color Flower Formation in Plants.
Front Plant Sci. 2016;6.

16. Johnson ET, Ryu S, Yi H, Shin B, Cheong H, Choi G. Alteration of a
single amino acid changes the substrate specificity of dihydroflavonol 4-
reductase. Plant J Cell Mol Biol. 2001;25:325–33.
https://doi.org/10.1046/j.1365-313x.2001.00962.x.

17. Choudhary N, Pucker B. Conserved amino acid residues and gene
expression patterns associated with the substrate preferences of the
competing enzymes FLS and DFR. PLOS ONE. 2024;19:e0305837.
https://doi.org/10.1371/journal.pone.0305837.

18. Weisshaar B, Jenkins GI. Phenylpropanoid biosynthesis and its
regulation. Curr Opin Plant Biol. 1998;1:251–7.
https://doi.org/10.1016/s1369-5266(98)80113-1.

19. Li S. Transcriptional control of flavonoid biosynthesis. Plant Signal
Behav. 2014;9:e27522. https://doi.org/10.4161/psb.27522.

20. Xu W, Dubos C, Lepiniec L. Transcriptional control of flavonoid
biosynthesis by MYB–bHLH–WDR complexes. Trends Plant Sci.
2015;20:176–85. https://doi.org/10.1016/j.tplants.2014.12.001.

21. Lloyd A, Brockman A, Aguirre L, Campbell A, Bean A, Cantero A, et al.
Advances in the MYB–bHLH–WD Repeat (MBW) Pigment Regulatory
Model: Addition of a WRKY Factor and Co-option of an Anthocyanin MYB
for Betalain Regulation. Plant Cell Physiol. 2017;58:1431–41.
https://doi.org/10.1093/pcp/pcx075.

22. Ramsay NA, Glover BJ. MYB-bHLH-WD40 protein complex and the
evolution of cellular diversity. Trends Plant Sci. 2005;10:63–70.
https://doi.org/10.1016/j.tplants.2004.12.011.

23. Gonzalez A, Zhao M, Leavitt JM, Lloyd AM. Regulation of the
anthocyanin biosynthetic pathway by the TTG1/bHLH/Myb transcriptional
complex in Arabidopsis seedlings. Plant J. 2008;53:814–27.
https://doi.org/10.1111/j.1365-313X.2007.03373.x.

24. Feller A, Machemer K, Braun EL, Grotewold E. Evolutionary and
comparative analysis of MYB and bHLH plant transcription factors. Plant J.
2011;66:94–116. https://doi.org/10.1111/j.1365-313X.2010.04459.x.

25. Albert NW, Butelli E, Moss SMA, Piazza P, Waite CN, Schwinn KE, et
al. Discrete bHLH transcription factors play functionally overlapping roles in
pigmentation patterning in flowers of Antirrhinum majus. New Phytol.
2021;231:849–63. https://doi.org/10.1111/nph.17142.

26. Airoldi CA, Hearn TJ, Brockington SF, Webb AAR, Glover BJ. TTG1
proteins regulate circadian activity as well as epidermal cell fate and
pigmentation. Nat Plants. 2019;5:1145–53. https://doi.org/10.1038/s41477-
019-0544-3.

27. Gonzalez A, Brown M, Hatlestad G, Akhavan N, Smith T, Hembd A, et
al. TTG2 controls the developmental regulation of seed coat tannins in
Arabidopsis by regulating vacuolar transport steps in the proanthocyanidin
pathway. Dev Biol. 2016;419:54–63.
https://doi.org/10.1016/j.ydbio.2016.03.031.

28. Borevitz JO, Xia Y, Blount J, Dixon RA, Lamb C. Activation Tagging
Identifies a Conserved MYB Regulator of Phenylpropanoid Biosynthesis.
Plant Cell. 2000;12:2383–93.

29. Tohge T, Nishiyama Y, Hirai MY, Yano M, Nakajima J, Awazuhara M,
et al. Functional genomics by integrated analysis of metabolome and
transcriptome of Arabidopsis plants over-expressing an MYB transcription

factor. Plant J Cell Mol Biol. 2005;42:218–35.
https://doi.org/10.1111/j.1365-313X.2005.02371.x.

30. Albert NW, Davies KM, Lewis DH, Zhang H, Montefiori M, Brendolise C, et al. A conserved network of transcriptional activators and repressors regulates anthocyanin pigmentation in eudicots. Plant Cell. 2014;26:962–80. https://doi.org/10.1105/tpc.113.122069.

31. Quattrocchio F, Wing J, Woude K van der, Souer E, Vetten N de, Mol J, et al. Molecular Analysis of the anthocyanin2 Gene of Petunia and Its Role in the Evolution of Flower Color. Plant Cell. 1999;11:1433–44. https://doi.org/10.1105/tpc.11.8.1433.

32. Streisfeld MA, Young WN, Sobel JM. Divergent selection drives genetic differentiation in an R2R3-MYB transcription factor that contributes to incipient speciation in Mimulus aurantiacus. PLoS Genet. 2013;9:e1003385. https://doi.org/10.1371/journal.pgen.1003385.

33. Wheeler LC, Walker JF, Ng J, Deanna R, Dunbar-Wallis A, Backes A, et al. Transcription Factors Evolve Faster Than Their Structural Gene Targets in the Flavonoid Pigment Pathway. Mol Biol Evol. 2022;39:msac044. https://doi.org/10.1093/molbev/msac044.

34. Marin-Recinos MF, Pucker B. Genetic factors explaining anthocyanin pigmentation differences. BMC Plant Biol. 2024;24:627. https://doi.org/10.1186/s12870-024-05316-w.

35. Pucker B, Walker-Hale N, Dzurlic J, Yim WC, Cushman JC, Crum A, et al. Multiple mechanisms explain loss of anthocyanins from betalain-pigmented Caryophyllales, including repeated wholesale loss of a key anthocyanidin synthesis enzyme. New Phytol. 2024;241:471–89. https://doi.org/10.1111/nph.19341.

36. Zufall RA, Rausher MD. Genetic changes associated with floral adaptation restrict future evolutionary potential. Nature. 2004;428:847–50. https://doi.org/10.1038/nature02489.

37. Del Valle JC, Alcalde-Eon C, Escribano-Bailón MT, Buide ML, Whittall JB, Narbona E. Stability of petal color polymorphism: the significance of anthocyanin accumulation in photosynthetic tissues. BMC Plant Biol. 2019;19:496. https://doi.org/10.1186/s12870-019-2082-6.

38. Thomas MM, Rudall PJ, Ellis AG, Savolainen V, Glover BJ.
Development of a complex floral trait: The pollinator-attracting petal spots
of the beetle daisy, Gorteria diffusa (Asteraceae). Am J Bot. 2009;96:2184–
96. https://doi.org/10.3732/ajb.0900079.

39. Fairnie ALM, Yeo MTS, Gatti S, Chan E, Travaglia V, Walker JF, et al.
Eco-Evo-Devo of petal pigmentation patterning. Essays Biochem.
2022;66:753–68. https://doi.org/10.1042/EBC20220051.

40. Martins TR, Berg JJ, Blinka S, Rausher MD, Baum DA. Precise spatio-
temporal regulation of the anthocyanin biosynthetic pathway leads to petal
spot formation in Clarkia gracilis (Onagraceae). New Phytol.
2013;197:958–69. https://doi.org/10.1111/nph.12062.

41. Wang Z, Li X, Chen M, Yang L, Zhang Y. Molecular and Metabolic
Insights into Anthocyanin Biosynthesis for Spot Formation on Lilium
leichtlinii var. maximowiczii Flower Petals. Int J Mol Sci. 2023;24:1844.
https://doi.org/10.3390/ijms24031844.

42. Ding B, Patterson EL, Holalu SV, Li J, Johnson GA, Stanley LE, et al.
Two MYB Proteins in a Self-Organizing Activator-Inhibitor System Produce
Spotted Pigmentation Patterns. Curr Biol CB. 2020;30:802-814.e8.
https://doi.org/10.1016/j.cub.2019.12.067.

43. Zhang Y, Cheng Y, Ya H, Xu S, Han J. Transcriptome sequencing of
purple petal spot region in tree peony reveals differentially expressed
anthocyanin structural genes. Front Plant Sci. 2015;6.

44. Yuan Y-W, Rebocho AB, Sagawa JM, Stanley LE, Bradshaw HD.
Competition between anthocyanin and flavonol biosynthesis produces
spatial pattern variation of floral pigments between Mimulus species. Proc
Natl Acad Sci. 2016;113:2448–53.
https://doi.org/10.1073/pnas.1515294113.

45. Lin R-C, Rausher MD. R2R3-MYB genes control petal pigmentation
patterning in Clarkia gracilis ssp. sonomensis (Onagraceae). New Phytol.
2021;229:1147–62. https://doi.org/10.1111/nph.16908.

46. Sasaki K, Takahashi T. A flavonoid from Brassica rapa flower as the
UV-absorbing nectar guide. Phytochemistry. 2002;61:339–43.
https://doi.org/10.1016/s0031-9422(02)00237-6.

47. Medel R, Botto-Mahan C, Kalin-Arroyo M. Pollinator-Mediated Selection
on the Nectar Guide Phenotype in the Andean Monkey Flower, Mimulus
luteus. Ecology. 2003;84:1721–32.

48. Lunau K, Fieselmann G, Heuschen B, van de Loo A. Visual targeting of
components of floral colour patterns in flower-naïve bumblebees (Bombus
terrestris; Apidae). Naturwissenschaften. 2006;93:325–8.
https://doi.org/10.1007/s00114-006-0105-2.

49. Davies KM, Albert NW, Schwinn KE, Davies KM, Albert NW, Schwinn
KE. From landing lights to mimicry: the molecular regulation of flower
colouration and mechanisms for pigmentation patterning. Funct Plant Biol.
2012;39:619–38. https://doi.org/10.1071/FP12195.

50. Rudall PJ. Weird and wonderful foxgloves. 2016.
https://www.kew.org/read-and-watch/weird-and-wonderful-foxgloves.
Accessed 2 May 2024.

51. Keeble F, Pellew MC, Jones WN. The Inheritance Op Peloria and
Flowercolour in Foxgloves (digitalis Purpurea). New Phytol. 1910;9:68–77.
https://doi.org/10.1111/j.1469-8137.1910.tb05554.x.

52. Bradley D, Carpenter R, Copsey L, Vincent C, Rothstein S, Coen E.
Control of inflorescence architecture in Antirrhinum. Nature. 1996;379:791–
7. https://doi.org/10.1038/379791a0.

53. Shannon S, Meeks-Wagner DR. A Mutation in the Arabidopsis TFL1
Gene Affects Inflorescence Meristem Development. Plant Cell.
1991;3:877–92. https://doi.org/10.2307/3869152.

54. Bradley D, Ratcliffe O, Vincent C, Carpenter R, Coen E. Inflorescence
Commitment and Architecture in Arabidopsis. Science. 1997;275:80–3.
https://doi.org/10.1126/science.275.5296.80.

55. Alvarez J, Guli CL, Yu X-H, Smyth DR. terminal flower: a gene affecting
inflorescence development in Arabidopsis thaliana. Plant J. 1992;2:103–16.
https://doi.org/10.1111/j.1365-313X.1992.00103.x.

56. Pnueli L, Gutfinger T, Hareven D, Ben-Naim O, Ron N, Adir N, et al.
Tomato SP-Interacting Proteins Define a Conserved Signaling System That
Regulates Shoot Architecture and Flowering. Plant Cell. 2001;13:2687–
702. https://doi.org/10.1105/tpc.010293.

57. Siadjeu C, Pucker B, Viehöver P, Albach DC, Weisshaar B. High
Contiguity de novo Genome Sequence Assembly of Trifoliate Yam
(Dioscorea dumetorum) Using Long Read Sequencing. Genes.
2020;11:274. https://doi.org/10.3390/genes11030274.

58. Wolff K, Friedhoff R, Schwarzer F, Pucker B. Data literacy in genome
research. J Integr Bioinforma. 2023;20. https://doi.org/10.1515/jib-2023-
0033.

59. Li H. Minimap2: pairwise alignment for nucleotide sequences.
Bioinformatics. 2018;34:3094–100.
https://doi.org/10.1093/bioinformatics/bty191.

60. GrandOmics. NextDenovo. 2023.

61. Shafin K, Pesout T, Lorig-Roach R, Haukness M, Olsen HE, Bosworth
C, et al. Nanopore sequencing and the Shasta toolkit enable efficient de
novo assembly of eleven human genomes. Nat Biotechnol. 2020;38:1044–
53. https://doi.org/10.1038/s41587-020-0503-6.

62. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-
prone reads using repeat graphs. Nat Biotechnol. 2019;37:540–6.
https://doi.org/10.1038/s41587-019-0072-8.

63. GrandOmics. NextPolish. 2023.

64. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The
Sequence Alignment/Map format and SAMtools. Bioinformatics.
2009;25:2078–9. https://doi.org/10.1093/bioinformatics/btp352.

65. Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B.
AUGUSTUS: ab initio prediction of alternative transcripts. Nucleic Acids
Res. 2006;34 suppl_2:W435–9. https://doi.org/10.1093/nar/gkl200.

66. Keller O, Kollmar M, Stanke M, Waack S. A novel hybrid gene
prediction method employing protein multiple sequence alignments.
Bioinforma Oxf Engl. 2011;27:757–63.
https://doi.org/10.1093/bioinformatics/btr010.

67. NCBI. sra-tools. 2020.

68. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al.
STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29:15–21.
https://doi.org/10.1093/bioinformatics/bts635.

69. Dobin A, Gingeras TR. Mapping RNA-seq Reads with STAR. Curr
Protoc Bioinforma. 2015;51:11.14.1-11.14.19.
https://doi.org/10.1002/0471250953.bi1114s51.

70. Haak M, Vinke S, Keller W, Droste J, Rückert C, Kalinowski J, et al.
High Quality de Novo Transcriptome Assembly of Croton tiglium. Front Mol
Biosci. 2018;5. https://doi.org/10.3389/fmolb.2018.00062.

71. Keilwagen J, Wenk M, Erickson JL, Schattat MH, Grau J, Hartung F.
Using intron position conservation for homology-based gene prediction.
Nucleic Acids Res. 2016;44:e89. https://doi.org/10.1093/nar/gkw092.

72. Keilwagen J, Hartung F, Grau J. GeMoMa: Homology-Based Gene
Prediction Utilizing Intron Position Conservation and RNA-seq Data.
Methods Mol Biol Clifton NJ. 2019;1962:161–77.
https://doi.org/10.1007/978-1-4939-9173-0_9.

73. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM.
BUSCO: assessing genome assembly and annotation completeness with
single-copy orthologs. Bioinformatics. 2015;31:3210–2.
https://doi.org/10.1093/bioinformatics/btv351.

74. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO
Update: Novel and Streamlined Workflows along with Broader and Deeper
Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral
Genomes. Mol Biol Evol. 2021;38:4647–54.
https://doi.org/10.1093/molbev/msab199.

75. Pucker B, Holtgräwe D, Weisshaar B. Consideration of non-canonical
splice sites improves gene prediction on the Arabidopsis thaliana
Niederzenz-1 genome sequence. BMC Res Notes. 2017;10:667.
https://doi.org/10.1186/s13104-017-2985-y.

76. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local
alignment search tool. J Mol Biol. 1990;215:403–10.
https://doi.org/10.1016/S0022-2836(05)80360-2.

77. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et
al. Gapped BLAST and PSI-BLAST: a new generation of protein database
search programs. Nucleic Acids Res. 1997;25:3389–402.

78. Pucker B, Iorizzo M. Apiaceae FNS I originated from F3H through
tandem gene duplication. PLOS ONE. 2023;18:e0280155.
https://doi.org/10.1371/journal.pone.0280155.

79. Rempel A, Choudhary N, Pucker B. KIPEs3: Automatic annotation of
biosynthesis pathways. PLOS ONE. 2023;18:e0294342.
https://doi.org/10.1371/journal.pone.0294342.

80. Pucker B. Automatic identification and annotation of MYB gene family
members in plants. BMC Genomics. 2022;23:220.
https://doi.org/10.1186/s12864-022-08452-5.

81. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability. Mol Biol Evol.
2013;30:772–80. https://doi.org/10.1093/molbev/mst010.

82. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-
Likelihood Trees for Large Alignments. PLOS ONE. 2010;5:e9490.
https://doi.org/10.1371/journal.pone.0009490.

83. Thoben C, Pucker B. Automatic annotation of the bHLH gene family in
plants. BMC Genomics. 2023;24:780. https://doi.org/10.1186/s12864-023-
09877-2.

84. Edgar RC. Muscle5: High-accuracy alignment ensembles enable
unbiased assessments of sequence homology and phylogeny. Nat
Commun. 2022;13:6968. https://doi.org/10.1038/s41467-022-34630-w.

85. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic
RNA-seq quantification. Nat Biotechnol. 2016;34:525–7.
https://doi.org/10.1038/nbt.3519.

86. Robinson JT, Thorvaldsdottir H, Turner D, Mesirov JP. igv.js: an
embeddable JavaScript implementation of the Integrative Genomics Viewer
(IGV). Bioinformatics. 2023;39:btac830.
https://doi.org/10.1093/bioinformatics/btac830.

87. Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B. An Arabidopsis thaliana T-DNA mutagenized population (GABI-Kat) for flanking sequence tag-based reverse genetics. Plant Mol Biol. 2003;53:247–59. https://doi.org/10.1023/B:PLAN.0000009297.37235.4a.

88. Pucker B, Friedhoff R, Horz J, Wolff K. Genome sequence and annotation of Digitalis purpurea. 2023. https://doi.org/10.24355/dbbs.084-202311130950-0.

89. Horz J, Wolff K, Friedhoff R, Pucker B. Digitalis purpurea genome sequence and annotation V2. 2025. https://doi.org/10.60507/FK2/4KUKXI.

90. Forkmann G, Heller W, Grisebach H. Anthocyanin Biosynthesis in Flowers of Matthiola incana Flavanone 3-and Flavonoid 3′-Hydroxylases. Z Für Naturforschung C. 1980;35:691–5. https://doi.org/10.1515/znc-1980-9-1004.

91. Ferrer J-L, Jez JM, Bowman ME, Dixon RA, Noel JP. Structure of chalcone synthase and the molecular basis of plant polyketide biosynthesis. Nat Struct Biol. 1999;6:775–84. https://doi.org/10.1038/11553.

92. Jez JM, Bowman ME, Dixon RA, Noel JP. Structure and mechanism of the evolutionarily unique plant enzyme chalcone isomerase. Nat Struct Biol. 2000;7:786–91. https://doi.org/10.1038/79025.

93. Brugliera F, Barri-Rewell G, Holton TA, Mason JG. Isolation and characterization of a flavonoid 3′-hydroxylase cDNA clone corresponding to the Ht1 locus of Petunia hybrida. Plant J. 2002;19:441–51. https://doi.org/10.1046/j.1365-313X.1999.00539.x.

94. Saito K, Yonekura-Sakakibara K, Nakabayashi R, Higashi Y, Yamazaki M, Tohge T, et al. The flavonoid biosynthetic pathway in Arabidopsis: Structural and genetic diversity. Plant Physiol Biochem. 2013;72:21–34. https://doi.org/10.1016/j.plaphy.2013.02.001.

95. Abrahams S, Lee E, Walker AR, Tanner GJ, Larkin PJ, Ashton AR. The Arabidopsis TDS4 gene encodes leucoanthocyanidin dioxygenase (LDOX) and is essential for proanthocyanidin synthesis and vacuole development. Plant J. 2003;35:624–36. https://doi.org/10.1046/j.1365-313X.2003.01834.x.

96. Kim S, Binzel ML, Yoo KS, Park S, Pike LM. Pink (P), a new locus
responsible for a pink trait in onions (Allium cepa) resulting from natural
mutations of anthocyanidin synthase. Mol Genet Genomics. 2004;272:18–
27. https://doi.org/10.1007/s00438-004-1041-5.

97. Jiang T, Zhang M, Wen C, Xie X, Tian W, Wen S, et al. Integrated
metabolomic and transcriptomic analysis of the anthocyanin regulatory
networks in Salvia miltiorrhiza Bge. flowers. BMC Plant Biol. 2020;20:349.
https://doi.org/10.1186/s12870-020-02553-7.

98. McClintock B. The Origin and Behavior of Mutable Loci in Maize. Proc
Natl Acad Sci U S A. 1950;36:344–55.

99. McClintock B. INDUCTION OF INSTABILITY AT SELECTED LOCI IN
MAIZE. Genetics. 1953;38:579–99.
https://doi.org/10.1093/genetics/38.6.579.

100. Owen CR, Bradshaw HD. Induced mutations affecting pollinator
choice in Mimulus lewisii (Phrymaceae). Arthropod-Plant Interact.
2011;5:235–44. https://doi.org/10.1007/s11829-011-9133-8.

101. Abid MA, Wei Y, Meng Z, Wang Y, Ye Y, Wang Y, et al. Increasing
floral visitation and hybrid seed production mediated by beauty mark in
Gossypium hirsutum. Plant Biotechnol J. 2022;20:1274–84.
https://doi.org/10.1111/pbi.13805.

102. Richter R, Dietz A, Foster J, Spaethe J, Stöckl A. Flower patterns
improve foraging efficiency in bumblebees by guiding approach flight and
landing. Funct Ecol. 2023;37:763–77. https://doi.org/10.1111/1365-
2435.14262.

103. Johnson SD, Midgley JJ. Fly pollination of Gorteria diffusa
(Asteraceae), and a possible mimetic function for dark spots on the
capitulum. Am J Bot. 1997;84:429–36. https://doi.org/10.2307/2446018.

104. de Jager ML, Willis-Jones E, Critchley S, Glover BJ. The impact of
floral spot and ring markings on pollinator foraging dynamics. Evol Ecol.
2017;31:193–204. https://doi.org/10.1007/s10682-016-9852-5.