Title: Predicting fitness related traits using gene expression and machine
learning

Authors: Georgia A. Henry[1]* & John R. Stinchcombe[1,2,3]

[1] Department of Ecology & Evolutionary Biology, University of Toronto, Toronto, Ontario,
Canada

[2] Koffler Scientific Reserve at Joker's Hill, University of Toronto, King, Ontario, Canada

[3] Swedish Collegium for Advanced Study, Uppsala University, Uppsala, Sweden
9

Corresponding author: georgiahenry5@gmail.com

Author contributions: G.A.H. and J.R.S contributed to the design, analysis and writing of the
manuscript. G.A.H. performed the data collection and implemented analyses.

Conflict of Interest Statement: The authors declare no conflict of interest.

Data Availability: The data underlying this article available through the NCBI Sequence Read
Archive https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1030903/.

Abstract

Evolution by natural selection occurs at its most basic through the change in frequencies of alleles; connecting those genomic targets to phenotypic selection is an important goal for evolutionary biology in the genomics era. The relative abundance of gene products expressed in a tissue can be considered a phenotype intermediate to the genes and genomic regulatory elements themselves, and more traditionally measured macroscopic phenotypic traits such as flowering time, size, or growth. The high-dimensionality, low sample size nature of transcriptomic sequence data is a double-edged sword, however, as it provides abundant information but makes traditional statistics difficult. Machine learning has many features which handle high-dimensional data well and is thus useful in genetic sequence applications. Here we examined the association of fitness-components with gene expression data in *Ipomoea hederacea* (Ivyleaf Morning Glory) grown under field conditions. We combine the results of two different machine learning approaches and find evidence that expression of photosynthesis-related genes is likely under selection. We also find that genes related to stress and light response were overall important in predicting fitness. With this study we demonstrate the utility of machine learning models for smaller samples, and their potential application for understanding natural selection.

Introduction

Natural selection is a ubiquitous evolutionary force which acts on the phenotypes of individuals in a population, resulting in genetic changes if heritable variation exists for the traits under selection. Understanding the shape and strength of natural selection typically requires observers to choose specific traits to measure which may or may not be the subject of selection

(Lande and Arnold 1983). The type of traits studied has been found to influence the strength

and temporal constancy of selection, with fecundity traits experiencing more consistent and

strong selection than survivorship (Hoekstra et al. 2001; Kingsolver et al. 2001). Trait

correlations and pleiotropy can additionally make detecting selection statistically difficult due to

their potential to reduce power (Hersch and Phillips 2004). With high-throughput phenotyping,

such as metabolomics or gene expression profiling, investigators often obtain a vast number of

highly correlated traits, making traditional phenotypic selection analysis difficult or impossible.

Machine learning modelling and regularization techniques, which can alleviate issues of

multicollinearity and overfitting due to an overabundance of explanatory variables, now make it

possible for investigators to avoid having to decide which traits may be important to selection,

and instead observe them from the data. Here, we use machine learning modelling to estimate

natural selection on gene expression, using estimates of fitness and gene expression obtained

from a field experiment with the Ivyleaf morning glory (*Ipomoea hederacea*).

Gene expression can be viewed as an intermediate between the genome and traditionally

observable traits, and reflects both the underlying genetic makeup and the environment (Liao

and Weng 2015; Josephs 2021). Quantifying gene expression through RNA sequencing of

tissue yields a less biased sampling of phenotypic traits compared to an observer's opinions on

what might be an important trait (Rockman and Kruglyak 2006; Josephs 2021), although

investigators must still decide on growth or rearing environments, as well as the tissues and

developmental stages to be sampled. RNA-sequence data is also highly multivariate, in that it

captures a sample of all the genes being expressed in a tissue at the moment of sampling,

yielding a broad representation of many phenotypes simultaneously. Although gene expression

was initially expected to largely be under stabilizing selection (e.g, from mutation accumulation

studies, Rifkin et al. 2005; also see Gilad et al. 2006), recent results suggest that there may

directional selection on genes' expression, especially when individuals are farther from the

fitness optima. For example, Groen et al. (2020) measured selection on gene expression under

drought and standard field conditions in domesticated rice (*Oryza sativa*). Using univariate

approaches, they were able to estimate selection differentials for individual genes' expression,

finding generally weak selection. Selection was stronger under drought conditions, with earlier

flowering time, modulated apparently by a single gene, allowing plants to "escape" the effects of

drought (Groen et al. 2020). They were additionally able to identify significant selection on

genes related to photosynthesis and growth, using a multivariate approach where they

regressed fitness on principal component (PC) scores that summarized expression of many

genes at once. Several investigators have constructed gene co-expression modules (Palakurty

et al. 2018, Josephs et al. 2020, Brown and Kelly 2021), and then related multivariate estimates

of expression of the entire module to phenotypic traits that are either indicative of overall plant

performance or size, or are traits likely to be under natural selection such as components of life

history. These experiments have found the expression of multiple genes in a module to be

significantly associated with traits likely to be under natural selection in populations, thus

implicating the expression levels of genes in the modules as potential targets of natural

selection. Collectively, these studies illustrate how measuring selection on gene expression may

facilitate identifying genes or pathways which are important for fitness (Price et al. 2022).

The dominant method of characterizing contemporary phenotypic selection is the Lande-Arnold

(1983) regression approach, and variants of it (Kingsolver and Schemske 1991, Rausher 1992).

In this approach, estimates of relative fitness are regressed on multiple, potentially correlated

phenotypes. The resulting coefficients, termed selection gradients, measure the direct effect of

the phenotype on relative fitness, accounting for the effects of other potentially correlated traits

included in the model (Lande and Arnold 1983). While this approach has yielded thousands of

estimates of natural selection (see Kingsolver et al. 2012), it faces fundamental limitations with

gene expression data. Transcriptomic sequencing data is typically plagued by the "small n, large

p" problem, where the number of "features" (estimates of expression) far exceeds the sample

size. As a consequence, the multiple regression approach to measuring selection is untenable,

as there are vastly more phenotypes than observations. Past studies seeking to measure

contemporary selection on gene expression have thus used data reduction techniques such as

PCA (Groen et al. 2020) or the construction of gene coexpression modules, whose overall

expression is itself summarized with PCA (eigen-expression; Josephs et al. 2020, Brown and

Kelly 2021). While these approaches allow estimates of selection on PC scores – of either the

whole transcriptome or a co-expression module – they do not allow direct or indirect selection

on an individual gene's expression to be distinguished. Consequently, one of the dominant

benefits of the Lande-Arnold (1983) approach is lost. Perhaps because of the "small n, large p"

problem or the difficulty of interpreting selection on PC scores (Chong et al. 2018), we have

dramatically fewer estimates of phenotypic selection on gene expression, especially compared

to morphology, behavior, and life history.

Machine learning tools are often well-suited to handling the "curse of dimensionality" due to

intrinsic and extrinsic regularization and are thus useful in the analysis of sequence data

(Schrider and Kern 2018; James et al. 2021), and may be useful for measuring selection on

multivariate measures of gene expression. Machine learning typically includes some type of loss

function which serves to balance under- and over-fitting of the model, and as such observations

with extreme values are not overly influential. The data is split into a training set (which is used

to fit the model) and a testing set of data, which can then be used to evaluate the model

performance, which can buffer against some overfitting by reducing the amount of statistical

noise being fit to the model. Additionally, feature selection either through manual evaluation and

pruning (as in Principal Component Regression), or through intrinsic model behaviour (as in

LASSO regression or Decision Trees), can alleviate problems due to too many explanatory

variables. Regularized estimates are biased (i.e., no longer unbiased, least squares estimates)

but often lead to more accurate overall prediction (see Morrissey 2014, Sztepanacz and Houle

2024). Generally, machine learning tools focus on optimizing prediction rather than estimation of

parameter values within a model (Schrider and Kern 2018; James et al. 2021; Greener et al.

2022) and thus are valuable for understanding complex nonlinear relationships between

variables. While there has been some exploration of using regularized regression to estimate

phenotypic selection (Morrissey 2014, Sztepanacz and Houle 2024), to date applications of

machine learning to studies of phenotypic selection, especially for gene expression, have been

limited.

We used gene expression data from 96 *Ipomoea hederacea* individuals grown under natural

field conditions to detect important fitness-related genes using various machine learning

methods. We examine the union of unsupervised and supervised approaches. Unsupervised

machine learning models are those in which the observations are not "labeled" with a response

variable, and supervised models in contrast use observations which are "labeled" with a

response variable. We first use principal component analysis on gene expression as an

unsupervised dimensionality reduction method followed by linear regression of relative fitness

on principal component scores. We then trained a supervised classification model to predict

whether an individual set seed based on our gene expression data. In combining disparate

methods, we sought to find consistently important candidate genes or biological processes

which are likely contributing to differences in fitness among individuals in the field. We

investigated the predictive ability of these machine learning approaches and extracted the

genes whose expression contributed most to the model fit (i.e. those associated most strongly

with fitness differences). We then used Gene Ontology (GO) analyses to deepen our

understanding of the types of genes which were significantly enriched in each model and the

biological processes commonly found to be important between the models.

Results

Selection differentials for gene expression

We initially estimated selection differentials for gene expression, using two analyses. First, for

each gene, we estimated the difference in expression between individuals that set seed (the

after-selection cohort, z^*) and the total population, regardless of seed set (the before selection

cohort, z). We found a wide distribution of selection differentials centered on zero (Fig. 1A),

suggesting that total selection to increase or decrease expression appears equally common,

and that some genes are under strong selection to increase or decrease expression. While

these analyses match the structure of the classification algorithms, we used for machine

learning (*see below*), they ignore the differential fecundity of individuals that set seed. We

secondly estimated selection differentials as the covariance between expression and relative

fitness, which includes differential fecundity (Fig. 1B), finding much the same pattern as in the

binary fitness selection differentials, although the distribution is slightly broader. The two

estimates of selection differentials were similar (Fig. 1C), indicating that differential fecundity is a

key component of selection on expression. Because selection differentials measure total

selection on a phenotype and cannot distinguish direct selection or indirect selection via

correlated traits, we next turned to selection gradients for gene expression.

162
Figure 1: Selection differentials for gene expression calculated using the binary fitness (set seed
or did not set seed) and using the relative fitness in number of set seed. A) Histogram of the
binary fitness selection differentials for standardized gene expression counts of 2753 genes. B)
Histogram of the selection differentials as the covariance between relative fitness and
standardized gene expression counts. C) The relationship between the selection differentials
calculated using the two methods for each gene. The binary fitness selection differentials have a
slightly narrower distribution but are highly correlated with the continuous fitness selection
differential (Pearson's correlation coefficient = 0.85, p < 0.001).

Unsupervised modeling

The Principal Component Analysis (PCA) transformed the 2753 standardized gene expression

counts from 96 individuals into 96 linear combinations of the standardized gene expression

measures. The first 10 principal components (PCs) describe 50.62% of the variation in gene

expression, and 68 PCs are required to explain >90% of the variation (recall that PCs are in

descending order of the variance in gene expression they explain). We then regressed relative

fitness on each PC individually to get the univariate regression p-value which we used to

evaluate the model fit with varying numbers of PCs as predictor variables. We assessed model

fits across different p-value cut-offs.Our best model included a total of 61 PCs with individual p-

values <0.60, which minimized the RMSE (root mean-squared error) and MAE (mean absolute

error), while maximizing the R-squared value (Figure S1). We note that these 61 PCs are *not*

the first 61, and that the overall model which minimizes the RMSE and MAE contains PC axes

that are not, on their own, significant predictors of fitness (Figure S2). The *caret* package, which

we used to perform our repeat cross-validation regressions, uses the correlation between the

predicted and observed response values as the R-squared value (Kuhn 2008). The model R-

squared was 0.549, with an RMSE of 1.719 and MAE of 1.381, and included 61 PCs (Table 1).

We then transformed the regression coefficients for the PCs used in the final model back to the

standard-deviation scaled gene expression counts using the rotations from the PCA (Chong et

al. 2018) to evaluate the strength and distribution of gene expression selection gradients (β).

We also calculated the standard error of the selection gradient for each gene's expression

(following Chong et al. 2018), and then evaluated whether the absolute value of the selection

gradient was greater than 1.96 × its standard error (SE), following the standard formula for a

95% confidence interval (Sokal and Rohlf 1995). Many (~59%) of the absolute selection

gradients were greater than 1.96 × SE, suggesting that most of our estimates are significantly

greater than sampling error, within the multivariate space spanned by these 61 PC axes. The

distribution of β's was centered on zero, with an approximately normal distribution, which

suggests that there is directional selection for both increased and decreased gene expression

(Fig. 2). Overall, the magnitude of β's was very small: an order of magnitude smaller than the

significant β's estimated for macroscopic traits in the same experiment, as reported by Henry

and Stinchcombe (2023; Fig. 2B). The only β for macroscopic, traditional traits that was within

the range of the gene expression selection gradients was for anther-stigma distance, which was

not significant. Given that these selection gradients are on individual genes and not the

combined outcome of many genes, as is the case for the quantitative traits studied by Henry

and Stinchcombe (2023), we expected the directional selection gradients on each gene's

expression individually to be small. In addition, if any of the expressed genes are themselves

causally associated with phenotypic traits under selection (i.e., expression → trait → fitness), we

expect a small estimate for expression's net effect on fitness (expression → fitness), due to it

being the product of two small numbers.

Figure 2: Distribution of selection gradients for gene expression of the 2753 genes. A) Selection
gradients shaded by whether they were greater than 1.96 x their standard error. B) The same
distribution of selection gradients plotted alongside the selection gradients of macroscopic
phenotypes. C) Selection gradients plotted as their absolute values, yellow representing the
negatively signed selection gradients, and blue representing the positively signed selection
gradients. D) Absolute selection gradients that are greater than 1.96 x their standard error.

223 **Table 1:** Metrics of model fit for both supervised and unsupervised methods. Gradient boosting
224 metrics are calculated from the testing data only. PCA regression metrics are from repeat five-
225 fold cross-validation, and R-squared is calculated as the correlation between the observed and
226 predicted data.

Model	Balanced Accuracy	Sensitivity	RMSE	R-squared
Gradient Boosting	0.737	0.545	-	-
PCA regression	-	-	1.719	0.549

Supervised modeling

We tested a variety of hyperparameters through a grid search with five-fold cross validation for

our supervised model. Our supervised machine learning model was a gradient boosting

classifier, which fits weakly predictive decision trees, progressively adjusting the parameter

weights to improve classification performance of misclassified observations. We tuned the

hyperparameters of the model using a grid search with five-fold cross validation of the data. The

best fitting model used a binomial deviance loss function, and the decision split quality was

determined by Friedman adjusted mean-squared error (Friedman 2001). The models were

regularized by using stochastic gradient boosting, which randomly subsamples the test data

each iteration to reduce overfitting in the final ensemble model. We used 1000 estimators, a

constant learning rate of 1.0, and subsampled 40% of the data each iteration. The final model

had an out-of-sample balanced accuracy score of 0.737, with sensitivity to the successful seed

set class of 0.545 (Table 1, Figure S3). We extracted the 278 most important features based on

their Gini importance from the model, with a cutoff of > 0.000677798, as the next 197 values

were tied.

GO Analysis

Once we obtained a subset of most important genes for each model, we performed BLAST

searches on gene sets using an *Arabidopsis thaliana* genome (Lamesch et al. 2012) as a

reference, and then mapped and annotated GO terms. We generated combined GO maps for

our gene subsets, filtering out intermediates, for each model. We found 11 GO terms common

across our two models, in addition to the higher level "biological process" term (Table 2). The

shared terms are involved in developmental and metabolism, and various stimulus responses.

These common terms highlight the importance in the response to both plant enemies (defense

response, defense response to other organisms, and response to biotic stimulus), and the

abiotic environment (response to light stimulus, and development terms). More broadly, non-

identical but related GO terms common across the models were associated with defense and

stress response (GO:0006950, GO:0009737, GO:0051707, GO:0009620), metabolism

(GO:0051171, GO:0060255, GO:0016071, GO:0080090, GO:0034641,

GO:0006518,GO:0008152, GO:0032787, GO:0044238, GO:0016070, GO:0071704,

GO:0006629) and reproductive structure development (GO:0048608, GO:0099402,

GO:0009653, GO:0022414, GO:0048731, GO:0009791) (see Supplemental Material, Table S2

– Table S3).

266 **Table 2:** Shared GO terms from GO mapping of gene subsets. Gene subsets were composed
267 of the most important genes in each mode, which were included in a BLAST search and
268 annotated with GO terms. Response terms dominate, along with developmental and metabolic
269 terms. The GO term for light response is also shared.

GO ID	GO Name
GO:0009791	post-embryonic development
GO:0009987	cellular process
GO:0050896	response to stimulus
GO:0044238	primary metabolic process
GO:0010468	regulation of gene expression
GO:0098542	defense response to other organism
GO:0006952	defense response
GO:0051716	cellular response to stimulus
GO:0009416	response to light stimulus
GO:0009607	response to biotic stimulus
GO:0009725	response to hormone

We then used *goseq* to adjust for count bias in our gene set to determine which GO terms were

significantly enriched in each model set, using $p < 0.10$ as our threshold. Two GO terms were

shared between our models, "seed maturation" (GO:0010431) and "vegetative to reproductive

phase transition of meristem" (GO:0010228) The two sets also included similar terms related to

photosynthesis, seed maturation and dormancy, stress, and heavy metals (see Supplemental

Material, Table S4 – Table S5).

Shared Important Genes

There were 29 shared genes in the sets of most important genes for each model. For the 29

genes that are among the top ranked in the PC regression and identified by the gradient

boosting classifier, we observed that the values of selection differentials (median = 0.17, range

= [-0.44, 0.79]) and selection gradients (median = 0.01, range = -0.04, 0.03]) for these genes

were bimodal, with fewer instances of values close to zero relative to the distribution of all of the

genes that we studied (Fig 1, Fig 2, Fig. S5, Fig. S6). These data suggest that the gradient

boosting classifier is identifying genes for further study and exploration that show the strongest

patterns of phenotypic selection. Of particular note was that in this top 29 genes identified by

each method, GO terms related to heavy metals were apparent in each. Consequently, we next

turned to the heavy metal concentrations of the soils at our field site.

Heavy Metal Analysis of Soils

None of the metals in the screen (Table S8) were above government agricultural guidelines

(Ontario Ministry of the Environment 2011) and were largely below the average for agricultural

fields in the area (Frank et al. 1976). While our sample of plant lines was from several states,

heavy metal concentrations in the field were also below the state-wide average for North

Carolina which is in the heart of *I. hederacea's* native range (Smith et al. 2008). Collectively,

these data suggest that neither an induced response to contaminated soil at our field site, nor

long term adaptation to heavy metal soils in the native range, are contributors to the importance

of the expression of LOC109174332 in predicting relative fitness. While heavy-metal associated

plant proteins (HIPP) have clearly been found to be associated with heavy-metal tolerance and

detoxification (hence their name), they have also demonstrated responses to drought and cold

tolerance (Barth et al. 2009; de Abreu-Neto et al. 2013; Zschiesche et al. 2015; Zheng et al.

2023), which interact with plant growth and development (Zschiesche et al. 2015; Guo et al.

2021).

Discussion

Connecting gene expression to fitness components can elucidate which genes are the targets of selection, but the high dimensionality, low sample size nature of gene expression data makes this goal computationally and statistically challenging. Machine learning (ML) techniques are well-suited to this sort of complex task, as feature selection, regularization to prevent overfitting (and thus poor prediction out of sample), and nonlinear functions are key aspects of many ML algorithms. In this study we successfully used gene expression in leaf tissue to predict the reproductive success of 96 *Ipomoea hederacea* individuals using statistical learning approaches. We find that after appropriate preprocessing of the data, a reasonable level of model accuracy can be attained using both supervised classification algorithms and semi-supervised regression, with our classification model predicting correctly over 70% of the test cases. Similarly, unsupervised dimensionality reduction followed by regression, had a correlation of 0.55 between the predicted and observed fitness values. Below, we discuss our results in light of the typical strength of selection gradients for gene expression, the insight of multiple analysis approaches, and GO categorization for understanding the relationship between expression and fitness components.

Distribution and strength of selection on gene expression

We found appreciable evidence for phenotypic selection on gene expression, using variants of the same methods that have provided such a rich picture of selection on life history, phenology, morphology, and sexually selected characters (Hoekstra et al. 2001; Kingsolver et al. 2001; Kingsolver et al. 2012). There are, however, several caveats associated with our analyses. First, there is uncertainty in our estimates of selection. While we estimated standard errors, and 95%

confidence intervals, for the selection gradients, the standard errors are only relevant in the

portion of the data described by the PC axes included in our analyses; the formula for a

confidence interval carries with it normality assumptions. Second, as with all selection analyses,

there is the potential of missing traits not included in the model: it may be that expression of

other genes (at other time points, or in other tissues) or other macroscopic traits are the true

targets of selection, and that our selection estimates would change with their inclusion in our

models. A knottier challenge is the possibility that both gene expression and relative fitness

were responding to some unknown variable, and that the associations we detected were

environmentally induced covariances (Rausher 1992; Stinchcombe et al. 2002). While we

attempted to minimize this possibility using a spatially replicated, randomized design– and we

detected little evidence of environmental covariances in estimates of selection on macroscopic

traits (Henry and Stinchcombe 2023) – we cannot formally exclude the possibility. Finally, our

estimate of fitness – lifetime seed production produced by an annual, largely selfing plant– omits

other fitness components, such as germination success and over-winter seed survival in the

seed bank. Many of these challenges are shared, by and large, by most estimates of phenotypic

selection that have been gathered by evolutionary biologists. The problem of missing traits is

inherent to the approach; many estimates of selection use fitness components or aspects of

performance (e.g., size) rather than true fitness. Finally, most of the thousands of estimates of

phenotypic selection in the literature (Hoekstra et al. 2001; Kingsolver et al. 2001; Kingsolver et

al. 2011; Kingsolver et al. 20012; Siepielski et al. 2009; Siepelski et al. 2013) are also likely

susceptible to environmental covariances. Replicated sampling of genotypes is one of the best

approaches for addressing the environmental covariance problem, but in the context of gene

expression, this is likely to be an expensive proposition.

With these caveats in mind, our results indicate that phenotypic selection on gene expression

can be detected using moderate sample sizes, even in a non-model species. We used principal

component analysis (an unsupervised machine learning approach) along with a repeat 5-fold

cross validation approach to linear regression (a supervised machine learning approach) to

estimate selection gradients. By back-transforming the selection gradients for the principal

component scores to return the selection gradients for each individual gene's expression, we

can improve interpretability and further investigate the most relevant genes. We found that

directional selection gradients for gene expression were overall symmetrically distributed around

zero, and thus selection for both increasing and decreasing gene expression were

approximately equal. The strength of selection was an order of magnitude lower than that of

other traits measured in this field experiment (Henry and Stinchcombe 2023). Our finding was

not unexpected, given low estimates of selection differentials in univariate analyses by Groen

(2020) and by intuition — combined gene expression over time leads to higher level quantitative

traits, and thus the individual expression components experience a fraction of the selection

pressure compared to the total, end-point phenotype. We believe these results represent the

first selection gradients for gene expression in the Lande-Arnold framework and illustrate the

merits of PC regression for estimating selection on high-dimensional traits.

Leveraging information from machine learning approaches

Both of our models had moderate predictive accuracy, which demonstrates the general

applicability of machine learning in understanding selection on gene expression. The ensemble

method of classification was able to predict with a moderately high degree of sensitivity given

our limited data and had a balanced accuracy score of 0.737. Similarly, although not directly

comparable, the regression model was able to explain a moderate amount of variation in

relative fitness and had an r-squared value of 0.549. The noise inherent in tissue-level gene

expression data paired with its high dimensionality and intrinsic modeling error can reduce

confidence in the repeatability of any one analysis. However, comparing the results from

disparate approaches, each optimized to reduce overfitting, improves the certainty in the overall

importance of genes and biological processes which emerge repeatedly. For example, we found

that genes identified as important for differentiating individuals that set seed or did not set seed

through a gradient boosting classifier showed bimodal patterns of selection differentials and

gradients, with either strong positive or negative coefficients. Beyond confirming that the

expression of these genes is under strong phenotypic selection, our data suggest that a

combined approach of using machine learning to identify important genes or features, in concert

with estimating parameters for evolutionary prediction (selection differentials and gradients) may

be a useful approach for understanding selection and evolution of gene expression.

Implications from GO analysis

We found common GO terms through the union of the most influential genes in each of the

models. These common terms suggest types of genes and gene networks whose expression is

directly associated with fitness, and thus potentially experiencing selection. The terms which

were common across all models from the GO mapping results were largely related to seed

development, stress and defense responses, as well metabolic processes and light response.

Growth rate is important for fitness, and nonlinear selection analysis has revealed that it has a

slightly compensatory relationship with early flowering time (see Henry and Stinchcombe 2023),

which is itself strongly selected for in *Ipomoea hederacea* grown in field environments

(Simonsen and Stinchcombe 2010; Campitelli and Stinchcombe 2013b, Henry and Stinchcombe

2023). Metabolic and developmental processes are thus not unexpected terms to be relevant for

predicting fitness related traits. Additionally stress and defense response genes were commonly

important, suggesting that the individuals better able to respond to biotic and abiotic stressors

such as water deficiency and fungal pathogens may be better able to successfully reproduce.

One of the genes important across all both models was a heavy-metal associated isoprenylated

plant protein (LOC109174332) which, along with GO terms associated with abiotic stress lead

us to test the site soil. However, the analysis of heavy metals in the soils of our field site suggest

that, if anything, heavy metal concentrations are lower than regional averages or the typical

soils from its native range, suggesting neither an induced response to metals nor long-term

adaptation. GO enrichment analysis also suggests that genes related to iron ion starvation are

more common than expected in the sets of influential genes from two of our models. Thus, it

seems most likely that the significance of LOC109174332 lies in its role in mediating the stress

responses and plant development in the field. These data suggest that expression responses to

environmental challenges such as water deficiency and biotic interactions may be important for

fitness in *Ipomoea hederacea* in the field.

Reproductive organ development GO terms were also found to be important in both models,

which, given the observed strong selection on flowering time in this and other systems, might be

expected. We found only two GO terms significantly enriched in our model subsets: seed

maturation and vegetative to reproductive phase transition of meristem. The field site for this

experiment is 400 km north of the observed northern range limit and the significance of these

GO terms, along with shared non-identical GO terms associated with photosynthesis potentially

highlights the importance of photosynthetic genes relevant for photoperiodic cues in

development and flowering time (Shimizu et al. 2015). Our results here complement those of

Groen et al. (2020), who also detected selection on photosynthesis in a field selection

experiment on gene expression in rice under stressful environmental conditions.

Future applications and conclusions

Here we have demonstrated the applicability of machine learning in detecting genes whose

expression is associated with fitness in the field, despite a relatively small sample. The

utilization of machine learning for associating genes with fitness related traits whilst including

environmental influences has obvious applications for agricultural crop improvements. Finding

genetic targets for increased seed or biomass production is a bedrock of modern plant breeding

(Moose and Mumm 2008; International Wheat Genome Sequencing Consortium (IWGSC)

2018). Additionally, evaluating combinations of genetic features which have a concerted impact

on fitness may be used for index selection, which may provide greater gains than improvement

at single loci. More generally, similar applications of machine learning with gene expression

could assist in identifying or validating associated higher-level phenotypes for trait-based

studies or selection, especially for non-model organisms. Machine learning is a wide and

developing field, with many potential applications in improving understanding of genetic and

transcriptomic sequencing data in evolutionary biology. Despite noisy data, there is still enough

signal of fitness differences that models were moderately successful with reasonable and

affordable sample sizes, suggesting that measuring selection on gene expression with these

and related tools is a promising approach. Overall, machine learning algorithms, and their

combined results, are effective in understanding the selective importance of biological pathways

and gene expression in a multivariate context, in understanding the strength and distribution of

selection across the transcriptome and can be harnessed to find candidate genes and

quantitative traits of interest for further study.

Methods

Study species and Natural History

We collected tissue from *Ipomoea hederacea* (Convolvulaceae), an annual vine commonly

distributed throughout the eastern USA in agricultural fields, roadsides and other disturbed

habitats. It is found as far north as New Jersey and southern Pennsylvania and ranges into

Mexico. It exhibits latitudinal clines in both Mendelian and quantitative genetic traits (Bright and

Rausher 2008; Simonsen and Stinchcombe 2010; Campitelli and Stinchcombe 2013a; Stock et

al. 2014) and neutral genetic patterns consistent with metapopulation dynamics (Campitelli and

Stinchcombe 2014). Germination typically occurs in early summer, and once plants begin to

flower, they continue to do so until frost.

Field Experimental Design and Sampling

For this study we subsampled tissue for gene expression from a larger field experiment (Henry

and Stinchcombe 2023) where seeds were generated by self-fertilization in a common

greenhouse environment. On 20 July 2021, we germinated scarified seeds into Pro-Mix BX

mycorrhizae soil in 4" peat pots in a glasshouse at ambient temperature and light. Three days

later, we transplanted pots into a recently ploughed old field at the Koffler Scientific Reserve

(www.ksr.utoronto.ca), with pots placed into the ground flush with field soil; we soaked the

individuals with water and provided no further or additional supplementation. We sampled tissue

from 100 individuals from 56 populations (1-3 individuals per population, mean = 1.80),

gathered from 10 states in *I. hederacea's* eastern North American distribution, spanning ~7° of

latitude (33.017681° to 40.340767° N). The plants were left to grow naturally among emergent

weeds and/or die naturally. A killing frost ended the experiment on 27 October 2021. For further

details on the field experimental design and environment, see Henry and Stinchcombe (2023).

Based on preliminary results (*see Results*), we decided to sample soils at our field site for a

panel of heavy metals. As a consequence of the time required for initial data analysis, soils were

collected 1 year after the field experiment concluded, from the same field, which had been

plowed on the same schedule as the year of our experiment. We collected soil from across five

transects, homogenizing the soil sampled within each transect, and sent the samples to the

Agricultural and Food Laboratories at the University of Guelph for evaluation. Levels of arsenic,

cobalt, chromium, cadmium, copper, mercury, nickel, lead, and zinc were quantified.

Tissue Collection and Extraction

We collected leaf tissue a single time, on 29 September 2021, 71 days after sowing. We chose

this time point to coincide with the approximate midpoint of the life cycle when slightly more than

half of plants were flowering. The spatial block from which we sampled leaf tissue started with

342 seeds planted; at the time of sampling 33 plants had died (9.6%), and 37 had never

germinated (10.8%). From the remaining available, we chose individuals using stratified random

sampling across the population of origin. We did not have *a priori* hypotheses about particular

genes, or that expression in leaf tissue (versus roots, meristems, vascular tissue, etc) was

specifically important in determining fitness, but rather sought to obtain a multivariate sample of

the transcriptome at the approximate midpoint of the life cycle. In this way, our estimates of

gene expression traits are analogous to other studies of phenotypic selection that measure a

suite of correlated traits at one moment in time (e.g., anti-herbivory traits, size, branch

architecture, and phenology). We sampled whole leaves approximately 2.5 cm in diameter and

placed them into RNAse-free 2 mL microtubes with a 6mm sterile glass bead. We immediately

submerged the samples in liquid nitrogen and transferred them into a cooler containing dry ice

and excess liquid nitrogen until tissue collection was complete. We stored samples in a -80 °C

freezer until we performed extractions in November 2021. We extracted mRNA using Qiagen

Plant RNEasy Extraction kits and followed the standard protocol, using a TissueLyser to

prepare samples. Genome Quebec performed NEB mRNA stranded library preparations and

Illumina library quality control, prior to sequencing using paired-end Illumina mRNA sequencing

(NovaSeq6000 S4 PE100 Sequencing). Of the 100 samples collected, 97 were successfully

sequenced, representing 55 populations. Of those 97 individuals, 58 had flowered by leaf tissue

collection, 86 had flowered by the end of the experiment and 26 had set seed by the end of the

experiment, approximately 1 month later (27 October 2021). We examined the samples for

quality and contamination using FastQC, summarized with MultiQC. The mean Phred score

across all reads and samples was above 35, and no primer contamination was detected. The

mean number of reads per sample was 31.7 million (range: 18-45.4 million).

Data processing

To align the raw reads we performed a two-step STAR alignment (Dobin et al. 2013) using the

Ipomoea nil genome (Hoshino et al. 2016); *I. nil* is the sister species to *I. hederacea* (Miller et al.

1999; One Thousand Plant Transcriptomes Initiative 2019), with variable inter-crossability

between the two, mediated by style length (Esserman 2012). The genome indexes for *I. nil* were

generated with STAR default parameters except for the 'genomeSAindexNbases' parameter,

which was reduced from the default 14 to 13, given the size of the genome (Dobin et al. 2013).

On average 92% of the reads were mapped successfully to the *I. nil* genome. We then used

gffread (Pertea and Pertea 2020) to generate a transcript annotation file for use in Salmon

(Patro et al. 2017), which we ran to quantify gene counts from the aligned samples. As we

needed to compare across samples, we imported our quantified gene counts into R using

tximport to compile the samples into a single data frame and transformed the read counts using

the Trimmed-M Means transformation (Robinson and Oshlack 2010) using the *edgeR* package

(Robinson et al. 2010; McCarthy et al. 2012; Chen et al. 2016). We then filtered out genes with

very low expression where we would not expect to be able to detect differences between

individuals that never flowered and those which flowered and successfully set seed; we did so

using the *filterByExpr()* function in the edgeR package (Chen et al. 2016), which reduced the

number of genes in our dataset to 2753. As we had two groups (set seed and did not set seed),

and our set seed group had 26 individuals, we defined "very low expression" as genes where 10

counts per million were observed less in less than 19 samples (the function in edgeR finds this

value as 70% of the smallest group size).

It is common to correct estimates of gene expression for batch effects or effects due to the time

of day of collection (Leek et al. 2012; see, e.g., Josephs et al. 2020). We elected not to do so for

several related reasons. First, plants were planted in the field in randomized order, and as such

tissue collection was in random order with respect to population of origin, genotype, and other

known and unknown features of the samples. Second, our goal is to evaluate the relationship

between gene expression, which may include time-of-day effects on the order of minutes to

hours, and relative fitness, estimated from the cumulative number of seeds set by the end of the

experiment, four weeks later; fitness itself reflects ~100 days of development, life history, and

growth. Rather than focusing on individual genes or results from single models, we concentrate

on broad patterns in the intersection of different types of models, each implemented with

permutation testing and cross-validation. Finally, preliminary analysis suggested that only one of

96 principal components of gene expression showed a significant rank correlation with order of

collection; this PC was not predictive of relative fitness, and was not included in our analyses

(*see below*).

Analytical Methods

Unsupervised modeling

We first performed principal component regression for dimensionality reduction without filtering

out features. We standardized the data such that the transformed gene expression counts such

that each gene had a standard deviation = 1 and mean = 0. We removed one outlier, which had

set 86 seeds (mean seed set = 1.75, range = $[0 - 24]$, excluding the outlier), to improve

prediction. We calculated relative fitness for the remaining 96 individuals, by dividing the

individual seed set by the mean seed set. Individuals that set no seeds were retained, having

relative fitness of zero (Figure S4).

We performed a principal component analysis on the transformed and standardized data using

prcomp from the *stats* package in R (R Core Team 2022). To determine which principal

components (PCs) to include in the model we took a supervised regression approach, as lower

PCs may still possess variation which may predict relative fitness well (Chong et al. 2018). We

eliminated the last PC as the variance it described was approaching zero and was likely to

introduce multicollinearity (Jolliffe 2002).

We first ran simple linear regression with relative fitness regressed on the PC scores of each

individual and compiled the p-values of each principal component regression. We then sorted

the PCs based on those p-values and used a 100-repeat 5-fold cross-validation regression with

PCs of increasing p-values to determine the optimal number of PCs to include. We used the

train function in the *caret* package to do so, saving the model parameters and performance

measure of each iteration. We evaluated the error rates (both root mean squared and absolute

error) and R-squared values of the models to determine the best model. Our model fitting

procedure retained PCs based on their ability to predict relative fitness, and as such, PCs that

explained relatively little variation in gene expression (the trailing principal components) but

predicted fitness well were retained. We continued to add PCs until the R-squared, mean

absolute error (MAE) and root mean squared error (RMSE) stopped improving.

To evaluate the importance of each gene we back-transformed the regression coefficients using

the PCA rotations which resulted in the selection gradients (Chong et al. 2018), which account

for selection on the expression of other genes. We then sorted the genes by their absolute

coefficients to determine the 300 genes which demonstrated the strongest selection gradients,

and thus were most influential in predicting relative fitness.

Supervised modeling

We performed supervised analyses in Python using scikit-learn (Pedregosa et al. 2011). We

removed the outlier and split the data into a training set (60%, or 57 samples) and test set (40%,

or 39 samples). We again standardized the data to remove differences due to variance in

expression among genes, we mean-centered and scaled the training data and then transformed

the test data using the mean and standard deviation of the training data. For our supervised

analyses we used classification algorithms, and as such classed our samples as having set

seed (and thus having fitness > 0) or not (and having a fitness = 0).

We used a grid search to tune the hyperparameters (such as learning rate and number of

iterations) of our models, which runs models with each combination of selected model

hyperparameter values (the "grid") and compares the fit of each model. We started with coarser

grids (a wider range of sparse hyperparameter options) to search a larger parameter space and

tuning further with finer-scale grids. To manage overfitting in the model we used five-fold cross-

validation during the grid search on the training data. Model score was calculated using the

balanced accuracy score, as we had far fewer samples which set seed and balanced accuracy

accounts for uneven class sizes. The balanced accuracy score is the average accuracy score of

each group, that is, the fraction of true positive results out of the total positive cases and the

fraction of the true negative results out of the total negative cases. In our case, the "positive"

case is the "set seed" class, and the "negative" case is the "no seed set" class. We then tested

the model on the 40% of the data withheld for testing and examined out-of-sample balanced

accuracy and sensitivity of the less-common class (individuals that successfully set seed).

Sensitivity is calculated as the true positive predicted results divided by the total positive cases.

For our second classifier we used an ensemble method, which trains a large number of "weak

learners" or models that are only marginally better than random, taken together to produce a

"committee" model. We used a gradient tree boosting classifier, which is an ensemble method

that uses some number of decision trees as weak learners, progressively adjusting weights to

increase the importance of poorly classified observations from the previous tree, to build a final

model weighted by the accuracy of each base model (Hastie et al. 2009). From this model we

were able to directly extract the importance of genes, which is given as the Gini importance of

the features in the model. We selected the top 278 features, as there were 197 ties in the

importance of genes below this threshold.

GO analysis

We then sought to characterize the types of genes which were important in predicting "success"

(i.e., having set seed) in each of the models individually and which types were common among

all models, using Gene Ontology (GO) biological terms. We also performed GO enrichment

analysis on the subsets of genes from our analyses. To do so we used Blast2Go (Conesa et al.

2005), first to perform BLAST using blastx-fast of the genes to an *Arabidopsis thaliana* genome

(Lamesch et al. 2012), then EMBL-EBI Interpro scan (Cantelli et al. 2022) using default options,

followed by mapping the associated biological GO terms and annotation by Blast2Go. We

subset the most important genes in each of the models, as described above and for each

subset we repeated the BLAST, mapping and annotation of GO terms. We constructed GO term

mapping for each set and extracted the union to evaluate common GO terms. We then used

goseq (Young et al. 2010) correcting for read count, comparing the entire set of genes from our

samples to use as a reference for GO enrichment analyses of each of the model sets, using

Fisher's Exact Test with a p-value of 0.05 as a filter to generate a set of over and under

expressed GO terms. Again, we took the union of the results to look at commonly enriched GO

terms.

Acknowledgements

We gratefully acknowledge financial support from NSERC Canada Discovery Grants (JRS, GAH), Weis-Zimmerman Graduate Fellowships in field biology (GAH), and the Swedish Collegium for Advanced Study (JRS). Comments from Stephen Wright, Jacqueline Sztepanacz, Art Weis, Megan Bontrager, and Joel McGlothlin improved the manuscript. We thank 3 anonymous reviewers and Kirk Lohmueller for helpful comments. Computational support was provided by the Digital Research Alliance of Canada. We thank Kate Brown and Radana Molnarova for support in the field at the Koffler Scientific Reserve.

References

Bright, K. L., and M. D. Rausher. 2008. Natural Selection on a Leaf-Shape Polymorphism in the Ivyleaf Morning Glory (*Ipomoea hederacea*). Evolution 62:1978–1990.

Brown, K. E., and J. K. Kelly. 2022. Genome-wide association mapping of transcriptome variation in Mimulus guttatus indicates differing patterns of selection on cis- versus trans-acting mutations. Genetics 220.

Campitelli, B. E., and J. R. Stinchcombe. 2013a. Natural selection maintains a single-locus leaf shape cline in Ivyleaf morning glory, *Ipomoea hederacea*. Mol. Ecol. 22:552–564.

Campitelli, B. E., and J. R. Stinchcombe. 2014. Population dynamics and evolutionary history of the weedy vine *Ipomoea hederacea* in North America. G3 4:1407–1416.

Campitelli, B. E., and J. R. Stinchcombe. 2013b. Testing potential selective agents acting on leaf shape in *Ipomoea hederacea*: predictions based on an adaptive leaf shape cline. Ecol. Evol. 3:2409–2423.

Cantelli, G., A. Bateman, C. Brooksbank, A. I. Petrov, R. S. Malik-Sheriff, M. Ide-Smith, H.

Hermjakob, P. Flicek, R. Apweiler, E. Birney, and J. McEntyre. 2022. The European

Bioinformatics Institute (EMBL-EBI) in 2021. Nucleic Acids Res. 50:D11–D19.

Chen, Y., A. T. L. Lun, and G. K. Smyth. 2016. From reads to genes to pathways:

differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-

likelihood pipeline. F1000Res. 5:1438.

Chong, V. K., H. F. Fung, and J. R. Stinchcombe. 2018. A note on measuring natural

selection on principal component scores. Evol Lett 2:272–280.

Conesa, A., S. Götz, J. M. García-Gómez, J. Terol, M. Talón, and M. Robles. 2005.

Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics

research. Bioinformatics 21:3674–3676.

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M.

Chaisson, and T. R. Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics

29:15–21.

Eserman, L. A. 2012. Taxonomy and crossing relationships in a small group of morning

glories (Ipomoea section Pharbitis). MSc Thesis, Southeastern Louisiana University, Hammond,

Louisiana, USA.

Friedman, J. H. 2001. Greedy function approximation: a gradient boosting machine. The

Annals of Statistics 29:1189–1232.

Gilad, Y., A. Oshlack, and S. A. Rifkin. 2006. Natural selection on gene expression. Trends

Genet. 22:456–461.

Glorot, X., A. Bordes, and Y. Bengio. 2011. Deep sparse rectifier neural networks. Pp.

315–323 *in* Proc. 14th Int. Conf. Artif. Intell. Statis.

Greener, J. G., S. M. Kandathil, L. Moffat, and D. T. Jones. 2022. A guide to machine

learning for biologists. Nat. Rev. Mol. Cell Biol. 23:40–55.

Groen, S. C., I. Ćalić, Z. Joly-Lopez, A. E. Platts, J. Y. Choi, M. Natividad, K. Dorph, W. M.

Mauck 3rd, B. Bracken, C. L. U. Cabral, A. Kumar, R. O. Torres, R. Satija, G. Vergara, A.

Henry, S. J. Franks, and M. D. Purugganan. 2020. The strength and pattern of natural

selection on gene expression in rice. Nature 578:572–576.

Hastie, T., R. Tibshirani, and J. Friedman. 2009. The Elements of Statistical Learning.

Springer New York.

Henry, G. A., and J. R. Stinchcombe. 2022. Strong selection is poorly aligned with genetic

variation in *Ipomoea hederacea*: implications for divergence and constraint.

Hersch, E. I., and P. C. Phillips. 2004. Power and potential bias in field studies of natural

selection. Evolution 58:479–485.

Hoekstra, H. E., J. M. Hoekstra, D. Berrigan, S. N. Vignieri, A. Hoang, C. E. Hill, P. Beerli,

and J. G. Kingsolver. 2001. Strength and tempo of directional selection in the wild. Proc. Natl.

Acad. Sci. U. S. A. 98:9157–9160.

Hoshino, A., V. Jayakumar, E. Nitasaka, A. Toyoda, H. Noguchi, T. Itoh, T. Shin-I, Y.

Minakuchi, Y. Koda, A. J. Nagano, M. Yasugi, M. N. Honjo, H. Kudoh, M. Seki, A. Kamiya, T.

Shiraki, P. Carninci, E. Asamizu, H. Nishide, S. Tanaka, K.-I. Park, Y. Morita, K. Yokoyama, I.

Uchiyama, Y. Tanaka, S. Tabata, K. Shinozaki, Y. Hayashizaki, Y. Kohara, Y. Suzuki, S.

Sugano, A. Fujiyama, S. Iida, and Y. Sakakibara. 2016. Genome sequence and analysis of the

Japanese morning glory *Ipomoea nil*. Nat. Commun. 7:13295.

International Wheat Genome Sequencing Consortium (IWGSC). 2018. Shifting the limits in

wheat research and breeding using a fully annotated reference genome. Science 361.

James, G., D. Witten, T. Hastie, and R. Tibshirani. 2021. Introduction to Statistical

Learning: With Applications in R. Springer.

Jolliffe, I. T. 2002. Principal Component Analysis. Springer New York.

Josephs, E. B., Y. W. Lee, C. W. Wood, D. J. Schoen, S. I. Wright, and J. R. Stinchcombe.

2020. The Evolutionary Forces Shaping Cis- and Trans-Regulation of Gene Expression within a

Population of Outcrossing Plants. Mol. Biol. Evol. 37:2386–2393.

Josephs, E. B. 2021. Gene expression links genotype and phenotype during rapid

adaptation.

Kingsolver, J. G., and D. W. Schemske. 1991. Path analyses of selection. Trends Ecol.

Evol. 6:276–280.

Kingsolver, J. G., H. E. Hoekstra, J. M. Hoekstra, D. Berrigan, S. N. Vignieri, C. E. Hill, A.

Hoang, P. Gibert, and P. Beerli. 2001. The strength of phenotypic selection in natural

populations. Am. Nat. 157:245–261.

Kingsolver, J. G., and S. E. Diamond. 2011. Phenotypic selection in natural populations:

what limits directional selection? Am. Nat. 177:346–357.

Kingsolver, J. G., S. E. Diamond, A. M. Siepielski, and S. M. Carlson. 2012. Synthetic

analyses of phenotypic selection in natural populations: lessons, limitations and future

directions. Evol. Ecol. 26:1101–1118.

Kuhn, M. 2008. Building Predictive Models in R Using the caret Package. J. Stat. Softw.

28:1–26.

Lamesch, P., T. Z. Berardini, D. Li, D. Swarbreck, C. Wilks, R. Sasidharan, R. Muller, K.

Dreher, D. L. Alexander, M. Garcia-Hernandez, A. S. Karthikeyan, C. H. Lee, W. D. Nelson, L.

Ploetz, S. Singh, A. Wensel, and E. Huala. 2012. The *Arabidopsis* Information Resource (TAIR):

improved gene annotation and new tools. Nucleic Acids Res. 40:D1202–10.

Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters.

Evolution 37:1210–1226.

Leek, J. T., W. E. Johnson, H. S. Parker, A. E. Jaffe, and J. D. Storey. 2012. The sva

package for removing batch effects and other unwanted variation in high-throughput

experiments. Bioinformatics 28:882–883.

Liao, B.-Y., and M.-P. Weng. 2015. Unraveling the association between mRNA

expressions and mutant phenotypes in a genome-wide assessment of mice. Proc. Natl. Acad.

Sci. U. S. A. 112:4707–4712.

McCarthy, D. J., Y. Chen, and G. K. Smyth. 2012. Differential expression analysis of

multifactor RNA-Seq experiments with respect to biological variation. Nucleic Acids Res.

40:4288–4297.

Miller, R. E., M. D. Rausher, and P. S. Manos. 1999. Phylogenetic systematics of Ipomoea

(Convolvulaceae) based on ITS and waxy sequences. Syst. Bot. 24:209–227.

Moose, S. P., and R. H. Mumm. 2008. Molecular plant breeding as the foundation for 21st

century crop improvement. Plant Physiol. 147:969–977.

Morrissey, M. B. 2014. In search of the best methods for multivariate selection analysis.

Methods in Ecology and Evolution 5:1095–1109.

One Thousand Plant Transcriptomes Initiative. 2019. One thousand plant transcriptomes and

the phylogenomics of green plants. Nature 574:679–685.

Palakurty, S. X., J. R. Stinchcombe, and M. E. Afkhami. 2018. Cooperation and

coexpression: How coexpression networks shift in response to multiple mutualists. Mol. Ecol.

27:1860–1873.

Patro, R., G. Duggal, M. I. Love, R. A. Irizarry, and C. Kingsford. 2017. Salmon provides

fast and bias-aware quantification of transcript expression. Nat. Methods 14:417–419.

Pedregosa, F., G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, A.

Müller, J. Nothman, G. Louppe, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A.

Passos, D. Cournapeau, M. Brucher, M. Perrot, and É. Duchesnay. 2011. Scikit-learn: Machine

Learning in Python. Journal of Machine Learning Research 12:2825–2830.

Pertea, G., and M. Pertea. 2020. GFF Utilities: GffRead and GffCompare. F1000Res. 9.

Price, P. D., D. H. Palmer Droguett, J. A. Taylor, D. W. Kim, E. S. Place, T. F. Rogers, J. E.

Mank, C. R. Cooney, and A. E. Wright. 2022. Detecting signatures of selection on gene

expression. Nat. Ecol. Evol. 6:1035–1045.

Rausher, M. D. 1992. The measurement of selection on quantitative traits: Biases due to

environmental covariances between traits and fitness. Evolution 46:616–626.

R Core Team. 2022. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

Rifkin, S. A., D. Houle, J. Kim, and K. P. White. 2005. A mutation accumulation assay

reveals a broad capacity for rapid evolution of gene expression. Nature 438:220–223.

Robinson, M. D., D. J. McCarthy, and G. K. Smyth. 2010. edgeR: a Bioconductor package

for differential expression analysis of digital gene expression data. Bioinformatics 26:139–140.

Robinson, M. D., and A. Oshlack. 2010. A scaling normalization method for differential

expression analysis of RNA-seq data. Genome Biol. 11:R25.

Rockman, M. V., and L. Kruglyak. 2006. Genetics of global gene expression. Nat. Rev.

Genet. 7:862–872.

Schrider, D. R., and A. D. Kern. 2018. Supervised machine learning for population

genetics: a new paradigm. Trends Genet. 34:301–312.

Shimizu, H., K. Katayama, T. Koto, K. Torii, T. Araki, and M. Endo. 2015. Decentralized

circadian clocks process thermal and photoperiodic cues in specific tissues. Nat Plants 1:15163.

Siepielski, A. M., J. DiBattista, and S. M. Carlson. 2009. It's about time: the temporal

dynamics of phenotypic selection in the wild. Ecol. Lett. 12:1261–1276.

Siepielski, A. M., K. M. Gotanda, M. B. Morrissey, S. E. Diamond, J. D. DiBattista, and S.

M. Carlson. 2013. The spatial patterns of directional phenotypic selection. Ecol. Lett. 16:1382–

1392.

Simonsen, A. K., and J. R. Stinchcombe. 2010. Quantifying Evolutionary Genetic

Constraints in the Ivyleaf Morning Glory, *Ipomoea hederacea*. Int. J. Plant Sci. 171:972–986.

Smith, D. B., W. F. Cannon, L. G. Woodruff, F. Solano, J. E. Kilburn, and D. L. Fey. 2008.

USGS Data Series 801: Geochemical and Mineralogical Data for Soils of the Conterminous

United States.

Sokal, R. R., and F. J. Rohlf. 1995. Biometry: the principles and practice of statistics in

biological research. W.H. Freeman and Co., New York, NY, USA.

Stinchcombe, J. R., M. T. Rutter, D. S. Burdick, P. Tiffin, M. D. Rausher, and R. Mauricio.

2002. Testing for environmentally induced bias in phenotypic estimates of natural selection:

theory and practice. Am. Nat. 160:511–523.

Stock, A. J., B. E. Campitelli, and J. R. Stinchcombe. 2014. Quantitative genetic variance

and multivariate clines in the Ivyleaf morning glory, *Ipomoea hederacea*. Philos. Trans. R. Soc.

Lond. B Biol. Sci. 369:20130259.

Sun, S., B. Dong, and Q. Zou. 2021. Revisiting genome-wide association studies from

statistical modelling to machine learning. Brief. Bioinform. 22.

Sztepanacz, J. L., and D. Houle. 2024. Regularized regression can improve estimates of

multivariate selection in the face of multicollinearity and limited data. Evol Lett 8:361–373.

Oxford Academic.

Young, M. D., M. J. Wakefield, G. K. Smyth, and A. Oshlack. 2010. Gene ontology analysis

for RNA-seq: accounting for selection bias. Genome Biol. 11:R14.