Resistify - A rapid and accurate annotation tool to identify NLRs and study their genomic organisation

Moray Smith[1,2] – moray.smith@hutton.ac.uk *corresponding*

John T. Jones[1,2] – john.jones@hutton.ac.uk

Ingo Hein[1,3] – ingo.hein@hutton.ac.uk

[1] Cell and Molecular Sciences Department, The James Hutton Institute, Invergowrie, Dundee, DD2 5DA

[2] School of Biology, University of St Andrews, St Andrews, KY16 9ST

[3] School of Life Sciences, The University of Dundee, Dundee

Abstract

Background

Nucleotide-binding domain Leucine-rich Repeat (NLR) proteins are a key component of the plant innate immune system. In plant genomes NLRs exhibit considerable presence/absence variation and sequence diversity. Recent advances in sequencing technologies have made the generation of high-quality novel plant genome assemblies considerably more straightforward. Accurately identifying NLRs from these genomes is a prerequisite for improving our understanding of NLRs and identifying potential novel sources of disease resistance.

Results

Whilst several tools have been developed to predict NLRs, they are hampered by low

accuracy, speed, and availability. Here, the NLR annotation tool Resistify is presented.

Resistify is an easy-to-use, rapid, and accurate tool to identify and classify NLRs from protein

sequences. Applying Resistify to the RefPlantNLR database demonstrates that it can

correctly identify NLRs from a diverse range of species.

Applying Resistify in combination with tools to identify transposable elements to a panel of

Solanaceae genomes reveals a previously undescribed association between NLRs and

Helitron transposable elements.

Conclusion

Resistify can rapidly identify NLRs within plant genomes and provides accurate structural

classifications. Its ease of use and accessibility allows easy integration into bioinformatic

workflows and projects, enhancing the study of this important group of genes. Applying

Resistify to a *Solanaceae* pangenome reveals an undescribed association between NLRs and

transposable elements.

Availability: https://github.com/SwiftSeal/resistify

Keywords

NLRs; Plant disease resistance; transposable elements; Helitrons; Solanum

Background

The plant innate immune system is an essential part of plant disease resistance. A key

component of this are the Nucleotide-binding domain Leucine-rich Repeat (NLR) proteins –

an abundant and diverse family present in plant and animal genomes. NLRs act as

intracellular immune receptors which elicit an immune response after the detection of

pathogen derived factors or their activity (Biezen and Jones, 1998). Since their discovery,

hundreds of functional NLRs have been cloned and many sequences have been identified

(Kourelis and van der Hoorn, 2018). The abundance of NLRs and their sequence diversity

shows considerable variation across the plant kingdom (Barragan and Weigel, 2021).

One driving force of NLR diversification is thought to be the proliferation of transposable

elements (TEs). In *Arabidopsis*, sites of high TE insertion frequency have been previously

identified to contain NLRs (Quadrana *et al.*, 2016), and NLRs exhibiting elevated

intraspecific diversity are in close proximity to TEs (Sutherland *et al.*, 2023). Of the TEs, LTR-

retrotransposons are the most prolific in plant genomes and are a major factor in genome

expansion (Lee and Kim, 2014). Strikingly, the *Capsicum annuum* genome exhibits a marked

expansion of NLRs nested within LTR-retrotransposons, and this effect is seen in other

Solanaceae including tomato and potato (Seong *et al.*, 2020). Other transposable elements,

such as the Helitrons, can also capture and transport genes across the plant genome (Lai *et al.*, 2005).

Canonical NLRs have a modular organisation. Central to the NLR structure is the highly

conserved NB-ARC domain which regulates the activity of the protein. Downstream of this is

the Leucine-Rich-Repeat (LRR) domain which has a role in ligand binding and auto-inhibition.

Upstream of the NB-ARC domain is variable. Often, there is either a coiled-coil (CC),

Resistance to Powdery Mildew 8-like (RPW8), or Toll/Interleukin-1 receptor/Resistance

protein (TIR) domain. Together, this modular structure means that NLRs can be separated

into different classifications: CNL, RNL, TNL, or NL if lacking a conserved N-terminal domain.

Recently, additional NLR associated signatures have been identified. For example, the post-

LRR C-terminal jelly-roll/Ig-like domain (C-JID) (Martin *et al.*, 2020) may be present in TNLs

and appears to bind with pathogen effectors, thereby contributing to the immune response.

Similarly, other NLRs have integrated decoys to detect pathogen activities. In terms of

initiating cell death through the formation of a resistosome, CNLs rely on an N-terminal

MADA motif, a hallmark of NRCs - NLRs which act as intermediaries in the NLR induced

immune response or NLRs that function independently of NRCs (Adachi *et al.*, 2019).

Due to their conservation and modular structure, NLRs lend themselves to automated

identification and classification. To date, several tools have been developed to achieve this

including DRAGO2, NLGenomeSweeper, NLR-Annotator, RGAugury, RRGPredictor, and

NLRtracker (table 1) (Li *et al.*, 2016; Osuna-Cruz *et al.*, 2018; Santana Silva and Micheli,

2020; Steuernagel *et al.*, 2020; Toda *et al.*, 2020; Kourelis *et al.*, 2021). The development

of the RefPlantNLR database has allowed the benchmarking of these tools against the

sequences of functionally characterised NLRs (Kourelis *et al.*, 2021). Performance varied

between tools with NLRtracker being the most sensitive and accurate. All tools perform well

at the identification of TIR, RPW8, and NB-ARC domains which are highly conserved, but are

often less accurate in predicting CC domains which are more variable and frequently missed

by InterProScan (Jones *et al.*, 2014). Additional to the currently available NLR classification

tools is NLRexpress – a set of machine learning predictors for CC, TIR, NB-ARC, and LRR

motifs (Martin *et al.*, 2022). Although NLRexpress does not directly identify or classify NLRs,

it is well suited for rapidly and accurately screening large sets of sequences for NLR-

associated motifs. One strength of NLRexpress is the CC-associated extended EDVID model

which performs strongly in identifying this challenging motif.

Table 1: A summary of the currently available NLR annotation tools. Adapted from (Kourelis

et al., 2021).

Tool	Input data	Method	Output	Distribution
DRAGO2	Protein, Transcript	Custom HMMs	Classification, Domains	Online only, API available
NLGenomeSweeper	Transcript, Genomic	InterProScan, MUSCLE, TransDecoder, BLAST, HMMER	Classification, Genome position, GFF annotation	GitHub, Manual dependency installation
NLR-Annotator	Transcript, Genomic	MEME motifs	Classification, Genome position, GFF annotation	GitHub
RGAugury	Transcript, Genomic	InterProScan	Classification,	GitHub,

			Genome position, GFF annotation	Online or local webservice, Docker container
RRGPredictor	Protein, Transcript	InterProScan	Classification	GitHub, Manual dependency installation
NLRtracker	Protein, Transcript	InterProScan, HMMER, MEME motifs	Classification, NB-ARC sequence, Domains, GFF annotation	GitHub, Manual dependency installation
NLRexpress	Protein	HMMER, Scikit-learn	Motif position, Motif sequence	GitHub, Online or local, Conda environment provided

One drawback with the currently available tools is the reliance on InterProScan as the

backend for domain annotation. InterProScan is designed as a comprehensive and

generalised domain annotation tool. As a result, each input sequence is scanned against

several databases each containing in total more than 180,000 protein signatures, the vast

majority of which are not NLR associated. Compounding this, NLRs only represent a fraction

of a plant proteome resulting in unnecessary searches against non-NLR sequences. In

addition, domains common to NLRs are present in InterProScan databases with differing

levels of curation which often results in overlapping or fragmented annotations. These must

be parsed, particularly the LRR domain which is represented by multiple InterProScan

signatures. The large databases that InterProScan relies upon must also be downloaded

which provides an additional challenge for installation or automated deployment.

Software that is readily available with minimum manual configuration is a requirement for

genome assembly and annotation projects. Projects may use an array of different tools,

making manual installation and dependency resolution time-consuming. Dependency

resolution solutions such as Conda, and automated deployment of tools in pipelines with

Snakemake or Nextflow are becoming increasingly common (Di Tommaso *et al.*, 2017;

Mölder *et al.*, 2021). Additionally, bioinformatics research is often restricted to shared High-

Performance Clusters (HPCs) where users have limited privileges and where storage is at a

premium. Tools that require root privileges, are distributed as web services, or require large

databases are often not conducive for genomics projects.

Here, the new NLR annotation tool Resistify is presented that overcomes some of the

constraints of currently available tools. We show that this tool accurately predicts NLR

sequences from diverse plant using a range of well curated sources and can be used for

pangenome analysis of NLRs in *Solanaceous* genomes. In addition, we use Resistify in

combination with the EDTA tool to investigate the genomic organisation of NLRs and their

association with transposable elements.

Methods

Resistify is implemented in python as a command line executable. First, Resistify executes a

hmmsearch of input protein sequences against a custom database of HMMs derived from

curated Pfam entries (table 2). These models are used to identify CC, RPW8, TIR, and NB-ARC

domains, as well as C-JID and MADA motifs. Domains of the same type are merged if they

overlap or are within 100 amino acids of each other. Preliminary testing in development

showed that this is necessary to overcome NB-ARC domain annotations which can become

split. MADA motifs are restricted to the N-terminal of the sequence. Using this evidence,

proteins are initially classified as belonging to either a CN, RN, TN, or N classification.

Proteins which do not have any evidence of an NB-ARC domain are discarded at this stage.

Table 2: HMM models including in the initial hmmsearch stage of Resistify.

Domain	Source
NB-ARC	https://www.ebi.ac.uk/interpro/entry/pfam/PF00931/
TIR	https://www.ebi.ac.uk/interpro/entry/pfam/PF01582/
TIR	https://www.ebi.ac.uk/interpro/entry/pfam/PF13676/
RPW8	https://www.ebi.ac.uk/interpro/entry/pfam/PF05659/
CC	https://www.ebi.ac.uk/interpro/entry/pfam/PF18052/
C-JID	https://www.ebi.ac.uk/interpro/entry/pfam/PF20160/
MADA	https://cdn.elifesciences.org/articles/49956/elife-49956-supp2-v2.hmm

Following this, the filtered set of NB-ARC containing proteins is screened for NLR-associated

motifs with the NLRexpress models. Sequences are then reclassified using the following

logic. If a protein belongs to class N (i.e., does not have any evidence of an upstream CC,

RPW8, or TIR domain) it is scanned for upstream TIR or CC motifs. As motif searches can be

more promiscuous, restricting motif searches to this condition prevents them from

interfering with non-ambiguous NLR classification. The sequence is then screened for LxLxxL

motifs to define the LRR domain. Following a previous definition, an LRR domain is

annotated if four or more LxLxxL motifs are identified with inter-motif gaps of less than 75

amino acids [@martin_nlrscape_2023]. Gaps larger than 75 amino acids are predicted to be

a break in the LRR domain and so the LRR annotation process is restarted from that position

onwards. If less than four motifs exist across the whole sequence this process is skipped.

This combined evidence is then integrated into the domain annotation data. The domains

are sorted by start position and a "domain string" is formed. For example, if the sorted

domains took the order TIR, NB-ARC, LRR, C-JID, then the domain string would be TNLj.

Alternatively, a canonical CNL with a MADA motif would take the form mCNL. The domain

strings are searched for substrings CNL, RNL, TNL, or NL and classified accordingly.

Sequences are further classified on the basis of whether they contain an N-terminal bound

MADA motif or CJID domain. Additionally, a count of the unique occurrences of the

conserved NB-ARC motifs is made as an indicator of NB-ARC integrity.

The primary output of Resistify is a table detailing the NLRs identified and the specific

classification for each sequence. The complete domain string, classification, NB-ARC motif

count, and MADA and CJID status are listed. A complete list of all annotations and

NLRexpress motifs are given as separate tables. Additionally, all NB-ARC domain sequences

are extracted and provided as a FASTA file.

Resistify is provided as a single executable available for download on GitHub, PyPi, and

Conda. All databases and models are distributed with the executable so that manual setup is

not required. All output files are placed in a nameable output directory for easy integration

into automated workflows. Temporary files are also handled internally to reduce output

clutter. Resistify requires scikit-learn v0.24.2 and hmmer v>=3.0.

RefPlantNLR benchmarking

174   Protein sequences of the RefPlantNLR database members were retrieved and used as input

175   for Resistify with default settings (Kourelis *et al.*, 2021). Resistify classifications were

176   compared directly in R v4.3.2 with tidyverse v2.0.0. Any sequence where the Resistify

177   classification did not exactly match with the provided RefPlantNLR structure was taken as a

178   potential misclassification.

Araport11 benchmarking

181   The latest release of Araport11 representative gene model protein sequences were

182   downloaded from TAIR and used as input for Resistify with default settings(Lamesch *et al.*,

183   2012). A phylogenetic tree was built from the Resistify-extracted NB-ARC domain sequences

184   with mafft v7.52.0 and fasttree v2.1.11 with default settings.

Pangenome pipeline

A Snakemake workflow was developed to predict genes and NLRs in a Solanum pangenome comprised of chromosome-scale genomes. Snakemake was executed in a mamba v1.4.2 environment with snakemake v7.32.3 and cookiecutter v1.7.3 (Mölder *et al.*, 2021). Genes were predicted *de-novo* from the genome alone using Helixer v0.3.2 with the land_plant_v0.3_a_0080.h5 model (Holst *et al.*, 2023). Protein sequences were extracted using AGAT v1.2.0 and used as input for Resistify (Dainat *et al.*, 2023). Ortholog analysis was carried out using OrthoFinder v2.5.5 using all predicted protein sequences (Emms and Kelly, 2019). Transposable elements were annotated with the latest GitHub release of EDTA (Ou *et al.*, 2019).

Gene models were translated to bed format using AGAT v1.2.0 and overlaps with intact transposable elements were identified using bedtools v2.31.1 with the command bedtools intersect using the options -f 0.9 -wo (Quinlan and Hall, 2010). To identify previously characterised NLRs, protein sequences of a subset of *Solanum*-originating NLRs in the RefPlantNLR database were queried against each genome with blastp v2.15.0 (Camacho *et al.*, 2009). The full pipeline and all post-hoc analysis are available at https://github.com/SwiftSeal/pangenomics/.

Results

207 Performance against RefPlantNLR

To evaluate the performance of Resistify (figure 1), it was applied against the RefPlantNLR

database - a curated set of 415 previously cloned NLRs from a diverse range of species.

NLRtracker was also applied to the RefPlantNLR database for comparison.

Figure 1: The Resistify program.

In the default mode, only three RefPlantNLR entries were not identified by Resistify -

AtNRG1.3 which carries only an LRR domain and *Pb1* and *RXL* which have significantly

truncated NB-ARC domain that are not listed in the RefPlantNLR database. However, if

required, these genes can be identified with Resistify using –ultra mode which skips the

initial filtering stage and reports and sequence which contains at least one NLRexpress motif.

Sequences are reported as an unmerged string of NLR motifs. Consequently, *AtNRG1.3*, *Pb1*,

and *RXL* are reported as NNLLLLLLL, CNNNLLLLLLLLLLL, and CNNNLLLLLLLLL respectively.

The largest source of variation between RefPlantNLR and Resistify classification was 29 NLRs

which had an NL structure according to RefPlantNLR but CNL according to Resistify. All of

these belonged to CNL-associated subclasses which are known to have CC domains that are

challenging to identify. This included 15 CC_{G10}-NLRs including *Pvr4*, *Tsw*, *RPS2*, *RPS5*, *SUT1*

and *SUMM2* which have previously been noted for their lack of the CC-associated EDVID

motif (Lee *et al.*, 2021). Others included six members of the *Pm5* locus which despite not

containing a CC domain in RefPlantNLR, have previously been identified to contain CC-like

domains (Xie *et al.*, 2020). This analysis demonstrates that Resistify is highly sensitive at

retrieving canonical NLRs and accurately describing their structure.

As NLRtracker does not exclude NLRs without NB-ARC domains, *AtNRG1.3*, *Pb1* and *RXL*

were identified and classified as "CC-NLR or CCR-NLR or CCG10-NLR", "CC-NLR", and "CC-

NLR" respectively. NLRtracker relies on the NB-ARC associated RNBS-D motif to classify NLRs

as "CC-NLR or CCR-NLR or CCG10-NLR". As a result, 26 of the conflicting CNLs identified by

Resistify were classified as "CC-NLR or CCR-NLR or CCG10-NLR" despite failing to identify a

CC domain. The exception of this was for *SpNBS-LRR*, *Rpp1-R1*, and *Rpp4C4* which were

classified as "UNDETERMINED".

Reliance on RNBS-D also led to classification of *NtTPN1* as "CC-NLR or CCR-NLR or CCG10-

NLR". Whilst *NtTPN1* is a CNL ortholog, it lacks any upstream domain and is structurally an

NL. Unexpectedly, the TNL *DSC1* was misclassified as "CC-NLR" despite having a domain

structure of "(TIR)(NBARC)(LRR)(CJID)" according to NLRtracker.

In summary, Resistify performs well at identifying canonical NLRs from a diverse range of

species. In default mode, it does not assign genes as NLRs with extremely truncated or

entirely absent NB-ARC domains, unlike NLRtracker which reports any sequence with NLR-

associated domains. However, this can be replicated in Resistify with the –ultra mode. Thus,

Resistify's structure-based classification method is well suited for correctly classifying NLRs,

including members of challenging CNL subclasses.

Performance against the Araport11 proteome

To assess the performance of Resistify across a well characterised and annotated genome,

the representative gene models of Araport11 were analysed (Cheng *et al.*, 2017). In total,

Resistify identified 166 NLRs - the majority of which were TNLs and CNLs (figure 2a). Of the

CNLs, 25% had a MADA motif, and 44.4% and 41.2% of NLs and TNLs had C-JID domain

respectively. Partial NLRs either without an N-terminal or LRR domain were also identified.

Figure 2: Resistify applied to Araport11.

a) The number of NLRs belonging to each identified classification. b) The number of NLRs

grouped by number of identified unique NB-ARC associated motifs. c) A phylogenetic tree of

NLRs based on the Resistify extracted NB-ARC domain sequences.

Manual inspection of the motifs within these sequences confirmed that they were not due

to a failure to identify these elements. The majority of NLRs carried all nine conserved NB-

ARC motifs - those with fewer increasingly belonged to partial NLR classifications (figure 2b).

NLRs with as few as two of the conserved NB-ARC motifs were successfully identified. A

phylogenetic tree of the Resistify-identified NB-ARC domains independently validated the

classifications and allowed the placement of ambiguous NL or N classified sequences into

subclasses (figure 2c).

NLRtracker identified an additional 48 sequences, however these sequences did not contain

any NB-ARC domain annotation according to either tool. Three sequences - AT4G19060.1,

AT4G19060.1, and AT5G45440.1 - were not identified by NLRtracker. According to Resistify

these contained a single NB-ARC domain each, with 5, 6, and 6 NB-ARC motifs respectively.

Overall, Resistify performs well at identifying and classifying NLRs from whole proteomes

and successfully retrieves NB-ARC domain annotations for phylogenetic analyses.

Application against an example workflow

To demonstrate how Resistify might be implemented to identify novel resistance genes, a

pangenome experiment was performed. Eighteen contiguous *Solanum* genomes were

downloaded and processed with a simple workflow which predicts genes de novo, identifies

orthologues, and classifies NLRs with Resistify (Tang *et al.*, 2022; Li *et al.*, 2023). For gene

annotations, the recently developed tool Helixer was selected for its near reference quality

predictions and lack of requirement for repeat-masking which is known to result in false-

negative NLR annotations (Bayer, Edwards and Batley, 2018; Holst *et al.*, 2023). The

workflow also predicts transposable elements with EDTA. The *C. annuum* genome was

included as an outgroup, but also because recent analysis has suggested a substantial

expansion of NLRs in this species associated with LTR transposable elements (Kim *et al.*,

2017).

Predicted gene content varied between genomes from 29,223 in *S. habrochaites* to 61,015 in

the larger *C. annuum* genome (table S1). Transposable element content ranged from 55.6%

in *S. chmielewskii* to 76.8% in *C. annuum*. Whilst there was no significant difference in total

TE content between tuber-bearing and non-tuber-bearing genomes ($p = 0.644$), there was a

significant increase in the number of intact TEs reported by EDTA in the genomes of tuber-

bearing species ($p = 0.008$).

In total, 8144 NLRs were identified across all genomes (figure 3). CNLs were the most

abundant classification of NLR identified, ranging from 84 in *S. habrochaites* to 422 in *S.*

tuberosum (group tuberosum RH10-15). There was a notable expansion of NLRs in tuber-

bearing *Solanum* species. This is in agreement with the previous observation that potato-

bearing *Solanum* species have an expansion of tuber expressed NLRs (Tang *et al.*, 2022).

This effect does not correlate with the proportion of genome occupied by transposable

elements of any classification (figure S1).

Figure 3: NLRs identified across the *Solanum* pangenome.

In total, 38,590 orthogroups were identified, of which 687 (1.8%) contained NLRs. The

distribution of orthogroups showed a clear divide between core and shell/cloud orthogroups

within the pangenome with the majority of NLR orthogroups existing within the shell/cloud

(fig 4a). This reflects the relatively large width of the pangenome, which captures NLR

variation over a Genus level. Species specific NLRomes also exhibit an abundance of cloud

orthographs, reflecting the high variability of NLRs in genomes (Van de Weyer *et al.*, 2019).

Many (48.1%, n = 371) orthogroups were classified as containing N or NL NLRs (figure 4b).

Whilst these classifications are less abundant (24.6%, n = 2081), they were often associated

with smaller orthogroups contributing to this inflation.

Figure 4: NLR-containing orthogroups.

a) The number of orthogroups shared between genomes. b) The number of orthogroups

according to NLR classification. c) The number of orthologs of known *Solanaceae* NLRs

identified in each genome.

The distribution of previously identified *Solanaceae* NLR orthologs across the pangenome

was examined (figure 4c). In agreement with previous findings, members of the NRC group

remained relatively stable across the pangenome except for NRC4a, NRC4b, and NRC6 which

were expanded in the tuber-bearing genomes. By contrast, an expansion of *Mi-1* and *Hero*

was evident in the non-tuber bearing and tuber bearing genomes respectively. At least one

ortholog was identified in each genome for all genomes.

It has previously been reported that there is a vast expansion of NLRs in the *C. annum*

genome due to retroduplication; NLRs nested within LTRs represented a large proportion

(~13%) of NLRs within the genome, and this effect is also seen in tomato (8%) and potato

(18%) (Kim *et al.*, 2017). To explore whether this effect could be linked to the expansion of

NLRs within the tuberising members of *Solanum*, a similar analysis was repeated. Intact TEs

were identified and considered to interact with NLRs if they covered >90% of the gene

annotation.

Unexpectedly, the effect could not be replicated and across all genomes only five intact LTRs

were identified as containing NLRs (figure 5a). None were identified in *C. annuum* but

instead in the *S. tuberosum* group Phureja and clone RH. The putative retrotransposed NLRs

within these all belonged to the same orthogroup (OG0000639) which was expanded in both

Phureja and RH, but not *S. stenotomum*.

Figure 5: Helitrons associated with NLRs.

a) The number of intact TEs overlapping with NLRs across the pangenome. b) The 5' motif of

NLR overlapping Helitrons. c) The 3' motif of NLR overlapping Helitrons. d) The *NRC1* locus

and its association with Helitrons across four genomes. Helitron (blue) and LTR (orange)

transposable elements are highlighted.

By contrast, a proliferation of Helitron-associated NLRs was identified in the tuber-bearing

genomes. All predicted Helitrons carried the expected 5'-TC…CTRR-3' signature as well as a

GC-rich region in the 3' terminal (figure 5b,c). An example of an NLR which may have

undergone Helitron-associated duplication is *NRC1* (figure 5d) which is present as a single

copy except in both *S. tuberosum* group Phureja genomes and *S. verrucosum*. Close

examination of the *NRC1* locus reveals a duplication event in *S. tuberosum* group Phureja

where one *NRC1* gene is nested entirely within a predicted Helitron. In *S. tuberosum* clone

RH which only has one copy of *NRC1*, *NRC1* is similarly nested within a Helitron although this

appears to be extended to a distant Helitron terminator past the site of duplication in group

Phureja. In *S. stenotomum*, only one *NRC1* is present and no Helitron predicted.

Interestingly, *S. verrucosum* has two copies of *NRC1* and whilst it does not have a nested

Helitron copy, the leftmost *NRC1* copy does have a short predicted Helitron upstream of its

start position in the same orientation as the other Helitron elements. In this case, the two

copies of *NRC1* appear to have been further separated by an intact LTR insertion which has

resulted in an additional gene prediction.

Discussion

Resistify is presented as a highly accurate and easy-to-use tool to aid the identification of

novel resistance genes. Applying Resistify against the RefPlantNLR database demonstrates

that it is effective at identifying NLRs from a diverse range of plant genomes. It is highly

sensitive at classifying challenging CNL families which often evade other NLR classification

tools due to the inclusion of the performant NLRexpress motif models. Members of the

CC_{G10}-NLR clade have previously been described as lacking the CC-associated EDVID motif

and are often predicted as having an NL structure in the RefPlantNLR database (Lee *et al.*,

2021). By utilising NLRexpress' highly sensitive extended EDVID motif model, Resistify

successfully classifies these elusive NLR clades, highlighting it as the most sensitive NLR

classifier to date.

Resistify can be easily integrated into workflows and is scalable to large pangenome

experiments. As high-quality genome assemblies become more common, providing

accessible tools for the study of NLRs will be crucial for fully appreciating their diversity and

identifying novel sources of resistance. The recent releases of chromosome level assemblies

for potato and tomato pangenomes are a valuable resource for understanding the diversity

of NLRs within *Solanum* (Tang *et al.*, 2022; Li *et al.*, 2023). An expansion of NLRs including

essential families such as the NRCs is apparent in tuber-bearing species.

As Resistify relies on an initial NB-ARC domain search to reduce the search space prior to

NLRexpress motif identification, NLRs with very truncated or entirely absent NB-ARC

domains can be missed. To resolve this, an additional mode is provided which does not

perform initial NB-ARC domain filtering which identifies sequences with any NLR motifs.

Unlike NLRtracker, Resistify does not search for noncanonical integrated domains (Kroj *et al.*,

2016). As integrated domains are widespread and often critical to NLR function, their

identification and analysis are an important factor in studies of NLR diversity (Barragan and

Weigel, 2021). Pairing Resistify with a secondary sweep for these domains with NLRtracker

or InterProScan would permit this analysis whilst also benefiting from a reduced search

space and high-quality NLR classifications from Resistify.

An unexpected finding of this study was the failure to replicate the previous observation of

an abundance of LTR associated NLRs across *Solanaceae* (Kim *et al.*, 2017). Differences in

gene and TE annotation and NLR classification likely contributed to this. Although Helixer

predicted ~70% more genes in *C. annuum* than were identified in the previous study, this did

not translate to an increase in NLRs identified. Instead, only 673 NLRs were identified here in

comparison to the previous estimate of 835. Previously, a tBLASTn search of the NB-ARC

domain followed by ORF identification and BLASTP searches against GenBank to classify

NLRs was used to identify NLRs in the genome (Seo *et al.*, 2016). This method may result in

more false positives in comparison to identifying NLRs directly from gene annotations. Here,

EDTA was selected for identifying TEs which has an improved sensitivity and selectivity for

identifying intact LTRs in comparison to LTRHarvest which was used previously. Although

Helixer does not require repeat masking which can result in false negative NLR annotations,

annotations used in training the model (which may have relied upon repeat masking) might

introduce TE-avoidant behaviour (Bayer, Edwards and Batley, 2018). However, the

identification of several LTR-associated NLRs indicates that this is unlikely to be the source of

the difference in the result.

The identification of Helitron-associated NLR expansion has not been previously reported.

Helitrons are challenging to identify due to their lack of structural elements and as a result

the pipeline used in this study is known to overestimate Helitron density (Baril, Imrie and

Hayward, 2022). However, a large part of this overestimation is likely due to EDTA

reannotating the genome with HelitronScanner predictions, which itself suffers from a high

false positive rate, leading to a proliferation of fragmented Helitron annotations across the

genome. For this study, only intact Helitrons which passed EDTAs stringent filter were

considered. As a result, all predicted Helitrons had the required structural elements for

activity. Further validation would be required to determine if Helitron association is a valid

mechanism of NLR expansion. The Helitron/*NRC1* relationship highlighted here would be a

good starting point for unpicking this mechanism.

Conclusions

Resistify is presented as a new tool to aide in the study of NLRs in plant genomes. It provides

high-quality structural classifications for canonical NLRs. Applying Resistify to the

RefPlantNLR database demonstrates that it can correctly classify NLRs from a diverse range

of plant species. Applying Resistify to a panel of *Solanaceae* genomes demonstrates its

suitability for studying NLRs in large datasets. This analysis revealed functionally important

NLR families undergoing expansion in tuberising *Solanum* species. Unexpectedly, combining

this with transposable element annotations did not identify a previous description of NLRs

embedded within LTR retrotransposons in *Solanaceae* genomes. Instead, an undescribed

association between NLRs and Helitron transposable elements in genomes with expanded

NLR inventories was revealed.

Acknowledgements

445 The authors acknowledge the Research/Scientific Computing teams at The James Hutton

446 Institute and NIAB for providing computational resources and technical support for the "UK's

447 Crop Diversity Bioinformatics HPC" (BBSRC grant BB/S019669/1), use of which has

448 contributed to the results reported within this paper. The authors also acknowledge Peter

449 Cock for his support with creating the conda distribution of Resistify.

Funding

This work was supported by the Rural & Environment Science & Analytical Services (RESAS) Division of the Scottish Government through the project JHI-B1-1, the Biotechnology and Biological Sciences Research Council (BBSRC) through awards BB/S015663/1 and BB/X009068/1. MS was supported through the East of Scotland Bioscience Doctoral Training Partnership (EASTBIO DTP), funded by the BBSRC award BB/T00875X/1.

Availability of data and materials

Resistify is available at https://github.com/SwiftSeal/resistify. All data analysis workflows and scripts are available at https://github.com/SwiftSeal/pangenomics. The RefPlantNLR database was accessed from its source publication (Kourelis *et al.*, 2021). The genomes used in this study were accessed from their source publications (Tang *et al.*, 2022; Li *et al.*, 2023).

References

Adachi, H. *et al.* (2019) 'An N-terminal motif in NLR immune receptors is functionally conserved across distantly related plant species', *eLife.* Edited by J.-M. Zhou, D. Weigel, and J.-M. Zhou, 8, p. e49956. Available at: https://doi.org/10.7554/eLife.49956.

Baril, T., Imrie, R.M. and Hayward, A. (2022) 'Earl Grey: a fully automated user-friendly transposable element annotation and analysis pipeline'. bioRxiv, p. 2022.06.30.498289. Available at: https://doi.org/10.1101/2022.06.30.498289.

Barragan, A.C. and Weigel, D. (2021) 'Plant NLR diversity: the known unknowns of pan-

NLRomes', *The Plant Cell*, 33(4), pp. 814–831. Available at:

https://doi.org/10.1093/plcell/koaa002.

Bayer, P.E., Edwards, D. and Batley, J. (2018) 'Bias in resistance gene prediction due to repeat

masking', *Nature Plants*, 4(10), pp. 762–765. Available at: https://doi.org/10.1038/s41477-

018-0264-0.

Biezen, E.A.V.D. and Jones, J.D.G. (1998) 'Plant disease-resistance proteins and the gene-for-

gene concept', *Trends in Biochemical Sciences*, 23(12), pp. 454–456. Available at:

https://doi.org/10.1016/S0968-0004(98)01311-5.

Camacho, C. *et al.* (2009) 'BLAST+: architecture and applications', *BMC Bioinformatics*, 10(1),

p. 421. Available at: https://doi.org/10.1186/1471-2105-10-421.

Cheng, C.-Y. *et al.* (2017) 'Araport11: a complete reannotation of the Arabidopsis thaliana

reference genome', *The Plant Journal*, 89(4), pp. 789–804. Available at:

https://doi.org/10.1111/tpj.13415.

Dainat, J. *et al.* (2023) 'NBISweden/AGAT: AGAT-v1.2.0'. Zenodo. Available at:

https://doi.org/10.5281/zenodo.8178877.

Di Tommaso, P. *et al.* (2017) 'Nextflow enables reproducible computational workflows',

Nature Biotechnology, 35(4), pp. 316–319. Available at: https://doi.org/10.1038/nbt.3820.

Emms, D.M. and Kelly, S. (2019) 'OrthoFinder: phylogenetic orthology inference for

comparative genomics', *Genome Biology*, 20(1), pp. 1–14. Available at:

https://doi.org/10.1186/s13059-019-1832-y.

Holst, F. *et al.* (2023) 'Helixer–de novo Prediction of Primary Eukaryotic Gene Models

Combining Deep Learning and a Hidden Markov Model'. bioRxiv, p. 2023.02.06.527280.

Available at: https://doi.org/10.1101/2023.02.06.527280.

Jones, P. *et al.* (2014) 'InterProScan 5: genome-scale protein function classification',

Bioinformatics, 30(9), pp. 1236–1240. Available at:

https://doi.org/10.1093/bioinformatics/btu031.

Kim, S. *et al.* (2017) 'New reference genome sequences of hot pepper reveal the massive

evolution of plant disease-resistance genes by retroduplication', *Genome Biology*, 18(1), p.

210. Available at: https://doi.org/10.1186/s13059-017-1341-9.

Kourelis, J. *et al.* (2021) 'RefPlantNLR is a comprehensive collection of experimentally

validated plant disease resistance proteins from the NLR family', *PLOS Biology*, 19(10), p.

e3001124. Available at: https://doi.org/10.1371/journal.pbio.3001124.

Kourelis, J. and van der Hoorn, R.A.L. (2018) 'Defended to the Nines: 25 Years of Resistance

Gene Cloning Identifies Nine Mechanisms for R Protein Function', *The Plant Cell*, 30(2), pp.

285–299. Available at: https://doi.org/10.1105/tpc.17.00579.

Kroj, T. *et al.* (2016) 'Integration of decoy domains derived from protein targets of pathogen

effectors into plant immune receptors is widespread', *New Phytologist*, 210(2), pp. 618–626.

Available at: https://doi.org/10.1111/nph.13869.

Lai, J. *et al.* (2005) 'Gene movement by Helitron transposons contributes to the haplotype

variability of maize', *Proceedings of the National Academy of Sciences*, 102(25), pp. 9068–

9073. Available at: https://doi.org/10.1073/pnas.0502923102.

Lamesch, P. *et al.* (2012) 'The Arabidopsis Information Resource (TAIR): improved gene

annotation and new tools', *Nucleic Acids Research*, 40(D1), pp. D1202–D1210. Available at:

https://doi.org/10.1093/nar/gkr1090.

Lee, H. *et al.* (2021) 'Genome-wide functional analysis of hot pepper immune receptors

reveals an autonomous NLR clade in seed plants', *The New Phytologist*, 229(1), pp. 532–547.

Available at: https://doi.org/10.1111/nph.16878.

Lee, S.-I. and Kim, N.-S. (2014) 'Transposable Elements and Genome Size Variations in Plants',

Genomics & Informatics, 12(3), pp. 87–97. Available at:

https://doi.org/10.5808/GI.2014.12.3.87.

Li, N. *et al.* (2023) 'Super-pangenome analyses highlight genomic diversity and structural

variation across wild and cultivated tomato species', *Nature Genetics*, 55(5), pp. 852–860.

Available at: https://doi.org/10.1038/s41588-023-01340-y.

Li, P. *et al.* (2016) 'RGAugury: a pipeline for genome-wide prediction of resistance gene

analogs (RGAs) in plants', *BMC Genomics*, 17(1), p. 852. Available at:

https://doi.org/10.1186/s12864-016-3197-x.

Martin, E.C. *et al.* (2022) 'NLRexpress—A bundle of machine learning motif predictors—

Reveals motif stability underlying plant Nod-like receptors diversity', *Frontiers in Plant*

Science, 13. Available at: https://www.frontiersin.org/articles/10.3389/fpls.2022.975888

(Accessed: 23 October 2023).

Martin, R. *et al.* (2020) 'Structure of the activated ROQ1 resistosome directly recognizing the

pathogen effector XopQ', *Science*, 370(6521), p. eabd9993. Available at:

https://doi.org/10.1126/science.abd9993.

Mölder, F. *et al.* (2021) 'Sustainable data analysis with Snakemake'. F1000Research. Available

at: https://doi.org/10.12688/f1000research.29032.2.

Osuna-Cruz, C.M. *et al.* (2018) 'PRGdb 3.0: a comprehensive platform for prediction and

analysis of plant disease resistance genes', *Nucleic Acids Research*, 46(D1), pp. D1197–

D1201. Available at: https://doi.org/10.1093/nar/gkx1119.

Ou, S. *et al.* (2019) 'Benchmarking transposable element annotation methods for creation of

a streamlined, comprehensive pipeline', *Genome Biology*, 20(1), p. 275. Available at:

https://doi.org/10.1186/s13059-019-1905-y.

Quadrana, L. *et al.* (2016) 'The Arabidopsis thaliana mobilome and its impact at the species

level', *eLife*. Edited by D. Zilberman, 5, p. e15716. Available at:

https://doi.org/10.7554/eLife.15716.

Quinlan, A.R. and Hall, I.M. (2010) 'BEDTools: a flexible suite of utilities for comparing

genomic features', *Bioinformatics*, 26(6), pp. 841–842. Available at:

https://doi.org/10.1093/bioinformatics/btq033.

Santana Silva, R.J. and Micheli, F. (2020) 'RRGPredictor, a set-theory-based tool for predicting

pathogen-associated molecular pattern receptors (PRRs) and resistance (R) proteins from

plants', *Genomics*, 112(3), pp. 2666–2676. Available at:

https://doi.org/10.1016/j.ygeno.2020.03.001.

Seo, E. *et al.* (2016) 'Genome-Wide Comparative Analyses Reveal the Dynamic Evolution of

Nucleotide-Binding Leucine-Rich Repeat Gene Family among Solanaceae Plants', *Frontiers in*

Plant Science, 7. Available at: https://www.frontiersin.org/articles/10.3389/fpls.2016.01205

(Accessed: 20 November 2023).

Seong, K. *et al.* (2020) 'Evolution of NLR resistance genes with noncanonical N-terminal

domains in wild tomato species', *New Phytologist*, 227(5), pp. 1530–1543. Available at:

https://doi.org/10.1111/nph.16628.

Steuernagel, B. *et al.* (2020) 'The NLR-Annotator Tool Enables Annotation of the Intracellular

Immune Receptor Repertoire1 [OPEN]', *Plant Physiology*, 183(2), pp. 468–482. Available at:

https://doi.org/10.1104/pp.19.01273.

Sutherland, C.A. *et al.* (2023) 'High intraspecies allelic diversity in Arabidopsis NLR immune

receptors is associated with distinct genomic and epigenomic features'. bioRxiv, p.

2023.01.12.523861. Available at: https://doi.org/10.1101/2023.01.12.523861.

Tang, D. *et al.* (2022) 'Genome evolution and diversity of wild and cultivated potatoes',

Nature, 606(7914), pp. 535–541. Available at: https://doi.org/10.1038/s41586-022-04822-x.

Toda, N. *et al.* (2020) 'NLGenomeSweeper: A Tool for Genome-Wide NBS-LRR Resistance

Gene Identification', *Genes*, 11(3), p. 333. Available at:

https://doi.org/10.3390/genes11030333.

Van de Weyer, A.-L. *et al.* (2019) 'A Species-Wide Inventory of NLR Genes and Alleles in

Arabidopsis thaliana', *Cell*, 178(5), pp. 1260-1272.e14. Available at:

https://doi.org/10.1016/j.cell.2019.07.038.

Xie, J. *et al.* (2020) 'A rare single nucleotide variant in Pm5e confers powdery mildew

resistance in common wheat', *New Phytologist*, 228(3), pp. 1011–1026. Available at:

https://doi.org/10.1111/nph.16762.

578