Hosts Manipulate Lifestyle Switch and Pathogenicity Heterogeneity of Opportunistic Pathogens in the Single-cell Resolution

Ziguang Wang,[1,2,6] Shuai Li,[3,6] Sheng Zhang,[1] Tianyu Zhang,[4] Yujie Wu,[1] Anqi Liu,[1] Kui Wang,[1] Xiaowen Ji,[1] Haiqun Cao,[1] Eng-King Tan,[5] Yongcheng Wang,[4,*] Yirong Wang,[3,*] Wei Liu,[1,*]

[1]School of Plant Protection, Anhui Agricultural University; Anhui Province Key Laboratory of Crop Integrated Pest Management, Hefei, China

[2]First Clinical Medical College, Mudanjiang Medical College, Mudanjiang, China

[3]Bioinformatics Center, College of Biology, Hunan University, Changsha, China

[4]Liangzhu Laboratory, Zhejiang University, Hangzhou, China.

[5]Department of Neurology, National Neuroscience Institute, Singapore General Hospital Campus, Singapore.

[6]These authors contributed equally
[*]Corresponding author: yongcheng@zju.edu.cn (Y.W.), wangyirong@hnu.edu.cn (Y.W.), liuwei5@ahau.edu.cn (W.L.).

Abstract Host-microbe interactions are virtually bidirectional, but how the host affects their symbionts is poorly understood. Here, we report that the host is a critical modulator to regulate the lifestyle switch and pathogenicity heterogeneity of the opportunistic pathogens *Serratia marcescens* utilizing the *Drosophila* and bacterium model system. First, we find that *Drosophila* larvae efficiently outcompete *S. marcescens* and typically drive a bacterial switch from pathogenicity to commensalism toward the fly. Furthermore, *Drosophila* larvae reshape the transcriptomic and metabolic profiles of *S. marcescens* characterized by a lifestyle switch. More important, the host alters pathogenicity and heterogeneity of *S. marcescens* in the single-cell resolution. Finally, we find that larvae-derived AMPs are required to recapitulate the response of *S. marcescens* to larvae. Altogether, our findings provide an insight into the pivotal roles of the host in harnessing the life history and heterogeneity of symbiotic bacterial cells, advancing knowledge of the reciprocal relationships between the host and microbiome.

Introduction

All metazoans ranging from insects to humans harbor a plethora of microbes referred to as the microbiome. The microbiome plays a pivotal role in host physiology and pathophysiology, with some species conferring benefits to the host and others causing damage (Delannoy-Bruno et al., 2021; Lynch and Hsiao, 2019; Morais et al., 2021). Over the past decades, tremendous effort, including ours (Jia et al., 2021; Liu et al., 2022; Liu et al., 2017), has been devoted to understanding the impact of microbial strains or more complex communities on their hosts. In fact, interactions between the host and microbiome are mutually bidirectional (Backhed et al., 2005; Jiang et al., 2019), conferring benefits to the host and microbial sides. Nevertheless, the knowledge of the effect of the host on the resident microbial community is still in its infancy. Studies reveal that hosts play a crucial role in shaping the assembly and composition of their unique microbiota (Muegge et al., 2011; Olm et al., 2022; Valeri and Endres, 2021). However, environmental fluctuations frequently happen at much shorter time scales that preclude bacterial adaptation by genetic mutation or species displacement. To cope with these situations, microbial communities have developed sophisticated transcriptional reprogramming to globally regulate gene expression (Avraham et al., 2015; Becattini et al., 2021; Prescott and Decho, 2020). Pathobionts routinely sense and specialize on host-derived substrates, execute context-dependent transitions from harmful to commensal states and generate the host-association continuum (Barak-Gavish et al., 2023; Proctor et al., 2023; Somvanshi et al., 2012). In this regard, the molecular mechanism by which the host restrictively controls gene transcription and metabolism of their symbionts remains much undefined.

Microbial populations frequently aggregate in several dozen micrometres and stochastically undergo specific differentiation into subpopulations with strikingly distinct properties, employing a strategy known as bet-hedging to survive rapid changes in the environment (Eldar and Elowitz, 2010; Raj and van Oudenaarden, 2008). As such, bacteria manifest heterogeneous transcriptional profiles, leading to phenotypic heterogeneity among individual bacterial cells (Dar et al., 2021; Gasch et al., 2017). Traditionally, gene expression of bacterial cells has been investigated in bulk or on a population level by mixing and averaging mRNA simultaneously from sorts of cells. Single-cell transcriptomics is revolutionizing the analysis of phenotypic cell-to-cell variation in eukaryotes, enabling us to explore the heterogeneity heretofore hidden within population behavior (Klein et al., 2015; Mancio-Silva et al., 2022; Perez et al., 2022). Recently,

droplet-based high-throughput bacterial single-cell RNA sequencing methods have been developed and applied to study antibiotic-associated heterogeneous cellular states(Ma et al., 2023; Xu et al., 2023). We took advantage of this approach to tackle the challenge that the host induces transcriptionally distinct subpopulations of target bacteria during the interaction, contributing to a full appreciation of the phenotypic heterogeneity within symbionts.

Drosophila has a relatively simple bacterial community of commensal and opportunistic pathogens, typically predominated by 5 to 20 bacterial species. All *Drosophila* symbionts are facultative bacteria that alternate between free-living and host-associated lifestyles. *Serratia marcescens* is an opportunistic pathogen and generates a pink pigment prodigiosin that is characteristically oscillatory to metabolic activities, making it a visible bioindicator of *S. marcescens* metabolism. In this regard, *Drosophila* has provided a promising model to study symbiosis, dysbiosis (Chandler et al., 2011; Kim et al., 2020).

To address these issues, we used a reductionist approach in which germ-free (GF) *Drosophila* was re-associated with *S. marcescens* (Kuraishi et al., 2011). *S. marcescens* generates a pink pigment prodigiosin that is characteristically oscillatory to metabolic activities, making it a visible bioindicator of *S. marcescens* metabolism. We found that *Drosophila* larvae sufficiently outcompeted *S. marcescens*, and resulted in shifts in the transcriptomic and metabolomic profile in bulk and single-cell resolution, providing a robust paradigm to further study the host-microbe interaction with the *Drosophila* model.

Results

***Drosophila* larvae alter surface topography and population size of symbionts**

All *Drosophila* symbionts are facultative bacteria that alternate between free-living and host-associated lifestyles. Free-living bacteria initially form small microcolonies extending from the substratum, continue to expand, coalesce, and eventually generate a surface slick (a biofilm-like cell mat) that is visible to the naked (Koo and Yamada, 2016). Fascinatingly, the topography of surface slick was typically altered by fly colonization. It was grey and partially segmented by weak flies (Figure 1C), and even pale and intact associated with infertile flies (Figure 1D). However, the surface slick associated with strong wild-type flies was corn-like yellow, and the mature layer of the bacterial community was completely broken (Figure 1A). The destruction process was further exacerbated by overcrowded larvae that liquefied the upper food layer,

leading to a yellow aqueous layer (Figure 1B). These results indicate that *Drosophila* plays a

critical role in altering the topography of bacterial community on *Drosophila* media (Deines et

al., 2020). Our data showed that the cultivable bacterial loads associated with weak or infertile

flies were dramatically higher than those of strong stocks (Figure 1E), indicating that the host

diminishes the bacterial loading of the shared niche.

Given that *Drosophila* larvae and adults coexist in the bottles, we sought to determine

which were directly responsible for these alterations. Similarly, bacteria loadings were

substantially higher in the medium associated with males and virgin females (without progenies)

than in the control (Figure 1F), suggesting that larvae mainly cause a decline in bacterial

population size. Consistently, the 50-day-aged flies that deposited few eggs mimicked weak flies

with grey surface slick and a decline in bacterial loading. Moreover, transplanted larvae directly

caused a dose-dependent decline in the overall bacterial loading in the habitat (Figure 1G). These

findings demonstrate that *Drosophila* larvae play a critical role in outcompeting their symbionts

in the shared habitat.

***Drosophila* larvae antagonize *S. marcescens* in the niche**

To experimentally characterize the impact of the host on the symbiont, we applied a reductionist

approach in which *Drosophila* mono-associated with *S. marcescens* was generated as depicted in

Figure 2A. Crawling larvae (~96 h post oviposition) were concurrently transferred to fly food

vials with 10^7 CFU bacterial load. We found that *S. marcescens* alone formed a pink surface

slick in the medium over time (Figure 2B, top). The color intensity of surface slicks was initially

accumulated, peaked at 24 h timepoint post-inoculation, but gradually faded thereafter. We

quantified the optical density of prodigiosin inside surface slicks with the spectrophotometer as

described (Kalivoda et al., 2010; Pan et al., 2020). The time-course amount of prodigiosin was

congruent with the color density visible to the naked eye above (Figure 2C). Compared to *S.*

marcescens alone, the color intensity of surface slicks was significantly dampened by larval

transplantations (Figure 2B, bottom). As expected, the amount of prodigiosin in co-culture

substantially declined compared with that of *S. marcescens* alone. Next, we examined bacterial

titers in the food at regular intervals. Our result showed that single *S. marcescens* grew more

rapidly than coexisting *S. marcescens*, and the population of single *S. marcescens* reached the

plateau value within 1 d (Figure 2D), indicating that the niche possesses a finite carrying

capacity of bacteria. However, the population of coexisting *S. marcescens* reached the plateau

value within 2 d. Moreover, the number of CFUs was predominantly suppressed by larva

transplantation at the initial phases of colonization (within 72 h post-inoculation) compared to

that of *S. marcescens* alone (Figure 2D), consistent with the result that larvae thwarted the

number of total mixed bacteria in the diet (Figure 2B). Noteworthily, the number of CFUs of *S.*

marcescens alone was lower than *S. marcescens* in co-cultures at the late stage (at 96 h post-

inoculation), likely that bacteria rapidly exhausted their nutritional resources and underwent

ecological suicide (Ratzke et al., 2018). These results suggest that the larva acts as a potential

competitor that efficiently prevents the outgrowth of *S. marcescens* in the habitat. To verify it,

different numbers of larvae were added to vials inoculated with *S. marcescens*. Indeed, the more

were larvae, the less the colour intensity of the surface slick (Figure 2-figure supplement 1A, B).

In addition, the population size of *S. marcescens* in food was in restricted control by larvae in a

dosage-dependent manner (Figure 2-figure supplement 1C). Altogether, the cumulating results

suggested that *Drosophila* larva has a competitive advantage in the habitat, and acts as a critical

regulator of their symbionts.

Drosophila larvae modulate the pathogen-to-commensal transition of *S. marcescens*

S. marcescens is a *Drosophila* pathobiont with the potential to switch between commensalism

and pathogenicity toward the host. We set out to examine the *S. marcescens* lifestyle switch from

pathogenicity to commensalism by assessing the survival of flies challenged with fly medium

processed single or coexisting *S. marcescens*, respectively. Our data showed that flies challenged

with *S. marcescens* alone manifested higher mortality than flies with *S. marcescens* in co-culture

(Figure 2E), suggesting that larva antagonizes the pathogenicity of their symbionts. In addition,

the expression of virulent factors was thwarted by larvae (Figure 2F). However, *S. marcescens*

more efficiently sustained optimal larval development upon nutrient scarcity than both axenicity

and commensal *Lactobacillus plantarum* (Matos et al., 2017; Storelli et al., 2011) (Figure 2-

figure supplement 1D), indicating that larvae-associated *S. marcescens* could impart fitness

benefits to their hosts. The cell wall of Gram-negative bacteria is composed of a single

layer of peptidoglycan surrounded by a membranous structure, which may function as a

virulence factor (Pan et al., 2022). Indeed, transmission electron microscopy confirmed a

significant reduction in the width of the cell wall of *S. marcescens* in co-culture compared to that

of *S. marcescens* alone (Figure 2G). Additionally, we observed that the expression of genes

related to cell wall biosynthesis was dampened by larva transplantation (Figure 2H). Taken

together, these results suggest that *Drosophila* larvae modulate *S. marcescens* lifestyle from
pathogenicity to commensalism toward the host.

Presumably, surface slicks can be destroyed by mechanical force from larva crawling and
burrowing (Dufrene and Persat, 2020). To tackle this issue, we agitated fly food to imitate larva
locomotion (Figure 2-figure supplement 2A). Indeed, agitation decreased colour intensity,
prodigiosin yield and population size in fly food (Figure 2-figure supplement 2B-D), implying
that mechanical force accounts for the negative regulation of *S. marcescens*. However, larval
transplantation resulted in a much more robust decline in colour intensity of surface slicks,
prodigiosin yield and population size than agitation alone. Of note, the surface of the slick with
agitation appeared lighter than that of larvae, possibly due to a stratification of prodigiosin
following agitation. In addition, the expression of prodigiosin synthesis genes was thwarted by
the agitation (Figure 2-figure supplement 2E). These results indicated that larva-derived
biofactors and/or synergism of force and biofactors primarily confer the inhibition of *S.*
marcescens (see the later results).

***Drosophila* enforces bacterial global transcriptional and metabolic adaptation to the host**
In order to understand the molecular basis for the bacterial lifestyle switch from pathogenicity to
commensalism in response to *Drosophila* larvae, bulk RNA-seq analysis was applied to bacterial
cells 24 h after inoculation. We devised an approach to efficiently collect bacteria from the agar
fly food as described in Methods. As shown in Figure 3A, Principal Component Analysis (PCA)
showed a larvae-dependent separation to *S. marcescens* alone on the first principal component of
variance. Compared to single *S. marcescens*, larvae-associated *S. marcescens* exhibited
significant upregulation of 360 genes and downregulation of 439 genes (Figure 3B and
Supplementary table 2), respectively. Functional annotations of the differentially expressed
genes (DEGs) were assigned using the Kyoto Encyclopedia of Genes and Genomes (KEGG).
KEGG pathway enrichment analysis highlighted that the most differentially upregulated DEGs
of coexisting *S. marcescens* were related to bacterial proliferation and growth, including
ribosome, translation factors, transcription, DNA replication proteins, and energy metabolism
(Figure 3D), implying that larval transplantations favored long-term maintenance of *S.*
marcescens in the diet. By contrast, the most differentially downregulated DEGs of single *S.*
marcescens were related to bacterial pathogenicity, including transporters, excretion, quorum
sensing, and exosome (Figure 3C). To validate our findings in bulk RNA-seq analysis,

quantitative PCR was used to examine the expression of predicted genes involved in
bacterial proliferation and pathogenicity. Indeed, most predicted genes associated with
pathogenicity were downregulated in single *S. marcescens* (Figure 3E), while genes associated
with bacterial proliferation were upregulated in response to larva colonization (Figure 3F).
Overall, *Drosophila* forms a long-term symbiosis with the bacterial community by harnessing
bacterial global transcription.

Next, we reasoned that the host could further elicit changes in bacteria metabolic networks
using untargeted metabolomics. The data showed that 91 metabolites were identified and their
concentrations were quantified (Supplementary table 3). Metabolomic profiles discriminated
larva-associated *S. marcescens* from *S. marcescens* alone (Figure 4A), indicating that the host
reshapes the global metabolic profile of bacterial cells. We found that larvae-associated *S.*
marcescens displayed significant upregulation of 22 metabolites and downregulation of 69
metabolites (Figure 4B). Unsupervised hierarchical clustering of the metabolome revealed
distinct clusters of metabolites in *S. marcescens* alone versus co-culture (Figure 4C). Moreover,
we detected a significant decrease in amino acid metabolism, phosphotransferase system, and
ABC transporters in *S. marcescens* in co-culture compared to *S. marcescens* alone (Figure 4D),
suggesting that the host suppresses differentiation of *S. marcescens* into the population with
pathogenicity. By contrast, the most differentially upregulated metabolites of *S. marcescens* in
co-culture were related to the biosynthesis of fatty acids and unsaturated fatty acids, and the
pentose phosphate pathway (Figure 3D). Taken together, these results suggest that the host
affects the global metabolic profile of symbiotic cells and drives the switch of the bacterial
lifestyle from pathogenicity to commensalism.

The co-expression network analysis of transcriptome and metabolome revealed that 42
differentially expressed metabolites were found to be related to the pathogen-to-commensal
switch, including ribosome, transcription, DNA replication, energy metabolism, ABC
transporters, phosphotransferase system, quorum sensing and exosome (Figure 4-figure
supplement 3A). Serotonin, sorbitol and lactobionic acid were related to the pathogen-to-
commensal switch (p value<0.05). To validate these results, we perturbed the *S. marcescens*
extracellular environment by adding predicted metabolites to the fly food. Indeed, we found that
serotonin, sorbitol and lactobionic acid efficiently reduced the prodigiosin yield and CFUs
(Figure 4-figure supplement 3B, C), highlighting the high quality of the prediction. Moreover, *S.*

marcescens perturbed with sorbitol manifested less virulent to flies (Figure 4-figure supplement

3D) and the impaired transcription of virulence-related genes (Figure 4-figure supplement 3E),

resembling the lifestyle of co-culturing *S. marcescens.* Collectively, these findings suggest that

Drosophila enforces bacterial transcriptional and metabolic adaptation to the host.

Larvae modulate pathogenicity heterogeneity of *S. marcescens*

Isogenic microbial populations can generate transcriptional variation across individual cells,

thereby prompting us to analyze bacterial populations at the single-cell resolution. We

characterized the transcriptome of individual *S. marcescens* cells by implementing bacterial

single-cell RNA sequencing on a platform available. To validate this approach, we first

determined the capacity of this platform to distinguish bacterial populations of heat-shocked *S.*

marcescens grown in a liquid medium as previously described (Dar et al., 2021). Indeed, the

reults provide strong evidence that the bacterial single-cell RNA sequencing approach used was

sufficient and reliable to capture transcriptional responses to heat shock (Figure 5-figure

supplement 4).

Next, we speculated about the heterogeneous transcriptional response of *S. marcescens*

challenged with larvae as well as mechanical force. The growth of *S. marcescens* was well

understood in fly food conditions as described in Figure 1, so we collected *S. marcescens* at the

late exponential phase when certain bacterial cells manifested their differentiation and

pathogenicity. We captured 2800 cells in total and detected a median of 198 genes per cell for

control, 333 for force, and 175 for larvae (Figure 5A). Strikingly, bacterial cells from each group

tended to form three distinct clusters that correspond to the agitated, larva-associated and control

cultures could be visualized by graph-based clustering of their gene expression profiles (Figure

5B). This result suggested that larval transplantation, as well as agitation, induce significant

changes in gene expression patterns, consistent with the bulk RNA-seq result that larvae caused

the shift in bacterial global transcription using Seurat (Kuchina et al., 2021). Next, we attempted

to partition the total bacterial population of *S. marcescens* into subpopulations with diverse

predicted functional capabilities by the algorithmic grouping of cell-expression profiles. As

shown in Figure 5C, we further partitioned the three groups into 7 subclusters (Subcluster 0-6)

representing subpopulations with distinct expression profiles, while the gene expression in each

cluster was continuous. Namely, the sampled populations of *S. marcescens* alone and in

coculture were respectively partitioned into three coexisting subgroups (Subcluster 0,1, and 2 for

co-culturing *S. marcescens* and Subcluster 3, 4, and 6 for single *S. marcescens*), implicating that

phenotypic diversity is a general feature of bacteria in clonal populations.

To understand the heterogeneity of gene expression patterns, we identified the DEGs

associated with pathogenicity to characterize the transcriptional profile (Supplementary table 4),

and ranked marker genes in each cluster (Figure 5D). We selected pathogenicity-related DEGs,

including *livI* (ABC transporter), *oppA* (quorum sensing), *secY* (secretion system), and *fp*

(virulence), and charted the feature expression of them in each cluster in low-dimensional space

(Figure 5E, F and I, J). We found most genes involved in pathogenicity displayed a substantial

decline upon larvae, which was consistent with the previous findings of reduced pathogenicity of

coexisting *S. marcescens*. Moreover, we found that *livI* exhibited gradually higher expression

down along Subcluster 3 to Subcluster 6 (Figure 5G), implicating the potentially hierarchical

virulence-regulatory network in these three subclusters. Analogously, *oppA*, *secY* and *fp*

exhibited a similar expression pattern in these three subclusters (Figure 5H, K and L). Given that

increased pathogenicity along three subclusters. It was likely that Subcluster 6 could be

developed from Subcluster 4, thereby inferring that Subcluster 6 could be identified as

subpopulations of single *S. marcescens* in a bona fide pathogen. Intriguingly, many virulence-

associated genes were highly expressed in Subclusters 5, implying that mechanical force could

not attenuate the virulence of *S. marcescens*. We tested it by assessing the survival of flies

challenged with processed fly medium from agitated *S. marcescens*. Indeed, flies treated with

agitated *S. marcescens* displayed a significantly shorter lifetime than flies with coculturing *S.*

marcescens (Figure 5-figure supplement 5A). Taken together, our results demonstrated the

presence of pathogenicity heterogeneity of *S. marcescens* at the single-microbe-level

transcriptional landscape.

Larvae modulate growth heterogeneity of *S. marcescens*

We showed that coexisting *S. marcescens* were related to growth, so we continued to identify the

DEGs associated with bacterial propagation to characterize the transcriptional profile

(Supplementary table 4), and ranked marker genes in each cluster (Figure 6A). Among DEGs of

ribosome between single *S. marcescens* and co-culturing *S. marcescens*, *rpsL* encodes the highly

conserved rps12 protein of the ribosomal accuracy center, while, tryptophan, biosynthesized by

trpD among DEGs of DNA replication, is an essential nutrient and serves as a building block for

protein synthesis and DNA replication. As anticipated, most genes involved in growth showed an

evident increase in larvae. Interestingly, *rpsL* and *trpD* similarly exhibited gradually higher

expression up along Subcluster 0 to Subcluster 1 (Figure 6B), indicating the potentially

hierarchical growth-regulatory network in these three subclusters. Taken together, our results

demonstrated the presence of bacterial propagation heterogeneity of *S. marcescens* at the single-

microbe-level transcriptional landscape.

Nitrogen is the fundamental element of nucleic acids and proteins. In the clusters

corresponding to larvae (clusters 0, 1 and 2), we observe peak expression of genes involved in

nitrogen metabolism, such as P-II family nitrogen regulator (*glnK*) and glutamate-ammonia

ligase (glnA; Figure 6-figure supplement 5B, D). *glnK* plays a critical role in regulating the

activity of glutamine synthetase (e.g. *glnA*), which promotes glutamine synthesis. Glutamine is

the most abundant non-essential amino acid, and serves as nitrogen and carbon sources for cell

growth and differentiation. Moreover, these three subclusters showed a significant increase in the

activity of genes involved in urea transport and utilization, such as urea ABC transporter

substrate-binding protein (*urtA*), allophanate hydrolase (*atzF*) and urea carboxylase (*uca*). *urtA*

accounts for the transport of urea, and then *atzF* and *uca* synergize to convert urea to ammonia

that can be used for glutamine biosynthesis. The regulatory schematic is shown in Figure 6J.

Additionally, glutamine synthesis needs a large amount of energy, so we then turned to carbon

metabolism. Indeed, we observed higher expression of phosphoglycerate kinase (*pgk*) and

bifunctional acetaldehyde-CoA/alcohol dehydrogenase (*adhE*) that participate in glycosis in the

subclusters 0, 1 and 2 (Figure 6G and Figure 6-figure supplement 5C). Moreover, we found that

pgk and *adhE* exhibited gradually higher expression in these three subclusters relative to

subclusters with single *S. marcescens* (Figure 6I and Figure 6-figure supplement 5E), suggesting

that bacteria could establish a dedicated replicative niche to efficiently replicate. Altogether,

these findings that the host globally results in transcriptional reprogramming of single-cell *S.*

marcescens, facilitating the cooperation among individual cells.

Larvae-derived AMPs efficiently antagonize *S. marcescens*

Our results showed that the host keeps strict control over the pathobiont, we next asked whether

one or multiple compound(s) secreted by the host could recapitulate this phenomenon. Indeed,

the data showed that diets with intestinal excreta exhibited a modest but significant decline in

color intensity of the slick, prodigiosin yield and population size compared to control (Figure

7A-C), in line with the result above (Figure 2-figure supplement 2C, D). Larval excreta contains

short antimicrobial peptides (AMPs) that efficiently resist a variety of pathogens (Marra et al.,

2021), prompting us to whether AMPs could recapitulate the response of *S. marcescens* to the

presence of *Drosophila* larvae. To refine this concept, AMPs were supplemented to fly food vials

with 10^7 CFU *S. marcescens* at the same time. We found that AMPs suppressed color intensity of

the slick, prodigiosin yield and population size in a dosage-dependent manner, indicating that

AMPs, to a lesser extent, played a role in monitoring *S. marcescens*. To verify it, we turned to a

compound mutant strain lacking *Defensin*, *Cecropins* (4 genes), *Drosocin*, *Diptericins* (2 genes),

Attacins (4 genes), *Metchnikowin*, and *Drosomycin*, referred to as "*ΔAMP*"(Hanson et al., 2019),

allowing direct investigation of their role in the lifestyle transition of the pathobiont. Indeed,

ΔAMP recapitulated an increment in color intensity of surface slick, prodigiosin yield and

population size compared to wild-type counterparts (Figure 7D-F). Moreover, the recapitulation

in color intensity of slick, prodigiosin yield and population size was substantially attenuated and

even abolished by the addition of AMPs, suggesting that much of the inhibition of the

microbiome can be ascribed to AMPs. Subsequently, quantitative PCR was employed to

confirm the expression of altered genes as depicted above (Figure 7G, H). Together, our

data demonstrated that larva-derived AMPs were a key factor that accounts for the restrictive

control over diet microbes.

Discussion

The important role of the host in affecting microbial physiology and behavior is only beginning

to be appreciated. In the current study, we found that *Drosophila* larvae act as a competitive

regulator that prevents *S. marcescens* outgrowth and antagonizes its pathogenicity. *Drosophila*

larvae monitored the transcriptomic and metabolomic profile of *S. marcescens*, characterized by

the lifestyle switch from mutualism to pathogenicity toward the fly. More importantly, we

highlight that the host alters the single-cell transcriptomic atlas of *S. marcescens* and phenotypic

heterogeneity in bacterial populations.

Drosophila adults are routinely attracted to oviposit their eggs on rotting fruits that possess

both commensal and pathogenic microbes (Liu et al., 2017; Wong et al., 2017). Extensive

attention has been dedicated to the essential roles of symbiotic bacteria in modulating the

physiology of their animal partner, or to the molecular mechanisms underlying physiological

benefits to their host. However, it's necessary to explore both sides of symbioses to obtain a

more complete understanding of the host-bacteria interaction. *S. marcescens* encounters a cost
associated with symbiosis, as the population size for it is diminished in the shared niche by the
larvae (Figure 2). Our findings highlight that larvae alleviate intraspecies competition of *S.*
marcescens through population size control at the initial stage, thereby promoting the fitness of
symbionts in the long maintenance. Taking into account these findings, the competition model is
more plausible to the larvae-pathogen system where larvae efficiently prevent the outgrowth of
the potential pathogenic bacterial community (Burns et al., 2017; Foster et al., 2017).
Interspecies compete for nearly the same or similar nutrients and space for their growth and
maintenance in the habitat, making *Drosophila* larvae a potential competitor for *S. marcescens*.
Intriguingly, a previous study reported the contrasting result that *Drosophila* farms its
commensal *L. plantarum*, and is required to optimize the extraction of dietary nutrients and
sustain the growth of their symbionts upon chronic undernutrition (Storelli et al., 2018). To
reconcile the two contradictions, another model of interspecies cooperation is suggested.
Consequently, larval transplantations favored long-term maintenance of *L. plantarum* on the diet,
and benefits from symbiosis override the costs of the initial competition in the long run,
conferring the beneficial effect of larval presence on bacterial maintenance. This paradox
possibly reflects a strategy developed by *Drosophila* to preserve its own fitness. Potential
pathogens differ from commensal bacteria, because they generate secondary metabolites
threatening insect life at the static stage of growth (Figure 2E). If the larvae obtain benefits to
promote their development and growth, they must exert strict control over them through
antimicrobial peptides and reactive oxygen species (Sharp and Foster, 2022). We found that
larvae-derived AMPs efficiently antagonized *S. marcescens* (Figure 7), thereby recapitulating the
response of *S. marcescens* to *Drosophila* larvae. Yet, the full complement of larva-derived
factors required for bacterial controls as well as their mode of functions remains elusive, paving
the way to seek the evolutionary-conserved animal factors essential to maintain symbiosis.
Taken together, our findings of the host on microbial communities would improve our
understanding of the ecology of host-symbiont interactions.
Owing to their saprophagous foraging behavior, *Drosophila* has to cope with many
potential pathogens in the environment (Black et al., 2018). We're appreciating the profound
influence of the host on the resident microbial community, but the underlying molecular
mechanisms by which the host is potentially involved in the perpetuation of host-bacteria

symbiosis are still poorly understood. Our result showed that *S. marcescens* versatilely displayed
transcriptional and metabolomic adaptations to *Drosophila* larvae (Figure 3 and Figure 4),
avoiding being competitors when they are associated with the host. Indeed, *S. marcescens*
rapidly reached the plateau and exhausted its nutritional resources, and then generated secondary
metabolites that could endanger insect life (Figure 2). Consistently, our transcriptome data
suggest the pathogen-to-commensal transition of *S. marcescens* by the presence of larvae,
including ABC transporters, phosphotransferase system, quorum sensing and exosome. In gram-
negative bacteria, these exporters transport lipids and polysaccharides from the cytoplasm to the
periplasm, or certain substances that need to be extruded from the cell, including surface
components of the bacterial cell (e.g., capsular polysaccharides, lipopolysaccharides, and
teichoic acid), proteins involved in bacterial pathogenesis (e.g., hemolysis), heme, hydrolytic
enzymes, competence factors, toxins, bacteriocins, peptide antibiotics, and siderophores. In
addition, quorum sensing, used by pathogens in disease and infection, regulates gene expression
following population density through autoinducers, allowing bacteria populations to
communicate and coordinate group behavior (Mukherjee and Bassler, 2019). The findings
suggest that the microbiota responds to host physiology by altering gene expression and
metabolism (Penterman et al., 2014). Consistently, a study observed that commensal bacteria
calibrate their transcriptional and metabolic output to different systemic inflammatory responses
(Fyhrquist et al., 2019). Taken together, these findings suggest that larvae elicit a set of
expressed genes in their harmful symbionts, favoring the shift from a pathogenic to a commensal
stage.

Most microbial communities consist of a genetically diverse assembly of different
organisms, and the level of genetic diversity plays an important part in community properties and
functions (Davies et al., 2022). However, genetically identical microbial cells can show different
behaviors, including differences in growth speed, gene expression and metabolism, and
biological diversity can arise at a lower level of biological units (Ackermann, 2015; Avraham et
al., 2015). Fortunately, the recent development of methods to interrogate populations on the
single-cell level has dramatically altered our understanding of cellular heterogeneity by
providing much greater resolution of different cell types and cell states (Hare et al., 2021; Imdahl
et al., 2020; McNulty et al., 2023). By applying this technique, we indeed observed that single
cells differ from each other with respect to gene expression even when genetic and

environmental differences between cells are reduced as much as possible in a shaking liquid
medium (Figure 5-figure supplement 4). Recent development in single-cell techniques facilitates
to reveal that distinct bacterial subpopulations contribute unique colonization and growth
strategies to infection sites (Llorens-Rico et al., 2022). For example, the host can drive
Salmonella phenotypic heterogeneity at the single-cell level throughout the course of infection,
highlighting how variation in gene expression, and metabolic activity contribute to overall
bacterial success (Tsai and Coombes, 2019). Consistently, we found that the host changes the
transcriptomic atlas of *S. marcescens* individual cells, and attenuates phenotypic heterogeneity of
virulent factors (Figure 5). To date, transcriptional approaches-profiling the host, the pathogen,
or both-have been employed to uncover substantial molecular details about the host and bacterial
factors that underlie infection outcomes. During the interaction, the niche environment
diversifies into 3D areas with varying degrees of growth conditions (Koo and Yamada, 2016). In
a single population, both in vitro and in vivo, *S. typhimurium* has been shown to display
significant cell-to-cell variation in attributes such as growth rate, expression of virulence factors,
and sensitivity to antibiotics (Claudi et al., 2014). With the development of this approach, it is
possible to identify and modify the transcription of certain target bacterial cells with the
expression of virulence factors in order to selectively treat human diseases in the gut
microenvironment.

Utilizing the *Drosophila* model system, we revealed a natural ecological phenomenon
whereby the host was a prerequisite for regulating the population size and lifecycle switch of
indigenous bacteria. It's of importance to understand the ecological and evolutionary processes
that shape host-associated microbial communities. Little is known about the effect of the host on
the commensal bacteria, and it would be also interesting to investigate whether the host and
commensal bacteria could synergize to antagonize the pathogenicity of potential pathogens.
Future studies that evaluate the molecular mechanism underlying the effect of the host on
microbial communities would improve our understanding of host-symbiont co-evolution in
nature.

Materials and methods
Fly culture and stocks
Drosophila flies were routinely reared and kept at the condition of 25°C, 55%-65% humidity with a 12 h: 12 h light-dark circle unless otherwise noted. The Canton-S strain was used as the wild-type fly in this work. Delta *14 AMP* mutant was generated as described (Hanson et al., 2019) and kindly gifted by Dr. Zhai in Hunan Normal University. The fly was raised on standard cornmeal–sugar–agar medium (1 liter): 105 g dextrose, 7.5 g agar, 26 g yeast, 50 g cornmeal, 0.25 g Sodium benzoate (Sigma Aldrich) dissolved in 8.5 ml 95% ethanol and 1.9 ml propionic acid (99%, Mallinckrodt Baker).

Generation of germ-free and gnotobiotic flies
Germ-free flies were generated as described (Jia et al., 2021). In brief, fresh embryos within 8 hours post egg-laying were collected from agar media with 1.5% grape juice, rinsed with ddH$_2$O, and transferred into 1.5 ml Eppendorf tubes. Diluted sanitizer Walch (1:30), 2.5% sodium hypochloride (Sigma Aldrich), 75% ethanol, and sterile PBS containing 0.01% TritonX-100 were successively applied to bleach embryos. The germ-free embryos were cultivated in autoclaved fly food with 10% yeast. Axenia of germ-free larvae was normally tested by plating the larval homogenates on nutrient agar plates (peptone 10 g/L; beef extract powder 3 g/L; NaCl 5 g/L; agar 15 g/L). Gnotobiotic flies were generated by inoculating bacterial strains to germ-free flies.

Survival assay
Newly eclosed female flies were collected and transferred into vials (15 flies/vial), and each group contained 6 vials with a total of 180 flies in each group based on two replicates for each. Before bacterial treatment, vials were pre-inoculated with *S. marcescens* alone and *S. marcescens* with 40 larvae, and incubated at 25°C for 24 h for culture. The flies were then transferred to vials processed with *S. marcescens* alone and *S. marcescens* in-coculture every 3 days. The number of dead flies was counted each day, and the proportion of surviving flies was calculated at each time point of the experiment. Experiments were independently replicated twice.

Bacteria culture and CFU counting
All the material to manipulate bacteria was sterilized before usage. The strains of *Serratia marcescens* with the Genbank accession number CP053378 and *Lactobacillus plantarum* with the Genbank accession number KY038178 were used. *S. marcescens* and *L. plantarum* were cultured in LB and MRS broth medium at 30°C, respectively. Bacteria cells were harvested by centrifugation (5000 rpm, 3 min), washed twice in 1×PBS, and resuspended in 1×PBS to obtain 10^8 cells/ml (OD$_{595}$ = 1). The bacterial suspension (20 μl) was supplemented to a vial with autoclaved fly food, and crawling GF larvae were then added to vials to generate a host-microbe interaction model. For the heat-shock experiment, the culture was transferred to a 45°C incubator at 12 h timepoint after postinoculation, and kept for 15 min. Heat-shocked bacteria were used for bacterial single-cell RNA seq below.

Bacterial load quantification
Fly food in vials was agitated and homogenated with 5 ml ddH$_2$O. Bacterial load was assessed by plating tenfold serial dilutions of the food homogenates on LB agar plates and incubating the plates at 30°C for 16 h. The numbers of CFUs were counted, and expressed as the total number of living bacteria per vial.

Prodigiosin production assay
The determination of the prodigiosin yield of bacteria was carried out with acidified ethanol and absorbance measurement as previously described (Kalivoda et al., 2010; Pan et al., 2020). The relative concentration of prodigiosin produced by solid-grown cultures was quantified as follows.

475   Samples were added to 1.2 ml acidified ethanol (4% 1 M HCl in ethanol) to extract prodigiosin
476   from the culture for 10 min. Cell debris and impurities were removed by centrifugation at 13000
477   rpm for 5 min, and the supernatant was transferred to a cuvette for measurement of absorbance at
478   534 nm. Prodigiosin production of samples was calculated as the optical density at 534 nm.
479   Experiments were independently replicated four times.

Transmission electron microscopy

Transmission electron microscopy (TEM) was conducted to observe the structures of the
bacterial cells as previously depicted (Morgelin, 2017). Briefly, bacterial cells were collected
after 24-h incubation and then fixed in 2.5% glutaraldehyde for 5 h. The cells were dehydrated in
a gradient series of ethanol solutions from 30 to 100% by incubation. For TEM analysis, the
samples were treated with acetone and embedded in epoxy resin. Thin sections (70 nm) were cut
with a diamond knife mounted on a Leica UC-7 ultramicrotome and collected on carbon-coated
Cu grids. TEM observation was performed with a JEOL JEM1400 TEM at 200 kV. Micrograph
films were digitally acquired at high resolution with EMSIS Morada G3 (Winey et al., 2014).

Real time-PCR analysis

RT-qPCR assay was performed as described previously (Liu et al., 2022). In brief, to assess the
expression levels of prodigiosin synthesis-related genes, extracellular polysaccharide production-
related genes, and partial down-regulated and up-regulated genes, the cultures of different groups
of *S. marcescens* were collected after 24 h incubation. The collected cells were then subjected to
total RNA extraction using a Bacteria RNA Extraction Kit (Vazyme, China). After treating the
total bacterial RNA with DNase I (Vazyme, China), the concentration and quantity of the total
bacterial RNA were determined using a NanoDrop spectrophotometer (Thermo Scientific), and
0.6 µg of the total bacterial RNA was subjected to reverse transcription using the HiScript III All-
in-one RT SuperMix Kit (Vazyme). The mixture was subjected to RT-qPCR analysis using the
ChamQ Universal SYBR qPCR master mix kit (Vazyme) in a CFX96™ Real-Time System
(BioRad, Hercules, CA, USA). RNA from three biological replicates were analyzed and four
technical replicates were performed. The relative expression values were calculated using the
following formula: $\triangle Ct=$ Ct (target gene) - Ct (reference gene), and the relative expression was
equal to $2^{-\triangle\triangle Ct}$. The 16S rRNA protein-encoding gene was used as an internal control. The
primers used for RT-qPCR analysis are listed in Supplementary table 1. Experiments were
independently replicated three times.

Bacterial bulk RNA Sequencing

To analyze the effect of larvae on the transcriptome in *S. marcescens*, the cultures of different
groups were collected after 24 h incubation. 2 ml ice-cold 1×PBS were added to the vial for 5
min to suspend bacteria cells, and then suspended bacteria cells were removed food remainder by
centrifugation (900 rpm, 3 min), washed twice in 1×PBS. Bacteria cells were harvested by
centrifugation (5000 rpm, 4 min), and washed twice in 1×PBS. The collected cells were then
subjected to total RNA extraction using a Bacteria RNA Extraction Kit (Vazyme, China). The
integrity, concentration and quantity of the bacterial RNA were determined using a NanoDrop
spectrophotometer (Thermo Scientific) and a 1% agarose gel. 1 µg total RNA with RIN value
above 7 was used for following strand-specific library construction using VAHTS Universal V8
RNA-seq Library Prep Kit for Illumina (Vazyme, China). Library quality was evaluated using an
Agilent 2100 Bioanalyzer. The sequences were sequenced and processed using Novaseq-PE150
by the company (Novogene, Beijing, China).

For annotation, the genome of *S. marcescens FY* was used as a reference. The significant
differentially expressed genes were determined in different groups using the DESeq software

(DESeq 2), with the standards of P-value ≤0.05, and fold change |log2Ratio|≥1.2. KEGG KAAS
database was used to annotate the genes with significantly differential expression based on their
functions using the BLAST.

Metabolomics analysis

Untargeted metabolomics was performed using a modified version of a previously reported
protocol(Tsugawa et al., 2015). In brief, L-2-chlorophenylalanine (0.06 mg/ml) dissolved in
methanol was taken as internal standard, the samples were porfomed to GC-MS analysis. QC
samples were prepared by mixing aliquots of all samples to be a pooled sample. Samples were
analyzed on an Agilent 7890B gas chromatography system coupled to an Agilent 5977A MSD
system (Agilent Technologies Inc., CA, USA). The temperature of the MS quadrupole and ion
source (electron impact) was set to 150℃ and 230℃, respectively. The collision energy was 70
eV. Mass spectrometric data were acquired in a full-scan mode (m/z50-500). For data processing,
MZmine 2 and MultiQuant software programs were used.

Bacterial single-cell RNA-Seq and analysis

Bacterial single-cell RNA sequencing was carried out on a commercially available platform
(M20 Genomics Company) (Xu et al., 2023). In brief, half a million *S. marcescens* cells were
collected by centrifuging at 4℃. The supernatant was removed and cell pellets were resuspended
in 2 ml ice-cold 4% formaldehyde (Sigma, 47608) and incubated with shaking overnight at 4℃.
The fixative was removed by centrifuging at 4000 rpm for 5 min at 4℃ and cells were washed
twice with 1 mL PBS-TRI (1x PBS supplemented with 0.1% Tween-20 and 0.1 U/mL Murine
RNase inhibitor (Vazyme, R301)). The supernatant was removed and cells were resuspended in
200 µl PBS-TRI (1x PBS supplemented with 0.1% Tween-20 and 0.2 U/mL Murine RNase
inhibitor (Vazyme, R 301)). The cell suspensions were counted with a Moxi cell counter and
diluted according to the manufacturer's instructions to obtain single-cell.

The bacterial single-cell RNA-Seq library was prepared according to the protocol of
VITAPilote® kit(M20 Genomics, R20114124). In situ reverse transcription of bacteria was
performed with random primers and the resulting cDNA fragment was added with adaptor. The
droplet barcoding for a single bacterium was performed on VITACruiser® Single Cell
Partitioning System(M20 Genomics, Hangzhou, China). Bacteria, DNA extension reaction mix,
and hydrogel barcoded beads were encapsulated using the VITACruiser. The aqueous phase
containing cDNAs was purified with magnetic beads. The cDNAs were amplified by PCR, and
purified with magnetic beads. All products were pooled to construct a standard sequencing
library. Sequencing was done on a PE150 (Illumina), and raw reads were aligned against the *S.*
marcescens FY genome and counted by StarSolo followed by secondary analysis in the
Annotated Data Format. Sequencing data was further analyzed using Seurat (v.4.3.0.1 with
default parameters except where indicated). Cells were filtered to retain only those with at least
100 genes. Genes were also screened to remove genes expressed only in fewer than five cells.
Then, We first log-transformed the data using the NormalizeData function, and selected the 2000
most variable genes using FindVariableFeatures. Then, we z-scored these highly-variable genes
using ʹScaleDataʹ. Next, we performed linear dimensionality reduction using principal
component analysis (PCA) down to 50 dimensions (ʹRunPCAʹ). Points in this embedding were
used to construct UMAP plots and find neighbours for clustering. FindClusters was used to run
the Louvain clustering algorithm and generate clusters. We confirmed that the clusters
highlighted in the main text appeared consistently for a range of resolutions from 0.5 to 1.5.
DEG was performed using the FindMarkers function with a log-fold change cutoff of 0.25.

Statistical analysis

Statistical analysis is performed using GraphPad Prism 9.0 and indicated inside each figure legend. The layout of all figures used Adobe Illustrator 2022. Experimental flies were tested at the same condition, and all data are collected from at least two independent experiments. D'Agostino–Pearson normality test was used to verify the normal distribution of data. If normally distributed, a two-tailed Student's t-test was used to compare two groups, and one-way ANOVA followed by Tukey's multiple comparisons test was used for comparisons of multiple groups. If not normally distributed, a two-tailed Mann-Whitney U-test was performed to compare two groups of samples, while Kruskal–Wallis test followed by Dunn's multiple comparisons test was used for multiple comparisons among three or more groups.

Acknowledgements

The authors would like to thank all members of Dr. Liu's and Dr. Wang's labs for discussion. We appreciate S. Wang, G. Wang, Y. Pan, Z. Zhai and Y. Zhang for their critical comments on the manuscript. We thank the Bloomington Stock Center for providing fly stocks, and Dr. Zongzhao Zhai for fly stocks. This work was supported by the National Natural Science Foundation of China (31501175), Natural Science Foundation of Anhui Province (2308085MC74) and Talents in Anhui Agricultural University (RC342201) to W.L., and the Ministry of Science and Technology of the People's Republic of China (2022YFE0132000), the Natural Science Foundation of Hunan Province (2022JJ40048), and the Fundamental Research Funds for the Central Universities of China (531118010546) to Y.W.

Additional information

Funding

Funder	Grant reference number	Author
National Natural Science Foundation of China	31501175	Wei Liu
Natural Science Foundation of Anhui Province	2308085MC74	Wei Liu
Talents in Anhui Agricultural University	RC342201	Wei Liu
Ministry of Science and Technology of the People's Republic of China	2022YFE0132000	Yirong Wang
Natural Science Foundation of Hunan Province	2022JJ40048	Yirong Wang
Fundamental Research Funds for the Central Universities of China	531118010546	Yirong Wang
The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.		

Author contributions

Z.W., H.C., E.T., Y.W., Y.W. and W.L. conceptualized the study, designed experiments, and interpreted all the data. Z.W. performed bacterial counting, survival assay, bull RNA-seq runs, TEM, metabolomics runs, bacterial single-cell RNA-seq and RT-PCR, with help from S.Z., Y.W. and A.L.; Y.W. contributed to bacterial single-cell RNA-seq. S.L. and S.Z. wrote codes and analyzed data of bull RNA-seq, metabolomics, and bacterial single-cell RNA-seq. Z.W., Y.W.

597 and W.L. wrote the manuscript with input from all authors. Y.W., Y.W. and W.L. conceived and
598 supervised the project.
599

Data availability.
The authors declare that all data supporting the findings of this study are available within this
article. The bulk RNA-seq data and single-cell RNA-seq date can be accessed at the GEO:
GSE232120 and GSE232484. Metabolomics data can be found under
MetaboLights:MTBLS7962. Further information and requests for resources and reagents should
be directed to and will be fulfilled by the lead contact, Dr. Wei Liu (liuwei5@ahau.edu.cn).
606

References

Ackermann, M. 2015. A functional perspective on phenotypic heterogeneity in microorganisms. *Nat Rev Microbiol* **13**, 497-508. DOI: https://doi.org/10.1038/nrmicro3491, PMID: 26145732.

Avraham, R., Haseley, N., Brown, D., Penaranda, C., Jijon, H.B., Trombetta, J.J., Satija, R., Shalek, A.K., Xavier, R.J., Regev, A.*, et al.* 2015. Pathogen Cell-to-Cell Variability Drives Heterogeneity in Host Immune Responses. *Cell* **162**, 1309-1321. DOI: https://doi.org/10.1016/j.cell.2015.08.027, PMID: 26343579.

Backhed, F., Ley, R.E., Sonnenburg, J.L., Peterson, D.A., and Gordon, J.I. 2005. Host-bacterial mutualism in the human intestine. *Science* **307**, 1915-1920. DOI: https://doi.org/10.1126/science.1104816, PMID: 15790844.

Barak-Gavish, N., Dassa, B., Kuhlisch, C., Nussbaum, I., Brandis, A., Rosenberg, G., Avraham, R., and Vardi, A. 2023. Bacterial lifestyle switch in response to algal metabolites. *Elife* **12**. DOI: https://doi.org/10.7554/eLife.84400, PMID: 36691727.

Becattini, S., Sorbara, M.T., Kim, S.G., Littmann, E.L., Dong, Q., Walsh, G., Wright, R., Amoretti, L., Fontana, E., Hohl, T.M.*, et al.* 2021. Rapid transcriptional and metabolic adaptation of intestinal microbes to host immune activation. *Cell Host Microbe* **29**, 378-393 e375. DOI: https://doi.org/10.1016/j.chom.2021.01.003, PMID: 33539766.

Black, E.P., Hinrichs, G.J., Barcay, S.J., and Gardner, D.B. 2018. Fruit Flies as Potential Vectors of Foodborne Illness. *J Food Prot* **81**, 509-514. DOI: https://doi.org/10.4315/0362-028X.JFP-17-255, PMID: 29474149.

Burns, A.R., Miller, E., Agarwal, M., Rolig, A.S., Milligan-Myhre, K., Seredick, S., Guillemin, K., and Bohannan, B.J.M. 2017. Interhost dispersal alters microbiome assembly and can overwhelm host innate immunity in an experimental zebrafish model. *Proc Natl Acad Sci U S A* **114**, 11181-11186. DOI: https://doi.org/10.1073/pnas.1702511114, PMID: 28973938.

Chandler, J.A., Lang, J.M., Bhatnagar, S., Eisen, J.A., and Kopp, A. 2011. Bacterial communities of diverse Drosophila species: ecological context of a host-microbe model system. *PLoS Genet* **7**, e1002272. DOI: https://doi.org/10.1371/journal.pgen.1002272, PMID: 21966276.

Claudi, B., Sprote, P., Chirkova, A., Personnic, N., Zankl, J., Schurmann, N., Schmidt, A., and Bumann, D. 2014. Phenotypic variation of Salmonella in host tissues delays eradication by antimicrobial chemotherapy. *Cell* **158**, 722-733. DOI: https://doi.org/10.1016/j.cell.2014.06.045, PMID: 25126781.

Dar, D., Dar, N., Cai, L., and Newman, D.K. 2021. Spatial transcriptomics of planktonic and sessile bacterial populations at single-cell resolution. *Science* **373**. DOI: https://doi.org/10.1126/science.abi4882, PMID: 34385369.

Davies, C.S., Worsley, S.F., Maher, K.H., Komdeur, J., Burke, T., Dugdale, H.L., and Richardson, D.S. 2022. Immunogenetic variation shapes the gut microbiome in a natural vertebrate population. *Microbiome* **10**, 41. DOI: https://doi.org/10.1186/s40168-022-01233-y, PMID: 35256003.

Deines, P., Hammerschmidt, K., and Bosch, T.C.G. 2020. Microbial Species Coexistence Depends on the Host Environment. *mBio* **11**. DOI: https://doi.org/10.1128/mBio.00807-20, PMID: 32694139.

Delannoy-Bruno, O., Desai, C., Raman, A.S., Chen, R.Y., Hibberd, M.C., Cheng, J., Han, N., Castillo, J.J., Couture, G., Lebrilla, C.B.*, et al.* 2021. Evaluating microbiome-directed fibre snacks in

gnotobiotic mice and humans. *Nature* **595**, 91-95. DOI: https://doi.org/10.1038/s41586-021-03671-4, PMID: 34163075.

Dufrene, Y.F., and Persat, A. 2020. Mechanomicrobiology: how bacteria sense and respond to forces. *Nat Rev Microbiol* **18**, 227-240. DOI: https://doi.org/10.1038/s41579-019-0314-2, PMID: 31959911.

Eldar, A., and Elowitz, M.B. 2010. Functional roles for noise in genetic circuits. *Nature* **467**, 167-173. DOI: https://doi.org/10.1038/nature09326, PMID: 20829787.

Foster, K.R., Schluter, J., Coyte, K.Z., and Rakoff-Nahoum, S. 2017. The evolution of the host microbiome as an ecosystem on a leash. *Nature* **548**, 43-51. DOI: https://doi.org/10.1038/nature23292, PMID: 28770836.

Fyhrquist, N., Muirhead, G., Prast-Nielsen, S., Jeanmougin, M., Olah, P., Skoog, T., Jules-Clement, G., Feld, M., Barrientos-Somarribas, M., Sinkko, H.*, et al.* 2019. Microbe-host interplay in atopic dermatitis and psoriasis. *Nat Commun* **10**, 4703. DOI: https://doi.org/10.1038/s41467-019-12253-y, PMID: 31619666.

Gasch, A.P., Yu, F.B., Hose, J., Escalante, L.E., Place, M., Bacher, R., Kanbar, J., Ciobanu, D., Sandor, L., Grigoriev, I.V.*, et al.* 2017. Single-cell RNA sequencing reveals intrinsic and extrinsic regulatory heterogeneity in yeast responding to stress. *PLoS Biol* **15**, e2004050. DOI: https://doi.org/10.1371/journal.pbio.2004050, PMID: 29240790.

Hanson, M.A., Dostalova, A., Ceroni, C., Poidevin, M., Kondo, S., and Lemaitre, B. 2019. Synergy and remarkable specificity of antimicrobial peptides in vivo using a systematic knockout approach. *Elife* **8**. DOI: https://doi.org/10.7554/eLife.44341, PMID: 30803481.

Hare, P.J., LaGree, T.J., Byrd, B.A., DeMarco, A.M., and Mok, W.W.K. 2021. Single-Cell Technologies to Study Phenotypic Heterogeneity and Bacterial Persisters. *Microorganisms* **9**. DOI: https://doi.org/10.3390/microorganisms9112277, PMID: 34835403.

Imdahl, F., Vafadarnejad, E., Homberger, C., Saliba, A.E., and Vogel, J. 2020. Single-cell RNA-sequencing reports growth-condition-specific global transcriptomes of individual bacteria. *Nat Microbiol* **5**, 1202-1206. DOI: https://doi.org/10.1038/s41564-020-0774-1, PMID: 32807892.

Jia, Y., Jin, S., Hu, K., Geng, L., Han, C., Kang, R., Pang, Y., Ling, E., Tan, E.K., Pan, Y.*, et al.* 2021. Gut microbiome modulates Drosophila aggression through octopamine signaling. *Nat Commun* **12**, 2698. DOI: https://doi.org/10.1038/s41467-021-23041-y, PMID: 33976215.

Jiang, X., Hall, A.B., Arthur, T.D., Plichta, D.R., Covington, C.T., Poyet, M., Crothers, J., Moses, P.L., Tolonen, A.C., Vlamakis, H.*, et al.* 2019. Invertible promoters mediate bacterial phase variation, antibiotic resistance, and host adaptation in the gut. *Science* **363**, 181-187. DOI: https://doi.org/10.1126/science.aau5238, PMID: 30630933.

Kalivoda, E.J., Stella, N.A., Aston, M.A., Fender, J.E., Thompson, P.P., Kowalski, R.P., and Shanks, R.M. 2010. Cyclic AMP negatively regulates prodigiosin production by Serratia marcescens. *Res Microbiol* **161**, 158-167. DOI: https://doi.org/10.1016/j.resmic.2009.12.004, PMID: 20045458.

Kim, E.K., Lee, K.A., Hyeon, D.Y., Kyung, M., Jun, K.Y., Seo, S.H., Hwang, D., Kwon, Y., and Lee, W.J. 2020. Bacterial Nucleoside Catabolism Controls Quorum Sensing and Commensal-to-Pathogen Transition in the Drosophila Gut. *Cell Host Microbe* **27**, 345-357 e346. DOI: https://doi.org/10.1016/j.chom.2020.01.025, PMID: 32078802.

Klein, A.M., Mazutis, L., Akartuna, I., Tallapragada, N., Veres, A., Li, V., Peshkin, L., Weitz, D.A., and Kirschner, M.W. 2015. Droplet barcoding for single-cell transcriptomics applied to

embryonic stem cells. *Cell* **161**, 1187-1201. DOI: https://doi.org/10.1016/j.cell.2015.04.044,
PMID: 26000487.

Koo, H., and Yamada, K.M. 2016. Dynamic cell-matrix interactions modulate microbial biofilm
and tissue 3D microenvironments. *Curr Opin Cell Biol* **42**, 102-112. DOI:
https://doi.org/10.1016/j.ceb.2016.05.005, PMID: 27257751.

Kuchina, A., Brettner, L.M., Paleologu, L., Roco, C.M., Rosenberg, A.B., Carignano, A., Kibler, R.,
Hirano, M., DePaolo, R.W., and Seelig, G. 2021. Microbial single-cell RNA sequencing by split-
pool barcoding. *Science* **371**, 798-+. DOI: https://doi.org/ARTN eaba5257
10.1126/science.aba5257, PMID: WOS:000619664700044.

Kuraishi, T., Binggeli, O., Opota, O., Buchon, N., and Lemaitre, B. 2011. Genetic evidence for a
protective role of the peritrophic matrix against intestinal bacterial infection in Drosophila
melanogaster. *Proc Natl Acad Sci U S A* **108**, 15966-15971. DOI:
https://doi.org/10.1073/pnas.1105994108, PMID: 21896728.

Liu, W., Lim, K.L., and Tan, E.K. 2022. Intestine-derived alpha-synuclein initiates and aggravates
pathogenesis of Parkinson's disease in Drosophila. *Transl Neurodegener* **11**, 44. DOI:
https://doi.org/10.1186/s40035-022-00318-w, PMID: 36253844.

Liu, W., Zhang, K., Li, Y., Su, W., Hu, K., and Jin, S. 2017. Enterococci Mediate the Oviposition
Preference of Drosophila melanogaster through Sucrose Catabolism. *Sci Rep* **7**, 13420. DOI:
https://doi.org/10.1038/s41598-017-13705-5, PMID: 29044155.

Llorens-Rico, V., Simcock, J.A., Huys, G.R.B., and Raes, J. 2022. Single-cell approaches in human
microbiome research. *Cell* **185**, 2725-2738. DOI: https://doi.org/10.1016/j.cell.2022.06.040,
PMID: 35868276.

Lynch, J.B., and Hsiao, E.Y. 2019. Microbiomes as sources of emergent host phenotypes. *Science*
365, 1405-1409. DOI: https://doi.org/10.1126/science.aay0240, PMID: 31604267.

Ma, P., Amemiya, H.M., He, L.L., Gandhi, S.J., Nicol, R., Bhattacharyya, R.P., Smillie, C.S., and
Hung, D.T. 2023. Bacterial droplet-based single-cell RNA-seq reveals antibiotic-associated
heterogeneous cellular states. *Cell* **186**, 877-891 e814. DOI:
https://doi.org/10.1016/j.cell.2023.01.002, PMID: 36708705.

Mancio-Silva, L., Gural, N., Real, E., Wadsworth, M.H., 2nd, Butty, V.L., March, S., Nerurkar, N.,
Hughes, T.K., Roobsoong, W., Fleming, H.E.*, et al.* 2022. A single-cell liver atlas of Plasmodium
vivax infection. *Cell Host Microbe* **30**, 1048-1060 e1045. DOI:
https://doi.org/10.1016/j.chom.2022.03.034, PMID: 35443155.

Marra, A., Hanson, M.A., Kondo, S., Erkosar, B., and Lemaitre, B. 2021. Drosophila Antimicrobial
Peptides and Lysozymes Regulate Gut Microbiota Composition and Abundance. *mBio* **12**,
e0082421. DOI: https://doi.org/10.1128/mBio.00824-21, PMID: 34253067.

Matos, R.C., Schwarzer, M., Gervais, H., Courtin, P., Joncour, P., Gillet, B., Ma, D., Bulteau, A.L.,
Martino, M.E., Hughes, S.*, et al.* 2017. D-Alanylation of teichoic acids contributes to
Lactobacillus plantarum-mediated Drosophila growth during chronic undernutrition. *Nat*
Microbiol **2**, 1635-1647. DOI: https://doi.org/10.1038/s41564-017-0038-x, PMID: 28993620.

McNulty, R., Sritharan, D., Pahng, S.H., Meisch, J.P., Liu, S., Brennan, M.A., Saxer, G., Hormoz, S.,
and Rosenthal, A.Z. 2023. Probe-based bacterial single-cell RNA sequencing predicts toxin
regulation. *Nat Microbiol*. DOI: https://doi.org/10.1038/s41564-023-01348-4, PMID: 37012420.

Morais, L.H., Schreiber, H.L.t., and Mazmanian, S.K. 2021. The gut microbiota-brain axis in behaviour and brain disorders. *Nat Rev Microbiol* **19**, 241-255. DOI: https://doi.org/10.1038/s41579-020-00460-0, PMID: 33093662.

Morgelin, M. 2017. Negative Staining and Transmission Electron Microscopy of Bacterial Surface Structures. *Methods Mol Biol* **1535**, 211-217. DOI: https://doi.org/10.1007/978-1-4939-6673-8_13, PMID: 27914081.

Muegge, B.D., Kuczynski, J., Knights, D., Clemente, J.C., Gonzalez, A., Fontana, L., Henrissat, B., Knight, R., and Gordon, J.I. 2011. Diet drives convergence in gut microbiome functions across mammalian phylogeny and within humans. *Science* **332**, 970-974. DOI: https://doi.org/10.1126/science.1198719, PMID: 21596990.

Mukherjee, S., and Bassler, B.L. 2019. Bacterial quorum sensing in complex and dynamically changing environments. *Nat Rev Microbiol* **17**, 371-382. DOI: https://doi.org/10.1038/s41579-019-0186-5, PMID: 30944413.

Olm, M.R., Dahan, D., Carter, M.M., Merrill, B.D., Yu, F.B., Jain, S., Meng, X., Tripathi, S., Wastyk, H., Neff, N.*, et al.* 2022. Robust variation in infant gut microbiome assembly across a spectrum of lifestyles. *Science* **376**, 1220-1223. DOI: https://doi.org/10.1126/science.abj2972, PMID: 35679413.

Pan, X., Sun, C., Tang, M., You, J., Osire, T., Zhao, Y., Xu, M., Zhang, X., Shao, M., Yang, S.*, et al.* 2020. LysR-Type Transcriptional Regulator MetR Controls Prodigiosin Production, Methionine Biosynthesis, Cell Motility, H(2)O(2) Tolerance, Heat Tolerance, and Exopolysaccharide Synthesis in Serratia marcescens. *Appl Environ Microbiol* **86**. DOI: https://doi.org/10.1128/AEM.02241-19, PMID: 31791952.

Pan, X., Tang, M., You, J., Osire, T., Sun, C., Fu, W., Yi, G., Yang, T., Yang, S.T., and Rao, Z. 2022. PsrA is a novel regulator contributes to antibiotic synthesis, bacterial virulence, cell motility and extracellular polysaccharides production in Serratia marcescens. *Nucleic Acids Res* **50**, 127-148. DOI: https://doi.org/10.1093/nar/gkab1186, PMID: 34893884.

Penterman, J., Abo, R.P., De Nisco, N.J., Arnold, M.F., Longhi, R., Zanda, M., and Walker, G.C. 2014. Host plant peptides elicit a transcriptional response to control the Sinorhizobium meliloti cell cycle during symbiosis. *Proc Natl Acad Sci U S A* **111**, 3561-3566. DOI: https://doi.org/10.1073/pnas.1400450111, PMID: 24501120.

Perez, R.K., Gordon, M.G., Subramaniam, M., Kim, M.C., Hartoularos, G.C., Targ, S., Sun, Y., Ogorodnikov, A., Bueno, R., Lu, A.*, et al.* 2022. Single-cell RNA-seq reveals cell type-specific molecular and genetic associations to lupus. *Science* **376**, eabf1970. DOI: https://doi.org/10.1126/science.abf1970, PMID: 35389781.

Prescott, R.D., and Decho, A.W. 2020. Flexibility and Adaptability of Quorum Sensing in Nature. *Trends Microbiol* **28**, 436-444. DOI: https://doi.org/10.1016/j.tim.2019.12.004, PMID: 32001099.

Proctor, D.M., Drummond, R.A., Lionakis, M.S., and Segre, J.A. 2023. One population, multiple lifestyles: Commensalism and pathogenesis in the human mycobiome. *Cell Host Microbe* **31**, 539-553. DOI: https://doi.org/10.1016/j.chom.2023.02.010, PMID: 37054674.

Raj, A., and van Oudenaarden, A. 2008. Nature, nurture, or chance: stochastic gene expression and its consequences. *Cell* **135**, 216-226. DOI: https://doi.org/10.1016/j.cell.2008.09.050, PMID: 18957198.

Ratzke, C., Denk, J., and Gore, J. 2018. Ecological suicide in microbes. *Nat Ecol Evol* **2**, 867-872. DOI: https://doi.org/10.1038/s41559-018-0535-1, PMID: 29662223.

Sharp, C., and Foster, K.R. 2022. Host control and the evolution of cooperation in host
microbiomes. *Nat Commun* **13**, 3567. DOI: https://doi.org/10.1038/s41467-022-30971-8, PMID:
35732630.

Somvanshi, V.S., Sloup, R.E., Crawford, J.M., Martin, A.R., Heidt, A.J., Kim, K.S., Clardy, J., and
Ciche, T.A. 2012. A single promoter inversion switches Photorhabdus between pathogenic and
mutualistic states. *Science* **337**, 88-93. DOI: https://doi.org/10.1126/science.1216641, PMID:
22767929.

Storelli, G., Defaye, A., Erkosar, B., Hols, P., Royet, J., and Leulier, F. 2011. Lactobacillus
plantarum promotes Drosophila systemic growth by modulating hormonal signals through TOR-
dependent nutrient sensing. *Cell Metab* **14**, 403-414. DOI:
https://doi.org/10.1016/j.cmet.2011.07.012, PMID: 21907145.

Storelli, G., Strigini, M., Grenier, T., Bozonnet, L., Schwarzer, M., Daniel, C., Matos, R., and
Leulier, F. 2018. Drosophila Perpetuates Nutritional Mutualism by Promoting the Fitness of Its
Intestinal Symbiont Lactobacillus plantarum. *Cell Metab* **27**, 362-377 e368. DOI:
https://doi.org/10.1016/j.cmet.2017.11.011, PMID: 29290388.

Tsai, C.N., and Coombes, B.K. 2019. The Role of the Host in Driving Phenotypic Heterogeneity in
Salmonella. *Trends Microbiol* **27**, 508-523. DOI: https://doi.org/10.1016/j.tim.2019.01.004,
PMID: 30755344.

Tsugawa, H., Cajka, T., Kind, T., Ma, Y., Higgins, B., Ikeda, K., Kanazawa, M., VanderGheynst, J.,
Fiehn, O., and Arita, M. 2015. MS-DIAL: data-independent MS/MS deconvolution for
comprehensive metabolome analysis. *Nat Methods* **12**, 523-526. DOI:
https://doi.org/10.1038/nmeth.3393, PMID: 25938372.

Valeri, F., and Endres, K. 2021. How biological sex of the host shapes its gut microbiota. *Front
Neuroendocrinol* **61**, 100912. DOI: https://doi.org/10.1016/j.yfrne.2021.100912, PMID:
33713673.

Winey, M., Meehl, J.B., O'Toole, E.T., and Giddings, T.H., Jr. 2014. Conventional transmission
electron microscopy. *Mol Biol Cell* **25**, 319-323. DOI: https://doi.org/10.1091/mbc.E12-12-0863,
PMID: 24482357.

Wong, A.C., Wang, Q.P., Morimoto, J., Senior, A.M., Lihoreau, M., Neely, G.G., Simpson, S.J.,
and Ponton, F. 2017. Gut Microbiota Modifies Olfactory-Guided Microbial Preferences and
Foraging Decisions in Drosophila. *Curr Biol* **27**, 2397-2404 e2394. DOI:
https://doi.org/10.1016/j.cub.2017.07.022, PMID: 28756953.

Xu, Z., Wang, Y., Sheng, K., Rosenthal, R., Liu, N., Hua, X., Zhang, T., Chen, J., Song, M., Lv, Y., *et
al.* 2023. Droplet-based high-throughput single microbe RNA sequencing by smRandom-seq.
Nat Commun **14**, 5130. DOI: https://doi.org/10.1038/s41467-023-40137-9, PMID: 37612289.

816
Figure 1. *Drosophila* larvae shape topographies and carrying capacities of bacterial community.
(**A-D**) Representative images of sticky "biofilm-like" formation on the surface of the sugar-corn-
yeast medium whereby *Drosophila* flies with differential robustness were raised. The
topographies of surface slick are differentially deconstructed and segmented by flies with
different robust flies. (**E**) Bacterial loads of the diet associated with strong, crowed, weak and
infertile flies, respectively. (**F**) Bacterial loads of the diet associated with male, virgin and aged
flies, respectively. (**G**) Bacterial loads of the diet associated with *Drosophila* larvae in a dosage-
dependent manner. n = 6 for each. Error bars indicate SEM. All variables have different letters,
and they are significantly different ($p < 0.01$). Kruskal–Wallis test followed by Dunn's multiple
comparisons test.
827

Figure 2. *Drosophila* larvae outcompete *S. marcescens* in the diet. (**A**) A diagram of a reductionist approach to investigate the role of *Drosophila* in regulating the physiology and behavior of *S. marcescens*. Top: Germ-free *Drosophila* larvae were generated by successive sterilization of fresh eggs with sanitizer Walch, sodium hypochloride (SH), ethanol, and PBS containing 0.01% TritonX-100T (PBST). Bottom: *S. marcescens* was cultured in a liquid

medium and re-inoculated to fly cornmeal food after washing with PBS buffer. In the meantime,
GF crawling larvae were transferred to the fly medium in the shared vials with *S. marcescens*. (**B**)
Representative images of surface slick inoculated with *S. marcescens* alone and with *S.*
marcescens over time. (**C**) The prodigiosin production of *S. marcescens* alone and in coculture at
different time points. Prodigiosin production was assessed with the spectrometer at OD_{534}. (**D**)
The bacterial load of *S. marcescens* alone and in coculture in the time course. (**E**) The survival
rate of adult flies challenged with *S. marcescens* alone and in coculture. Single and coculturing *S.*
marcescens were obtained after 24-h incubation as described in Figure 1A, and the percentage of
living female flies treated with *S. marcescens* alone and in coculture was calculated to monitor
lifespan. n = 180 for each. (**F**) RT-qPCR analysis of the expression levels of virulence-related
genes of with *S. marcescens* alone and in coculture. n = 3 for each. (**G**) Transmission electron
microscopy of *S. marcescens* alone and in coculture. Scale bars: 400 nm (left panel) or 200 nm
(right panel). (**H**) RT-qPCR analysis of the expression levels of extracellular polysaccharide
production-related genes in the control and larvae groups. n = 3 for each. Error bars indicate
SEM. All variables have different letters, and they are significantly different ($p < 0.01$). Kruskal–
Wallis test followed by Dunn's multiple comparisons test.

Figure 3. *Drosophila* larvae adjust bacterial global transcriptional adaptation to the host. (**A**)
Principal coordinate analysis (PCA) of unweighted, jack-knifed UniFrac distances of the
transcriptional profile of *S. marcescens* alone and with larvae. PC1, principal coordinate 1; PC2,
principal coordinate 2. Scattered dots in different colors represent samples from different
experimental groups. n = 4-5. (**B**) Volcano plot comparing gene expression profiles of *S.*
marcescens alone and with larvae after 24 h of incubation. X-axis represents the log2-
transformed value of gene expression change folds between larvae and control groups. Y-axis
represents the logarithmic transformation value of gene expression levels in *S. marcescens*.
Genes belonging to different pathways are represented by different colored shapes as indicated.
▽ depicts genes significantly upregulated in *S. marcescens* with larvae compared to *S.*

marcescens alone (log2 fold change < 1; adjusted $p < 0.01$), and △ depicts genes significantly

downregulated in *S. marcescens* with larvae (log2 fold change < 1; adjusted $p < 0.01$) compared

to *S. marcescens* alone. ○ depicts genes without significant alteration compared to *S. marcescens*

alone. (**C**, **D**) Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis of the significantly

downregulated and upregulated genes in *S. marcescens* with larvae compared to *S. marcescens*

alone. (**E**, **F**) RT-qPCR analysis of the expression levels of downregulated and upregulated genes

in the control and larvae groups. n = 3 for each. Error bars indicate SEM.

868
Figure 4. *Drosophila* larvae affect the global metabolic profile of *S. marcescens*. (**A**) Principal
coordinate analysis (PCA) of unweighted, jack-knifed UniFrac distances of metabolic profile of
S. marcescens alone and with larvae. PC1, principal coordinate 1; PC2, principal coordinate 2.
Scattered dots in different colors represent samples from different experimental groups. n = 5 for
each. (**B**) Volcano plot comparing metabolic profiles of between control and larvae groups after
24 h of incubation. X-axis represents the log2-transformed value of gene expression change folds
between larvae and control groups. Y-axis represents the logarithmic transformation value of
gene expression levels in *S. marcescens*. Red dots depict genes significantly upregulated in *S.*

marcescens with larvae compared to *S. marcescens* alone (log2 fold change < 1; adjusted p <
0.01), and blue dots depict genes significantly downregulated in *S. marcescens* with larvae (log2
fold change < 1; adjusted p < 0.01) compared to *S. marcescens* alone. Grey dots depict genes
without significant alteration compared to *S. marcescens* alone. (**C**) The distinct clusters of
metabolites in *S. marcescens* alone versus co-culture. (**D**, **E**) KEGG pathway analysis of the
significantly downregulated and upregulated metabolites in *S. marcescens* with larvae compared
to *S. marcescens* alone.

Figure 5. Pathogenicity heterogeneity of *S. marcescens*. (**A**) mRNA gene counts per cell for *S. marcescens* alone, with force and with larvae. Each dot represents a bacterial cell of *S. marcescens*. (**B**) Joint UMAP two-dimensional analysis showing that are distinct clusters among *S. marcescens* alone, with force and with larvae. (**C**) The cell subpopulation among the control, force and larvae groups. There were three distinct subpopulations in the control and force groups. (**D**) Mean expression levels of genes involved in ABC transporter, Quorum sensing, Secretion system, Two-component system, LPS and Peptidoglycan biosynthesis and Virulence-related genes in different subclusters. The shape of each dot indicates the proportion of cells in the cluster, while the color indicates the average activity normalized from 0 % to 100 % across all clusters. (**E**, **F**) The expression of a representative gene of ABC transporter and Quorum sensing was highlighted on the UMAP. The red color bars represent the normalized expression of a gene across all cells analyzed. (**G**, **H**) Violin plots of *livI* and *oppA* gene in different subclusters. Each dot represents a single cell and the shapes represent the expression distribution. (**I**, **J**) The expression of a representative gene of Secretion system and Virulence-related genes was highlighted on the UMAP. The red color bars represent the normalized expression of a gene across all cells analyzed. (**K**, **L**) Violin plots of *secY* and *fp* gene in different subclusters. Each dot represents a single cell and the shapes represent the expression distribution.

Figure 6. Growth heterogeneity of *S. marcescens*. (**A**) Mean expression levels of genes involved in Ribsome, DNA replication, Nitrogen metabolism and Carbon metabolism in different subclusters. The shape of each dot indicates the proportion of cells in the cluster, while the color indicates the average activity normalized from 0 % to 100 % across all clusters. (**B**, **C**) The expression of a representative gene of Ribsome and DNA replication was highlighted on the UMAP. The red color bars represent the normalized expression of a gene across all cells analyzed.

(**D**, **E**) Violin plots of *rpsL* and *trpD* genes in different subclusters. Each dot represents a single cell and the shapes represent the expression distribution. (**F**, **G**) The expression of two representative genes of Nitrogen metabolism and Carbon metabolism was highlighted on the UMAP. The red color bars represent the normalized expression of a gene across all cells analyzed.

(**H**, **I**) Violin plots of *glnK*, *glnA, pgk* and *adhE* genes in different subclusters. Each dot represents a single cell and the shapes represent the expression distribution. (**J**) Schematic of the pathogenicity and commensalism regulatory pathway.

Figure 7. Larvae-derived AMPs antagonize *S. marcescens.* (**A**) Representative images of surface slick with *S. marcescens* alone, with larvae, with secreta and with AMPs. (**B**) The prodigiosin production of *S. marcescens* alone, with larvae, with secreta and with AMPs. (**C**) Bacterial loads of *S. marcescens* alone, with larvae, with secreta and with AMPs. (**D**) Representative images of surface slick with *S. marcescens* alone, with wild-type larvae, with *ΔAMP14* larvae, with AMPs and with *ΔAMP14* larvae+AMPs. (**E**) The prodigiosin production of *S. marcescens* alone, with wild-type larvae, with *ΔAMP14* larvae, with AMPs and with *ΔAMP14* larvae+AMPs. (**F**) Bacterial loads of with *S. marcescens* alone, with wild-type larvae, with *ΔAMP14* larvae, with AMPs and with *ΔAMP14* larvae+AMPs. (**G**, **H**) RT-qPCR analysis of the expression levels of downregulated and upregulated genes in the *S. marcescens* alone, with wild-type larvae, with *ΔAMP14*larvae, with AMPs and with *ΔAMP14* larvae+AMPs. For B-C and E-F, n = 6 for each. Fo G-H, n = 3 for each. Error bars indicate SEM. All variables have different letters, and they are significantly different ($p < 0.01$). Kruskal–Wallis test followed by Dunn's multiple comparisons test.

figure supplement 1. *Drosophila* larvae modulate *S. marcescens* lifestyle switch**. (A)** Representative images of surface slick inoculated with *S. marcescens* in coculture with different numbers of crawling larvae. (**B**) The prodigiosin production of *S. marcescens* in coculture with a series of crawling larvae. n = 6 for each. (**C**) The bacterial load of *S. marcescens* in coculture with different numbers of crawling larvae. n = 6 for each. (**D**) *S. marcescens* promoted the development of GF larvae. Developmental timing of GF, *L. plantarum*-, and *S. marcescens*-associated *Drosophila* was assessed on the poor diet with 0.5% yeast. The cumulative percentage of the pupation emergence is shown over time. n = 60 for each. Error bars indicate SEM. All variables have different letters, and they are significantly different ($p < 0.01$). Kruskal–Wallis test followed by Dunn's multiple comparisons test.

944
figure supplement 2. Mechanical force didn't contribute to *S. marcescens* lifestyle. (**A**) The
schematic illustration of *S. marcescens* alone, with force and with larvae. Single and coculturing
S. marcescens were generated as described in Figure 1A. In the meantime, *S. marcescens* with
force were agitated using sterile glass sticks at the 2-h interval. (**B**) Surface slicks associated with
S. marcescens alone, with force and with larvae. (**C**) The prodigiosin production of *S.*
marcescens alone, with force and with larvae. n = 6 for each. (**D**) Bacterial loads of *S.*
marcescens alone, with force and with larvae. n = 6 for each. (**E**) RT-qPCR analysis of the
expression levels of the *pigA, pigC, pigM* and *pigI* genes of *S. marcescens* alone, with force and
with larvae. n = 3 for each. Error bars indicate SEM. All variables have different letters, they are
significantly different ($p < 0.01$). Kruskal–Wallis test followed by Dunn's multiple comparisons
test.

figure supplement 3. Interaction network analysis of transcriptome and metabolome. (**A**)
Interaction network analysis of transcriptome and metabolome of *S. marcescens* alone and with
larvae. 42 differentially expressed metabolites (red circles) are related to differentially expressed

genes involved in the ribosome (light brown), transcription (dark brown), DNA replication (light purple), energy metabolism (dark green), ABC transporters (dark blue), phosphotransferase system (light blue), quorum sensing (dark purple) and exosome (light blue). (**B**) The prodigiosin production of *S. marcescens* alone and with predicted metabolites. (**C**) Bacterial loads of *S. marcescens* alone and with predicted metabolites. n = 6 for each. (**D**) The survival rate of adult flies challenged with *S. marcescens* alone and with metabolites. (**E**) RT-qPCR analysis of the expression levels of virulence-related genes of with *S. marcescens* alone and with metabolites. n = 3 for each. Error bars indicate SEM. All variables have different letters, and they are significantly different ($p < 0.01$). Kruskal–Wallis test followed by Dunn's multiple comparisons test.

970
figure supplement 4. Heat-induced phenotypic heterogeneity of *S. marcescens.* (**A**) mRNA gene
counts per cell for the control and heat-shock groups. Each dot represents a bacterial cell of *S.
marcescens.* (**B**) Joint UMAP two-dimensional analysis of *S. marcescens* showing that are
distinct clusters between the control and heat-shock groups. (**C**) The cell subpopulation typing of
the control and heat-shock groups. (**E**, **F**) The expression of two representative genes of heat
shock and stress response was highlighted on the UMAP. The red color bars represent the
normalized expression of a gene across all cells analyzed.

978
figure supplement 5. (**A**) The survival rate of adult flies challenged with *S. marcescens* alone,
with agitation and in coculture. *S. marcescens* with alone, with agitation and in coculture were
obtained after 24-h incubation as described in Figure 1A, and the percentage of living female
flies was calculated to monitor lifespan. n = 180 for each. (**B**, **C**) The expression of two
representative genes of Nitrogen metabolism and Carbon metabolism was highlighted on the
UMAP. The red color bars represent the normalized expression of a gene across all cells
analyzed. (**D**, **E**) Violin plots of *glnK*, *glnA*, *pgk* and *adhE* genes in different subclusters. Each
dot represents a single cell and the shapes represent the expression distribution.