Circadian rhythms of macrophages are altered by the acidic pH of the tumor microenvironment.

Amelia M. Knudsen-Clark[1], Daniel Mwangi[2], Juliana Cazarin[2], Lauren M. Hablitz[3], Minsoo Kim[1], Brian J.

Altman[2,4*].

[1]Department of Microbiology and Immunology, [2]Department of Biomedical Genetics, [3]Center for

Translational Neuromedicine, and [4]Wilmot Cancer Institute, University of Rochester Medical Center,

Rochester, New York, USA.

*Corresponding author.

E-mail: Brian_Altman@URMC.rochester.edu

Abstract

Macrophages are prime therapeutic targets due to their pro-tumorigenic and immunosuppressive

functions in tumors, but varying efficacy of therapeutic approaches targeting macrophages highlights our

incomplete understanding of how the tumor microenvironment (TME) can influence regulation of

macrophages. The circadian clock is a key internal regulator of macrophage function, but how circadian

rhythms of macrophages may be influenced by the tumor microenvironment remains unknown. We

found that conditions associated with the TME such as polarizing stimuli, acidic pH, and elevated lactate

concentrations can each alter circadian rhythms in macrophages. Circadian rhythms were enhanced in

pro-resolution macrophages but suppressed in pro-inflammatory macrophages, while acidic pH had

divergent effects on circadian rhythms depending on macrophage phenotype. While cyclic AMP (cAMP)

has been reported to play a role in macrophage response to acidic pH, our results indicate that pH-driven

changes in circadian rhythms are not mediated solely by the cAMP signaling pathway. Remarkably, clock

correlation distance analysis of tumor-associated macrophages (TAMs) revealed evidence of circadian

disorder in TAMs. This is the first report providing evidence that circadian rhythms of macrophages are

altered within the TME. Our data suggest that heterogeneity in circadian rhythms at the population level

may underlie this circadian disorder. Finally, we sought to determine how circadian regulation of

macrophages impacts tumorigenesis, and found that tumor growth was suppressed when macrophages

had a functional circadian clock. Our work demonstrates a novel mechanism by which the tumor

microenvironment can influence macrophage biology through altering circadian rhythms, and the

contribution of circadian rhythms in macrophages to suppressing tumor growth.

Introduction

Tumor-associated macrophages (TAMs) are one of the most abundant leukocytes found in solid tumors, with high intra-tumoral TAM density generally associated with poor clinical outcome [1-3]. This is consistent with the largely pro-tumorigenic role of macrophages within tumors[4]. Macrophages are highly plastic professional phagocytes whose ability to sense and respond to the environment makes them uniquely equipped to protect tissue integrity under normal homeostatic conditions[5, 6]. However, within the chronically inflamed tumor microenvironment (TME), failure to resolve the inflammation can lead to uncontrolled secretion of tissue repair factors by TAMs, promoting tumor growth and metastatic capacity[7-9]. At the same time, conditions in the TME can drive TAMs to suppress potentially anti-tumorigenic inflammatory activity through various mechanisms including secretion of anti-inflammatory cytokines and expression of checkpoint inhibitors such as PD-L1, promoting immune suppression[4, 10-16].

TAMs are known to suppress the response to many standard of care treatments through their pro-tumorigenic and immunosuppressive functions, making them a prime therapeutic target[17]. However, we still have an incomplete understanding of how the TME influences macrophages, limiting our ability to target them; this is highlighted by the varying efficacy of therapeutic approaches used to target macrophages[18]. This is thought to be due in part to the significant phenotypical heterogeneity of TAMs within tumors[18-20]. Evidence suggests this heterogeneity is due to the ability of macrophages to sense and adapt to the local microenvironment, which varies within tumors depending on several factors including the distance from blood vessels and neighboring cells[21-24]. Indeed, various conditions in the TME have been shown to influence macrophage phenotype and function[4].

In particular, poor vascularization of solid tumors leads to inefficient delivery of oxygen, creating regions

of hypoxia[25]. The hypoxic response promotes enhanced glycolytic activity of cells within the region,

which, coupled with poor tissue drainage as a result of leaky vasculature, results in elevated levels of

protons and lactate, acidifying the microenvironment[26, 27]. Of conditions in the TME, it has been well

appreciated that acidic pH can promote a pro-resolution phenotype, thereby contributing to the pro-

tumorigenic and immunosuppressive functions of macrophages within tumors[28-30].

The myriad ways in which the TME can impact regulation of macrophages remain to be fully elucidated.

Circadian rhythms are a key regulatory system that is present in almost all cells of the body, and are an

understudied facet of macrophage biology[31]. Acidic pH is a condition commonly associated with the

TME that has been shown to alter circadian rhythms in cell lines[32]; however, whether pH influences

circadian rhythms in macrophages remain unknown.

Circadian rhythms are 24-hour rhythms that impart oscillations in the levels of circadian-regulated gene

transcripts and proteins in a tissue- and cell-specific manner, resulting in time-of-day-dependent variation

in many cellular processes[33, 34]. The molecular clock, which we will refer to as the circadian clock, drives

these rhythms in cells through a cell-autonomous transcription/translation feedback loop, which is

controlled in part by the transcription factor BMAL1[31]. Circadian clocks are synchronized by signals sent

out from the central circadian clock housed within the suprachiasmatic nucleus of the hypothalamus,

which entrains circadian clocks to the time of day[31]. This allows for the temporal coordination of cells

in spatially distinct tissues, although how the local microenvironment influences circadian rhythms

remains to be elucidated.

All leukocytes tested to date have functional circadian clocks[35-42]. As such, nearly every arm of the

immune response (both innate and adaptive) is subject to circadian regulation[35, 37, 43]. Time-of-day-

dependent regulation of immune responses is achieved though temporal gating of response to stimuli,

effector function, and cell trafficking[42, 44-48], all of which promote coordination between the multiple

phases of the immune response[40, 42, 49-53].

Key aspects of macrophage function are subject to circadian regulation, including cytokine secretion and

phagocytosis[36, 44, 45, 51, 54]. This results in a time-of-day-dependent macrophage response to

stimuli, which modulates the magnitude of the resulting adaptive immune response and determines

disease progression[44, 51, 55]. Circadian regulation of macrophages is of particular interest given

recent evidence of time-of-day variation in the frequency of TAMs expressing surface markers associated

with phenotype[56-58]. A promising application of such circadian variation was made evident in

leveraging observations of circadian frequency in TAMs expressing immune checkpoint blockade (ICB)

target PD-1 to increase efficacy of PD-1/PD-L1 ICB therapy by timing treatment to the time of day when

PD-1+ TAMs were most frequent[57, 59]. This suggests that leveraging time-of-day variations in

therapeutic targets could be promising avenue to increase efficacy, highlighting the importance of

understanding how circadian rhythms of macrophages may be influenced by conditions in the TME,

which remains unclear.

In this work, we present evidence that circadian rhythms of macrophages are altered in the TME. We

uncover a novel way in which two conditions within the TME, acidic pH and lactate, can influence

macrophage biology through modulation of circadian rhythms. We also find that macrophages of

different phenotypes have distinct circadian rhythms. Remarkably, we found evidence of circadian

disorder in tumor-associated macrophages, indicating that circadian rhythms are altered in macrophages

within the TME. Furthermore, our data suggest that heterogeneity in circadian rhythms at the

population level may underlie the observed circadian disorder. This work elucidates a novel way in which

the TME can alter macrophage biology, and represents the first steps to understanding how the tumor

microenvironment can alter circadian rhythms of immune cells such as macrophages.

Results

Macrophages of different phenotypes exhibit different circadian rhythms.

As macrophages are a phenotypically heterogeneous population in the TME, we first sought to

understand whether diversity in macrophage phenotype could translate to diversity in circadian rhythms

of macrophages. To this end, we used the well-established *in vitro* polarization models of IL-4 and IL-13-

stimulated (pro-resolution, often referred to as 'M2' and used as a model of pro-tumorigenic

macrophages) and IFNγ and LPS-stimulated (pro-inflammatory, often referred to as 'M1' and broadly

viewed as anti-tumorigenic) macrophages. Arginase 1 (ARG1; *Arg1*) and Inducible Nitric Oxide Synthase

(iNOS; *Nos2*) are upregulated in pro-resolution and pro-inflammatory macrophages, respectively, and

thus are commonly used as markers of the pro-resolution or pro-inflammatory phenotype in

macrophages[60]. Consistent with previous studies, we found that *Arg1* was induced in IL-4 and IL-13-

stimulated macrophages, but not IFNγ and LPS-stimulated macrophages; in contrast, *Nos2* was induced

in IFNγ and LPS-stimulated macrophages, but not IL-4 and IL-13-stimulated macrophages

(Supplementary Figure 1). This indicates that macrophages stimulated with IL-4 and IL-13 were polarized

toward a pro-resolution phenotype, while macrophages stimulated with IFNγ and LPS were polarized

toward a pro-inflammatory phenotype.

Circadian rhythms of macrophages were measured by monitoring PER2, a key component of the

circadian clock, via the rhythmic activity of the PER2-Luciferase (PER2-Luc) fusion protein in a live cell

LumiCycle luminometer (Supplementary Figure 2A)[61]. Bone marrow-derived macrophages (BMDMs)

were generated from bone marrow of mice expressing PER2-Luc. Following differentiation, the circadian

clocks of BMDMs were synchronized, and rhythms were observed for up to 4 days (Figure 2B).

To determine whether phenotype can influence circadian rhythms in macrophages, BMDMs were

cultured in the presence or absence of polarizing stimuli, and rhythms were observed by LumiCycle

(Figure 1). The amplitude of rhythms is the magnitude of change between the peak and the trough, and

is indicative of the strength of rhythms[62, 63]. Amplitude of rhythms was suppressed in pro-

inflammatory macrophages compared to unstimulated macrophages. In contrast, amplitude of rhythms

in pro-resolution macrophages was enhanced. This suggests that rhythms are suppressed in pro-

inflammatory macrophages but enhanced in pro-resolution macrophages, which agrees with previous

observations[64, 65]. Period is the amount of time it takes to complete one full oscillation, and is

indicative of how fast the circadian clock is running[66]. Compared to unstimulated macrophages, period

was lengthened in pro-resolution macrophages but shortened in pro-inflammatory macrophages. This In

line with others' observations, and suggests that the clock runs slower in pro-resolution macrophages

but runs faster in pro-inflammatory macrophages[64, 65].

Interestingly, we observed differences in damping of rhythms in polarized macrophages. Damping is

measured as the number of days required for the amplitude of rhythms to decrease by 30% of the first

cycle[67]. Damping of rhythms in most free-running cell populations, cultured in the absence of external

synchronizing stimuli, occurs naturally as the circadian clocks of individual cells in the population become

desynchronized from each other; thus, damping can be indicative of desynchrony within a

population[68]. The damping rate increases as the number of days required for rhythms to damp

decreases; conversely, as damping rate decreases as the number of days required for rhythms to damp

increases. We observed increased rate of damping in pro-inflammatory macrophages compared to

unstimulated macrophages, indicating that population-level rhythms were maintained for a shorter

period of time in pro-inflammatory macrophages. In contrast, damping rate was decreased in pro-

resolution macrophages, indicating that population-level rhythms were maintained for longer in pro-

resolution macrophages. These data suggest that pro-inflammatory macrophages may have an impaired

ability to maintain synchrony, while pro-resolution macrophages may have an enhanced ability to

maintain synchrony.

Collectively, these data suggest that pro-inflammatory macrophages have weaker rhythms and impaired

ability to maintain synchrony, while pro-resolution macrophages have enhanced rhythms and an

increased ability to maintain synchrony. This is evidence that macrophages of different phenotypes have

distinct circadian rhythms, suggesting that diversity of macrophage phenotype may lead to diversity in

macrophage circadian rhythms.

Acidic pH alters circadian rhythms of macrophages.

The TME has previously been shown to be acidic, with a pH ranging from 6.8 to 6.3; much more acidic

than the typical pH in blood and healthy tissue of 7.3-7.4[69-71]. There has been a previous report in

which acidic pH can alter circadian rhythms, but whether this applies to macrophages remains

unknown[32]. Thus, we cultured BMDMs under conditions of varying pH within a range that mimics that

found within the TME (pH7.4-pH6.5). As macrophages are a heterogeneous population in the TME, we

assessed the influence of acidic pH on rhythms of unstimulated, pro-resolution, and pro-inflammatory

macrophages. In line with previous observations, macrophages cultured at pH 6.5 were polarized toward

a pro-resolution phenotype, characterized by increased expression of *Arg1* and *Vegf* compared to

macrophages cultured at pH 7.4 (Supplementary Figure 3A). Pro-inflammatory macrophages cultured at

pH 6.5 had decreased expression of *Nos2* (iNOS) compared to those cultured at pH 7.4, suggesting that

an acidic pH of 6.5 both promotes a pro-resolution phenotype and suppresses a pro-inflammatory

phenotype.

It has been observed that inducible cyclic AMP early repressor (*Icer*), an isoform of cyclic AMP (cAMP)-

response modulator (*Crem*), is upregulated downstream of acid-sensing in macrophages, and has been

used as a "biomarker" for macrophages exposed to acidic conditions in tumors. We observed induction

of *Icer* in unstimulated and pro-resolution macrophages cultured at pH 6.5 compared to pH 7.4,

indicating that these macrophages were sensing acidic conditions (Supplementary Figure 3B). In line with

previous observations that *Icer* is induced downstream of LPS-driven TLR4 signaling, *Icer* was

upregulated in pro-inflammatory macrophages compared to unstimulated macrophages even at neutral

pH 7.4[72]. Although *Icer* was not further upregulated in pro-inflammatory macrophages at pH 6.5

compared to pH 7.4, *Nos2* was suppressed at pH 6.5 compared to pH 7.4, suggesting that pro-

inflammatory macrophages responded to acidic pH. In all, these data confirm that macrophages of

various phenotypes can sense and respond to acidic conditions within the range of pH found in the TME.

To determine whether an acidic microenvironment can influence circadian rhythms in macrophages, we

assessed rhythms of unstimulated, pro-resolution, and pro-inflammatory macrophages under normal

and acidic conditions. To this end, BMDMs were polarized toward a pro-resolution or a pro-inflammatory

phenotype, or left unstimulated, and cultured in media at a normal pH of 7.4 or at acidic pH of 6.8 or 6.5;

PER2-Luc rhythms were then observed by LumiCycle. In unstimulated and pro-resolution BMDMs, lower

pH led to enhanced amplitude, a shortening in period, and increased damping rate of rhythms at pH 6.8

and pH 6.5 relative to neutral pH 7.4 (Figure 2A,B). This suggests that in unstimulated and pro-resolution

macrophages, acidic pH can strengthen rhythms by enhancing amplitude and speed up the circadian

clock, but may impair ability to maintain synchrony. Notably, changes in amplitude and period occurred

in a dose-dependent fashion as pH decreased, indicating that rhythms are altered in a pH-dependent

manner. In contrast, pro-inflammatory macrophages cultured at pH 6.5 exhibited suppressed amplitude,

elongated period, and decreased damping rate of rhythms compared to those cultured at pH 7.4 (Figure

2C). This suggests that in pro-inflammatory macrophages, acidic pH can weaken rhythms by decreasing

amplitude and slow down the speed of the clock, but may promote the ability to maintain synchrony.

Low pH was also observed to alter the expression of the circadian clock genes *Per2*, *Cry1*, and *Nr1d1*

(REV-ERBα) over time across different macrophage phenotypes (Figure 2D,E,F). These data indicate that

exposure to acidic pH can induce changes in circadian rhythms of macrophages. Interestingly, while

rhythms of unstimulated and pro-resolution macrophages are enhanced under acidic pH despite

increased damping rate, rhythms of pro-inflammatory macrophages are suppressed under acidic

conditions but have improved damping rate. This suggests that acidic pH modulates rhythms differently

in macrophage of different phenotypes. The observation that acidic pH can enhance rhythms is

particularly interesting, given that acidic pH is a stressful condition that can compromise macrophage

survival (Supplementary Figure 4)[30].

While BMDMs are a widely used model for studying macrophages *in vitro*, there are biological

differences between BMDMs generated in culture and tissue-resident macrophages differentiated *in*

vivo. Thus, we sought to determine whether our observations of pH-induced changes in rhythms were

relevant to macrophages differentiated *in vivo*. To this end, we harvested peritoneal macrophages from

mice expressing PER2-Luc in the morning at ZT0 (6 AM) or in the evening at ZT12 (6 PM). Peritoneal

macrophages were cultured in media at neutral pH of 7.4 or acidic pH of 6.5 and observed by LumiCycle.

Recapitulating our results in BMDMs, peritoneal macrophages exhibited increased amplitude, decreased

period, and increase rate of damping at pH 6.5 compared to pH 7.4 (Figure 3A). These data indicate that

pH-dependent changes in circadian rhythms are relevant to *in vivo*-differentiated macrophages.

Circadian rhythms confer time-of-day variability in response to stimuli. As we have observed that acidic

pH can influence circadian rhythms of macrophages, we sought to understand if macrophages may be

more or less susceptible to pH-induced changes in rhythms depending on time of day of exposure. To

this end, we compared the magnitude of change in amplitude, period, and damping in peritoneal

macrophages when exposed to acidic pH 6.5 compared to neutral pH 7.4 at different times of day (Figure

3B). We observed no significant difference in the pH-driven change in amplitude, period, or damping in

rhythms of peritoneal macrophages taken in the morning at ZT0 compared to those taken in the evening

at ZT12. This indicates that the influence of pH on rhythms of macrophages was similar when exposed to

acidic pH in the morning or in the evening, which suggests that macrophages are similarly susceptible to

pH-induced changes in rhythms regardless of time of day of exposure.

Lactate alters circadian rhythms of macrophages in a manner distinct from acidic pH.

Elevated lactate concentrations often co-localize to regions of high acidity, due to the export of both

protons and lactate by glycolytic cells[73-75]. In tumors, concentration of lactate has been observed to

be present in concentrations of 30mM, elevated over typical lactate levels in blood and healthy tissue of

1.5-3mM[76]. There are previous reports that lactic acid can promote polarization of macrophages

toward a pro-resolution phenotype[77]. Thus, we sought to understand if lactate may be a feature of the

TME capable of influencing circadian rhythms of macrophages in addition to acidic pH. To this end, we

cultured BMDMs in the presence or absence of 25 mM sodium-L-lactate. In line with previous

observations, BMDMs exposed to lactate had elevated levels of *Vegf*; however, we did not observe

significant elevation of *Arg1* (Figure 4A)[77].

We then cultured BMDMs at normal pH 7.4 or acidic pH 6.5, in the presence or absence of 25 mM

sodium-L-lactate, and monitored circadian rhythms (Figure 4). Rhythms of BMDMs at pH 7.4 exposed to

25 mM sodium-L-lactate had elongated period and decreased damping time compared to BMDMs

cultured at pH 7.4 without sodium-L-lactate (Figure 4B). This suggests that lactate can slow down the

circadian clock and may impair the ability of macrophages to maintain synchrony. Interestingly, these

changes in rhythms are different from those observed in under acidic conditions, indicating that lactate

can modulate circadian rhythms in macrophages in a manner distinct from acidic pH.

As previously observed, macrophages exposed to acidic pH 6.5 exhibited increased amplitude, shortened

period, and increased damping rate of circadian rhythms. When BMDMs were exposed to both acidic pH

and elevated lactate, the increased amplitude observed at pH 6.5 is maintained; however, the shortened

period observed at pH 6.5 is lost, with period lengthened in BMDMs cultured in 25 mM sodium-l-lactate.

The increased damping rate of rhythms in BMDMs cultured at pH 6.5 compared to pH 7.4 is maintained,

and is further dampened by exposure to 25 mM sodium-l-lactate. These data indicate that changes in

rhythms associated with acidic conditions persisted when co-exposed to elevated lactate. In all, these

data indicate that concentration of lactate similar to that present in the TME can influence circadian

rhythms of macrophages. Lactate altered rhythms differently than acidic pH, and when macrophages

were exposed to acidic pH and lactate together, rhythms were further altered. This suggests that when

macrophages are exposed to multiple conditions capable of altering circadian rhythms, each condition

may contribute to a combined effect on rhythms that differs from its individual impact.

Induction of cAMP signaling alone is not sufficient to fully drive changes in circadian rhythms

associated with acidic pH.

Evidence in the literature suggests that acidic pH is primarily sensed by macrophages via certain G

protein-coupled receptors (GPCRs), inducing an increase in intracellular cAMP that drives downstream

signaling through the cAMP pathway[29]. Transcriptional changes downstream of cAMP signaling

subsequently promotes a pro-resolution phenotype[29, 78, 79]. Downstream of cAMP signaling,

transcription of the *Crem* isoform *Icer* is induced, which has been used as a "biomarker" for

macrophages exposed to acidic conditions in tumors[29]. In line with previous reports, we have

observed induction of *Icer* in macrophages under acidic pH, suggesting that cAMP signaling is being

induced under acidic conditions (Supplementary Figure 2A)[29]. This occurs as early as 2 hours,

concurrent with changes in rhythms, which are observed by 6 hours following exposure to acidic

conditions. It has been shown that induction of cAMP signaling alone is sufficient to drive a pro-

resolution phenotype in macrophages similar to that observed under acidic conditions[29, 78].

Additionally, cAMP signaling has been previously observed to modulate circadian rhythms in SCN and rat

fibroblasts[80, 81]. Thus, we sought to understand if the cAMP signaling pathway may be mediating the

pH-induced changes in circadian rhythms in macrophages.

We first sought to determine if elevation in intracellular cAMP alone was sufficient to drive changes in

rhythms observed in macrophages under acidic conditions. To this end, we treated macrophages with

forskolin, an adenylyl cyclase activator that stimulates production of cAMP, or IBMX, which drives

accumulation of cAMP through inhibition of phosphodiesterases (PDEs). We used a range of doses

similar to those previously shown to induce cAMP signaling in macrophages in the literature[29, 80, 82].

Treatment with either forskolin or IBMX increased amplitude of rhythms in macrophages, but not to the

same magnitude as acidic pH, and did not result in a changed period (Figure 5). Moreover, amplitude of

rhythms was not altered in forskolin- or IBMX-treated macrophages at pH 6.5, indicating neither forskolin

treatment nor IBMX treatment had any additional effect on rhythms under acidic conditions. These data

indicate that in macrophages, cAMP signaling alone induces enhanced amplitude of rhythms similar to

low pH, but the magnitude of this change is far less; additionally, period, which is altered under acidic

conditions, remains unchanged. This suggests that cAMP signaling alone may contribute to but is not

sufficient to fully recapitulate the changes in rhythms observed under acidic conditions.

Adenylyl cyclase inhibitor MDL-12330A can suppress pH-mediated changes in amplitude of circadian

rhythms and pro-resolution phenotype without suppressing cAMP signaling.

To further test whether pH-induced changes in rhythms are mediated by cAMP signaling, we treated

BMDMs with MDL-12330A (henceforth referred to as MDL-12), an adenylyl cyclase inhibitor which has

previously been shown to suppress cAMP signaling in macrophages under acidic conditions[29]. As

previously observed, amplitude of rhythms was enhanced in macrophages cultured at pH 6.5, while

period was shortened and damping rate of rhythms was increased. When BMDMs cultured at pH 6.5

were treated with MDL-12, the elevated amplitude of rhythms observed at pH 6.5 was suppressed

(Figure 6A). Notably, this occurred in a dose-dependent manner, indicating that this is a drug-dependent

effect. Importantly, rhythms of MDL-12-treated macrophages at pH 7.4 had similar amplitude to vehicle-

treated macrophages at pH 7.4. This suggests that the inhibitory effect of MDL-12 on pH-induced

enhancement of amplitude in macrophage rhythms was specific to signaling occurring under acidic

conditions. However, MDL-12 treatment of macrophages at pH 7.4 resulted in shortened period and

decreased damping rate compared to vehicle-treated macrophages, suggesting that MDL-12 treatment

had additional effects on rhythms. Despite this, MDL-12-mediated suppression of pH-driven changes in

amplitude, but not period or damping, suggests that the pH-driven changes in these different

parameters of rhythms may occur through different pathways. Interestingly, although the adenylyl

cyclase inhibition by MDL-12 is reported to be irreversible, we found that pretreatment up to 2 hours is

not sufficient to suppress pH-induced changes in amplitude. Only when macrophages continued to be

cultured with MDL-12 while exposed to acidic conditions was amplitude suppressed (Supplementary

Figure 5). Meanwhile, co-treating cells with acidic pH and MDL12 without any pre-treatment was

sufficient to suppress elevation of amplitude under acidic conditions (Figure 6A).

Evidence suggests that acidic pH signals through the cAMP pathway to promote a pro-resolution

phenotype in macrophages, with induction of *Icer* occurring directly downstream of cAMP signaling[29].

Despite preventing changes in amplitude under acidic pH, MDL12 treatment at the dose and treatment

schedule used does not suppress induction of *Icer* in macrophages under acidic conditions (Figure 6B).

However, induction of *Arg1* expression in macrophages under acidic conditions was suppressed by MDL-

12. This suggests that at the dose and treatment strategy used, MDL-12 partially suppresses the

response of macrophages to acidic pH by suppressing the pH-driven polarization toward a pro-resolution

phenotype and changes in amplitude.

To further investigate how MDL-12 was influencing cAMP signaling at the dose and treatment strategy

used, we evaluated phosphorylation of cyclic AMP-response element binding protein (CREB).

Phosphorylation of CREB occurs downstream of cAMP and has commonly been used as a readout to

assess induction of cAMP production in macrophages[78, 82]. In line with evidence in the literature that

exposure to acidic pH drives an increase in intracellular cAMP in macrophages[29], we observed that

downstream phosphorylation of CREB was elevated in macrophages exposed to acidic pH compared to

those in non-acidic conditions (Figure 6C). Unexpectedly, pCREB levels remained elevated in BMDMs at

pH 6.5 despite treatment with MDL-12, indicating that pH-driven phosphorylation of CREB was not

suppressed by MDL12 treatment. In fact, pCREB was elevated in MDL-12-treated BMDMS at pH 7.4,

suggesting that MDL-12 treatment alone induced phosphorylation of CREB. This is particularly surprising

considering that amplitude was not altered in MDL-12-treated macrophages at neutral pH 7.4 despite

elevated pCREB. This suggests that some elements of the cAMP signaling pathway, such as pCREB, may

be divorced from the pH-induced changes in rhythms. Collectively, our data indicate that while the cAMP

signaling pathway is induced under acidic conditions, pH-induced changes in rhythms may not be

attributed to cAMP signaling alone, as MDL-12 treatment suppressed pH-induced changes in amplitude

of rhythms, but not period or damping, without suppressing signaling through the cAMP pathway.

There is evidence of circadian disorder in tumor-associated macrophages, which may be due to

heterogeneity in circadian rhythms within the TAM population.

As we have observed that acidic pH at levels commonly observed in the TME can alter circadian rhythms

in macrophages *in vitro* and *ex vivo*, we next sought to investigate whether circadian rhythms can be

altered in the TME *in vivo*. Using publicly available data, we analyzed gene expression of tumor-

associated macrophages isolated from LLC (Lewis Lung carcinoma) tumors[83]. In line with previous

observations, TAMs had elevated expression of *Arg1* relative to peritoneal macrophages (Figure 7A).

Expression of *Crem*, which encodes *Icer*, was also elevated in TAMs, indicating that these TAMs were

exposed to acidic conditions within the TME (Figure 7A)[29].

To understand the status of the circadian clock in TAMs, we performed clock correlation distance (CCD)

analysis. This analysis has previously been used to assess functionality of the circadian clock in whole

tumor and in normal tissue[84]. As the circadian clock is comprised of a series of

transcription/translation feedback loops, gene expression is highly organized in a functional, intact clock,

with core clock genes existing in levels relative to each other irrespective of the time of day. In a

synchronized population of cells, this ordered relationship is maintained at the population level, which

can be visualized in a heatmap. Using previously published RNA-sequencing data of mouse peritoneal

macrophages[44], we found that gene co-expression of clock genes was ordered in a population of wild

type macrophages with functional clocks and intact circadian rhythms (Figure 7B). In contrast, clock gene

co-expression is disordered in BMAL1 KO macrophages with a genetic disruption of the circadian clock,

leading to disruption of circadian rhythms (Supplementary Figure 6). Clock correlation distance analysis

revealed that, similar to the BMAL1 KO peritoneal macrophages, the co-expression relationship between

the core circadian clock genes in TAMs is significantly more disordered than that of WT peritoneal

macrophages (Figure 7C). This indicates that there is population-level disorder in the circadian rhythms

of tumor-associated macrophages, suggesting that circadian rhythms are indeed altered in macrophages

within the TME.

Circadian disorder assessed by CCD has previously been used to infer disruption of circadian

rhythms[85]. Indeed, we observed that genetic disruption of circadian rhythms by BMAL1 KO resulted in

a disordered clock, as observed in peritoneal macrophages (Figure 7B). However, since CCD is a

population-level analysis, heterogeneity of rhythms, as observed in a desynchronous cell population,

rather than disruption of rhythms, may also underlie the circadian disorder observed by CCD.

Heterogeneity in macrophage phenotype, exposure to acidic pH, and lactate are all factors present in the

TME and relevant to tumor-associated macrophages. We have observed that each of these factors can

alter circadian rhythms in macrophages, both alone and in combination with each other. Thus, we sought

to understand if heterogeneity in macrophage rhythms could be contributing to the disorder in clock

gene co-expression and poor CCD score indicative of population-level disorder in TAM rhythms.

To address this, we examined if differences in rhythms of macrophages within a population might

contribute to population-level disorder as measured by CCD. To this end, we used publicly available data

of peritoneal macrophages taken at different times of day in four-hour intervals across two days[36]. We

then constructed four different sample groups in which samples were pooled according to time of day of

harvest. As a control population for synchronized cell population with homogenous rhythms, samples

taken at the same time of day were pooled. We then modeled a progressively desynchronized

population with increased differences in phase of rhythms by pooling samples that were taken four

hours apart, eight hours apart, or twelve hours apart (Figure 8A). CCD was then performed on these four

populations (Figure 8B). CCD score worsened as populations became increasingly desynchronized, with

the 12hr desynchronized population having a significantly worse CCD score than synchronized,

homogenous macrophage population (Figure 8C). This indicates that as circadian rhythms of individual

macrophages within a population become more different from each other, circadian disorder increases

at the population-level. This analysis suggests that heterogeneity in rhythms, as observed with

desynchrony, may underlie population-level disorder of the circadian clock as measured by CCD.

Collectively, these data suggest that rhythms are disordered in tumor-associated macrophages, and that

heterogeneity in macrophage rhythms within the TAM population may underlie this population-level

disorder in circadian rhythms of TAMs observed by CCD.

Circadian rhythms of macrophages can influence tumor growth in a murine model of pancreatic

cancer.

We then sought to determine how circadian rhythms in tumor-associated macrophages may influence

tumor growth in KCKO, which is a murine model of pancreatic ductal adenocarcinoma (PDAC)[86, 87]. To

this end, we used a genetic disruption of the circadian clock in macrophages. While myeloid-specific

genetic mouse models are useful, they are not macrophage-specific; thus, co-injection experiments are

commonly used to determine macrophage-specific roles[77, 88, 89]. Thus, we co-injected BMDMs from

WT or BMAL1 KO mice along with KCKO cells into WT mice, and tumor growth was measured. We saw a

significant increase in the growth of tumors co-injected with BMAL1 KO macrophages compared to those

co-injected with WT macrophages (Figure 9). These results suggest intact circadian rhythms of

macrophages can restrain tumor growth, in agreement with similar published findings in a murine model

of melanoma[88].

Discussion

Macrophages experience altered environmental conditions within the tumor microenvironment, but

how these may affect macrophage circadian rhythms remains unclear. Here we assessed whether

circadian rhythms are altered in macrophages within the TME. To this end, we explored whether

conditions commonly associated with the tumor microenvironment could influence circadian rhythms in

macrophages. As TAMs are phenotypically heterogenous, we first assessed circadian rhythms in

macrophages polarized toward different phenotypes. We found that polarization state affects circadian

rhythmicity, with pro-inflammatory macrophages exhibiting far weaker rhythms that pro-resolution

macrophages (Figure 1). We then modeled acidic conditions in the TME by exposing macrophages to pH

and lactate levels similar to those found in the TME, and found that low pH in particular dramatically

altered the rhythms of macrophages (Figures 2-4). Changes in cAMP signaling may contribute to these

changes in rhythmicity, but low pH induced alterations far beyond what is observed by enhancing cAMP

signaling pharmacologically (Figure 5). While the adenylyl-cyclase inhibitor MDL-12 largely rescues the

changes in amplitude observed in low pH, our data suggest that a pathway other than canonical cAMP

signaling may be involved in this effect (Figure 6). Finally, we assessed the status of the circadian clock in

tumor-associated macrophages, the potential contribution of heterogeneity in circadian rhythms to

population-level rhythms, and assessed whether the circadian regulation of macrophages impacts tumor

growth. Our results indicate that macrophage rhythms as a whole are disordered within tumors (Figure

7), and that heterogeneity in rhythms within the tumor-associated macrophage population may underlie

this observed circadian disorder (Figure 8). We further demonstrated that the intact macrophage

circadian clock can suppress tumor growth (Figure 9). Overall, our results for the first time demonstrate

that exposure of macrophages to conditions associated with the tumor microenvironment can influence

circadian rhythms, a key aspect of macrophage biology.

A critical question in understanding the role of circadian rhythms in macrophage biology is determining

how different polarization states of macrophages affect their internal circadian rhythms. This is

especially important considering that tumor-associated macrophages are a highly heterogeneous

population. Our data indicate that compared to unstimulated macrophages, rhythms are enhanced in

pro-resolution macrophages, characterized by increased amplitude and improved ability to maintain

synchrony; in contrast, rhythms are suppressed in pro-inflammatory macrophages, characterized by

decreased amplitude and impaired ability to maintain synchrony (Figure 1). These agree with the

findings of others which have shown that polarizing stimuli alone and in combination with each other

can alter rhythms differently in macrophages[64, 65]. In a tumor, macrophages exist along a continuum

of polarization states and phenotypes[18-21, 24]. Thus, while our characterizations of rhythms in *in vitro*-

polarized macrophages provide a foundation for understanding how phenotype affects circadian

rhythms of macrophages, further experiments will be needed to assess macrophages across the full

spectrum of phenotypes. Indeed, alteration of rhythms may be just as highly variable and context-

dependent as phenotype itself.

In addition to polarizing stimuli, tumor-associated macrophages are exposed to a variety of conditions

within the tumor microenvironment that may alter their circadian rhythms. We observed that exposure

to acidic pH alters rhythms in macrophages, increasing amplitude of pro-resolution macrophages but

suppressing amplitude of pro-inflammatory macrophages (Figure 2). This indicates that pH affects

rhythms differently depending on phenotype, hinting at additional layers of complexity in how the

environment could contribute to changes in circadian rhythms. Even further changes in rhythms were

observed when macrophages were exposed to lactate in conjunction with acidic pH (Figure 4). These

observations suggest that the combination of stimuli present in the microenvironment such as lactate

and low pH, as well as various polarizing stimuli, can each contribute to modulate rhythms, resulting in

highly context-dependent changes in circadian rhythms of macrophages based on the

microenvironment. As macrophages are highly plastic and are exquisitely capable of sensing and

responding to their environment, one could reason that changes in circadian rhythms, and downstream

circadian regulation, are a mechanism by which macrophages can adopt different programs to respond

to their environment.

It was previously observed that acidic pH can disrupt circadian rhythms in cell lines[32]. However, while

acidic pH alters rhythms in macrophages, it does not ablate them. This suggests that the influence of

acidic pH on circadian rhythms can vary between cell types. pH-induced circadian disruption was found

to be driven by inhibition of mTORC1 activity in cell lines, and there was evidence to suggest that

mTORC1 activity was sensitive to pH in T cells [90]. Thus, the role of mTORC1 activity in mediating pH-

driven changes in circadian rhythms of macrophages will be a topic of future investigation.

The mechanism through which acidic pH can modulate the circadian clock in macrophages remains

unclear. Evidence in the literature suggests that acidic pH promotes a pro-resolution phenotype in

macrophages by driving signaling through the cAMP pathway[29]. It has previously been shown that

cAMP signaling can modulate the circadian clock[81]. However, our data indicate that cAMP signaling is

not fully sufficient to confer pH-mediated changes in circadian rhythms in macrophages. Treatment with

forskolin or IBMX enhanced amplitude of oscillation, but was not sufficient to fully recapitulate pH-

induced changes in rhythms (Figure 5). Interestingly, treatment with MDL-12, commonly known as an

inhibitor of adenylyl cyclase[29, 91], resulted in suppression of pH-induced changes in amplitude of

circadian rhythms and alleviated pH-mediated polarization toward a pro-resolution phenotype,

measured by *Arg1* expression (Figure 6). However, our data indicate that treatment with MDL-12 at the

dose and treatment strategy used did not inhibit signaling through the cAMP signaling pathway, as *Icer*

induction and phosphorylation of CREB were not suppressed. While MDL-12 is commonly used as an

adenylyl cyclase inhibitor, it has also been documented to have inhibitory activity toward

phosphodiesterases (PDEs) and the import of calcium into the cytosol through various mechanisms[92,

93]. This is of particular interest, as calcium signaling has also been shown to be capable of modulating

the circadian clock[94]. Furthermore, while acid-sensing through GPCRs have been the most well-

characterized pathways in macrophages, there remain additional ways in which acidic pH can be sensed

by macrophages such as acid-sensing ion channels[95, 96]. Further work is required to understand the

signaling pathways through which pH can influence macrophage phenotype and circadian rhythms.

Taken together, our findings indicate that both macrophage polarization state and exposure to

environmental conditions such as low pH and lactate each affect the circadian rhythms of macrophages.

One prominent question that remains elusive is whether circadian rhythms of macrophages are

governed by phenotype, or whether phenotype is subject to control of circadian rhythms. Alternately,

the relationship between the two could very well be bi-directional. Acidic pH promotes a pro-resolution

phenotype and suppresses pro-inflammatory activity as early as 2 hours post-exposure (Supplementary

Figure 3), with changes in rhythms observed by 12 hours (Figure 2). This suggests that pH-driven changes

in phenotype may precede changes in rhythms. How much the pH-driven changes in rhythms is

influenced directly by pH sensing, or indirectly through the influence of pH on phenotype, remains

unknown. Certainly, polarization toward a pro-resolution phenotype and exposure to acidic pH each

have different effects on circadian rhythms of macrophages; while polarization of macrophages toward a

pro-resolution phenotype increases amplitude and decreases damping rate of circadian rhythms,

exposure to acidic pH not only increases amplitude, but shortens period and increases damping rate of

rhythms (Figures 1, 2). Additionally, acidic pH drives changes in rhythms of pro-resolution macrophages

similar to that of unstimulated macrophages, suggesting that the signaling driving these changes in

rhythms are not redundant between the two conditions. This suggests that acidic pH-promoted

polarization toward a pro-resolution phenotype does not on its own account for the pH-driven changes

in rhythms. Studies have shown that circadian rhythms are regulated by metabolism[97], and it is well-

appreciated that macrophages of distinct phenotypes prioritize different metabolic pathways[98]. Thus,

the metabolic needs of macrophages of different phenotypes likely influences the response to pH and

other metabolically stressful conditions in the TME, and downstream effects on circadian rhythms.

Indeed, we observed that exposure to acidic pH had an opposite effect on rhythms of pro-inflammatory

macrophages compared to pro-resolution macrophages (Figure 2). Further work is required to

understand the interplay between macrophage phenotype, metabolism, and circadian rhythms.

We observed that acidic pH appears to enhance circadian rhythms of unstimulated and pro-resolution

macrophages, and we and others have shown evidence that macrophages are exposed to an acidic

environment within the TME[28, 29]. However, analysis of TAMs by clock correlation distance (CCD)

presents evidence that rhythms are disordered in TAMs (Figure 7). CCD is one of the most practical tools

currently available to assess circadian rhythms due to its ability to assess rhythms independent of time

of day and without the need for a circadian time series, which is often not available in publicly available

data from mice and humans[84]. However, CCD has some limitations, primary of which is that it is a

measure of population-level circadian rhythms. There is currently a major barrier in the circadian field in

the ability to measure circadian rhythms at the single-cell level. Some recent work has defined single-

cell rhythms and uncovered a surprising amount of heterogeneity in mouse and fly tissues[99-101]**.** A full

measure of single-cell circadian dynamics within a tissue must be assessed to facilitate studies of

rhythms *in vivo* where cell populations are highly heterogeneous, such as within the tumor

microenvironment or at a site of infection. Indeed, we present evidence that heterogeneity of circadian

rhythms within a given population may underlie this observed circadian disorder (Figure 8). We have

shown that various stimuli can alter rhythms of macrophages in a complex and contributing manner,

including polarizing stimuli, acidic pH, and lactate. TGFβ is produced by a variety of cells within the TME,

and was recently identified as a signal that can modulate circadian rhythms[102, 103]. Thus, it is

conceivable that, in addition to acidic pH, other stimuli in the TME are influencing circadian rhythms to

drive population-level disorder that we are observing by CCD. Notably, we cannot rule out that the

disorder observed by CCD may be indicative of circadian disruption.

In light of our observations that conditions associated with the TME can alter circadian rhythms in

macrophages, it becomes increasingly important to understand the relevance of macrophage rhythms to

their function in tumors. Data from our lab and others suggest that disruption of the macrophage-

intrinsic circadian clock accelerates tumor growth, indicating that circadian regulation of macrophages is

tumor-suppressive[88]. This agrees with complementary findings that behavioral disruption of circadian

rhythms in mice (through chronic jetlag) disrupts tumor macrophage circadian rhythms and accelerates

tumor growth[56]**.** It remains unclear whether this is through the pro-tumorigenic functions of

macrophages such as extracellular matrix remodeling or angiogenesis, through suppression of the anti-

tumor immune response, or a combination of both functions. Further work will be needed to tease apart

these distinctions.

Whereas much work has been done to characterize how macrophages are regulated within the TME, the

impact of the TME on circadian rhythms of macrophages remained elusive. Our work uncovers a novel

way in which conditions associated with the TME can influence macrophage biology through modulation

of circadian rhythms. While the majority of studies investigating the circadian regulation of macrophages

have been conducted studying macrophages under homeostatic conditions or in response to acute

inflammation[36, 38, 44, 45, 104], our work contributes to an emerging body of evidence that the tissue

microenvironment can influence circadian rhythms[102]. This is increasingly important when considering

the role of circadian rhythms in immune responses at sites of ongoing, chronic inflammation where the

microenvironment is altered, such as within tumors. In identifying factors within the TME that can

modulate circadian rhythms of macrophages and uncovering evidence of circadian disorder within

tumor-associated macrophages, this work lays the foundation for further studies aimed at understanding

how the TME can influence the function of tumor-associated macrophages through modulation of

circadian rhythms.

Limitations of the Study

Our observations of rhythms in macrophages of different phenotypes are limited by *in vitro* polarization

models. It is important to note that while our data suggest that pro-inflammatory macrophages have

suppressed rhythms and increased rate of desynchrony, it remains unclear the extent to which these

findings apply to the range of pro-inflammatory macrophages found *in vivo*. We use IFNγ and LPS co-

treatment *in vitro* to model a pro-inflammatory macrophage phenotype that is commonly referred to as

'M1', but under inflammatory conditions *in vivo*, macrophages are exposed to a variety of stimuli that

result in a spectrum of phenotypes, each highly context-dependent. The same is true for for 'M2';

different tissue microenvironment are different and pro-resolution macrophages exist in a spectrum.

Rhythms were heavily suppressed in pro-inflammatory macrophages, which makes analysis of rhythm

parameters in pro-inflammatory macrophages more challenging as amplitude and signal reaches limit of

detection. We assessed parameters of oscillation using LumiCycle Analysis, a method commonly used in

the circadian field[64, 105]. Our observations of changes in amplitude and period in pro-inflammatory

macrophages compared to unstimulated macrophages agrees with the literature, where these changes

in rhythms have been observed using LumiCycle as well as by mRNA[64, 65]. This supports the validity

and reproducibility of our observations despite the challenges of observing and analyzing rhythms of

pro-inflammatory macrophages.

Methods

Animals

Mice were maintained in individually ventilated cages with bedding and nesting material in a

temperature-controlled, pathogen-free environment in the animal care facility at the University of

Rochester. All animal protocols were approved by the University of Rochester Committee of Animal

Resources (UCAR). All experiments were performed in compliance with the NIH- and University of

Rochester-approved guidelines for the use and care of animals, as well as recommendations in the Guide

for the Care and Use of Laboratory Animals of the National Research Council[106]. Mice were housed on

a 12:12 light dark cycle. In some cases, to ease timepoint collection, mice were housed under reverse

lighting conditions in a 12:12 dark light cycle for at least 2 weeks prior to use in experiments. Mice used

for experiments were between the ages of 8-14 weeks old; both male and female mice were used. Mice

were euthanized humanely prior to harvesting peritoneal macrophages or bone marrow.

Previously characterized mice with a myeloid-specific deletion of BMAL1 (LysM-cre$^{+/-}$Bmal1$^{flox/flox}$;

referred to as BMAL1 KO mice)[45] in a C57BL/6 background were generated by crossing LysM-cre

mice[107] with Bmal1$^{flox/flox}$ mice[108]. These mice were further crossed with PER2-Luc mice[61] to

generate BMAL1 KO or wild-type control mice (LysM-cre$^{-/-}$Bmal1$^{flox/flox}$; referred to as WT) expressing

PER2-Luc. PER2-Luc (strain #006852), LysM-cre (strain #004781), and Bmal1$^{flox/flox}$(strain #007668) mice

used for breeding to generate WT and BMAL1 KO mice were purchased from the Jackson Laboratory.

Differentiation and culture of bone marrow-derived macrophages

Bone marrow-derived macrophages (BMDMs) were generated from bone marrow isolated from WT

mice using a well-established protocol for differentiation of BMDMs over 7 days[109, 110]. In brief, bone

marrow cells were seeded at 200,000 cells/mL on non-tissue culture treated-plates in BMDM

Differentiation Media: RPMI (Corning, CAT#MT10040CV) supplemented with 20% (v/v) L929 supernatant

and 10% (v/v) heat-inactivated (HI) fetal bovine serum (FBS) (Cytiva, CAT#SH30396.03), supplemented

with 100 U/mL Penicillin-Streptomycin (Gibco, CAT# 15140122). Cells were grown at 37°C in air enriched

with 5% CO_2. On day 3, additional BMDM Differentiation Media was added to cells. On day 6 of the

differentiation protocol, BMDMs were seeded at $1.2*10^6$ cells/mL and left in BMDM Differentiation

Media, and kept at 37°C in air enriched with 5% CO_2. On day 7, BMDM Differentiation Media was

removed and BMDMs were synchronized.

To synchronize BMDMs, BMDMs were first serum starved for 24 hours in serum-free media (RPMI,

supplemented with 100 U/mL Penicillin-Streptomycin); BMDMs were then subjected to serum shock by

replacing serum-free media with RPMI supplemented with 50% (v/V) HI horse serum (Corning,

CAT#35030CV) at 37^oC in air enriched with 5% CO_2. At the end of this synchronization protocol, media

was replaced with Atmospheric Media, which has been formulated for use at atmospheric CO_2 levels and

enhanced pH stability by increasing buffering capacity at low pH[32]: RPMI (Corning, CAT#50-020-PC),

25mM HEPES (Gibco, CAT#15630080), 25mM PIPES (Sigma, CAT#P1851), supplemented with 10% (v/v)

HI FBS and 100 U/mL Penicillin-Streptomycin. Atmospheric Media was adjusted to pH 7.4, 6.8, or 6.5

with NaOH and filter-sterilized.

BMDMs cultured in Atmospheric Media at pH 7.4, 6.8, or 6.5 were either left unstimulated or were

polarized toward a pro-resolution ('M2') or pro-inflammatory phenotype ('M1') by addition of 10 ng/mL

IL-4 (PeproTech, CAT#214-14) and 10 ng/mL IL-13 (PeproTech, CAT# 210-13), or 50 ng/mL IFNγ

(PeproTech, CAT#315-05) and 100 ng/mL LPS (Invitrogen, CAT#00497693), respectively. For lactate

experiments, sodium-L-lactate (Sigma, CAT#L7022) or vehicle was added to Atmospheric Media for 25

mM sodium-L-lactate or 0 mM sodium-L-lactate in Atmospheric Media. For interrogation of cAMP

signaling pathway, BMDMs were cultured in Atmospheric Media at pH 7.4 or 6.5 with vehicle or 5, 10, or

15 µM MDL-12330A (Sigma, CAT#M182). For phenocopy experiments (Figure 5), BMDMs were not

synchronized prior to the experiment. BMDMs were cultured in Atmospheric Media at pH 7.4 or 6.5 with

vehicle or 20, 40, or 80 µM IBMX (Sigma, CAT#I5879) or forskolin (Sigma, CAT#344270). For LumiCycle

experiments, 100mM D-luciferin was added to Atmospheric Media at 1:1000 for 100µM D-luciferin

(Promega, CAT#E1602). Cells cultured in Atmospheric Media were kept at 37^oC in atmospheric

conditions, and were either monitored over time by LumiCycle or harvested for RNA or protein at the

time points indicated.

Isolation and culture of peritoneal macrophages

Peritoneal exudate cells were harvested from mice as previously published[111]. To isolate peritoneal

macrophages, peritoneal exudate cells were seeded at $1.2*10^6$ cells/mL in RPMI/10% HI FBS

supplemented with 100U/mL Penicillin-Streptomycin and left at 37^oC for 1 hour, after which non-

adherent cells were rinsed off[110]. Peritoneal macrophages were then cultured in Atmospheric Media

at pH 7.4 or 6.5 with 100μM D-luciferin, and kept at 37^oC in atmospheric conditions.

Quantification of circadian rhythm parameters

Using the Lumicycle Analysis program version 2.701 (Actimetrics), raw data was fitted to a linear

baseline, and the baseline-subtracted data was fitted to a damped sin wave from which period and

damping were calculated[105]. Amplitude was calculated from baseline-subtracted data by subtracting

the bioluminescent values of the first peak from the first trough as previously published[64].

Quantitative PCR

Cells were lysed and RNA was isolated using the E.Z.N.A. HP Total RNA Kit (Omega BioTek, CAT#R6812-

02). RNA was reverse transcribed to cDNA using the ABI Reverse Transcription Reagents system, using

oligo dT for priming (Applied Biosystems, CAT#N8080234). qPCR was performed with cDNA using

PerfeCTa SYBR Green FastMix (QuantaBio, CAT#95074-05K) and with the Quant Studio 5 quantitative PCR

machines (Applied Biosystems). Triplicate technical replicates were performed, outlier replicates (defined

as being more than 1 Ct away from other two replicates) were discarded, and relative mRNA was

normalized to *Tbp* and assessed by the ΔΔCt. The following primers were used: *Tbp* (5'-

CCAGAACTGAAAATCAACGCAG-3', 5'-TGTATCTACCGTGAATCTTGGC-3') (IDT, Mm.PT.39a.22214839), *Arg1*

(5'-CTCCAAGCCAAAGTCCTTAGAG-3', 5'-AGGAGCTGTCATTAGGGACATC-3')[112], *Vegf* (5'-

CCACGACAGAAGGAGAGCAGAAGTCC-3', 5'-CGTTACAGCAGCCTGCACAGCG-3')[77], *Nos2* (5'-

GCTTCTGGTCGATGTCATGAG-3', 5'-TCCACCAGGAGATGTTGAAC-3'), *Icer* (5'-ATGGCTGTAACTGGAGATGAA-

3', 5'-GTGGCAAAGCAGTAGTAGGA-3')[29], *Per2* (5'-TGAGGTAGATAGCCCAGGAG-3,' 5'-

GCTATGAAGCGCCTAGAATCC-3') (IDT, Mm.PT.58.5594166), *Cry1* (5'-GCTATGCTCCTGGAGAGAACG T-3,' 5'-

TGTCCCCGTGAGCATAGTGTAA-3')[113], *Nr1d1* (5'-GAGCCACTAGAGCCAATGTAG-3,' 5'-

CCAGTTTGAATGACCGCTTTC-3') (IDT, Mm.PT.58.17472803).

Western blot

Cells were lysed using the M-Per lysis reagent (Thermo Scientific, CAT#78501), supplemented with

protease and phosphatase inhibitor cocktail (1:100; Sigma, CAT#PPC1010) and phosphatase inhibitor

cocktail 2 (1:50; Sigma, CAT#P5726), with 200μM deferoxamine (Sigma, CAT#D9533). Lysates were

incubated on ice for 1 hour, then centrifuged at 17,000 xg to pellet out debris; supernatant was

collected. Protein was quantified using the Bio-Rad DC Protein Assay Kit (Bio-Rad, CAT#5000112), and

lysates of equal concentration were prepared and run by SDS-PAGE on Bio-Rad Criterion 4–15% 26-well

gradient gel (Bio-Rad, CAT#5678095). Gels were transferred using the Trans-Blot Turbo system (Bio-Rad)

to nitrocellulose membranes (Bio-Rad CAT#1704271).

The following primary antibody was used: rabbit anti-p-CREB (Ser133, Ser129) (Invitrogen, CAT#44-

297G). The following secondary antibody was used: goat anti-rabbit Alexa Fluor 680 (Invitrogen,

CAT#A21109). Of note, two different anti-CREB antibodies were tested (Cell Signaling, CAT#9197 and

Invitrogen, CAT#35-0900) in combination with the appropriate secondary antibody, but neither revealed

bands at the correct molecular weight for CREB protein. Membranes were digitally imaged using a

ChemiDoc MP (Bio-Rad) and uniformly contrasted. Total protein was imaged by Stain-Free imaging

technology (Bio-Rad) and used as loading control. To visualize total protein, image of entire membrane

was shrunk to match the size of pCREB.

Survival under acidic pH

BMDMs were seeded, in triplicate, at $1.2*10^6$ cells/mL in a 96-well plate. BMDMs were synchronized,

then cultured in Atmospheric Media at pH 7.4, 6.8, or 6.5 containing 10 ng/mL IL-4 and 10 ng/mL IL-13,

or 50 ng/mL IFNγ and 100 ng/mL LPS, or vehicle for unstimulated control. BMDMs were fixed at 1, 2, and

3 days later. BMDMs were stained with DAPI, and plates were imaged using a Celigo S. Number of nuclei

per well was enumerated using Celigo software to quantify the number of adherent BMDMs after time in

culture under acidic conditions as a readout of survival.

Tumor growth

Mice were anesthetized via inhalation of 4 vol% isoflurane in 100% oxygen at a flow rate of 4 L/min prior

to injection. Following application of 70% ethanol to the site of injection, with $1*10^6$ WT or BMAL1 KO

macrophages and $1*10^6$ KCKO cells in 100 uL saline were subcutaneously co-injected in the flank of WT

mice. In line with previously published co-injection tumor experiments, mice were injected with

macrophages at a 1:1 ratio[77, 88]. Tumor growth was measured by caliper, and volume was calculated

by the modified ellipsoidal formula: tumor volume = $0.5*(length*width^2)$[114]. Mice were euthanized

when there was ulcer formation or when tumor size reached a diameter of 20mm.

Processing and analysis of publicly available gene expression data

FASTQ files from a previously published analysis of peritoneal macrophages from WT or BMAL1 KO mice

were downloaded from EMBL- European Bioinformatics Institute Array Express (accession #E-MTAB-

8411)[44]. For these samples, multiple files of the same run were concatenated before mapping. FASTQ

files from a previous published study of tumor-associated macrophages were downloaded from NCBI

GEO (accession #GSE188549)[83]. Both sets of samples were mapped to transcripts using Salmon 1.3.0

in mapping-based mode using a decoy-aware transcription built from the Gencode M25 GRCm38

primary assembly mouse genome and M25 mouse transcriptome[115]. Single-end mapping was used

for GSE188549 samples and paired-end mapping was used for E-MTAB-8411. All transcripts were then

collapsed to gene-level using Tximport v1.14.2with the Gencode M25 transcriptome, and genes were

annotated with symbols using the Ensembl GRCm38.100 transcriptome annotations[116]. Transcripts

per million (TPM) outputted from Tximport were used for downstream analyses. CEL files from a

previously published time series analysis of peritoneal macrophages from WT mice were downloaded

from NCBI GEO (accession #GSE25585)[36]. Microarray data was imported and analyzed from CEL files

using the packages affy and Limma, and genes were annotated with symbols using the University of

Michigan Brain Array Custom CDF v25.0 for the Mouse Gene 1.0 ST Array[117, 118]. Clock correlation

distance analysis was performed as previously described[84].

Data sharing

Raw data will be provided by uploading to FigShare, as will R code used for data analysis.

Acknowledgements

We would like to thank Dr. Jim Miller (Department of Microbiology and Immunology, University of

Rochester, NY) and Dr. Paul Brookes (Department of Anesthesiology and Perioperative Medicine,

University of Rochester, NY) for their intellectual contributions. All visual illustrations were made using

BioRender. The project described was supported by Award Number T32AI007285 (to AMKC) from the

National Institute of Allergy and Infectious Diseases, Grant Number T32 GM135134 (to AMKC) from the

National Institutes of Health, Grant Number R00CA204593 (to BJA) from the National Cancer Institute of

the National Institutes of Health, and the Wilmot Predoctoral Cancer Research Fellowship (to AMKC)

from the Wilmot Cancer Institute.

Author Contributions

Conceptualization: Amelia M. Knudsen-Clark, Brian J. Altman

Data Curation: Amelia M. Knudsen-Clark, Brian J. Altman

Formal analysis: Amelia M. Knudsen-Clark, Brian J. Altman, Lauren M. Hablitz

Funding acquisition: Brian J. Altman, Amelia M. Knudsen-Clark

Investigation: Amelia M. Knudsen-Clark, Juliana Cazarin, Daniel Mwangi

Methodology: Amelia M. Knudsen-Clark, Brian J. Altman, Minsoo Kim

Resources: Brian J. Altman

Software: Brian J. Altman, Amelia M. Knudsen-Clark, Daniel Mwangi

Supervision: Brian J. Altman

Validation: Amelia M. Knudsen-Clark

Writing – original draft: Amelia M. Knudsen-Clark

Writing – review and editing: Amelia M. Knudsen-Clark, Juliana Cazarin, Brian J. Altman, Daniel Mwangi,

Minsoo Kim, Lauren M. Hablitz.

Conflict of Interest

772   The authors declare no conflict of interest.

Figure Legends

Figure 1. Macrophages of different phenotypes have distinct circadian rhythms. Bone marrow-derived

macrophages (BMDMs) were obtained from C57BL/6 mice expressing PER2-Luc. The circadian clocks of

BMDMs were synchronized by a 24-hour period of serum starvation in media with 0% serum, followed

by a 2-hour period of serum shock in media with 50% serum. Luciferase activity of BMDMs stimulated

with either 10 ng/mL IL-4 and 10 ng/mL IL-13, or 50 ng/mL IFNγ and 100 ng/mL LPS; or left unstimulated

was monitored in real time by LumiCycle. Data was baseline-subtracted using the running average.

Oscillation parameters of BMDMs were measured by LumiCycle Analysis. Shown are mean and standard

error of the mean (SEM), n=5 biological replicates. Statistical significance determined by unpaired two-

tailed T test with Welch's correction; *, $p < 0.05$; **, $p<0.005$; ***, $p<0.0005$. Data shown are

representative of 2 independent experiments.

Figure 2. Acidic pH alters circadian rhythms of bone marrow-derived macrophages *in vitro*. Bone

marrow-derived macrophages (BMDMs) were obtained from C57BL/6 mice expressing PER2-Luc. The

circadian clocks of BMDMs were synchronized by a 24-hour period of serum starvation in media with 0%

serum, followed by a 2-hour period of serum shock in media with 50% serum. BMDMs were then

cultured in media with neutral pH 7.4 or acidic media with pH 6.8 or 6.5, and stimulated with either 10

ng/mL IL-4 and 10 ng/mL IL-13 (B, E), or 50 ng/mL IFNγ and 100 ng/mL LPS (C, F); or left unstimulated (A,

D). Luciferase activity was monitored in real time by LumiCycle (A, B, C). Shown are mean and SEM, n=2

biological replicates, representative of 2 independent experiments. Data from both experiments was

baseline-subtracted using the running average, and oscillation parameters were measured by LumiCycle

Analysis. Shown are mean and SEM, n=5 biological replicates. In parallel, RNA was collected at 12, 16, 20,

and 24 hours post-synchronization, and qt-PCR was performed to assess oscillation of transcripts

encoding core clock proteins in macrophages under acidic conditions (D, E, F). Shown are mean and SEM,

n=3 biological replicates. Data shown are representative of 2 independent experiments. Statistical

significance determined by unpaired two-tailed T test with Welch's correction; *, p < 0.05; **, p<0.005;

***, p<0.0005.

Figure 3. Acidic pH alters circadian rhythms of bone marrow-derived macrophages *ex vivo* at

temporally distinct times of day. Peritoneal macrophages were obtained at ZT0 or ZT12 from C57BL/6

mice expressing PER2-Luc and cultured in media with neutral pH 7.4 or acidic pH 6.5. Luciferase activity

was monitored in real time by LumiCycle (A). Shown are mean and SEM, n=2 biological replicates,

representative of 2 independent experiments. Data from both experiments was baseline-subtracted

using the running average, and oscillation parameters were measured by LumiCycle Analysis; shown are

mean and SEM, n=4 biological replicates. Statistical significance determined by unpaired two-tailed T test

with Welch's correction. The magnitude of change in circadian oscillation parameters between

macrophages at pH 7.4 and pH 6.5 was compared between peritoneal macrophages taken at ZT 0 or ZT

12 (B). Shown is the mean and SEM, n=4; data pooled from the 2 independent experiments in (A).

Statistical significance determined by paired two-tailed T test with Welch's correction; *, p < 0.05; **,

p<0.005; ***, p<0.0005.

Figure 4. Lactate alters circadian rhythms in macrophages, both alone and in conjunction with acidic

pH. Bone marrow-derived macrophages (BMDMs) were obtained from C57BL/6 mice expressing PER2-

Luc. The circadian clocks of BMDMs were synchronized by a 24-hour period of serum starvation in media

with 0% serum, followed by a 2-hour period of serum shock in media with 50% serum. BMDMs were

then cultured in media with neutral pH 7.4 or acidic pH 6.5, supplemented with 0 mM or 25 mM sodium-

L-lactate. RNA was collected at 6 hours post-treatment, and expression of pro-resolution phenotype

markers *Arg1* or *Vegf* was quantified by qtPCR (A). Shown is the mean and SEM, n=6 biological replicates,

data pooled from 2 independent experiments. Luciferase activity was monitored in real time by

LumiCycle (B). Shown is the mean and SEM, n=4 biological replicates; data representative of 2

independent experiments. Data from both experiments was baseline-subtracted using the running

average, and oscillation parameters were measured by LumiCycle Analysis. Shown is the mean and SEM,

n=7-10 biological replicates. Statistical significance determined by unpaired two-tailed T test with

Welch's correction; *, $p < 0.05$; **, $p<0.005$; ***, $p<0.0005$; ****, $p<0.00005$.

Figure 5. Treatment with Forskolin or IBMX does not fully recapitulate pH-driven changes in circadian

rhythms of macrophages. Bone marrow-derived macrophages (BMDMs) were obtained from C57BL/6

mice expressing PER2-Luc. BMDMs were cultured in media with neutral pH 7.4 or acidic pH 6.5, and

treated with vehicle or 20, 40 or 80 uM Forsokolin (A), or treated with vehicle or 20, 40, 80 uM IBMX (B).

Luciferase activity was monitored in real time by LumiCycle. Data was baseline-subtracted using the

running average. Shown is the mean, n=2 biological replicates. Data shown are representative of 2

independent experiments.

Figure 6. Treatment with MDL-12 suppresses pH-induced changes in amplitude of circadian rhythms

without suppressing signaling through the cAMP signaling pathway under acidic conditions. Bone

marrow-derived macrophages (BMDMs) were obtained from C57BL/6 mice expressing PER2-Luc. The

circadian clocks of BMDMs were synchronized by a 24-hour period of serum starvation in media with 0%

serum, followed by a 2-hour period of serum shock in media with 50% serum. BMDMs were then

cultured in media with neutral pH 7.4 or acidic pH 6.5, and treated with vehicle or 5, 10, or 15uM MDL-

12. Luciferase activity was monitored in real time by LumiCycle. Data was baseline-subtracted using the

running average, and oscillation parameters were measured by LumiCycle Analysis (A). Shown is the

mean, n=4 biological replicates. Expression of genes associated with acid sensing (*Icer*) and pro-

resolution phenotype (*Arg1*) was measured (B). Shown is the mean and SEM; n=3 biological replicates.

Phosphorylation of CREB was assessed (C). Statistical significance determined by unpaired two-tailed T

test with Welch's correction; *, $p < 0.05$; **, $p<0.005$; ***, $p<0.0005$; ****, $p<0.00005$. Data shown are

representative of 2-3 independent experiments.

Figure 7. Clock correlation distance (CCD) analysis provides evidence of circadian disorder in tumor-

associated macrophages. RNAseq datasets of WT peritoneal macrophages (n=5), BMAL1 KO peritoneal

macrophages (n=4), and tumor-associated macrophages (TAMs, n=10) were analyzed for expression of

Arg1 and *Crem* (A). Statistical significance determined by one-way ANOVA with Turkey's multiple

comparison test; ****, $p<0.00005$. Clock correlation distance (CCD) analysis was performed (B) and

statistical analysis to compare CCD scores was performed by calculating delta CCD; *, $p<0.05$ (C).

Figure 8. Heterogeneity in circadian rhythms of cells within a population can lead to circadian disorder

observed by CCD. Increasingly desynchronized populations were modeled using an RNAseq data set of

WT peritoneal macrophages (n=12) taken at 4-hour intervals across two days. A schematic of the

populations used in experimental design (A). Clock correlation distance (CCD) analysis was performed (B)

and statistical analysis to compare CCD scores was performed by calculating delta CCD (C).

Figure 9. A functional circadian clock in macrophages can influence tumor growth in a murine model of

PDAC. Bone marrow-derived macrophages (BMDMs) obtained from WT or BMAL1 KO mice were

subcutaneously co-injected with KCKO cells into the flank of WT mice. Tumor growth was measured by

caliper. Shown is the mean and SEM; n=20 individual mice, 10 male and 10 female. Statistical significance

determined at each time point by unpaired two-tailed T test with Welch's correction; *, $p < 0.05$; **,

$p<0.005$; ***, $p<0.0005$. Data shown are representative of 2 independent experiments.

Supplemental Figure 1. Polarization of macrophages toward pro-resolution or pro-inflammatory

phenotype. Bone marrow-derived macrophages (BMDMs) were derived from C57BL/6 mice expressing

PER2-Luc. The circadian clocks of BMDMs were synchronized by a 24-hour period of serum starvation in

media with 0% serum, followed by a 2-hour period of serum shock in media with 50% serum. BMDMs

were then stimulated with either 10 ng/mL IL-4 and 10 ng/mL IL-13, or 50 ng/mL IFNγ and 100 ng/mL

LPS; or left unstimulated. RNA was collected at 6 hours post-synchronization, and qt-PCR was performed

to assess expression of genes encoding Arginase 1 (*Arg1*) and iNOS (*Nos2*). Shown are mean and

standard error of the mean (SEM), n=3 biological replicates. Statistical significance determined by

unpaired two-tailed T test with Welch's correction; *, $p < 0.05$; **, ***, $p<0.0005$. Data shown are

representative of 2 independent experiments.

Supplemental Figure 2. The PER2-Luciferase reporter system enables real-time monitoring of circadian

rhythms of macrophages. A schematic of the PER2-Luciferase (PER2-Luc) luciferase reporter system (A).

Bone marrow-derived macrophages (BMDMs) were derived from C57BL/6 mice expressing PER2-Luc.

The circadian clocks of BMDMs were synchronized by a 24-hour period of serum starvation in media with

0% serum, followed by a 2-hour period of serum shock in media with 50% serum. BMDMs were then

cultured in RPMI/10% FBS supplemented with D-luciferin, and luciferase activity was monitored in real

time by LumiCycle (B). Shown are mean and SEM; n=2 biological replicates. Data shown are

representative of 2 independent experiments.

Supplemental Figure 3. Macrophages sense and respond to an acidic extracellular environment when

cultured *in vitro* in media with acidic pH. Bone marrow-derived macrophages (BMDMs) were obtained

from C57BL/6 mice expressing PER2-Luc. BMDMs were cultured in media with pH 7.4 or acidic media

with pH 6.5, and stimulated with either 10 ng/mL IL-4 and 10 ng/mL IL-13, or 50 ng/mL IFNγ and 100

ng/mL LPS; or left unstimulated. RNA was collected at 2 hours post-treatment, and qt-PCR was

performed to assess expression of genes associated with phenotype (A) or acid sensing (B) in

macrophages. Shown are mean and SEM, n=3 biological replicates. Statistical significance determined by

two-tailed T test with Welch's correction; *, p < 0.05; **, p<0.005; ***, p<0.0005; ****, p<0.00005. Data

shown are representative of 2 independent experiments.

Supplementary Figure 4. Survival of macrophages under acidic pH. Bone marrow-derived macrophages

(BMDMs) were obtained from C57BL/6 mice expressing PER2-Luc. The circadian clocks of BMDMs were

synchronized by a 24-hour period of serum starvation in media with 0% serum, followed by a 2-hour

period of serum shock in media with 50% serum. BMDMs were then cultured in media with pH 7.4 or

acidic media with pH 6.8 or 6.5, and stimulated with either 10 ng/mL IL-4 and 10 ng/mL IL-13, or 50

ng/mL IFNγ and 100 ng/mL LPS; or left unstimulated. Cells were fixed at 1, 2, and 3 days post-treatment

and stained with DAPI. Number of nuclei was counted using Celigo to determine the number of adherent

cells. Shown are mean and SEM, n=3 biological replicates. Statistical significance determined by multiple

unpaired T test with Welch's correction; *, p < 0.05; **, p<0.005; ***, p<0.0005. Data shown are

representative of 2 independent experiments.

Supplementary Figure 5. Pre-treatment with 15 uM MDL-12 for up to 2 hours is not necessary to

suppress pH-driven changes in amplitude of circadian rhythms. Bone marrow-derived macrophages

(BMDMs) were obtained from C57BL/6 mice expressing PER2-Luc. The circadian clocks of BMDMs were

synchronized by a 24-hour period of serum starvation in media with 0% serum, followed by a 2-hour

period of serum shock in media with 50% serum. BMDMs were then cultured in media with neutral pH

7.4 or acidic pH 6.5, and treated with vehicle or 15uM MDL-12 (A). Alternately, BMDMs were cultured at

pH 7.4 and pre-treated with 15uM MDL-12 for 30 minutes (B, C) or 2 hours (D, E), after which media was

removed and cells were cultured in media at pH 7.4 or pH 6.5 in the presence (C, E) or absence (B, D) of

MDL-12. Luciferase activity was monitored in real time by LumiCycle. Shown is the mean, n=2 biological

replicates. Data for cells that received no pre-treatment (A) was overlaid on plots of data where cells

received pre-treatment (B-E) to allow for comparison of changes in rhythms.

Supplementary Figure 6. Macrophages from BMAL1 KO mice have disrupted circadian rhythms. Levels

of BMAL1 in bone marrow-derived macrophages (BMDMs) from WT or BMAL1 KO mice were assessed

by western blot (A). To confirm functional disruption of the circadian clock, peritoneal macrophages or

BMDMs were obtained from WT or BMAL1 KO mice expressing PER2-Luc and cultured in vitro with D-

luciferin. Luciferase activity was monitored in real time by LumiCycle (B). Shown is the mean and SEM,

n=2, representative of 2 independent experiments.

References

1.   Cassetta, L. and J.W. Pollard, *Targeting macrophages: therapeutic approaches in cancer.* Nat Rev
Drug Discov, 2018. **17**(12): p. 887-904.
2.   Zhang, Q.W., et al., *Prognostic significance of tumor-associated macrophages in solid tumor: a*
meta-analysis of the literature. PLoS One, 2012. **7**(12): p. e50946.
3.   Gentles, A.J., et al., *The prognostic landscape of genes and infiltrating immune cells across*
human cancers. Nat Med, 2015. **21**(8): p. 938-945.
4.   DeNardo, D.G. and B. Ruffell, *Macrophages as regulators of tumour immunity and*
immunotherapy. Nat Rev Immunol, 2019.
5.   Mosser, D.M. and J.P. Edwards, *Exploring the full spectrum of macrophage activation.* Nat Rev
Immunol, 2008. **8**(12): p. 958-69.
6.   Murray, P.J. and T.A. Wynn, *Protective and pathogenic functions of macrophage subsets.* Nat Rev
Immunol, 2011. **11**(11): p. 723-37.
7.   Coussens, L.M. and Z. Werb, *Inflammation and cancer.* Nature, 2002. **420**(6917): p. 860-7.
8.   Murray, P.J., *Nonresolving macrophage-mediated inflammation in malignancy.* Febs j, 2018.
285(4): p. 641-653.
9.   Schoppmann, S.F., et al., *Tumor-associated macrophages express lymphatic endothelial growth*
factors and are related to peritumoral lymphangiogenesis. Am J Pathol, 2002. **161**(3): p. 947-56.
10.   Huber, R., et al., *Tumour hypoxia promotes melanoma growth and metastasis via High Mobility*
Group Box-1 and M2-like macrophages. Sci Rep, 2016. **6**: p. 29914.
11.   Graham, D.K., et al., *The TAM family: phosphatidylserine sensing receptor tyrosine kinases gone*
awry in cancer. Nat Rev Cancer, 2014. **14**(12): p. 769-85.
12.   Cook, R.S., et al., *MerTK inhibition in tumor leukocytes decreases tumor growth and metastasis.* J
Clin Invest, 2013. **123**(8): p. 3231-42.
13.   Roberts, A.W., et al., *Tissue-Resident Macrophages Are Locally Programmed for Silent Clearance*
of Apoptotic Cells. Immunity, 2017. **47**(5): p. 913-927.e6.
14.   Fadok, V.A., et al., *Macrophages that have ingested apoptotic cells in vitro inhibit*
proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF-
beta, PGE2, and PAF. J Clin Invest, 1998. **101**(4): p. 890-8.
15.   Strassmann, G., et al., *Evidence for the involvement of interleukin 10 in the differential*
deactivation of murine peritoneal macrophages by prostaglandin E2. J Exp Med, 1994. **180**(6): p.
2365-70.
16.   Kuang, D.M., et al., *Activated monocytes in peritumoral stroma of hepatocellular carcinoma*
foster immune privilege and disease progression through PD-L1. J Exp Med, 2009. **206**(6): p.
1327-37.
17.   Ruffell, B. and L.M. Coussens, *Macrophages and therapeutic resistance in cancer.* Cancer Cell,
2015. **27**(4): p. 462-72.
18.   Mantovani, A., et al., *Macrophages as tools and targets in cancer therapy.* Nature Reviews Drug
Discovery, 2022. **21**(11): p. 799-820.
19.   Cuccarese, M.F., et al., *Heterogeneity of macrophage infiltration and therapeutic response in*
lung carcinoma revealed by 3D organ imaging. Nat Commun, 2017. **8**: p. 14293.
20.   Chevrier, S., et al., *An Immune Atlas of Clear Cell Renal Cell Carcinoma.* Cell, 2017. **169**(4): p. 736-
749.e18.

21.   Laviron, M., et al., *Tumor-associated macrophage heterogeneity is driven by tissue territories in*
breast cancer. Cell Rep, 2022. **39**(8): p. 110865.

22.   Nalio Ramos, R., et al., *Tissue-resident FOLR2(+) macrophages associate with CD8(+) T cell*
infiltration in human breast cancer. Cell, 2022. **185**(7): p. 1189-1207.e25.

23.   Yano, S., et al., *Spatial-temporal FUCCI imaging of each cell in a tumor demonstrates locational*
dependence of cell cycle dynamics and chemoresponsiveness. Cell Cycle, 2014. **13**(13): p. 2110-9.

24.   Huang, Y.K., et al., *Macrophage spatial heterogeneity in gastric cancer defined by multiplex*
immunohistochemistry. Nat Commun, 2019. **10**(1): p. 3928.

25.   Lyssiotis, C.A. and A.C. Kimmelman, *Metabolic Interactions in the Tumor Microenvironment.*
Trends Cell Biol, 2017. **27**(11): p. 863-875.

26.   Raghunand, N., R.A. Gatenby, and R.J. Gillies, *Microenvironmental and cellular consequences of*
altered blood flow in tumours. Br J Radiol, 2003. **76 Spec No 1**: p. S11-22.

27.   Gillies, R.J., et al., *Targeting acidity in cancer and diabetes.* Biochim Biophys Acta Rev Cancer,
2019. **1871**(2): p. 273-280.

28.   El-Kenawi, A., et al., *Acidity promotes tumour progression by altering macrophage phenotype in*
prostate cancer. Br J Cancer, 2019. **121**(7): p. 556-566.

29.   Bohn, T., et al., *Tumor immunoevasion via acidosis-dependent induction of regulatory tumor-*
associated macrophages. Nat Immunol, 2018. **19**(12): p. 1319-1329.

30.   Jiang, W., et al., *Extracellular Acidity Reprograms Macrophage Metabolism and Innate*
Responsiveness. The Journal of Immunology, 2021. **206**(12): p. 3021-3031.

31.   Partch, C.L., C.B. Green, and J.S. Takahashi, *Molecular architecture of the mammalian circadian*
clock. Trends Cell Biol, 2014. **24**(2): p. 90-9.

32.   Walton, Z.E., et al., *Acid Suspends the Circadian Clock in Hypoxia through Inhibition of mTOR.*
Cell, 2018. **174**(1): p. 72-87.e32.

33.   Mure, L.S., et al., *Diurnal transcriptome atlas of a primate across major neural and peripheral*
tissues. Science, 2018. **359**(6381).

34.   Zhang, R., et al., *A circadian gene expression atlas in mammals: implications for biology and*
medicine. Proc Natl Acad Sci U S A, 2014. **111**(45): p. 16219-24.

35.   Haspel, J.A., et al., *Perfect timing: circadian rhythms, sleep, and immunity - an NIH workshop*
summary. JCI Insight, 2020. **5**(1).

36.   Keller, M., et al., *A circadian clock in macrophages controls inflammatory immune responses.*
Proc Natl Acad Sci U S A, 2009. **106**(50): p. 21407-12.

37.   Silver, A.C., et al., *Circadian expression of clock genes in mouse macrophages, dendritic cells, and*
B cells. Brain Behav Immun, 2012. **26**(3): p. 407-13.

38.   Nguyen, K.D., et al., *Circadian gene Bmal1 regulates diurnal oscillations of Ly6C(hi) inflammatory*
monocytes. Science, 2013. **341**(6153): p. 1483-8.

39.   Baumann, A., et al., *The circadian clock is functional in eosinophils and mast cells.* Immunology,
2013. **140**(4): p. 465-74.

40.   Adrover, J.M., et al., *A Neutrophil Timer Coordinates Immune Defense and Vascular Protection.*
Immunity, 2019. **50**(2): p. 390-402.e10.

41.   Arjona, A. and D.K. Sarkar, *Circadian oscillations of clock genes, cytolytic factors, and cytokines in*
rat NK cells. J Immunol, 2005. **174**(12): p. 7618-24.

42.   Druzd, D., et al., *Lymphocyte Circadian Clocks Control Lymph Node Trafficking and Adaptive*
Immune Responses. Immunity, 2017. **46**(1): p. 120-132.

43.   Scheiermann, C., et al., *Clocking in to immunity.* Nat Rev Immunol, 2018. **18**(7): p. 423-437.

44.   Kitchen, G.B., et al., *The clock gene Bmal1 inhibits macrophage motility, phagocytosis, and*
impairs defense against pneumonia. Proc Natl Acad Sci U S A, 2020.

45.   Gibbs, J.E., et al., *The nuclear receptor REV-ERBalpha mediates circadian regulation of innate*
immunity through selective regulation of inflammatory cytokines. Proc Natl Acad Sci U S A, 2012.
109(2): p. 582-7.
46.   He, W., et al., *Circadian Expression of Migratory Factors Establishes Lineage-Specific Signatures*
that Guide the Homing of Leukocyte Subsets to Tissues. Immunity, 2018. **49**(6): p. 1175-1190.e7.
47.   Nobis, C.C., et al., *The circadian clock of CD8 T cells modulates their early response to vaccination*
and the rhythmicity of related signaling pathways. Proc Natl Acad Sci U S A, 2019. **116**(40): p.
20077-20086.
48.   Fortier, E.E., et al., *Circadian variation of the response of T cells to antigen.* J Immunol, 2011.
187(12): p. 6291-300.
49.   Beam, C.A., et al., *Synchronization of the Normal Human Peripheral Immune System: A*
Comprehensive Circadian Systems Immunology Analysis. Sci Rep, 2020. **10**(1): p. 672.
50.   Mazzoccoli, G., et al., *A timetable of 24-hour patterns for human lymphocyte subpopulations.* J
Biol Regul Homeost Agents, 2011. **25**(3): p. 387-95.
51.   Silver, A.C., et al., *The circadian clock controls toll-like receptor 9-mediated innate and adaptive*
immunity. Immunity, 2012. **36**(2): p. 251-61.
52.   Phillips, A.C., et al., *Preliminary evidence that morning vaccination is associated with an*
enhanced antibody response in men. Psychophysiology, 2008. **45**(4): p. 663-6.
53.   Long, J.E., et al., *Morning vaccination enhances antibody response over afternoon vaccination: A*
cluster-randomised trial. Vaccine, 2016. **34**(24): p. 2679-85.
54.   Curtis, A.M., et al., *Circadian control of innate immunity in macrophages by miR-155 targeting*
Bmal1. Proc Natl Acad Sci U S A, 2015. **112**(23): p. 7231-6.
55.   Deng, W., et al., *The Circadian Clock Controls Immune Checkpoint Pathway in Sepsis.* Cell Rep,
2018. **24**(2): p. 366-378.
56.   Aiello, I., et al., *Circadian disruption promotes tumor-immune microenvironment remodeling*
favoring tumor cell proliferation. Sci Adv, 2020. **6**(42).
57.   Tsuruta, A., et al., *Diurnal Expression of PD-1 on Tumor-Associated Macrophages Underlies the*
Dosing Time-Dependent Antitumor Effects of the PD-1/PD-L1 Inhibitor BMS-1 in B16/BL6
Melanoma-Bearing Mice. Mol Cancer Res, 2022. **20**(6): p. 972-982.
58.   Strauss, L., et al., *Targeted deletion of PD-1 in myeloid cells induces antitumor immunity.* Sci
Immunol, 2020. **5**(43).
59.   Qian, D.C., et al., *Effect of immunotherapy time-of-day infusion on overall survival among*
patients with advanced melanoma in the USA (MEMOIR): a propensity score-matched analysis of
a single-centre, longitudinal study. Lancet Oncol, 2021. **22**(12): p. 1777-1786.
60.   Martinez, F.O. and S. Gordon, *The M1 and M2 paradigm of macrophage activation: time for*
reassessment. F1000Prime Rep, 2014. **6**: p. 13.
61.   Yoo, S.H., et al., *PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals*
persistent circadian oscillations in mouse peripheral tissues. Proc Natl Acad Sci U S A, 2004.
101(15): p. 5339-46.
62.   Rodheim, K., C. Jung, and K. Wright, *016 Associations between Circadian Melatonin and*
Temperature Amplitudes during Constant Routine. Sleep, 2021. **44**(Supplement_2): p. A8-A8.
63.   Wu, G., et al., *Normalized coefficient of variation (nCV): a method to evaluate circadian clock*
robustness in population scale data. Bioinformatics, 2021. **37**(23): p. 4581-4583.
64.   Chen, S., et al., *A Pro- and Anti-inflammatory Axis Modulates the Macrophage Circadian Clock.*
Front Immunol, 2020. **11**: p. 867.
65.   Lellupitiyage Don, S.S., et al., *Macrophage circadian rhythms are differentially affected based on*
stimuli. Integr Biol (Camb), 2022. **14**(3): p. 62-75.

66. Pilorz, V., et al., *A Novel Mechanism Controlling Resetting Speed of the Circadian Clock to*
Environmental Stimuli. Current Biology, 2014. **24**(7): p. 766-773.
67. Abe, M., et al., *Circadian rhythms in isolated brain regions.* J Neurosci, 2002. **22**(1): p. 350-6.
68. Gaspar, L.S., et al., *The importance of determining circadian parameters in pharmacological*
studies. British Journal of Pharmacology, 2019. **176**(16): p. 2827-2847.
69. Boedtkjer, E. and S.F. Pedersen, *The Acidic Tumor Microenvironment as a Driver of Cancer.*
Annual Review of Physiology, 2020. **82**(1): p. 103-126.
70. Gillies, R.J., Z. Liu, and Z. Bhujwalla, *31P-MRS measurements of extracellular pH of tumors using*
3-aminopropylphosphonate. Am J Physiol, 1994. **267**(1 Pt 1): p. C195-203.
71. Estrella, V., et al., *Acidity Generated by the Tumor Microenvironment Drives Local Invasion.*
Cancer Research, 2013. **73**(5): p. 1524-1535.
72. Lv, S., et al., *A negative feedback loop of ICER and NF-κB regulates TLR signaling in innate*
immune responses. Cell Death Differ, 2017. **24**(3): p. 492-499.
73. Kraut, J.A. and N.E. Madias, *Lactic acidosis.* N Engl J Med, 2014. **371**(24): p. 2309-19.
74. Pastorekova, S. and R.J. Gillies, *The role of carbonic anhydrase IX in cancer development: links to*
hypoxia, acidosis, and beyond. Cancer Metastasis Rev, 2019. **38**(1-2): p. 65-77.
75. Swietach, P., *What is pH regulation, and why do cancer cells need it?* Cancer Metastasis Rev,
2019. **38**(1-2): p. 5-15.
76. de la Cruz-López, K.G., et al., *Lactate in the Regulation of Tumor Microenvironment and*
Therapeutic Approaches. Front Oncol, 2019. **9**: p. 1143.
77. Colegio, O.R., et al., *Functional polarization of tumour-associated macrophages by tumour-*
derived lactic acid. Nature, 2014. **513**(7519): p. 559-63.
78. Polumuri, S., D.J. Perkins, and S.N. Vogel, *cAMP levels regulate macrophage alternative*
activation marker expression. Innate Immun, 2021. **27**(2): p. 133-142.
79. Tavares, L.P., et al., *Blame the signaling: Role of cAMP for the resolution of inflammation.*
Pharmacol Res, 2020. **159**: p. 105030.
80. Yagita, K. and H. Okamura, *Forskolin induces circadian gene expression of rPer1, rPer2 and dbp in*
mammalian rat-1 fibroblasts. FEBS Lett, 2000. **465**(1): p. 79-82.
81. O'Neill, J.S., et al., *cAMP-dependent signaling as a core component of the mammalian circadian*
pacemaker. Science, 2008. **320**(5878): p. 949-53.
82. Misra, U.K. and S.V. Pizzo, *Coordinate regulation of forskolin-induced cellular proliferation in*
macrophages by protein kinase A/cAMP-response element-binding protein (CREB) and Epac1-
Rap1 signaling: effects of silencing CREB gene expression on Akt activation. J Biol Chem, 2005.
280(46): p. 38276-89.
83. Geeraerts, X., et al., *Macrophage Metabolism As Therapeutic Target for Cancer, Atherosclerosis,*
and Obesity. Front Immunol, 2017. **8**: p. 289.
84. Shilts, J., G. Chen, and J.J. Hughey, *Evidence for widespread dysregulation of circadian clock*
progression in human cancer. PeerJ, 2018. **6**: p. e4327.
85. Pariollaud, M., et al., *Circadian disruption enhances HSF1 signaling and tumorigenesis in Kras-*
driven lung cancer. Science Advances, 2022. **8**(39): p. eabo1123.
86. Besmer, D.M., et al., *Pancreatic ductal adenocarcinoma mice lacking mucin 1 have a profound*
defect in tumor growth and metastasis. Cancer Res, 2011. **71**(13): p. 4432-42.
87. Kidiyoor, A., et al., *Pancreatic Cancer Cells Isolated from Muc1-Null Tumors Favor the Generation*
of a Mature Less Suppressive MDSC Population. Front Immunol, 2014. **5**: p. 67.
88. Alexander, R.K., et al., *Bmal1 integrates mitochondrial metabolism and macrophage activation.*
eLife, 2020. **9**: p. e54090.
89. Lee, C.-C., et al., *Macrophage-secreted interleukin-35 regulates cancer cell plasticity to facilitate*
metastatic colonization. Nature Communications, 2018. **9**(1): p. 3763.

90.   Walton, Z.E., R.C. Brooks, and C.V. Dang, *mTOR Senses Intracellular pH through Lysosome*
1117   *Dispersion from RHEB.* Bioessays, 2019: p. e1800265.
1118   91.   Guellaen, G., et al., *RMI 12330 A, an inhibitor of adenylate cyclase in rat liver.* Biochim Biophys
1119   Acta, 1977. **484**(2): p. 465-75.
1120   92.   Hunt, N.H. and T. Evans, *RMI 12330A, an inhibitor of cyclic nucleotide phosphodiesterases and*
1121   *adenylate cyclase in kidney preparations.* Biochim Biophys Acta, 1980. **613**(2): p. 499-506.
1122   93.   van Rossum, D.B., et al., *Ca2+ entry mediated by store depletion, S-nitrosylation, and TRP3*
1123   *channels. Comparison of coupling and function.* J Biol Chem, 2000. **275**(37): p. 28562-8.
1124   94.   O'Neill, J.S. and A.B. Reddy, *The essential role of cAMP/Ca2+ signalling in mammalian circadian*
1125   *timekeeping.* Biochem Soc Trans, 2012. **40**(1): p. 44-50.
1126   95.   Ni, L., et al., *Identification and Function of Acid-sensing Ion Channels in RAW 264.7 Macrophage*
1127   *Cells.* Curr Med Sci, 2018. **38**(3): p. 436-442.
1128   96.   Selezneva, A., A.J. Gibb, and D. Willis, *The contribution of ion channels to shaping macrophage*
1129   *behaviour.* Front Pharmacol, 2022. **13**: p. 970234.
1130   97.   Putker, M., et al., *Mammalian Circadian Period, But Not Phase and Amplitude, Is Robust Against*
1131   *Redox and Metabolic Perturbations.* Antioxidants & Redox Signaling, 2017. **28**(7): p. 507-520.
1132   98.   Viola, A., et al., *The Metabolic Signature of Macrophage Responses.* Front Immunol, 2019. **10**: p.
1133   1462.
1134   99.   Morris, E.L., et al., *Single-cell transcriptomics of suprachiasmatic nuclei reveal a Prokineticin-*
1135   *driven circadian network.* Embo j, 2021. **40**(20): p. e108614.
1136   100.   Ma, D., et al., *A transcriptomic taxonomy of Drosophila circadian neurons around the clock.* Elife,
1137   2021. **10**.
1138   101.   Auerbach, B.J., G.A. FitzGerald, and M. Li, *Tempo: an unsupervised Bayesian algorithm for*
1139   *circadian phase inference in single-cell transcriptomics.* Nat Commun, 2022. **13**(1): p. 6580.
1140   102.   Finger, A.M., et al., *Intercellular coupling between peripheral circadian oscillators by TGF-β*
1141   *signaling.* Sci Adv, 2021. **7**(30).
1142   103.   Massagué, J., *TGFb in Cancer.* Cell, 2008. **134**(2): p. 215-230.
1143   104.   Geiger, S.S., et al., *Daily variation in macrophage phagocytosis is clock-independent and*
1144   *dispensable for cytokine production.* Immunology, 2019.
1145   105.   Ramanathan, C., et al., *Cell Type-Specific Functions of Period Genes Revealed by Novel Adipocyte*
1146   *and Hepatocyte Circadian Clock Models.* PLOS Genetics, 2014. **10**(4): p. e1004244.
1147   106.   Committee, N.R.C., *Guide for the Care and Use of Laboratory Animals*, in *Guide for the Care and*
1148   *Use of Laboratory Animals*. 2011, National Academies Press (US)
1149   Copyright © 2011, National Academy of Sciences.: Washington (DC).
1150   107.   Clausen, B.E., et al., *Conditional gene targeting in macrophages and granulocytes using LysMcre*
1151   *mice.* Transgenic Res, 1999. **8**(4): p. 265-77.
1152   108.   Storch, K.F., et al., *Intrinsic circadian clock of the mammalian retina: importance for retinal*
1153   *processing of visual information.* Cell, 2007. **130**(4): p. 730-741.
1154   109.   Trouplin, V., et al., *Bone marrow-derived macrophage production.* J Vis Exp, 2013(81): p. e50966.
1155   110.   Gonçalves, R. and D.M. Mosser, *The Isolation and Characterization of Murine Macrophages.*
1156   Current Protocols in Immunology, 2015. **111**(1): p. 14.1.1-14.1.16.
1157   111.   Ray, A. and B.N. Dittel, *Isolation of mouse peritoneal cavity cells.* J Vis Exp, 2010(35).
1158   112.   Liu, P.S., et al., *alpha-ketoglutarate orchestrates macrophage activation through metabolic and*
1159   *epigenetic reprogramming.* Nat Immunol, 2017. **18**(9): p. 985-994.
1160   113.   Huber, A.L., et al., *CRY2 and FBXL3 Cooperatively Degrade c-MYC.* Mol Cell, 2016. **64**(4): p. 774-
1161   789.
1162   114.   Tomayko, M.M. and C.P. Reynolds, *Determination of subcutaneous tumor size in athymic (nude)*
1163   *mice.* Cancer Chemother Pharmacol, 1989. **24**(3): p. 148-54.

1164   115.   Patro, R., et al., *Salmon provides fast and bias-aware quantification of transcript expression.* Nat
1165   Methods, 2017. **14**(4): p. 417-419.
1166   116.   Soneson, C., M.I. Love, and M.D. Robinson, *Differential analyses for RNA-seq: transcript-level*
1167   *estimates improve gene-level inferences.* F1000Res, 2015. **4**: p. 1521.
1168   117.   Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-sequencing and*
1169   *microarray studies.* Nucleic Acids Res, 2015. **43**(7): p. e47.
1170   118.   Dai, M., et al., *Evolving gene/transcript definitions significantly alter the interpretation of*
1171   *GeneChip data.* Nucleic Acids Res, 2005. **33**(20): p. e175.
1172

Fig 1

Fig 2

Fig 3

A

B

Fig 4

A

B

Fig 5

A

Forskolin Treatment

Legend	
—— pH7.4_veh_Forskolin	···· pH6.5_veh_Forskolin
—— pH7.4_20uM_Forskolin	···· pH6.5_20uM_Forskolin
—— pH7.4_40uM_Forskolin	···· pH6.5_40uM_Forskolin
—— pH7.4_80uM_Forskolin	···· pH6.5_80uM_Forskolin

B

IBMX Treatment

Legend	
—— pH7.4_veh_IBMX	···· pH6.5_veh_IBMX
—— pH7.4_20uM_IBMX	···· pH6.5_20uM_IBMX
—— pH7.4_40uM_IBMX	···· pH6.5_40uM_IBMX
—— pH7.4_80uM_IBMX	···· pH6.5_80uM_IBMX

Fig 6

A

Amplitude / Period / Damping graphs

pH7.4, vehicle — black solid
pH7.4, 5uM MDL-12 — red solid
pH7.4, 10uM MDL-12 — orange solid
pH7.4, 15uM MDL-12 — yellow solid

pH6.5, vehicle — black dotted
pH6.5, 5uM MDL-12 — red dotted
pH6.5, 10uM MDL-12 — orange dotted
pH6.5, 15uM MDL-12 — yellow dotted

■ pH 7.4   ■ pH 6.5

B

Icer   *Arg1*

■ pH7.4, vehicle   ■ pH6.5, vehicle
☐ pH7.4, 15uM MDL-12 ☐ pH6.5, 15uM MDL-12

C

| | pH 7.4 0 uM MDL-12 | pH 7.4 15 uM MDL-12 | pH 6.5 0 uM MDL-12 | pH 6.5 15 uM MDL-12 |

CT (min) 0 10 60 120 360 ...

kDa 50
37

p-CREB (43 kDa)

Total Protein

Fig 7

Fig 8

A

Synchronized	4-Hour Desynchronized	8-Hour Desynchronized	12-Hour Desynchronized

Population

Average

B

Synchronized	4-Hour Desynchronized	8-Hour Desynchronized	12-Hour Desynchronized

Core Clock Genes (y-axis): Arntl, Npas2, Clock, Cry1, Cry2, Nr1d1, Nr1d2, Per1, Per2, Per3, Dbp, Tef

Core Clock Genes (x-axis): Arntl, Npas2, Clock, Cry1, Cry2, Nr1d1, Nr1d2, Per1, Per2, Per3, Dbp, Tef

Co-Expression Value

1.0
0.5
0.0
-0.5
-1.0

C

More Disordered

More Ordered

CCD Score

*

Synchronized, 4hr Desynchronized, 8hr Desynchronized, 12hr Desynchronized

Fig 9