The ELF3-BBX24/BBX25-PIF4 module controls thermosensory growth in Arabidopsis

Bidhan Chandra Malakar[1], Shivani Singh[1], Vikas Garhwal[1,#], Rajanesh Chandramohan[1,#], Gouranga Upadhyaya[1], and Sreeramaiah N. Gangappa[1,2,*]

Affiliations:

[1]Department of Biological Sciences, Indian Institute of Science Education and Research Kolkata, Mohanpur 741246, West Bengal, India

[2]Centre for Climate and Environmental Studies, Indian Institute of Science Education and Research Kolkata, Mohanpur 741246, West Bengal, India

*Author for correspondence: ngsreeram@iiserkol.ac.in (S.N.G)

[#]Authors contributed equally

The author responsible for the distribution of data and materials integral to the findings presented in this article as per the policy described in the Instructions for Authors is Sreeramaiah N. Gangappa (ngsreeram@iiserkol.ac.in)

ABSTRACT

Temperature serves as a crucial environmental cue governing the growth and
adaptation of plants in their natural habitat. PHYTOCHROME INTERACTING
FACTOR 4 (PIF4) is a central regulator that promotes thermomorphogenesis in
Arabidopsis. Understanding its precise regulation is critical for optimal
thermomorphogenic growth. Here, we identified two BBX proteins, BBX24 and
BBX25, as novel components of the PIF4-mediated thermosensory pathway and act
to promote warm temperature-mediated growth. The *bbx24* and *bbx25* single and
double mutants showed moderate to strong temperature-insensitive hypocotyl and
cotyledon growth. Warm temperature induces *BBX24* and *BBX25* mRNA expression
and protein accumulation. Genetic and biochemical analysis revealed that
BBX24/BBX25 promotes PIF4-mediated thermosensory growth by counteracting a
key component of the evening complex, ELF3. While ELF3 inhibits *BBX24*/*BBX25*
gene expression at low ambient temperatures in the evening, warm temperature-
mediated inhibition of ELF3 activity results in enhanced BBX24/BBX25 activity.
Moreover, BBX24/25 inhibit ELF3 function through direct physical interaction and
likely relieves repression on PIF4, enhancing its activity and thermomorphogenesis.
Together, this study unravels ELF3-BBX24/BBX25-PIF4 as a key regulatory module
that controls growth and development under varying temperature cues.

Keywords: Thermomorphogenesis, BBX24, BBX25, PIF4, ELF3, gene regulation

68 **INTRODUCTION**

Ecological cues such as light and temperature are key in regulating plant growth, architecture, and fitness (Casal and Balasubramanian, 2019; Casal and Questa, 2018; Franklin et al., 2014; Legris et al., 2019; Lippmann et al., 2019; Quint et al., 2023). Plants have evolved sophisticated and multifaceted mechanisms to perceive, integrate, and process these cues (Krahmer and Fankhauser, 2023; Legris *et al.*, 2019). Because plants experience these conflicting signals, they must constantly sense and integrate them to optimise growth and acclimate to local environments. Warm ambient temperatures strongly influence plant growth by accelerating vegetative and reproductive transitions, resulting in shortened life-cycle duration (Casal and Balasubramanian, 2019; Crawford et al., 2012; Koini et al., 2009; Krahmer and Fankhauser, 2023; Kumar et al., 2012; Quint *et al.*, 2023). However, warm temperatures strongly compromise immunity and severely affect seed set and yield (Burroughs et al., 2023; Koester et al., 2014; Zhao et al., 2017). Increasing seasonal growth temperature due to global warming impairs plant growth and yield, threatening farmers' livelihood and food security (Peng et al., 2004; Zhao *et al.*, 2017). Therefore, understanding how plants sense and integrate temperature cues is critical to delineating thermosensory signalling mechanisms to breed climate-resilient crops.

The light signaling pathway consists of photoreceptors and downstream signaling molecules that have been shown to play a pivotal role in sensing and integrating temperature cues (Legris *et al.*, 2019; Li et al., 2016). Key studies revealed that red-light absorbing phytochrome B (phyB) senses temperature cues (Jung et al., 2016; Legris et al., 2016). PhyB exists in two interconvertible physiologically active (Pfr) and inactive (Pr) forms. Warm temperature favours the conversion of Pfr to Pr form, reducing the phyB activity (Legris *et al.*, 2016). The phyB targets a group of growth-promoting transcription factors, PHYTOCHROME INTERACTING FACTORS (PIFs), for phosphorylation and degradation in a red-light-dependent manner (de Lucas et al., 2008; Huq and Quail, 2002; Lorrain et al., 2008). Among PIFs, PIF4 and PIF7 have been shown to mediate thermomorphogenesis, respectively, under SD and LD photoperiods (Burko et al., 2022; Chung et al., 2020; Delker et al., 2022; Fiorucci et al., 2020; Koini *et al.*, 2009; Kumar *et al.*, 2012). PIF4 is essential for

warm temperature-induced hypocotyl and petiole growth, rosette development,
hyponasty, and flowering (Casal and Balasubramanian, 2019; Favero et al., 2020;
Koini *et al.*, 2009; Kumar *et al.*, 2012; Oh et al., 2012; Quint *et al.*, 2023; Quint et al.,
2016; Zhao and Bao, 2021). PIF4-mediated thermomorphogenesis is linked to the
activation of genes involved in auxin/brassinosteroid biosynthesis and signaling
(Franklin et al., 2011; Ibanez et al., 2018; Krahmer and Fankhauser, 2023; Sun et
al., 2012), cell elongation (Gangappa and Kumar, 2017) and flowering(Kumar *et al.*,
2012). Warm temperature-mediated inhibition of phyB activity increases PIF4
function, resulting in thermosensory growth (Legris *et al.*, 2016). Unlike phyB, the
blue light receptor Cryptochrome 1 (CRY1) inhibits thermomorphogenesis through
PIF4 sequestration(Ma et al., 2016). The central oscillator components of the
circadian clock, including the evening complex (EC), play a crucial role in controlling
thermomorphogenesis by regulating PIF4 function. The EARLY FLOWERING 3
(ELF3), a critical component of the evening complex (EC) and a thermosensor,
inhibits *PIF4* transcription and also sequesters it by forming heterodimers (Box et al.,
2015; Jung et al., 2020; Nieto et al., 2015; Nusinow et al., 2011). Moreover, upstream
regulatory components such as CONSTITUTIVE PHOTOMORPHOGENIC 1
(COP1), DE-ETIOLATED 1 (DET1) and SUPPRESSOR OF PHYTOCHROME A-
105 1 (SPA1) proteins promote PIF4-mediated thermomorphogenic growth by
stabilizing PIF4 protein (Gangappa and Kumar, 2017; Lee et al., 2020; Park et al.,
2017). Recent studies also reveal that the HEMERA/REGULATOR OF the
CHLOROPLAST BIOGENESIS (HMR/RCB) module works cooperatively to stabilize
PIF4 during the day and promote thermomorphogenesis (Qiu et al., 2019; Qiu et al.,
2021). The Heat shock transcription factor (HSF) family members, HSFA1a/b/d/e,
interact with PIF4 and stabilize it during light by interfering with PIF4-phyB
interaction, resulting in thermomorphogenesis during the day (Tan et al., 2023a).

The B-BOX containing zinc-finger (BBX) proteins are highly conserved transcription
factors across eukaryotes that regulate gene expression by directly binding to DNA
or as transcriptional co-activators or co-repressors through protein-protein
interaction. BBX proteins have emerged as crucial integrators of various
environmental cues (Ding et al., 2018; Gangappa and Botto, 2014; Khanna et al.,

2009; Song et al., 2020; Vaishak et al., 2019; Yuan et al., 2021). The BBX24 and BBX25 are close homologs that function as transcriptional co-repressors of HY5 to inhibit photomorphogenesis (Gangappa et al., 2013; Job et al., 2018). Additionally, BBX24 functions in various signaling pathways, including UV-B signaling (Huang et al., 2022; Jiang et al., 2012), shade-avoidance response (Crocco et al., 2015; Saura-Sanchez et al., 2023) and flowering (Li et al., 2014). However, the role of BBX24 and BBX25 in temperature-mediated regulation of growth and development remains unknown. Here, we report that BBX24/BBX25 as key regulators of PIF4-mediated thermosensory growth. We found that *BBX24*/*BBX25* gene expression and protein levels are enhanced at warm temperatures, further enhancing PIF4 protein accumulation and thermomorphogenesis. Our study demonstrates that ELF3 functions upstream of BBX24/BBX25, inhibiting their gene expression during the evening hours under low temperatures. However, warm temperature-mediated inhibition of ELF3 activity results in increased BBX24/BBX25 function and, in turn, enhances PIF4-mediated thermomorphogenic growth. Additionally, BBX24/BBX25 inhibit ELF3 protein accumulation resulting in the release of the negative inhibitory effect of ELF3 on PIF4 protein and transcript accumulation. Together, this study uncovers a novel regulatory mechanism by which BBX24/BBX25 promote PIF4-mediated thermosensory growth via antagonizing ELF3 function.

RESULTS

BBX24 and BBX25 are essential for warm temperature-mediated growth

To unravel the possible role of BBX24 and BBX25 in temperature-mediated regulation of hypocotyl growth, we analyzed the hypocotyl growth response of wild-type (Col-0) and *bbx24* and *bbx25* mutants grown under 22°C and 27°C. Our data revealed that *bbx24* and *bbx25* had moderate but significantly shorter hypocotyls at 22°C and 27°C than Col-0 both under short-day (SD) and long-day (LD) photoperiods (Figure 1A-1C). Compared to Col-0, *bbx24* showed stronger inhibition in hypocotyl growth than the *bbx25* mutant at 27°C (Figure 1A-1C). Interestingly, the *bbx24bbx25* double mutant had a much shorter hypocotyl length at 27°C than the

single mutants (Figure 1A-1C), suggesting their additive function in promoting warm
temperature-mediated hypocotyl growth. Contrary to mutants, the *BBX24*
overexpression (hereafter referred to as *BBX24-OE*) transgenic line showed
enhanced thermosensory hypocotyl growth as they had significantly longer
hypocotyls than Col-0 at 22°C and 27°C both under SD and LD (Figure 1D-1F). To
further understand whether overexpression of *BBX25* has a similar effect to *BBX24-*
OE in promoting hypocotyl growth, we generated multiple transgenic lines
overexpressing *BBX25* (Supplemental Figure S1A and 1B). Like *BBX24-OE* lines,
transgenic lines overexpressing *BBX25* showed exaggerated hypocotyl growth
compared to Col-0 at 22°C and 27°C, under SD and LD (Figure 1D-1F;
Supplemental Figure S1C-S1E). Warm temperature-mediated hypocotyl elongation
is attributed to increased cell elongation, which is due to increased growth-promoting
hormones such as auxin and brassinosteroid (Franklin *et al.*, 2011; Ibanez *et al.*,
2018; Oh *et al.*, 2012). To know if BBX24/BBX25-mediated hypocotyl growth is due
to increased cell elongation, we analyzed the cell length of hypocotyl epidermal cells
from seedlings grown at 22°C and 27°C under SD. Our data reveal that the hypocotyl
epidermal cell length was significantly longer in Col-0 at 27°C than at 22°C (Figure
1G). However, in the *bbx24bbx25* double mutant*,* the cell length was strongly
inhibited (Figure 1G). Contrarily, *BBX24* and *BBX25* overexpression lines had
significantly longer cells at both 22°C and 27°C than Col-0 (Figure 1G). Together,
these results suggest that BBX24/BBX25-mediated hypocotyl growth in response to
warm temperature is linked to an increase in cell elongation of hypocotyl epidermal
cells.

Warm temperature-mediated hypocotyl growth is linked to enhanced accumulation
of auxin and brassinosteroid (BR) growth hormones (Franklin *et al.*, 2011; Gray et
al., 1998; Ibanez *et al.*, 2018; Krahmer and Fankhauser, 2023; Oh *et al.*, 2012;
Stavang et al., 2009). As *bbx24bbx25* double mutant has extremely short hypocotyls
under warm temperatures, we wanted to see if exogenous application of auxin or BR
can rescue this phenotype. Our data suggest that in response to a synthetic auxin
picloram (PIC) or epibrassinolide (eBL) the short hypocotyl phenotype of
bbx24bbx25 was partially recovered with a higher % hypocotyl growth at 27°C

compared to Col-0 (Supplemental Figure S2A-F). In contrast, *BBX24-OE* and

BBX25-OE lines showed enhanced sensitivity to both the hormones with a reduced

% hypocotyl growth compared to Col-0 (Supplemental Figure S2A-F). Moreover, we

also found that overexpression of *BBX24* and *BBX25* enhanced the promoter activity

of *DR5*, a synthetic auxin reporter (Supplemental Figure S2G). Contrastingly, when

we treated the *BBX24-OE* and *BBX25-OE* seedlings with auxin inhibitor 2,3,5-

triiodobenzoic acid (TIBA, an auxin transport inhibitor) and a BR biosynthetic

inhibitor, Propiconazole (PPZ), they showed enhanced sensitivity to the inhibitor

similar to Col-0 (Supplemental Figure S3A-F). Thus, these results suggest that

BBX24/BBX25 indeed promote auxin/BR governed hypocotyl growth in response to

warmth.

BBX24/BBX25 function is necessary for the thermoresponsive gene

expression

Consistent with the hypocotyl data, our qRT-PCR results from SD-grown seedlings

harvested at ZT23 revealed that auxin biosynthetic gene *YUCCA8* (*YUC8*), cell-wall

elongation gene *XYLOGLUCAN ENDOTRANSGLYCOSYLASE 7* (*XTR7*), auxin

signaling gene *INDOLE ACETIC ACID19* (*IAA19*), BR signaling gene

PACLOBUTRAZOL RESISTANCE5 (*PRE5*) showed significant downregulation in

the *bbx24bbx25* mutant at 27°C, while moderately reduced at 22°C, compared to

Col-0 (Figure 1H). However, in the *BBX24* and *BBX25* overexpression lines, these

genes showed significant upregulation both at 22°C and 27°C (Figure 1H).

Additionally, our qPCR data from SD-grown seedlings harvested during daytime at

ZT3 revealed that expression of growth and hormone signaling genes such as

YUC8, *DWARF4* (*DWF4*), *Gibberellin 3-oxidases* (*GA3OX1*), *IAA19*, *PRE5*,

SMALL AUXIN *UP-REGULATED RNA 21* (*SAUR21*), and genes involved in cell

elongation, such as *EXPANSIN 8* (*EXP8*) and *XTR7* showed significant

downregulation in the *bbx24bbx25* double mutant while upregulated in the

overexpression lines than Col-0, at 22°C and 27°C (Supplemental Figure S4A and

B). Together, these data confirm that BBX24/BBX25 induce the expression of

growth-related genes in response to warm temperatures both at night and during the

day.

BBX24/BBX25 gene expression and protein levels are enhanced in response

to warm temperatures

As BBX24/BBX25 promote warm temperature-mediated hypocotyl growth, we

wanted to see if warm temperature regulates their gene expression and protein

abundance. The qRT-PCR data from six-day-old seedlings revealed that *BBX24* and

BBX25 transcript levels showed significant upregulation at 27°C compared to 22°C

both at the end of the night (ZT23) and during daytime (ZT3) (Supplemental Figure

S5A). Further, when we monitored the expression of *BBX24* and *BBX25* for an

extended period during the day into early night, we found that *BBX24* and *BBX25*

expressions peak early in the day but their expression reduced around evening and

early night, both at 22°C and 27°C (Figure 2A). Interestingly, their expression is

significantly upregulated at 27°C than at 22°C both under SD (Figure 2A) and LD

(Supplemental Figure S5B and C) conditions. Moreover, immunoblot analysis of the

35S::BBX24-GFP line revealed that BBX24 protein levels was elevated at 27°C

compared to 22°C, specifically at ZT2 (Supplemental Figure S5D). Further, when we

monitored BBX24 protein accumulation at different ZT times, we found that at ZT0,

BBX24 protein was hardly detectable, and the protein stabilized in response to light

and reached the peak at ZT2 and then reduced at ZT4 and ZT8 at 22°C (Figure 2B).

However, at 27°C, BBX24 protein abundance increased nearly two-fold over 22°C

at ZT2, and the protein level was largely comparable between 22°C and 27°C at ZT4

and ZT8 (Figure 2B). Similarly, immunoblot analysis of the *35S::Myc-BBX25*

transgenic line revealed that at ZT2, BBX25 protein levels was also elevated at 27°C

compared to 22°C under SD (Supplemental Figure S5E). Specifically, BBX25 protein

was more stable at ZT2 and ZT4 at 27°C compared to 22°C (Figure 2C). Unlike

BBX24, BBX25 was detectable at the end of the night (ZT23), but its levels was not

altered at 27°C (Figure 2C). Consistent with the increased accumulation of BBX24

protein in response to 27°C, microscopic analysis of *35S:BBX24:GFP* seedlings

revealed that BBX24-GFP was more prominent in the hypocotyl region as relative

fluorescence of GFP was significantly increased at 27°C than at 22°C (Figure 2D-
2E), while no fluorescence was detected in Col-0 (Supplemental Figure S5F).
Subsequently, we also observed an increased number of cells with BBX24-GFP
fluorescence in seedlings grown at 27°C compared to 22°C (Figure 2F). Together,
these results suggest that warm temperatures promote BBX24 and BBX25 gene
expression and protein accumulation.

BBX24/BBX25 are essential for the accumulation of PIF4 protein in response
to warm temperature

As our above results indicate that BBX24/BBX25 are required for the optimal
expression of growth and hormone-related genes in response to warm
temperatures, we wanted to know if this is due to differential PIF4 protein
abundance. Our immunoblot data from SD-grown seedlings at 22°C revealed that at
the end of the night (ZT23), PIF4 protein levels was reduced in the *bbx24bbx25*
double mutant than Col-0, while during the day time it was more or less similar to
Col-0 (Figure 3A). Interestingly, at 27°C, the PIF4 protein levels was strongly
reduced in the *bbx24bbx25* double mutant at the end of the night (ZT23) and during
the day (ZT2-ZT8) compared to Col-0 (Figure 3A). However, in Col-0, PIF4 levels
stayed very high at the end of the night (ZT23) and throughout the day (ZT2-ZT8)
(Figure 3A). Supporting this, when 22°C grown seedlings under SD were treated
with 27°C for short durations (0, 2, 4, 8h), the PIF4 protein levels in Col-0 increased
with the duration of warmth treatment (Figure 3B). However, in the *bbx24bbx25*,
PIF4 fails to accumulate in response to warm and its levels stayed very low than Col-
0 (Figure 3B). Consistently, overexpression of *BBX24* and *BBX25* also resulted in
increase in PIF4 protein levels at 22°C and 27°C compared to Col-0 (Figure 3C).
BBX24/BBX25 was shown to inhibit photomorphogenesis by interfering with the HY5
transcriptional activity (Gangappa *et al.*, 2013). Therefore, we wanted to see if
BBX24/BBX25-mediated thermosensory hypocotyl growth is due to altered HY5
protein status. Our immunoblot data suggests that HY5 protein levels were
comparable between *bbx24bbx25* double mutant and Col-0 both at 22°C and 27°C
(Figure 3D), confirming that BBX24/BBX25-mediated thermosensory hypocotyl

growth is not due to reduced HY5 levels but due to enhanced PIF4 protein accumulation. Together, these results suggest that the BBX24/BBX25 function is crucial for promoting PIF4 protein abundance in response to warmth both during night and daytime.

Next, we wanted to see if BBX24/BBX25 also affects *PIF4* at the transcriptional level; we checked its transcript levels and found that *PIF4* expression is moderately but significantly downregulated in the *bbx24bbx25* mutant while upregulated in *BBX24* and *BBX25* overexpression transgenic lines at the end of the night (ZT23), both at 22°C and 27°C under SD (Supplemental Figure S6A). However, the regulation of *PIF4* gene expression by BBX24/BBX25 is likely indirect, as our ChIP analysis followed by qPCR data revealed that BBX24/BBX25 did not bind to the *PIF4* promoter (Supplemental Figure S6B-D). Furthermore, ChIP analysis also indicates that BBX24/BBX25 do not bind to the promoters of PIF4 targeted growth-responsive genes (Supplemental Figure S6E-G).

Further, when we treated five-day-old seedlings with proteasomal inhibitor MG132, we found that protein stability increased in the *bbx24bbx25* double mutant compared to mock-treated seedlings (Figure 3E). However, it was still visibly reduced compared to MG132-treated Col-0 seedlings (Figure 3E and 3F). Consistent with this, when we treated seedlings with a translation inhibitor, cycloheximide (CHX), we found that PIF4 degradation was significantly faster in the *bbx24bbx25* mutant, specifically at 60 and 120 min of CHX treatment, compared to Col-0 (Figure 3G and 3H). Together, these data suggest that BBX24 and BBX25 play a crucial role in *PIF4* transcript accumulation and also in maintaining its protein abundance at the posttranslational level.

BBX24/BBX25-mediated hypocotyl elongation in response to warmth is dependent on PIF4

As BBX24/BBX25 mediated warm temperature response is linked to enhanced PIF4 protein accumulation, we wanted to check the genetic interaction of BBX24/BBX25 with PIF4. We introduced *pif4* mutation into *BBX24-OE* and *BBX25-OE* lines through genetic crosses and generated homozygous lines. Analysis of hypocotyl length

revealed that the exaggerated hypocotyl growth of the *BBX24-OE* line was strongly

suppressed in the *pif4* mutant background as the hypocotyl length of *BBX24-OE pif4*

was close to the *pif4* mutant (Supplemental Figure S7A and B). Similarly, the *BBX25-*

OE pif4 hypocotyl growth was markedly suppressed compared to the *BBX25-OE* line

(Supplemental Figure S7A and B). These results confirm that BBX24/BBX25-

mediated thermosensory hypocotyl growth strongly depends on the PIF4, implying

that BBX24/BBX25 probably acts upstream of PIF4.

BBX24/BBX25 function is essential for PIF4-mediated thermosensory growth

Consistent with the above results that BBX24/BBX25 are critical for maintaining PIF4

protein abundance, our genetic analysis also revealed that BBX24/BBX25 are

necessary for PIF4-mediated thermosensory growth (Figure 4). We have introduced

bbx24, *bbx25* and *bbx24bbx25* mutations in the *pPIF4:PIF4-FLAG* transgenic line,

which overexpresses PIF4 (Gangappa et al., 2017) (hereafter referred to as *PIF4-*

OE) (Supplemental Figure S8A) and analyzed their hypocotyl phenotypes. The

bbx24 or *bbx25* single mutants showed moderate suppression, while the

bbx24bbx25 strongly suppressed the exaggerated hypocotyl and cotyledon

phenotype of the *PIF4-OE* line at 22°C and 27°C under SD (Figure 4A-4D;

Supplemental Figure S8B-E). Similar to SD, under LD 22°C, *bbx24* and *bbx25*

mutations showed additive effects in suppressing the cotyledon phenotype of the

PIF4-OE line (Supplemental Figure S8F and G). Consistent with the growth

suppression, *bbx24bbx25* strongly suppressed the expression of growth-promoting

genes such as *YUC8*, *IAA19*, *XTR7*, *SAUR22* and *PRE5* in the *PIF4-OE* line

(Supplemental Figure S9A).

Notably, in line with the reduced growth and gene expression phenotype, the PIF4

protein levels was moderately reduced in the single mutants but almost undetectable

in the double mutant background both at the end of the night (Figure 4E) and during

the day (Supplemental Figure S9B). Similarly, when 22°C SD grown seedlings were

exposed to short-term warmth (27°C) for 24h, the PIF4 protein levels was markedly

reduced in both *bbx24 PIF4-OE* and *bbx25 PIF4-OE* compared to *PIF4-OE* line

(Figure 4F). Interestingly, the PIF4 protein was almost undetectable in the

bbx24bbx25 PIF4-OE mutant (Figure 4F). Consistent with the unstable nature of the

PIF4 protein in the *bbx24bbx25* double mutant, we found that the *PIF4* transcript

accumulation was drastically reduced in the *bbx24bbx25* double mutant than Col-0,

both at 22°C and 27°C (Supplemental Figure S9C). Also, in line with the reduced

PIF4 mRNA accumulation in the *bbx24bbx25* double mutant observed above,

treating *bbx24bbx25* double mutants with MG132 only partially stabilized the PIF4

protein (Figure 4G) further confirms that BBX24/BBX25-mediated PIF4 protein

accumulation is manifested both at the transcription and posttranslational level.

In line with the role of BBX24/BBX25 in facilitating PIF4 protein accumulation,

overexpression of *BBX24* or *BBX25* in the *PIF4-OE* background resulted in further

enhancement in the hypocotyl length while they further suppressed the small

cotyledon phenotype of *PIF4-OE* in SD (Supplemental Figure S9D-G) and LD

(Supplemental Figure S10A-D) conditions. The *PIF4-OE BBX24-OE* and *PIF4-OE*

BBX25-OE double overexpression transgenic lines showed etiolated growth with

very long hypocotyl and half-opened or closed cotyledons, especially at 27°C, which

is a characteristic phenotype of genotypes with very high PIF4 activity (Supplemental

Figure S9D and E). In line with the exaggerated growth phenotypes of the double

overexpression transgenic lines, immunoblotting data revealed that when 22°C LD

grown seedlings were exposed to 27°C for 4h, PIF4 protein levels was elevated in

the *PIF4-OE BBX24-OE* and *PIF4-OE BBX25-OE* transgenic seedlings compared

to *PIF4-OE* (Supplemental Figure S10E). The *PIF4* transcript levels in these *BBX24-*

OE PIF4-OE and *BBX25-OE PIF4-OE* transgenic seedlings was significantly more

than the Col-0 but lower than the *PIF4-OE* line, both at 22°C and 27°C under SD

(Supplemental Figure S10F), which is likely due to feedback regulation to regulate

PIF4 threshold levels.

BBX24/BBX25 physically interact with PIF4

Next, to know if BBX24/BBX25-mediated PIF4 protein stabilization at the

posttranslational level requires direct physical interactions, we carried out a Co-

immunoprecipitation (Co-IP) assay. When BBX24-GFP protein was pulled down

using an anti-GFP antibody, we could detect PIF4 protein in the immunoprecipitated

complex using a native PIF4 antibody (Figure 4H), while in Col-0 (used as a negative control), we didn't detect PIF4 protein, suggesting that BBX24 physically interacts with PIF4 in vivo (Figure 4H). Similarly, when we pulled down Myc-BBX25 using an anti-cMyc antibody, we could detect the PIF4 protein in the immunoprecipitated complex (Figure 4I) but not in Col-0, implying that BBX25 physically interacts with PIF4 in vivo (Figure 4I). Consistent with Co-IP data, Bimolecular Fluorescent Complementation (BiFC) assays from onion epidermal cells also revealed BBX24 and BBX25 physically interacts with PIF4 in planta (Figure 4J). To further confirm these interactions, we also performed yeast two-hybrid (Y2H) assays by cloning BBX24 and BBX25 with GAL4 DNA-binding domain (DBD) and PIF4 with Activation domain (AD). Our Y2H assay revealed that BBX24/PIF4 and BBX25/PIF4 co-transformed colonies were able to grow on the minimal media lacking Histidine and Adenine and were also able to activate the production of α-galactosidase, which converts 5-Bromo-4-chloro-3-indolyl α-D-galactopyranoside (X-α-Gal) substrate into and blue colour (Figure 4K). Together, these shreds of evidence confirm that BBX24 and BBX25 physically interact with PIF4 in vivo.

ELF3 inhibits *BBX24/BBX25* gene expression during dawn in a temperature-dependent manner

ELF3, ELF4 and LUX ARRHYTHMO (LUX) form the Evening Complex (EC). ELF3 is the key component of EC as it bridges ELF4 and LUX (Nusinow *et al.*, 2011) and is a crucial negative regulator of thermosensory signaling (Box *et al.*, 2015; Jung *et al.*, 2020; Nieto *et al.*, 2015). ELF3 strongly inhibits hypocotyl growth under optimal temperatures by directly inhibiting PIF4 transcription and its transcriptional activity through direct physical association and sequestration (Nieto *et al.*, 2015; Nusinow *et al.,* 2011). Our above results show that *BBX24/BBX25* mRNA expression peaks at the beginning of the day and is drastically reduced during the end of the day through to early night. However, their expression was significantly elevated under warm temperatures, even during the evening, as shown above (Figure 2A). To know if ELF3 inhibits *BBX24/BBX25* gene expression in the evening, we analyzed gene expression through qPCR from Ws (Wild-type) and *elf3* mutant seedlings harvested

at ZT6 and ZT10 grown in SD. The expression of *BBX24* and *BBX25* was

comparable between Ws and *elf3* mutant at ZT6 (Figure 5A and 5B). Interestingly,

their expression is significantly increased in *elf3* mutant at ZT10 compared to Ws,

both at 22°C and 27°C (Figure 5A and 5B), suggesting the plausible role of ELF3 in

inhibiting *BBX24/BBX25* gene expression during the evening when the EC activity

peaks.

LUX is a DNA-binding component of EC and has been shown to bind to conserved

cis-acting variable motifs such as *AGATTCGA*, *AGATACGC*, *ATATTCGA*,

AAGATCTT and *GGATCCGA* (Chow et al., 2012; Helfer et al., 2011; Nusinow *et al.*,

2011; Silva et al., 2020). We looked for potential LUX binding sites to further

understand if ELF3 (a component of EC) can directly bind to *BBX24* and *BBX25*

promoters. Interestingly, the *BBX24* and *BBX25* promoters harbour potential LUX

binding motifs, *GATACG* and *AAGATCG*, respectively (Figure 5C and D), which are

similar to the previously reported motifs bound by LUX (Silva *et al.*, 2020). We carried

out a Chromatin Immunoprecipitation (ChIP) assay using a *35S::ELF3-YFP*

transgenic line expressing stable protein at ZT10 under 22°C and 27°C under SD

(Supplemental Figure 11A). Our ChIP qPCR results revealed a significant

enrichment of ELF3 onto the *BBX24* and *BBX25* promoters at the region containing

the LUX-Binding site in the IP samples compared to mock (preimmune serum)

(Figure 5E and 5F). However, the binding of ELF3 to *BBX24* and *BBX25* promoters

was strongly inhibited at 27°C grown seedlings (Figure 5E and 5F). This is also

consistent with the earlier reports that EC binding to the target DNA was

compromised at warm temperatures (27°C) (Silva *et al.*, 2020). In summary, our

study demonstrates *BBX24/BBX25* as novel targets of EC for transcriptional

repression at lower temperatures to inhibit growth, while warm temperature-

mediated inhibition of ELF3 activity relieves this repression.

BBX24/BBX25 genetically act downstream to ELF3

Our above results suggest that BBX24/BBX25 have an impact on PIF4 at both

protein and transcript levels. ELF3, a potent negative regulator of PIF4, inhibits *PIF4*

transcription and also forms heterodimers and sequesters it (Box *et al.*, 2015; Nieto

et al., 2015; Nusinow *et al.*, 2011). As ELF3 binds to the *BBX24*/*BBX25* promoter to

suppress their gene expression, we hypothesize that ELF3 could be the possible

target through which BBX24/BBX25 regulates PIF4 functions. To address this, we

performed epistatic interactions by generating *elf3bbx24*, *elf3bbx25* and

elf3bbx24bbx25 mutants. Measurement of hypocotyl length of double, triple and

parental genotypes along with WT (segregated from Col-0xWs) suggests that while

bbx24 moderately suppressed, *bbx25* showed weaker suppression of long hypocotyl

phenotype (Supplemental Figure S11B and C) and smaller cotyledon area of *elf3*

mutant (Supplemental Figure S11D and E). However, *bbx24bbx25* could strongly

suppress the long hypocotyl phenotype of *elf3* mutant, as *bbx24bbx25elf3* hypocotyl

length was significantly shorter (Figure 6A and 6B). In contrast, the cotyledon area

of *bbx24bbx25elf3* was significantly bigger than *elf3* mutant (Supplemental Figure

S11F and G), suggesting that *elf3* growth phenotypes are strongly dependent on

BBX24/BBX25.

As our genetic and molecular data reveal that BBX24/BBX25 probably function

downstream to ELF3 but upstream to PIF4, we wanted to see if the suppression of

elf3 growth phenotype by *bbx24bbx25* is due to reduced PIF4 protein levels.

Interestingly, our immunoblot data from six-days-old 22°C SD-grown seedlings or

22°C SD-grown seedlings exposed to 27°C for 24h indeed showed that PIF4 protein

levels was markedly reduced in the *bbx24bbx25elf3* triple mutant compared to *elf3*

mutant both at 22°C and 27°C (Figure 6C). Moreover, increased PIF4 accumulation

in *elf3* mutant at ZT23 and ZT8 in 22°C and 27°C grown seedlings was found to be

dependent on BBX24/BBX25, as PIF4 protein levels was strikingly reduced in the

bbx24bbx25elf3 triple mutant (Supplemental Figure 12A). The increased PIF4

protein levels in the *elf3* mutant is likely due to increased accumulation of *PIF4*

transcript. In line with this, we found that the *PIF4* transcript levels was significantly

reduced in the *bbx24bbx25elf3* triple mutant compared to *elf3* (Supplemental Figure

S12B). Consistent with the suppression of *elf3* phenotype by *bbx24bbx25*, the

expression of PIF4-regulated growth genes such as *XTR7*, *IAA19*, *PRE5,* and *IAA29*

are significantly downregulated in the *bbx24bbx25elf3* mutant background than *elf3*

(Supplemental Figure S12C). Together, these results confirm that BBX24/BBX25-

mediated inhibition of ELF3 function is necessary for optimal PIF4 gene expression
and protein accumulation.

Consistent with the role of BBX24/BBX25 in thermosensory hypocotyl and cotyledon
growth, we also found that they play an important role in later developmental stages,
such as petiole elongation and flowering. We found that long petiole and early
flowering phenotypes of *PIF4-OE* and *elf3* mutants were strongly suppressed by
bbx24bbx25 mutations both under SD (Supplemental Figure S13) and LD
(Supplemental Figure S14) conditions.

BBX24/BBX25 antagonizes ELF3 to facilitate PIF4 function

As our above data suggest that BBX24/BBX25 functions antagonistic to ELF3, we
hypothesized that BBX24/BBX25 might inhibit ELF3 activity to promote PIF4
functions. To further test this, we introduced *BBX24-OE* and *BBX25-OE* transgenes
into the *ELF3-OE* background, generated *BBX24-OE ELF3-OE* and *BBX25-OE*
ELF3-OE double homozygous transgenic lines and analyzed their hypocotyl
phenotype. Measurement of hypocotyl length from SD-grown seedlings at 22°C and
27°C revealed that overexpression of *BBX24* or *BBX25* significantly suppressed the
short hypocotyl phenotype of the *ELF3-OE* transgenic line (Supplemental Figure
S15A and B). Consistent with this, our immunoblot data using ELF3 specific antibody
(Supplemental Figure S15D) revealed that ELF3 protein level is elevated in the
bbx24bbx25 double mutant but markedly reduced in the *BBX24* and *BBX25*
overexpression lines compared to Col-0 (Figure 6D). In contrast to the effect of
bbx24bbx25 mutations in increasing ELF3 protein levels, *ELF3* gene expression was
not altered in the *bbx24bbx25* double mutant compared to Col-0 (Supplemental
Figure 15C). Further, consistent with the suppression of short hypocotyl length of
ELF3-OE line by *BBX24-OE* and *BBX25-OE*, we found that, overexpression of
BBX24 and *BBX25* strongly recovered the reduced PIF4 protein levels of *ELF3-OE*
line (Figure 6E) indicating negative inhibitory effects of BBX24/BBX25 on ELF3 to
facilitate PIF4 protein accumulation.

Next, to know if BBX24/BBX25-mediated inhibition of ELF3 function requires direct
physical interactions, we carried out a Co-IP assay. We used *35S:BBX24-GFP*

transgenic lines and pulled down BBX24-GFP from the total protein extract using an

anti-GFP antibody. In the BBX24-GFP immunoprecipitated complex, we could detect

ELF3 as revealed by immunoblotting using a native anti-ELF3 antibody, but not in

the Col-0 (Figure 6F). Similarly, when we pulled down from the total protein extract

of *35S:Myc-BBX25* transgenic line, we could detect ELF3 in the immunoprecipitated

complex but not in Col-0 (Figure 6G), indicating that BBX24 and BBX25 physically

interact with ELF3 in vivo. To support these physical interactions, we performed a

BiFC assay and found that BBX24 and BBX25 physically interact with ELF3 in planta

in onion epidermal cells (Figure 6H, Supplemental Figure S15E). When we co-

transformed BBX24-nYFP/ELF3-cYFP and BBX25-nYFP/ELF3-cYFP constructs,

we observed reconstitution of uniform YFP fluorescence in the nucleus (Figure 6H).

Interestingly, we also observed two or more discrete condensates in the nucleus

(Figure 6H; Supplemental Figure 15F). Furthermore, Y2H assays, using ELF3 as

bait and BBX24 and BBX25 as prey, found that BBX24 and BBX25 physically

interact with ELF3 (Figure 6I). The BD-BBX24/AD-ELF3 and BD-BBX25/AD-ELF3

combinations of constructs, when transformed onto Y2H-Gold reporter strain, they

grew on the synthetic minimal media lacking Histidine and Adenine and also, these

colonies developed blue colour in the presence of X-α-Gal substrate (Figure 6I).

Together, our genetic and biochemical data confirm that BBX24/BBX25 inhibit ELF3

function to promote PIF4-mediated thermomorphogenic growth.

DISCUSSION

Rising global temperatures due to climate change profoundly affect plant health and

pose a significant threat to crop yields (Burroughs *et al.*, 2023; Koester *et al.*, 2014;

Peng *et al.*, 2004; Zhao *et al.*, 2017). While plants have evolved intricate signaling

mechanisms to sense, integrate, and adapt to temperature fluctuations, the

underlying signaling mechanisms are poorly understood. As PIF4 is the central

component of thermomorphogenesis, maintaining its protein abundance is crucial in

achieving optimal thermomorphogenic growth both during the night and daytime.

BBX proteins have emerged as crucial signaling nodes for orchestrating adaptive

responses to diverse environmental signals such as light, shade, and abiotic

stresses (Ding *et al.*, 2018; Gangappa and Botto, 2014; Podolec et al., 2022; Song

et al., 2020; Tripathi et al., 2017; Vaishak *et al.*, 2019; Wang et al., 2014; Zhao et al.,

2020). This study discovers that BBX24/BBX25 are novel and essential components

of the thermosensory signaling pathway and act to promote PIF4 function by

inhibiting its critical repressor, ELF3 (Figure 7).

Our study demonstrates that BBX24/BBX25 play a positive regulatory role in

temperature-responsive hypocotyl growth (Figure 1). BBX24/BBX25 functions

exhibit significant overlap, as evidenced by a more pronounced reduction in cell

length and hypocotyl growth in the *bbx24bbx25* double mutant than in the *bbx24* and

bbx25 single mutants at warm temperature, as opposed to *BBX24-OE* and *BBX25-*

OE overexpression lines that promoted cell elongation and hypocotyl growth (Figure

1). Consistent with the growth phenotypes, *bbx24bbx25* double mutants had

reduced expression of PIF4 target genes such as *XTR7*, *YUC8*, *EXP8*, *IAA19*, etc.,

involved in hormone biosynthesis and signaling both at night and during the day

(Figure 1; Supplemental Figure S4), which is in line with the reduced PIF4 protein

accumulation in response to warm temperatures (Figure 3). Contrary to *bbx24bbx25*

mutants, the enhanced thermomorphogenic response and gene expression of the

BBX24-OE and *BBX25-OE* lines, was consistent with the increased PIF4 protein

levels (Figure 3C). Our genetic analysis demonstrates that BBX24/BBX25-mediated

thermomorphogenic response is directed through PIF4 (Supplemental Figure S7).

In agreement with the BBX24/BBX25 acting through PIF4, the genetic analysis also

revealed that the absence of *BBX24* and *BBX25* strongly suppress the robust growth

and gene expression phenotypes of the transgenic line overexpressing *PIF4-FLAG*,

which is due to the reduced PIF4-FLAG protein accumulation in the single mutants,

while complete absence of PIF4-FLAG protein in the *bbx24bbx25* background

(Figure 4). The lack of PIF4-FLAG protein was somewhat surprising compared to

native PIF4 protein. The possible reason could be the enhanced degradation of

PIF4-FLAG protein over native PIF4 protein.

Several thermosensory signaling components such as HMR (HEMERA), RCB

(REGULATOR OF CHLOROPLAST BIOGENESIS), and HSFA1s (HEAT SHOCK

TRANSCRIPTION FACTOR 1) have been shown to stabilize PIF4 and promote

daytime mediated thermomorphogenic growth (Qiu *et al.*, 2021; Tan et al., 2023b).

Meanwhile, COP1/DET1/SPA, upstream regulators of PIF4, promote PIF4 protein

accumulation during the night (Gangappa and Kumar, 2017). Our study reveals that

BBX24/BBX25 serve as novel upstream regulators, essential for maintaining optimal

PIF4 levels at night and also during the day to promote thermomorphogenesis

(Figure 1 and 3). As BBX24/BBX25 and COP1/DET1 promote PIF4 accumulation,

their possible functional connection in maintaining optimal PIF4 protein levels in the

dark period needs further investigation. While BBX24/BBX25 largely contribute to

PIF4 protein accumulation by enhancing *PIF4* gene expression, likely, they are also

protecting PIF4 protein from its negative regulators at the post posttranslational level,

as revealed from our MG132 and CHX treatment experiments (Figure 3E-3H). The

physical interactions between BBX24/BBX25 and PIF4 (Figure 4) also provide

evidence that these proteins likely play a crucial role in safeguarding PIF4 from its

repressors.

Previous studies have shown that ELF3 acts upstream of PIF4 and inhibits its

function both at the transcription and at the protein level (Box *et al.*, 2015; Nieto *et*

al., 2015; Nusinow *et al.*, 2011). Interestingly, our data demonstrate that the *elf3*

mutant growth phenotypes, gene expression, and PIF4 protein accumulation

strongly depend on BBX24/BBX25, suggesting that BBX24/BBX25 likely act

downstream to ELF3 but upstream to PIF4 (Figure 6). Our RT-qPCR and ChIP data

further support that ELF3 inhibits *BBX24* and *BBX25* gene expression during the

evening at 22°C, which is nullified at 27°C (Figure 5), due to warm temperature-

mediated inactivation of ELF3 activity (Jung *et al.*, 2020). Nevertheless, our

immunoblot data revealed that BBX24/BBX25 inhibit ELF3 protein accumulation at

warm temperatures (Figure 6D). Therefore, in addition to warm temperature-

mediated inhibition of ELF3 activity through liquid-liquid phase separation (LLPS),

BBX24/BBX25 inhibit ELF3 function by reducing ELF3 accumulation. Moreover, the

physical association of BBX24/BBX25 could also sequester with ELF3, resulting in

enhanced PIF4 function (Figure 6). This is also supported by genetic data, wherein

BBX24-OE and *BBX25-OE* significantly suppressed the short hypocotyl phenotype

and reduced PIF4 accumulation in the *ELF3-OE* transgenic line (Figure 6E;

Supplemental Figure S15A and B). Therefore, we propose that constitutive

thermomorphogenic response and high PIF4 activity in the *elf3* mutant is not solely

attributed to the alleviation of transcriptional repression on PIF4 but also due to the

concurrent increase in BBX24/BBX25 activity, and these dual mechanisms together

potentiate thermomorphogenic response (Figure 6; Supplemental Figure S11 and

S12).

Consistent with the role of BBX24/BBX25 in promoting PIF4 function, the PIF4-

mediated thermosensory petiole growth and flowering are also dependent on PIF4

both under SD and LD photoperiods, highlighting their role beyond seedling

development (Supplemental Figure S13 and S14). These pieces of evidence reveal

that BBX24/BBX25 proteins act as key regulators of growth and development in

response to varying light and temperature cues by coordinating clock and external

cues. In summary, we propose the ELF3-BBX24/BBX25-PIF4 as a novel regulatory

module (Figure 7), critical for integrating temperature and light cues to regulate

thermosensory growth and development in Arabidopsis.

METHODS

Plant Material and Growth Conditions

All the lines used here are in Columbia-0 (Col-0) background except *elf3*-4, which is

in Wassilewskija (WS) background. The genotypes *bbx24-1* (Gangappa *et al.*, 2013),

bbx25-2 (Gangappa *et al.*, 2013) *35S::BBX24-GFP*(Job *et al.*, 2018), *pPIF4*::*PIF4-*

FLAG (Gangappa *et al.*, 2017), *pif4-101* (Lorrain *et al.*, 2008), and elf3-4 {Dixon,

2011 #2021), have been described previously. For hypocotyl experiments, seeds

were surface sterilized with 70% ethanol containing 0.5% Triton X-100 for 10 minutes

and plated on half-strength Murashige and Skoog (MS) medium (Sigma-Aldrich)

supplemented with 1% sucrose, 0.5 mM MES (pH-5.7) and 0.8% agar following

stratification at 4°C for four-days. For the hormone experiments, seeds were plated

on MS media containing different concentrations of hormones or their inhibitors.

Upon seed germination at 22°C long day (16h Light/8h Dark) growth chamber

(Percival Scientific) seedlings were transferred to respective 22°C and 27°C short

day (8h Light/16h Dark) conditions or as stated in the experimental conditions with 70% humidity and light intensity of 100 µmol m^{-2} sec^{-1}. Five days after germination (six-day-old), seedlings were taken for the hypocotyl length and cotyledon area measurements using ImageJ software. At least 25 seedlings for each genotype were taken for the measurements.

Generation of *BBX25* overexpression transgenic lines

For the generation of *35S::Myc-BBX25* overexpressing transgenic lines, the full-length BBX25 coding sequence was amplified from 22°C LD grown Col-0 cDNA and cloned into the pENTR vector through pENTR™ Directional TOPO® Cloning Kits (Invitrogen), which is then subcloned into the destination vector pGWB618 (Nakamura et al., 2010) using LR cloning kit (Invitrogen). The binary expression cassette is then transformed into *Agrobacterium tumefaciens* GV3101 strain. Agrobacterium harbouring the transgene were transfected into Col-0 through the floral dip method. Positive transformants were selected on ½ MS containing 7.5 ug/ml glufosinate-ammonium (BASTA). Four independent lines showing 3:1 segregation on BASTA were chosen for further analysis. In the T$_3$ generation, these lines were selected based on transgene and protein expression analysis for further analysis.

Generation of double and triple mutants

For the generation of double mutants *bbx24-1bbx25-2*, *bbx24-1 PIF4-OE*, *bbx25-2 PIF4-OE*, *BBX24OE pif4-101*, *BBX25-OE pif4-101*, *bbx24-1 elf3-4*, *bbx25-2 elf3-4*, *BBX24-OE PIF4-OE*, *BBX25-OE PIF4-OE*, *BBX24-OE ELF3-OE* and *BBX25-OE ELF3-OE* we crossed *bbx24-1* and *bbx25-2* with each other and *bbx24-1*, *bbx25-2*, *BBX24-OE* and *BBX25-OE* with the respective genotypes. For the generation of triple mutants *bbx24-1 bbx25-2 PIF4-OE* and *bbx24-1 bbx25-2 elf3-4* we crossed homozygous double mutant of *bbx24-1 PIF4-FLAG* with *bbx25-2 PIF4-FLAG* and *bbx24-1elf3-4* with *bbx25-2elf3-4*, respectively. After crossing, we got F$_1$ seeds that grew for one more generation to get F$_2$ seeds. Approximately 20-30 F$_2$ seedlings growing on antibiotic selection were picked up where homozygosity was confirmed through PCR-based genotyping method using gene-specific and T-

DNA specific primers. T-DNA-specific primers used were SALK LB1.3 for *bbx24-1*,
SAIL LB3 for *pif4-101*, and *spm1* for *bbx25-2*. For *elf3-4* genotyping, we amplified
PCR fragments using dCAPS primers followed by AluI restriction digestion. To
confirm the homozygosity of *BBX24-OE*, *BBX25-OE*, *PIF4-FLAG* and *ELF3-OE*, we
first confirmed the presence of CDS using PCR, followed by the survival of seedlings
on an antibiotic selection plate in the next generation. F_3 generation seedlings
showing 100% survival on double antibiotic selection were used for further
experiments. Primers used for genotyping are listed in Table S1.

Gene expression analysis

Approximately 100 mg seedlings from the indicated genotypes and growth
conditions were harvested, frozen in liquid nitrogen, and stored at -80°C until further
processing. For RNA extraction, seedlings were ground to a fine powder in liquid
nitrogen, and total RNA was isolated using an RNeasy plant mini kit (Qiagen)
following the kit protocol with on-column DNase I digestion (Qiagen) for 30 min. at
RT. 1.5 µg of total RNA was converted into cDNA using SuperScript® III First-Strand
Synthesis kit (Invitrogen) by taking Oligo(dT)$_{20}$ primers following manufacturer's
instructions. For RT-qPCR, nuclease-free water-diluted cDNA was mixed with
PowerUp™ SYBR™ Green Master Mix (ThermoFisher Scientific) and gene-specific
primers. Each gene RT-qPCR reaction was set in triplicate with a Biorad CFX96
Touch Real-Time PCR Detection System or an Applied Biosystems^{TM} 7500 Fast
Real-Time PCR System. The transcript level of each gene was calculated relative to
that of EF1α. RT-qPCR results were analyzed in fold change using the $2^{-\Delta\Delta Ct}$ method.

Protein extraction and Immunoblotting

For western blotting, approximately 100-150 mg tissue of six-day-old *Arabidopsis*
seedlings grown in different experimental conditions were harvested in a
microcentrifuge tube, snap-frozen and ground to fine powder in liquid nitrogen. Total
protein was extracted in 250 µl of protein extraction buffer containing 50 mM Tris-
HCl pH 8.0, 150 mM NaCl, 10% glycerol, 5mM DTT, 1% NP-40, 0.5mM PMSF, and
1X EDTA free Protease Inhibitor Cocktail (Roche).
After resuspending in the extraction buffer, crude protein samples were cleared by

centrifuging at 15,000 rpm for 10 min at 4°C. Protein concentration was estimated

using Bradford reagent (Biorad). Approximately 30 µg of total protein was boiled at

95°C for 5 minutes in 5x SDS gel loading dye (100 mM Tris-HCL pH 7.5, 12%

glycerol, 4% SDS, 200 mM DTT, 40 mM β-ME and 0.001% bromophenol blue),

separated via SDS-PAGE and transferred to the PVDF membrane (Millipore, Sigma)

by wet transfer method.

For protein detection, blots were blocked with 5% skim milk in TBST for 2h at RT,

incubated overnight at 4°C with respective primary antibodies, followed by

incubation in 1:10,000 dilutions of horseradish peroxidase-conjugated rabbit anti-

goat (Thermo Fisher Scientific, 811620) or goat anti-rabbit (Abcam, ab205718)

secondary antibodies at room temperature for 1 h. Primary antibodies used were

polyclonal goat anti-PIF4 antibodies (Agrisera, AS16 3955, 1:2500), polyclonal rabbit

anti-GFP antibodies 1:7500 (Abcam, ab290), monoclonal rabbit anti-GFP (Invitrogen

G10362, 1:500) monoclonal HRP conjugated mouse anti-myc (HRP) antibodies

(Abcam, ab62928, 1:5000), monoclonal HRP conjugated mouse anti-FLAG

antibodies (Abcam, ab49763, 1:5000), polyclonal rabbit anti-HY5 antibodies

(Agrisera, AS12 1867, 1:2500), polyclonal rabbit anti-ELF3 antibodies (Agrisera,

AS18 4168, 1:2000) polyclonal rabbit anti-actin antibodies (Abcam, ab197345,

1:10,000), and Rabbit polyclonal anti-Histone H3 antibodies, 1:10,000 (abcam,

ab1791). The chemiluminescence signal was detected using the SuperSignal West

Femto kit (Thermo Fisher Scientific) and Syngene gel-doc system. Protein band

intensities were measured using the ImageJ software (https://imagej.nih.gov/).

MG132 and CHX treatment

Mg132 and CHX treatments were carried out as reported (Zhou et al., 2021) with

minor modifications. For MG132 treatment, five-day-old *Arabidopsis* seedlings were

transferred to liquid MS medium containing 50µM MG132 or DMSO and then

incubated for 8 h or 16 h before tissues were harvested for immunoblot. To

determine the degradation rate of PIF4 protein by blocking de novo protein

synthesis, five-day-old 22°C SD grown seedlings treated with MG132 as above

before shifting to 27°C for 8hrs were washed five times and then transferred to liquid

MS medium supplemented with 200µM CHX followed by tissue harvest at different

time points for western blot analysis.

707   Chromatin Immunoprecipitation (ChIP) assay

ChIP assays were performed as previously described (Yamaguchi et al., 2014) with

modifications. Approximately 1.5 g of *35S::ELF3-YFP*(*elf3-4*) seedlings grown for

five days after germination at 22°C SD and 27°C SD were harvested and cross-

linked with 30 ml of 1% formaldehyde in 1X PBS for 2 x 10 mins by vacuum

infiltration. Crosslinking was quenched by adding 125 mM glycine under vacuum for

another 5 min. Fixed seedlings were rinsed with 30 ml 1X PBS twice, dried on stacks

of filter paper, frozen in liquid nitrogen and stored at -80°C. Frozen samples were

crushed in liquid N_2 until they became light green and homogenized in 30 ml of

extraction buffer I (10 Tris-HCL pH 8, 0.4M sucrose, 10 mM MgCl2, 5mM β-

Mercaptoethanol, 2 mM PMSF and 1X protease inhibitor cocktail). Lysate was

passed through 2 layers of Miracloth to remove debris and centrifuged at 1000 x g

for 20 minutes at 4°C to pellet down cell organelles. Pellet resuspended in 5ml of

extraction buffer II (10 mM Tris-HCL pH 8, 0.25M sucrose, 10 mM MgCl2, 1% Triton

X-100, 5mM β-Mercaptoethanol, 2 mM PMSF and 1X protease inhibitor cocktail) to

burst other cell organelles and centrifuged at 1000 x g for 10mins at 4°C. The

deposited pellet was then dissolved in 300 µl of extraction buffer III (10 mM Tris-HCL

pH 8, 1.7 M sucrose, 2mM MgCl2, 0.15% Triton X-100, 5mM β-Mercaptoethanol,

2mM PMSF and 1X protease inhibitor cocktail) which was carefully layered at the

top of 600 µl extraction buffer III and centrifuged at 16,000 x g for 1hr at 4°C to

concentrate nuclei. The nuclear pellet was dissolved in 300 µl of sonication buffer,

and chromatin was sheared by sonicating for 20 cycles for 30 sec. ON/OFF.

Sonicated samples were cleared by centrifuging at 12000 x g for 10 mins. 1/10th of

the sonicated sample was kept at -20°C as input, and the rest was diluted 10-fold in

ChIP dilution buffer for immunoprecipitation. Protein-DNA complexes were

immunoprecipitated by incubating the diluted chromatin overnight at 4°C with 15 µl

of Dynabeads Protein-G (Invitrogen) equilibrated with 2 µg of anti-GFP antibody

(Takara Bio, 632381). Beads were washed for 2 x 10 mins in low salt wash buffer, 1

x 10 mins in high salt wash buffer, 1 x 10 min in LiCl wash buffer, and 2 x 10 min in

TE buffer. DNA from input and IP were eluted by incubation with 10% chelex 100-

resin (Biorad) at 95°C for 10mins followed by 20 µg of Proteinase K treatment at

37°C for 30 min. The supernatant was collected after centrifuging at 16000 x g for 2

min., followed by the precipitation of DNA through the phenol-chloroform extraction

method. Precipitated DNA was dissolved in 50 µl of nuclease-free water, and 2 µl

was used per reaction for chip-qPCR.

For the ChIP assay of BBX24/BBX25 binding on *PIF4* and its target genes promoter,

we have used six-day-old *35S:BBX24-GFP* and *35S:Myc-BBX25* seedlings grown

at 27°C and tissues were harvested at ZT2 and ZT4, respectively. Further, we have

processed the tissues in a similar way as described above for ELF3 ChIP analysis.

Primers used for ChIP-qPCR analysis are listed in Supplemental Table 1.

Yeast two-hybrid (Y2H) assay

Y2H assay was performed following the Matchmaker® Gold Yeast Two-Hybrid

System User Manual (Takara Bio). Briefly, full-length coding sequences of *BBX24*

and *BBX25* were cloned into the bait vector pGBKT7 containing DNA-Binding

Domain (DBD), while PIF4 and ELF3 were cloned into the prey vector pGADT7

having Activation Domain (AD). For checking the protein-protein interactions,

different combinations of bait and prey constructs were co-transformed into Y2H

Gold yeast strain (Clontech) using Yeastmaker™ Yeast Transformation kit

(Clontech), and transformants were selected on double dropout medium SD/-Leu/-

Trp at 30°C for three-days. For confirmation of protein-protein interactions, overnight

grown double dropout screened yeast cells were diluted to an OD_{600} of 0.5 with 4X

serial dilutions were spotted on quadruple dropout medium SD/-Leu/-Trp/-Ade/-His

supplemented with X-alpha-gal. Plates were grown at 30°C for 72 hr before taking

the picture.

Co-Immunoprecipitation

Coimmunoprecipitation assays were performed using Dynabeads™ Protein G
Immunoprecipitation Kit (Invitrogen). Approximately 500 mg of six-day-old seedlings
grown at 27°C for BBX24/BBX25-PIF4 and at 22°C for BBX24/BBX25-ELF3
immunoprecipitation, harvested at ZT2 and ZT8 were used, respectively. Harvested
seedlings were grounded in liquid nitrogen, and the powder was homogenized in 1
ml of co-immunoprecipitation buffer containing 50 mM Tris-HCL pH-8, 100 mM NaCl,
10% glycerol, 2mM DTT, 1mM EDTA, 0.1% NP-40, 1mM PMSF, and 1X protease
inhibitor cocktail. Crude protein extract was cleared by two rounds of centrifugation
at 13000 rpm for 10 mins at 4°C. An aliquot of 100 µl was boiled in 5X SDS loading
dye as an input. The remaining 900 µl protein lysate was incubated with 50 µl
Dynabeads (Invitrogen) equilibrated with anti-GFP or anti-cMyc antibody at 4°C for
6 h. Then, beads were washed, and immunoprecipitated proteins were eluted by
boiling the beads at 70°C for 10 minutes in 1X Laemmli buffer. Then, input and
immunoprecipitated protein samples were subjected to SDS-PAGE for
immunodetection. For immunoprecipitation assay of BBX24-GFP/Myc-BBX25-PIF4
polyclonal rabbit anti-GFP (Abcam, ab290, 1:5000), monoclonal mouse anti-myc
(Abcam, ab62928, 1:5000) and polyclonal goat anti-PIF4 antibodies (Agrisera
AS163955, 1:2500) were used. For BBX24-GFP/Myc-BBX25-ELF3
immunoprecipitation, monoclonal rabbit anti-GFP (Invitrogen G10362, 1:500),
monoclonal mouse anti-myc (Abcam, ab62928, 1:5000) and polyclonal rabbit anti-
ELF3 antibodies (Agrisera, AS18 4168, 1:2000) were used.

Differential Interference Contrast (DIC) Microscopy for cell length
measurement

For histological analysis, 6-day-old whole seedlings were fixed and cleared using
serial dilution of ethanol (15%, 50%, 70%, 96% and 100%), each for 15 min in a 24-
well plate. The seedlings were then kept in a cold room at 4° C overnight. On a
subsequent day, seedlings were rehydrated in an ethanol gradient (100%, 96%,
70%, 50% and 15%), each step for 15 min., after which those were washed with 0.2
M sodium phosphate buffer (pH=7). Aniline blue (Sigma-Aldrich, Germany) staining
solution was prepared as 0.1% w/v stock solution in 0.2 M sodium phosphate buffer.

Seedlings were stained with a 1:20 dilution of aniline blue stock solution with 0.2M

sodium phosphate buffer. Samples were incubated overnight at room temperature.

Seedlings were mounted onto chloral hydrate solution (chloral hydrate: glycerol:

Milli-Q water:: 8:1:2), and hypocotyls were observed using an Olympus IX81 inverted

microscope under differential interference contrast optics. Images were acquired

with cellSens Standard Software (Olympus).

Micrographs of six seedlings were obtained for each genotype, and their cell length

was considered under two different temperature conditions, 22°C and 27°C short-

day conditions. The cell length of hypocotyls was defined as the average number of

cortical cells crossing the central region of hypocotyls- the number of cells varies

with the genotype and the growth condition. Only the cells extending in the non-

dividing files were counted.

Bimolecular Fluorescent Complementation (BiFC) assay

For the BiFC assays, the *CDS* of *BBX24*, *BBX25*, *PIF4* and *ELF3* in the pENTR-

D/TOPO vectors have been recombined into pSPYNE-nYFP and pSPYCE-cYFP

vectors using LR-Clonase (Invitrogen) to obtain the fusion proteins. *BBX24* and

BBX25 were cloned into the pSPYNE vector, whereas *PIF4* and *ELF3* were cloned

in the pSPYCE vector. Sequence-verified constructs were then mobilized into

Agrobacterium tumefaciens (strain GV3101) with the help of the freeze-thaw

method. *Agrobacterium*-mediated *in planta* transient transformation onto onion bulb

as reported(Xu et al., 2014) The p19 protein of the tomato bushy stunt virus has

been used to subdue gene silencing. For infiltration, nYFP-BBX24/c-YFP-PIF4,

nYFP-BBX25/cYFP-PIF4, nYFP-BBX24/c-YFP-ELF3 and nYFP-BBX25/cYFP-ELF3

along with the p19 combination, were used with a final concentration of OD_{600} 0.6:

0.6: 1.0. After infiltration, onion bulbs were kept at 28±1 °C in the dark for 72 h. The

infected adaxial epidermis of the bulb scale was peeled and visualized with an SP8

confocal microscope (Leica, Germany) using 40x objective (ex: 488 nm for YFP, ex:

405 nm for DAPI) and deconvoluted using Leica Lightning software. Nuclei were

counterstained with DAPI (1 µg/ml). A combination of nYFP-BBX24/cYFP, nYFP-

BBX25/cYFP, nYFP/cYFP, nYFP/cYFP-PIF4 and nYFP/cYFP-ELF3 were used as

the negative controls. For the BBX24/BBX25-ELF3 BiFC assay, an Olympus IX81

epifluorescence microscope was used, and images were acquired using ImageJ Fiji

software.

Quantification of GFP fluorescence using epifluorescence microscope

GFP fluorescence of BBX24-GFP was captured using an Olympus IX81 inverted

microscope equipped with an LED lamp. Approximately ten seedlings were taken for

imaging after five days of growth in the respective conditions. The BBX24-GFP

fusion protein was excited with a 488 nm light, and the emitted fluorescence signal

was detected between 500nm and 561nm range. Images were acquired, and

fluorescence intensity was quantified using ImageJ's Fiji software. The mean

fluorescence intensity was calculated by marking the area with a freehand selection

tool. Subsequently, relative fluorescence intensity was calculated against 27°C

seedlings.

GUS Histochemical staining

For GUS-staining six-day-old 22°C and 27°C SD grown F1 seedlings of *DR5:GUS,*

BBX24-OEDR5:GUS and *BBX25-OEDR5:GUS* seedlings were harvested at ZT2

and fixed in 50mM Sodium phosphate buffer containing 2% formaldehyde and 0.05%

Triton X-100. After fixation, the seedlings were appropriately cleaned by washing

them three times with 50 mM phosphate buffer. To detect the GUS activity, seedlings

were submerged and vacuum infiltrated for 4mins in 0.5 mg/ml 5-bromo-4chloro-3-

indoyl glucuronide (X-Gluc) solution containing 50mM sodium phosphate buffer,

5mm Na2-EDTA, 1mM potassium ferrocyanide and 1mM potassium ferricyanide

following incubation at 37°C for overnight. The next day, seedlings were destained

with 75% and 95% ethanol for 3 hours each at 37°C and observed under a

stereomicroscope.

Quantification and statistical analysis

849 Hypocotyl length, cotyledon area, and protein band intensities were quantified using

850 ImageJ software (https://imagej.nih.gov/). The statistically significant difference in

the data sets was analyzed using GraphPad Prism version 8.0 (https://www.graphpad.com/). One-way ANOVA or Two-way ANOVA Tukey test was used for multiple groups of observations. Students' t-test was used to determine the significance of two groups of observations. Details of the statistical analysis can be found in the figure legends.

FUNDING

This study was supported by grants from the Department of Biotechnology (Ramalingaswami Fellowship grant, BT/RLF/Re-entry/ 28/2017), Science and Engineering Research Board (Start-up Research Grant, SRG/2019/000446), Intramural grant (IISER Kolkata), Ministry of Education (MoE/STARS-1/416) Government of India.

ACKNOWLEDGEMENTS

We thank Dr. Vinod Kumar, Prof. Salome Prat and Prof. Javier F. Botto, respectively, for the *pPIF4:PIF4-FLAG*, *35S::ELF3-YFP*(*elf3-4*) and *35S::BBX24-GFP* transgenic lines seeds. We thank all the members of Gangappa lab for fruitful discussions. B.C.M., S.S. and V.G. acknowledge the Council of Scientific and Industrial Research, Department of Biotechnology, and University Grants Commission, respectively, for their doctoral fellowships. G.U acknowledges the Postdoctoral Fellowship from IISER Kolkata (PDF/DBS/2021/025) and Science & Engineering Research Board (SERB-PDF/2022/000529). R.C. acknowledges the KVPY Fellowship from the Department of Science and Technology, Govt. of India.

AUTHOR CONTRIBUTIONS

S.N.G. conceptualized, designed and supervised the study, analyzed data, and wrote the first and subsequent drafts of the manuscript with inputs from all the authors. B.C.M. designed the study, planned and conducted most experiments, collected results, analyzed the data, and helped write the manuscript draft. S.S. generated *elf3* and *DR5:GUS*-related genetic lines, conducted phenotypic analysis, and measured cell length. V.G. assisted in raising *pif4-101*-related genetic lines with phenotypic analysis. R.C. contributed to generating novel genetic materials and

880 conducted experiments. G.U. carried out BiFC experiments. All the authors have
881 read and approved the final version of the manuscript.

DATA AVAILABILITY

883 The data presented in this study is available in the main figures and the
884 supplementary material.

SUPPLEMENTAL INFORMATION

Supplemental Figure S1. Molecular and phenotypic analysis of BBX25
overexpression transgenic lines.

Supplemental Figure S2. BBX24/BBX25 mediated thermosensory hypocotyl
growth is linked to growth-promoting hormones.

Supplemental Figure S3. BBX24/BBX25 overexpression lines show differential
sensitivity to Auxin and BR inhibitors.

Supplemental Figure S4. BBX24/BBX25 regulates the expression of hormone and
growth-related genes during the daytime.

Supplemental Figure S5. Warm temperatures promote BBX24/BBX25 transcript
and protein accumulation.

Supplemental Figure S6. ChIP analysis for BBX24/BBX25 enrichment on *PIF4* and
its target genes promoter.

Supplemental Figure S7. BBX24/BBX25 mediated thermosensory hypocotyl
growth depends on PIF4.

Supplemental Figure S8. BBX24/BBX25 are essential for PIF4-mediated regulation
of hypocotyl and cotyledon growth in response to warm temperatures.

Supplemental Figure S9. BBX24/BBX25 functions are required for PIF4-governed
hypocotyl and cotyledon growth at higher temperatures.

Supplemental Figure S10. Regulation of hypocotyl and cotyledon growth by PIF4
requires BBX24/BBX25 under LD photoperiod.

Supplemental Figure S11. ELF3 genetically interacts with BBX24 and BBX25 to
control thermosensory hypocotyl and cotyledon growth**.**

Supplemental Fi. S12. BBX24/BBX25 inhibit ELF3 function to promote PIF4-
governed thermosensory growth.

Supplemental Figure S13. ELF3-BBX24/BBX25-PIF4 module controls the
thermosensory flowering and petiole growth under SD photoperiod.

Supplemental Figure S14. BBX24/BBX25 is required to manifest petiole growth
and flowering of the *PIF4-OE* transgenic line and *elf3* mutants under LD photoperiod.

Supplemental Fig S15. BBX24/BBX25 antagonize ELF3 function.

Supplemental Table S1. List of oligonucleotides used in this study.

916
917 # REFERENCES
918

Box, M.S., Huang, B.E., Domijan, M., Jaeger, K.E., Khattak, A.K., Yoo, S.J.,
Sedivy, E.L., Jones, D.M., Hearn, T.J., Webb, A.A.R., et al. (2015). ELF3 controls
thermoresponsive growth in Arabidopsis. Curr Biol **25**:194-199.
10.1016/j.cub.2014.10.076.

Burko, Y., Willige, B.C., Seluzicki, A., Novák, O., Ljung, K., and Chory, J. (2022).
PIF7 is a master regulator of thermomorphogenesis in shade. Nat Commun **13**:4942.
10.1038/s41467-022-32585-6.

Burroughs, C.H., Montes, C.M., Moller, C.A., Mitchell, N.G., Michael, A.M., Peng,
B., Kimm, H., Pederson, T.L., Lipka, A.E., Bernacchi, C.J., et al. (2023).
Reductions in leaf area index, pod production, seed size, and harvest index drive
yield loss to high temperatures in soybean. J Exp Bot **74**:1629-1641.
10.1093/jxb/erac503.

Casal, J.J., and Questa, J.I. (2018). Light and temperature cues: multitasking
receptors and transcriptional integrators. New Phytol **217**:1029-1034.
10.1111/nph.14890.

Casal, J.J., and Balasubramanian, S. (2019). Thermomorphogenesis. Annu Rev
Plant Biol **70**:321-346. 10.1146/annurev-arplant-050718-095919.

Chow, B.Y., Helfer, A., Nusinow, D.A., and Kay, S.A. (2012). ELF3 recruitment to
the PRR9 promoter requires other Evening Complex members in the Arabidopsis
circadian clock. Plant Signal Behav **7**:170-173. 10.4161/psb.18766.

Chung, B.Y.W., Balcerowicz, M., Di Antonio, M., Jaeger, K.E., Geng, F.,
Franaszek, K., Marriott, P., Brierley, I., Firth, A.E., and Wigge, P.A. (2020). An

RNA thermoswitch regulates daytime growth in Arabidopsis. Nat Plants **6**:522-532.
10.1038/s41477-020-0633-3.

Crawford, A.J., McLachlan, D.H., Hetherington, A.M., and Franklin, K.A. (2012).
High temperature exposure increases plant cooling capacity. Curr Biol **22**:R396-397.
10.1016/j.cub.2012.03.044.

Crocco, C.D., Locascio, A., Escudero, C.M., Alabadi, D., Blazquez, M.A., and
Botto, J.F. (2015). The transcriptional regulator BBX24 impairs DELLA activity to
promote shade avoidance in Arabidopsis thaliana. Nat Commun **6**:6202.
10.1038/ncomms7202.

de Lucas, M., Daviere, J.M., Rodriguez-Falcon, M., Pontin, M., Iglesias-Pedraz,
J.M., Lorrain, S., Fankhauser, C., Blazquez, M.A., Titarenko, E., and Prat, S.
(2008). A molecular framework for light and gibberellin control of cell elongation.
Nature **451**:480-484. 10.1038/nature06520.

Delker, C., Quint, M., and Wigge, P.A. (2022). Recent advances in understanding
thermomorphogenesis signaling. Curr Opin Plant Biol **68**:102231.
10.1016/j.pbi.2022.102231.

Ding, L., Wang, S., Song, Z.T., Jiang, Y., Han, J.J., Lu, S.J., Li, L., and Liu, J.X.
(2018). Two B-Box Domain Proteins, BBX18 and BBX23, Interact with ELF3 and
Regulate Thermomorphogenesis in Arabidopsis. Cell Rep **25**:1718-1728 e1714.
10.1016/j.celrep.2018.10.060.

Favero, D.S., Kawamura, A., Shibata, M., Takebayashi, A., Jung, J.H., Suzuki,
T., Jaeger, K.E., Ishida, T., Iwase, A., Wigge, P.A., et al. (2020). AT-Hook
Transcription Factors Restrict Petiole Growth by Antagonizing PIFs. Curr Biol
30:1454-1466 e1456. 10.1016/j.cub.2020.02.017.

Fiorucci, A.S., Galvão, V.C., Ince, Y., Boccaccini, A., Goyal, A., Allenbach
Petrolati, L., Trevisan, M., and Fankhauser, C. (2020). PHYTOCHROME
INTERACTING FACTOR 7 is important for early responses to elevated temperature
in Arabidopsis seedlings. New Phytol **226**:50-58. 10.1111/nph.16316.

Franklin, K.A., Toledo-Ortiz, G., Pyott, D.E., and Halliday, K.J. (2014). Interaction
of light and temperature signalling. J Exp Bot **65**:2859-2871. 10.1093/jxb/eru059.

Franklin, K.A., Lee, S.H., Patel, D., Kumar, S.V., Spartz, A.K., Gu, C., Ye, S., Yu,
P., Breen, G., Cohen, J.D., et al. (2011). Phytochrome-interacting factor 4 (PIF4)
regulates auxin biosynthesis at high temperature. Proc Natl Acad Sci U S A
108:20231-20235. 10.1073/pnas.1110682108.

Gangappa, S.N., and Botto, J.F. (2014). The BBX family of plant transcription
factors. Trends in Plant Science **19**:460-470. 10.1016/j.tplants.2014.01.010.

Gangappa, S.N., and Kumar, S.V. (2017). DET1 and HY5 Control PIF4-Mediated
Thermosensory Elongation Growth through Distinct Mechanisms. Cell Rep **18**:344-
351. 10.1016/j.celrep.2016.12.046.

Gangappa, S.N., Berriri, S., and Kumar, S.V. (2017). PIF4 Coordinates
Thermosensory Growth and Immunity in Arabidopsis. Curr Biol **27**:243-249.
10.1016/j.cub.2016.11.012.

Gangappa, S.N., Crocco, C.D., Johansson, H., Datta, S., Hettiarachchi, C.,
Holm, M., and Botto, J.F. (2013). The Arabidopsis B-BOX protein BBX25 interacts
with HY5, negatively regulating BBX22 expression to suppress seedling
photomorphogenesis. Plant Cell **25**:1243-1257. 10.1105/tpc.113.109751.

Gray, W.M., Ostin, A., Sandberg, G., Romano, C.P., and Estelle, M. (1998). High
temperature promotes auxin-mediated hypocotyl elongation in Arabidopsis. Proc
Natl Acad Sci U S A **95**:7197-7202. 10.1073/pnas.95.12.7197.

Helfer, A., Nusinow, D.A., Chow, B.Y., Gehrke, A.R., Bulyk, M.L., and Kay, S.A.
(2011). LUX ARRHYTHMO encodes a nighttime repressor of circadian gene
expression in the Arabidopsis core clock. Curr Biol **21**:126-133.
10.1016/j.cub.2010.12.021.

Huang, Y., Xiong, H., Xie, Y., Lyu, S., Miao, T., Li, T., Lyu, G., and Li, S. (2022).
BBX24 Interacts with DELLA to Regulate UV-B-Induced Photomorphogenesis in
Arabidopsis thaliana. Int J Mol Sci **23**10.3390/ijms23137386.

Huq, E., and Quail, P.H. (2002). PIF4, a phytochrome-interacting bHLH factor,
functions as a negative regulator of phytochrome B signaling in Arabidopsis. EMBO
J **21**:2441-2450. 10.1093/emboj/21.10.2441.

Ibanez, C., Delker, C., Martinez, C., Burstenbinder, K., Janitza, P., Lippmann,
R., Ludwig, W., Sun, H., James, G.V., Klecker, M., et al. (2018). Brassinosteroids
Dominate Hormonal Regulation of Plant Thermomorphogenesis via BZR1. Curr Biol
28:303-310 e303. 10.1016/j.cub.2017.11.077.

Jiang, L., Wang, Y., Li, Q.F., Bjorn, L.O., He, J.X., and Li, S.S. (2012). Arabidopsis
STO/BBX24 negatively regulates UV-B signaling by interacting with COP1 and
repressing HY5 transcriptional activity. Cell Res **22**:1046-1057. 10.1038/cr.2012.34.

Job, N., Yadukrishnan, P., Bursch, K., Datta, S., and Johansson, H. (2018). Two
B-Box Proteins Regulate Photomorphogenesis by Oppositely Modulating HY5
through their Diverse C-Terminal Domains. Plant Physiol **176**:2963-2976.
10.1104/pp.17.00856.

Jung, J.H., Barbosa, A.D., Hutin, S., Kumita, J.R., Gao, M., Derwort, D., Silva,
C.S., Lai, X., Pierre, E., Geng, F., et al. (2020). A prion-like domain in ELF3
functions as a thermosensor in Arabidopsis. Nature **585**:256-260. 10.1038/s41586-
020-2644-7.

Jung, J.H., Domijan, M., Klose, C., Biswas, S., Ezer, D., Gao, M.J., Khattak, A.K.,
Box, M.S., Charoensawan, V., Cortijo, S., et al. (2016). Phytochromes function as
thermosensors in. Science **354**:886-889. 10.1126/science.aaf6005.

Khanna, R., Kronmiller, B., Maszle, D.R., Coupland, G., Holm, M., Mizuno, T.,
and Wu, S.H. (2009). The Arabidopsis B-box zinc finger family. Plant Cell **21**:3416-
3420. 10.1105/tpc.109.069088.

Koester, R.P., Skoneczka, J.A., Cary, T.R., Diers, B.W., and Ainsworth, E.A.
(2014). Historical gains in soybean (Glycine max Merr.) seed yield are driven by
linear increases in light interception, energy conversion, and partitioning efficiencies.
J Exp Bot **65**:3311-3321. 10.1093/jxb/eru187.

Koini, M.A., Alvey, L., Allen, T., Tilley, C.A., Harberd, N.P., Whitelam, G.C., and
Franklin, K.A. (2009). High temperature-mediated adaptations in plant architecture
require the bHLH transcription factor PIF4. Curr Biol **19**:408-413.
10.1016/j.cub.2009.01.046.

Krahmer, J., and Fankhauser, C. (2023). Environmental Control of Hypocotyl
Elongation. Annu Rev Plant Biol 10.1146/annurev-arplant-062923-023852.

Kumar, S.V., Lucyshyn, D., Jaeger, K.E., Alos, E., Alvey, E., Harberd, N.P., and
Wigge, P.A. (2012). Transcription factor PIF4 controls the thermosensory activation
of flowering. Nature **484**:242-245. 10.1038/nature10928.

Lee, S., Paik, I., and Huq, E. (2020). SPAs promote thermomorphogenesis by
regulating the phyB-PIF4 module in Arabidopsis. Development
14710.1242/dev.189233.

Legris, M., Ince, Y.C., and Fankhauser, C. (2019). Molecular mechanisms
underlying phytochrome-controlled morphogenesis in plants. Nat Commun **10**:5219.
10.1038/s41467-019-13045-0.

Legris, M., Klose, C., Burgie, E.S., Rojas, C.C., Neme, M., Hiltbrunner, A.,
Wigge, P.A., Schafer, E., Vierstra, R.D., and Casal, J.J. (2016). Phytochrome B
integrates light and temperature signals in Arabidopsis. Science **354**:897-900.
10.1126/science.aaf5656.

Li, F., Sun, J., Wang, D., Bai, S., Clarke, A.K., and Holm, M. (2014). The B-box
family gene STO (BBX24) in Arabidopsis thaliana regulates flowering time in
different pathways. PLoS One **9**:e87544. 10.1371/journal.pone.0087544.

Li, K.L., Yu, R.B., Fan, L.M., Wei, N., Chen, H.D., and Deng, X.W. (2016). DELLA-
mediated PIF degradation contributes to coordination of light and gibberellin
signalling in. Nature Communications **7**ARTN 11868

10.1038/ncomms11868.

Lippmann, R., Babben, S., Menger, A., Delker, C., and Quint, M. (2019).
Development of Wild and Cultivated Plants under Global Warming Conditions. Curr
Biol **29**:R1326-R1338. 10.1016/j.cub.2019.10.016.

Lorrain, S., Allen, T., Duek, P.D., Whitelam, G.C., and Fankhauser, C. (2008).
Phytochrome-mediated inhibition of shade avoidance involves degradation of
growth-promoting bHLH transcription factors. Plant J **53**:312-323. 10.1111/j.1365-
313X.2007.03341.x.

Ma, D., Li, X., Guo, Y., Chu, J., Fang, S., Yan, C., Noel, J.P., and Liu, H. (2016).
Cryptochrome 1 interacts with PIF4 to regulate high temperature-mediated hypocotyl
elongation in response to blue light. Proc Natl Acad Sci U S A **113**:224-229.
10.1073/pnas.1511437113.

Nakamura, S., Mano, S., Tanaka, Y., Ohnishi, M., Nakamori, C., Araki, M., Niwa,
T., Nishimura, M., Kaminaka, H., Nakagawa, T., et al. (2010). Gateway Binary
Vectors with the Bialaphos Resistance Gene,

, as a Selection Marker for Plant Transformation. Biosci Biotech Bioch **74**:1315-1319.
10.1271/bbb.100184.

Nieto, C., Lopez-Salmeron, V., Daviere, J.M., and Prat, S. (2015). ELF3-PIF4
interaction regulates plant growth independently of the Evening Complex. Curr Biol
25:187-193. 10.1016/j.cub.2014.10.070.

Nusinow, D.A., Helfer, A., Hamilton, E.E., King, J.J., Imaizumi, T., Schultz, T.F.,
Farre, E.M., and Kay, S.A. (2011). The ELF4-ELF3-LUX complex links the circadian
clock to diurnal control of hypocotyl growth. Nature **475**:398-402.
10.1038/nature10182.

Oh, E., Zhu, J.Y., and Wang, Z.Y. (2012). Interaction between BZR1 and PIF4
integrates brassinosteroid and environmental responses. Nat Cell Biol **14**:802-809.
10.1038/ncb2545.

Park, Y.J., Lee, H.J., Ha, J.H., Kim, J.Y., and Park, C.M. (2017). COP1 conveys
warm temperature information to hypocotyl thermomorphogenesis. New Phytol
215:269-280. 10.1111/nph.14581.

Peng, S., Huang, J., Sheehy, J.E., Laza, R.C., Visperas, R.M., Zhong, X.,
Centeno, G.S., Khush, G.S., and Cassman, K.G. (2004). Rice yields decline with
higher night temperature from global warming. Proc Natl Acad Sci U S A **101**:9971-
9975. 10.1073/pnas.0403720101.

Podolec, R., Wagnon, T.B., Leonardelli, M., Johansson, H., and Ulm, R. (2022).
Arabidopsis B-box transcription factors BBX20-22 promote UVR8 photoreceptor-
mediated UV-B responses. Plant J **111**:422-439. 10.1111/tpj.15806.

Qiu, Y., Li, M., Kim, R.J., Moore, C.M., and Chen, M. (2019). Daytime temperature
is sensed by phytochrome B in Arabidopsis through a transcriptional activator
HEMERA. Nat Commun **10**:140. 10.1038/s41467-018-08059-z.

Qiu, Y., Pasoreck, E.K., Yoo, C.Y., He, J., Wang, H., Bajracharya, A., Li, M.,
Larsen, H.D., Cheung, S., and Chen, M. (2021). RCB initiates Arabidopsis
thermomorphogenesis by stabilizing the thermoregulator PIF4 in the daytime. Nat
Commun **12**:2042. 10.1038/s41467-021-22313-x.

Quint, M., Delker, C., Franklin, K.A., Wigge, P.A., Halliday, K.J., and van Zanten,
M. (2016). Molecular and genetic control of plant thermomorphogenesis. Nat Plants
2:15190. 10.1038/nplants.2015.190.

Quint, M., Delker, C., Balasubramanian, S., Balcerowicz, M., Casal, J.J.,
Castroverde, C.D.M., Chen, M., Chen, X., De Smet, I., Fankhauser, C., et al.
(2023). 25 Years of thermomorphogenesis research: milestones and perspectives.
Trends Plant Sci **28**:1098-1100. 10.1016/j.tplants.2023.07.001.

Saura-Sanchez, M., Chiriotto, T.S., Cascales, J., Gomez-Ocampo, G.,
Hernandez-Garcia, J., Li, Z., Pruneda-Paz, J.L., Blazquez, M.A., and Botto, J.F.
(2023). BBX24 Interacts with JAZ3 to Promote Growth by Reducing DELLA Activity
in Shade Avoidance. Plant Cell Physiol **64**:474-485. 10.1093/pcp/pcad011.

Silva, C.S., Nayak, A., Lai, X., Hutin, S., Hugouvieux, V., Jung, J.H., Lopez-
Vidriero, I., Franco-Zorrilla, J.M., Panigrahi, K.C.S., Nanao, M.H., et al. (2020).
Molecular mechanisms of Evening Complex activity in Arabidopsis. Proc Natl Acad
Sci U S A **117**:6901-6909. 10.1073/pnas.1920972117.

Song, Z.Q., Bian, Y.T., Liu, J.J., Sun, Y., and Xu, D. (2020). B-box proteins: Pivotal
players in light-mediated development in plants. Journal of Integrative Plant Biology
62:1293-1309. 10.1111/jipb.12935.

Stavang, J.A., Gallego-Bartolome, J., Gomez, M.D., Yoshida, S., Asami, T.,
Olsen, J.E., Garcia-Martinez, J.L., Alabadi, D., and Blazquez, M.A. (2009).
Hormonal regulation of temperature-induced growth in Arabidopsis. Plant J **60**:589-
601. 10.1111/j.1365-313X.2009.03983.x.

Sun, J., Qi, L., Li, Y., Chu, J., and Li, C. (2012). PIF4-mediated activation of
YUCCA8 expression integrates temperature into the auxin pathway in regulating
arabidopsis hypocotyl growth. PLoS Genet **8**:e1002594.
10.1371/journal.pgen.1002594.

Tan, W., Chen, J., Yue, X., Chai, S., Liu, W., Li, C., Yang, F., Gao, Y., Gutiérrez
Rodríguez, L., Resco de Dios, V., et al. (2023a). The heat response regulators
HSFA1s promote Arabidopsis thermomorphogenesis via stabilizing PIF4 during the
day. Sci Adv **9**:eadh1738. 10.1126/sciadv.adh1738.

Tan, W., Chen, J., Yue, X., Chai, S., Liu, W., Li, C., Yang, F., Gao, Y., Gutierrez
Rodriguez, L., Resco de Dios, V., et al. (2023b). The heat response regulators
HSFA1s promote Arabidopsis thermomorphogenesis via stabilizing PIF4 during the
day. Sci Adv **9**:eadh1738. 10.1126/sciadv.adh1738.

Tripathi, P., Carvallo, M., Hamilton, E.E., Preuss, S., and Kay, S.A. (2017).
Arabidopsis B-BOX32 interacts with CONSTANS-LIKE3 to regulate flowering. Proc
Natl Acad Sci U S A **114**:172-177. 10.1073/pnas.1616459114.

Vaishak, K.P., Yadukrishnan, P., Bakshi, S., Kushwaha, A.K., Ramachandran,
H., Job, N., Babu, D., and Datta, S. (2019). The B-box bridge between light and
hormones in plants. J Photoch Photobio B **191**:164-174.
10.1016/j.jphotobiol.2018.12.021.

Wang, C.Q., Guthrie, C., Sarmast, M.K., and Dehesh, K. (2014). BBX19 interacts
with CONSTANS to repress FLOWERING LOCUS T transcription, defining a
flowering time checkpoint in Arabidopsis. Plant Cell **26**:3589-3602.
10.1105/tpc.114.130252.

Xu, K., Huang, X., Wu, M., Wang, Y., Chang, Y., Liu, K., Zhang, J., Zhang, Y.,
Zhang, F., Yi, L., et al. (2014). A rapid, highly efficient and economical method of
Agrobacterium-mediated in planta transient transformation in living onion epidermis.
PLoS One **9**:e83556. 10.1371/journal.pone.0083556.

Yamaguchi, N., Winter, C.M., Wu, M.F., Kwon, C.S., William, D.A., and Wagner,
D. (2014). PROTOCOLS: Chromatin Immunoprecipitation from Arabidopsis Tissues.
Arabidopsis Book **12**:e0170. 10.1199/tab.0170.

Yuan, L., Yu, Y.J., Liu, M.M., Song, Y., Li, H.M., Sun, J.Q., Wang, Q., Xie, Q.G.,
Wang, L., and Xu, X.D. (2021). BBX19 fine-tunes the circadian rhythm by interacting
with PSEUDO-RESPONSE REGULATOR proteins to facilitate their repressive
effect on morning-phased clock genes. Plant Cell **33**:2602-2617.
10.1093/plcell/koab133.

Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D.B., Huang, Y., Huang, M., Yao, Y.,
Bassu, S., Ciais, P., et al. (2017). Temperature increase reduces global yields of
major crops in four independent estimates. Proc Natl Acad Sci U S A **114**:9326-
9331. 10.1073/pnas.1701762114.

Zhao, H., and Bao, Y. (2021). PIF4: Integrator of light and temperature cues in plant
growth. Plant Sci **313**:111086. 10.1016/j.plantsci.2021.111086.

Zhao, X., Heng, Y., Wang, X., Deng, X.W., and Xu, D. (2020). A Positive Feedback
Loop of BBX11-BBX21-HY5 Promotes Photomorphogenic Development in
Arabidopsis. Plant Commun **1**:100045. 10.1016/j.xplc.2020.100045.

Zhou, Y., Park, S.-H., Soh, M.Y., and Chua, N.-H. (2021). Ubiquitin-specific
proteases UBP12 and UBP13 promote shade avoidance response by enhancing
PIF7 stability. Proceedings of the National Academy of Sciences **118**:e2103633118.
doi:10.1073/pnas.210363311

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184 **FIGURES AND FIGURE LEGENDS**

1185

1186
1187
Figure 1. BBX24 and BBX25 promote thermosensory hypocotyl growth and
gene expression.
(A) Representative images of six-day-old Col-0, *bbx24*, *bbx25*, and *bbx24bbx25*
seedlings grown at 22°C and 27°C under short-day (SD, upper panel) and long-day
(LD, lower panel) photoperiods.
(B-C) Hypocotyl length measurements (mean ± SD; n ≥ 20) of SD (**b**) and LD (**c**)
grown seedlings that are shown in **A**.
(D) Representative seedling images of Col-0, *BBX24-OE*, and *BBX25-OE* lines (line
#16) were grown at 22°C and 27°C for six days under SD (upper panel) and LD
(lower panel) conditions.
(E and F) Hypocotyl length (mean ± SD; n ≥ 20) of SD (**E**) and LD (**F**) grown seedlings
shown in (**D**).
(G) Cell length (mean ± SD; n ≥ 15) of hypocotyl epidermal cells from indicated
genotypes grown at 22°C and 27°C for six days under SD.
(H) qRT-PCR analysis (mean ± SD of three biological replicates) showing relative
transcript levels of thermoresponsive genes *YUC8*, *XTR7*, *IAA19* and *PRE5* in the

indicated genotypes that were grown for six days at 22°C and 27°C under SD, and
the samples were harvested at ZT23. *EF1α* was used as an internal control.
In Figs. **B**, **C**, **E-G**, and **h**, different lowercase letters, shown above the bars, denote
statistically significant differences between the samples (Two-way ANOVA followed
by Tukey's post-hoc test, $p < 0.05$). Numbers indicate fold changes between 22°C
and 27°C. All the experiments were repeated at least three times, and similar results
were obtained.   In Figure **A** and **D**, the scale bar represents 1 mm and 2 mm,
respectively. All the experiments were repeated three times, and similar results were
observed.
1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

Figure 2. Warm temperatures promote BBX24/BBX25 transcript and protein accumulation.

(A) qRT-PCR analysis shows *BBX24* and *BBX25* expression in six-day-old Col-0 seedlings grown at 22°C and 27°C under SD conditions, and samples were collected at indicated time points. *EF1α* was used as a loading control, and the transcript level of each sample was normalized to Col-0 22°C ZT0 time point. Error bars depict the mean ± SD of three biological replicates. Asterisks denote statistically significant differences as determined by Student's t-test, **P<0.01, ***P<0.001.

(B) Immunoblots showing BBX24-GFP protein level in six-days-old *35S::BBX24-GFP* seedlings grown at 22°C and 27°C under SD photoperiod and samples were harvested at different time points as indicated.

(C) Immunoblot showing Myc-BBX25 protein status in six-days-old *35S::Myc-BBX25* seedlings grown at 22°C and 27°C under SD, and samples were harvested at different time points as indicated.

In Figure **B** and **C**, 27°C grown Col-0 seedlings harvested at ZT2 were used as a negative control. Actin served as a loading control and was used for normalization. Relative protein levels (shown underneath each immunoblot) were calculated by

setting 22°C grown Col-0 ZT2 to 1. The experiments were repeated three times, and similar results were obtained.

(D-F) Representative images (**D**), relative GFP fluorescence intensity (**E**) and number of cells showing the fluorescence (**F**) in hypocotyls of *35S:BBX24-GFP* transgenic seedlings grown for six days after germination at 22°C and 27°C under short-day conditions. At least ten seedlings have been captured for the GFP fluorescence. Asterisks denote statistically significant differences as determined by Student's t-test, ***P<0.001. All the experiments were repeated three times, and similar results were observed.

1262

Figure 3. BBX24/BBX25 promotes the accumulation of PIF4 protein in response to warm temperatures.

(A) Immunoblot analysis of PIF4 protein levels in six-day-old Col-0 and *bbx24bbx25* seedlings grown diurnally at 22°C and 27°C under SD and samples were harvested at the indicated time (ZT) points.

(B) Immunoblot showing PIF4 protein levels in Col-0 and *bbx24bbx25* double mutant seedlings grown at 22°C SD for six days or 22°C SD grown seedlings exposed to 27°C for indicated time points. The *pifQ* mutant was used as a negative control.

(C) Immunoblot analysis of PIF4 protein level in six-day-old Col-0, *BBX24-OE*, and *BBX25-OE* seedlings grown at constant 22°C and 27°C under SD photoperiod, and tissue were harvested at ZT23.

(D) Immunoblot analysis of HY5 protein levels in six-day-old Col-0 and *bbx24bbx25* seedlings grown diurnally at 22°C and 27°C under SD and sampled at the indicated time (ZT) points.

(E and **F)** Immunoblot analysis (**E**) and relative band intensity of PIF4 protein levels (**F**) in six days-old 22°C SD grown Col-0 and *bbx24bbx25* double mutant seedlings shifted to 27°C for 24hrs with MG132 treatment for 16hrs before tissue harvest. Different lowercase letters, shown above the bars, denote statistically significant differences between the samples (Two-way ANOVA followed by Tukey's post-hoc test, $p < 0.05$).

(G and **H)** Western blot analysis (**G**) and relative band intensity of PIF4 protein degradation (**H**) in six-day-old 22°C SD grown Col-0 and *bbx24bbx25* double mutants seedlings treated with 50µM MG132 before being shifted to 27°C for 8hrs were washed five times and then transferred to liquid MS medium containing 200µM

CHX following tissue harvest at the indicated time points. Asterisks represent
statistically significant differences determined by the Student's t-test, *P<0.05,
**P<0.01.
Actin was used as a loading control in all the immunoblots. The relative PIF4 or HY5
protein level normalized to actin was shown underneath the respective immunoblot.
All the experiments were repeated three times, and similar results were observed.

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

Figure 4. BBX24/BBX25 are essential for PIF4-mediated hypocotyl elongation in response to warm ambient temperatures.

(**A** and **B**) Representative seedlings images (a) and hypocotyl length (b) of six-days-old Col-0*, bbx24bbx25, PIF4-OE*, and *bbx24bbx25PIF4-OE* seedlings grown at 22°C and 27°C under SD. The data shown is mean ± SD (n ≥ 20) seedlings. The scale bar represents 1 mm.

(**C** and **D**) Representative cotyledon images (c) and cotyledon area (d) of six-day-old indicated genotypes grown at 22°C and 27°C under SD. The data shown is mean ± SD (n ≥ 20) of. The scale bar represents 1 mm. In Figs. b and d, different lowercase letters, shown above the bars, denote statistically significant differences between the samples (Two-way ANOVA followed by Tukey's post-hoc test, p < 0.05). Numbers indicate fold changes between 22°C and 27°C.

(E and **F)** Immunoblot analysis of PIF4-FLAG protein levels in the indicated genotypes grown for six days at 22°C and 27°C, and samples were harvested at ZT23 (e), or 22°C SD grown five-day-old seedlings shifted to 27°C for 24h and sampled at ZT23 (f).

(G) Immunoblot showing PIF4-FLAG protein levels in the indicated genotypes grown under 22°C SD for six days were shifted to 27°C for 24 h, followed by MG132 treatment for 16 h before tissues were harvested. In Figs. **E-G**, Col-0 was used as a negative control. The relative PIF4 protein levels were normalized against Actin and are shown underneath each immunoblot. The normalized PIF4-FLAG protein levels in PIF4-OE at 22°C short day conditions were set to 1, and others were calculated as relative levels. We detected two bands of PIF4-FLAG, and the top band probably corresponds to the phosphorylated form of PIF4. Numbers indicate fold changes between 22°C and 27°C.

(H) Immunoblots of coimmunoprecipitation assays showing BBX24 interacts with PIF4 in planta. Total protein from six-day-old Col-0 and *35S::BBX24-GFP* seedlings grown at 27°C, sampled at ZT2, were used, and BBX24-GFP was pulled down using anti-GFP antibody coupled to magnetic Dynabeads. BBX24-GFP and PIF4 from input and pulldown fractions were detected by immunoblot using anti-GFP and native anti-PIF4 antibodies. Asterisk denote a non-specific band.

(I) Immunoblots of coimmunoprecipitation assay showing BBX25 interacts with PIF4 in planta. Total protein from six-day-old Col-0 and *35S::Myc-BBX25* seedlings grown at 27°C, sampled at ZT2, were used, and Myc-BBX25 was immunoprecipitated using an anti-myc antibody coupled to magnetic Dynabeads. Myc-BBX25 and PIF4 from input and pulldown fractions were detected by immunoblotting using anti-Myc and native anti-PIF4 antibodies.

(J) BiFC assays show the physical interaction of BBX24 and BBX25 with PIF4 in planta in onion epidermal cells. The adaxial epidermis of the onion bulb was co-infiltrated with nYFP, BBX24-nYFP, or BBX25-nYFP, and cYFP or PIF4-cYFP as indicated. Two days after incubation, onion peels were analysed for fluorescence in different combinations of co-infiltered samples.

(K) BBX24 and BBX25 physically interact with PIF4 in yeast. Overnight grown cultures of Y2H Gold yeast strain containing different combinations of bait (pGBKT7, pGBKT7-BBX24, and pGBKT7-BBX25) and prey (pGADT7 and pGADT7-PIF4) constructs diluted to an OD_{600} of 0.5 with 4x serial dilutions were spotted on double and quadruple dropout medium supplemented with X-alpha-gal. Pictures were taken after 72 h of incubation at 30°C. The experiments were repeated at least three times, and similar results were obtained. All the interactions were repeated three times, and similar results were observed.

1355

Figure 5. ELF3 represses *BBX24/BBX25* transcription during dawn while warm temperature dampens it.

(A and **B)** qRT-PCR analysis showing relative transcript levels of *BBX24* and *BBX25* in six-day-old seedlings of Ws and *elf3* grown at constant 22°C and 27°C SD conditions, sampled at ZT6 and ZT10. The expression level of each sample was normalized to Ws at 22°C of the respective time point, *and EF1α* was used as an internal control. The data shown is the mean ± SD, with Error bars depicting SD (n=3 biological replicates).

(C and **D)** Schematic representation of *BBX24* and *BBX25* promoters depicting LUX binding sites and the exact position of each PCR fragment used for analysis.

(E and **F)** ChIP-qPCR assay showing ELF3 enrichment onto *BBX24* and *BBX25* promoters in vivo. Six-day-old seedlings of *35S::ELF3-YFP* (*elf3-4*) grown at 22°C and 27°C SD conditions were sampled at ZT10 for cross-linking. For each amplicon, fold enrichment was calculated as a ratio between anti-GFP and mock (pre-immune

1371   serum). Genomic fragment towards the *3'UTR* of *BBX24* and *BBX25* was used as
1372   a negative control. The data shown is the (mean ± SD, n=2, two biological replicates.
1373   The experiment was repeated twice, and similar results were seen. Asterisks denote
1374   statistically significant differences as determined by Student's t-test, *P<0.05,
1375   **P<0.01.

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

Figure 6. BBX24/BBX25 antagonizes ELF3 to promote PIF4-mediated thermosensory growth.

(**A** and **B**) Representative seedling images and hypocotyl length of six-day-old segregated WT (Col-0xWs), *bbx24bbx25*, *elf3* and *bbx24bbx25elf3* seedlings grown at 22°C and 27°C under SD. The scale bar = 2 mm. In Figure b, different lowercase letters, shown above the bars, denote statistically significant differences between the samples (Two-way ANOVA followed by Tukey's post-hoc test, $p < 0.05$). Numbers indicate fold changes between 22°C and 27°C.

(C) Immunoblots showing PIF4 protein levels in WT (Col-0xWs), *bbx24bbx25*, *elf3* and *bbx24bbx25elf3* seedlings grown at 22°C for six days or transferred to 27°C for 24 hours and harvested at ZT23. Histone H3 was used as a loading control, and the relative levels of PIF4 protein normalized to Histone H3 are shown underneath each immunoblot.

(D) Immunoblot showing ELF3 protein levels in six-day-old Col-0, *bbx24bbx25*, *BBX24-OE* and BBX25-OE seedlings grown at constant 22°C and 27°C under short-day conditions. Actin was used as a loading control, and the relative ELF3 protein level was calculated against Col-0 at 22°C SD.

(E) Immunoblot analysis of PIF4 protein levels of the indicated genotypes was shifted to 27°C for 24 hours after growing for five days at 22°C SD conditions. Actin was used as a loading control, and the relative PIF4 protein levels shown underneath the immunoblot were calculated against Col-0.

(F and **G)** Co-immunoprecipitation assay showing in vivo protein-protein interactions between BBX24-ELF3 and BBX25-ELF3. Total protein from six-day-old Col-0, *35S:BBX24-GFP* and *35S:Myc-BBX25* seedlings grown at 22°C SD, harvested at ZT8, were used. The BBX24-GFP and BBX25-Myc proteins were pulled down using anti-GFP and anti-Myc antibodies coupled to magnetic Dynabeads. In the input and immunoprecipitated protein complex, ELF3 was detected using native anti-ELF3 antibodies. Similarly, all the proteins were detected using respective antibodies in the input and immunoprecipitated fractions. Asterisk denote a non-specific band.

(H) BiFC assays show the physical association of BBX24 and BBX25 with ELF3 in onion epidermal cells. The adaxial epidermis of the onion bulb was co-infiltrated with nYFP, BBX24-nYFP or BBX25-nYFP with cYFP or ELF3-cYFP, as indicated. After two days of incubation, onion peels were observed for fluorescence under a microscope in different co-infiltrate samples. Scale bar = 50 µm.

(I) Yeast two-hybrid assay showing BBX24 and BBX25 physical interaction with ELF3 in yeast. Y2H Gold strain harbouring different combinations of bait (pGBKT7, pGBKT7-BBX24, and pGBKT7-BBX25) and prey (pGADT7 and pGADT7-ELF3) constructs were grown overnight to an OD_{600} of 0.5 with 4x serial dilutions were dropped on double and quadruple dropout medium supplemented with X-alpha-gal. Pictures were captured after incubating the plates for three days at 30°C. The experiments were repeated at least three times, and similar results were obtained.

1441
1442
Figure 7. The molecular model depicting BBX24/BBX25-mediated regulation of thermosensory growth.

The model elucidates the molecular mechanism by which BBX24/BBX25 promotes thermosensory growth in Arabidopsis. In the absence of BBX24/BBX25, ELF3 exerts a robust inhibitory effect on PIF4 at the transcriptional and protein activity levels, resulting in compromised thermosensory responses. However, in the presence of BBX24/BBX25, they antagonize ELF3, relieving its inhibitory effect on PIF4, resulting in enhanced PIF4 function, and promoting warm temperature-mediated growth.

1451