1

Human iPSC-derived Committed Cardiac Progenitors Generate Cardiac Tissue Grafts in a Swine Ischemic Cardiomyopathy Model without Triggering Ventricular Arrhythmias

Amish N. Raval, MD[1,2]*, Eric G. Schmuck, PhD[1], Vladislav Leonov, PhD[1], Sushmita Roy, PhD[1], Yukihiro Saito, MD, PhD[1], Tianhua Zhou, MD[1], Hae Jae Jang, BS[1], Kaylee Wilker, BS[1], James Conklin, BS[2], Timothy A. Hacker, PhD[3], Chad Koonce, BS[4], Meghan Boyer, BS[4], Kristin Stack, BS[4], Ellen Hebron, BS[4], Scott K. Nagle, MD, PhD[5], Jianhua Zhang, PhD[1], Patrick C.H. Hsieh, MD, PhD[1,6,7], Timothy J. Kamp, MD, PhD[1,6]*

[1]Department of Medicine, Division of Cardiovascular Medicine, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA.

[2]Department of Biomedical Engineering, University of Wisconsin, Madison, USA

[3]Cardiovascular Research Center, University of Wisconsin-Madison School of Medicine and Public Health, Madison, USA

[4]Fujifilm Cellular Dynamics Inc., Madison, WI, USA

[5]Department of Radiology, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

[6]Department of Cell and Regenerative Biology, University of Wisconsin-Madison, USA

[7]Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan.

*To whom correspondence should be addressed:

Timothy J. Kamp, tjk@medicine.wisc.edu, 1111 Highland Ave., WIMR II 8459, Madison, WI 53705

Amish N Raval, anr@medicine.wisc.edu, H4/568, Box 3248 Clinical Science Center, 600 Highland Ave., Madison, WI 53792

<u>Short Title</u>: Committed Cardiac Progenitors for Cardiac Repair

<u>Word Count</u>: 8607

<u>Journal Subject Terms</u>: Cell Therapy, Stem cells, Heart Failure, Chronic Ischemic Heart Disease, Treatment.

ABSTRACT

Background: Intramyocardial injection of human pluripotent stem cell-derived cardiomyocytes following a myocardial infarction (MI) improves cardiac function in large animal models, but associated ventricular arrhythmias are major safety concern. We hypothesized that transendocardial injection of human induced pluripotent stem cell (hiPSC)-derived committed cardiac progenitor cells (CCPs), combined with cardiac fibroblast-derived extracellular matrix (cECM) to enhance cell retention, will generate cardiac tissue grafts improving contractility without triggering ventricular arrhythmias.

Methods: hiPSCs were differentiated using bioreactors and small molecules to produce committed cardiac progenitor cells (CCPs). MI was created using a coronary artery balloon occlusion and reperfusion model in Yucatan mini pigs. Four weeks later, epicardial needle injections of CCPs+cECM were performed in a small initial feasibility cohort (n=6), and then transendocardial injections of CCPs+cECM (n=14), CCPs alone (n=14), cECM alone (n=4) or vehicle control (n=13) into the peri-infarct region in a randomized cohort. Arrhythmias were evaluated using implanted event recorders. Magnetic resonance imaging (MRI) and invasive pressure-volume assessment were used to evaluate left ventricular anatomic and functional performance. Detailed histology was performed to detect and characterize human grafts.

Results: A scalable biomanufacturing protocol was developed generating CCPs which can efficiently differentiate into cardiomyocytes or endothelial cells in vitro. Intramyocardial delivery of CCPs to post-MI porcine hearts resulted in engraftment and differentiation of CCPs to form ventricular cardiomyocyte rich grafts. There was no significant difference in cardiac MRI-based measured cardiac volumes or function

between control, CCP and CCP+cECM groups; however, pressure-volume analysis

showed an improvement in dobutamine-stimulated functional reserve in CCP and

CCP+cECM groups. Delivery of CCPs did not result in tumors or ventricular arrhythmias.

Conclusions: Transendocardial delivery of CCPs with or without cECM into post-MI

porcine hearts resulted in comparable human cardiomyocyte grafts which did not improve

resting LV function but did improve stress-induced contractile reserve without triggering

ventricular arrhythmias.

Key words: Stem Cell, Heart Failure, Treatment, Trial, Catheter

INTRODUCTION

Currently approved medication and device therapies can blunt the progression of heart failure (HF) following myocardial infarction (MI), but the morbidity and mortality in post - MI survivors remain significant.[1,2] For patients who suffer large MIs, replacement of necrotic myocardium with fibrotic scar frequently leads to progressive heart failure, and there are currently no available treatments to replace lost cardiomyocytes and restore contractility. Therefore, cell-based therapies have been investigated for their potential to generate functional cardiac tissue in post-MI or ischemic cardiomyopathy animal models and patients. More than two decades of investigations and clinical trials with a variety of adult cell sources such as bone marrow mononuclear cells have produced no clear evidence of new functional heart muscle or consistent improvements in clinical outcomes.[3] In contrast, post-MI delivery of human pluripotent stem cell (PSC)-derived cardiomyocytes (CMs) from embryonic stem cells (ESCs) or induced pluripotent stem cells (iPSCs) has produced viable cardiomyocyte grafts in rat,[4-7] guinea pig,[8-10] swine,[11-14] and non-human primate[15-17] myocardium. However, large animal experimental studies using hPSC-CMs have been complicated by ventricular arrhythmias, relatively low cell retention in the myocardium and mixed functional results.

hPSC-derived cardiac progenitor cells are also being investigated for cardiac cell therapy in the post-MI failing heart,[18,19] but progenitor cells are a less studied cell preparation for therapy relative to hPSC-CMs. However, there are several features that make cardiac progenitors cells a promising cell source. Cardiac progenitor cells are smaller cells than hPSC-CMs and may be more amenable to manipulation and cell delivery while maintaining viability. The hPSC-derived cardiac progenitor cells are proliferative and therefore may generate larger hPSC-CM grafts per cell delivered. Importantly, cardiac

progenitor cells can differentiate to not only hPSC-CMs but also other cardiac cell

lineages such as endothelial cells and smooth muscle cells needed to generate viable

cardiac tissue grafts. The first in human heart failure treatment clinical trial using hPSCs

tested early stage hESC-derived progenitors delivered in a fibrin patch at the time of

coronary artery bypass surgery and demonstrated safety for this preparation.[18] However,

hPSCs undergo a dynamic range of cell state transitions in the process of differentiating

to terminal cardiac lineages which are variably referred to as cardiac progenitors, and the

specific characteristics of the given progenitor state chosen for cell therapy is critically

important. The optimal cardiac progenitor cell population isolated from differentiating

hPSCs and delivery strategy for cardiac repair remain to been determined.

Low cell retention is a vexing problem for all cardiac cell therapies. Our group previously

described a cardiac extracellular matrix (cECM) engineered biomaterial using high-

density culture-expanded cardiac fibroblasts isolated from human cadaveric hearts.

cECM is an acellular, uniquely fibronectin abundant, non-chemically cross-linked, non-

hydrogel scaffold with immunomodulatory and angiogenic properties that can be

formulated into particulates, reconstituted into solution, and administered as an

injectable[20,21]. cECM was shown to educate monocytes to a distinct population of

macrophages that exhibit anti-inflammatory and pro-angiogenic features,[21] and can

promote myocardial retention of mesenchymal stem cells in rodent heart disease models

when co-delivered epicardially via an open chest.[22,23] In comparison, a more clinically

appealing delivery approach is transendocardial injection (TEI). TEI uses a steerable,

needle tipped catheter that is introduced into the left ventricle via a femoral artery to

perform targeted intramuscular injections from the endocardial surface.

In this study, we tested whether human iPSC-derived committed cardiac progenitors

(CCPs) alone or with cECM could be safely delivered to post-MI failing porcine heart and

engraft to form cardiac tissue grafts that lead to improvements in functional outcomes.

We observed that the CCPs, regardless of the presence of cECM, successfully engrafted

and differentiated into cardiomyocytes, endothelial cells and fibroblasts, leading to the

formation of cardiac tissue grafts. Cardiac MRI did not show significant improvement in

chamber size or ejection fraction, but pressure-volume analysis demonstrated an

improvement in dobutamine-stimulated functional reserve by CCPs. Cell delivery did not

induce ventricular arrhythmias in the engrafted hearts.

METHODS

For detailed methods, please see the Supplemental Methods section in the Supplemental

Material. Data and study materials are available from the corresponding authors upon

reasonable request. All animal use in this study adhered to an approved protocol by the

institutional animal care and use committee at the University of Wisconsin – Madison,

protocol, M005512-R03-A01.

Statistical Analysis

Statistical analysis was carried out using GraphPad Prism version 9. MRI data was

analyzed with a 2-way ANOVA with a Sidak post-hoc analysis for multiple comparisons.

Pressure volume loops were analyzed with a One-Way ANOVA with a Bonferroni post-

hoc analysis. All data presented as mean +/- standard error of the mean unless

otherwise stated as standard deviation. Statistical significance was defined as $P<0.05$.

RESULTS

Production of human iPSC-derived CCPs

For scalable production of human iPSC-derived CCPs, iPSCs were propagated as

monolayer cultures in CELLSTACK culture chambers. Cells were singularized and

inoculated into PBS3 Bioreactors to form cellular aggregates that were differentiated in

suspension culture using an accelerated small molecule-based protocol employing

biphasic modulation of Wnt signaling along with application of a rho-kinase inhibitor and

TGF-β receptor inhibitor (**Figure 1A**). Flow cytometry analysis revealed a progressive

loss of expression of the epithelial marker, EpCAM, present on iPSCs and upregulation

of the mesodermal marker CD56 (NCAM) during the differentiation protocol (**Figure**

1B).[24] A KDR$^+$/PDGFRα$^+$ population emerges and peaks at day 4 consistent with the

generation of a cardiac progenitor population.[25,26] KDR expression subsequently

decreases on day 5 and 6 suggesting advancing maturation of these cardiac progenitors.

CCPs are identified as the CD56$^+$/PDGFRα$^+$ or CD56$^+$/CXCR4$^-$ population present on day

6. Average flow cytometry time course data from 20 runs demonstrate the reproducibility

of the differentiation protocol (**Figure 1B, Supplemental Tables 1-4)**. Analysis of gene

expression by quantitative RT-PCR over the 6-day protocol revealed loss of expression

pluripotency genes (*NANOG, SOX2, POU5F1)* and an increase in expression of cardiac

progenitor genes (*HAND2, GATA4, NKX2-5, PDGFRA, TBX5*) (**Figure 1C**). At day 6, the

aggregates are dissociated, cryopreserved and described as CCPs.

Single cell RNA-sequencing (scRNA-seq) was also performed to better characterize the

CCPs. Cryopreserved CCPs were recovered, and a total of 37,014 cells were analyzed

using the 10X Genomics Next GEM single cell 3' assay. Chromium single cell data output

from the 10X Genomics analysis pipeline were further analyzed using Seurat and

Cellxgene VIP software. Transcriptomic analyses of the CCPs identified 7 clusters, and

5 clusters were closely related (C0, C1, C2, C4, C6) (**Figure 2A)**. Analysis of differentially

expressed genes (DEGs) revealed cardiomyocyte-fated cells in clusters C0, C2, C4 and

C6 with expression of genes encoding cardiac transcription factors (*MEF2C, TBX5,*

NKX2-5), myofilaments (*ACTC1, MYL4, MYL7, MYH6*), EC coupling proteins (*RYR2*) and

ion channels (*CACNA1D, KCNIP4, SLC8A1*). These 4 cell populations account for 64%

of the total cells in the CCPs and were annotated cardiomyocyte progenitor (CMP)

(**Figure 2B,F**). Further analysis of the DEGs and the GO terms showed the C2 and C4

cell populations differentially express cell cycle marker genes *MKI67* and *TOP2A*. In

addition, C2 cells differentially express *CCNB1* (Cyclin B1) and *CENPA* in contrast to C4

cells which express *CDC6* and *FEN1* that are essential for DNA replication, indicating

capture of M phase and S phase proliferating CMPs, respectively (**Figure 2E,F,**

Supplement 2). The small cluster C6 has abundant expression of *TTN* and *RYR2*,

suggesting a differentiating cardiomyocyte (CM). Interestingly, cluster C1 that accounts

for 15% of the total CCPs has elevated expression of ribosomal genes including *RPLP1*,

RPS15 as well as genes for ATP synthase complex (*ATP5F1E*) and NADH

dehydrogenase (*NDUFS5*) in mitochondrial that are essential for protein translation and

metabolism along with lower expression of myofilament, EC coupling and ion channels

genes than the CMP populations, so we annotated this cell population as a cardiac

progenitor (CP) (**Figure 2A,F**). The C3 and C5 clusters account for 12% and 8.6% of the

total CCPs, respectively, and have distinct gene expression pattern from the CMP. The

C3 cells differentially express typical endodermal marker genes of EPCAM and FOXA2,

and C5 cells differentially express typical endothelial marker genes of CD34, CDH5

(CD144), and PECAM1 (CD31) (**Figure 2C,D,F)**. Overall, the scRNA-seq analysis

confirmed that the CCP preparation is composed primarily of cells fated to become CMs

(~80%) whereas a smaller populations of endothelial and endodermal cells are present.

In vitro potency of CCPs

To test the potency of the cryopreserved CCPs, cells were thawed and plated on vitronectin substrate in RPMI+B27 medium and characterized after 7 days of culture. After 7 days in culture in RPMI+B27, the cells differentiated into contracting sheets of iPSC-derived cardiomyocytes (iPSC-CMs) (Video supplement 1). Flow cytometry demonstrated that upon thawing 9.3±4.0% of the cells expressed low levels of sarcomeric α-actinin (SAA), but after 7 days of culture 95.9±0.8% of the differentiated cells expressed SAA consistent with differentiation primarily to iPSC-CMs (**Figure 3A**). Immunolabeling of the differentiated cells confirmed that nearly all cells were iPSC-CMs which expressed Nxk2.5, SAA, cardiac troponin I (cTnI), and cardiac troponin T (cTnT) (**Figure 3B**). Optical mapping of the contracting monolayers of iPSC-CMs loaded with a voltage-sensitive fluorescent dye, FluoVolt™, demonstrated optical action potentials with an average spontaneous rate of 1.16±0.02 Hz (n=4). Point stimulation of the cultures over a range of frequencies (1.5-3.0 Hz) showed electrical coupling between the iPSC-CMs with uniform conduction through the monolayers and conduction velocities (CV) ranging from 9.2±0.2 to 13.4±0.4 cm/s (**Figure 3C,D**). The optical action potential durations progressively shortened at higher rates of stimulation consistent with rate adaptation of cardiac action potentials (**Figure 3C,D).** Single iPSC-CMs (n=27) were characterized with perforated patch clamp recording and demonstrated action potentials when stimulated at 1 Hz with an average maximum diastolic potential (MDP) of -72.5±1.1 mV, action potential amplitude (APA) of 107.0±1.7 mV and dV/dt_{max} of 112.1±14.5 V/s **(Figure 3E, Supplemental Table 5)**. These results show that thawed CCPs plated in basal RPMI+B27 medium without additional growth factors or small molecules rapidly

differentiate predominantly to iPSC-CMs and exhibit electrophysiological properties typical of early developing human cardiomyocytes.

To determine the potency of CCPs to differentiate into iPSC-derived endothelial cells (iPSC-ECs), thawed CCPs were plated in an endothelial medium, Vasculife LS1020™. After 12 days in culture, flow cytometry demonstrated that 88.0±1.3% of cells were CD31$^+$/CD144$^+$ iPSC-ECs (**Figure 3F**). Further characterization of the CD31$^+$/CD144$^+$ population shows that 15.2±1.2% of the cells express CXCR4, a marker of arterial endothelial cells (**Figure S1**).[27,28] Immunolabeling of the day 12 cultures also demonstrated expression of CD31 in a monolayer of human iPSC-ECs (**Figure 3F**). To test the functional properties of the iPSC-ECs, a matrigel tube formation assay was carried out on replated day 12 cells. One day after replating, the iPSC-ECs organized into an interconnected network of tubes in Matrigel (**Figure 3G**). Thus, in medium optimized for EC culture, CCPs readily differentiate to iPSC-ECs.

<u>CCPs engraft in swine infarcted myocardium and differentiate to cardiac lineages</u>

A total of 70 pigs underwent LAD balloon occlusion reperfusion, of which 52 survived the post-MI period to treatment (**Figure S2**). Of the remaining 52 pigs, 6 pigs were evaluated in the roll-in feasibility cohort which underwent open-chest epicardial injections of CCPs+cECM after 4 weeks post-MI. Pigs were sacrificed 2 weeks later, and hearts were sectioned for histopathology. The other 46 pigs were randomly assigned to one of four treatment groups: CCP+cECM, CCP alone, cECM alone or Control (5% Flexbumin). Entry of pigs into the cECM alone cohort was stopped after only 3 pigs were treated due to lack of a clear trend toward improved ejection fraction and limitations on animal

numbers. Thus, the functional studies focused on comparing 3 groups: CCPs,

CCPs+cECM, and control.

In the feasibility roll-in cohort, pigs were followed for 2 weeks post cell delivery, and the

presence of engrafted human cells was assessed by immunolabeling for human nuclear

antigen, Ku80, or human mitochondria. All 6 pigs demonstrated Ku80 positive human cell

grafts that were composed largely of cTnI co-labeled cells; however, cTnI labeling

intensity was less than observed in the native pig myocardium (**Figure 4A**). Trichrome

staining of an adjacent section revealed the corresponding human cell grafts adjacent to

the native myocardium (**Figure 4B**). Human cell grafts were identified within the scar and

in adjacent native myocardium (**Figure 4C-E**). The grafts were composed predominantly

of human cardiomyocytes based on the cTnI immunolabeling, but the sarcomeric pattern

and density of labeling was consistent with immature cardiomyocytes. In addition, using

co-labeling with an antibody detecting human mitochondria and an antibody for CD31, we

detected human endothelial cells in the grafts (**Figure 4F-I**). These results demonstrate

that the CCPs can survive following injection into the post-MI porcine heart and

differentiate into cardiomyocytes and endothelial cells to form cardiac tissue grafts.

In the randomized phase, pigs were sacrificed at 4 weeks post-treatment, although a

small subset of pigs were survived up to 8 weeks after transendocardial delivery

(**Supplemental Table 6** for summary of randomized pigs and analysis). In pigs receiving

CCPs with or without cECM, hearts were systematically blocked, and sections were

immunolabeled for human Ku80 and cTnI. Human grafts were identified in 13/14 (92.9%)

of pigs in each the CCPs and CCPs+cECM cohorts. The graft area observed on the

histological sections were comparable between the CCP and CCP+cECM groups (0.68

+/- 0.25 mm^2 and 0.37 +/- 0.168 mm^2, respectively, p=0.30, supplemental **Figure S4D**).

Furthermore, the number of graft areas detected was comparable between the CCP and

CCP+cECM groups (5.8±2.2 and 5.2±1.6, p=0.57, **Supplemental Figure S4E**)

Macroscopic graft size was not measured given the blocking strategy used to enable

detailed histological characterization of the large porcine hearts. In the animals sacrificed

4 weeks after CCP delivery, we quantitatively analyzed the cellular composition of the

grafts by co-labeling for lineage specific markers and Ku80. The vast majority of Ku80$^+$

cells co-labeled with cTnI (~90%, **Figure 5A,D**) indicating cardiomyocytes. Less than 1%

of the Ku80$^+$ cells were endothelial cells based on CD31 labeling (**Figure 5B,D)** and 1-

2% of the Ku80$^+$ cells were vimentin positive stromal cells (**Figure 5C,D**). Both CCP and

CCP+cECM groups showed similar cellular compositions of the grafts. A small fraction

(<10%) of the cellular composition of grafts could be other cell types not evaluated for

including possible residual CCPs, which were not assessed given the lack of a unique

marker to distinguish them from native myocardium. There were no areas of

osteochondral differentiation.[29] Grafts were present in both CCP and CCP+cECM animals

up to the latest time point sampled, 8 weeks after delivery.

<u>CCP-derived grafts show progressive maturation of ventricular cardiomyocytes</u>

To assess the chamber-specificity of the differentiated cardiomyocytes in the grafts and

to examine for evidence of maturation, we immunolabeled sections with an antibody

recognizing the ventricular-specific myosin light chain, MLC2v at 2-, 4- and 8-weeks post-

transplant. Most of the human cells identified using an anti-human mitochondrial antibody

expressed MLC2V at all time points tested (**Figure 5G-L, Figure S5**). There was

evidence of progressive sarcomere organization and maturation based on the pattern of

MLC2v labeling from 2 weeks to 8 weeks, although the cells remained smaller and with

less sarcomere organization than adjacent adult porcine myocardium. Together these

results demonstrate that transendocardial delivery of CCPs in the post-MI swine heart leads to formation of cardiac grafts predominantly composed of human ventricular cardiomyocytes with occasional human endothelial cells and stromal cells. The grafts persist for the 8 weeks of observation in this immunosuppressed swine model and show evidence of progressive cardiomyocyte maturation.

Cardiac MRI assessment

Twenty-two pigs were evaluable for MRI in the three experimental groups, whereas 19 pigs were excluded from the MRI analysis due to: LVEF >45% (n=1), infarct size <8% of LV (n=1), hemodynamically significant opportunistic sepsis/infection (n=13), and technical issues with the MRI acquisition (n=4) (see **Supplemental Figure 1, Supplemental Table 6**). Prior to CCP delivery, there was no significant difference in baseline post-MI left ventricular ejection fraction: CCPs alone (n=5) 43.8 +/- 4.3%, and CCPs+cECM (n=9) 39.9 +/- 4.1%, and Control (n=8) 44.8 +/- 4.2%. Four weeks following transendocardial delivery, no statistically significant change in ejection fraction was observed for any of the three groups, but CCPs trended to improvement (p=0.08), while CCPs+cECM (p=0.3) and control injections (p=0.99) showed no clear improvement (**Figure 6A,B**). Similarly, there was no statistically significant difference between groups in left ventricular end-diastolic volume and left ventricular end-systolic volume following intramyocardial injection (**Figure 6C,D**). Late gadolinium enhancement-determined infarct volume was not significantly changed in any of the three groups (**Figure 6E,F**).

Pressure-volume hemodynamic analysis shows CCPs improve EF and contractility reserve

Invasive hemodynamic pressure-volume (PV) studies were performed at the endpoint of the study for each animal (**Figure 7A-C)**. Only pigs meeting the MRI inclusion criteria above (EF<45% or infarct >8% of LV, free of infection) and sacrificed at the 4-week timepoint were considered potentially evaluable by PV loop analysis. Technical problems during PV-loop acquisition or poor-quality loops caused the exclusion of 3 additional animals. Comparison of resting ejection fractions from the hemodynamic studies demonstrated a significantly higher ejection fraction in the CCP (46.1 ±1.8%) group compared to Control (37.1 ±2.0%) (p=0.007), and the CCP+cECM (43.3 ±1.3%) group showed a trend to improvement (**Figure 7D**). End systolic volume was significantly reduced in CCP (49.1 +/-3.0 ml) and CCP+cECM (51.3 +/-1.7 ml) groups compared to Control (75.3 +/-5.6 mL) (p=0.0004 ANOVA; p=0.001 and p<0.0001 respectively) (**Figure 7E**). End diastolic volume was also significantly reduced in CCP (91.0 +/-5.1 ml) and CCP+cECM (90.5 +/-2.9 ml) groups compared to the Control group (119.3 +/-6.0 ml) (p=0.0006 ANOVA; p=0.003 and p=0.001 respectively) (**Figure 7F)**. There was no significant difference in end systolic elastance between CCP, CCP+cECM and Control groups at baseline (p=0.13 ANOVA). Following dobutamine challenge to assess contractile reserve, a significant increase in elastance was observed in all groups (CCP alone (8.4 +/-0.9) and CCPs+cECM (4.5 +/-0.7) control (4.4 +/-0.7) (compared to baseline (p=0.0002, p=0.004, and 0.04 respectively) (**Figure 7G**). The CCP alone group showed significantly greater elastance in the presence of dobutamine relative to Control (p<0.006) and CCP+cECM (p=0.007).

<u>Safety evaluation shows no CCP-induced ventricular tachycardia</u>

Safety assessment for the study included evaluation of procedure related complications, gross or histological tumor formation, and arrhythmias. In all treated pigs, there were no

procedure related adverse events such as cardiac perforation, MI, stroke, or limb

ischemia. Continuous electrocardiographic monitoring for significant tachycardia (>150

bpm for greater than 24 beats) using an implanted cardiac monitor (REVEAL LINQ™)

was performed from the time of the initial MI until pigs were sacrificed, in both the initial

feasibility cohort and the randomized cohort. There was only one episode of sustained

ventricular tachycardia (>30 seconds) that self-terminated in one randomized cohort pig

treated with CCP+cECM on post-injection day 2 (**Figure 8B**). This animal (4680) also

exhibited post-MI ventricular tachycardia prior to cell delivery. There was one brief run of

ventricular tachycardia (10 s) detected in a CCP treated pig on post-injection day 11

(**Figure 8 A,E**). Otherwise, there were no episodes of ventricular tachycardia or

ventricular fibrillation detected in any of the pigs treated with CCPs followed for up to 8

weeks. Immunolabeling for the gap junction protein, connexin 43 (Cx43), did show

relatively sparse expression of Cx43 within the graft relative to native myocardium and

occasionally between graft cardiomyocytes and native myocardium (**Figure 8F)**. Finally,

necropsy did not reveal evidence of gross tumors in any animal, and there was no cardiac

histological evidence of tumor formation observed.

DISCUSSION

In this preclinical study, we tested the ability of minimally invasive TEI delivery of a novel

human iPSC cell product, CCPs, alongside a retention biomaterial to remuscularize the

porcine post-MI myocardium. The major findings are: i) transplanted CCPs differentiate

to form grafts composed primarily of cTnI and MLC2v positive ventricular cardiomyocyte

and occasional human CD31 positive endothelial cells, ii) neither transplant of CCPs or

CCPs+cECM significantly improved cardiac MRI measured ejection fraction or cardiac

volumes, but the CCP and CCP+cECM groups demonstrated significantly greater

contractile reserve relative to control by invasive hemodynamic assessment, iii) CCP +/-

cECM administration was feasible without adverse safety outcomes including no tumor

formation and no ventricular arrhythmias, and iv) human iPSC-CMs in grafts displayed

progressive maturation but remained structurally immature at 8 weeks post injection.

Together, these results demonstrate that CCPs hold promise as a therapy for the post-

MI heart.

The concept of using cardiac progenitor cells for post-MI heart failure treatment has been

investigated in a variety of contexts. The adult heart does not contain a pool of stem or

progenitor cells with potency to generate new heart muscle,[30,31] but hPSCs provide an

accessible and clinically relevant source of cardiac progenitor cells. However, hPSC-

derived cardiac progenitor cell populations are heterogenous and dynamic during

differentiation akin to cardiac development in which different progenitor populations

gradually give rise to distinct heart regions and structures. An early study using *in vitro*

engineered heart tissue comparing mouse ESC-CMs and ESC-derived Flk1$^+$/PDGFRα$^+$

cardiac progenitor cells demonstrated that the cardiac progenitor cells differentiated to

CMs and improved the electrical coupling in the engineered heart tissue more effectively

than the ESC-CMs.[32] Using similar cell preparations from hESCs, a study comparing

hESC-CMs to KDR$^+$/PDGFRα$^+$ cardiovascular progenitor post-MI in a nude rat model

showed comparable grafts and functional effects from the two populations.[33] A

developmentally earlier mesodermal SSEA1$^+$ cell population derived from rhesus ESCs

was shown to engraft and differentiate to cardiomyocytes, populating 20% of the scar

region in the post-MI rhesus heart.[34] However, a more recent study of the xenogeneic

transplantation of hESC-derived SSEA1$^+$ cells post-MI nonhuman primate hearts found

no human cells remaining at 140 days, although an improvement in cardiac function was

observed and attributed to decreased apoptosis of endogenous cardiomyocytes.[35]

Likewise, transplantation of hESC-derived SSEA1[+] cells in fibrin patches in a rat MI model

showed improvement in LV function despite lack of survival of the transplanted cells,

suggesting a paracrine mechanism of action.[36] Studies using an intermediate stage

cardiac progenitor population first identified by high levels of Isl1 expression showed in

rhesus heart slice models that the cardiac progenitors are highly migratory and can track

to areas of cryoinjury.[37] In post-MI porcine model, the same cardiac progenitors engrafted

and formed cardiomyocyte rich grafts, but there was no statistically significant

improvement in MRI measures of LV function after 12 weeks.[37] Yap et al.,[19] provided

evidence that late-stage or committed cardiac progenitors can be effectively delivered at

the time of surgically induced MI, engraft and differentiate to relevant cardiac lineages

forming cardiomyocytes rich grafts lasting at least 2 months with evidence for

improvement in EF from cardiac MRI.[19] However, the study by Yap et al., found evidence

for ventricular arrhythmias following cell delivery in approximately half of the treated

animals.[19] In the present study, we used late stage CCPs, but unlike prior studies they

were differentiated from hiPSCs rather than hESCs. Furthermore, although scRNA-seq

analysis showed heterogeneity in our CCP preparation, more than 80% of the cells were

cardiomyocyte-fated in contrast to the Yap et al. which identified 33% of the cells as

cardiac progenitors.[19] The present study used clinically relevant TEI delivery one month

following infarct in contrast to prior studies using surgical, epicardial injection in the acute

or subacute phase of MI. Comparable to earlier studies, we observed engraftment and

differentiation to form cardiac muscle grafts, but there was not an improvement in cardiac

function on cardiac MRI evaluation despite an improvement in dobutamine recruitable

functional reserve. The mechanism underlying the improvement in functional reserve

could reflect a less advanced stage of heart failure in which β-adrenergic signaling is less

desensitized which may be due to a paracrine mechanism or a direct contribution of

engrafted cells unmasked by β-adrenergic stimulation. In summary, studies using hPSC-

derived cardiac progenitor cells in animal models post-MI demonstrate the ability of the

cells to engraft and differentiate with variable evidence for improvement in cardiac

function and risk of ventricular arrhythmias.

Significant safety concerns due to ventricular arrhythmias following delivery of hPSC-CMs

by epicardial injection in large animal post-MI heart failure models have arisen, although

the arrhythmias are generally transient, starting within 1 week after transplant and lasting

approximately one month.[11,15,17] Current evidence suggests that these ventricular

arrhythmias are the result of rapid intrinsic automaticity arising from a subpopulation of

engrafted CMs rather than reentry mechanisms.[13,16] The sustained ventricular

tachycardia has been tolerated in nonhuman primate models following cell therapy, but

in porcine models, there has been significant mortality observed in treated animals.[11,38]

Pharmacological treatment with ivabradine and amiodarone reduced the engraftment

arrhythmia burden but did not eliminate it.[38] Given that abnormal automaticity is the

presumed basis for these ventricular arrhythmias, gene editing was used to alter

expression of 4 different ion channels and transporters to eliminate automaticity.[14] This

approach greatly reduced engraftment arrhythmias, although there was still an early

period of sustained ventricular tachycardia. The arrhythmic risk for cardiac progenitor

cells is less clear as some of the large animal studies did not directly monitor for

arrhythmias.[34] However, Yap et al. observed that approximately half of the treated pigs

with CCPs developed sustained ventricular tachycardia starting 9 days or later and lasting

at least a month and in one animal up to week 12, which appears slightly delayed relative

to the hPSC-CM studies. This potentially reflects the additional time needed for

differentiation of CCPs. In contrast, the present study observed one relatively short

episode of nonsustained ventricular tachycardia in all of the CCP and CCP+cECM treated

animals which was in an animal exhibiting runs of ventricular tachycardia prior to cell

therapy. The mechanisms underlying the minimal arrhythmic burden in the present CCP

study will require further investigation, and potential considerations include differences in

the cell population, TEI vs. epicardial delivery, graft size or localization, and time after MI

for therapy. Our working model is that gradually differentiating CMs from the CCPs

integrate into the native myocardium in smaller, distributed endocardial grafts entrained

by the surrounding myocardium without coalescing into large grafts. Future mechanistic

studies of electrical integration of the predominantly endocardial cellular grafts resulting

from TEI will require alternative approaches from previously successful surface optical

mapping of whole heart epicardial grafts.[16] Furthermore, differences in arrhythmia

surveillance approaches may contribute to some differences in observed arrhythmia

burden between studies as investigations using implanted loop recorders such as this

one, rather than continuous telemetry, will not detect slower ventricular tachycardia (<150

bpm in this study). Ultimately identifying the key factors that eliminate or minimize the

arrhythmia risk is necessary for this therapy to advance successfully to the clinic.

Combining cell products with matrix preparations or synthetic hydrogels has been

demonstrated in previous studies to improve cell retention and survival in post-MI hearts.

For example, the pro-survival cocktail used in several studies includes the complex matrix

preparation, Matrigel™.[5] Gelatin hydrogels have been used successfully in the

transplantation of iPSC-CM cardiomyocytes spheroids in post-MI animal models.[39] In this

study, we tested the addition of an injectable formulation of cECM to the CCP product,

but this did not result in improved functional benefits at the cECM dose delivered;

however, we did find an increase in the number of epicardial grafts on the surface of the

heart in the cECM+CCP group relative to CCP alone. This may be related to cECM

helping retain any cells that get into pericardial space. How this potential differential

localization of grafts impacts long-term outcomes will require additional study. In addition,

cECM has been proposed to have favorable immunomodulatory effects via alternatively

activated macrophage education,[21] which could theoretically make local, ischemic tissue

more hospitable for co-transplanted cells. However, these favorable influences may be

blunted in this immunosuppressed animal model. Overall, the tested delivery of cECM did

not improve functional outcomes in this study, but additional studies are needed to

determine the optimally formulated and dosed delivery vehicle to promote cell survival

and engraftment.

TEI delivery has emerged as an appealing, minimally invasive approach for direct

intramyocardial transplantation of cells, biomaterials and other biologic agents in a

targeted fashion. In a recent pooled analysis, this delivery approach was observed to be

overall safe in >1800 patients injected in clinical trials worldwide[40]. In this study, the

NOGA/Myostar (Biosense Webster/J&J) mapping and TEI catheter platform was used to

perform targeted injections; however, this platform was recently discontinued from

production by the manufacturer. Alternative cardiac mapping approaches to facilitate

accurate targeting with TEI, such as co-registered multimodality imaging, have been

described but remain investigational.[41] Similarly, alternative investigational TEI catheters,

such as the helical needle-tipped "Helix" catheter (BioCardia, Sunnyvale, CA), are being

actively used in active adult cell therapy trials[42,43]. These novel cardiac visualization

methods and catheter designs may pave the way to help inject a new generation of

therapeutic cells designed to remuscularize the damaged heart.

Overall, the optimal hPSC cell product and delivery strategy to remuscularize the heart

remains to be determined. Even a combination of cell products may be preferable such

as hiPSC-derived endothelial cells and hiPSC-CMs to promote cell survival and

revascularization.[44] Ground breaking phase 1 clinical trials are testing hPSC-CMs to treat

heart failure employing epicardial injections,[45] transcatheter injections (NCT04982081)

and epicardial patches (NCT04945018, NCT04396899) are underway world-wide as well

as a completed trial demonstrating the safety of hESC-SSEA1[+] progenitor cells

transplanted at the time of coronary artery bypass grafting surgery.[18] Despite these

developments, important unanswered questions remain including concerns about

immunogenicity, repeat dosing possibilities, electromechanical integration,

arrhythmogenicity, and whether minimally invasive transcatheter administration will be

feasible for wider adoption of this therapy.

Nevertheless, early phase clinical trials are underway testing the safety of delivering

hPSC-CMs in patients with heart failure with reduced ejection fraction either by direct

intramuscular injection or transplantation of cellularized sheets to the epicardium

[clinicaltrials.gov NCT03763136; NCT04945018; NCT03763136; NCT04696328;

NCT04945018].

<u>Limitations</u>

Following cardiac transplantation of the CCPs, we identified the progeny of the CCPs to

predominantly include cardiomyocytes and more rare endothelial cells and fibroblasts;

however, we cannot exclude other small populations of cell types including residual

CCPs. We did not observe tumor formation which suggests that the cell preparation did

not include pluripotent stem cells capable of forming teratomas. The dosing of CCPs was

based on the literature and pilot studies of epicardial injected CCPs, so improvements in

dosing and repeat dosing could positively impact outcomes. The current study did not

provide long-term follow-up of the animals, in part due to the challenge of opportunistic

infections in the setting aggressive immunosuppression. There was qualitative evidence

of maturation over 2 months based on myofilament organization, but no quantitative

measure of maturation was made. Infections also caused a reduction in the planned

sample size. Electrical integration of the cells was not directly examined in this study, but

it will be an important subject of future research to define the mechanisms leading to lack

of arrhythmias in our study relative to hPSC-CMs studies.

Acknowledgements:

Sincere thank you to Allison Rodgers, Everett Temme, Sarah El Meanawy and Vivian

Hacker for animal handling and procedures, and Emily Ferris for MRI acquisition and

analyses.

Sources of Funding

Funding support provided by:

1. National Institutes of Health 5U01HL148690-02

2. National Institutes of Health T32 HL007936

3. Fujifilm Cellular Dynamics, Inc, Madison, WI

4. Cellular Logistics, Inc., Madison, WI

5. University of Wisconsin-Madison

Disclosures

ANR: Research or educational grants from Fujifilm Cellular Dynamics, Siemens Healthcare, BioCardia. Consultant for Blue Rock Therapeutics and Novo Nordisk. Co-founder with equity interest in Cellular Logistics, Inc.

EGS: Co-founder with equity interest in Cellular Logistics, Inc.

CK, MB, KS, and EH are employees of Fujifilm Cellular Dynamics, Inc.

Figure 1. Differentiation of human iPSCs to CCPs. A) Differentiation schematic to

manufacture CCPs with images showing starting aggregates of human iPSCs and final
aggregates of CCPs. **B)** Time course of differentation characterizing cell surface protein
expression of CD56, KDR, PDGFRα, and EpCAM using flow cytometry from a
representative experiment with average data from 7-20 different runs, error bars ±S.D. **C)**
Gene expression of pluripotent and cardiac-specific genes at day 0 (iPSCs) and day 6
(CCPs) relative to the ribosomal protein *RPL32*. Error bars ±S.E.M., ***$p<0.005$ by t-test
comparison.

Figure 2. Single cell RNA-sequencing analysis of CCPs. A) UMAP projection for CCPs identifies 7 cell clusters annotated based on differential gene expression. **B-E)** Embedding plots for expression of cardiac marker genes in CMP, endothelial cell, endodermal cell, and proliferating cell, respectively. Scale is log2 fold change. **F)** Dot plot

showing expression of representative marker genes used to annotate the clusters. CMP: cardiomyocyte progenitor; CP: cardiac progenitor; CM: cardiomyocyte.

Figure 3. **In vitro differentiation of CCPs to iPSC-CMs and iPSC-ECs**. **A)** Flow cytometry analysis of CCPs immediately after thaw and following 7 d of culture on vitronectin in RPMI+B27 for sarcomeric α-actinin (SAA) expression. **B)** Immunolabeling of differentiated CCPs after 7 d of culture in RPMI+B27 for cardiomyoctye markers Nkx2.5, SAA, cardiac troponin I (cTnI) and cardiac troponin T (cTnT). **C)** Optical action potentials measured at 1.5, 2.0, and 3.0 Hz stimulation from differentiated iPSC-CM monolayers loaded with Fluovolt™ with conduction maps shown to right. **D)** Average data from optical action potentials for conduction velocity (CV) and action potential duration at 50% (APD50) and 80% (APD80) of repolarization. **E)** Average perforated patch clamp recorded action potential in red and SD in blue for 27 iPSC-CMs. **F)** Flow cytometry for endothelial markers CD31 and CD144 after 12 days culture of CCPs in Vasculife LS1020

medium and immunolabeling (bottom left panel) for CD31. **F)** Dissociated day 12 cells in Vasculife LS1020 plated for one day in undiluted Matrigel form tubes demonstrated by phase contrast micrograph.

Figure 4. iPSC-CCPs engraft and differentiate to cardiomyocytes and endothelial cells 2 weeks after delivery in post-MI swine heart. A) Epifluorescence image of porcine heart section with human graft identified by Ku80 immunolabeling and cardiomyoctyes by cTnI immunolabeling. **B)** Light microscopic image of adjacent histological section stained with Mason's Trichrome. **C-E)** Immunofluorescence images of human graft in porcine myocardium immunolabeled as indicated identifies human cardiomyocytes. **F-I)** Immunofluorescence image of another human graft in porcine

554 myocardium labeled with anti-Ku80 and anti-CD31 to identify human endothelial cells in
555 the graft.

Figure 5. CCP graft composition and maturation. A) Immunofluorescence image of 4-week CCP graft dual immunolabeled for cTnT (cardiomyocytes) and Ku80+ to detect human cells with DAPI nuclear labeling. **B)** Immunofluorescence image of 4-week CCP graft dual immunolabeled for human CD31 (endothelial cells) and Ku80+ with DAPI

indicating nuclei. Arrowhead indicates example of human endothelial cells (CD31$^+$ and
Ku80$^+$). **C)** Immunofluorescence image of 4-week CCP graft dual immunolabeled for
vimentin (stromal/fibroblasts cells) and Ku80$^+$ to identify human cells. Arrowhead
indicates example of human stromal cell (vimentin$^+$ and Ku80$^+$). **D)** Summary data for
graft composition of cardiomyocytes (cTnT), endothelial cells (CD31), and
stromal/fibroblast (vimentin) per animal (% of total Ku80$^+$ cells) at 4-weeks after CCP or
CCP+cECM delivery. Bar is average and error bars are SEM, n=4-9. There are no
significant differences between CCPs and CCP+cECM for any of the markers. **G-L)**
Confocal immunofluorescence images of CCP-derived grafts at time points indicated
post-delivery immunolabeled for ventricular-specific MLC2v and human-specific
mitochondrial antibody with DAPI to identify nuclei. There is progressive maturation of the
MLC2v myofilament pattern with increased organization and density of myofilaments with
time. Upper row scale bar 200 μm, and lower row expanded inset from indicated region
in white box with scale bar 10 μm. Individual channels for panels J,K,L are in
supplemental Figure S5.

Figure 6. Cardiac MRI analysis. A) Representative diastolic (upper) and systolic (lower) cardiac cine MR images from Control, CCP only and CCP+cECM treated animals. B-D) Comparisons of baseline and 4 weeks after treatment of LVEF, LVEDV and LVESV showed no statistically significant differences between groups. E) Representative cardiac MRI viability images. F) Comparisons of baseline and 4 weeks MRI infarct size showed no significant differences.

CCP = committed cardiac progenitor cells, cECM = cardiac extracellular matrix, LVEF = left ventricular ejection fraction, LVEDV = left ventricular end diastolic volume, LVESV = left ventricular end systolic volume.

Figure 7. Invasive hemodynamic analysis demonstrates CCP and CCP+cECM show reduced left ventricular volumes and improved contractile reserve. A-C) Representative pressure-volume loop recordings of baseline-dobutamine stress from Control, CCP and CCP-cECM treated pigs at 4 weeks following treatment. **D)** LVEF was significantly greater in the CCP group compared to Control. **E)** LVEDV was significantly reduced in CCP and CCP+ECM relative to Control. **F)** LVESV was significantly reduced in CCP and CCPP+ECM groups relative to Control. **G)** End systolic elastance determined from pressure-volume recordings at baseline and after dobutamine show increased contractile reserve in the CCP and CCP+cECM groups.

CCP = committed cardiac progenitor cells, cECM = cardiac extracellular matrix, LVEF = left ventricular ejection fraction, LVEDV = left ventricular end diastolic volume, LVESV = left ventricular end systolic volume *p≤0.05, **p≤0.01, ***p≤0.0001

 Figure 8. **CCP delivery does not result in ventricular tachycardia. A-D)** Plots show

the daily cumulative time each animal is in ventricular tachycardia (defined as >24 beats
at rate >150 bpm) per day for each of the 4 randomized groups with recordings starting
the day of MI and day 0 being the day of treatment. **E)** Representative ECG recording of
the onset of a 10 s run of ventricular tachycardia detected animal 507 on day 11. S
indicates sinus complex and V indicates ventricular complex. CCP = committed cardiac
progenitor cells, cECM = cardiac extracellular matrix. **F)** Confocal immunofluorescence
image of a graft demonstrating Cx43 labeling within host myocardium, in graft and
between graft and host using antibodies to against cTNI, Ku80, and Cx43 with DAPI
nuclear labeling. Region in white box is at increased magnification to right.

REFERENCES

1. Tsao CW, Aday AW, Almarzooq ZI, Alonso A, Beaton AZ, Bittencourt MS, Boehme AK, Buxton AE, Carson AP, Commodore-Mensah Y, et al. Heart Disease and Stroke Statistics-2022 Update: A Report From the American Heart Association. *Circulation*. 2022;145:e153-e639. doi: 10.1161/CIR.0000000000001052

2. Rana JS, Khan SS, Lloyd-Jones DM, Sidney S. Changes in Mortality in Top 10 Causes of Death from 2011 to 2018. *J Gen Intern Med*. 2021;36:2517-2518. doi: 10.1007/s11606-020-06070-z

3. Zhang J, Bolli R, Garry DJ, Marban E, Menasche P, Zimmermann WH, Kamp TJ, Wu JC, Dzau VJ. Basic and Translational Research in Cardiac Repair and Regeneration: JACC State-of-the-Art Review. *J Am Coll Cardiol*. 2021;78:2092-2105. doi: 10.1016/j.jacc.2021.09.019

4. Riegler J, Tiburcy M, Ebert A, Tzatzalos E, Raaz U, Abilez OJ, Shen Q, Kooreman NG, Neofytou E, Chen VC, et al. Human Engineered Heart Muscles Engraft and Survive Long Term in a Rodent Myocardial Infarction Model. *Circ Res*. 2015;117:720-730. doi: 10.1161/CIRCRESAHA.115.306985

5. Laflamme MA, Chen KY, Naumova AV, Muskheli V, Fugate JA, Dupras SK, Reinecke H, Xu C, Hassanipour M, Police S, et al. Cardiomyocytes derived from human embryonic stem cells in pro-survival factors enhance function of infarcted rat hearts. *Nat Biotechnol*. 2007;25:1015-1024. doi: nbt1327 [pii] 10.1038/nbt1327

6. Caspi O, Huber I, Kehat I, Habib M, Arbel G, Gepstein A, Yankelson L, Aronson D, Beyar R, Gepstein L. Transplantation of human embryonic stem cell-derived cardiomyocytes improves myocardial performance in infarcted rat hearts. *J Am Coll Cardiol*. 2007;50:1884-1893. doi: 10.1016/j.jacc.2007.07.054

7. Park SJ, Kim RY, Park BW, Lee S, Choi SW, Park JH, Choi JJ, Kim SW, Jang J, Cho DW, et al. Dual stem cell therapy synergistically improves cardiac function and vascular regeneration following myocardial infarction. *Nature communications*. 2019;10:3123. doi: 10.1038/s41467-019-11091-2

8. Weinberger F, Breckwoldt K, Pecha S, Kelly A, Geertz B, Starbatty J, Yorgan T, Cheng KH, Lessmann K, Stolen T, et al. Cardiac repair in guinea pigs with human engineered heart tissue from induced pluripotent stem cells. *Science translational medicine*. 2016;8:363ra148. doi: 10.1126/scitranslmed.aaf8781

9. Shiba Y, Fernandes S, Zhu WZ, Filice D, Muskheli V, Kim J, Palpant NJ, Gantz J, Moyes KW, Reinecke H, et al. Human ES-cell-derived cardiomyocytes electrically couple and suppress arrhythmias in injured hearts. *Nature*. 2012;489:322-325. doi: 10.1038/nature11317

10. Dhahri W, Sadikov Valdman T, Wilkinson D, Pereira E, Ceylan E, Andharia N, Qiang B, Masoudpour H, Wulkan F, Quesnel E, et al. In Vitro Matured Human Pluripotent Stem Cell-Derived Cardiomyocytes Form Grafts With Enhanced Structure and Function in Injured Hearts. *Circulation*. 2022;145:1412-1426. doi: 10.1161/CIRCULATIONAHA.121.053563

11. Romagnuolo R, Masoudpour H, Porta-Sanchez A, Qiang B, Barry J, Laskary A, Qi X, Masse S, Magtibay K, Kawajiri H, et al. Human Embryonic Stem Cell-Derived Cardiomyocytes Regenerate the Infarcted Pig Heart but Induce Ventricular Tachyarrhythmias. *Stem cell reports*. 2019;12:967-981. doi: 10.1016/j.stemcr.2019.04.005

12. Zhao M, Nakada Y, Wei Y, Bian W, Chu Y, Borovjagin AV, Xie M, Zhu W, Nguyen T, Zhou Y, et al. Cyclin D2 Overexpression Enhances the Efficacy of Human Induced Pluripotent Stem Cell-Derived Cardiomyocytes for Myocardial Repair in a Swine Model of Myocardial Infarction. *Circulation*. 2021;144:210-228. doi: 10.1161/CIRCULATIONAHA.120.049497

13. Selvakumar D, Clayton ZE, Prowse A, Dingwall S, Kim SK, Reyes L, George J, Shah H, Chen S, Leung HHL, et al. Cellular heterogeneity of pluripotent stem cell-derived cardiomyocyte grafts is mechanistically linked to treatable arrhythmias. *Nat Cardiovasc Res*. 2024;3:145-165. doi: 10.1038/s44161-023-00419-3

655   14.   Marchiano S, Nakamura K, Reinecke H, Neidig L, Lai M, Kadota S, Perbellini F, Yang X, Klaiman
656   JM, Blakely LP, et al. Gene editing to prevent ventricular arrhythmias associated with
657   cardiomyocyte cell therapy. *Cell Stem Cell*. 2023;30:396-414 e399. doi:
658   10.1016/j.stem.2023.03.010
659   15.   Chong JJ, Yang X, Don CW, Minami E, Liu YW, Weyers JJ, Mahoney WM, Van Biber B, Cook SM,
660   Palpant NJ, et al. Human embryonic-stem-cell-derived cardiomyocytes regenerate non-human
661   primate hearts. *Nature*. 2014;510:273-277. doi: 10.1038/nature13233
662   16.   Liu YW, Chen B, Yang X, Fugate JA, Kalucki FA, Futakuchi-Tsuchida A, Couture L, Vogel KW, Astley
663   CA, Baldessari A, et al. Human embryonic stem cell-derived cardiomyocytes restore function in
664   infarcted hearts of non-human primates. *Nat Biotechnol*. 2018;36:597-605. doi:
665   10.1038/nbt.4162
666   17.   Shiba Y, Gomibuchi T, Seto T, Wada Y, Ichimura H, Tanaka Y, Ogasawara T, Okada K, Shiba N,
667   Sakamoto K, et al. Allogeneic transplantation of iPS cell-derived cardiomyocytes regenerates
668   primate hearts. *Nature*. 2016;538:388-391. doi: 10.1038/nature19815
669   18.   Menasche P, Vanneaux V, Hagege A, Bel A, Cholley B, Parouchev A, Cacciapuoti I, Al-Daccak R,
670   Benhamouda N, Blons H, et al. Transplantation of Human Embryonic Stem Cell-Derived
671   Cardiovascular Progenitors for Severe Ischemic Left Ventricular Dysfunction. *J Am Coll Cardiol*.
672   2018;71:429-438. doi: 10.1016/j.jacc.2017.11.047
673   19.   Yap L, Chong LY, Tan C, Adusumalli S, Seow M, Guo J, Cai Z, Loo SJ, Lim E, Tan RS, et al.
674   Pluripotent stem cell-derived committed cardiac progenitors remuscularize damaged ischemic
675   hearts and improve their function in pigs. *NPJ Regen Med*. 2023;8:26. doi: 10.1038/s41536-023-
676   00302-6
677   20.   Schmuck EG, Roy S, Dhillon A, Walker S, Spinali K, Colevas S, Zhou T, Chhabra G, Liu Y, Kader
678   Sagar MA, et al. Cultured cardiac fibroblasts and myofibroblasts express Sushi Containing
679   Domain 2 and assemble a unique fibronectin rich matrix. *Exp Cell Res*. 2021;399:112489. doi:
680   10.1016/j.yexcr.2021.112489
681   21.   Roy S, Spinali K, Schmuck EG, Kink JA, Hematti P, Raval AN. Cardiac fibroblast derived matrix-
682   educated macrophages express VEGF and IL-6, and recruit mesenchymal stromal cells. *J
683   Immunol Regen Med*. 2020;10. doi: 10.1016/j.regen.2020.100033
684   22.   Schmuck EG, Mulligan JD, Ertel RL, Kouris NA, Ogle BM, Raval AN, Saupe KW. Cardiac fibroblast-
685   derived 3D extracellular matrix seeded with mesenchymal stem cells as a novel device to
686   transfer cells to the ischemic myocardium. *Cardiovascular engineering and technology*.
687   2014;5:119-131. doi: 10.1007/s13239-013-0167-1
688   23.   Schmuck EG, Hacker TA, Schreier DA, Chesler NC, Wang Z. Beneficial effects of mesenchymal
689   stem cell delivery via a novel cardiac bioscaffold on right ventricles of pulmonary arterial
690   hypertensive rats. *Am J Physiol Heart Circ Physiol*. 2019;316:H1005-H1013. doi:
691   10.1152/ajpheart.00091.2018
692   24.   Evseenko D, Zhu Y, Schenke-Layland K, Kuo J, Latour B, Ge S, Scholes J, Dravid G, Li X, MacLellan
693   WR, et al. Mapping the first stages of mesoderm commitment during differentiation of human
694   embryonic stem cells. *Proc Natl Acad Sci U S A*. 2010;107:13742-13747. doi:
695   10.1073/pnas.1002077107
696   25.   Kattman SJ, Witty AD, Gagliardi M, Dubois NC, Niapour M, Hotta A, Ellis J, Keller G. Stage-specific
697   optimization of activin/nodal and BMP signaling promotes cardiac differentiation of mouse and
698   human pluripotent stem cell lines. *Cell Stem Cell*. 2011;8:228-240. doi: S1934-5909(10)00703-4
699   [pii] 10.1016/j.stem.2010.12.008

700   26.   Zhang J, Tao R, Campbell KF, Carvalho JL, Ruiz EC, Kim GC, Schmuck EG, Raval AN, da Rocha AM,
701   Herron TJ, et al. Functional cardiac fibroblasts derived from human pluripotent stem cells via
702   second heart field progenitors. *Nature communications*. 2019;10:2238. doi: 10.1038/s41467-
703   019-09831-5

27. Ivins S, Chappell J, Vernay B, Suntharalingham J, Martineau A, Mohun TJ, Scambler PJ. The
CXCL12/CXCR4 Axis Plays a Critical Role in Coronary Artery Development. *Developmental cell*.
2015;33:455-468. doi: 10.1016/j.devcel.2015.03.026
28. Zhang J, Chu LF, Hou Z, Schwartz MP, Hacker T, Vickerman V, Swanson S, Leng N, Nguyen BK,
Elwell A, et al. Functional characterization of human pluripotent stem cell-derived arterial
endothelial cells. *Proc Natl Acad Sci U S A*. 2017;114:E6072-E6078. doi:
10.1073/pnas.1702295114
29. Jebran AF, Seidler T, Tiburcy M, Daskalaki M, Kutschka I, Fujita B, Ensminger S, Bremmer F,
Moussavi A, Yang H, et al. Engineered heart muscle allografts for heart repair in primates and
humans. *Nature*. 2025;639:503-511. doi: 10.1038/s41586-024-08463-0
30. Sultana N, Zhang L, Yan J, Chen J, Cai W, Razzaque S, Jeong D, Sheng W, Bu L, Xu M, et al.
Resident c-kit(+) cells in the heart are not cardiac stem cells. *Nature communications*.
2015;6:8701. doi: 10.1038/ncomms9701
31. van Berlo JH, Kanisicak O, Maillet M, Vagnozzi RJ, Karch J, Lin SC, Middleton RC, Marban E,
Molkentin JD. c-kit+ cells minimally contribute cardiomyocytes to the heart. *Nature*.
2014;509:337-341. doi: 10.1038/nature13309
32. Song H, Yoon C, Kattman SJ, Dengler J, Masse S, Thavaratnam T, Gewarges M, Nanthakumar K,
Rubart M, Keller GM, et al. Interrogating functional integration between injected pluripotent
stem cell-derived cells and surrogate cardiac tissue. *Proc Natl Acad Sci U S A*. 2010;107:3329-
3334. doi: 10.1073/pnas.0905729106
33. Fernandes S, Chong JJH, Paige SL, Iwata M, Torok-Storb B, Keller G, Reinecke H, Murry CE.
Comparison of Human Embryonic Stem Cell-Derived Cardiomyocytes, Cardiovascular
Progenitors, and Bone Marrow Mononuclear Cells for Cardiac Repair. *Stem cell reports*.
2015;5:753-762. doi: 10.1016/j.stemcr.2015.09.011
34. Blin G, Nury D, Stefanovic S, Neri T, Guillevic O, Brinon B, Bellamy V, Rucker-Martin C, Barbry P,
Bel A, et al. A purified population of multipotent cardiovascular progenitors derived from
primate pluripotent stem cells engrafts in postmyocardial infarcted nonhuman primates. *J Clin
Invest*. 2010;120:1125-1139. doi: 40120 [pii]10.1172/JCI40120

35. Zhu K, Wu Q, Ni C, Zhang P, Zhong Z, Wu Y, Wang Y, Xu Y, Kong M, Cheng H, et al. Lack of
Remuscularization Following Transplantation of Human Embryonic Stem Cell-Derived
Cardiovascular Progenitor Cells in Infarcted Nonhuman Primates. *Circ Res*. 2018;122:958-969.
doi: 10.1161/CIRCRESAHA.117.311578
36. Bellamy V, Vanneaux V, Bel A, Nemetalla H, Emmanuelle Boitard S, Farouz Y, Joanne P, Perier
MC, Robidel E, Mandet C, et al. Long-term functional benefits of human embryonic stem cell-
derived cardiac progenitors embedded into a fibrin scaffold. *J Heart Lung Transplant*.
2015;34:1198-1207. doi: 10.1016/j.healun.2014.10.008
37. Poch CM, Foo KS, De Angelis MT, Jennbacken K, Santamaria G, Bahr A, Wang QD, Reiter F,
Hornaschewitz N, Zawada D, et al. Migratory and anti-fibrotic programmes define the
regenerative potential of human cardiac progenitors. *Nature cell biology*. 2022;24:659-671. doi:
10.1038/s41556-022-00899-8
38. Nakamura K, Neidig LE, Yang X, Weber GJ, El-Nachef D, Tsuchida H, Dupras S, Kalucki FA,
Jayabalu A, Futakuchi-Tsuchida A, et al. Pharmacologic therapy for engraftment arrhythmia
induced by transplantation of human cardiomyocytes. *Stem cell reports*. 2021;16:2473-2487.
doi: 10.1016/j.stemcr.2021.08.005
39. Kawaguchi S, Soma Y, Nakajima K, Kanazawa H, Tohyama S, Tabei R, Hirano A, Handa N, Yamada
Y, Okuda S, et al. Intramyocardial Transplantation of Human iPS Cell-Derived Cardiac Spheroids
Improves Cardiac Function in Heart Failure Animals. *JACC Basic Transl Sci*. 2021;6:239-254. doi:
10.1016/j.jacbts.2020.11.017

40. Raval AN, Pepine CJ. Clinical Safety Profile of Transendocardial Catheter Injection Systems: A
Plea for Uniform Reporting. *Cardiovasc Revasc Med*. 2021;22:100-108. doi:
10.1016/j.carrev.2020.06.031
41. Tomkowiak MT, Klein AJ, Vigen KK, Hacker TA, Speidel MA, VanLysel MS, Raval AN. Targeted
transendocardial therapeutic delivery guided by MRI-x-ray image fusion. *Catheter Cardiovasc*
Interv. 2011;78:468-478. doi: 10.1002/ccd.22901
42. Raval AN, Cook TD, Duckers HJ, Johnston PV, Traverse JH, Abraham WT, Altman PA, Pepine CJ.
The CardiAMP Heart Failure trial: A randomized controlled pivotal trial of high-dose autologous
bone marrow mononuclear cells using the CardiAMP cell therapy system in patients with post-
myocardial infarction heart failure: Trial rationale and study design. *Am Heart J*. 2018;201:141-
148. doi: 10.1016/j.ahj.2018.03.016
43. Raval AN, Johnston PV, Duckers HJ, Cook TD, Traverse JH, Altman PA, Dhingra R, Hematti P,
Borrello I, Anderson RD, et al. Point of care, bone marrow mononuclear cell therapy in ischemic
heart failure patients personalized for cell potency: 12-month feasibility results from CardiAMP
heart failure roll-in cohort. *Int J Cardiol*. 2021;326:131-138. doi: 10.1016/j.ijcard.2020.10.043
44. Cheng YC, Hsieh ML, Lin CJ, Chang CMC, Huang CY, Puntney R, Wu Moy A, Ting CY, Herr Chan DZ,
Nicholson MW, et al. Combined Treatment of Human Induced Pluripotent Stem Cell-Derived
Cardiomyocytes and Endothelial Cells Regenerate the Infarcted Heart in Mice and Non-Human
Primates. *Circulation*. 2023;148:1395-1409. doi: 10.1161/CIRCULATIONAHA.122.061736
45. Zhang H, Xue Y, Pan T, Zhu X, Chong H, Xu C, Fan F, Cao H, Zhang B, Pan J, et al. Epicardial
injection of allogeneic human-induced-pluripotent stem cell-derived cardiomyocytes in patients
with advanced heart failure: protocol for a phase I/IIa dose-escalation clinical trial. *BMJ Open*.
2022;12:e056264. doi: 10.1136/bmjopen-2021-056264