1   TITLE: Inferring a role for programmed cell death during the origin and evolution of wing
2   polyphenism in ants
3
4
5

6   Lisa Hanna[1], Brendon E. Boudinot[2,3], Jürgen Liebig[4], and Ehab Abouheif[1]*
7
8
9   [1]Department of Biology, McGill University, Montreal, QC, Canada
10
11   [2]Department of Entomology, National Museum of Natural History, Smithsonian
12   Institution, Washington, D.C., U.S.A.
13
14   [3]Entomology III, Division of Terrestrial Zoology, Senckenberg Naturmuseum Frankfurt,
15   Germany.
16
17   [4]School of Life Sciences, Arizona State University, Tempe, Arizona, U.S.A.
18
19   *Corresponding author: Ehab Abouheif (ehab.abouheif@mcgill.ca)

40   Keywords: apoptosis, ants, wing polyphenism, gene regulatory network, wing imaginal disc,
41   major evolutionary transitions, ponerines, formicines, ancestral state reconstruction

42   **ABSTRACT**
43
Major evolutionary transitions in individuality, such as the evolution of multicellularity, occur
when solitary individuals unite to form a single replicating organism with a division of labor
between constituent individuals. Programmed Cell Death (PCD) has been proposed to play an
important role during the origin and evolution of major transitions. Yet, it remains unclear to
what extent PCD was involved in the major evolutionary transition to eusociality in ants, where
solitary individuals united to form eusocial colonies with a division of labour between
morphologically distinct queen and worker castes. Wing polyphenism is one of the principle
morphological differences between queen and worker castes within a colony and is defined as
the ability of an egg (same genome) to develop either into a queen with wings or into a worker
that is completely wingless in response to environmental cues, such as nutrition and
temperature. Because wing polyphenism evolved once at the origin of ants, it was likely
intimately linked with the origin and evolution of eusociality. Here we test whether PCD plays a
role in wing polyphenism in species from across the ant phylogeny encompassing taxa with
both ancestral-like and derived characteristics. We show that PCD, mediated by the apoptosis
pathway, is present in the rudimentary wing discs of worker larvae in 15 out of the 16 species
tested. Using ancestral state reconstruction, we infer the presence of PCD in the last common
ancestor of the poneroid and formicoid clades as well as all extant ants. Based on these results,
our analysis predicts a role for PCD in regulating wing polyphenism during the evolutionary
origin of ants. Implications for the role PCD in major evolutionary transitions to individuality are
discussed.
64
65

INTRODUCTION

Major evolutionary transitions in individuality occur when solitary individuals unite to form a single replicating individual. During the history of life, such transitions include the evolution of eukaryotic cells from the integration of solitary prokaryotic cells, the integration of single-celled organisms to form multicellular organisms, the integration of solitary organisms to form eusocial colonies, and the integration of distantly-related organisms to form obligate endosymbioses (Buss, 2014; Maynard-Smith & Szathmary, 1997; West et al., 2015). Despite great advances made in understanding the evolutionary principles underlying these major transitions, the developmental mechanisms that facilitated these events remain poorly explored.

Recent advances in understanding the molecular and developmental processes during major transitions to multicellularity, has led to the proposal that programmed cell death (PCD) is a mechanism that plays an essential and general role in the emergence of major evolutionary transitions (Durand et al., 2019; Huettenbrenner et al., 2003). Unlike other modes of cell death, PCD is a genetically regulated process that occurs under specific physiological conditions (Galluzzi et al., 2018). PCD has ancient origins and is frequently found in unicellular eukaryotes and prokaryotes in addition to multicellular life (Ameisen, 2002). PCD is thought to play a role in facilitating major transitions by enhancing cooperation, formation of a morphological division of labor, and mediation of communication and conflict (Durand et al., 2019; Huettenbrenner et al., 2003; Libby & Ratcliff, 2014). For example, in some unicellular clonal or aggregate colonies, some individuals undergo PCD under stress conditions to provide nutrition to other members of the clonal group hence enhancing the survival and growth of the clones and thereby promoting group living – a precursor to multicellularity (Durand et al., 2014; Durand et al., 2011; Orellana et al., 2013). Moreover, experimental studies inducing multicellularity in yeast showed that the multicellular clusters fracture (via some cells undergoing PCD) allowing them to propagate into daughter cells (Ratcliff et al., 2012). Finally, in honeybees, PCD functions to degenerate ovaries in worker-destined larvae but not in queen-destined larvae (Durand et al., 2019; Hartfelder & Steinbruck, 1997). This led Durand et al. (2019) to propose that PCD may have played a role in the origin of reproductive division of labor in bees, and therefore, in a major transition to eusociality.

Here we test whether or not PCD contributed to the major evolutionary transition to eusociality in ants (family Formicidae) – a group that evolved eusociality only once (Boudinot, Richter, et al., 2022; Holldobler & Wilson, 1990). Concomitant with the evolution of eusociality, ants evolved a wing polyphenism, in which an egg can develop either into a winged queen or wingless worker in response to environmental cues, such as nutrition and temperature, that in turn influence hormone levels such as juvenile hormone (Hanna & Abouheif, 2021; Penick et al., 2012; Wheeler, 1986) (Fig. 1a). Wing polyphenism is a universal trait in ants (excepting for secondary losses) that evolved once at the origin of the group (Boudinot, Khouri, et al., 2022; Boudinot, Richter, et al., 2022; Hanna & Abouheif, 2021; Peeters, 2012; Perrichot et al., 2008). Ecological factors might have led to selection for a wing polyphenism in ants, where the wingless worker caste engaged in ground foraging and nesting while the winged reproductive

caste participated in nuptial flights. In the early lineages of ants, queens and workers are
thought to have similar reproductive capacities, and the reproductive division of labor primarily
occurs through behavioral regulation (Peeters, 1991). Therefore, the early evolution of wingless
workers may have played an important role in the evolution of eusociality by limiting worker's
participation in nuptial flight and dispersal, which in turn limited the ability of workers to mate
and reproduce while also facilitating cooperation in brood rearing and foraging. Here, we ask if
PCD contributed to a major evolutionary transition in eusociality by playing a key role in wing
polyphenism in ants by degenerating the wings in the worker caste and limiting worker
dispersal.
119
Individuals in the worker caste of ants, although completely wingless as adults, transiently
develop wing discs during larval development (Abouheif & Wray, 2002; Dewitz, 1878;
Rajakumar et al., 2018; Wheeler & Nijhout, 1981) (Fig. 1b–d, blue arrowheads). These wing
discs are therefore considered as rudimentary organs, which Hall (2003) defines as 'an
embryonic primordium of a more fully developed feature (the insect wing) found in an ancestor
(a winged solitary wasp)'. Examining the highly conserved wing GRN in queen wing discs and
worker rudimentary wing discs in derived ant species has revealed that the wing GRN is
conserved between winged reproductive ants and other winged insects, but is interrupted in
the wingless worker castes of different ant species (Abouheif & Wray, 2002; Shbailat &
Abouheif, 2013; Shbailat et al., 2010). However, in two Amblyoponine species from the genus
Mystrium, which are thought to reflect the ancestral characteristics of ants during the origin of
eusociality, no interruptions were detected (Behague et al., 2018). This finding led Behague et
al (2018) to propose that apoptosis (the most common genetic pathway for PCD) was co-opted
to eliminate wing discs in the wingless worker caste during the origin of eusociality in ants.
Although apoptosis has been previously detected in the worker rudimentary wing discs of two
ant species (Sameshima et al., 2004; Shbailat et al., 2010), its presence across the ants and at
the origin of wing polyphenism has yet to be tested.
137
The main goal of our study is to test for the role of PCD in wing polyphenism across the ant
phylogeny. We do this by focusing on the two major clades of ants: the 'poneroid' and
'formicoid' clades. Species in the poneroid clade, like in the genus *Mystrium*, are of key
importance to understanding the mechanism that may have been present in the early history of
wing polyphenism, since these species exhibit several characteristics that are generally
considered ancestral within ants (Burchill & Moreau, 2016; Keller & Peeters, 2022; Peeters,
1997; Wilson & Holldobler, 2005). Such characteristics are low queen-worker dimorphism, small
colony size, and predatory behavior (Keller & Peeters, 2022; Wilson & Holldobler, 2005). In
contrast, species in the formicoid clade largely exhibit derived social characteristics, such as
high queen-worker dimorphism and subcaste diversification (Holldobler & Wilson, 1990;
Wilson, 1953) We first test for the presence of apoptosis markers across a series of
developmental stages in *Harpegnathos saltator*, a focal species with ancestral characteristics
belonging to the poneroid clade. Second, we test for the presence of apoptosis markers in early
and late stage larvae in an additional 15 species across the ant phylogeny, belonging to both
the poneroid and formicoid clades. Finally, we conduct an ancestral state reconstruction
analysis to infer whether or not PCD in wing polyphenism is present at ancestral nodes and at

the base of all extant ants. Studying the mechanism(s) operating during the origin of wing
polyphenism, and therefore, during or near the evolution of eusociality, will provide us with
insight on molecular and developmental mechanisms contributing to the emergence of a major
transitions in individuality during evolutionary history.
158
159 **RESULTS**
160
In *H. saltator*, we first tested for the presence of PCD in the rudimentary wing discs of worker-
destined larvae using the TUNEL assay, a marker of apoptotic cell death that detects DNA
fragmentation (Kyrylkova et al., 2012). We focused on the last larval instar (terminal stage)
beginning from the early-mid to the larval-pupal transition (Fig. 2). We found small amounts of
DNA degradation at the early and mid-periods in both the rudimentary wing pouch and hinge
(Fig. 2a', b' white arrowheads; Fig. 1b). Subsequently, during the terminal period, the amount of
DNA degradation increases substantially (Fig. 2c', d') and by the larval-pupal transition, TUNEL
signal is present at high levels throughout the entire rudimentary wing disc (Fig. 2e'). We then
tested for the presence of PCD in the wing discs of male-destined larvae, which will develop
into adult males with fully functional wings. We found that during the early and mid-period, the
amount and intensity of the TUNEL signal in the wing discs in male-destined larvae is much
lower as compared to that in the rudimentary wing discs of worker-destined larvae at similar
stages (Fig. 2f'–i' white arrowheads, compare with Fig. 2a'–d'). Moreover, during the larval-
pupal transition stage, where TUNEL expression is at its highest in the worker rudimentary wing
discs (Fig. 2e'), we find no signs of DNA degradation in the male wing discs (Fig. 2j'). Together,
these results suggest that the strong levels of DNA degradation in the worker rudimentary wing
discs relative to male wing discs is responsible for degrading the left-over rudiment after
growth has ceased, leading to the emergence of wingless worker adults. In contrast, the
presence of some DNA degradation in the male wing discs suggest that it may play a role in
shaping the final adult wings in males.
181
We next tested for PCD (TUNEL) in the rudimentary wing discs of worker-destined larvae across
15 additional species from five major subfamilies of ants (Fig. 3; Supp. 1). We then conducted
an ancestral state estimation analysis using data from all species tested to infer whether PCD
was present or absent in the rudimentary wing discs of workers in the last common ancestor of
ants. Like in *H. saltator*, we found positive TUNEL signal in the rudimentary wing discs during
late larval development in two other species belonging to the poneroid clade: *Odontomachus*
brunneus (Ponerinae) and *Stigmatomma pallipes* (Amblyoponinae) (Fig. 3a–c'). Moreover, we
observed a strong positive TUNEL signal in late-stage rudimentary worker wing discs in 12 out
of the 13 species we sampled from the formicoid clade (Fig. 3; Supp. 1). In 12 formicoid species,
TUNEL signal is predominantly present at high levels during the larval-pupal transition stage
(Fig. 3e', f', g', h', i', j', k'; Supp. 1), and in four species, it is also detected as early as the mid
period during last larval stage (Fig. 3f, h, i, j, arrowheads). *Tetramorium immigrans*
(Myrmicinae), which possesses very small rudimentary wing discs, is the only species where we
do not observe TUNEL signal at any stage examined (Fig. 3k–k'). Our ancestral state estimation
(under the equal rates model)) using data from all species tested strongly supports the presence
of PCD (>95% confidence) as being ancestral to all living ants, including the poneroformicine

clade (= poneroids + formicoids) (Supp. 2). Our inference remains robust after accounting for:
(1) uncertainty in the presence or absence of PCD in species from missing ant subfamilies; (2)
the absence of data for *Martialis* and Leptanillinae (the sister group to all other Formicidae);
and (3) different taxon sampling regimes (Supp. 2). Altogether, our experimental data and
ancestral state estimation analyses infer role for PCD in the regulation of wing polyphenism in
the last common ancestor of all extant ants.
204
Finally, we tested which specific forms of PCD are active in rudimentary wing discs in workers of
three ant species that belong to three different subfamilies. Although the TUNEL assay is a
general marker for apoptotic cell death, we sought to confirm the presence of apoptosis by
testing for the presence of the active form of caspase 3, the human ortholog of the *Drosophila*
ICE (Drice) protein. (Xu et al., 2009). We found strong cleaved-caspase 3 expression in the
rudimentary wing discs in workers of: *H. saltator* (Ponerinae), *Camponotus floridanus*
(Formicinae), and *Pheidole dentata* (Myrmicinae). In worker-destined larvae of *H. saltator* and
C. floridanus, cleaved-caspase 3 expression in the rudimentary wing disc starts around mid-
terminal larval development and spreads throughout the remaining larval stage (Fig. 4a–a', c–
c'). In *P. dentata*, we observe cleaved-caspase 3 expression only towards the late stages of
terminal development (Fig. 4d–d'). To test for the presence of autophagy, a type of PCD
characterized by the presence of vesicles (Yan et al., 2020), we used two different markers: atg8
and LC3A/B. The rudimentary wing discs of *H. saltator* and *P. dentata* workers show no
detectable signs of autophagy (Fig.4 b–b', e–e'). These experiments confirm that PCD in the
rudimentary wing discs of workers across ants is carried out via the apoptosis pathway.
220
221   **DISCUSSION**
222
PCD is a process that has been repeatedly co-opted to play a role in the regulation of
alternative phenotypes, such as polyphenism and sexual dimorphism, across the tree of Life
(Hanna & Abouheif, 2023). Here we show that PCD has again been co-opted to play a role in the
regulation of wing polyphenism in ants, a nearly universal trait in the group that evolved
concomitantly with eusociality in ants. We demonstrate that PCD through apoptosis, is strongly
expressed in late-stage rudimentary wing discs of workers in species sampled from across the
phylogeny of ants (Fig. 5 green circles). The presence of apoptosis in the rudimentary wing discs
of 15 out of 16 species we sampled across the poneroid and formicoid clades strongly suggests
that this process evolved once and is conserved across ants. Using formal ancestral state
estimation, we infer that PCD is ancestral to the poneroformicines (= poneroid + formicoid
clades) and the extant Formicidae as a whole (Fig. 6 green circles). This predicts PCD is present
and operating in the 'basal-most' subfamilies (Leptanillinae and Martialinae; see, e.g.,
Romiguier et al. (2022), for which wing polyphenism in females is also known to occur
(Borowiec et al., 2011; Chen et al., 2017; Hsu et al., 2017). In this context, rediscovery of
Martialinae will be of particular value, given that only two workers and a few males have been
collected to date (Boudinot, 2015; Rabeling et al., 2008). Furthermore, the occurrence of wing
polyphenism in females has been recently confirmed to be present among the extinct stem
lineages of ants (Boudinot, Khouri, et al., 2022), which supports PCD as a derived condition for
all Formicidae, extant and extinct (Fig. 5 dotted green circle). Future developmental studies

focusing on species in the Leptanillinae will be able to confirm the presence of PCD during the
split of all extant ants.
244
PCD has contributed to the emergence of major evolutionary transitions in Earth's history by
playing key roles in communication, conflict mediation, division of labor, and cooperation
(Durand et al., 2019). Here we show that PCD very likely played a role in the origin of wing
polyphenism in ants, and therefore, in the major evolutionary transition to eusociality in ant. By
helping to generate a wingless worker caste, PCD may have been critical in reinforcing the
reproductive division of labor in ancestral ant societies by limiting dispersal of workers, and
therefore, their participation in mating flights. Multiple ecological, evolutionary, and
developmental factors influenced the evolution of eusociality in ants and in other eusocial
organisms. Here, we show that PCD may have been one of these factors during the origin of
eusociality in ants.
255
Finally, many wasp taxa that are closely related to ants evolved a sexual wing dimorphism,
where the females are wingless or short winged and the males are fully winged (e.g., XXX), or
evolved wing polyphenism in females (*e.g.*, some Ichneumonidae, Cynipidae, Chalcidoidea, and
Bethylidae). Given the overwhelming role of PCD in polyphenism and sexual dimorphisms
across the tree of life (Hanna & Abouheif, 2023), we predict that it may be operating in wasp
wing development to generate differences between the males and females or between female
morphs within a species. This is a question of great interest because if PCD is present in these
taxa, it would raise the intriguing possibility that wing polyphenism in ants may have originated
from a latent potential for sexual wing dimorphism and polyphenism.
265
266   **MATERIALS AND METHODS**
267
Ant collection and colony care
Ant colonies were housed in plastic boxes lined with either fluon or talc powder. Artificial nests
were constructed using glass test tubes half-filled with water and plugged with cotton. All
colonies were maintained at 25 °C, 70% humidity and a 12h day:night cycle. Ants were fed a
combination of mealworms, crickets, fruit flies, fruits, and Bhatkar-Whitcomb diet (Bhatkar &
Whitcomb, 1970). *Harpegnathos saltator* colonies were housed in plaster nests and fed live
crickets only 2-3 times per week.
275
Ant colonies were obtained from the following locals. *Stigmatomma pallipes* were collected at
McGill Gault Nature Reserve (Quebec, Canada). *Lasius niger, Tetramorium immigrans,*
Aphaenogaster rudis, and *Aphaenogaster picea* were collected at Mont Royal Park and McGill
Gault Nature Reserve (Quebec, Canada). *Odontomachus brunneus* and *Camponotus floridanus*
were collected at Gainesville (Florida, USA). *Pheidole dentata* were collected at Gainesville
(Florida, USA) and Austin (Texas, USA). *Pheidole hyatti* were collected from Tempe (Arizona,
USA). *Pheidole pallidula* were collected from Lyon (France). *Pheidole noda, Tetraponera*
rufonigra, and *Polyrhachis rastellata* were collected from Mae Tang (Chiang Mai, Thailand). *Atta*
colombica and *Acromyrmex echinatior* were collected from Gamboa (Panama). *Harpegnathos*

saltator colonies were propagated in the laboratory since the original collection in 1999 from
different locations in India.
287
288 **Immunohistochemistry, hybridization chain reaction, and TUNEL stainings**
289
290 **Fixation and Dissection**
291 Terminal stage larvae were collected and fixed as previously described by Shbailat and Abouheif
292 (2013). Fixed larvae were dissected using a Zeiss Discovery V12 stereomicroscope to expose the
293 wing discs and remove any obstructing fat tissue.
294
Immunohistochemistry
The following primary antibodies were used for immunohistochemistry: anti-cleaved caspase-3
(1:100-1:200, Cell Signaling Technology, #9661) and anti-LC3A/B (1:100, Cell Signaling
Technology, #4108). For all immunohistochemistry assays, fluorescent secondary anti-rabbit
polyclonal Alexa Fluor-555 (AbCam) antibody was used at 1:500 dilution to detect the primary
antibody, according to Khila and Abouheif (2008).
301
Hybridization Chain Reaction
To detect mRNA expression of *H. saltator atg8* we utilized the hybridization chain reaction
(HCR) methodology (Choi et al., 2018). Probe sets, amplifiers, and buffers were purchased from
Molecular Instruments, Inc. Procedure followed based on the Protocols for HCR^{TM} RNA-FISH
(v3.0) (generic sample in solution) acquired from molecularinstruments.com. Prior to
procedure, samples were fixed and dissected as described above, rehydrated in 25%, 50%, 75%
and 100% PTw (1X PBS; 0.1% Tween 20) and permeabilized in PTw and 2% Triton X-100 for 1
hour. Larvae were post-fixed in 4% formaldehyde in SSCT (5X sodium chloride sodium citrate;
0.1% Tween 20) for 2 hours and then washed 3 times with SSCT before moving into glycerol.
311
TUNEL
To detect programmed cell death in the wing discs of all species, we used the TUNEL (Terminal
deoxynucleotidyl transferase dUTP nick end labeling) assay. We used the *In Situ* Cell Death
Detection Kit, TMR red (Roche). Larval samples were fixed, dissected and stored in 100%
methanol in -30 °C prior to procedure. Dissected larvae were then rehydrated in 25%, 50%,
75%, and 100% PTw (1X PBS; 0.1% Tween 20). Samples were then washed 3 times (10 min
each) in freshly made PBT (1X PBS; 0.1% Triton; 0.1% BSA) and permeabilized in PTw and 2%
Triton X-100 for 1 hour, followed by 5 min washes in PBT. Positive and negative controls were
incubated in DNase buffer at 37°C for 10 minutes and subsequently washed with PBT for 5 min
(Supp. 3). Enzyme mix and label solution were added to the samples and the positive controls
(only label solution was added to negative controls) and incubated in the dark at 37 °C for 1
hour followed by 3 washes in PTw 5 min each. Samples were stained with DAPI (1:1000) for 1
hour to overnight and followed by gradual moving into glycerol. Samples were stored in final
concentration of 85% glycerol/DAPI before mounting and imaging.
326
327 **Imaging**

328 Fluorescent images of larval rudimentary wing discs of all species utilized in this study were
329 taken using a Zeiss AxioImager Z1 microscope.
330
Ancestral state estimation
To reconstruct the ancestral conditions of PCD, we conducted a series of ancestral state
estimation (ASEs) analyses given prior and new experimental data. To ensure that our estimates
were based on well-supported phylogenetic results, we used the favored phylogram (P) and
chronogram (C) of Romiguier et al. (2022), particularly as a recent simulation study has
demonstrated the importance of branch lengths for ancestral state modeling (Wilson et al.,
2022). We pruned the two trees (P, C) to three alternative taxon sets: (P1, C1) the first included
only those terminals that match species with experimental data (17 taxa), (P2, C2) the second
included experimental terminals plus *Martialis* and a leptanilline (22 taxa), and (P3, C3) the
third included all ant terminals from the Romiguier et al. (2022) dataset in order to account for
phylogenetic uncertainty (74 taxa). Because we had surplus experimental sampling of
Aphaenogaster and *Pheidole* with identical results, we included only one species of the former
and two species of the latter, as this matched the phylogenomic sampling of Romiguier et al.
(2022). We also excluded non-ant outgroups as there is no applicable experimental data, thus
insufficient information to parameterize a phylogenetic model that includes these taxa. As the
'all rates different' model overfit the data, resulting in complete statistical separation, we only
report results under the 'equal rates' model.
348

349 **ACKNOWLEDGMENTS**
350 We thank members of the Abouheif lab for comments. This work was supported by a Natural
351 Sciences and Engineering Research Council (NSERC) Discovery Grant to E.A., and a Doctoral
352 Fellowship from Fonds de Recherche du Quebec-Nature et Technologies (FRQNT) to L.H.

353
354

355　Figure 1

371
Figure 1. Wing polyphenism in the ponerine ant *H. saltator*: (a) environmental cues determine the development of winged queens (wings indicated in yellow) or wingless workers and reproductive workers 'gamergates'. (b) a cartoon illustration of *H. saltator* larva during terminal development indicating the location of the wing discs (blue arrowheads) and the general axes and regions of the wing disc in ants. Dissected terminal stage larvae in *H. saltator* stained with DAPI showing: (c) wing discs in winged males and (d) rudimentary wing discs in wingless female workers (blue arrowheads).

Figure 2

Figure 2. Programmed cell death in the wing discs of wingless workers and winged male larvae of *H. saltator*. (a-e) Larval developmental series in wingless female workers from early/mid terminal stage to larval-pupal transition stained with DAPI. (a'-e') a magnified view of a wing disc from the larva in the panel directly above showing TUNEL signal (red) and DAPI (blue). White arrowheads in (a', b') indicate area with TUNEL expression. (f-j) DAPI staining of a larval developmental series of winged males during terminal larval development corresponding in stage to those shown in the workers. (f'-j') a magnified view of a wing disc from the larva in the panel directly above showing TUNEL signal (red) and DAPI (blue). White arrowheads in (a', b' and f'-j') indicate region where TUNEL is expressed. Yellow dashed boxes in (a-e) and (f-j) indicate the wing disc magnified in the panels below. Images in panels (a-e) and (f-j) are to scale. Images in panels (a'- e') and (f'-j') are not to scale. All wing discs are oriented with distal axis on left and proximal axis on right (see Figure 1b).

Figure 3

Figure 3. Programmed cell death in two stages of rudimentary wing disc development in larvae of wingless workers across the ant phylogeny. DAPI (blue) and TUNEL (red) signal in rudimentary wing discs of larvae in mid-terminal stage (left) and larval-prepupal transition stage (right). White arrowheads in panels (a, b, c, f, h, i, j) indicate region where TUNEL is expressed. Light black lines on the left mark phylogenetic relationships between the species represented. Subfamily labels are marked on the phylogenetic tree, species labels are marked on the left of the panels and clade labels are marked on the right-hand side of the panels. Images are not to scale. Phylogenetic relationship based on Borowiec et al (2019).

Figure 4

Figure 4. Programmed cell death is carried out via the apoptosis pathway. Expression of DAPI
(blue) and cleaved-caspase 3 (red) during mid (left) and late terminal larval (right) stages in *H.*
saltator (a-a'), and soldier larvae of *C. floridanus* (c-c') and *P. dentata* (d-d'). Expression of DAPI
(blue) and autophagy marker *atg8* or LC3A/B (red) in rudimentary wing discs of *H. saltator* (b-
b') and soldier larvae of *P. dentata* (e-e').

Figure 5. Programmed cell death in wing polyphenism across the ant phylogeny. Wing
polyphenism (blue circle) and eusociality (pink circle) evolved concomitantly in ants
(Formicidae, gray large box). Wing polyphenism and eusociality are found in some wasp
species/groups. Outgroups for Formicidae are for demonstration purposes only and many
branches in the Hymenoptera are excluded. Summary of PCD presence/absence found by this
study for poneroid species (purple box) and formicoid species (orange box) are listed on the
right. Characters indicate species with: presence of PCD (green circle), absence of PCD (red
circle), PCD untested (gray circles). Gray X indicates species where testing is not possible. PCD
regulation of wing polyphenism is predicted for all extant ants based on Ancestral State
Estimations (ASE; green circles in nodes). ASE values for all tested models yielded over 95%
confidence. PCD regulation of wing polyphenism at the origin of ants is proposed (dashed green
circle). Branch length do not indicate evolutionary time. Phylogenetic relationships within the
Formicidae are based on Boudinot, Khouri, et al. (2022); Moreau (2008); Ward et al. (2016).

446   Supplemental 1

450   **Supplemental 1.** Programmed cell death in additional ant species from the subfamily
451   Myrmicinae. For all panels, a rudimentary wing discs at the late terminal larval stage is shown
452   stained in DAPI (blue) and TUNEL (red).

454   Supplemental 2
455

456
457
Supplemental 2. Ancestral State Estimations for presence and absence of PCD. Estimates based
only on species where assays for PCD have been performed are shown using a phylogram (a)
and chronogram (a') modeling. Estimates based on species where assays for PCD have been
performed as well as the addition of basal outgroups *Martialis* and leptanilline are shown using
a phylogram (b) and chronogram (b') modeling. Estimates based on 74 ant taxa including those
without experimental data are shown using a phylogram (c) and chronogram (c') modeling.
Filled blue circles indicate estimates for presence of PCD; open circles indicate estimates for
absence of PCD, open circle indicate species where PCD has not been examined. All models
yielded over 95% confidence.
467

468 Supplemental 3
469

470
471
472 **Supplemental 3.** Positive and negative TUNEL controls shown for three ant species. DAPI (blue),
473 TUNEL (red).
474

REFERENCES

Abouheif, E., & Wray, G. A. (2002, Jul 12). Evolution of the gene network underlying wing polyphenism in ants. *Science, 297*(5579), 249-252. https://doi.org/10.1126/science.1071468

Ameisen, J. C. (2002, Apr). On the origin, evolution, and nature of programmed cell death: a timeline of four billion years. *Cell Death Differ, 9*(4), 367-393. https://doi.org/10.1038/sj.cdd.4400950

Behague, J., Fisher, B. L., Peronnet, R., Rajakumar, R., Abouheif, E., & Molet, M. (2018, Mar). Lack of interruption of the gene network underlying wing polyphenism in an early-branching ant genus. *J Exp Zool B Mol Dev Evol, 330*(2), 109-117. https://doi.org/10.1002/jez.b.22794

Bhatkar, A., & Whitcomb, W. H. (1970). Artificial diet for rearing various species of ants. *The Florida Entomologist, 53*(4), 229-232.

Borowiec, M. L., Schulz, A., Alpert, G. D., & Baňař, P. (2011). Discovery of the worker caste and descriptions of two new species of Anomalomyrma (Hymenoptera: Formicidae: Leptanillinae) with unique abdominal morphology. *Zootaxa, 2810*, 1-14.

Boudinot, B. E. (2015). Contributions to the knowledge of Formicidae (Hymenoptera, Aculeata): a new diagnosis of the family, the first global male-based key to subfamilies, and a treatment of early branching lineages. *European Journal of Taxonomy, 120*.

Boudinot, B. E., Khouri, Z., Richter, A., Griebenow, Z. H., van de Kamp, T., Perrichot, V., & Barden, P. (2022). Evolution and systematics of the Aculeata and kin (Hymenoptera), with emphasis on the ants (Formicoidea: †@@@idae fam. nov., Formicidae). *bioRxiv*. https://doi.org/https://doi.org/10.1101/2022.02.20.480183

Boudinot, B. E., Richter, A., Katzke, J., Chaul, J. C. M., Keller, R. A., Economo, E. P., Beutel, R. G., & Yamamoto, S. (2022). Evidence for the evolution of eusociality in stem ants and a systematic revision of †Gerontoformica (Hymenoptera: Formicidae). *zoological Journal of the Linnean Society, 195*(4), 1355-1389.

Burchill, A. T., & Moreau, C. S. (2016). Colony size evolution in ants: macroevolutionary trends. *Ins. Soc, 63*, 291-298. https://doi.org/https://doi.org/10.1007/s00040-016-0465-3

Buss, L. W. (2014). *The Evolution of Individuality*. Princeton University Press.

518 Chen, Z.-L., F.-M., S., & Zhou, S.-Y. (2017). First record of the monotypic genus Opamyrma
519   (Hymenoptera: Formicidae) from China. *Far Eastern Entomologist, 335*, 7-11.
520
521 Choi, H. M. T., Schwarzkopf, M., Fornace, M. E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha,
522   A., & Pierce, N. A. (2018, Jun 26). Third-generation in situ hybridization chain reaction:
523   multiplexed, quantitative, sensitive, versatile, robust. *Development, 145*(12).
524   https://doi.org/10.1242/dev.165753
525
526 Dewitz, H. (1878). Beiträge zur postembryonalen Gliedmassenbildung bei den Insecten.
527   *Zeitschrift für wissenschaftliche Zoologie, 30*, 78-105.
528
529 Durand, P. M., Barreto Filho, M. M., & Michod, R. E. (2019, Dec). Cell Death in Evolutionary
530   Transitions in Individuality. *Yale J Biol Med, 92*(4), 651-662.
531   https://www.ncbi.nlm.nih.gov/pubmed/31866780
532
533 Durand, P. M., Choudhury, R., Rashidi, A., & Michod, R. E. (2014, Feb). Programmed death in a
534   unicellular organism has species-specific fitness effects. *Biol Lett, 10*(2), 20131088.
535   https://doi.org/10.1098/rsbl.2013.1088
536
537 Durand, P. M., Rashidi, A., & Michod, R. E. (2011, Feb). How an organism dies affects the fitness
538   of its neighbors. *Am Nat, 177*(2), 224-232. https://doi.org/10.1086/657686
539
Galluzzi, L., Vitale, I., Aaronson, S. A., Abrams, J. M., Adam, D., Agostinis, P., Alnemri, E. S.,
Altucci, L., Amelio, I., Andrews, D. W., Annicchiarico-Petruzzelli, M., Antonov, A. V.,
Arama, E., Baehrecke, E. H., Barlev, N. A., Bazan, N. G., Bernassola, F., Bertrand, M. J. M.,
Bianchi, K., Blagosklonny, M. V., Blomgren, K., Borner, C., Boya, P., Brenner, C.,
Campanella, M., Candi, E., Carmona-Gutierrez, D., Cecconi, F., Chan, F. K., Chandel, N. S.,
Cheng, E. H., Chipuk, J. E., Cidlowski, J. A., Ciechanover, A., Cohen, G. M., Conrad, M.,
Cubillos-Ruiz, J. R., Czabotar, P. E., D'Angiolella, V., Dawson, T. M., Dawson, V. L., De
Laurenzi, V., De Maria, R., Debatin, K. M., DeBerardinis, R. J., Deshmukh, M., Di Daniele,
N., Di Virgilio, F., Dixit, V. M., Dixon, S. J., Duckett, C. S., Dynlacht, B. D., El-Deiry, W. S.,
Elrod, J. W., Fimia, G. M., Fulda, S., Garcia-Saez, A. J., Garg, A. D., Garrido, C.,
Gavathiotis, E., Golstein, P., Gottlieb, E., Green, D. R., Greene, L. A., Gronemeyer, H.,
Gross, A., Hajnoczky, G., Hardwick, J. M., Harris, I. S., Hengartner, M. O., Hetz, C., Ichijo,
H., Jaattela, M., Joseph, B., Jost, P. J., Juin, P. P., Kaiser, W. J., Karin, M., Kaufmann, T.,
Kepp, O., Kimchi, A., Kitsis, R. N., Klionsky, D. J., Knight, R. A., Kumar, S., Lee, S. W.,
Lemasters, J. J., Levine, B., Linkermann, A., Lipton, S. A., Lockshin, R. A., Lopez-Otin, C.,
Lowe, S. W., Luedde, T., Lugli, E., MacFarlane, M., Madeo, F., Malewicz, M., Malorni, W.,
Manic, G., Marine, J. C., Martin, S. J., Martinou, J. C., Medema, J. P., Mehlen, P., Meier,
P., Melino, S., Miao, E. A., Molkentin, J. D., Moll, U. M., Munoz-Pinedo, C., Nagata, S.,
Nunez, G., Oberst, A., Oren, M., Overholtzer, M., Pagano, M., Panaretakis, T., Pasparakis,
M., Penninger, J. M., Pereira, D. M., Pervaiz, S., Peter, M. E., Piacentini, M., Pinton, P.,
Prehn, J. H. M., Puthalakath, H., Rabinovich, G. A., Rehm, M., Rizzuto, R., Rodrigues, C.
M. P., Rubinsztein, D. C., Rudel, T., Ryan, K. M., Sayan, E., Scorrano, L., Shao, F., Shi, Y.,

Silke, J., Simon, H. U., Sistigu, A., Stockwell, B. R., Strasser, A., Szabadkai, G., Tait, S. W. G., Tang, D., Tavernarakis, N., Thorburn, A., Tsujimoto, Y., Turk, B., Vanden Berghe, T., Vandenabeele, P., Vander Heiden, M. G., Villunger, A., Virgin, H. W., Vousden, K. H., Vucic, D., Wagner, E. F., Walczak, H., Wallach, D., Wang, Y., Wells, J. A., Wood, W., Yuan, J., Zakeri, Z., Zhivotovsky, B., Zitvogel, L., Melino, G., & Kroemer, G. (2018, Mar). Molecular mechanisms of cell death: recommendations of the Nomenclature Committee on Cell Death 2018. *Cell Death Differ, 25*(3), 486-541. https://doi.org/10.1038/s41418-017-0012-4

Hall, B. K. (2003, Aug). Descent with modification: the unity underlying homology and homoplasy as seen through an analysis of development and evolution. *Biol Rev Camb Philos Soc, 78*(3), 409-433. https://doi.org/10.1017/s1464793102006097

Hanna, L., & Abouheif, E. (2021). The origin of wing polyphenism in ants: An eco-evo-devo perspective. *Curr Top Dev Biol, 141*, 279-336. https://doi.org/10.1016/bs.ctdb.2020.12.004

Hanna, L., & Abouheif, E. (2023, Aug). Deep conservation and co-option of programmed cell death facilitates evolution of alternative phenotypes at multiple biological levels. *Semin Cell Dev Biol, 145*, 28-41. https://doi.org/10.1016/j.semcdb.2022.05.024

Hartfelder, K., & Steinbruck, G. (1997). Germ cell cluster formation and cell death are alternatives in caste-specific differentiation of the larval honey bee ovary. *Invertebrate Reproduction and Development, 31*(1-3), 237-250.

Holldobler, B., & Wilson, E. O. (1990). *The Ants*. Harvard University Press.

Hsu, P. W., Hsu, F. C., Hsiao, Y., & Lin, C. C. (2017, May 15). Taxonomic notes on the genus Protanilla (Hymenoptera: Formicidae: Leptanillinae) from Taiwan. *Zootaxa, 4268*(1), 117-130. https://doi.org/10.11646/zootaxa.4268.1.7

Huettenbrenner, S., Maier, S., Leisser, C., Polgar, D., Strasser, S., Grusch, M., & Krupitza, G. (2003, Jun). The evolution of cell death programs as prerequisites of multicellularity. *Mutat Res, 543*(3), 235-249. https://doi.org/10.1016/s1383-5742(02)00110-2

[Record #678 is using a reference type undefined in this output style.]

Khila, A., & Abouheif, E. (2008, Nov 18). Reproductive constraint is a developmental mechanism that maintains social harmony in advanced ant societies. *Proc Natl Acad Sci U S A, 105*(46), 17884-17889. https://doi.org/10.1073/pnas.0807351105

Kyrylkova, K., Kyryachenko, S., Leid, M., & Kioussi, C. (2012). Detection of Apoptosis by TUNEL Assay. In C. kioussi (Ed.), *Odontogenesis. Methods in Molecular Biology (Methods and Protocols)* (Vol. 887). Humana Press.

Libby, E., & Ratcliff, W. C. (2014, Oct 24). Evolution. Ratcheting the evolution of multicellularity.
Science, 346(6208), 426-427. https://doi.org/10.1126/science.1262053

Maynard-Smith, J., & Szathmary, E. (1997). *The Major Transitions in Evolution*. Oxford
University Press.

Moreau, C. S. (2008, Jul). Unraveling the evolutionary history of the hyperdiverse ant genus
Pheidole (Hymenoptera: Formicidae). *Mol Phylogenet Evol, 48*(1), 224-239.
https://doi.org/10.1016/j.ympev.2008.02.020

Orellana, M. V., Pang, W. L., Durand, P. M., Whitehead, K., & Baliga, N. S. (2013). A role for
programmed cell death in the microbial loop. *PLoS One, 8*(5), e62595.
https://doi.org/10.1371/journal.pone.0062595

Peeters, C. (1991). The occurrence of sexual reproduction among ant workers. *Biological
Journal of the Linnean Society, 44*, 141-152.

Peeters, C. (1997). Morphologically 'primitive' ants: comparative review of social characters,
and the importance of queen-worker dimorphism. In J. C. Choe & B. J. Crespi (Eds.), *The
Evolution of Social Behavior in Insects and Arachnids*. Cambridge University Press.

Peeters, C. (2012). Convergent evolution of wingless reproductives across all subfamilies of
ants, and sporadic loss of winged queens (Hymenoptera: Formicidae). *Myrmecological
News, 16*, 75-91.

Penick, C. A., Prager, S. S., & Liebig, J. (2012, Dec). Juvenile hormone induces queen
development in late-stage larvae of the ant Harpegnathos saltator. *J Insect Physiol,
58*(12), 1643-1649. https://doi.org/10.1016/j.jinsphys.2012.10.004

Perrichot, V., Lacau, S., Neraudeau, D., & Nel, A. (2008, Feb). Fossil evidence for the early ant
evolution. *Naturwissenschaften, 95*(2), 85-90. https://doi.org/10.1007/s00114-007-
0301-8

Rabeling, C., Brown, J. M., & Verhaagh, M. (2008, Sep 30). Newly discovered sister lineage sheds
light on early ant evolution. *Proc Natl Acad Sci U S A, 105*(39), 14913-14917.
https://doi.org/10.1073/pnas.0806187105

Rajakumar, R., Koch, S., Couture, M., Fave, M. J., Lillico-Ouachour, A., Chen, T., De Blasis, G.,
Rajakumar, A., Ouellette, D., & Abouheif, E. (2018, Oct). Social regulation of a
rudimentary organ generates complex worker-caste systems in ants. *Nature, 562*(7728),
574-577. https://doi.org/10.1038/s41586-018-0613-1

Ratcliff, W. C., Denison, R. F., Borrello, M., & Travisano, M. (2012, Jan 31). Experimental
evolution of multicellularity. *Proc Natl Acad Sci U S A, 109*(5), 1595-1600.
https://doi.org/10.1073/pnas.1115323109
652

653   Romiguier, J., Borowiec, M. L., Weyna, A., Helleu, Q., Loire, E., La Mendola, C., Rabeling, C.,
654   Fisher, B. L., Ward, P. S., & Keller, L. (2022, Jul 11). Ant phylogenomics reveals a natural
655   selection hotspot preceding the origin of complex eusociality. *Curr Biol, 32*(13), 2942-
656   2947 e2944. https://doi.org/10.1016/j.cub.2022.05.001
657

658   Sameshima, S. Y., Miura, T., & Matsumoto, T. (2004, Sep-Oct). Wing disc development during
659   caste differentiation in the ant Pheidole megacephala (Hymenoptera: Formicidae). *Evol
660   Dev, 6*(5), 336-341. https://doi.org/10.1111/j.1525-142X.2004.04041.x
661

662   Shbailat, S. J., & Abouheif, E. (2013, Mar). The wing-patterning network in the wingless castes of
663   Myrmicine and Formicine ant species is a mix of evolutionarily labile and non-labile
664   genes. *J Exp Zool B Mol Dev Evol, 320*(2), 74-83. https://doi.org/10.1002/jez.b.22482
665

666   Shbailat, S. J., Khila, A., & Abouheif, E. (2010, Nov-Dec). Correlations between spatiotemporal
667   changes in gene expression and apoptosis underlie wing polyphenism in the ant
668   Pheidole morrisi. *Evol Dev, 12*(6), 580-591. https://doi.org/10.1111/j.1525-
669   142X.2010.00443.x
670

671   Ward, P. S., Blaimer, B. B., & Fisher, B. L. (2016, Feb 2). A revised phylogenetic classification of
672   the ant subfamily Formicinae (Hymenoptera: Formicidae), with resurrection of the
673   genera Colobopsis and Dinomyrmex. *Zootaxa, 4072*(3), 343-357.
674   https://doi.org/10.11646/zootaxa.4072.3.4
675

676   West, S. A., Fisher, R. M., Gardner, A., & Kiers, E. T. (2015, Aug 18). Major evolutionary
677   transitions in individuality. *Proc Natl Acad Sci U S A, 112*(33), 10112-10119.
678   https://doi.org/10.1073/pnas.1421402112
679

680   Wheeler, D. E. (1986). Developmental and physiological determinants of caste in social
681   hymenoptera: evolutionary implications. *The American Naturalist, 128*(1), 13-34.
682

683   Wheeler, D. E., & Nijhout, H. F. (1981). Imaginal wing discs in larvae of the soldier caste of
684   Pheidole bicarinata vinelandica Forel (Hymenoptera : Formicidae). *International Journal
685   of Insect Morphology and Embryology, 10*(2), 131-139.
686   https://doi.org/https://doi.org/10.1016/S0020-7322(81)80018-9
687

688   Wilson, E. O. (1953). The origin and evolution of polymorphism in ants. *The Quarterly Review of
689   Biology*, 136-156.
690

691   Wilson, E. O., & Holldobler, B. (2005, May 24). The rise of the ants: a phylogenetic and
692   ecological explanation. *Proc Natl Acad Sci U S A, 102*(21), 7411-7414.
693   https://doi.org/10.1073/pnas.0502264102
694

695   Wilson, J. D., Mongiardino Koch, N., & Ramírez, M. J. (2022). Chronogram or phylogram for
696   ancestral state estimation? Model-fit statistics indicate the branch lengths underlying a
697   binary character's evolution. *Methods in Ecology and Evolution, 13*(8), 1679-1689.
698

699   Xu, D., Woodfield, S. E., Lee, T. V., Fan, Y., Antonio, C., & Bergmann, A. (2009, Jan-Mar). Genetic
700   control of programmed cell death (apoptosis) in Drosophila. *Fly (Austin), 3*(1), 78-90.
701   https://doi.org/10.4161/fly.3.1.7800
702

703   Yan, G., Elbadawi, M., & Efferth, T. (2020). Multiple cell death modalities and their key features
704   (Review). *World Academy of Sciences Journal, 2*, 39-48.
705
706
707