Feed-forward stimulation of CAMK2 by the oncogenic pseudokinase PEAK1 generates

a therapeutically 'actionable' signalling axis in triple negative breast cancer

Xue Yang[1,2], Xiuquan Ma[1,2], Tianyue Zhao[1,2], David R Croucher[3,4], Elizabeth V. Nguyen[1,2],

Kimberley C. Clark[1,2], Changyuan Hu[1,2], Sharissa L Latham[3,4], Charles Bayly-Jones[1,2], Viet

Chi Bao Nguyen[5], Srikanth Budnar[1,2], Sung-Young Shin[1,2], Lan K. Nguyen[1,2], Thomas R.

Cotton[1,2], Anderly C. Chüeh[1,2], Terry C C Lim Kam Sian[1,2], Margaret M. Stratton[5], Andrew

M. Ellisdon[1,2] and Roger J Daly[1,2,#].

[1]Cancer Program, Monash Biomedicine Discovery Institute, Clayton, VIC 3800, Australia.

[2]Department of Biochemistry and Molecular Biology, Monash University, Melbourne, VIC

3800, Australia.

[3]Garvan Institute of Medical Research, Darlinghurst, NSW 2010, Australia.

[4]St Vincent's Clinical School, Faculty of Medicine, UNSW Sydney, Darlinghurst, NSW 2010,

Australia.

[5]Department of Biochemistry and Molecular Biology, University of Massachusetts, Amherst,

MA 01003, USA

[#]Correspondence: roger.daly@monash.edu

Summary

The PEAK family of pseudokinases, comprising PEAK1-3, are signalling scaffolds that play

oncogenic roles in several poor prognosis human cancers, including triple negative breast

cancer (TNBC). However, therapeutic targeting of pseudokinases is challenging due to their

lack of catalytic activity. To address this, we screened for PEAK1 effectors by affinity

purification and mass spectrometry, identifying calcium/calmodulin-dependent protein kinase

2 (CAMK2)D and CAMK2G. PEAK1 promoted CAMK2D/G activation in TNBC cells via a

novel feed-forward mechanism involving PEAK1/PLCγ1/Ca^{2+} signalling and direct binding

via a consensus CAMK2 interaction motif in the PEAK1 N-terminus. In turn, CAMK2

phosphorylated PEAK1 to enhance association with PEAK2, which is critical for PEAK1

oncogenic signalling. To achieve pharmacologic targeting of PEAK1/CAMK2, we repurposed

RA306, a second generation CAMK2 inhibitor under pre-clinical development for treatment

of cardiovascular disease. RA306 demonstrated on-target activity against CAMK2 in TNBC

cells and inhibited PEAK1-enhanced migration and invasion *in vitro*. Moreover, RA306

significantly attenuated TNBC xenograft growth and blocked metastasis in a manner mirrored

by CRISPR-mediated PEAK1 ablation. Overall, these studies establish PEAK1 as a critical

cell signalling nexus, identify a novel mechanism for regulation of Ca^{2+} signalling and its

integration with tyrosine kinase signals, and identify CAMK2 as a therapeutically 'actionable'

target downstream of PEAK1.

Introduction

Pseudokinases play critical roles in normal cellular and developmental processes as well as in human diseases such as cancer, despite lacking the kinase activity of their *bona fide* kinase counterparts. This reflects the ability of pseudokinases to function as allosteric regulators and/or scaffolds that assemble protein signalling complexes[1, 2]. Due to their pathological functions, pseudokinases are attracting increasing attention as therapeutic targets[3].

The PEAK family of pseudokinases, comprising PEAK1 (previously known as SgK269), PEAK2 (Pragmin and SgK223) and PEAK3 (C19orf35) share a conserved molecular architecture and sequence homology, and function as scaffolds to mediate and regulate signalling downstream of particular growth factor receptors and integrins[4, 5]. PEAK1 is a critical regulator of EGFR signalling, mediating a qualitative shift in signal output over time from mitogenic/survival signalling to that promoting cell migration/invasion[16]. In addition, PEAK1 is strongly implicated in progression of a variety of cancers. For example, in breast cancer, PEAK1 promotes cell proliferation, epithelial-mesenchyme transition (EMT), aberrant acinar morphology in 3D culture and xenograft growth and metastasis[6, 7, 8], in non small cell lung cancer it enhances migration, invasion, EMT and experimental metastasis[9], while in pancreatic cancer models it promotes resistance to specific therapies[10]. Of particular interest, high PEAK1 expression occurs in triple negative breast cancer (TNBC) and pancreatic cancer[6, 10], two cancers associated with poor prognosis and limited targeted therapeutic options. PEAK2 and PEAK3 also enhance cell migration and invasion and are overexpressed in specific malignancies[4, 11, 12, 13, 14, 15].

All PEAK family members exhibit a C-terminal pseudokinase domain flanked by a split helical

dimerization (SHED) domain, and N-terminal extensions of contrasting length that harbour

recruitment sites for specific SH2, SH3 and PTB domain-containing effectors. These effectors

include Grb2 (bound by PEAK1 and PEAK3), Shc1 (PEAK1), C-terminal Src kinase (Csk)

(PEAK2) and ASAP1, Cbl and PYK2 (PEAK3)[4, 5]. An important characteristic of the PEAK

family is their ability to form homotypic and heterotypic complexes[4, 16]. Dimerization of the

PEAKs is mediated by the α-helical SHED domain that forms a highly conserved dimer

interface driven by hydrophobic interactions[17, 18], while higher order oligomers can also form

via pseudokinase domain interfaces[18]. Dimerization is critical to the signalling potential of

PEAK scaffolds[4, 12, 18]. Furthermore, since the repertoire of known interactors differs across

this family of scaffolds, heterotypic versus homotypic association acts to diversify signal

output[4, 16]. For example, PEAK1 requires PEAK2 to efficiently activate Stat3 and promote

migration in MCF-10A cells[16, 19], while signalling by PEAK3 is modulated both quantitatively

and temporally by the presence of PEAK1 and PEAK2[4].

In this study, we addressed how homo- and heterotypic association of PEAK1 and PEAK2

regulates signal output by defining the interactomes of the corresponding dimeric complexes.

Amongst the identified binding interactions, the association of PEAK1 with

calcium/calmodulin-dependent protein kinase 2 (CAMK2)D and CAMK2G was of particular

interest, given the known roles of specific CAMK2 isoforms in many human cancers, including

breast cancer, and the therapeutic 'actionability' of these serine/threonine kinases[20, 21].

Importantly, we demonstrate potent, feed-forward activation of CAMK2 by PEAK1,

highlighting PEAK1 as a critical regulator and integrator of tyrosine kinase and Ca^{2+} signals.

In addition, we identify CAMK2D and CAMK2G as novel downstream effectors of PEAK1 in

TNBC and 'repurposing' of the second generation CAMK2 inhibitor RA306 as a potential

therapeutic strategy against oncogenic PEAK1 signalling in poor prognosis human cancers

such as TNBC.

Results

Characterization of dimer-specific interactomes of PEAK1 and PEAK2 by BiCAP

To characterize the interactomes of PEAK1 and PEAK2 homodimers, and the PEAK1/PEAK2

heterodimer, we applied bimolecular complementation affinity purification coupled with

tandem mass spectrometry (BiCAP-MS/MS) (Supp. Fig. 1A)[22]. Specific dimeric complexes

were affinity purified and associated proteins identified by LC-MS/MS in data-independent

acquisition mode (Methods, Fig. 1A-C, Supp. Table 1). Strong correlation between triplicate

biological replicates was observed (mean sample correlation scores: Vector = 0.78; PEAK1

homodimer = 0.85; PEAK2 homodimer = 0.94; PEAK1/2 heterodimer = 0.94) (Supp. Fig.

1B). Grb2, a well-characterized PEAK1 binding partner[6, 16] bound the PEAK1 homodimer and

PEAK1/PEAK2 heterodimer, but not the PEAK2 homodimer, consistent with our previous

results that PEAK1 bridges Grb2 to PEAK2[16]. Two protein phosphatase 1 (PP1) family

members, PPP1CA and PPP1CC were previously identified as PEAK1 interactors in a

traditional immunoaffinity based-MS/MS screen[19]. However, their selectivity towards PEAK1

versus PEAK2 was unclear. Here, like Grb2, they were identified as interactors of the PEAK1

homodimer and PEAK1/PEAK2 heterodimer, with another PP1 family member, PPP1CB,

associating with the PEAK1 homodimer. Of interest given a recent paper reporting a key

regulatory role for 14-3-3 proteins in regulating PEAK3 effector recruitment and biological

activity[23], multiple 14-3-3 isoforms bound both homodimers and the heterodimer (Supp. Table

1). Further novel interactors identified included the RAC1 guanine nucleotide exchange factor

(GEF) FARP1 and the MAPK pathway antagonist SPRY4, which were selective for the

PEAK2 homodimer. In addition, Ca^{2+}/calmodulin-dependent protein kinase 2 (CAMK2)D and

G associated with the PEAK1 and PEAK2 homodimers and the PEAK1/PEAK2 heterodimer,

while CAMK2B was recruited by the PEAK2 homodimer and PEAK1/PEAK2 heterodimer

(Figs. 1A-C). Bioinformatic analysis of the interactomes for each of the PEAK complexes in

order to identify enriched pathways/processes identified calcium-related pathways that

included CAMK2 enzymes as being enriched (specifically 'postsynaptic signalling' for

PEAK1 and PEAK2 homotypic complexes and both 'myometrial' and 'cardiac signalling' for

all 3 complexes) (Supp. Table 2).

Overall, these data provided novel and important insights into the binding selectivity of

PEAK1/2 complexes and identified specific CAMK2 family members as novel interactors of

these pseudokinase scaffolds.

Characterization of CAMK2 interaction with PEAK1 and PEAK2

To confirm the association of specific CAMK2 family members with PEAK1 and PEAK2, we

focused on CAMK2D and CAMK2G, since these kinases associated with all of the homo- and

heterotypic PEAK complexes analysed by BiCAP-MS/MS. Interaction of these CAMK2s with

endogenous PEAK1 and PEAK2 was interrogated by proximity ligation assays in MDA-MB-

231 breast cancer cells (Fig. 1D-F). Using this technique, the interactions between PEAK1 and

CAMK2D or G could be detected and were significantly decreased when PEAK1 was knocked

down using siRNA (Fig. 1D-E). Similar results were observed for the interactions between

PEAK2 and CAMK2D or G (Fig. 1D,F). These data demonstrate that CAMK2D and G interact

with both PEAK1 and PEAK2 *in vivo,* at endogenous expression levels.

Regulation of CAMK2 activity by PEAK1

The known overexpression of PEAK1 in breast cancer[6, 7] led us to interrogate the functional

consequences of PEAK1/CAMK2 interaction. Upon an increase in intracellular Ca^{2+}, binding

of Ca^{2+}/calmodulin (CaM) to the CAMK2 regulatory segment leads to release of the regulatory

segment from the kinase domain, kinase activation, and phosphorylation of T286 (T287 in

CAMK2D and G). In the presence of T286 phosphorylation, the CAMK2 regulatory segment

can no longer mediate autoinhibition, and the activity of CAMK2 is Ca^{2+}/CaM autonomous [24].

Knockdown of PEAK1 in MDA-MB-231 TNBC cells did not affect the expression levels of

CAMK2D and G but significantly reduced their activation, as determined by phosphorylation

at T287 (Fig. 2A-B). In addition, overexpression of PEAK1 in these cells significantly

enhanced activation of these two CAMK2 isoforms (Fig. 2C-D). Positive regulation of

CAMK2D/G activity by PEAK1 was also observed in MDA-MB-468 TNBC cells (Supp. Fig.

2). Treatment of MDA-MB-231 cells with BAPTA/AM, a membrane permeable Ca^{2+} chelator

which blocks activation of CaM by Ca^{2+} [25] significantly reduced basal CAMK2 T287

phosphorylation and completely blocked the effect of PEAK1 on CAMK2 activation,

indicating that PEAK1-mediated CAMK2 activation is Ca^{2+}-dependent (Fig. 2E-F).

In light of this result, we determined whether PEAK1 can modulate Ca^{2+} levels and thereby

indirectly regulate CAMK2 activation. Application of Fluo-4 imaging revealed that PEAK1

overexpression significantly enhanced intracellular Ca^{2+} levels at steady state (Fig. 2G-H).

With regard to the mechanism underpinning PEAK1-regulated Ca^{2+} signalling, the known roles

of the PEAKs as regulators and/or substrates of specific tyrosine kinases[4, 5] led us to determine

whether PEAK1 overexpression impacts tyrosine phosphorylation and hence activation of

PLCγ1, leading to generation of IP_3 that triggers Ca^{2+} release from intracellular stores[26]. Indeed,

PEAK1 knockdown decreased tyrosine phosphorylation of PLCγ1 Y783 (Fig. 2I), while

PEAK1 overexpression enhanced PLCγ tyrosine phosphorylation on this site (Fig. 2J).

However, given that PEAK1 is a pseudokinase, PLCγ1 phosphorylation must be mediated by

a *bona fide* tyrosine kinase acting downstream of PEAK1. Since PEAK1 is a member of a

prominent Src family kinase (SFK) signalling network in TNBC[27] and PLCγ1 is a known

substrate of SFKs[28], we hypothesized that SFKs may mediate this role. Supporting this

hypothesis, treatment of MDA-MB-231 TNBC cells with the highly-selective SFK inhibitor

eCF506[29]   blocked PEAK1-enhanced PLCγ1 tyrosine phosphorylation (Supp. Fig. 3).

However, interrogation of the BiCAP datasets did not reveal association of specific SFKs or

PLCγ1 with any of the PEAK complexes, strongly suggesting that the regulatory mechanism

does not involve direct binding.

Structural requirements for PEAK1/2 interaction with CAMK2

To determine which regions of PEAK1 and PEAK2 mediate interaction with CAMK2D and

G, we utilized previously described deletion mutants of PEAK1 and PEAK2[16] that delete the

N-terminal region (ΔN-term), N1 α-helix (ΔN1) and a combination of the pseudokinase

domain and helices J-M (ΔCterm), as well as a new mutant comprising just the extreme N-

terminal region (1-324 amino acids) (N-term) (Fig. 3A). Co-immunoprecipitation of Flag-

tagged PEAK1 with endogenous CAMK2D could be demonstrated for WT, ΔN1 and ΔCterm

PEAK1 but not for the ΔNterm PEAK1 mutant (Fig. 3B). However, the Nterm region alone

did not associate with CAMK2D, indicating that it is required but not sufficient for interaction

with these CAMK2s (Fig. 3B). In addition, since the N1 and Cterm regions are required for

PEAK1 to undergo homotypic association and also heterotypic interaction with PEAK2[16],

these data also indicated that dimerization is not required for PEAK1 to associate with these

CAMK2s. Next, we generated a series of deletion mutants to more narrowly define the

CAMK2 binding region on PEAK1 (Fig. 3C). Association with CAMK2D was lost with the

Δ261-324 mutant (Fig. 3D). Furthermore, deletion of smaller regions spanning amino acids

261-300 or 301-324 also abolished CAMK2D association (Fig. 3D). Deletion of amino acids

301-324 also markedly reduced CAMK2G association (Fig. 3E).

Although PEAK1 and PEAK2 share similar overall domain architecture, the N-terminal

extensions, including the region of PEAK1 required for CAMK2 association (amino acids 261-

324), are poorly conserved[16], indicating that the CAMK2 binding mechanism for PEAK2 is

likely to differ. Consistent with this hypothesis, while CAMK2D association with PEAK2 was

significantly reduced with the PEAK2 ΔNterm mutant, the greatest effect was observed with

the ΔN1 and ΔCterm mutants, which markedly and almost completely abolished association,

respectively (Supp. Fig. 4A-C). Since the N1 and Cterm regions are required for PEAK2 to

undergo homotypic and heterotypic association[16], these data suggest that PEAK2 interaction

with these CAMK2s requires PEAK2 dimerization. Supporting this model, three point

mutations in PEAK2 (Supp. Fig. 4A) that abrogate dimerization[18] all led to diminished

association with CAMK2D (Supp. Fig. 4D). This contrasts with PEAK1, where the

dimerization-impaired ΔN1 and ΔCterm mutants retained association with the CAMK2s. In

addition, PEAK1 knockdown did not disrupt association of PEAK2 with CAMK2D (Supp. Fig.

4E), indicating that PEAK2 interaction with CAMK2D is not mediated via PEAK1, consistent

with the BiCAP data (Fig. 1B). Taken together, these data highlight contrasting structural

requirements for PEAK1 and PEAK2 association with CAMK2, with a greater contribution

from a discrete N-terminal region in the case of PEAK1, and a requirement for dimerization in

the case of PEAK2.

PEAK1 directly binds CAMK2 via a conserved motif to promote CAMK2 activation and PEAK1 phosphorylation

Since the PEAK1 Δ301-324 mutant was deficient in CAMK2 association (Fig. 3D-E), we determined the effect of this mutant on CAMK2 activation. Interestingly, this mutant was significantly impaired in its ability to activate CAMK2D/G as determined by phosphorylation at T287 (Fig 3F-G), indicating that PEAK1-mediated CAMK2 activation requires association of the two proteins in addition to a Ca^{2+} signal (Fig. 2E-F).

A recent report characterized how CAMK2 interacts with substrate binding partners via a conserved sequence motif in the binding partner, R/K-X-X-ϕ-X-R/K-X-X-S/T-ϕ (where ϕ represents a hydrophobic residue) that binds across a single continuous site on the CAMK2 kinase domain[30]. It also revealed that binding of the conserved CAMK2 interaction motif (CIM) competes with autoinhibition of CAMK2 by the regulatory segment of the kinase domain, explaining the ability of certain substrates, such as GluN2B, to lock CAMK2 in an active conformation[30]. Of note, the Rac GDP/GTP exchange factor TIAM1 exhibits a pseudosubstrate version of the conserved motif (with A rather than S/T), but is still able to bind and promote phosphorylation of itself on other sites[31]. To address whether PEAK1 may interact with CAMK2 via this mechanism, we searched the PEAK1 N-terminal region for the conserved motif. We first constructed a multiple sequence alignment across diverse species and subsequently subjected this to MEME analysis to identify conserved motifs. This analysis revealed that the PEAK1 amino acids 297-307 exhibit close similarity to the canonical binding motif but with an alanine at position 0 (amino acid 306), as observed with TIAM1 (Fig. 4A). Furthermore, the N-terminal region and the binding motif are highly conserved across species (Supp. Fig. 5).

The presence of a CIM between amino acids 297-307 is consistent with disruption of binding

by deletion of amino acids 261-300 and 301-324 (Fig. 3D-E) and the failure of the Δ301-324

mutant to activate CAMK2 (Fig 3F-G). To determine whether a synthetic peptide

corresponding to the PEAK1 CIM bound CAMK2, we applied isothermal calorimetry. These

data revealed a high affinity interaction with a mean dissociation constant (K_D) of 635 nM (Fig.

4B), comparable to that of the TIAM1 interacting region (1.1 μM)[30]. Modelling of the

PEAK1/CAMK2 interaction using AlphaFold predicts (pLDDT 90.7, pTM 0.899) that the

PEAK1 CIM interacts across the same surface of the CAMK2 kinase domain as other known

interactors[30], with key salt bridges formed between PEAK1 R303 (-3 position) and R297 (-9)

with glutamate residues in CAMK2 (Fig. 4C-F, Supp. Fig. 6). In addition, L301 is

accommodated in a hydrophobic pocket of CAMK2 (Fig. 4E). We also note the predicted

formation of salt bridges and electrostatic interactions by PEAK1 D308, D310, D311 and D314

across the surface of CAMK2 (Fig. 4F). Of note, while this modelling was undertaken with

CAMK2A, the binding site is conserved across all CAMK2 isoforms, as highlighted by

AlphaFold2 modelling and multiple sequence alignments of CAMK2A, B, D and G (Supp.

Fig. 6).

To test the importance of key residues within the PEAK1 CIM for CAMK2 activation, we

generated PEAK1 mutants with alanine substitutions at the R303 (-3), L301 (-5) and R297 (-

9) positions (termed AAA) and glutamate substitutions at R303 and R297 (EE).

Overexpression of these two mutants significantly enhanced intracellular Ca^{2+} levels (Fig. 5A-

B) and PLCγ1 tyrosine phosphorylation (Fig. 5C-D). However, the mutants lost the ability to

activate CAMK2 as determined by phosphorylation at T287 (Fig. 5E-G). These data indicate

that while the ability of PEAK1 to activate CAMK2 is calcium-dependent (Fig. 2E-F), PEAK1-

enhanced Ca^{2+} is not sufficient to increase CAMK2 T287 phosphorylation, and direct

association with CAMK2 is also required. The data also support the validity of the Alphafold

modelling.

CAMK2 binding to TIAM1 leads to sustained phosphorylation of TIAM1[31]. To determine

whether PEAK1 is a substrate for CAMK2, we utilized RA306, a 'next generation' CAMK2

inhibitor developed by Sanofi for chronic cardiovascular indications that is selective for

CAMK2D and G over other isoforms[32]. RA306 exhibited 'on-target' activity against CAMK2

in MDA-MB-231 cells (Supp. Fig. 7A) and significantly inhibited PEAK1 serine/threonine

phosphorylation (Fig. 6A-B). In addition, recombinant CAMK2 demonstrated *in vitro* kinase

activity towards PEAK1 that was detectable using a pAkt substrate antibody that also

recognizes the minimal CAMK2 consensus motif for phosphorylation (RXXS/T) or a pSer/Thr

antibody (Fig. 6C). Interestingly, PEAK1 phosphorylation was markedly reduced when the

PEAK1 AAA and EE CIM mutants were used as substrate, indicating that efficient

phosphorylation of PEAK1 by CAMK2 requires interaction of the two proteins via the PEAK1

CIM (Fig. 6C).

CAMK2 phosphorylates PEAK1 to regulate the PEAK1 interactome

In order to characterize the impact of CAMK2-mediated phosphorylation on PEAK1

scaffolding potential, we compared the interactomes of wild-type (WT) PEAK1 and the AAA

and EE CIM mutants *in vivo*. We utilized this approach since it more selectively addresses the

impact of CAMK2-mediated PEAK1 phosphorylation, while RA306 treatment would exert a

more general inhibition of CAMK2-mediated cellular phosphorylation. Binding of the adaptors

Crk-L and Grb2, which associate with PEAK1 via SH3 and SH2 domain-mediated interactions,

respectively[23,9], was unaffected by the CIM mutations (Supp. Fig. 7B). In contrast, Western

blotting of PEAK1 IPs revealed significantly reduced association of 14-3-3 proteins with the

two mutants (Supp. Fig. 7C-D). Since 14-3-3 binding to client proteins is classically

phosphorylation-dependent, this indicated that impaired CAMK2 activation by the CIM

mutants leads to decreased phosphorylation of PEAK1 on 14-3-3 binding sites. Next, we

assayed heterotypic association of PEAK1 with PEAK2, a critical determinant of signal output,

with PEAK2 binding being required for PEAK1 to promote cell migration and oncogenic

signalling[16]. Importantly, whereas heterotypic association of wildtype PEAK1 with PEAK2 in

MDA-MB-231 TNBC cells could be readily detected, this was markedly reduced for the CIM

mutants (Figure 6D). Overall, these data demonstrate that CAMK2-mediated PEAK1

phosphorylation is a critical regulator of PEAK1-mediated protein-protein interactions and

signalling.

PEAK1/CAMK2 signalling is a therapeutically 'actionable' target in TNBC

The finding that CAMK2 and PEAK1 reciprocally regulate led us to interrogate how

expression of these proteins is related to breast cancer patient survival. Interestingly, combined

high expression of PEAK1 and CAMK2D was significantly associated with poor overall

survival, while a non-significant trend was detected for the PEAK1/CAMK2G combination

(Supp. Fig. 8A-C). Of note, this analysis emphasized the importance of high expression of both

PEAK1 and CAMK2D (Supp. Fig. 8B), consistent with a co-activating PEAK1/CAMK2D

'cassette' promoting breast cancer progression. To gain further insights into the combined role

of PEAK1 and CAMK2/G in breast cancer, we interrogated their association with distant

metastasis-free survival (DMFS). However, this was only possible with a meta-cohort derived

from certain publicly-available datasets (see Methods) and for the PEAK1/CAMK2G

combination. Nevertheless, this revealed a significant association between high

PEAK1/CAMK2G and reduced DMFS (Supp. Fig. 8D).

Next, we characterized the effects of PEAK1/CAMK2 signalling on biological endpoints in

TNBC cells. Previously we reported that CRISPR-mediated PEAK1 KO in MDA-MB-231

cells significantly reduced anchorage-dependent proliferation as well as tumoursphere

formation in low attachment plates[8]. Interestingly, treatment with RA306 resulted in a dose-

dependent reduction in viability of these cells under anchorage-dependent and -independent

conditions (high and low attachment plates, respectively) (Supp. Fig. 9A-B). These effects

appeared to be mediated via decreased proliferation, as PEAK1 KO or RA306 did not induce

apoptosis (Supp. Fig. 9C-D). Taken together, these data indicate that both PEAK1 and CAMK2

contribute to proliferative signalling in TNBC.

Next, we probed the role of PEAK1/CAMK2 in cell migration/invasion. Knockdown of either

CAMK2D or G blocked the effect of PEAK1 overexpression on cell migration and invasion

(Fig. 7A-B, Supp. Fig. 10A) and RA306 treatment blocked PEAK1-promoted migration in

both MDA-MB-231 and -468 cells (Fig. 7C, Supp. Fig. 10B-C) and PEAK1-promoted invasion

in MDA-MB-231 cells (Supp. Fig. 10D). Furthermore, the Δ301-324 PEAK1 mutant, which

cannot bind or activate CAMK2 (Fig. 3D-G), was unable to enhance TNBC cell migration and

invasion (Fig. 7D, Supp. Fig. 10E). Overall, these data indicate a requirement for CAMK2

expression and/or activation, as well as PEAK1/CAMK2 binding, in regulation of cell

migration and invasion by PEAK1.

We then evaluated RA306 in an *in vivo* model of TNBC tumour growth and metastasis.

Previously, we demonstrated that PEAK1 gene knock-out significantly inhibited growth and

lung metastasis of orthotopic MDA-MB-231 TNBC xenografts[8]. To test RA306, we utilized

the highly metastatic variant of this cell line, MDA-MB-231-HM[33], so that we could readily

assay metastatic spread from the primary tumour. In addition, we generated PEAK1-deficient

MDA-MB-231-HM cells via CRISPR editing (Supp. Fig. 11A) to compare the effect of RA306

with PEAK1 knock-out and also the combined effect of these two manipulations. On target

activity of RA306 could be detected in the xenografts and PEAK1 ablation also reduced

CAMK2 activation (Supp. Fig. 11B-C). RA306 treatment significantly impaired primary

tumour growth (Fig. 7E). Importantly, the degree of growth inhibition was similar to that

observed upon PEAK1 gene knock-out and combining RA306 with gene knock-out did not

enhance this further, strongly suggesting that RA306 treatment and PEAK1 knock-out are

targeting the same oncogenic signalling axis (Fig. 7E). Furthermore, we also allowed a separate

cohort of primary tumours to grow to the same mean size (Supp Fig 11D), resected them and

then monitored metastasis. Strikingly, RA306 blocked metastasis to a similar extent to PEAK1

knock-out (Fig 7F). Since PEAK1 KO or RA306 treatment impacts cell proliferation, migration

and invasion but not apoptosis[8] (Fig. 7, Supp. Fig. 9-10), the effect on primary tumour growth

is likely mediated via reduced proliferation while the reduced metastatic growth must be

independent of primary tumour size and reflect a decreased ability to metastasize to the lung

and/or reduced proliferation at this secondary site. Overall, these data identify RA306 as a

potential therapeutic strategy against the PEAK1/CAMK2 axis in TNBC and potentially other

human cancers.

Discussion

The pseudokinase scaffold PEAK1 has emerged as a critical regulator of intracellular

signalling, transducing and integrating pleiotropic signals downstream of a variety of growth

factor receptors and integrins. For example, PEAK1 promotes Ras activation via Grb2

binding[6], and couples to paxillin regulation at focal adhesions via Crk[8]. In addition, it can

indirectly impact Csk and Stat3 via association with PEAK2[16] and ASAP1, PYK2 and Cbl via

PEAK3[4]. Our discovery that PEAK1 can also promote Ca^{2+} signals via PLCγ1 and moreover,

binds and sustains activation of the Ca^{2+} effector CAMK2 greatly expands the signalling

repertoire of this scaffold and cements its role as a fundamental cell signalling nexus. It also

identifies PEAK1 as the first member of the kinase superfamily to sustain activation of

CAMK2 via direct binding and the first oncoprotein to target CAMK2 via this mechanism.

This expansion of the known signalling functions of PEAK1 calls for a re-evaluation of its role

in physiological processes linked to CAMK2 signalling including learning and memory,

cardiac function and osteogenic differentiation[20]. As discussed later, it also highlights a

potential strategy for therapeutically targeting the oncogenic activity of PEAK1.

Our findings lead to a model whereby PEAK1 enhances and prolongs CAMK2 activation and

signalling by two mechanisms that together provide a 'feed-forward' regulatory process for

CAMK2 (Figure 8). First, PEAK1 overexpression leads to enhanced and more sustained Ca^{2+}

signals in cells, reflecting increased tyrosine phosphorylation and hence activation of PLCγ1

which is driven in a SFK-dependent manner. While the role of SFKs in this pathway is

consistent with PEAK1 functioning within a SFK signalling network in TNBC[27], the detailed

mechanism currently remains unclear, and based on our BiCAP data, does not appear to involve

direct binding of PEAK1/2 to SFKs or PLCγ1. In this regard, it differs from the direct binding

and regulation of the tyrosine kinases Csk and PYK2 by PEAK2 and PEAK3, respectively[12,

34]. In terms of the SFKs responsible, TNBCs express several SFKs including Src, Yes, Fyn and

Lyn, but expression of the latter is particularly high, and since Lyn also regulates PEAK1

tyrosine phosphorylation, it represents an interesting candidate for further investigation[6, 27, 35].

The enhanced Ca^{2+} signal generated by this PEAK1/SFK/PLCγ1/Ca^{2+} pathway leads to a

greater pool of activated CAMK2. Second, PEAK1 binds directly to the Ca^{2+}-activated

CAMK2 via the conserved CIM and this has two effects – first, it leads to sustained CAMK2

activation, as determined by T287 phosphorylation, and second, it promotes CAMK2-mediated

PEAK1 phosphorylation.

Promotion of CAMK2 T287 phosphorylation and PEAK1 phosphorylation by binding of the

PEAK1 CIM to CAMK2 can be explained by a model that incorporates two key characteristics

of this enzyme/interactor system - first, the known ability of CAMK2 to form a large

holoenzyme comprising 12-14 monomers [24] and second, PEAK1 CIM binding 'locking'

CAMK2 in an active state with the αD helix rotated outward[30] (Figure 8). In this model,

repeated cycles of association/disassociation of one PEAK1 CIM to different CAMK2 subunits

within the complex leads to sustained activation of a subset of subunits, allowing continued

autophosphorylation on T287, which occurs *in trans* [24]and phosphorylation of PEAK1.

However, we accept that alternative, or contributory, mechanisms are conformational changes

within the holoenzyme induced by PEAK1 binding to one monomer that promote more

widespread enzyme activation and autophosphorylation, and T287 being protected from

phosphatase action by PEAK1 binding to CAMK2. In addition, CIM-mediated recruitment of

PEAK1 to one monomer within the Ca^{2+}-activated oligomer leads to phosphorylation of

PEAK1 by other activated monomers within the complex. This would be facilitated by the

relatively large number of consensus CAMK2 phosphorylation sites present on PEAK1 (41

R/KXXS/T sites in total). Importantly, a precedent for a reciprocally-activating interaction of

CAMK2 with a binding partner exhibiting a pseudosubstrate CIM is provided by

CAMK2/TIAM1. Here, binding of the TIAM1 CIM to CAMK2 leads to sustained

phosphorylation of TIAM1 by CAMK2[31].

An interesting question is the stoichiometry of the resulting PEAK1/CAMK2 complexes.

While CAMK2 forms a large holoenzyme complex of 12-14 subunits, PEAK1 can form dimers

via the SHED domain and also higher order oligomers via pseudokinase domain interfaces[18].

These properties may facilitate formation of assemblies with CAMK2 oligomers. In addition,

the localization of the PEAK1 CIM towards the N-terminus of a largely disordered extension

provides considerable flexibility for assembly of large multimeric protein complexes.

Interestingly, in contrast to PEAK1, PEAK2 lacks a consensus CIM, and its dimerization

potential is essential for CAMK2 association. While the impact of PEAK2 on CAMK2

regulation requires further characterization, one possibility is that PEAK2 contributes to

assembly of higher-order complexes of PEAK1/2 with CAMK2.

While PEAK1 is known to be regulated by site-selective tyrosine phosphorylation[6, 19, 36], our

study provides the first demonstration of positive regulation of PEAK1 by serine/threonine

phosphorylation. Furthermore, we determine that like tyrosine phosphorylation, this regulates

the PEAK1 interactome. In particular, the two PEAK1 CIM mutants, which are poor substrates

for CAMK2, exhibit decreased association with PEAK2. This provides one explanation for the

role of CAMK2 in PEAK1-enhanced migration, since heterotypic interaction with PEAK2 is

required for this process[16]. Conceivably, CAMK2-mediated phosphorylation may regulate

PEAK1 conformation and accessibility of the dimerization and/or oligomerization interfaces.

Another protein that exhibits significantly reduced binding to the PEAK1 CIM mutants is 14-

3-3. This is consistent with the overlap between the minimal CAMK2 phosphorylation site

consensus (R/KXXS/T) and the 14-3-3 mode 1 motif for binding, RXXpS/TXP. In the case of

PEAK3, 14-3-3 binding to phosphorylated PEAK3 S69 impairs Grb2 and Crk recruitment to

the scaffold, and the PEAK3 S69A mutant exhibits markedly enhanced ability to promote cell

migration[23]. With PEAK1, the overall effect of CAMK2-mediated phosphorylation on PEAK1-

promoted migration/invasion is positive, so 14-3-3 binding may downregulate binding of

negative regulatory factors, and/or increase association with positive ones, such as PEAK2.

Reflecting their role in a variety of disease states, pseudokinases have emerged as potential

therapeutic targets, albeit challenging ones given their lack of catalytic activity. Possible

therapeutic intervention strategies include the use of small molecule allosteric regulators,

therapeutic antibodies for cell surface receptor pseudokinases such as HER3, targeted

degradation approaches eg using PROTACs, and pharmacologic modulation of 'actionable'

downstream effector pathways[3]. The oncogenic pseudokinase PEAK1 presents a particular

conundrum, since it does not bind ATP or GTP, the dimerization interface is relatively large

and difficult to target using small molecule drugs, and PEAK1 is an intracellular protein and

therefore not accessible by standard therapeutic antibody approaches[5]. However, in this study

we identify a novel strategy for targeting PEAK1 signalling that exploits the functional

dependency of this kinase on a readily 'actionable' *bona fide* kinase, CAMK2. Importantly,

supporting therapeutic targeting of this serine/threonine kinase, CAMK2 isoforms are

overexpressed in a variety of human cancers, including breast cancer, where they promote cell

proliferation, migration and invasion[20, 21]. They are also positively associated with metastasis.

In an animal model of prostate cancer metastasis, genetic ablation of all CAMK2 family

members significantly reduced lymph node metastasis[37]. Expression, as well as activation, of

CAMK2 is significantly higher in lymph node metastases than primary breast cancers, and high

expression of the CAMK2 family is associated with worse DMFS in TNBC[21]. In addition, brain

metastatic colonization of TNBC is associated with CAMK2 signaling[38]. Our work now

provides novel insights into the oncogenic mechanism of CAMK2, with a key role being

CAMK2-mediated phosphorylation of PEAK1. This promotes heterodimerization of PEAK1

with PEAK2, a process required for PEAK1 to promote cell migration, presumably because

PEAK2 recruits/activates critical effectors such as Csk and Stat3 [16]. In addition, the activated CAMK2 may phosphorylate other downstream targets that promote metastasis, such as the Rho family GEF TIAM1 [31], and regulate the focal adhesion proteins FAK and paxillin[39].

In general, characterization of the oncogenic role of CAMK2 has been compromised by a poor understanding of regulatory mechanisms, with functional studies often utilizing constitutively active mutants, as well as a reliance on tool compounds for CAMK2 inhibition, such as KN93, that have numerous off-target effects[40]. Our study addresses these issues by identifying the oncogenic pseudokinase PEAK1 as a novel feed-forward activator of CAMK2 and the second generation CAMK2D/G inhibitor RA306 as a suitable targeted agent to block PEAK1/CAMK2 signalling in TNBC and other cancers where PEAK1 is involved, such as pancreatic cancer. Importantly, RA306 avoids central nervous system side-effects and exhibits good oral bioavailability, highlighting its potential for oncology re-purposing[32]. In addition, the identification of a reciprocally-activating PEAK1/CAMK2 complex indicates that these two proteins could be used as predictive biomarkers to guide patient stratification for anti-CAMK2 treatment. Overall, our work identifies an unsuspected interaction across the kinase superfamily with significant implications for our understanding of cell signalling in normal and disease states and precision targeting of PEAK1 and CAMK2.

Methods

Plasmids and siRNAs

The retroviral expression vector pRetroX-IRES-DsRedExpress (DsR) containing N-terminal HA-tagged and C-terminal FLAG/Myc-tagged PEAK1 as well as its deletion mutants including ΔNterm, ΔCterm and ΔN1, together with the expression vector pMIG containing N-terminal HA-tagged and C-terminal FLAG-tagged PEAK2 as well as PEAK2 site mutants (L955A, L966A, W1382A) were previously generated in our laboratory[16, 18]. pRetroX-IRES-

DsRedExpress plasmid containing Flag-tagged PEAK1 N terminal (residues 1–324) deleted

and other mutants (Δ1-260, Δ261-300, Δ261-324, Δ301-324, R297A/L301A/R303A and

R297E/R303E) were purchased from GenScript.

pDEST-V1 (Addgene plasmid 73637) and pDEST-V2 (Addgene plasmid 73638) vectors were

assembled from various recombinant and synthetic components as described previously[22].

Vectors expressing V1- or V2-labeled fusions of PEAK1 or PEAK2 were generated by

recombination cloning from pDONR223-PEAK1 and pDONR223-PEAK2 constructs into

pDEST-V1 or pDEST-V2 destination vectors using Gateway LR Clonase enzyme mix (Life

Technologies) according to the manufacturer's instructions and verified by sequencing. The

expression vector encoding full-length Venus fluorescent protein was a gift from S. Michnick

(University of Montreal).

CAMK2A kinase domain (residues 7-274) was cloned into a pET vector containing an N-

terminal 6×His followed by a SUMO tag. A D135N mutation was introduced by site-directed

mutagenesis.

Sequences of siRNAs were PEAK1 (ON-TARGETplus Human PEAK1 siRNA, cat. LQ-

005339-00-0010), PEAK2 (MISSION Human PEAK2 siRNAs, cat. #1:

SASI_Hs02_00316033, #2: SASI_Hs02_00316034, #3: SASI_Hs02_00316035, #4:

SASI_Hs02_00316036), CAMK2D (siGENOME Human CAMK2D siRNAs, cat. #1: D-

004042-02-0002, #2: D-004042-03-0002, #3: D-004042-04-0002, #4: D-004042-05-0002),

CAMK2G (siGENOME Human CAMK2G siRNAs, cat. #1: D-004536-01-0005, #2: D-

004536-03-0005, #3: D-004536-05-0005, #4: D-004536-06-0005). Negative control siRNA

was ON-TARGET plus Non-targeting pool (Dharmacon, D-001810-10-20).

Antibodies

The following antibodies were purchased from Santa Cruz Biotechnology (Dallas, TX):

PEAK1 (cat. sc-100403), β-actin (cat. sc-69879) and Pan 14-3-3 (cat. sc-1657). CAMK2D (cat.

ab181052) and CAMK2G (cat. ab201966) were purchased from Abcam. Phospho-Akt

Substrate (RXXS*/T*) (cat. 9614), T-CAMK2 (cat. 4436), CAMK2-pT287 (cat. 12716), PLC-

γ (cat. 5690S), PLC-γ-pY783 (cat.14008S) and cleaved PARP Asp 214 (cat. 9546) were

purchased from Cell Signaling Technology. Flag (cat. F1804 and F7425), and α-tubulin (cat.

T5168) were purchased from Sigma. The following antibodies were also used: CAMK2D

(Genetex, cat. GTX111401), HA (Roche Applied Science, cat. 11867423001). Phospho-

Ser/Thr antibody (cat. 612548) was purchased from BD Transduction Laboratories. The

custom-made PEAK2 (SgK223) antibody was generated by Cambridge Research

Biochemicals Ltd. by standard procedures and described previously[14].

Inhibitors and chemicals

RA306 was custom synthesized by Reagency (cat. RGNCY-0117). The SFK inhibitor eCF506

was obtained from Professor Margaret Frame (Edinburgh Cancer Research Centre). Captisol

(cat. A508148) was purchased from AmBeed. BAPTA-AM (cat. A1076) and DMSO (cat.

d8418) were purchased from Sigma. Doxorubicin-HCl was obtained from Selleckchem (cat.

S1208).

Cell lines and Tissue Culture

HEK293T cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM, Gibco,

cat. 1200046) supplemented with 10% (v/v) fetal bovine serum (FBS; Fisher Biotech, cat. S-

FBS-US-015). MDA-MB-231 parental cells were obtained from EG&G Mason Research

Institute, Worcester, MA. MDA-MB-231 cells stably expressing the murine ecotropic receptor

(MDA-MB-231_EcoR) were a kind gift from A/Prof Maija Kohonen-Corish at the Garvan

Institute of Medical Research. MDA-MB-468 cells were obtained from the American Type

Culture Collection (ATCC). The highly metastatic MDA-MB-231 cell line (MDA-MB-

231_HM) was originally established by Prof. Zhou Ou from Fudan University Shanghai Cancer

Center in China. The MDA-MB-231_HM cell line expressing luciferase/mCherry was a kind

gift from A/Prof. Erica Sloan at Monash University. In this paper, MDA-MB-231_HM refers

to the cell line expressing luciferase/mCherry.

Breast cancer cell lines were maintained in RPMI-1640 medium (Life Technologies, cat.

11875119 and Gibco, cat. 31800-089) supplemented with 10% FBS (Fisher Biotec, cat. S-FBS-

US-015), 10 µg/mL Actrapid penfill insulin (Clifford Hallam Healthcare), and 20 mM HEPES

(Life Technologies, cat. 15630080). All cell lines were authenticated by short tandem repeat

polymorphism, single-nucleotide polymorphism, fingerprint analyses and underwent routine

mycoplasma testing by PCR.

Cell Lysis, Immunoprecipitation, and Immunoblotting

Cell lysates for immunoblotting and immunoprecipitation (IP) were prepared using

radioimmunoprecipitation (RIPA) buffer and normal lysis buffer (NLB), respectively[41]. For IP

of overexpressed Flag- or HA-tagged proteins, cell lysates were incubated with either anti-

FLAG M2 affinity-agarose beads (Sigma, cat. A2220) or red anti-HA affinity gel (Sigma, cat.

E6779) at 4°C for 2 - 4 h with end-to-end rotation. After the incubation, the beads were washed

extensively with ice-cold lysis buffer, the immune complexes were eluted with SDS-PAGE

loading buffer, and subjected to Western blotting analysis, which was performed following

SDS-polyacrylamide gel electrophoresis using 5% stacking gels and 8%, 10% or 12%

separating gels. In order to ensure that phospho and total antibodies were used at non-limiting

concentrations, Western blot analyses were undertaken on dilution series of cell lysates.

Generation of PEAK1 knock-out cells

Stable MDA-MB-231_HM PEAK1 knockout cells were generated using the Synthego

Immortalized Cell Nucleofection Protocol as per manufacturer's instructions (Synthego).

Briefly, ribonucleoprotein (RNP) complexes were assembled by combining Cas9 2NLS

(Synthego), human PEAK1 sgRNA (Gene Knockout Kit v2, Synthego) and Nucleofector

SolutionTM L + supplement (Synthego) and incubating at room temperature for 10 mins. RNP

complexes were combined with cells in NucleocuvetteTM strips immediately prior to

electroporation. Cells were electroporated using the 4D-X Core unit (Lonza) using setting CH-

125. Cells were immediately resuspended in normal culture medium following electroporation.

Cell lysates were collected 72h post-electroporation to confirm PEAK1 knockout efficiency by

Western blot.

Transfections

SMARTpool siRNA or individual siRNA duplexes were applied to cells using DharmaFECT

(Dharmacon, Lafayette, CO) transfection reagent according to the manufacturer's instructions.

Plasmid transfections were performed using Lipofectamine 3000 (Invitrogen) according to the

manufacturer's instructions.

BiCAP purification

This was performed as previously described[22]. Briefly, HEK293T cells were transfected with

the appropriate BiCAP constructs. Cells were lysed with ice-cold lysis buffer comprising 50

mM Tris (pH 7.4), 150 mM NaCl, 1 mM EDTA, 1% (v/v) TritonX-100 supplemented with

fresh EDTA-free protease inhibitor cocktail and 0.2 mM sodium orthovanadate. Samples were

cleared by centrifugation at 18,000 g for 10 min at 4 °C to remove cellular debris. Total protein

concentration was determined by Bradford assay using Bio-Rad Protein Assay Dye Reagent

Concentrate (Bio-Rad Laboratories) according to the manufacturer's instructions. To purify

BiCAP dimers (or Venus alone as a negative control for nonspecific interactions), 1 mg of total

lysate was incubated with 35 μl of GFP-Trap_A agarose beads slurry (ChromoTek GmbH) for

2 h at 4°C with end-to-end rotation. Beads were extensively washed with wash buffer [10 mM

Tris (pH 7.5), 150 mM NaCl, 0.5 mM EDTA] and subjected to on-bead digestion with trypsin,

and the resulting peptides were analyzed by mass spectrometry (MS). Samples from each

experimental condition were subjected to data-dependent acquisition (DDA) to construct a

spectral library. Three biological replicates for each condition were then analysed by data-

independent acquisition (DIA) to characterize the corresponding interactomes. Conditions for

DDA and DIA are detailed below.

Mass spectrometry-based proteome analysis

MS analysis was based on an UltiMate 3000 RSLC nano-LC system (Thermo Fisher Scientific)

coupled to a Q Exactive Plus Mass Spectrometer (Thermo Fisher Scientific). The peptides were

loaded via a Pepmap100 trap column, 100 μm × 2 cm nanoviper, (Thermo Fisher Scientific),

eluted and separated on a Pepmap100 RSLC analytical column, 75 μm × 50 cm, nanoViper,

C18 (Thermo Fisher Scientific). For each injection, 1 μg of peptides was loaded onto the

column and eluted using increasing concentrations of buffer B (80% acetonitrile / 0.1% formic

acid) at 250 nL/min over 150 min. The eluent was nebulized and ionized using a

nanoelectrospray source (ThermoFisher Scientific) with a distal-coated fused silica emitter

(Trajan, Ringwood, Victoria, Australia). The capillary voltage was set at 1.7 kV.

To generate spectral libraries for proteomics, the MS was firstly operated in DDA mode to

automatically switch between full MS scans and tandem MS/MS scans. Survey full scan MS

spectra (m/z 375–1700) were acquired in the Orbitrap with 70 000 resolution after

accumulation of ions to a 1×10^6 target value with a maximum injection time of 120 ms. Up to

12 of the most intensely charged ions (z ≥ +2) were sequentially isolated and fragmented in the

collision cell by higher-energy collisional dissociation with the following parameters, fixed

injection time of 120 ms, 35 000 resolution and automatic gain control target of 5×10^5.

Dynamic exclusion was set to 15 s.

The DIA mode consisted of a MS1 scan (scan range from 400 to 1220 m/z, at resolution of

35 000, an AGC target of 5×10^6 and a maximum ion injection time of 120 ms) followed by

MS2 scans with 60 DIA windows at a resolution of 35 000, an AGC target of 3×10^6 with

automatic injection time, isolation window of 30 m/z. The stepped collision energies were

22.5%, 25%, and 27.5%. The spectra were recorded in profile type.

Data Processing

The DDA spectra were analyzed with MaxQuant (version 1.5.2.8) [42] against the human UniProt

fasta database (v2015–08, 20,210 entries) using default settings. The spectral library was then

generated in Spectronaut® software (version 8, Biognosys, Schlieren, Switzerland) and

normalized to iRT peptides. The DIA data were firstly transformed into htrms format and

analyzed with Spectronaut® using default settings. The MS data were searched against the same

human UniProt database and the aforementioned spectral library. The false discovery rate

(FDR) was set to 1% at peptide and protein level. Perseus (version 2019) was used to normalize

the data and impute for missing values [43]. Candidate interacting proteins were defined by

applying the following criteria: p value <0.05 by two-tailed *t*-test and a log2 fold change of ≥

1.5 against the Venus control. Pathway enrichment was undertaken using the WikiPathways

Database.

Protein expression and purification

CAMK2A D135N kinase domain was recombinantly expressed in BL21(DE3) cells

(Millipore) as previously described[30]. Protein expression was induced for ~6 h by 1 mM

isopropyl β-a-D-1-thiogalactopyranoside (IPTG; GoldBio) in an 18 °C shaker. Cell pellets were

resuspended and lysed in Buffer A [25 mM Tris pH 8.5, 50 mM KCl, 40 mM imidazole, 10%

glycerol] with protease inhibitor mixture [0.2 mM AEBSF, 5.0 μM leupeptin, 1 μg/ml

pepstatin, 1 μg/ml aprotinin, 0.1 mg/ml trypsin inhibitor, 0.5 mM benzamidine] and DNase (1

μg/ml; Sigma). Protein purification was performed using an ÄKTA pure chromatography

system at 4 °C. After pelleting the cell debris, clarified cell lysate was loaded onto two 5 mL

HisTrap FF Ni-NTA Sepharose columns (GE) and eluted with a combination of 50% buffer A

and 50% buffer B [25 mM Tris-HCl pH 8.5, 150 mM KCl, 1 M imidazole, 10% glycerol] for

a final concentration of 0.5 M imidazole. Protein was desalted using a HiPrep 26/10 desalting

column (GE), and His-SUMO tags were cleaved with Ulp1 protease overnight at 4 °C in buffer

C [25 mM Tris-HCl pH 8.5, 40 mM KCl, 40 mM imidazole, 2 mM tris(2-carboxyethyl)

phosphine (TCEP), 10% glycerol]. Cleaved tags were separated by a subtractive Ni-NTA step

followed by an anion exchange step performed with a 5-ml HiTrap Q-FF (GE) and protein was

eluted with a KCl gradient. Eluted proteins were concentrated and further purified in gel

filtration buffer [25 mM tris-HCl pH 8.0, 150 mM KCl, 1 mM TCEP, 10% glycerol] using a

Superdex 75 10/300 GL size exclusion column (GE). Pure fractions were then concentrated,

aliquoted, flash-frozen in liquid nitrogen, and stored at −80 °C until further use.

Peptide synthesis

The PEAK1 peptide used for ITC was synthesized to >95% purity with acetylation at the N-

terminus and amidation at the C-terminus (Genscript). The peptide sequence was as follows:

KKWNTIPLRNKSLQRICAVDYDDSYD.

Isothermal titration calorimetry

ITC data were obtained using a Malvern Auto-iTC200 calorimeter (UMass Amherst). Peptide

was dissolved in gel filtration buffer with 6% v/v DMSO and concentration was determined

using $\varepsilon280 = 8370$ M-1cm-1. CAMK2 kinase domain was diluted in gel filtration buffer with

5.6% v/v DMSO and concentration was determined with $\varepsilon280 = 43850$ M-1cm-1. Titrations

were performed with the peptide as the titrant. Blank titrations were performed by injecting the

peptide into gel filtration buffer. All titrations were performed at 20 °C using the standard 19-

injection method where one injection of 0.4 μL is followed by 18 x 2 μL injections with 150 s

between injections. The stir speed was set to 750 rpm, and the reference power was set to 10.

Data were buffer-subtracted and analyzed using MicroCal ITC Origin add-on dissociation

model v1.00 using the one-site fitting model.

Comparative modelling of PEAK1(291-320):CAMK2

Several candidate structural models of PEAK1(291-320):CAMK2 were generated by

AlphaFold-Multimer[44, 45] (v3) using the ColabFold[46] (v1.5.1) community implementation.

Multiple sequence alignments and template searches were performed by MMseqs[44] and

HHsearch using default settings. Custom templates of CAMK2 in complex with known tight

binders were required to achieve models with acceptable metrics of pLTTD and pTM scores.

These interactions were manually curated from crystal structures determined by Özden et al[30]

(crystal symmetry mates were discarded). Six recycle rounds were conducted and final models

were ranked according to pTMscore. The best scoring models were relaxed with Amber[47] to

minimize steric clashes.

Cross species multiple sequence alignment of PEAK1 N-terminus

To generate a multiple sequence alignment, two rounds of PSI-BLAST[48] were performed

against the non-redundant clustered database[49] (to increase taxonomic depth) using human

PEAK1 as a query sequence. A cutoff of expectation value of 1×10^{-3} was used to select

sequences for subsequent rounds and alignment. Finally, the top 50 and top 250 sequences

(sorted by expectation value) were subjected to alignment in Clustal Omega[50, 51]. The full

sequence range for the top 250 sequences was subjected to a MEME[52] search to identify

consensus motifs.

Proximity ligation assay (PLA)

PLA was performed to detect protein-protein interactions in MDA-MB-231 cells using

Duolink™ In Situ Red Starter Kit (Sigma, cat. DUO92101). All the reagents, including probes,

detection reagents and wash buffers were provided in this kit. On day 1, 6×10^{4} cells were seeded

per well in a 48-well plate and incubated at 37 °C overnight. On day 2, cells were gently washed

3 times with 1x phosphate-buffered saline [PBS; NaCl (136 mM), KCl (2.6 mM), Na2HPO4

(10 mM), KH2PO4 (1.76 mM)], and then fixed in 4% (v/v) paraformaldehyde (VWR, cat.

ALFA43368.9M) for 30 min on ice. Following that, cells were washed with 1xPBS again and

permeabilized with 0.2% Triton X-100 (Sigma, cat. T8787) for 3 min at room temperature.

After another 1xPBS washing step, cells were incubated with blocking solution for 30 min at

37 °C, following which, cells were incubated with two primary antibodies from different

species against the targeted proteins overnight at 4 °C. On day 3, cells were washed with 1x

Tris buffered saline (TBS) solution with the detergent Tween-20 [TBS-T; Tris base (50 mM),

NaCl (150 mM), pH 7.6, Tween-20 (0.01% v/v)], then PLA probes were added to the cells and

incubated for 60 min at 37 °C. Following ligation and amplification steps, samples were

mounted with Mounting Medium with DAPI (Invitrogen, cat. P-36931). Cells were

photographed using a confocal microscope and dot numbers were analyzed using ImageJ

software (Version 1.52).

Calcium Assay

MDA-MB-231 EcoR cells expressing DsR empty vector, wildtype PEAK1 or PEAK1-CIM mutants were generated by retroviral-mediated transduction as previously described[53]. Cells were seeded into FluoroDish cell culture dishes (World Precision Instruments, cat. FD35-100) and cultured overnight. The medium was then replaced with starvation medium (RPMI-1640 medium supplemented with 0.4% FBS, 10 μg/mL Actrapid penfill insulin, 20 mM HEPES) supplemented with 5 μM Calcium indicator Fluo-4 AM (Invitrogen, cat. F14217). Cells were then incubated for 60 min at 37°C. The dye-loading media was removed and replaced with starvation medium without Fluo-4 AM. Images were acquired with a fluorescence microscope (Leica DMi8) using the FITC filter set with excitation peaking at 495 nm and emission peaking at 519 nm. The cellular fluorescence was quantitated using ImageJ software. A total of >200 cells were quantified per condition in each biological replicate. The background was subtracted from the cell signal to plot fluorescence intensity.

Proliferation assays

These were performed using the CellTiter-Glo® 2.0 Cell Viability Assay (Promega – G9241) as per the manufacturer's instructions. Briefly, MDA-MB-231 HM cells were harvested and seeded into High Attachment (HA) (Falcon, FAL353072) and ultra-Low attachment (LA) (Corning, CLS3474) plates at a density of 1000 cells/well. After 16-18h, Day 0 assays were undertaken and the cells were treated either with vehicle control (DMSO) or RA306 at varying concentrations. After 120 h, assays were performed on replicates of vehicle control and RA306-treated cells.

Transwell Migration/Invasion Assay

Transwell migration and invasion assays were essentially as previously described[54]. Transwell

migration chambers with 8-μm pores were purchased from Corning (cat. CLS3464-48EA) and

MERCK Millipore (cat. MCEP24H48). Matrigel invasion chambers pre-coated with Matrigel

were purchased from Corning (cat. 354483). Prior to seeding, cells were treated with

mitomycin C at 10μg/ml for 1 h to prevent cell division.

In vitro kinase assay

HEK293T cells were transiently transfected with plasmids encoding wildtype PEAK1 or the

CAMK2-interaction motif mutants. Cell lysates were prepared using RIPA buffer[41]

supplemented with 12.5 mM β-glycerol phosphate. Cell lysates were then incubated with anti-

FLAG M2 affinity-agarose beads (Sigma, cat. A2220) at 4°C for 2 h with end-to-end rotation.

After the incubation, the beads were washed extensively with ice-cold RIPA buffer and kinase

reaction buffer [Tris (pH 7.5) 40 mM, $MgCl_2$ 10 mM, $CaCl_2$ 0.5 mM]. After the last wash,

beads were incubated in kinase reaction buffer supplemented with 100 ng recombinant

CAMK2D (Invitrogen, cat. PV3373), 100 μM recombinant calmodulin (Enzo, cat. BML-

SE325-0001) and 2 mM Adenosine 5'-triphosphate (ATP) disodium salt hydrate (Sigma-

Aldrich, cat. A1852-1VL) for 30 min at 37 °C. Following incubation, the immune complexes

were eluted with SDS-PAGE loading buffer and subjected to Western blotting analysis.

Animal Studies

All procedures involving mice were conducted in accordance with National Health and

Medical Research Council (NHMRC) regulations on the use and care of experimental animals

and the study protocol approved by the Monash University Animal Ethics Committee.

Power calculations performed indicated group sizes of n = 8 per treatment group were required

to detect a significant difference in tumor growth inhibition of 50% between treated and control

(significance level: 0.05; power: 90%). All *in vivo* experiments were performed with 6-week-

old female BALB/c athymic nude mice obtained from the Animal Resources Centre (Canning

Vale, Australia) and housed in sterile conditions at Monash Animal Research Platform.

Throughout the study, each cage was randomly assigned to the experimental groups without

considering any other variable. Mice were randomly divided into four equal groups, cas9-

control cells with captisol (control), cas9-control cells with RA306, cas9-sgPEAK1 cells with

captisol and cas9-sgPEAK1 cells with RA306.

Xenograft model

Mice (n=40) were injected with 2×10^5 MDA-MB-231_HM Cas9 control or sgPEAK1 stable

knockdown cells suspended in 20μl PBS into the fourth mammary fat pad. One day post-

injection, RA306 (30mg/kg) or captisol-only control was administered via daily oral gavage

until the end of the experiment. To measure tumour burden, mice were injected with luciferin

(150 mg/kg) by an intraperitoneal route. Tumour burden was measured weekly using the AMI-

HTX imaging system (Spectral Instruments Imaging). Results are presented as mean +/- SEM

of total BLI signal (photons/s).

Tumour metastasis model

Mice (n=40) were injected with 2×10^5 MDA-MB-231_HM Cas9 control or sgPEAK1 stable

knockdown cells suspended in 20μl PBS into the fourth mammary fat pad. One day post-

injection, RA306 (30mg/kg) or captisol-only control was administered via daily oral gavage

until the end of the experiment. When the primary tumours from each group reached ~ 100

mm^3 in size, they were resected. Tumour metastasis to secondary sites was then measured

weekly using the AMI-HTX imaging system (Spectral Instruments Imaging). Organs were

dissected immediately following the final BLI image and re-imaged in the AMI-HTX ex vivo

to allow for direct measurement of organ-specific tumour burden via BLI. Results are presented

as mean tumour luciferase intensities +/- SEM of total BLI signal (photons/s).

Tumour tissue homogenization

Following dissection at the experimental endpoint, tumour tissues were transferred

immediately to dry ice. Tumour tissues were cut into small pieces and added to a 1.5ml tube

containing RIPA buffer and Zirconia tissue disruption beads (Research Products International,

cat. 9835) then homogenized for 10s at low speed and 30s at high speed at room temperature.

To prepare samples for Western blotting, homogenized samples were centrifuged for 20 min

at 20,000g at 4°C. The supernatant was then collected and the protein concentration determined

using either Bradford Ultra Protein Assay Reagent (Expedeon, cat. EP-BFU05L) or Pierce TM

BCA Protein Assay Reagents (Thermo Scientific, cat. 23225). The protein concentration of

each sample was subsequently adjusted using RIPA buffer, and SDS-PAGE loading buffer was

added prior to Western blotting analysis.

Association of PEAK1 and CAMK2D/G with overall survival and distant metastasis-free

survival

For overall survival analysis, mRNA expression, mutation profile and associated overall

survival data from 2509 breast cancer patients as part of the METABRIC trial [55] were

downloaded from the cBioPortal for Cancer Genomics portal (https://www.cbioportal.org/).

Breast cancer patients were divided and classified into three main groups each having either

low, or medium, or high expression of *CAMK2D, CAMK2G* or *PEAK1*, individually. Main

groups were then further categorised into four sub-groups exhibiting different combined

expression patterns: *i) CAMK2D* and *PEAK1* both high, *ii) CAMK2D* and *PEAK1* both low,

iii) CAMK2G and *PEAK1* both high, *iv) CAMK2G* and *PEAK1* both low, *v) CAMK2D* low and

PEAK1 high, *vi) CAMK2D* high and *PEAK1* low. Survival analyses comparing overall survival

between these sub-groups were subsequently performed using a Log-rank test (with p < 0.05

considered significant). The Log-rank test statistics and survival curves were generated using

Kaplan-Meier estimate and implemented using the Logrank package [55] in MATLAB 2023a.

To determine association with distant metastasis-free survival (DMFS), mRNA expression and

DMFS data were obtained for the following cohorts in the Gene Expression Omnibus (GEO)

database: Princess Margaret Cancer Centre (database accesssions GSE6532 and GSE9195),

Veridex LLC (GSE12093) and Bayer Technology Services GmbH (GSE11121). CAMK2D

data were unavailable in most datasets; therefore, this analysis focused on PEAK1 and

CAMK2G. To create a more robust and larger meta-dataset we applied z-score standardization

to the gene expression data of individual cohorts and then combined all cohort data to generate

a 527 patient meta-cohort. Subsequently, breast cancer patients were divided into low or high

expression groups for PEAK1 and CAMK2G, individually, based on the 33% percentile of

gene expression (bottom 33% vs top 67%). Finally, patient groups were categorized into two

subgroups based on combined expression patterns: *i)* PEAK1 and CAMK2G both low, and *ii)*

PEAK1 and CAMK2G both high. The DMFS was compared between these two subgroups

using a log-rank test (p < 0.05 considered significant). Log-rank test statistics and survival

curves were generated using the *survival* (version 3.5.8) and *survminer* (version 0.4.9)

packages in R.

Statistical analyses

Experimental data were subject to appropriate statistical analyses as detailed in the

corresponding figure legends.

Data availability

The mass spectrometry proteomics data for the spectral library and DIA analysis have been

deposited to the ProteomeXchange Consortium via the PRIDE[56] partner repository with the

dataset identifier PXD044872.

Reviewer account details:

Username: reviewer_pxd044872@ebi.ac.uk

Password: LHFHdDGH

The remaining data are available within the Article, Supplementary Information or Source Data

file.

References

1. Reiterer V, Eyers PA, Farhan H. Day of the dead: pseudokinases and
pseudophosphatases in physiology and disease. *Trends Cell Biol* **24**, 489-505 (2014).
850

851 2. Zeqiraj E, van Aalten DM. Pseudokinases-remnants of evolution or key allosteric
852 regulators? *Current opinion in structural biology* **20**, 772-781 (2010).
853

854 3. Kung JE, Jura N. Prospects for pharmacological targeting of pseudokinases. *Nat Rev*
855 *Drug Discov* **18**, 501-526 (2019).
856

857 4. Hou J*, et al.* Distinct PEAK3 interactors and outputs expand the signaling potential of
858 the PEAK pseudokinase family. *Sci Signal* **15**, eabj3554 (2022).
859

860 5. Patel O, Roy MJ, Murphy JM, Lucet IS. The PEAK family of pseudokinases, their
861 role in cell signalling and cancer. *FEBS J* **287**, 4183-4197 (2020).
862

863 6. Croucher DR*, et al.* Involvement of Lyn and the atypical kinase SgK269/PEAK1 in a
864 basal breast cancer signaling pathway. *Cancer Res* **73**, 1969-1980 (2013).
865

866 7. Agajanian M*, et al.* PEAK1 Acts as a Molecular Switch to Regulate Context-
867 Dependent TGFbeta Responses in Breast Cancer. *PLoS One* **10**, e0135748 (2015).
868

869 8. Abu-Thuraia A*, et al.* AXL confers cell migration and invasion by hijacking a
870 PEAK1-regulated focal adhesion protein network. *Nature communications* **11**, 1-20
871 (2020).
872

873 9. Ding C*, et al.* Overexpression of PEAK1 contributes to epithelial-mesenchymal
874 transition and tumor metastasis in lung cancer through modulating ERK1/2 and JAK2
875 signaling. *Cell Death Dis* **9**, 802 (2018).
876

877 10. Kelber JA, *et al.* KRas induces a Src/PEAK1/ErbB2 kinase amplification loop that
878 drives metastatic growth and therapy resistance in pancreatic cancer. *Cancer Res* **72**,
879 2554-2564 (2012).
880
881 11. Leroy C, *et al.* Quantitative phosphoproteomics reveals a cluster of tyrosine kinases
882 that mediates SRC invasive activity in advanced colon carcinoma cells. *Cancer Res*
883 **69**, 2279-2286 (2009).
884
885 12. Ounoughene Y, *et al.* SHED-Dependent Oncogenic Signaling of the PEAK3 Pseudo-
886 Kinase. *Cancers (Basel)* **13**, (2021).
887
888 13. Senda Y, Murata-Kamiya N, Hatakeyama M. C-terminal Src kinase-mediated EPIYA
889 phosphorylation of Pragmin creates a feed-forward C-terminal Src kinase activation
890 loop that promotes cell motility. *Cancer Sci* **107**, 972-980 (2016).
891
892 14. Tactacan CM, *et al.* The pseudokinase SgK223 promotes invasion of pancreatic
893 ductal epithelial cells through JAK1/Stat3 signaling. *Mol Cancer* **14**, 139 (2015).
894
895 15. Lecointre C, *et al.* Oncogenic Signalling of PEAK2 Pseudokinase in Colon Cancer.
896 *Cancers (Basel)* **14**, (2022).
897
898 16. Liu L, *et al.* Homo- and Heterotypic Association Regulates Signaling by the
899 SgK269/PEAK1 and SgK223 Pseudokinases. *J Biol Chem* **291**, 21571-21583 (2016).
900
901 17. Ha BH, Boggon TJ. The crystal structure of pseudokinase PEAK1 (Sugen kinase 269)
902 reveals an unusual catalytic cleft and a novel mode of kinase fold dimerization. *J Biol*
903 *Chem* **293**, 1642-1650 (2018).
904
905 18. Patel O, *et al.* Structure of SgK223 pseudokinase reveals novel mechanisms of
906 homotypic and heterotypic association. *Nat Commun* **8**, 1157 (2017).
907
908 19. Zheng Y, *et al.* Temporal regulation of EGF signalling networks by the scaffold
909 protein Shc1. *Nature* **499**, 166-171 (2013).
910
911 20. Brzozowski JS, Skelding KA. The Multi-Functional Calcium/Calmodulin Stimulated
912 Protein Kinase (CaMK) Family: Emerging Targets for Anti-Cancer Therapeutic
913 Intervention. *Pharmaceuticals (Basel)* **12**, (2019).
914
915 21. Chi M, *et al.* Phosphorylation of calcium/calmodulin-stimulated protein kinase II at
916 T286 enhances invasion and migration of human breast cancer cells. *Sci Rep* **6**, 33132
917 (2016).
918
919 22. Croucher DR, *et al.* Bimolecular complementation affinity purification (BiCAP)
920 reveals dimer-specific protein interactions for ERBB2 dimers. *Sci Signal* **9**, ra69
921 (2016).
922
923 23. Roy MJ, *et al.* Structural mapping of PEAK pseudokinase interactions identifies 14-3-
924 3 as a molecular switch for PEAK3 signaling. *Nat Commun* **14**, 3542 (2023).
925

926 24. Bhattacharyya M, Karandur D, Kuriyan J. Structural Insights into the Regulation of
927   Ca(2+)/Calmodulin-Dependent Protein Kinase II (CaMKII). *Cold Spring Harb*
928   *Perspect Biol* **12**, (2020).
929

930 25. Tsien RY. A non-disruptive technique for loading calcium buffers and indicators into
931   cells. *Nature* **290**, 527-528 (1981).
932

933 26. Mandal S, Bandyopadhyay S, Tyagi K, Roy A. Recent advances in understanding the
934   molecular role of phosphoinositide-specific phospholipase C gamma 1 as an emerging
935   onco-driver and novel therapeutic target in human carcinogenesis. *Biochim Biophys*
936   *Acta Rev Cancer* **1876**, 188619 (2021).
937

938 27. Hochgrafe F, *et al.* Tyrosine phosphorylation profiling reveals the signaling network
939   characteristics of Basal breast cancer cells. *Cancer Res* **70**, 9391-9401 (2010).
940

941 28. Jones NP, Peak J, Brader S, Eccles SA, Katan M. PLCgamma1 is essential for early
942   events in integrin signalling required for cell motility. *J Cell Sci* **118**, 2695-2706
943   (2005).
944

945 29. Temps C, *et al.* A Conformation Selective Mode of Inhibiting SRC Improves Drug
946   Efficacy and Tolerability. *Cancer Res* **81**, 5438-5450 (2021).
947

948 30. Ozden C, *et al.* CaMKII binds both substrates and activators at the active site. *Cell*
949   *Rep* **40**, 111064 (2022).
950

951 31. Saneyoshi T, *et al.* Reciprocal Activation within a Kinase-Effector Complex
952   Underlying Persistence of Structural LTP. *Neuron* **102**, 1199-1210 e1196 (2019).
953

954 32. Beauverger P, *et al.* Reversion of cardiac dysfunction by a novel orally available
955   calcium/calmodulin-dependent protein kinase II inhibitor, RA306, in a genetic model
956   of dilated cardiomyopathy. *Cardiovasc Res* **116**, 329-338 (2020).
957

958 33. Yang X, *et al.* The pseudokinase NRBP1 activates Rac1/Cdc42 via P-Rex1 to drive
959   oncogenic signalling in triple-negative breast cancer. *Oncogene* **42**, 833-847 (2023).
960

961 34. Lecointre C, *et al.* Dimerization of the Pragmin Pseudo-Kinase Regulates Protein
962   Tyrosine Phosphorylation. *Structure* **26**, 545-554 e544 (2018).
963

964 35. Sausgruber N, *et al.* Tyrosine phosphatase SHP2 increases cell motility in triple-
965   negative breast cancer through the activation of SRC-family kinases. *Oncogene* **34**,
966   2272-2278 (2015).
967

968 36. Bristow JM, Reno TA, Jo M, Gonias SL, Klemke RL. Dynamic phosphorylation of
969   tyrosine 665 in pseudopodium-enriched atypical kinase 1 (PEAK1) is essential for the
970   regulation of cell migration and focal adhesion turnover. *J Biol Chem* **288**, 123-131
971   (2013).
972

973 37. Yu G, *et al.* Organelle-Derived Acetyl-CoA Promotes Prostate Cancer Cell Survival,
974   Migration, and Metastasis via Activation of Calmodulin Kinase II. *Cancer Res* **78**,
975   2490-2502 (2018).

977 38. Ren D, *et al.* Targeting Brain-Adaptive Cancer Stem Cells Prohibits Brain Metastatic
978 Colonization of Triple-Negative Breast Cancer. *Cancer Res* **78**, 2052-2064 (2018).
979

980 39. Easley CA, Brown CM, Horwitz AF, Tombes RM. CaMK-II promotes focal adhesion
981 turnover and cell motility by inducing tyrosine dephosphorylation of FAK and
982 paxillin. *Cell Motil Cytoskeleton* **65**, 662-674 (2008).
983

984 40. Pellicena P, Schulman H. CaMKII inhibitors: from research tools to therapeutic
985 agents. *Front Pharmacol* **5**, 21 (2014).
986

987 41. Brummer T, *et al.* Increased proliferation and altered growth factor dependence of
988 human mammary epithelial cells overexpressing the Gab2 docking protein. *Journal of*
989 *Biological Chemistry* **281**, 626-637 (2006).
990

991 42. Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized
992 p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat*
993 *Biotechnol* **26**, 1367-1372 (2008).
994

995 43. Tyanova S, *et al.* The Perseus computational platform for comprehensive analysis of
996 (prote)omics data. *Nat Methods* **13**, 731-740 (2016).
997

998 44. Evans R, *et al.* Protein complex prediction with AlphaFold-Multimer. *bioRxiv*,
999 2021.2010.2004.463034 (2022).
1000

1001 45. Jumper J, *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*
1002 **596**, 583-589 (2021).
1003

1004 46. Mirdita M, Schutze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M.
1005 ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682 (2022).
1006

1007 47. Hornak V, Abel R, Okur A, Strockbine B, Roitberg A, Simmerling C. Comparison of
1008 multiple Amber force fields and development of improved protein backbone
1009 parameters. *Proteins* **65**, 712-725 (2006).
1010

1011 48. Altschul SF, *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein
1012 database search programs. *Nucleic Acids Res* **25**, 3389-3402 (1997).
1013

1014 49. Steinegger M, Soding J. MMseqs2 enables sensitive protein sequence searching for
1015 the analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028 (2017).
1016

1017 50. McWilliam H, *et al.* Analysis Tool Web Services from the EMBL-EBI. *Nucleic Acids*
1018 *Res* **41**, W597-600 (2013).
1019

1020 51. Sievers F, *et al.* Fast, scalable generation of high-quality protein multiple sequence
1021 alignments using Clustal Omega. *Mol Syst Biol* **7**, 539 (2011).
1022

1023 52. Bailey TL, Elkan C. Fitting a mixture model by expectation maximization to discover
1024 motifs in biopolymers. *Proc Int Conf Intell Syst Mol Biol* **2**, 28-36 (1994).
1025

1026 53. Croucher DR, Rickwood D, Tactacan CM, Musgrove EA, Daly RJ. Cortactin
1027   modulates RhoA activation and expression of Cip/Kip cyclin-dependent kinase
1028   inhibitors to promote cell cycle progression in 11q13-amplified head and neck
1029   squamous cell carcinoma cells. *Mol Cell Biol* **30**, 5057-5070 (2010).
1030
1031 54. Bennett HL, Brummer T, Jeanes A, Yap AS, Daly RJ. Gab2 and Src co-operate in
1032   human mammary epithelial cells to promote growth factor independence and
1033   disruption of acinar morphogenesis. *Oncogene* **27**, 2693-2704 (2008).
1034
1035 55. Pereira B*, et al.* The somatic mutation profiles of 2,433 breast cancers refines their
1036   genomic and transcriptomic landscapes. *Nat Commun* **7**, 11479 (2016).
1037
1038 56. Perez-Riverol Y*, et al.* The PRIDE database resources in 2022: a hub for mass
1039   spectrometry-based proteomics evidences. *Nucleic Acids Res* **50**, D543-D552 (2022).
1040
1041

Acknowledgements

The authors acknowledge the Monash Proteomics and Metabolomics Facility, Monash

University, for the provision of MS instrumentation, training and technical support, and

Monash Micro Imaging for assistance with the calcium imaging experiments. The work was

supported by research grants to RJD from the National Health and Medical Research Council

of Australia (Ideas Grant 2003599) and Australian Research Council (ARC) (DP220103638

and DP190103672) and to MMS (NIH R35GM145376). DRC acknowledges fellowship

support from the National Breast Cancer Foundation (NBCF Fellowship IIRS-20-032), AME

is a Victorian Department of Health and Human Services Victorian Cancer Agency Mid-Career

Fellow (MCRF21036) and CB is an ARC DECRA Fellow (DE240100992). VCBN is a

member of the Molecular and Cellular Biology Graduate Program, University of

Massachusetts, Amherst, MA 01003, USA and is supported by a fellowship from the

University of Massachusetts as part of the Chemistry-Biology Interface Training Program

(National Research Service Award T32 GM139789).

1056

Author Contributions

Conceptualization, RJD; Methodology, XY, XM, DRC, CB, SB, SS, LKN, ACC, MMS, AME,

RJD; Investigation, XY, XM, DRC, EVN, KCC, CH, SLL, TZ, CB, VCBN, SB, SS, TCCLKS;

Writing – Original Draft, XY, XM, RJD; Writing – Review and Editing, TRC, MMS, AME,

RJD; Visualization, XY, TRC, RJD; Funding acquisition, DRC, MMS, CB, AME, RJD;

Supervision, XM, RJD; Project administration, RJD.

1063

1064   **Declaration of interests**

1065   The authors declare that they have no competing interests.

1066

1067   **Figure Legends**

1068

Figure 1. Identification of specific CAMK2 isoforms as PEAK1/2 interactors via BiCAP-

MS/MS. A-C. Results of BiCAP-MS/MS screen. Volcano plots indicate enriched interactors

for the PEAK1 homotypic complex (A), PEAK2 homotypic complex (B) and PEAK1/PEAK2

heterotypic complex (C). The known PEAK1 interactor Grb2 and targets of particular interest,

including specific CAMK2 family members, are highlighted in blue. Dotted lines indicate fold

change and p value cut-offs. **D. Western blotting validation of PEAK1 and PEAK2**

knockdown. PEAK1 and PEAK2 were knocked down using siRNA pools, with successful

knockdown confirmed by Western blotting. **E-F. Validation of PEAK1/2 association with**

CAMK2D and CAMK2G by proximity ligation assays. Left panels, representative

proximity ligation assay (PLA) images from MDA-MB-231 cells for interaction of PEAK1 (E)

or PEAK2 (F) with CAMK2D and CAMK2G. Cells were transfected with either si-PEAK1

(E), si-PEAK2 (F) or si-Control. Right panels, quantification of protein interactions in each

cell. The data are expressed as dot number divided by cell number. In each condition, more

than 50 cells have been counted. Error bars represent the standard error of the mean from n=3

independent experiments. * = p < 0.05, ** = p < 0.01 by student's t-test.

1084

Figure 2. PEAK1-mediated regulation of CAMK2 activity in triple negative breast cancer

cells. A-D. Manipulation of PEAK1 expression modulates CAMK2 activity. PEAK1 was

knocked down using a siRNA (A-B) or overexpressed via transient transfection (C-D) in MDA-

MB-231 cells and autonomous activation of CAMK2 determined by Western blotting as

indicated. Blotting for 14-3-3 served as a loading control. Histograms indicate CAMK2 T287

phosphorylation normalized for total CAMK2 expression, with 'au' indicating arbitrary units.

E-F. PEAK1-promoted CAMK2 activation is calcium-dependent. MDA-MB-231 cells

were transfected with a PEAK1 construct in the presence or absence of 10 μM BAPTA-AM.

Cell lysates were Western blotted as indicated (E) and CAMK2 activation quantified as above

for CAMK2D/G combined (F), with data expressed relative to the DsR (vector)/DMSO control

which was arbitrarily set at 1. **G-H. Overexpression of PEAK1 increases intracellular Ca^{2+}.**

MDA-MB-231_EcoR cells stably overexpressing PEAK1 were loaded with calcium-sensitive

dye Fluo-4 AM and then imaged with a fluorescence microscope. Representative images from

3 independent experiments are shown (G) as well as quantification of Fluo-4 images (H),

presented as mean Fluo-4 fluorescence ± SEM. These are representative data from one

biological replicate from a total of three. **I-J. PEAK1 regulates tyrosine phosphorylation of**

PLCγ1. MDA-MB-231 cells were transfected with PEAK1 siRNA (I) or a PEAK1 expression

vector (J) and then cell lysates were Western blotted as indicated. The histograms indicate

Y783-phosphorylated PLCγ1 normalized for total PLCγ1 expression. For all panels, error bars

represent the standard error of the mean from n=3 independent experiments. NS indicates p >

0.05, * = p < 0.05, ** = p<0.01, *** = p<0.001, ****, p < 0.0001, by ratio paired t-test (B, D,

I, J), two-way ANOVA with Tukey's multiple comparisons test (F), or unpaired t-test (H).

1107

Figure 3. Mapping the CAMK2 binding region on PEAK1. A. Schematic structure of

PEAK1 and various truncation mutants. WT, wildtype. **B. Association of CAMK2D with**

specific PEAK1 deletion mutants. The indicated Flag-tagged PEAK1 deletion mutants were

expressed in HEK293T cells. Total cell lysates (TCL) and Flag IPs were subjected to Western

blotting as indicated. **C. Schematic representation of various PEAK1 N-terminal deletion**

mutants. Deleted regions of PEAK1 are indicated by red crosses. **D-E. Fine structure**

mapping of N-terminal regions required for the association of PEAK1 with CAMK2D

(D) or CAMK2G (E). Analysis was undertaken as in (B). The asterisk in 3D highlights

degradation products present for the Δ261-300 and Δ261-324 mutants. **F-G. PEAK1 binding**

to CAMK2 is required for CAMK2 activation. MDA-MB-231 cells were transfected with

WT PEAK1 or the Δ301-324 PEAK1 construct. Cell lysates were Western blotted as indicated

(F), with the dotted line in F indicating where the image was cropped to remove an irrelevant

lane. Raw data for the full image accompanies the manuscript. Relative CAMK2 activation,

as determined by T287 phosphorylation, was then quantified (G). Data are expressed relative

to vector control which was arbitrarily set at 1. All data in B, D and E are representative of at

least 2 independent experiments. In G, error bars represent the standard error of the mean from

n=3 independent experiments. NS indicates $p > 0.05$, * = $p < 0.05$ by one-way ANOVA with

Tukey's multiple comparisons test.

1126

1127 **Figure 4. Characterization of a conserved CAMK2 interaction motif (CIM) on PEAK1.**

1128 **A. Sequence alignment of the PEAK1 CIM with other CAMK2 binding partners**. The

1129 critical residues for CAMK2 interaction (R297, L301 and R303 in PEAK1) are highlighted in

1130 yellow. The canonical motif numbering (A306 being 0) is shown above the alignment. **B.**

1131 **Representative ITC measurement of CAMK2 kinase domain binding to PEAK1 CIM**

peptide. The mean Kd value from n=3 independent experiments is indicated. **C. Comparative modelling of the PEAK1/CAMK2 interaction by AlphaFold.** Surface representation of CAMK2 with PEAK1 amino acids 291-320 shown as ribbon (grey). The orientation is as in Supp Figure 6. CAMK2 is coloured by electrostatic charge. The PEAK1 binding site on CAMK2 is noticeably negative, with key arginine residues of PEAK1 mediating interactions with CAMK2. The N- and C-termini are labelled as N and C, respectively. **D. Surface representation of modelled PEAK1/CAMK2 interaction.** CAMK2 is grey and PEAK1 amino acids 291-320 are purple. **E-F. Focused view of key predicted molecular interactions.** Boxed regions from (D) illustrating predicted formation of salt bridges and electrostatic interactions (dotted lines) by specific PEAK1 residues across the surface of CAMK2. The canonical numbering of the PEAK1 motif is shown in white boxes.

Figure 5. The role of the PEAK1 CIM in CAMK2 activation. A-B. Effect of CIM mutation on intracellular Ca^{2+} levels. Constructs expressing WT PEAK1 or the indicated mutants were stably expressed in MDA-MB-231_EcoR cells via retroviral transduction. PEAK1 expression levels were determined by Western blot (A) and Ca^{2+} levels (B) assayed as in Figure 2G-H. Data in B represent mean Fluo-4 fluorescence $\pm$ SEM from one representative biological replicate from a total of two independent experiments. Statistical analysis utilized one way ANOVA with Dunnett's multiple comparisons test with **** indicating $p < 0.0001$. **C-D. Effect of CIM mutation on tyrosine phosphorylation of PLCγ1**. The indicated plasmids were transiently transfected into HEK293 cells and cell lysates Western blotted as indicated (C). PLCγ1 tyrosine phosphorylation was normalized for total expression (D). Data in D are expressed relative to the vector control which was arbitrarily set at 1 and represent the mean $\pm$ SEM of 3 independent biological replicates, * indicates $p<0.05$ by unpaired t-test. **E-G. Effect of CIM mutation on CAMK2 activation.** Plasmids encoding WT PEAK1 and the indicated

1157 mutants were transfected into MDA-MB-231 cells. Total cell lysates were Western blotted as

1158 indicated (E) and then relative CAMK2 activation was quantified (F-G). Data are expressed

1159 relative to the vector (DsR) control which was arbitrarily set at 1. NS indicates $p > 0.05$, ** =

1160 $p < 0.01$ and **** = $p < 0.0001$ by one-way ANOVA with Dunnett's multiple comparisons

1161 test.

1162

Figure 6. CAMK2 phosphorylates PEAK1 to regulate the PEAK1 interactome. A-B.

RA306 inhibits PEAK1 serine/threonine phosphorylation. MDA-MB-231_EcoR cells

stably expressing Flag-tagged PEAK1 were treated with RA306 (1 μM) for 24 h and cell lysates

were directly Western blotted or subject to immunoprecipitation with Flag antibody prior to

blotting as indicated (A). Relative serine/threonine phosphorylation of PEAK1, normalized for

total PEAK1, is shown in the histogram (B). **C. CAMK2 phosphorylates PEAK1 *in vitro*.**

Flag-tagged versions of WT and mutant PEAK1 proteins were expressed in HEK293T cells.

PEAK1 proteins were immunoprecipitated using anti-Flag antibody and following washing,

immediately treated with SDS-PAGE sample buffer (SB) (no incubation) or incubated in

kinase reaction buffer supplemented as indicated at 37°C for 30 min. Reactions were terminated

with SB. Samples were then Western blotted as indicated. The asterisk indicates a background

band present in all CAMK2D-incubated samples including the vector control while the

arrowhead highlights phosphorylated PEAK1. **D. Impact of CIM mutations on heterotypic**

association with PEAK2. Flag-tagged WT PEAK1 or PEAK1 mutants were expressed in

MDA-MB-231 cells and association with endogenous PEAK2 assayed by IP/Western. For

panel B, * = $p<0.05$ by ratio paired t-test.

1179

1180 **Figure 7. Role of CAMK2 in PEAK1-regulated biological endpoints in TNBC models. A.**

1181 **PEAK1 overexpression in combination with CAMK2D and CAMK2G knockdown in**

MDA-MB-231 cells. Cells were transfected with a PEAK1 plasmid in the presence or absence of siRNA-mediated CAMK2D or CAMK2G knockdown. Cell lysates were Western blotted as indicated. **B. Role of CAMK2D/G in PEAK1-promoted MDA-MB-231 cell migration.** Cells validated in (A) were subjected to a transwell migration assay. **C-D**, **Role of CAMK2 activation.** MDA-MB-231 cells were transfected with a PEAK1 plasmid and subject to transwell migration assays in the presence or absence of RA306 (C). Plasmids expressing WT PEAK1 and the Δ301-324 mutant that cannot activate CAMK2 were transiently transfected into MDA-MB-231 cells, and cells were subjected to a transwell migration assay (D). **E. Role of PEAK1/CAMK2 signalling in TNBC xenograft growth.** Control MDA-MB-231_HM cells (Cas9-control) or PEAK1 KO cells (Cas9-sgPEAK1) in combination with RA306 or vehicle control (captisol) were injected into the mammary fat pad. Left panel, representative whole body bioluminescent imaging (BLI) images from each group at different time points as indicated. Right panel, quantification of xenograft growth. Each treatment group exhibited $\geq$ 9 mice. **F. Role of PEAK1/CAMK2 signalling in TNBC metastasis.** Control MDA-MB-231_HM cells (Cas9-control) or PEAK1 KO cells (Cas9-sgPEAK1) in combination with RA306 or vehicle control (captisol) were injected into the mammary fat pad. At Day 19 post-injection when the primary tumour from each group reached $\sim$ 100 mm^3, they were resected, and tumour growth at secondary sites was measured. Left panel, representative whole body and *ex vivo* lung BLI images from each group at Day 72 post injection. Right panel, luciferase intensities of tumour metastasis to lung were generated from each group. Error bars represents the standard error of the mean from n=3 independent experiments (B-D) or from the individuals of each group (E & F). NS indicates $p > 0.05$, $* = p < 0.05$, $** = p < 0.01$, $*** = p < 0.001$, $**** = p < 0.0001$ by two-way ANOVA with Tukey's multiple comparisons test (B, C, E), one-way ANOVA with Tukey's multiple comparisons test (D), or one-way ANOVA with Dunnett's multiple comparisons test (F).

1207

Figure 8. Schematic representation of feed-forward regulation of CAMK2 by PEAK1.

PEAK1 enhances intracellular Ca^{2+} by promoting SFK-mediated tyrosine phosphorylation of

PLCγ1. This leads to classical activation of CAMK2 by Ca^{2+}/CaM and the αD helix rotates

out. The PEAK1 CIM then binds to the CAMK2 substrate binding site and competes with the

regulatory segment, effectively 'locking' CAMK2 in an active conformation. Sustained

phosphorylation of PEAK1 and CAMK2 itself (on T287) then occurs as the PEAK1 CIM

undergoes cycles of dissociation and re-association. However, other mechanisms may

contribute to the observed effects on CAMK2 activation, as indicated on the figure. CIM-

mediated recruitment of PEAK1 to the CAMK2 holoenzyme promotes serine/threonine

phosphorylation of PEAK1 which in turn positively regulates heterotypic association with

PEAK2.

1219

1220 **Supplementary Information Titles and Legends**

1221

1222 Supp.Table 1. BiCAP-MS/MS data for PEAK dimer-specific interactomes.

1223 Supp.Table 2. Pathway enrichment analysis for dimer-specific interactomes.

1224

Supp. Figure 1. Characterization of PEAK complex interactomes via BiCAP-MS/MS.

Supp. Figure 2. Manipulation of PEAK1 expression modulates CAMK2 activity in MDA-MB-

468 cells.

Supp. Figure 3. Effect of SFK inhibitor eCF506 on PLCγ1 Y783 phosphorylation.

Supp. Figure 4. Determination of the structural requirements for CAMK2 interaction with

PEAK2.

Supp. Figure 5. Sequence alignment of the PEAK1 CIM motif across diverse species.

Supp. Figure 6. Comparative modelling analysis by AlphaFold (to accompany Figure 4C-F).

Supp. Figure 7. Role of CAMK2 in regulating PEAK1 phosphorylation and the PEAK1
interactome.

Supp. Figure 8. Association of PEAK1 and CAMK2 expression with breast cancer patient
survival.

Supp. Figure 9. Effect of the CAMK2 inhibitor RA306 on cell proliferation and apoptosis.

Supp. Figure 10. Role of CAMK2 in PEAK1-regulated biological endpoints in TNBC models.

Supp. Figure 11. Genetic or pharmacological targeting of the PEAK1/CAMK2 axis reduces
TNBC development *in vivo*.

1241

1242

Figure 1.

Figure 1. Identification of specific CAMK2 isoforms as PEAK1/2 interactors via BiCAP-MS/MS. A-C. Results of BiCAP-MS/MS screen. Volcano plots indicate enriched interactors for the PEAK1 homotypic complex (A), PEAK2 homotypic complex (B) and PEAK1/PEAK2 heterotypic complex (C). The known PEAK1 interactor Grb2 and targets of particular interest, including specific CAMK2 family members, are highlighted in blue. Dotted lines indicate fold change and p value cut-offs. **D. Western blotting validation of PEAK1 and PEAK2 knockdown.** PEAK1 and PEAK2 were knocked down using siRNA pools, with successful knockdown confirmed by Western blotting. **E-F. Validation of PEAK1/2 association with CAMK2D and CAMK2G by proximity ligation assays.** Left panels, representative proximity ligation assay (PLA) images from MDA-MB-231 cells for interaction of PEAK1 (E) or PEAK2 (F) with CAMK2D and CAMK2G. Cells were transfected with either si-PEAK1 (E), si-PEAK2 (F) or si-Control. Right panels, quantification of protein interactions in each cell. The data are expressed as dot number divided by cell number. In each condition, more than 50 cells have been counted. Error bars represent the standard error of the mean from n=3 independent experiments. $* = p < 0.05$, $** = p < 0.01$ by student's t-test.

Figure 2.

Figure 2. PEAK1-mediated regulation of CAMK2 activity in triple negative breast cancer cells. A-D. Manipulation of PEAK1 expression modulates CAMK2 activity. PEAK1 was knocked down using a siRNA (A-B) or overexpressed via transient transfection (C-D) in MDA-MB-231 cells and autonomous activation of CAMK2 determined by Western blotting as indicated. Blotting for 14-3-3 served as a loading control. Histograms indicate CAMK2 T287 phosphorylation normalized for total CAMK2 expression, with 'au' indicating arbitrary units. **E-F. PEAK1-promoted CAMK2 activation is calcium-dependent.** MDA-MB-231 cells were transfected with a PEAK1 construct in the presence or absence of 10 μM BAPTA-AM. Cell lysates were Western blotted as indicated (E) and CAMK2 activation quantified as above for CAMK2D/G combined (F), with data expressed relative to the DsR (vector)/DMSO control which was arbitrarily set at 1. **G-H. Overexpression of PEAK1 increases intracellular Ca^{2+}.** MDA-MB-231_EcoR cells stably overexpressing PEAK1 were loaded with calcium-sensitive dye Fluo-4 AM and then imaged with a fluorescence microscope. Representative images from 3 independent experiments are shown (G) as well as quantification of Fluo-4 images (H), presented as mean Fluo-4 fluorescence ± SEM. These are representative data from one biological replicate from a total of three. **I-J. PEAK1 regulates tyrosine phosphorylation of PLCγ1.** MDA-MB-231 cells were transfected with PEAK1 siRNA (I) or a PEAK1 expression vector (J) and then cell lysates were Western blotted as indicated. The histograms indicate Y783-phosphorylated PLCγ1 normalized for total PLCγ1 expression. For all panels, error bars represent the standard error of the mean from n=3 independent experiments. NS indicates $p > 0.05$, * = $p < 0.05$, ** = $p<0.01$, *** = $p<0.001$, ****, $p < 0.0001$, by ratio paired t-test (B, D, I, J), two-way ANOVA with Tukey's multiple comparisons test (F), or unpaired t-test (H).

Figure 3.

Figure 3. Mapping the CAMK2 binding region on PEAK1. A. Schematic structure of PEAK1 and various truncation mutants. WT, wildtype. **B. Association of CAMK2D with specific PEAK1 deletion mutants.** The indicated Flag-tagged PEAK1 deletion mutants were expressed in HEK293T cells. Total cell lysates (TCL) and Flag IPs were subjected to Western blotting as indicated. **C. Schematic representation of various PEAK1 N-terminal deletion mutants.** Deleted regions of PEAK1 are indicated by red crosses. **D-E. Fine structure mapping of N-terminal regions required for the association of PEAK1 with CAMK2D (D) or CAMK2G (E).** Analysis was undertaken as in (B). The asterisk in 3D highlights degradation products present for the Δ261-300 and Δ261-324 mutants. **F-G. PEAK1 binding to CAMK2 is required for CAMK2 activation.** MDA-MB-231 cells were transfected with WT PEAK1 or the Δ301-324 PEAK1 construct. Cell lysates were Western blotted as indicated (F), with the dotted line in F indicating where the image was cropped to remove an irrelevant lane. Raw data for the full image accompanies the manuscript. Relative CAMK2 activation, as determined by T287 phosphorylation, was then quantified (G). Data are expressed relative to vector control which was arbitrarily set at 1. All data in B, D and E are representative of at least 2 independent experiments. In G, error bars represent the standard error of the mean from n=3 independent experiments. NS indicates $p > 0.05$, $* = p < 0.05$ by one-way ANOVA with Tukey's multiple comparisons test.

Figure 4.

Figure 4. Characterization of a conserved CAMK2 interaction motif (CIM) on PEAK1. A. Sequence alignment of the PEAK1 CIM with other CAMK2 binding partners. The critical residues for CAMK2 interaction (R297, L301 and R303 in PEAK1) are highlighted in yellow. The canonical motif numbering (A306 being 0) is shown above the alignment. **B. Representative ITC measurement of CAMK2 kinase domain binding to PEAK1 CIM peptide.** The mean Kd value from n=3 independent experiments is indicated. **C. Comparative modelling of the PEAK1/CAMK2 interaction by AlphaFold.** Surface representation of CAMK2 with PEAK1 amino acids 291-320 shown as ribbon (grey). The orientation is as in Supp Figure 6. CAMK2 is coloured by electrostatic charge. The PEAK1 binding site on CAMK2 is noticeably negative, with key arginine residues of PEAK1 mediating interactions with CAMK2. The N- and C-termini are labelled as N and C, respectively. **D. Surface representation of modelled PEAK1/CAMK2 interaction.** CAMK2 is grey and PEAK1 amino acids 291-320 are purple. **E-F. Focused view of key predicted molecular interactions.** Boxed regions from (D) illustrating predicted formation of salt bridges and electrostatic interactions (dotted lines) by specific PEAK1 residues across the surface of CAMK2. The canonical numbering of the PEAK1 motif is shown in white boxes.

Figure 5

Figure 5. The role of the PEAK1 CIM in CAMK2 activation. A-B. Effect of CIM mutation on intracellular Ca^{2+} levels. Constructs expressing WT PEAK1 or the indicated mutants were stably expressed in MDA-MB-231_EcoR cells via retroviral transduction. PEAK1 expression levels were determined by Western blot (A) and Ca^{2+} levels (B) assayed as in Figure 2G-H. Data in B represent mean Fluo-4 fluorescence ± SEM from one representative biological replicate from a total of two independent experiments. Statistical analysis utilized one way ANOVA with Dunnett's multiple comparisons test with **** indicating $p < 0.0001$. **C-D. Effect of CIM mutation on tyrosine phosphorylation of PLCγ1**. The indicated plasmids were transiently transfected into HEK293 cells and cell lysates Western blotted as indicated (C). PLCγ1 tyrosine phosphorylation was normalized for total expression (D). Data in D are expressed relative to the vector control which was arbitrarily set at 1 and represent the mean ± SEM of 3 independent biological replicates, * indicates p<0.05 by unpaired t-test. **E-G. Effect of CIM mutation on CAMK2 activation.** Plasmids encoding WT PEAK1 and the indicated mutants were transfected into MDA-MB-231 cells. Total cell lysates were Western blotted as indicated (E) and then relative CAMK2 activation was quantified (F-G). Data are expressed relative to the vector (DsR) control which was arbitrarily set at 1. NS indicates $p > 0.05$, ** = $p < 0.01$ and **** = $p < 0.0001$ by one-way ANOVA with Dunnett's multiple comparisons test.

Figure 6.

Figure 6. CAMK2 phosphorylates PEAK1 to regulate the PEAK1 interactome. A-B. RA306 inhibits PEAK1 serine/threonine phosphorylation. MDA-MB-231_EcoR cells stably expressing Flag-tagged PEAK1 were treated with RA306 (1 µM) for 24 h and cell lysates were directly Western blotted or subject to immunoprecipitation with Flag antibody prior to blotting as indicated (A). Relative serine/threonine phosphorylation of PEAK1, normalized for total PEAK1, is shown in the histogram (B). **C. CAMK2 phosphorylates PEAK1 *in vitro*.** Flag-tagged versions of WT and mutant PEAK1 proteins were expressed in HEK293T cells. PEAK1 proteins were immunoprecipitated using anti-Flag antibody and following washing, immediately treated with SDS-PAGE sample buffer (SB) (no incubation) or incubated in kinase reaction buffer supplemented as indicated at 37°C for 30 min. Reactions were terminated with SB. Samples were then Western blotted as indicated. The asterisk indicates a background band present in all CAMK2D-incubated samples including the vector control while the arrowhead highlights phosphorylated PEAK1. **D. Impact of CIM mutations on heterotypic association with PEAK2.** Flag-tagged WT PEAK1 or PEAK1 mutants were expressed in MDA-MB-231 cells and association with endogenous PEAK2 assayed by IP/Western. For panel B, * = p<0.05 by ratio paired t-test.

Figure 7

Figure 7. Role of CAMK2 in PEAK1-regulated biological endpoints in TNBC models. A. PEAK1 overexpression in combination with CAMK2D and CAMK2G knockdown in MDA-MB-231 cells. Cells were transfected with a PEAK1 plasmid in the presence or absence of siRNA-mediated CAMK2D or CAMK2G knockdown. Cell lysates were Western blotted as indicated. **B. Role of CAMK2D/G in PEAK1-promoted MDA-MB-231 cell migration.** Cells validated in (A) were subjected to a transwell migration assay. **C-D**, **Role of CAMK2 activation.** MDA-MB-231 cells were transfected with a PEAK1 plasmid and subject to transwell migration assays in the presence or absence of RA306 (C)**.** Plasmids expressing WT PEAK1 and the Δ301-324 mutant that cannot activate CAMK2 were transiently transfected into MDA-MB-231 cells, and cells were subjected to a transwell migration assay (D). **E. Role of PEAK1/CAMK2 signalling in TNBC xenograft growth.** Control MDA-MB-231_HM cells (Cas9-control) or PEAK1 KO cells (Cas9-sgPEAK1) in combination with RA306 or vehicle control (captisol) were injected into the mammary fat pad. Left panel, representative whole body bioluminescent imaging (BLI) images from each group at different time points as indicated. Right panel, quantification of xenograft growth. Each treatment group exhibited ≥ 9 mice. **F. Role of PEAK1/CAMK2 signalling in TNBC metastasis.** Control MDA-MB-231_HM cells (Cas9-control) or PEAK1 KO cells (Cas9-sgPEAK1) in combination with RA306 or vehicle control (captisol) were injected into the mammary fat pad. At Day 19 post-injection when the primary tumour from each group reached ~ 100 mm^3, they were resected, and tumour growth at secondary sites was measured. Left panel, representative whole body and *ex vivo* lung BLI images from each group at Day 72 post injection. Right panel, luciferase intensities of tumour metastasis to lung were generated from each group. Error bars represents the standard error of the mean from n=3 independent experiments (B-D) or from the individuals of each group (E & F). NS indicates $p > 0.05$, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, **** = $p < 0.0001$ by two-way ANOVA with Tukey's multiple comparisons test (B, C, E), one-way ANOVA with Tukey's multiple comparisons test (D), or one-way ANOVA with Dunnett's multiple comparisons test (F).

Figure 8.

Figure 8. Schematic representation of feed-forward regulation of CAMK2 by PEAK1.
PEAK1 enhances intracellular Ca^{2+} by promoting SFK-mediated tyrosine phosphorylation of
PLCγ1. This leads to classical activation of CAMK2 by Ca^{2+}/CaM and the αD helix rotates out.
The PEAK1 CIM then binds to the CAMK2 substrate binding site and competes with the
regulatory segment, effectively 'locking' CAMK2 in an active conformation. Sustained
phosphorylation of PEAK1 and CAMK2 itself (on T287) then occurs as the PEAK1 CIM
undergoes cycles of dissociation and re-association. However, other mechanisms may
contribute to the observed effects on CAMK2 activation, as indicated on the figure. CIM-
mediated recruitment of PEAK1 to the CAMK2 holoenzyme promotes serine/threonine
phosphorylation of PEAK1 which in turn positively regulates heterotypic association with
PEAK2.