Fusion, fission, and scrambling of the bilaterian genome in Bryozoa

Thomas D. Lewin,[1] Isabel Jiah-Yih Liao,[1] Mu-En Chen,[1] John D. D. Bishop,[2] Peter

W. H. Holland,[3] and Yi-Jyun Luo[1]*

[1]Biodiversity Research Center, Academia Sinica, Taipei, Taiwan

[2]Marine Biological Association, Plymouth, UK

[3]Department of Biology, University of Oxford, Oxford, UK

*Corresponding author: yjluo@gate.sinica.edu.tw

Running title: Genome scrambling in bryozoans

Abstract

Groups of orthologous genes are commonly found together on the same chromosome over
vast evolutionary distances. This extensive physical gene linkage, known as macrosynteny, is
seen between bilaterian phyla as divergent as Chordata, Echinodermata, Mollusca, and
Nemertea. Here, we report a unique pattern of genome evolution in Bryozoa, an understudied
phylum of colonial invertebrates. Using comparative genomics, we reconstruct the
chromosomal evolutionary history of five bryozoans. Multiple ancient chromosome fusions
followed by gene mixing led to the near-complete loss of bilaterian linkage groups in the
ancestor of extant bryozoans. A second wave of rearrangements, including chromosome
fission, then occurred independently in two bryozoan classes, further scrambling bryozoan
genomes. We also discover at least five derived chromosomal fusion events shared between
bryozoans and brachiopods, supporting the traditional but highly debated Lophophorata
hypothesis. Finally, we show that chromosome fusion and fission processes led to the
partitioning of genes from bryozoan Hox clusters onto multiple chromosomes. Our findings
demonstrate that the canonical bilaterian genome structure has been lost across all studied
representatives of an entire phylum; reveal that linkage group fission can occur very frequently
in specific lineages; and provide a powerful source of phylogenetic information.

Introduction

Advances in genome sequencing and assembly have in recent years made it possible to
compare the organization of entire genomes across diverse branches of the animal kingdom.
The emerging picture is one of a common set of metazoan genes, superimposed with
extensive gene duplication, gene loss, and gene novelty in each lineage, plus a surprising
degree of large-scale organizational similarity. Many groups of orthologous genes have been
retained on the same chromosome for hundreds of millions of years and remain together in
diverse and highly divergent lineages (Putnam et al. 2007, 2008; Simakov et al. 2013, 2022;
Schultz et al. 2023; Lin et al. 2024). This conservation of chromosome-scale gene linkages,
known as macrosynteny, is intriguing, as the selective pressures maintaining it and its
significance for genome function are still not fully understood.

The persistence of gene linkages enables the identification of orthologous
chromosomes between distantly related phyla. Furthermore, it permits the reconstruction of
ancestral linkage groups (ALGs)—hypothesized sets of orthologous genes located on the
same chromosome or chromosome arm in ancestral species (Simakov et al. 2022; Mackintosh
et al. 2023a). Lineage-specific ALGs have been identified in several animal groups, including
vertebrates (Sacerdot et al. 2018; Simakov et al. 2020) and Lepidoptera (butterflies and

moths) (Wright et al. 2024). Most strikingly, the discovery of 24 ALGs shared between spiralians and deuterostomes (Simakov et al. 2022; Marlétaz et al. 2023) revealed the genome structure of the most recent bilaterian common ancestor. These highly-conserved bilaterian blocks of orthologous genes are retained in species from phyla as divergent as chordates, echinoderms, molluscs, and annelids (Wang et al. 2017; Simakov et al. 2022; Marlétaz et al. 2023; Martín-Zamora et al. 2023; Lin et al. 2024), suggesting that there are substantial functional constraints to genome organization. There are lineages within phyla that have experienced genomic scrambling, including the breaking up or mixing of genes from different ALGs, such as planarians (Ivankovic et al. 2023), tunicates (Plessy et al. 2024) and clitellate annelids (Schultz et al. 2024; Vargas-Chávez et al. 2024; Lewin et al. 2024a). However, conservation of ALGs seems to be the norm at the phylum level. Despite this general conservation, fluctuations in chromosome number—often leading to ALG rearrangement—are a relatively frequent evolutionary process (White 1954; King 1995; Rice et al. 2015; Román-Palacios et al. 2021). In animals, most of these changes are attributed to inter-chromosomal rearrangements such as translocations and fusions, though they may also arise through whole genome duplication (Orr 1990; Gregory and Mable 2005; Moriyama and Koshiba-Takeuchi 2018; Muffato et al. 2023). Following chromosome fusion, orthologous genes from different ALGs may either be maintained in separate chromosome sections or shuffled and interspersed along the chromosome in a process known as 'fusion-with-mixing' (Simakov et al. 2022). Conversely, chromosomes can undergo fission, splitting into separate parts, but the division of ALGs through this process is rare (Simakov et al. 2022; Wright et al. 2024).

The interplay between the stability of orthologous gene sets and the dynamic nature of inter-chromosomal rearrangements shapes the genomic landscape. These rearrangements, while disrupting gene function and expression (Harewood and Fraser 2014) and altering recombination rates (Yoshida et al. 2023), also play a role in driving adaptation (Coyle and Kroll 2007; Dunham et al. 2002; Cheng et al. 2012) and contributing to macroevolutionary processes like speciation (Rieseberg 2001; de Vos et al. 2020; Berdan et al. 2023; Mackintosh et al. 2023b). This dynamic quality of chromosomal changes has made macrosynteny a powerful tool for phylogenetic studies (Steenwyk and King 2024). In particular, fusion-with-mixing events can be particularly powerful phylogenetic markers as the intermixing of genes from different ALGs is considered rare and irreversible. In light of this, macrosynteny was recently used to investigate early branching metazoan lineages (Schultz et al. 2023) and teleost evolution (Parey et al. 2023).

Even in the age of phylogenomics, when the amino acid sequences of thousands of deduced proteins are aligned and analyzed, there remains a small group of 'Problematica': taxa for which a robust phylogenetic placement remains tantalizingly out of reach (Jenner and Littlewood 2008; Jenner et al. 2009). Many of these taxa are rare or have few species. Yet

one phylum—the Bryozoa—stands out among Problematica as highly abundant, species-rich,
and ecologically important (Gordon and Costello 2016; Lombardi et al. 2020), increasingly as
alien invasive species (Loxton et al. 2017; Weaver et al. 2018; Pratt et al. 2022). Bryozoans
are benthic, suspension-feeding, aquatic invertebrates that are important ecosystem
engineers (Wood et al. 2012; Bock and Gordon 2013). They form colonies of individual zooids,
generally with calcified skeletons (Zhang et al. 2021), and each zooid can be morphologically
differentiated for specialized roles in feeding, brooding, or defense. Despite their unique
biological traits, bryozoans and bryozoan genomes remain relatively understudied (Orr et al.
2021; Liao et al. 2023). Several chromosome-scale assemblies have recently been made
available (Hoencamp et al. 2021; Bishop et al. 2023a, 2023b; Wood et al. 2023; Bishop et al.
2024), but these are without comparative analysis, and beyond an accepted placement within
Spiralia, the phylum's phylogenetic position remains unresolved (Jenner and Littlewood 2008;
Jenner et al. 2009; Khalturin et al. 2022; Liao et al. 2023) (Supplemental Fig. S1).

Here, we present a comparative genomic analysis of five bryozoan genomes, focusing
on genome rearrangements and chromosome evolution. Our study aims to (i) characterize the
inter-chromosomal rearrangements that have occurred with the phylum Bryozoa; (ii)
determine how the 24 bilaterian ALGs are organized in bryozoans; (iii) reconstruct the genome
organization of the last common ancestor of extant bryozoans; and (iv) use macrosynteny to
elucidate the phylogenetic position of Bryozoa.
104

Results

Chromosome-scale genomes of five bryozoans

To reconstruct the evolutionary history of bryozoan chromosomes, we first assembled a
dataset of five chromosome-scale genomes. Four assemblies, *Bugulina stolonifera* (BST)
(Wood et al. 2023), *Cryptosula pallasiana* (CPA) (Bishop et al. 2023b), *Membranipora*
membranacea (MME) (Bishop et al. 2023a), and *Watersipora subatra* (WSU) (Bishop et al.
2024), are marine bryozoans in the class Gymnolaemata, which accounts for approximately
90% of described bryozoan species (Bock and Gordon 2013) (Fig. 1A). The fifth, *Cristatella*
mucedo (CMU) (Hoencamp et al. 2021), is an outgroup belonging to the freshwater class
Phylactolaemata (Taylor and Waeschenbach 2015). Four of the five assemblies were
generated and made openly accessible by the Darwin Tree of Life (DToL) project (The Darwin
Tree of Life Project Consortium et al. 2022), sequenced to chromosome-level using PacBio
HiFi long reads and high-throughput chromosome conformation capture (Hi-C) scaffolding
(Wood et al. 2023; Bishop et al. 2023a, 2023b, 2024)**.**

119
Figure 1. Comparative genomics of five chromosome-scale bryozoan genomes. (*A*)
Relationships between the three classes of bryozoans. Class Phylactolaemata is the most
basal, containing just 86 species, all of which inhabit freshwater environments (Bock and
Gordon 2013). Stenolaemata comprises 543 species, all of which are marine.
Gymnolaemata is the most species-rich class, comprising 5,240 species. (*B*) Genome size
and gene model count of bryozoan genome assemblies used in this study. Dashed lines
mark mean values. The clade containing *W. subatra* and *Cry. pallasiana* has larger
genomes. (*C*) Tree of bryozoan species constructed with chromosome-scale genomes with
the scallop *Pecten maximus* as the outgroup. The tree was constructed using the maximum
likelihood method (LG+F+R6 model) with 1,000 bootstrap replicates in IQ-TREE, utilizing
994 orthologous genes. The scale bar represents the number of substitutions per site. (*D*)
Phylogenetic relationships among animal phyla, with dashed lines indicating areas of
uncertainty in their placement. Phylogeny based on (Telford et al. 2015) and (Liao et al.
2023). (*E*) Principal component analysis (PCA) conducted using orthologous gene group
counts as determined by OrthoFinder. Points are colored by their phylum, as indicated in
panel *D*. PCs 1 and 2 explain 10.0% and 9.4% of the variance in the dataset, respectively.
Variance ratios for the top 30 principal components are available as Supplemental Fig. S2.
Species abbreviations: Ava, *Adineta vaga*; Aqu, *Amphimedon queenslandica*; Aru, *Asterias*
rubens; Bfl, *Branchiostoma floridae*; Bmi, *Bolinopsis microptera*; Cel, *Caenorhabditis*
elegans; Cin, *Ciona intestinalis*; Cte, *Capitella teleta*; Dme, *Drosophila melanogaster*; Dpu,
Daphnia pulex; Dre, *Danio rerio*; Gfa, *Galaxea fascicularis*; Hmi, *Hofstenia miamia*; Hro,
Helobdella robusta; Hsa, *Homo sapiens*, Lan, *Lingula anatina*; Llo, *Lineus longissimus*; Lva,
Lytechinus variegatus; Nve, *Nematostella vectensis*; Obi, *Octopus bimaculoides*; Ofu,
Owenia fusiformis; Pau, *Phoronis australis*; Pma, *Pecten maximus*; Pvu, *Patella vulgata*;
Sko, *Saccoglossus kowalevskii*; Sma, *Schistosoma mansoni*; Sme, *Schmidtea*
mediterranea; Sro, *Symsagittifera roscoffensis*; Tad, *Trichoplax adhaerens*; Tca, *Tribolium*
castaneum; Xtr, *Xenopus tropicalis*.

We first performed comprehensive gene prediction using the BRAKER3 pipeline,
utilizing transcriptomic data as supporting evidence (Fig. 1B; Supplemental Tables S1–S3).
Benchmarking Using Single-Copy Orthologs (BUSCO) (Simão et al. 2015; Manni et al. 2021)
reveals a high degree of completeness in gene predictions while also indicating some losses
in bryozoans, as evidenced by completeness scores of 82.7% to 88.6% from the Metazoa
database (odb10) (Supplemental Table S1). Given accurate phylogenetic relationships are
key to inferring chromosome rearrangement events, we built a maximum likelihood
phylogenetic tree of the five bryozoan species using gene models obtained from these
chromosome-scale genomes (Fig. 1C); this tree is consistent with recent phylogenetic
analyses using mitochondrial and ribosomal data (Orr et al. 2022). Principal component
analysis of orthologous gene counts finds bryozoans to have a gene set broadly similar to that
of other lophotrochozoans (Fig. 1D,E; Supplemental Tables S4,S5).

Massive chromosome rearrangements within Bryozoa

Haploid chromosome number in our bryozoan dataset ranges from eight in *Cri. mucedo* to 12
in *Cry. pallasiana*. To investigate how the chromosomes of each species are related and
understand whether large genome restructuring events have occurred within the phylum
Bryozoa, we used OrthoFinder (Emms and Kelly 2019) to identify 4,242 single-copy orthologs
and mapped the position of these orthologs within each genome. We then constructed synteny
maps which we displayed as ribbon plots (Fig. 2) and Oxford dot plots (Supplemental Fig. S3).
Throughout our analysis of chromosome rearrangements, we apply the algebraic symbol
conventions established by Simakov et al., wherever applicable (Simakov et al. 2022).
Specifically, we use triple bar for one-to-one relationships, also known as syntenic
equivalences (≡); tensor for fusion with mixing (⊗); dot product for end-to-end fusion (●); down
arrow for centric insertion (⬇); and addition for fission (+) (Fig. 2A). However, although these
descriptions are effective for taxa with relatively conserved genomes, we found that the highly
derived nature of bryozoan genomes resulted in many chromosomes not fitting clearly into
these predefined categories. Furthermore, the notation may become excessively complex and
confusing when addressing chromosomes that have undergone numerous ALG fusions and
have intricate evolutionary histories. We therefore also utilize narrative descriptions, especially
for highly derived chromosomes.

Figure 2. Chromosome rearrangement in gymnolaemate and phylactolaemate bryozoans.
(*A*) Schematic representations of possible inter-chromosomal rearrangements. White bars
represent chromosomes; colored lines connect orthologous genes; the same color line
represents genes in the same ancestral linkage group. (*B*) Chromosome-scale gene linkage
between five bryozoan species revealed by ribbon plots. Horizontal bars represent
chromosomes. Vertical lines connect the genomic position of orthologous genes in each
genome. Chromosomes connected by many orthologs are inferred to be homologous (i.e.,
orthologous chromosomes). In gymnolaemate species, 11 ALGs *a* to *k* are highly conserved,
though the chromosome containing ALG *a* underwent a fission event to form chromosomes
CPA10 and CPA12 in *Cry. pallasiana*. These ALGs are not conserved in the
phylactolaemate species *Cri. mucedo.* Links between chromosomes sharing five or fewer
genes between chromosomes are removed for clarity but can be seen in the Oxford dot
plots. (*C*) Oxford dot plot revealing chromosome-scale gene linkage between the two most
closely related species in our dataset, *Cry. pallasiana* and *W. subatra*. Each axis represents
the entire length of the genome of one species. Gray lines separate chromosomes. Each
point represents a pair of orthologs placed by their ordinal position in each genome. Strings
of orthologs perpendicular to the main diagonal axis reveal clear intra-chromosomal
inversions (dark purple). (*D*) Single-chromosome Oxford dot plots for *Cry. pallasiana* and *W.
subatra* chromosomes with clear intra-chromosomal inversions (dark purple). (*E*) Ribbon plot
for *Cry. pallasiana* and *W. subatra* highlighting intra-chromosomal inversions (dark purple)
(*F*) Microsynteny mixing score versus divergence time for gymnolaemate bryozoan
chromosomes. Each point represents a chromosome. Microsynteny mixing score measures
how genes are shuffled within a chromosome by intra-chromosomal inversions (see

Materials and Methods): a higher score indicates more mixing. Mixing score increases with phylogenetic distance. Abbreviation: MYA, million years ago.

We used the chromosomal position of orthologous genes to identify orthologous chromosomes among the four gymnolaemate species. There is complete syntenic equivalence among three gymnolaemate bryozoans (*B. stolonifera*, *M. membranacea,* and *W. subatra*)*,* which all have a haploid chromosome number of 11, suggesting there have been no instances of chromosome fusion or fission (Fig. 2B). Based on gene position in these species, we assigned 11 gymnolaemate ALGs, labeled ALGs *a* to *k* (Fig. 2B). The final gymnolaemate species, *Cry. pallasiana*, has undergone one fission event to diverge from this state, with the chromosome containing gymnolaemate ALG *a* (*M. membranacea* chr1 ≡ *B. stolonifera* chr1 ≡ *W. subatra* chr1) splitting to form two smaller chromosomes (*Cry. pallasiana* chr10 + chr12) (Fig. 2B). Given the reported rarity of chromosome fission events in animal evolution, we verified that this is not an assembly error by generating a Hi-C contact map, which confirmed that the fission is genuine (Supplemental Fig. S4). This aside, chromosome structure is highly conserved between these species. In contrast, a striking lack of conservation is observed when the phylactolaemate bryozoan *Cri. mucedo* is compared. None of the 11 gymnolaemate ALGs are conserved, and most are completely dispersed and scrambled across the entire *Cri. mucedo* genome (Fig. 2B). For instance, we identified four gymnolaemate ALGs on seven of eight *Cri. mucedo* chromosomes (gymnolaemate ALGs *a*, *b*, *c*, and *g*) (Supplemental Fig. S5). This represents an almost complete loss of chromosome relationships between the two groups. Overall, genome structure is highly conserved within gymnolaemate bryozoans but large inter-chromosomal rearrangements have resulted in highly divergent genome structures between members of Gymnolaemata and Phylactolaemata.

Macrosynteny studies over long evolutionary distances often overlook the precise order of genes, known as microsynteny (Renwick 1971; Passarge et al. 1999). This is because intra-chromosomal rearrangements like inversions rapidly alter gene order, leading to weak microsyntenic signals which are difficult to trace between distantly related species. Within our dataset, there is minimal conservation of gene order between bryozoans with the exception of the two most closely related species, *W. subatra* and *Cry. pallasiana* (Supplemental Fig. S3), which are estimated to have diverged 60–90 million years ago (Orr et al. 2022). Conservation of microsynteny between these species permitted us to investigate the process by which intra-chromosomal rearrangements occur. By generating Oxford dot plots for whole genomes (Fig. 2C) and individual chromosomes (Fig. 2D), we identified six clear cases of large intra-chromosomal inversions between these two species (e.g., *W. subatra* chr1 vs. *Cry. pallasiana* chr12). We noticed many inversions with a complete lack of intermixing, suggesting that these inversions involved entire chromosome arms (e.g., *W. subatra* chr8 vs. *Cry. pallasiana* chr3)

(Fig. 2D,E). We developed a microsynteny mixing score (see Methods) to quantify the extent

of intra-chromosomal gene shuffling. Between the closely related species *W. subatra* and *Cry.*

pallasiana, mixing score varies greatly between chromosomes, suggesting genes are not

shuffled at equal rates on different chromosomes (Fig. 2F). In addition, the mixing score

increases with estimated divergence time in our dataset (Fig. 2F; Supplemental Tables

S6,S7). Overall, the mixing of genes within a chromosome increases with the divergence time

between species, and occurs at least in part due to large inversions.

Near-complete disruption of bilaterian ancestral linkage groups in Bryozoa

To facilitate synteny analysis using bilaterian ancestral linkage groups, we created a new

pipeline, SyntenyFinder (see Methods). SyntenyFinder integrates OrthoFinder and RIdeogram

and is capable of conducting large-scale comparisons. We first validated our synteny analysis

methods using bilaterians with established chromosome relationships. We applied the

bilaterian ALG chromosome assignment to the amphioxus *Branchiostoma floridae* and found

it to be consistent with previous work (Simakov et al. 2022), identifying bilaterian ALG fusions

on chr1 (A1⊗A2), chr2 (C1⊗J2), chr3 (C2●Q), and chr4 (O1●I) . This result confirms the

accuracy of our ortholog identification and macrosynteny methodologies.

We next investigated how bryozoan chromosomes relate to the conserved bilaterian

genome organization. Using only the bryozoan data, it remains uncertain whether the

Gymnolaemata, Phylactolaemata, or neither is similar to the ancestral state of bilaterians. To

resolve this, in addition to the amphioxus *B. floridae* (Simakov et al. 2020), we compared the

macrosynteny of bryozoans with two other bilaterians with conserved genome organization:

the nemertean *Lineus longissimus* (Kwiatkowski et al. 2021) and the mollusc *Pecten maximus*

(Kenny et al. 2020). Both *L. longissimus* and *P. maximus* belong to the Lophotrochozoa, the

same major protostome clade as bryozoans, while the deuterostome *B. floridae* serves as an

outgroup. We generated a dataset of 995 single-copy orthologs and associated each ortholog

with one of the 24 bilaterian ALGs described by Simakov et al. (Simakov et al. 2022)

(Supplemental Data S1). Synteny maps show that, despite their divergence at the base of the

Bilateria, the genome organizations of *B. floridae*, *P. maximus*, and *L. longissimus* exhibit high

levels of similarity, having largely preserved the ancestral bilaterian ALGs (Fig. 3;

Supplemental Fig. S6; Supplemental Table S8). Our analysis revealed that *P. maximus* and

L. longissimus share four fusion events that have also been recorded in annelids and

brachiopods (J2⊗L, O1⊗R, O2⊗K, and H⊗Q) (Simakov et al. 2022; Martín-Zamora et al.

2023), suggesting that they are ancestral to lophotrochozoans. In addition, we detected

several lineage-specific fusion events. Specifically, we identified a fusion on chr2 (B2⊗M) in

P. maximus; and a fusion on chr2 (C1⊗G) in *L. longissimus*. Overall, genome structures in

the deuterostome *B. floridae* and lophotrochozoans *P. maximus* and *L. longissimus* retain
most bilaterian ALGs alone on single chromosomes while each have a few lineage-specific
fusion events. These changes are modest, considering that these species have been diverging
for over half a billion years since the last common ancestor of bilaterians.

Figure 3. Extensive fusion-with-mixing of bilaterian ancestral linkage groups in bryozoans.
Chromosome-scale gene linkage between five bryozoan species and three outgroup
bilaterians: a nemertean, a mollusc, and a chordate. Horizontal bars represent
chromosomes. Vertical lines connect the genomic position of orthologous genes in each
genome. Lines are colored by the bilaterian ALGs defined by (Simakov et al. 2022). These
bilaterian ALGs are highly conserved in species ranging from the amphioxus *Branchiostoma
floridae* to the scallop *Pecten maximus* and the nemertean *Lineus longissimus*, as indicated
by asterisks, but are extensively scrambled in bryozoans.

Like *B. floridae*, *P. maximus*, and *L. longissimus*, bryozoan genomes contain genes
from all 24 bilaterian ALGs. However, in contrast to their highly conserved state in other
lophotrochozoans and deuterostomes, these bilaterian ALGs have been extensively combined
and mixed in the bryozoan lineage (Fig. 3; Supplemental Fig. S6; Supplemental Table S9).
We found that 18 out of the 24 bilaterian ALGs (75%) have been combined and mixed across
four gymnolaemate chromosomes, indicating that gymnolaemate bryozoans underwent
substantial irreversible fusion-with-mixing events followed by fragmentation of a single,

ancestrally fused chromosome into four distinct ones. We also detect other bryozoan-specific
fusions: ALGs A1 and C2 initially fused (A1⊗C2) and then divided into two chromosomes
(e.g., *M. membranacea* chr6 and chr8); ALG P joined with the lophotrochozoan fusion K⊗O2
(P⊗K⊗O2) and later separated into two chromosomes (e.g., *M. membranacea* chr9 and
chr10); ALGs F and N fused (F⊗N) and subsequently split into two chromosomes (e.g., *M.*
membranacea chr5 and chr11). One chromosome contains only ALG M genes (e.g., *M.*
membranacea chr7), although various genes from ALG M were translocated to other
chromosomes. The identification of the same chromosome structure across *M. membranacea*,
B. stolonifera, *W. subatra*, and *Cry. pallasiana* confirms that the observed genome scrambling
is not due to assembly errors but rather represents genuine evolutionary divergence. We also
found that the *Cri. mucedo* genome arrangement, while unique from other bryozoans, similarly
does not preserve the bilaterian ALGs on separate chromosomes. Massive rearrangements
have independently taken place within the lineage leading to *Cri. mucedo*, resulting in the
dispersion of bilaterian ALGs throughout its genome (Fig. 3). Overall, the genome
organizations of the two classes of bryozoans studied here are distinct from each other and
highly divergent from the bilaterian ancestral state.

The presence of several ALGs on two or four chromosomes in gymnolaemate
bryozoans led us to question whether the splitting of ALGs was due to whole genome
duplication (WGD) or chromosome duplications rather than chromosome fission. We tested
this using two complementary methods, each using *Metaphire vulgaris*, an annelid with a
known WGD (Jin et al. 2020), as a positive control. We first tested for whole genome
duplication using synonymous substitutions per site (K_s) distributions for paralogs within each
genome. The large peak in the distribution for *M. vulgaris* is caused by its WGD, and the
absence of similar large peaks within the K_s distributions of gymnolaemate bryozoans does
not support the hypothesis that a recent WGD occurred in this group (Supplemental Fig. S7).
There may be very shallow peaks at higher K_s values (K_s = 3 to 4) in the gymnolaemate
bryozoans which could indicate ancient WGD in this lineage, but the signal is very weak
(Supplemental Fig. S7). Given this uncertainty, we then used MCScanX (Wang et al. 2012) to
identify and align collinear blocks of genes within each genome. The presence of duplicated
gene blocks on chromosomes with genes from the same ALG would suggest that they were
formed by duplication rather than fission. Accordingly, strong pairwise relationships were
identified between duplicated chromosomes in *M. vulgaris.* However, in gymnolaemate
bryozoans, this analysis revealed zero or few such homologous blocks of genes, which
provides no evidence that they are derived from WGD or whole chromosome duplication
events (Supplemental Fig. S8). Finally, we used a Duplicate Genes Classifier (Wang et al.
2012) to determine the origin of duplicates found in the genomes. While 23.5% of *M. vulgaris*
genes are derived from WGD/segmental duplication events, only 0.4 to 3.6% of genes in the

four gymnolaemate species fall into this category (Supplemental Table S10). This again lends

no support to the idea of a WGD in gymnolaemate bryozoans. In the phylactolaemate *Cri.*

mucedo, we identified several sections of homology between *Cri. mucedo* chromosomes (1

with 2; 1 with 5; 4 with 6) (Supplemental Fig. S8) and several very small very small K_s peaks

(Supplemental Fig. S7), suggesting possible segmental duplications in this lineage. Overall,

we failed to find evidence that the presence of ALGs on two or four chromosomes in

gymnolaemate bryozoans is the product of WGD or large-scale chromosomal duplications,

though this cannot be conclusively ruled out.

The last common ancestor of extant bryozoans had a highly rearranged genome

Since the Gymnolaemata and Phylactolaemata have such distinct genome arrangements, we

sought to uncover the genomic structure of their ancestor, the last common ancestor of extant

bryozoan lineages. Did it maintain the general bilaterian organization or had rearrangements

already occurred? We leveraged the conserved gene linkages found in modern genomes to

reconstruct the ancestral state by searching for fusion and fission events that are shared by

both gymnolaemate and phylactolaemate species. Since these two classes diverged at the

base of the extant Bryozoa, inter-chromosomal rearrangements found in both sets of species

can be inferred to have been present in their ancestor. This implicitly assumes that they did

not arise multiple times independently, but the complex nature of the shared rearrangements

makes that scenario highly improbable. This analysis revealed the likely presence of six

linkage groups in the ancestor of living bryozoans, suggesting an ancestral karyotype of six

chromosomes (designated here as α, β, γ, δ, ε, and ζ) (Fig. 4A,B). Notably, bryozoan ALGs α

and β comprise 10 and eight bilaterian ALGs, respectively, indicating numerous events of

sequential fusion-with-mixing (Fig. 4C; Supplemental Fig. S9; Supplemental Table S11). Thus,

chromosomal rearrangements had fused and shuffled bilaterian ALGs extensively, even

before the extant clades of bryozoans diverged.

Figure 4. Reconstruction of ancestral genome architecture of extant bryozoans. (*A*)
Chromosome-scale gene linkage between *L. longissimus*, *M. membranacea,* and *Cri.
mucedo*. Links between orthologs are colored by their 6 bryozoan ALGs. Bryozoan ALGs are
inferred from sets of genes that are co-located on the same chromosome (region) in both
gymnolaemate and phylactolaemate bryozoans. *L. longissimus* is used as an outgroup
because it largely preserves the 24 bryozoan ALGs on separate chromosomes but possess
the four fusion events that are shared by lophotrochozoans (J2⊗L, O1⊗R, O2⊗K, and
H⊗Q). (*B*) Composition of bryozoan ALGs by bilaterian ALGs, alongside the reconstructed
bryozoan ancestral state (six chromosomes: α, β, γ, δ, ε, and ζ). Triangles indicate the
inclusion of several genes from the bilaterian ALG but the majority of genes from that ALG
are elsewhere. (*C*) Reconstruction of the genome rearrangements that formed the bryozoan
ancestral state. The top row shows bilaterian ALGs, and the bottom row shows bryozoan
ALGs. (*D*) Reconstruction of the genome rearrangements that led from the bryozoan
ancestral state to the extant gymnolaemate genomes. Solid black arrows indicate fusion-
with-mixing. Solid gray arrows indicate dispersal around the genome. Dashed arrows
indicate chromosome fission. (*E*) Reconstruction of the genome rearrangements that led
from the bryozoan ancestral state to the extant phylactolaemate genomes.

Based on this ancestral state of extant bryozoans, we reconstructed the events leading
to the observed genome structures in contemporary species (Fig. 4D). In the gymnolaemates,
the chromosomes containing ALGs α and β underwent fusion-with-mixing (α⊗β) to create one
enormous chromosome containing 18 bilaterian ALGs. This chromosome then underwent
sequential fission events to form four daughter chromosomes containing these 18 bilaterian
ALGs (e.g., *M. membranacea* chr1 to chr4). Meanwhile, bryozoan ALGs δ, ε, and ζ each

underwent a single fission event to form two separate chromosomes. ALG γ was maintained alone on one chromosome (chr7), but genes from this ALG also dispersed to all other chromosomes (Fig. 4D). The *Cri. mucedo* genome exhibits a structure more derived from the ancestral state, characterized by numerous fission and fusion events with limited mixing, resulting in chromosomes consisting of distinct segments derived from different bryozoan ALGs (Fig. 4E). Overall, the bryozoan common ancestor had a highly derived genome, and lineage-specific fusion and fission events have further modified the genomes of both gymnolaemate and phylactolaemate bryozoans.

Shared derived chromosomal rearrangements between bryozoans and brachiopods

Recent studies have proposed the application of macrosynteny analysis to phylogeny inference, arguing that the infrequency and irreversibility of fusion-with-mixing allows these events to serve as phylogenetic markers (Schultz et al. 2023; Parey et al. 2023; Steenwyk and King 2024). In light of the contentious debate surrounding the phylogenetic position of bryozoans, we sought to determine if macrosynteny could offer insights in this regard. Two contrasting hypotheses exist: (a) the Lophophorata hypothesis, which proposes a close relationship to phoronids and brachiopods (Nesnidal et al. 2013; Luo et al. 2017; Marlétaz et al. 2019; Laumer et al. 2019), and (b) the Polyzoa hypothesis, which proposes closer affiliation to Entoprocta and Cycliophora (Hejnol et al. 2009; Kocot et al. 2017; Khalturin et al. 2022) (Supplemental Fig. S1). Although no chromosome-scale genomes are available for Cycliophora, Entoprocta, or Phoronida species, we recently sequenced a chromosome-level genome for the brachiopod *Lingula anatina* (Lewin et al. 2024b). We first validated the quality of this assembly using genome statistics and found it to be highly contiguous (Supplemental Tables S12,S13), then verified that the chromosome number was supported by cytological karyotype studies (n = 10) (Nishizawa et al. 2010). Next, we checked the assembly by comparing chromosome structure to two species with highly conserved genomes: *P. maximus* and *L. longissimus* (Supplemental Fig. S10). The recovery of many ancestral bilaterian ALGs on singular chromosomes further supports that the genome is correctly assembled (Supplemental Fig. S10).

We then conducted a macrosynteny comparison between the bryozoan *M. membranacea*, representative of Gymnolaemata, and the brachiopod *L. anatina* to test for the existence of shared derived rare genomic changes in the form of chromosome fusion-with-mixing events. We identified nine fusion-with-mixing events shared between *L. anatina* and *M. membranacea* (Fig. 5A,B). Four of these events (J2⊗L, O1⊗R, O2⊗K, and H⊗Q) are also shared with annelids, nemerteans, and molluscs (Fig. 3) (Simakov et al. 2022; Martín-Zamora

et al. 2023). The remaining five fusion-with-mixing events are exclusive to brachiopods and bryozoans, making them shared derived characteristics unique to these groups (Fig. 5C, marked by ⊗ occurring outside of parentheses). This suggests that bryozoans and brachiopods are closely related, having inherited these changes from a recent common ancestor, supporting the Lophophorata hypothesis (Fig. 5D, scenario Type A). The changes could also be attributed to convergent evolution (Fig. 5E,F, Type B) or a mixture of inheritance from a common ancestor and convergent evolution (Fig. 5G, Type C), but inheritance from a recent common ancestor is the most parsimonious explanation, requiring the fewest fusion-with-mixing events.

Figure 5. Shared derived chromosome features between bryozoans and brachiopods support the Lophophorata hypothesis. (*A*) Chromosome-scale gene linkage between the bryozoan *M. membranacea* and the brachiopod *L. anatina*. Horizontal bars represent chromosomes. Vertical lines connect the genomic position of orthologous genes in each genome. Lines are colored by bilaterian ALGs. (*B*) Oxford dot plot showing shared ALG fusion events between *L. anatina* and *M. membranacea*. Each point represents a pair of orthologs placed by their ordinal position in each genome, colored by their bilaterian ALG. Black boxes highlight shared fusion-with-mixing events, where two ALGs appear together on the same chromosome in both species. (*C*) There are nine shared fusion-with-mixing events between *L. anatina* and *M. membranacea*. Several spiralian phyla share four of these events ('lophotrochozoan ancestral fusions'), marked by ⊗ within parentheses, but their timing is

uncertain. These events may, therefore, not be useful phylogenetic markers for the position
of bryozoans. Confirmed derived fusion-with-mixing events shared between bryozoans and
brachiopods but not present in molluscs and annelids are marked by ⊗ occurring outside of
parentheses. (*D–G*) Visual representation of four possible evolutionary scenarios that could
give rise to the five derived fusion-with-mixing events shared between bryozoans and
brachiopods (Types A, B, and C). An open circle marks the base of the Lophotrochozoa.
Black circles mark chromosome fusion-with-mixing events. (*D*) Scenario Type A:
Brachiopods, phoronids, and bryozoans are closely related, forming the Lophophorata. The
five fusion-with-mixing events shared between bryozoans and *L. anatina* occurred in the
ancestor of the Lophophorata. Total number of fusions required: 5. (*E*) Scenario Type B-1:
Bryozoans, Cycliophora, and Entoprocta are closely related, forming the Polyzoa. Bryozoans
are distantly related to brachiopods. The five fusion-with-mixing events shared between
bryozoans and *L. anatina* occurred independently in bryozoans and brachiopods. Total
number of fusions required: 10. (*F*) Scenario Type B-2: Brachiopods, phoronids, and
bryozoans are closely related, forming the Lophophorata. The five fusion-with-mixing events
shared between bryozoans and *L. anatina* occurred independently in bryozoans and
brachiopods. Total number of fusions required: 10. (*G*) Scenario Type C: Brachiopods,
phoronids, and bryozoans are closely related, forming the Lophophorata. Some of the five
fusion-with-mixing events shared between bryozoans and *L. anatina* occurred in the
ancestor of Lophophorata, while some occurred independently in bryozoans and
brachiopods. Total number of fusions required: 6–9.
465

Under the assumption that fusion-with-mixing events are irreversible, the Type A
scenario would be contradicted if any bryozoan, phoronid, or brachiopod were discovered
without the described fusion-with-mixing events. To test this, we examined whether the same
fusion events are present in *Cri. mucedo*, the most divergent bryozoan in our dataset. Despite
Cri. mucedo having a highly divergent chromosome organization, all five derived fusion-with-
mixing events shared by *M. membranacea* and *L. anatina* also occur together on the same
chromosomes in *Cri. mucedo* (Supplemental Fig. S11). Altogether, we observed nine shared
fusion chromosome events between bryozoans and brachiopods, with at least five being
derived fusion-with-mixing events and thus considered potential phylogenetic markers. These
events are notably absent in molluscs, annelids, and nemerteans, which we interpret as
indicating a close relationship between bryozoans and brachiopods, supporting the
Lophophorata hypothesis. In the future, sequencing a phoronid genome to chromosome level
will be a key test of this hypothesis, revealing whether the same fusions are found in this
phylum.
480

481 **Genome structure and Hox gene evolution**

482 In bilaterians, Hox genes are essential for body patterning and cell type specification (Hubert
483 and Wellik 2023), a role facilitated by their organization into tight clusters (Akam 1989; Graham

et al. 1989; Duboule and Dollé 1989). We explored whether extensive genome scrambling has impacted Hox cluster organization in bryozoans. We annotated the Hox genes of the five bryozoan species (Supplemental Fig. S12; Supplemental Data S2) and identified extensive gene loss in the bryozoan Hox gene set, which lacks *Hox1*, *Hox3*, *Scr*, *Antp*, *Lox2*, and *Post1* in all five species. Strikingly, the remaining Hox genes are distributed across two separate chromosomes in every assembly (Fig. 6A). In bilaterians such as *P. maximus* and *L. longissimus*, the Hox cluster is associated with the bilaterian ALG B2. This ALG is fragmented across four chromosomes (five in *Cry. pallasiana*) in each bryozoan genome, as a result of successive fusion and fission events (Fig. 6B). The main Hox cluster is located on orthologous chromosomes containing ALG B2 genes in each bryozoan genome assembly, while Hox genes separated from the cluster are also on a different orthologous chromosome that contains ALG B2 genes (Fig. 6C). This evidence suggests that the presence of Hox genes across two chromosomes in bryozoans is a unique case of partial cluster tandem duplication (*Hox4*, *Lox5*, *Lox4*, and *Post2* genes) followed by separation due to chromosome fission (Fig. 6D).

Figure 6. Evolution of the bryozoan Hox cluster. (*A*) Left: Representation of the main Hox cluster of bryozoans and two outgroups, the nemertean *L. longissimus* and the mollusc *P.*

maximus. Colored rectangles represent Hox genes. Horizontal black lines represent chromosomes. Right: Separate Hox genes located in different chromosomes. (*B*) Positions of ALG B2 genes (green) in bryozoan genomes. (*C*) Positions of Hox genes (purple) in bryozoan genomes. (*D*) Representation of the evolutionary events that resulted in observed bryozoan Hox gene arrangements. In the bryozoan ancestral lineage, chromosomes containing bilaterian ALGs A1, B1, B2 (including the Hox cluster), B3, C1, D, E, and G underwent sequential fusion events to form bryozoan ALG β; the Hox cluster underwent a partial duplication, copying genes *Hox4*, *Lox5*, *Lox4*, and *Post2*; and the duplicated *Lox4* gene was subsequently lost. In the lineage leading to *Cri. mucedo* (Phylactolaemata), the main cluster *Post2* was lost, and the chromosome containing ALG β underwent fission, leaving the main cluster on chr2 and duplicated genes on chr1. In the lineage leading to *M. membranacea* (Gymnolaemata), the duplicated *Lox5* and *Post2* were lost, and the chromosome containing ALG β underwent fission, leaving the main cluster on chr1 and duplicated genes on chr2.

Given the critical dependence of Hox gene regulation on genomic organization, we hypothesized that physical separation might alter gene expression. To investigate this, we analyzed the expression of bryozoan Hox paralogs located on different chromosomes using published RNA-sequencing data (Supplemental Fig. S13; Supplemental Table S14). In *Cri. mucedo*, both the *Hox4b* and *Lox5b* genes separated from the main cluster are expressed more highly than their main cluster counterparts, *Hox4a* and *Lox5a* (paired *t*-tests $p = 0.0009$ and $p = 0.0001$, respectively) (Supplemental Fig. S13). Therefore, the physical separation of Hox cluster genes by genome rearrangements is associated with differences in expression levels.

Discussion

Our study reveals dramatic rearrangements to bryozoan chromosomes that have extensively fragmented the 24 ALGs inherited by all bilaterians. This presents a stark contrast to reports of ALG conservation from other spiralians (Wang et al. 2017; Martín-Zamora et al. 2023), deuterostomes (Lin et al. 2024), and even non-bilaterians like cnidarians and ctenophores (Simakov et al. 2022; Schultz et al. 2023). We distinguish two distinct periods in which the bryozoan genome was drastically remodelled: the first generating the derived bryozoan ancestral state of just six ALGs from the 24 bilaterian ALGs, largely by chromosome fusion events, and the second independently modifying this state further in different bryozoan classes, including ALG fission events.

Three aspects of bryozoan genomes stand out as particularly remarkable. Firstly, the magnitude of chromosome fusions is notable, with 18 out of 24 (75%) bilaterian ALGs on the same chromosomes. Secondly, the occurrence of these massive fusion events across a phylum is exceptionally rare. Unlike dipterans and clitellate annelids, where such extensive

fusion events are specific to certain lineages (Zdobnov et al. 2005; Simakov et al. 2022; Schultz et al. 2024; Lewin et al. 2024a; Vargas-Chávez et al. 2024), our data indicate that these rearrangements are present across the entire extant bryozoan phylum. This suggests that massive rearrangements had already occurred by the time of the last common ancestor of living lineages. However, we note that much of bryozoan diversity is represented only in the fossil record. Six of the seven orders of Stenolaemata are long extinct and less than 40% of bryozoan genera have living representatives (Taylor and Waeschenbach 2015), while approximately 70% of described bryozoan species are fossil (Gordon and Costello 2016). Thirdly, many ALGs in bryozoans have been separated by ALG fissions. Major studies on animal chromosome evolution suggested that a general rule in metazoans is that ALGs are combined but not broken up (Simakov et al. 2022; Wright et al. 2024). In contrast, we identified numerous fissions occurring in the common ancestor of the four gymnolaemate species around 350 million years ago (Orr et al. 2022), with subsequent, lineage-specific fission in *Cry. pallasiana* less than 90 million years ago (Orr et al. 2022). This indicates a prolonged period during which fissions were evolutionarily tolerated, challenging the notion that they occur in brief, time-limited periods and are followed by evolutionary stasis (Simakov et al. 2022).

Why have bryozoans specifically undergone radical ALG fissions while many animal phyla retained the conserved ancestral state? Although our study doesn't offer a definitive answer, we can apply Mayr's classical framework (Mayr 1961) to outline the elements of a possible explanation. Firstly, we must identify the mechanism that generated these rearrangements (the proximate cause). Secondly, we must understand why these rearrangements conferred a fitness advantage and were thus favored by selection (the ultimate cause). The high level of rearrangement in bryozoans could be due to either (i) an increase in the rate of rearrangement events (the proximate cause), (ii) an increase in the rate of fixation of rearrangement events (the ultimate cause), or both. Loss of DNA repair machinery (Vargas-Chávez et al. 2024), transposable element invasions (Ahola et al. 2014; Höök et al. 2023), or environmental mutagenic agents (Schultz et al. 2024) have been proposed as potential proximate causes other systems, though additional studies are needed to test whether these were contributors in bryozoans.

The ultimate cause of large-scale genome restructuring events is particularly intriguing from an evolutionary standpoint. One possibility is that rearrangements are selectively neutral or weakly deleterious but are fixed at an increased rate due to genetic drift (Wright 1941; Mackintosh et al. 2023c). Notably, bryozoans are colonial and have an asexual stage in which they reproduce by budding (Schwaha 2020), and genetic drift and mutational load are elevated in asexual populations (Haag and Roze 2007). Alternatively, rearrangements may be adaptive and favored by positive selection, as has been suggested for loss of genome structure during the marine–freshwater transition in clitellate annelids (Sun et al. 2021; Schultz et al. 2024;

Vargas-Chávez et al. 2024; Lewin et al. 2024a). The transition to obligate colonialism is a

significant evolutionary innovation in bryozoans (Schwaha et al. 2020) that correlates with the

timing of genome restructuring, but further work is required to understand whether this, or

other bryozoan adaptations such as biomineralization (Taylor et al. 2015), were facilitated by

chromosomal rearrangement events.

A third possibility is that increased rate of fixation of rearrangements results from the

relaxation in bryozoans of strong evolutionary pressure that has been maintained in other

lineages. The role of conserved macrosyntenic gene linkages in genome functioning remains

largely unknown, but their high-fidelity maintenance in diverse lineages for over half a billion

years (Simakov et al. 2022) implies at least some importance. One idea is that this is due to

long-range interactions between distant elements on the same chromosome, which means

that gene linkages are significant to gene expression (Tolhuis et al. 2002; Miele and Dekker

2008; Dean 2011). A change in the relative importance of mechanisms of gene regulation may

therefore reduce the emphasis on long-range interactions and render conserved genome

organization obsolete. Our work highlights the need for studies relating gene regulation to

genome organization, which may be key to understanding the ultimate causes of the evolution

of highly derived genomic architectures.

Having elucidated the organization of bryozoan genomes, we questioned whether

macrosynteny could resolve their phylogenetic position. Substantial evidence has been

presented supporting both their association with Phoronida and Brachiopoda (Lophophorata)

(Nesnidal et al. 2013; Luo et al. 2017; Marlétaz et al. 2019; Laumer et al. 2019) and with

Entoprocta and Cycliophora (Polyzoa) (Hejnol et al. 2009; Kocot et al. 2017; Khalturin et al.

2022). Our study identifies nine fusion-with-mixing events shared between bryozoans and

brachiopods, of which five are confirmed as derived characters, supporting the Lophophorata

topology. Fusion-with-mixing events are essentially irreversible in evolution, but convergent

evolution cannot be ruled out (Steenwyk and King 2024). If these rare genomic changes are

specific to Lophotrochozoans (excluding Entoprocta and Cycliophora), then the presence of

at least nine shared derived fusion events in brachiopods and bryozoans would almost rule

out convergence, becoming a conclusive piece of evidence in this ongoing debate. The future

availability of chromosome-scale genomes from these groups is expected to provide clarity.

Looking further, evidence from macrosynteny is likely to become increasingly prominent in

discussions concerning phylogenetically challenging clades.

Overall, though macrosyntenic gene linkages are conserved across diverse bilaterian

lineages, members of the phylum Bryozoa possess highly derived genomes with scrambling

of these gene sets. Divergence has continued within the phylum, leading to distinct

organizational structures in two bryozoan classes, and ALG fission, previously

underappreciated in its importance, has occurred with high frequency. The addition of

information on the third bryozoan class (the Stenolaemata, represented by present-day cyclostomes) when chromosome-level genomes become available will add considerably to this picture. Furthermore, although we present comparative genomics data on Hox gene duplications, additional studies on the spatial expression patterns of bryozoan Hox genes are necessary to examine how genome rearrangements have affected their functions (Fuchs et al. 2011). We expect the bryozoan genomes annotated in this study to be crucial for the pursuit of such future research avenues.

Methods

Assembly acquisition and gene prediction

Chromosome-scale assemblies of *B. stolonifera* (GenBank accession no. GCA_935421135) (Wood et al. 2023), *Cry. pallasiana* (GCA_945261195) (Bishop et al. 2023b), *M. membranacea* (GCA_914767715) (Bishop et al. 2023a), and *W. subatra* (GCA_963576615) (Bishop et al. 2024) were sequenced as part of the Darwin Tree of Life project (The Darwin Tree of Life Project Consortium et al. 2022) and retrieved from the NCBI Datasets resource. A chromosome-scale assembly of *Cri. mucedo* (Hoencamp et al. 2021) was obtained from the NCBI Gene Expression Omnibus (GSE169088). None of the bryozoan assemblies used have publicly available gene models, so gene prediction was performed in-house. Repetitive elements were first annotated with RepeatModeler (v2.0.4) (Flynn et al. 2020) and masked using RepeatMasker (v4.1.5) in sensitive mode (Smit et al. 2015). Gene prediction and annotation were performed using the BRAKER3 pipeline (v3.0.3), incorporating hints from RNA sequencing (RNA-seq) data as aligned reads in bam format (Stanke et al. 2006, 2008; Li et al. 2009; Barnett et al. 2011; Lomsadze et al. 2014; Buchfink et al. 2015; Hoff et al. 2016, 2019; Gabriel et al. 2024).

BRAKER3 uses RNA-seq and genome data to train the gene prediction tools AUGUSTUS and GeneMark, producing highly supported gene transcripts. Raw RNA-seq reads for gene prediction (Supplemental Table S2) were sourced from the NCBI Sequence Read Archive (SRA) using the SRA Toolkit (v3.0.0) (Leinonen et al. 2011) with GNU parallel (Tange 2021). Transcriptomic data for *B. stolonifera*, *Cry. pallasiana*, *M. membranacea*, and *W. subatra* were generated from adult colonies collected by the Marine Biological Association, Plymouth, UK, in association with the Darwin Tree of Life project. For these four species, RNA extraction was performed on the same colony used for DNA sequencing to generate the reference assembly. For *B. stolonifera* and *M. membranacea,* this dataset was supplemented with independent datasets from developmental stages (Supplemental Table S2). RNA-seq data for *Cri. mucedo* was derived from mature adult colonies from a population different to that

from which the genome assembly was produced (Supplemental Table S2). Quality control for

RNA-seq data was performed with FastQC (Andrews and Others 2010). Reads were trimmed

with Trimmomatic (v0.39) (Bolger et al. 2014) and aligned to the genome with STAR (v2.7.10b)

(Dobin et al. 2013). BUSCO (v5.4.7) (Simão et al. 2015; Manni et al. 2021) was used to

evaluate the quality of gene prediction. Hox genes were identified with reciprocal tBLASTn

searches (Camacho et al. 2009), and orthology was confirmed using gene trees built in IQ-

TREE (v2.2.2.3) (Minh et al. 2020). Hox annotations are consistent with previous work, where

available (Saadi et al. 2023).

Phylogenetics

Single-copy orthologs were identified using OrthoFinder (v2.5.4) (Emms and Kelly 2019).

Alignments for each set of orthologs were created with MAFFT (v7.520) (Katoh et al. 2002;

Katoh and Standley 2013), trimmed with ClipKIT (v1.4.1) (Steenwyk et al. 2020), and

concatenated with PhyKIT (v1.11.7) (Steenwyk et al. 2021). IQ-TREE (v2.2.2.3) (Minh et al.

2020) was used to construct maximum likelihood trees with automated substitution model

selection using ModelFinder (Kalyaanamoorthy et al. 2017) and 1,000 UFBoot2 ultra-fast

bootstrap replicates (Hoang et al. 2018). Trees were rendered with iTOL (Letunic and Bork

2021). A tree of Hox protein sequences was generated using the same IQ-TREE method.

Macrosynteny analysis

We developed SyntenyFinder, a pipeline comprising a collection of Python and R scripts

designed to facilitate macrosynteny analysis using chromosome-level genome assemblies.

SyntenyFinder is a straightforward pipeline with two main functions: first, it uses OrthoFinder

(v2.5.4) (Emms and Kelly 2019) to identify single-copy orthologs in a dataset of protein

sequence files. Second, it utilizes a custom R script to generate chromosome idiograms using

RIdeogram (v0.2.2) (Hao et al. 2020) and to create dot plots showing the positions of

orthologous genes between two genomes. SyntenyFinder is made available on Github

(https://github.com/symgenoevolab/SyntenyFinder) and can be run in two modes: either

automatically downloading and running on annotated genomes from NCBI by their accessions,

or on local files parsed directly to the program. It requires three types of file as an input: (1)

genome sequence FASTA file; (2) gene positions in the format of a GTF file; and (3) a protein

sequence FASTA file.

For the purposes of this paper, two runs were performed: first with only the five

bryozoan species, and second adding *B. floridae* (Chordata), *P. maximus* (Mollusca), *L.

longissimus* (Nemertea) and *L. anatina* (Brachiopoda). In both cases, only orthologs which

were single-copy in all study species were used for downstream analysis. No gene loss or

duplication was permitted. In the second case, to track the genomic position of bilaterian

ALGs, we used an input dataset of the genes used to define the 24 bilaterian ALGs (Simakov
et al. 2022) and used OrthoFinder to find orthologs of these genes in bryozoan genomes. The
restriction to genes with known ALG relationships reduces the noise within the dataset.
Macrosynteny plots for this manuscript were created in R (v4.3.0) (R Core Team 2023). The
same set of genes is used for each comparison in both idiograms and dot plots of
macrosynteny. In idiograms, links between chromosomes with fewer than five shared genes
were removed for visual clarity. These represent single-gene or few-gene translocations. All
genes, including translocated genes, are shown in dot plots. For plotting purposes and to
ensure alignment with orthologous chromosomes, the orientations of chr1, chr2, chr4, chr6,
chr7, chr10, and chr11 in *W. subatra*, and chr2, chr3, chr7, chr9, chr10, chr11, chr12 in *Cry.*
pallasiana were reversed from their orientations as presented in published datasets.
699

Examination of large-scale duplications
Two complementary methods were used to determine whether segmental duplication or whole
genome duplication (WGD) events have occurred in the bryozoan lineage. In each case, we
use the annelid *M. vulgaris* as a positive control for a species with a known WGD (Jin et al.
2020; Lewin et al. 2024a; Vargas-Chávez et al. 2024). First, synonymous substitutions per
site (K_s) distributions were calculated with wgd (v.2.0.26) (Chen et al. 2024). Briefly, paralogs
were identified with 'wgd dmd' using Diamond (v2.0.14.152) (Buchfink et al. 2015, 2021) and
then 'wgd ksd' was run to calculate the K_s distribution using PAML (v4.9) (Yang 2007). The K_s
between two paralogs is an estimate of their divergence time. The null expectation with no
WGD is that the number of paralogs decreases exponentially as K_s increases. When a WGD
occurs, an excess of duplicates are created simultaneously, creating a peak in the distribution
(Tiley et al. 2018). Paralogs situated in collinear blocks in the genome, so-called 'anchor
duplicates', are more likely to be produced by WGD than dispersed paralogs and are therefore
more reliable for WGD detection (Chen et al. 2024). Anchor duplicates were identified using
'wgd syn' to implement the collinearity search algorithm of i-ADHoRe (v3.0.01) (Proost et al.
2011). K_s distributions using only anchor duplicates were then plotted with 'wgd syn' (Chen et
al. 2024).
Second, MCScanX (Wang et al. 2012) was used to identify collinear blocks of
duplicated genes within bryozoan genomes and to classify all duplicates within each genome
as dispersed, proximal, tandem or WGD/segmental. The presence of large homologous
regions within a genome containing collinear blocks of duplicated genes is suggestive of WGD.
The MCScanX output was plotted as a Circos plot (Krzywinski et al. 2009) using shinyCircos
(v2.0) (Wang et al. 2023).
723

Validation of chromosome fission

Given the absence of a high-quality Hi-C contact map for *Cry. pallasiana* (Bishop et al. 2023b),

the fission event in this species remained uncertain. To validate the fission event that led to

the formation of chromosomes CPA10 and CPA12, as shown in Fig. 2, we examined

chromatin interactions by mapping Hi-C data to the chromosome-level assembly. Hi-C reads

for *Cry. pallasiana* were downloaded from the NCBI SRA (accession ERR9866429) using the

SRA Toolkit. The SRA file was fetched with prefetch and converted to FASTQ format using

fasterq-dump. The Hi-C reads were then aligned to the genome assembly (accession

GCA_945261195.1) using BWA (v0.7.17-r1198) (Li 2013), following the Arima Hi-C mapping

pipeline (A160156 v03). A Hi-C contact map was generated using YaHS (v1.2a.1) (Zhou et al.

2023) and JuicerTools (v1.9.9), and visualized with JuiceBox (v1.11.08) (Durand et al. 2016).

Quantification of microsynteny mixing score

To assess the amount of intra-chromosomal gene shuffling, we first calculated Spearman's

rank correlation coefficient (ρ). This coefficient quantifies the strength and direction of

association between the ranked order of orthologous genes along orthologous chromosome

pairs. The Spearman's coefficient is given by the following equation:

$\rho = 1 - (6 * \Sigma d_i^2) / [n(n^2 - 1)]$　　　　　　　　　(1)

where ρ is the coefficient, d_i represents the difference in ranks between corresponding values

of the two variables, and *n* is the total number of observations. Subsequently, we defined the

synteny mixing score, an indicator of chromosomal rearrangement, using the equation:

748　　　　　　　　　　　　$1 - |\rho|$　　　　　　　　　　　　　(2)

where $|\rho|$ denotes the absolute value of the Spearman's coefficient. This metric allows the

synteny mixing score to vary from 0, indicative of no intra-chromosomal rearrangement and

thus a high degree of conserved microsynteny, to 1, representative of maximum

rearrangement, signifying considerable disruption in the original gene order.

Estimation of species divergence time

A phylogenetic tree, constructed above from single-copy orthologs extracted from five

bryozoan genomes and a mollusc outgroup (*P. maximus*), was used to estimate divergence

times. We calibrated this tree with contemporary molecular and fossil data (Orr et al. 2022),

focusing on the divergence between *P. maximus* and *Cri. mucedo* during the Cambrian Period

(541 to 488 million years ago) (Erwin et al. 2011), employing the ape package in R (Paradis

and Schliep 2018).

Quantification of gene expression

Gene expression was quantified in a published *Cri. mucedo* (PRJNA594616) RNA-seq datasets. Raw reads were obtained from the NCBI SRA using the SRA toolkit (Leinonen et al. 2011) and GNU parallel (Tange 2021). Hox gene expression was quantified with Salmon (v1.10.2) (Patro et al. 2017). *Hox4a*, *Hox4b*, *Lox5a*, and *Lox5b* gene expression levels were compared using *t*-tests with a Benjamini-Hochberg correction for multiple testing.

Data access

Gene models for all five bryozoan species have been deposited in Dryad (https://doi.org/10.5061/dryad.76hdr7t3f). Custom Python and R scripts used for macrosynteny analysis, calculating mixing scores, and estimating divergence times are available in our GitHub repository (https://github.com/symgenoevolab/SyntenyFinder) and as Supplemental Code.

Competing interest statement

The authors declare no competing interests.

Acknowledgments

This work was funded by a Royal Society Newton International Fellowship (NIF\R1\201315) and an Academia Sinica Career Development Award (AS-CDA-112-L06) to Y.-J.L. We acknowledge the efforts in sample collection by the Marine Biological Association (MBA)– Darwin Tree of Life collecting team: Patrick Adkins, Robert Mrowicki, Joanna Harley, and Helen Jenkins, as well as Christine Wood of the MBA. We thank the members of the Symbiosis Genomics & Evolution Lab for their assistance and support.

Author contributions

T.D.L. and Y.-J.L. designed research. T.D.L., I.J.-Y.L., M.-E.C., and Y.-J.L. performed research. T.D.L., I.J.-Y.L., J.D.D.B., and Y.-J.L. contributed new reagents/analytic tools. T.D.L. and Y.-J.L. analyzed data. T.D.L., P.W.H.H., and Y.-J.L. wrote the paper. All authors read and approved the final manuscript.

References

Ahola V, Lehtonen R, Somervuo P, Salmela L, Koskinen P, Rastas P, Välimäki N, Paulin L, Kvist J, Wahlberg N, et al. 2014. The Glanville fritillary genome retains an ancient karyotype and reveals selective chromosomal fusions in Lepidoptera. *Nat Commun* **5**: 4737.

Akam M. 1989. Hox and HOM: homologous gene clusters in insects and vertebrates. *Cell* **57**: 347–349.

Andrews S, Others. 2010. FastQC: a quality control tool for high throughput sequence data.

Barnett DW, Garrison EK, Quinlan AR, Strömberg MP, Marth GT. 2011. BamTools: a C++ API and toolkit for analyzing and managing BAM files. *Bioinformatics* **27**: 1691–1692.

Berdan EL, Aubier TG, Cozzolino S, Faria R, Feder JL, Giménez MD, Joron M, Searle JB, Mérot C. 2023. Structural variants and speciation: Multiple processes at play. *Cold Spring Harb Perspect Biol*. http://dx.doi.org/10.1101/cshperspect.a041446.

Bishop J, Adkins P, Wood C, Jenkins H, Marine Biological Association Genome Acquisition Lab, Darwin Tree of Life Barcoding collective, Wellcome Sanger Institute Tree of Life programme, Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective, Tree of Life Core Informatics collective, Darwin Tree of Life Consortium. 2023a. The genome sequence of the sea mat, *Membranipora membranacea* (Linnaeus, 1767). *Wellcome Open Res* **8**: 38.

Bishop J, Wood C, Adkins P, Jenkins H, Marine Biological Association Genome Acquisition Lab, Darwin Tree of Life Barcoding collective, Wellcome Sanger Institute Tree of Life programme, Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective, Tree of Life Core Informatics collective, Darwin Tree of Life Consortium. 2023b. The genome sequence of an encrusting bryozoan, *Cryptosula pallasiana* (Moll, 1803). *Wellcome Open Res* **8**: 128.

Bishop J, Wood CA, Marine Biological Association Genome Acquisition Lab, Darwin Tree of Life Barcoding collective, Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team, Wellcome Sanger Institute Scientific Operations: Sequencing Operations, Wellcome Sanger Institute Tree of Life Core Informatics team, Tree of Life Core Informatics collective, Darwin Tree of Life Consortium. 2024. The genome sequence of the red ripple bryozoan, *Watersipora subatra* (Ortmann, 1890). *Wellcome Open Res* **9**: 458.

Bock PE, Gordon DP. 2013. Phylum Bryozoa Ehrenberg, 1831. *Zootaxa* **3703**: 67–74.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**: 2114–2120.

Buchfink B, Reuter K, Drost H-G. 2021. Sensitive protein alignments at tree-of-life scale
using DIAMOND. *Nat Methods* **18**: 366–368.

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND.
Nat Methods **12**: 59–60.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. *BMC Bioinformatics* **10**: 421.

Cheng C, White BJ, Kamdem C, Mockaitis K, Costantini C, Hahn MW, Besansky NJ. 2012.
Ecological genomics of *Anopheles gambiae* along a latitudinal cline: a population-
resequencing approach. *Genetics* **190**: 1417–1432.

Chen H, Zwaenepoel A, Van de Peer Y. 2024. wgd v2: a suite of tools to uncover and date
ancient polyploidy and whole-genome duplication. *Bioinformatics* **40**: btae272.

Coyle S, Kroll E. 2007. Starvation induces genomic rearrangements and starvation-resilient
phenotypes in yeast. *Mol Biol Evol* **25**: 310–318.

Dean A. 2011. In the loop: long range chromatin interactions and gene regulation. *Brief
Funct Genomics* **10**: 3–10.

de Vos JM, Augustijnen H, Bätscher L, Lucek K. 2020. Speciation through chromosomal
fusion and fission in Lepidoptera. *Philos Trans R Soc Lond B Biol Sci* **375**: 20190539.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–
21.

Duboule D, Dollé P. 1989. The structural and functional organization of the murine HOX
gene family resembles that of *Drosophila* homeotic genes. *EMBO J* **8**: 1497–1505.

Dunham MJ, Badrane H, Ferea T, Adams J, Brown PO, Rosenzweig F, Botstein D. 2002.
Characteristic genome rearrangements in experimental evolution of Saccharomyces
cerevisiae. *Proc Natl Acad Sci USA* **99**: 16144–16149.

Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, Aiden EL. 2016.

Juicebox provides a visualization system for Hi-C contact maps with unlimited zoom.
Cell Syst **3**: 99–101.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol* **20**: 238.

Erwin DH, Laflamme M, Tweedt SM, Sperling EA, Pisani D, Peterson KJ. 2011. The
Cambrian conundrum: early divergence and later ecological success in the early history
of animals. *Science* **334**: 1091–1097.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. 2020.
RepeatModeler2 for automated genomic discovery of transposable element families.
Proc Natl Acad Sci USA **117**: 9451–9457.

Fuchs J, Martindale MQ, Hejnol A. 2011. Gene expression in bryozoan larvae suggest a
fundamental importance of pre-patterned blastemic cells in the bryozoan life-cycle.
Evodevo **2**: 13.

Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, Stanke M. 2024.
BRAKER3: Fully automated genome annotation using RNA-seq and protein evidence
with GeneMark-ETP, AUGUSTUS, and TSEBRA. *Genome Res* **34**: 769–777.

Gordon D, Costello M. 2016. Bryozoa—not a minor phylum. *NZ Sci Rev* **73**: 63–66.

Graham A, Papalopulu N, Krumlauf R. 1989. The murine and Drosophila homeobox gene
complexes have common features of organization and expression. *Cell* **57**: 367–378.

Gregory TR, Mable BK. 2005. CHAPTER 8 - Polyploidy in Animals. In *The Evolution of the*
Genome (ed. T.R. Gregory), pp. 427–517, Academic Press, Burlington.

Haag CR, Roze D. 2007. Genetic load in sexual and asexual diploids: segregation,
dominance and genetic drift. *Genetics* **176**: 1663–1678.

Hao Z, Lv D, Ge Y, Shi J, Weijers D, Yu G, Chen J. 2020. RIdeogram: drawing SVG
graphics to visualize and map genome-wide data on the idiograms. *PeerJ Comput Sci*
6: e251.

Harewood L, Fraser P. 2014. The impact of chromosomal rearrangements on regulation of
gene expression. *Hum Mol Genet* **23**: R76–82.

Hejnol A, Obst M, Stamatakis A, Ott M, Rouse GW, Edgecombe GD, Martinez P, Baguñà J,
Bailly X, Jondelius U, et al. 2009. Assessing the root of bilaterian animals with scalable

phylogenomic methods. *Proc Biol Sci* **276**: 4261–4270.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the
ultrafast bootstrap approximation. *Mol Biol Evol* **35**: 518–522.

Hoencamp C, Dudchenko O, Elbatsh AMO, Brahmachari S, Raaijmakers JA, van Schaik T,
Sedeño Cacciatore Á, Contessoto VG, van Heesbeen RGHP, van den Broek B, et al.
2021. 3D genomics across the tree of life reveals condensin II as a determinant of
architecture type. *Science* **372**: 984–989.

Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2016. BRAKER1: Unsupervised
RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS.
Bioinformatics **32**: 767–769.

Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. 2019. Whole-genome annotation with
BRAKER. *Methods Mol Biol* **1962**: 65–95.

Höök L, Näsvall K, Vila R, Wiklund C, Backström N. 2023. High-density linkage maps and
chromosome level genome assemblies unveil direction and frequency of extensive
structural rearrangements in wood white butterflies (*Leptidea* spp.). *Chromosome Res*
31: 2.

Hubert KA, Wellik DM. 2023. Hox genes in development and beyond. *Development* **150**.
http://dx.doi.org/10.1242/dev.192476.

Ivankovic M, Brand JN, Pandolfini L, Brown T, Pippel M, Rozanski A, Schubert T, Grohme
MA, Winkler S, Robledillo L, et al. 2023. A comparative analysis of planarian genomes
reveals regulatory conservation in the face of rapid structural divergence. *bioRxiv*
2023.12.22.572568. https://www.biorxiv.org/content/10.1101/2023.12.22.572568v1
(Accessed January 17, 2024).

Jenner RA, Littlewood DTJ. 2008. Problematica old and new. *Philos Trans R Soc Lond B*
Biol Sci **363**: 1503–1512.

Jenner RA, Timothy D, Littlewood J. 2009. Invertebrate Problematica: kinds, causes, and
solutions. http://dx.doi.org/10.1093/acprof:oso/9780199549429.003.0011 (Accessed
January 16, 2024).

Jin F, Zhou Z, Guo Q, Liang Z, Yang R, Jiang J, He Y, Zhao Q, Zhao Q. 2020. High-quality
genome assembly of *Metaphire vulgaris*. *PeerJ* **8**: e10313.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**: 587–589.

Katoh K, Misawa K, Kuma K-I, Miyata T. 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**: 3059–3066.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**: 772–780.

Kenny NJ, McCarthy SA, Dudchenko O, James K, Betteridge E, Corton C, Dolucan J, Mead
D, Oliver K, Omer AD, et al. 2020. The gene-rich genome of the scallop *Pecten*
maximus. *Gigascience* **9**: giaa037.

Khalturin K, Shunatova N, Shchenkov S, Sasakura Y, Kawamitsu M, Satoh N. 2022. Polyzoa
is back: The effect of complete gene sets on the placement of Ectoprocta and
Entoprocta. *Sci Adv* **8**: eabo4400.

King M. 1995. *Species Evolution: The Role of Chromosome Change*. Cambridge University
Press.

Kocot KM, Struck TH, Merkel J, Waits DS, Todt C, Brannock PM, Weese DA, Cannon JT,
Moroz LL, Lieb B, et al. 2017. Phylogenomics of Lophotrochozoa with consideration of
systematic error. *Syst Biol* **66**: 256–282.

Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA.
2009. Circos: an information aesthetic for comparative genomics. *Genome Res* **19**:
1639–1645.

Kwiatkowski D, Blaxter M, Darwin Tree of Life Barcoding collective, Wellcome Sanger
Institute Tree of Life programme, Wellcome Sanger Institute Scientific Operations: DNA
Pipelines collective, Tree of Life Core Informatics collective, Darwin Tree of Life
Consortium. 2021. The genome sequence of the bootlace worm, *Lineus longissimus*
(Gunnerus, 1770). *Wellcome Open Res* **6**: 272.

Laumer CE, Fernández R, Lemer S, Combosch D, Kocot KM, Riesgo A, Andrade SCS,
Sterrer W, Sørensen MV, Giribet G. 2019. Revisiting metazoan phylogeny with genomic
sampling of all phyla. *Proc Biol Sci* **286**: 20190831.

Leinonen R, Sugawara H, Shumway M, International Nucleotide Sequence Database
Collaboration. 2011. The sequence read archive. *Nucleic Acids Res* **39**: D19–21.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. *Nucleic Acids Res* **49**: W293–W296.

Lewin TD, Liao IJ-Y, Luo Y-J. 2024a. Annelid comparative genomics and the evolution of
massive lineage-specific genome rearrangement in bilaterians. *Mol Biol Evol* **41**:
msae172.

Lewin TD, Shimizu K, Liao IJ-Y, Chen M-E, Endo K, Satoh N, Holland PWH, Wong YH, Luo
Y-J. 2024b. Brachiopod genome unveils the evolution of the BMP–Chordin network in
bilaterian body patterning. *bioRxiv* 2024.05.28.596352.

Liao IJ-Y, Lu T-M, Chen M-E, Luo Y-J. 2023. Spiralian genomics and the evolution of animal
genome architecture. *Brief Funct Genomics* **22**: 498–508.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. *arXiv [q-bioGN]*. http://arxiv.org/abs/1303.3997.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
R. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–
2079.

Lin C-Y, Marlétaz F, Pérez-Posada A, Martínez-García PM, Schloissnig S, Peluso P,
Conception GT, Bump P, Chen Y-C, Chou C, et al. 2024. Chromosome-level genome
assemblies of 2 hemichordates provide new insights into deuterostome origin and
chromosome evolution. *PLoS Biol* **22**: e3002661.

Lombardi C, Taylor PD, Cocito S. 2020. Bryozoans: The "forgotten" bioconstructors. in
Perspectives on the marine animal forests of the world, pp. 193–217, Springer
International Publishing, Cham.

Lomsadze A, Burns PD, Borodovsky M. 2014. Integration of mapped RNA-Seq reads into
automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Res* **42**: e119.

Loxton J, Wood CA, Bishop JDD, Porter JS, Spencer Jones M, Nall CR. 2017. Distribution of
the invasive bryozoan Schizoporella japonica in Great Britain and Ireland and a review
of its European distribution. *Biol Invasions* **19**: 2225–2235.

Luo Y-J, Kanda M, Koyanagi R, Hisata K, Akiyama T, Sakamoto H, Sakamoto T, Satoh N.
2017. Nemertean and phoronid genomes reveal lophotrochozoan evolution and the
origin of bilaterian heads. *Nat Ecol Evol* **2**: 141–151.

Mackintosh A, de la Rosa PMG, Martin SH, Lohse K, Laetsch DR. 2023a. Inferring inter-
chromosomal rearrangements and ancestral linkage groups from synteny. *bioRxiv*
2023.09.17.558111. https://www.biorxiv.org/content/10.1101/2023.09.17.558111v1
(Accessed January 8, 2024).

Mackintosh A, Vila R, Laetsch DR, Hayward A, Martin SH, Lohse K. 2023b. Chromosome
fissions and fusions act as barriers to gene flow between *Brenthis* fritillary butterflies.
Mol Biol Evol **40**. http://dx.doi.org/10.1093/molbev/msad043.

Mackintosh A, Vila R, Martin SH, Setter D, Lohse K. 2023c. Do chromosome
rearrangements fix by genetic drift or natural selection? Insights from *Brenthis*
butterflies. *Mol Ecol*. http://dx.doi.org/10.1111/mec.17146.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO update: Novel
and streamlined workflows along with broader and deeper phylogenetic coverage for
scoring of eukaryotic, prokaryotic, and viral genomes. *Mol Biol Evol* **38**: 4647–4654.

Marlétaz F, Couloux A, Poulain J, Labadie K, Da Silva C, Mangenot S, Noel B, Poustka AJ,
Dru P, Pegueroles C, et al. 2023. Analysis of the *P. lividus* sea urchin genome highlights
contrasting trends of genomic and regulatory evolution in deuterostomes. *Cell Genom* **3**:
100295.

Marlétaz F, Peijnenburg KTCA, Goto T, Satoh N, Rokhsar DS. 2019. A new spiralian
phylogeny places the enigmatic arrow worms among gnathiferans. *Curr Biol* **29**: 312–
318.e3.

Martín-Zamora FM, Liang Y, Guynes K, Carrillo-Baltodano AM, Davies BE, Donnellan RD,
Tan Y, Moggioli G, Seudre O, Tran M, et al. 2023. Annelid functional genomics reveal
the origins of bilaterian life cycles. *Nature* **615**: 105–110.

Mayr E. 1961. Cause and effect in biology. *Science* **134**: 1501–1506.

Miele A, Dekker J. 2008. Long-range chromosomal interactions and gene regulation. *Mol*
Biosyst **4**: 1046–1057.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear
R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic inference in
the genomic era. *Mol Biol Evol* **37**: 1530–1534.

Moriyama Y, Koshiba-Takeuchi K. 2018. Significance of whole-genome duplications on the
emergence of evolutionary novelties. *Brief Funct Genomics* **17**: 329–338.

Muffato M, Louis A, Nguyen NTT, Lucas J, Berthelot C, Roest Crollius H. 2023.
Reconstruction of hundreds of reference ancestral genomes across the eukaryotic
kingdom. *Nat Ecol Evol* **7**: 355–366.

Nesnidal MP, Helmkampf M, Meyer A, Witek A, Bruchhaus I, Ebersberger I, Hankeln T, Lieb
B, Struck TH, Hausdorf B. 2013. New phylogenomic data support the monophyly of
Lophophorata and an Ectoproct-Phoronid clade and indicate that Polyzoa and
Kryptrochozoa are caused by systematic bias. *BMC Evol Biol* **13**: 253.

Nishizawa A, Sarashina I, Tsujimoto Y, Iijima M, Endo K. 2010. Artificial fertilization, early
development and chromosome numbers in the brachiopod *Lingula anatina*. *Special*
Papers in Palaeontology **84**: 309–316.

Orr HA. 1990. "Why polyploidy is rarer in animals than in plants" revisited. *Am Nat* **136**: 759–
770.

Orr RJS, Di Martino E, Ramsfjell MH, Gordon DP, Berning B, Chowdhury I, Craig S,
Cumming RL, Figuerola B, Florence W, et al. 2022. Paleozoic origins of cheilostome
bryozoans and their parental care inferred by a new genome-skimmed phylogeny. *Sci*
Adv **8**: eabm7452.

Orr RJS, Sannum MM, Boessenkool S, Di Martino E, Gordon DP, Mello HL, Obst M,
Ramsfjell MH, Smith AM, Liow LH. 2021. A molecular phylogeny of historical and
contemporary specimens of an under-studied micro-invertebrate group. *Ecol Evol* **11**:
309–320.

Paradis E, Schliep K. 2018. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**: 526–528.

Parey E, Louis A, Montfort J, Bouchez O, Roques C, Iampietro C, Lluch J, Castinel A,
Donnadieu C, Desvignes T, et al. 2023. Genome structures resolve the early
diversification of teleost fishes. *Science* **379**: 572–575.

Passarge E, Horsthemke B, Farber RA. 1999. Incorrect use of the term synteny. *Nat Genet*
23: 387.

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-
aware quantification of transcript expression. *Nat Methods* **14**: 417–419.

Plessy C, Mansfield MJ, Bliznina A, Masunaga A, West C, Tan Y, Liu AW, Grašič J, Del Río
Pisula MS, Sánchez-Serna G, et al. 2024. Extreme genome scrambling in marine

planktonic *Oikopleura dioica* cryptic species. *Genome Res* **34**: 426–440.

Pratt C, Denley D, Metaxas A. 2022. Ocean warming and multiple source populations
increase the threat of an invasive bryozoan to kelp beds in the northwest Atlantic
Ocean. *Mar Ecol Prog Ser* **695**: 65–81.

Proost S, Fostier J, De Witte D, Dhoedt B, Demeester P, Van de Peer Y, Vandepoele K.
2011. i-ADHoRe 3.0—fast and sensitive detection of genomic homology in extremely
large data sets. *Nucleic Acids Res* **40**: e11–e11.

Putnam NH, Butts T, Ferrier DEK, Furlong RF, Hellsten U, Kawashima T, Robinson-Rechavi
M, Shoguchi E, Terry A, Yu J-K, et al. 2008. The amphioxus genome and the evolution
of the chordate karyotype. *Nature* **453**: 1064–1071.

Putnam NH, Srivastava M, Hellsten U, Dirks B, Chapman J, Salamov A, Terry A, Shapiro H,
Lindquist E, Kapitonov VV, et al. 2007. Sea anemone genome reveals ancestral
eumetazoan gene repertoire and genomic organization. *Science* **317**: 86–94.

R Core Team. 2023. R: A language and environment for statistical computing. Vienna: R
foundation for statistical computing. *Vienna: R foundation for statistical computing*.
https://cir.nii.ac.jp/crid/1370576118723163397 (Accessed January 9, 2024).

Renwick JH. 1971. The mapping of human chromosomes. *Annu Rev Genet* **5**: 81–120.

Rice A, Glick L, Abadi S, Einhorn M, Kopelman NM, Salman-Minkov A, Mayzel J, Chay O,
Mayrose I. 2015. The Chromosome Counts Database (CCDB) - a community resource
of plant chromosome numbers. *New Phytol* **206**: 19–26.

Rieseberg LH. 2001. Chromosomal rearrangements and speciation. *Trends Ecol Evol* **16**:
351–358.

Román-Palacios C, Medina CA, Zhan SH, Barker MS. 2021. Animal chromosome counts
reveal a similar range of chromosome numbers but with less polyploidy in animals
compared to flowering plants. *J Evol Biol* **34**: 1333–1339.

Saadi AJ, de Oliveira AL, Kocot KM, Schwaha T. 2023. Genomic and transcriptomic survey
of bryozoan Hox and ParaHox genes with emphasis on phylactolaemate bryozoans.
BMC Genomics **24**: 711.

Sacerdot C, Louis A, Bon C, Berthelot C, Roest Crollius H. 2018. Chromosome evolution at
the origin of the ancestral vertebrate genome. *Genome Biol* **19**: 166.

Schultz DT, Haddock SHD, Bredeson JV, Green RE, Simakov O, Rokhsar DS. 2023.
Ancient gene linkages support ctenophores as sister to other animals. *Nature* **618**: 110–
117.

Schultz DT, Heath-Heckman EAC, Winchell CJ, Kuo D-H, Yu Y-S, Oberauer F, Kocot KM,
Cho S-J, Simakov O, Weisblat DA. 2024. Acceleration of genome rearrangement in
clitellate annelids. *bioRxiv* 2024.05.12.593736.
https://www.biorxiv.org/content/10.1101/2024.05.12.593736v1 (Accessed July 10,
2024).

Schwaha T. 2020. *Phylum Bryozoa*. ed. T. Schwaha. De Gruyter, Berlin, Germany

Schwaha TF, Ostrovsky AN, Wanninger A. 2020. Key novelties in the evolution of the
aquatic colonial phylum Bryozoa: evidence from soft body morphology. *Biol Rev Camb*
Philos Soc **95**: 696–729.

Simakov O, Bredeson J, Berkoff K, Marletaz F, Mitros T, Schultz DT, O'Connell BL, Dear P,
Martinez DE, Steele RE, et al. 2022. Deeply conserved synteny and the evolution of
metazoan chromosomes. *Sci Adv* **8**: eabi5884.

Simakov O, Marletaz F, Cho S-J, Edsinger-Gonzales E, Havlak P, Hellsten U, Kuo D-H,
Larsson T, Lv J, Arendt D, et al. 2013. Insights into bilaterian evolution from three
spiralian genomes. *Nature* **493**: 526–531.

Simakov O, Marlétaz F, Yue J-X, O'Connell B, Jenkins J, Brandt A, Calef R, Tung C-H,
Huang T-K, Schmutz J, et al. 2020. Deeply conserved synteny resolves early events in
vertebrate evolution. *Nat Ecol Evol* **4**: 820–830.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics **31**: 3210–3212.

Smit AFA, Hubley R, Green P. 2015. RepeatMasker Open-4.0.

Stanke M, Diekhans M, Baertsch R, Haussler D. 2008. Using native and syntenically
mapped cDNA alignments to improve de novo gene finding. *Bioinformatics* **24**: 637–
644.

Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction in eukaryotes
with a generalized hidden Markov model that uses hints from external sources. *BMC*
Bioinformatics **7**: 62.

Steenwyk JL, Buida TJ 3rd, Li Y, Shen X-X, Rokas A. 2020. ClipKIT: A multiple sequence
alignment trimming software for accurate phylogenomic inference. *PLoS Biol* **18**:
e3001007.

Steenwyk JL, Buida TJ, Labella AL, Li Y, Shen X-X, Rokas A. 2021. PhyKIT: a broadly
applicable UNIX shell toolkit for processing and analyzing phylogenomic data.
Bioinformatics **37**: 2325–2331.

Steenwyk JL, King N. 2024. The promise and pitfalls of synteny in phylogenomics. *PLoS Biol*
22: e3002632.

Sun Y, Sun J, Yang Y, Lan Y, Ip JC-H, Wong WC, Kwan YH, Zhang Y, Han Z, Qiu J-W, et
al. 2021. Genomic signatures supporting the symbiosis and formation of chitinous tube
in the deep-sea tubeworm *Paraescarpia echinospica*. *Mol Biol Evol* **38**: 4116–4134.

Tange O. 2021. GNU Parallel 2210822 ('Kabul'). *Zenodo*.
https://play.google.com/store/books/details?id=sKdSDwAAQBAJ.

Taylor PD, Lombardi C, Cocito S. 2015. Biomineralization in bryozoans: present, past and
future: Bryozoan biomineralization. *Biol Rev Camb Philos Soc* **90**: 1118–1150.

Taylor PD, Waeschenbach A. 2015. Phylogeny and diversification of bryozoans.
Palaeontology **58**: 585–599.

Telford MJ, Budd GE, Philippe H. 2015. Phylogenomic insights into animal evolution. *Curr*
Biol **25**: R876–87.

The Darwin Tree of Life Project Consortium, Blaxter M, Mieszkowska N, Palma FD, Holland
P, Durbin R, Richards T, Berriman M, Kersey P, Hollingsworth P, et al. 2022. Sequence
locally, think globally: The Darwin Tree of Life Project. *Proc Natl Acad Sci USA* **119**:
e2115642118.

Tiley GP, Barker MS, Burleigh JG. 2018. Assessing the performance of *Ks* plots for
detecting ancient whole genome duplications. *Genome Biol Evol* **10**: 2882–2898.

Tolhuis B, Palstra RJ, Splinter E, Grosveld F, de Laat W. 2002. Looping and interaction
between hypersensitive sites in the active beta-globin locus. *Mol Cell* **10**: 1453–1465.

Vargas-Chávez C, Benítez-Álvarez L, Martínez-Redondo GI, Álvarez-González L, Salces-
Ortiz J, Eleftheriadi K, Escudero N, Guiglielmoni N, Flot J-F, Novo M, et al. 2024. A
punctuated burst of massive genomic rearrangements by chromosome shattering and

the origin of non-marine annelids. *bioRxiv* 2024.05.16.594344.

https://www.biorxiv.org/content/10.1101/2024.05.16.594344v2.full (Accessed August 10,

2024).

Wang S, Zhang J, Jiao W, Li J, Xun X, Sun Y, Guo X, Huan P, Dong B, Zhang L, et al. 2017.

Scallop genome provides insights into evolution of bilaterian karyotype and

development. *Nat Ecol Evol* **1**: 120.

Wang Y, Jia L, Tian G, Dong Y, Zhang X, Zhou Z, Luo X, Li Y, Yao W. 2023. shinyCircos-

V2.0: Leveraging the creation of Circos plot with enhanced usability and advanced

features. *Imeta* **2**: e109.

Wang Y, Tang H, Debarry JD, Tan X, Li J, Wang X, Lee T-H, Jin H, Marler B, Guo H, et al.

2012. MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and

collinearity. *Nucleic Acids Res* **40**: e49.

Weaver H, Cook P, Bock P, Gordon D. 2018. *Australian Bryozoa Volume 1: Biology,*

Ecology and Natural History. Csiro Publishing.

White MJD. 1954. *Animal cytology & evolution*. Cambridge university press.

Wood ACL, Probert PK, Rowden AA, Smith AM. 2012. Complex habitat generated by marine

bryozoans: a review of its distribution, structure, diversity, threats and conservation.

Aquat Conserv **22**: 547–563.

Wood C, Bishop J, Adkins P, Jenkins H, Marine Biological Association Genome Acquisition

Lab, Wellcome Sanger Institute Tree of Life programme, Wellcome Sanger Institute

Scientific Operations: DNA Pipelines collective, Tree of Life Core Informatics collective,

Darwin Tree of Life Consortium. 2023. The genome sequence of an erect bryozoan,

Bugulina stolonifera (Ryland, 1960). *Wellcome Open Res* **8**: 26.

Wright CJ, Stevens L, Mackintosh A, Lawniczak M, Blaxter M. 2024. Comparative genomics

reveals the dynamics of chromosome evolution in Lepidoptera. *Nat Ecol Evol* **8**: 777–

790.

Wright S. 1941. On the probability of fixation of reciprocal translocations. *Am Nat* **75**: 513–

522.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**:

1586–1591.

Yoshida K, Rödelsperger C, Röseler W, Riebesell M, Sun S, Kikuchi T, Sommer RJ. 2023.
Chromosome fusions repatterned recombination rate and facilitated reproductive
isolation during *Pristionchus* nematode speciation. *Nat Ecol Evol* **7**: 424–439.

Zdobnov EM, von Mering C, Letunic I, Bork P. 2005. Consistency of genome-based methods
in measuring Metazoan evolution. *FEBS Lett* **579**: 3355–3361.

Zhang Z, Zhang Z, Ma J, Taylor PD, Strotz LC, Jacquet SM, Skovsted CB, Chen F, Han J,
Brock GA. 2021. Fossil evidence unveils an early Cambrian origin for Bryozoa. *Nature*
599: 251–255.

Zhou C, McCarthy SA, Durbin R. 2023. YaHS: yet another Hi-C scaffolding tool.
Bioinformatics **39**: btac808.